Title: Allocating Limited Surveillance Effort for Outbreak Detection of Endemic Foot and Mouth

Disease

Authors: Ariel Greiner[1,2], José Luis Herrera-Diestra[3], Michael Tildesley[4], Katriona Shea[1],

Matthew Ferrari[1]

[1]Center for Infectious Disease Dynamics, Pennsylvania State University, State College, PA

16801, USA

[2]Department of Biology, University of Oxford, Oxford OX1 3SZ, UK

[3]Department of Integrative Biology, The University of Texas at Austin, Austin, Texas, United

States of America

[4]Zeeman Institute for Systems Biology & Infectious Disease Epidemiology Research,

Mathematics Institute and School of Life Sciences, University of Warwick, Coventry, United

Kingdom

Abstract:

Foot and Mouth Disease (FMD) affects cloven-hoofed animals globally and has become a major

economic burden for many countries around the world. Countries that have had recent FMD

outbreaks are prohibited from exporting most meat products; this has major economic

consequences for farmers in those countries, particularly farmers that experience outbreaks or

are near outbreaks. Reducing the number of FMD outbreaks in countries where the disease is

endemic is an important challenge that could drastically improve the livelihoods of millions of

people. As a result, significant effort is expended on surveillance; but there is a concern that

uninformative surveillance strategies may waste resources that could be better used on control

management. Rapid detection through sentinel surveillance may be a useful tool to reduce the

scale and burden of outbreaks. In this study, we use an extensive outbreak and cattle shipment

network dataset from the Republic of Türkiye to test three possible strategies for sentinel

surveillance allocation that differ in their data requirements: ranging from low to high data

needs, we allocate limited surveillance to [1] farms that frequently send and receive shipments

of animals (Network Connectivity), [2] farms near other farms with past outbreaks (Spatial

Proximity) and [3] farms that receive many animals from other farms with past outbreaks

(Network Proximity). We determine that all of these surveillance methods find a similar number

of outbreaks – 2-4.5 times more outbreaks than were detected by surveying farms at random.

On average across surveillance efforts, the Network Proximity and Network Connectivity

methods each find a similar number of outbreaks and the Spatial Proximity method always finds

the fewest outbreaks. Since the Network Proximity method does not outperform the other

methods, these results indicate that incorporating both cattle shipment data and outbreak data

provides only marginal benefit over the less data-intensive surveillance allocation methods for

this objective. We also find that these methods all find more outbreaks when outbreaks are rare.

This is encouraging, as early detection is critical for outbreak management. Overall, since the

Spatial Proximity and Network Connectivity methods find a similar proportion of outbreaks, and

are less data-intensive than the Network Proximity method, countries with endemic FMD whose

resources are constrained could prioritize allocating sentinels based on whichever of those two

methods requires less additional data collection.

Keywords: Foot and Mouth Disease, network, endemic disease, epidemiology, surveillance,

management, Republic of Türkiye, cattle, livestock, agriculture

Introduction

Foot and Mouth Disease (FMD) is an acute systemic vesicular disease in cloven-hoofed

animals worldwide, particularly common in domesticated cattle, sheep and goats. Mortality from

FMD is rare but productivity is often severely reduced (1). In some countries like the Republic of

Türkiye, Tanzania and India, FMD is continuously circulating (endemic FMD), while others are

considered free of FMD (2). FMD can present a significant economic burden on countries

because meat export from endemic countries is severely restricted (3). Livestock farmers in

endemic countries risk severe loss of income, either due to reduced productivity of their

livestock due to the disease or because of imposed control policies such as culling ((4); (5)).

The negative effects of FMD are felt at the country-level as well, due to the trade barriers, costs

of measures to prevent and control the disease and general reduction in food security (5).

FMD spreads to nearby farms through direct contact with infected animals, through fomites on

surfaces (sharing of farm equipment, veterinarians or personnel) or by wind (rare) (1). It may

also spread to farms that receive livestock from infected farms through the shipment network, as

the livestock and the vehicles that transport them may carry and spread the disease (1). Thus,

the farms with the highest risk of an FMD outbreak are likely those that are near already

infected farms or are farms that receive cattle from already infected farms.

In epidemic settings, FMD has been controlled and eliminated through a mixture of movement

restrictions, targeted depopulation initiatives and targeted vaccination campaigns ((6);(7)).

Movement restrictions limit or prevent the movement of livestock between farms and to markets,

significantly reducing the rate of spread of FMD in a region. Movement restrictions are often

recommended to be broad (8), but some have called for them to be more targeted to reduce

excess loss of profit and livestock ((9);(10)). Targeted depopulation initiatives and vaccination

campaigns focus on livestock in farms with FMD outbreaks and farms that are a certain distance

away from farms with outbreaks (ring culling/vaccination) (6). Targeted depopulation initiatives

are often found to be the most effective and rely on the existence of effective surveillance

programs to find outbreaks quickly ((11);(12);(13)).

The vast majority of FMD cases occur in countries where FMD is endemic ((14);(15);(16)), and

designing effective control measures for such countries can reduce economic burden and

accelerate progress towards FMD-free status. The number of FMD outbreaks in countries with

endemic FMD fluctuates over time–e.g. in the Republic of Türkiye there were large outbreaks of

serotype A in 2006 and 2011 and serotype O in 2007 and 2010 (17). Control measures in

endemic countries have focused on reducing the prevalence and economic burden (5) of the

disease in all or parts of the country through shipment restrictions and mass (entire regions) and

ring vaccination campaigns ((7);(18)). In the Republic of Türkiye, where FMD is endemic, control

measures aided by an extensive surveillance program managed to reduce the prevalence of the

disease from 45% to 5% between 2008 and 2018 (19). However, many countries with endemic

FMD have less effective surveillance and thus are not able to target their control measures as

effectively ((7);(20)).

If one could monitor the farms with the highest risk of an FMD outbreak in countries with

endemic FMD, it would be easier to manage at low prevalence (i.e., before any particular

outbreak takes hold) (21). This would then delimit the necessary scope of control measures and

prevent FMD from spreading widely. Control measures are costly (20), so pursuing targeted

control measures at low prevalence is both more productive and more cost effective. This has

been demonstrated in FMD-free countries –Tildesley et al. (22) found that vaccinating near

known infected farms would have reduced the size of the 2001 FMD epidemic in the UK and

Schley et al.(9) found that, with sufficient information, targeted movement restrictions in the UK

would have been effective and led to fewer cattle deaths in the 2001 and 2007 epidemics.

Surveillance of farms will help detect outbreaks early, but as surveillance is expensive and time-

consuming, sentinel surveillance strategies are often prioritized. Under a sentinel surveillance

strategy, farms that are thought to be at high risk of infection are identified and asked to report

outbreaks (23). Thus, it is important to identify which farms should be preferentially allocated for

sentinel surveillance in countries with endemic FMD so that early outbreaks can be discovered

before FMD prevalence becomes too high. One method that can help find high-risk farms is to

model a country as a network of nodes (farms) and edges (shipment pathways) along which

diseases may transmit and then use network theory ((24);(25);(26);(27)) and past outbreak data

to determine which locations might be most at risk of outbreaks. Network theory has been used

before to find high risk farms in FMD-free countries experiencing FMD outbreaks. For example,

Dawson et al. (28) used animal movement data from the UK to determine that targeting nodes

with the highest number of animal movements allowed them to accurately predict the size and

spread of simulated FMD-like epidemics.

Much of the theory on how infectious processes spread on networks that informs our

understanding of which nodes are highest risk assumes an epidemic setting (29). It is less clear

which nodes are highest risk for outbreaks in an endemic setting (17). In this study, we analyze

FMD incidence in an endemic setting with a highly resolved network of livestock shipments and

outbreak incidence to evaluate strategies for efficient outbreak detection. We compare the

effectiveness of sentinel surveillance methods informed by shipment network and/or outbreak

incidence information to help identify high risk nodes for FMD in endemic settings.

We focus this study on the Republic of Türkiye, where FMD is endemic, and there is an

extensive surveillance system to help manage FMD outbreaks in domestic cattle. The Republic

of Türkiye's surveillance program records the start date and location (epiunit; defined as a

geographic area containing one or more farms) of all reported outbreaks and records which

cattle were shipped between which epiunits at which time. Guyver-Fletcher et al. (30) used

these data to show that shipment-mediated persistence of FMD was possible and Herrera-

Diestra et al. (17) used these data to show that epiunits that were more central in the network

had a higher risk of an outbreak – demonstrating the usefulness of networks for the persistence

of FMD and for identifying high risk epiunits in the Republic of Türkiye. With these data we can

compare sentinel surveillance allocation based on [1] network properties, [2] spatial proximity to

outbreaks and [3] network proximity to outbreaks.

In this study, we compare different sentinel surveillance allocation methods to determine how

countries with endemic FMD could monitor for FMD to prevent outbreaks from spreading widely.

We find that the three proposed surveillance methods were able to find many more outbreaks

than a non-data-informed (i.e., random) surveillance method that surveyed the same number of

epiunits. This indicates the value of using outbreak incidence data and shipment network data to

allocate limited sentinel surveillance effort and this study provides some general guidelines for

designing sentinel surveillance programs in countries with endemic FMD. Better allocation of

surveillance effort will help countries with endemic FMD better utilize their resources – allowing

for more efficient and less costly management protocols for controlling FMD outbreaks.

Methods

Overview

We used cattle shipment data and outbreak incidence data from January 2007 to July 2012

from the Republic of Türkiye to develop and compare general methods (Fig. 1) for allocating

limited sentinel surveillance effort for outbreak detection of endemic FMD. Using the cattle

shipment data (edgelists with information about the origin, destination, timing and frequency of

cattle shipments), we constructed overlapping two-month directed-weighted cattle shipment

networks (nodes = epiunits, edges = cattle shipment routes) based on the networks calculated

by Herrera-Diestra et al. (17) from the same dataset. We then used the outbreak incidence data,

the geographic epiunit locations and these networks to develop and compare three surveillance

methods (Fig. 1) across seven different levels of surveillance effort to investigate which

method(s) would be the most effective at finding recorded outbreaks and if effectiveness varied

by effort level.

Model System

FMD is continuously circulating in the Republic of Türkiye and, in response, the government has

developed an extensive surveillance program that monitors FMD outbreaks and tracks cattle

movement among farms. For this study, we used data from the Turkish Veterinary authorities,

facilitated by the European Commission for foot-and-mouth disease (EuFMD), who granted us

access to data from the TurkVet database. The data used in this study extended from January

1, 2007 to July 4, 2012 and covered 54,096 epiunits**,** 5,125 outbreaks and 14,261,447 cattle

shipments. The original dataset contained 55,193 rows of epiunits, but 1097 of those rows were

removed because they were empty or contained duplicate epiunits. Epiunits are geographically

defined regions of the Republic of Türkiye that contain comparable numbers of cattle to other

epiunits (31). Epiunits are considered the basic epidemiological units for recording FMD

outbreaks in the Republic of Türkiye (31). In the original dataset, only 40,746 epiunits had a

unique set of coordinates, 15,445 epiunits shared 2,095 coordinates. To avoid removing all of

these epiunits from the dataset, approximate coordinates were found for 2,445 epiunits through

geocoding using the Google Maps Application Programming Interface and coordinates for the

remaining 13,000 epiunits were chosen through restricted sampling from a uniform distribution

(to ensure that the epiunits remained in the correct district, as the district of each epiunit was

known) (additional details in (32)). This dataset describes when an outbreak was reported in a

particular epiunit and thus we define an outbreak at the epiunit level (i.e. every epiunit either has

an outbreak or does not have an outbreak at any given time). For each outbreak, this dataset

contains information on which epiunit experienced the outbreak, the start date of the outbreak

and the serotype of the outbreak. A and O serotypes of FMD are most common in this dataset

and have been continuously endemic in the Republic of Türkiye, while Asia-1 re-emerged in

2011 after being unobserved since 2001 (17). In our dataset, we observed 1479 serotype A

outbreaks, 1784 serotype O outbreaks, 455 Asia 1 outbreaks while 1458 outbreaks did not have

serotype information.

Network Construction

FMD has an incubation time of 2-14 days (33) and a serial interval of 8-9 days (34), thus it could

take up to twenty days for one cow to become infected with FMD and transmit it to another cow.

With that in mind, we constructed directed-weighted networks from two-month overlapping time

periods ((month *t*, month *t+1*) - i.e. network 1: month 1 + month 2, network 2: month 2 + month

3, etc.; *t* : {1-67}) of cattle shipment data. The two-month time period allows us to capture the

start of any outbreaks in month *t+1* that spread from, or were the result of, outbreaks in month *t*,

with allowance for uncertainty in exact start date of either outbreak. Thus, the time period of two

months allowed us to incorporate all of the network connections along which an outbreak that

started in month *t* could have spread to other epiunits. We look at overlapping two-month

networks so outbreaks in every possible month *t* and month *t+1* (within the time period, i.e.

January 2007 is never a month *t+1* and July 2012 is never a month *t*) are considered as both

potential start and end points of transmission.

We constructed these networks by summing up all of the cattle shipment events between every

ordered pair of epiunits *i* and *j* for each two-month time period (i.e. (*i,j*) =/= (*j,i*) where (*i,j*)

represents a shipment of cattle from epiunit i to epiunit j). The weight of each edge in the

network described the frequency of cattle shipments from epiunit *i* to epiunit *j* in that two-month

time period. Note that not all epiunits sent or received shipments of cattle in every two-month

time period (indeed, some did not report cattle shipments throughout the entire January 2007 to

July 2012 period) and so these networks varied in size.

Surveillance Methods

We tested three different surveillance methods (Network Connectivity, Spatial Proximity,

Network Proximity) that were all informed by different data (network information, epiunit location

or outbreak information) (Fig. 1) from the Republic of Türkiye, hence we will refer to these as

'Data-Informed Surveillance Methods'. In Fig. 1 we show a simple example of how we would

select the best two epiunits (out of the ten shown); using the Network Connectivity method we

select the two most highly connected epiunits with no consideration for outbreak location or start

date (Fig. 1a), with the Spatial Proximity method we would select the two epiunits in closest

spatial proximity to the epiunit with an outbreak at that time (Fig. 1b) and with the Network

Proximity method we would select the two epiunits that the epiunit with an outbreak at that time

sends cattle to most frequently (Fig. 1c). We chose to compare these three surveillance

methods as they made adequate use of the data at hand, were all relatively straightforward to

implement, relate to standard reactive control measures for FMD ((6);(7);(11);(12);(18)) and

have been proposed as methods for allocating sentinel surveillance locations in the past for

FMD and other diseases ((28);(35);(36);(37)). We also calculated the effectiveness of searching

randomly-selected epiunits for outbreaks in order to compare the data-informed surveillance

methods against a non-data informed surveillance method.

For each surveillance method we explored 7 different levels of surveillance effort (5%, 10%,

15%, 20%, 25%, 30%, 35%). A surveillance effort level of x means that we surveyed

approximately x% of all epiunits in the dataset using that surveillance method (not just those

found in the two-month network; supplementary material – Fig. S1). The specifics of how x% of

epiunits were selected is described below for each method.

<u>Random Method</u>

As a null model, we selected x% of epiunits at random 10,000 times (x = surveillance effort

level) and assessed how many of those epiunits had an outbreak at any point in our dataset,

and then recorded the mean value across the 10,000 replicates. This surveillance method was

not informed by geographic location data, outbreak data or cattle shipment network data.

<u>Network Connectivity Method</u>

Since FMD may transmit via the cattle shipment network in the Republic of Türkiye, we first

decided to survey the epiunits that were deemed most central in the network (Fig. 1a), as these

experience the most cattle shipments and thus would likely be at higher risk of an outbreak

((24);(25);(27)). We chose 'degree' (number of nodes connected to a focal node in the network)

as our connectivity metric because [1] it is simple to calculate and understand, [2] it forms the

basis of most other centrality metrics (betweenness centrality, eigenvector centrality, strength,

etc.; (38);(39)) and [3] because we found that it performed similarly to all of the other

connectivity metrics we tested, even those that used more information from the two-month

networks (supplementary material – Fig. S2).

To calculate the degree of each epiunit, we computed the number of different epiunits that every

epiunit sent or received cattle from during the two months in total, as summarized by each two-

month overlapping network. We calculated the degree of every node in every two-month

overlapping network and ranked the epiunits from highest degree to lowest degree (epiunits with

the same degree were ordered indiscriminately by taking the default ordering from igraph (40)

(Fig. 1a)). For each surveillance effort level x, we assessed how many of the x% top ranking

epiunits had an outbreak that started in month $t+1$ of each two-month time period.

Herrera-Diestra et al. (17) determined that epiunits that experienced outbreaks were more

central in the cattle shipment network that described all of the cattle shipments from January

2007 to July 2012. Thus, we additionally calculated the degree of each epiunit in the network

that described all of the cattle shipments from January 2007 to July 2012 ('All-Month Network

Connectivity Surveillance method'). We then quantified the proportion of outbreaks (that

occurred at any time in our dataset) detected in the x% of epiunits with the highest degree in

this network, for each surveillance effort level x. We note that the All-Month Network

Connectivity method includes cattle shipments, and thus edges, that may occur after a given

outbreak. Though unrealistic in practice, we present this analysis in the supplementary material

(Fig. S3) for comparison.

Spatial Proximity Method

The second data-informed surveillance method assessed the number of outbreaks that could be

detected through searching epiunits that were geographically close to epiunits that had

experienced recent outbreaks (Fig. 1b). This is a reasonable surveillance method to explore as

FMD is known to transmit between nearby farms through direct contact with infected animals or

through fomites (i.e. passed along through the sharing of equipment, sharing of veterinarians or

face-to-face interactions among humans and cattle in neighbouring epiunits) and many control

methods are based on this known transmission pathway ((6);(11);(12);(13)).

To search epiunits near epiunits with recent outbreaks, we recorded how many outbreaks with

start dates in month $t+1$ occurred in epiunits that were geographically close (within a certain

radius) to epiunits with outbreaks with start dates in month t (outbreak epiunit) (Fig. 1b). For a

given two-month network and level of surveillance effort, x, the radius chosen was the smallest

radius away from each outbreak epiunit that, when summed across all of the outbreak epiunits,

included at least x% of all epiunits (Fig. 1b). For each two-month network and level of

surveillance effort x, we increased the radius in increments of 1/50 degree units until it satisfied

this condition. For every surveillance effort level and every two-month network, one radius was

used to search around every outbreak epiunit.

Thus, if one month t had more outbreak epiunits than another month t, the radius chosen would

likely be smaller for the first month t than for the second at the same surveillance effort level.

This method ensured that we could survey a comparable number of epiunits across months and

across surveillance methods (see supplementary material – Fig. S1 for more details). This

method only used the outbreak data and the geographic locations of the epiunits and did not

account for the existence of network connections among epiunits.

<u>Network Proximity Method</u>

The final data-informed surveillance method assessed how many outbreaks could be detected

through searching epiunits that were closest, in network-space (calculated from the two-month

networks), to epiunits that had experienced outbreaks with start dates in month t (outbreak

epiunits, Fig. 1c). We define the closest in network-space epiunits as those epiunits that receive

the most cattle shipments directly from outbreak epiunits.

For every outbreak epiunit, we ranked the epiunits they sent cattle directly to (Level 1 epiunits,

i.e. first order neighbours) based on the weight of the edges that connected them to the

outbreak epiunit (i.e., number of shipment events from the outbreak epiunit to each level 1

epiunit; edges with more events have higher weight; ranked highest to lowest). If two such

epiunits had the same weight, their order was determined indiscriminately by taking the default

ordering from igraph (40). Once all of the Level 1 epiunits were considered, the epiunits

connected to the Level 1 epiunits (i.e. the Level 2 epiunits or second order neighbours) were

ranked in the same fashion. Additional levels of epiunits were considered as necessary for each

surveillance effort level, x.

For every surveillance effort level and every two-month network, one rank and level combination

was used to search around every outbreak epiunit. The rank and level combination chosen was

the smallest rank and level combination away from each outbreak epiunit that, when summed

across all of the outbreak epiunits, included at least x% of all epiunits (Fig. 1c). Thus, if one

month t had more outbreak epiunits than another month t, the rank and level combination

chosen would likely be smaller for the first month t than for the second at the same surveillance

effort level. We then assessed how many of the top x% of epiunits had an outbreak that started

in month $t+1$ of that two-month time period.

This method again ensured that we could survey a comparable number of epiunits across

months and across surveillance methods (see supplementary material – Fig. S1 for more

details). This method used both the outbreak data and the full weighted, directed cattle

shipment networks. Note that for certain two-month networks, less than 35% of the total epiunits

in the network could be reached from the outbreak epiunits. Thus at the 35% surveillance effort

level, for this surveillance method, there are some months that do not record a 'number of

outbreaks detected' and the average across months is N/A (see supplementary material – Fig.

S4, S5).

Type of Information	(a) Network Connectivity	(b) Spatial Proximity	(c) Network Proximity
Network	✓		✓
Outbreak		✓	✓
Epiunit Location		✓	

Fig 1. *Surveillance Methods Summary* - The three different data-informed surveillance methods are shown diagrammatically with an accompanying description of which types of information they are informed by. For each surveillance method, two magnifying glasses are shown on the epiunits that would have been selected by each surveillance method at a surveillance effort of 20% (2/10 epiunits). Epiunits with outbreaks on them in the diagram indicate epiunits with outbreaks with start dates in month t (i.e. 'outbreak epiunits'). The circle in (b) indicates the search radius chosen at a surveillance effort of 20%. Note that 'Network' and 'Outbreak' information are dynamic and require consistent collection while 'Epiunit Location' information (the latitude and longitude of the centroid of the epiunit) is static and does not change over time. 'Outbreak' information is a record of the epiunit and start date of a particular outbreak. 'Network' information records the start and end epiunit of all cattle shipment events.

Outbreak Detection

We calculated the number of outbreaks detected in month $t+1$ (at every month $t+1$, $t+1$:[2,67]) using each of these surveillance methods. Note that when we refer to the number or percentage of outbreaks detected in a 'month', we are referring to the number detected in month $t+1$ of the (month t, month $t+1$) two-month network. We also calculated the average number of outbreaks detected by each method at each surveillance effort level, across all months for which we had network and outbreak data (January 2007 to July 2012). Additionally, we assessed the number

of outbreaks of serotype O or serotype A detected by serotype-specific versions of each of

these surveillance methods, but since the serotype-specific results were similar to the non-

serotype-specific results we present those results in the supplementary material (Table S2, S3).

Results

Across Months

All three data-informed surveillance methods (Fig. 1) detect similar percentages of outbreaks at

each surveillance effort level but at higher surveillance effort levels the Network Proximity

method and Network Connectivity method detect 5-10% more outbreaks than the Spatial

Proximity method, averaged across time (January 2007 to July 2012, Fig. 2). At the lowest

surveillance effort level (5%), the data-informed surveillance methods detect four to five times

more outbreaks on average than were detected by the random surveillance method (Fig. 2). As

surveillance effort level is increased, the data-informed surveillance methods detect relatively

fewer outbreaks. At 30% surveillance effort level these methods only detect approximately two

times more outbreaks on average than the random surveillance method (Fig. 2). Across

surveillance effort levels, the Spatial Proximity method consistently finds the fewest outbreaks of

the three data-informed surveillance methods on average (Fig. 2). The Network Proximity

method and Network Connectivity method find a similar percentage of outbreaks across

surveillance effort levels on average (Fig. 2).

The Network Connectivity surveillance method computed on the two-month networks detected

more outbreaks on average across months than the All-Month Network Connectivity

surveillance method that was computed on the network generated from the entire time series

(January 2007 to July 2012), across all surveillance effort levels (supplementary material – Fig.

S3). The All-Month Network Connectivity surveillance method finds the smallest percentage of

outbreaks at low surveillance effort level, but the second most (on average) at the highest

surveillance effort level (supplementary material – Fig. S3).

Fig 2. *Performance of Surveillance Methods Across Surveillance Effort Levels* - The points represent the

average percent of outbreaks each surveillance method detected across all *t+1* months at each

surveillance effort level. The ribbons represent the variability across months (the interquartile range).

When we used the data-informed surveillance methods to search for only serotype A or

serotype O outbreaks, they detected a similar percentage of outbreaks averaged across months

as the non-serotype specific surveillance methods (supplementary material – Table S2, S3). In

general, the Network Connectivity method was the best at finding serotype A outbreaks and the

Network Proximity method was the best at finding serotype O outbreaks on average but the

differences in surveillance method performance are fairly minor (supplementary material – Table

S2, S3).

Month by Month

As surveillance effort level was increased, the three different data-informed surveillance

methods all detected more outbreaks per month (Fig. 3; supplementary material – Fig. S4). As

the number of outbreaks in a month decreased, the proportion of outbreaks detected generally

increased for all surveillance methods and across surveillance effort levels (Fig. 3, Fig. 4;

supplementary material – Fig. S4, Fig. S5). For example, when there were fewer than 15

outbreaks in a month, the surveillance methods detected up to 67% of outbreaks at a 5%

surveillance effort level (Fig. 3a) and up to 100% of outbreaks at a 35% surveillance effort level

(supplementary material – Fig. S4d) (across methods). In contrast, when there were more than

50 outbreaks in a month, surveillance only detected up to 44% of outbreaks at a 5%

surveillance effort level (Fig. 3a, Fig. 4a) and up to 89% of outbreaks at a 35% surveillance

effort level (supplementary material – Fig. S4d, Fig. S5d) (across methods). In general, the

three data-informed surveillance methods all showed fairly similar outbreak detection patterns to

each other (Fig. 3, supplementary material – Fig. S4).

Fig 3. *Month-by-Month Surveillance Method Performance* - The bars in each panel show the number of

outbreaks detected by each surveillance method at each surveillance effort level (5%, 15% 30%

respectively) at each *t+1* month. The black line shows the number of outbreaks reported in each *t+1*

month. The *t+1* months are ordered by decreasing number of outbreaks, with the left-most month having

the most outbreaks. Versions of these graphs that correspond to 10%, 20%, 25% and 35% surveillance

effort levels are in the supplementary material (Fig. S4).

Even though on average across months the Network Proximity method and Network

Connectivity method consistently detected more outbreaks than the Spatial Proximity Method

(Fig. 2), none of the data-informed surveillance methods consistently performed better than the

other two for all months across surveillance effort levels (Fig. 4, supplementary material – Fig.

S5). Also, no method was consistently better at detecting outbreaks as the number of outbreaks

per month decreased (Fig. 4, supplementary material – Fig. S5). As surveillance effort level

increased, the Network Connectivity method performed better in more months compared to the

Network Proximity method while the Spatial Proximity method performed better in the fewest

months at every surveillance effort level (Fig. 4, supplementary material – Fig. S5).

Fig 4. *Best Surveillance Methods Each Month* - Each panel shows the percent of outbreaks detected by

the data-informed surveillance method that detected the most outbreaks in each *t+1* month at each

surveillance effort level. The bars indicate the percent of outbreaks detected by the best surveillance

method and the horizontal black line indicates the percent of outbreaks detected by the Random

surveillance method at that surveillance effort level. The coloured bars indicate the data-informed

surveillance method that detected the most outbreaks at that surveillance effort level for that *t+1* month.

The *t+1* months are ordered by declining number of outbreaks, with the left-most month having the most

outbreaks. Versions of these graphs that correspond to 10%, 20%, 25% and 35% surveillance effort

levels are in the supplementary material (Fig. S5).

Discussion

All data-informed surveillance methods detected 2 to 4.5 times more outbreaks than the random

surveillance method. For example, at a 5% surveillance effort level the data-informed methods

detected roughly 20% of the outbreaks (on average), while the random surveillance method only

detected 20% of outbreaks when searching at a 20% surveillance effort level (Fig. 2). None of

the data-informed sentinel surveillance methods were consistently better across surveillance

effort levels or across months than any other method. However, we determined that surveillance

methods informed by network data (Network Connectivity, Network Proximity; Fig. 1) performed

better than the Spatial Proximity method on average across all surveillance effort levels (Fig. 2;

especially at high surveillance effort levels) and across months (Fig. 4; supplementary material

– Fig. S5). As surveillance effort level increased, the surveillance method that relied exclusively

on network properties (Network Connectivity method) improved in comparison to the other

methods (Fig. 2). Gilbert et al. (41) determined that the spread of FMD in the Republic of

Türkiye had become more associated with long-range transportation of animals than with short-

range transmission; a finding that these results support. The surveillance method that used both

network and outbreak information ('Network Proximity'; Fig. 1) was on average the most

effective at small surveillance effort levels, but at larger surveillance effort levels it was not much

better than the less informed methods and it also became limited by the availability of edges in

the network (at 35% surveillance effort level; supplementary material – Fig. S4d, Fig. S5d).

These results imply that knowing where and when outbreaks happen is less crucial when more

epiunits are under surveillance (Fig. 2). Lastly, we determined that all of the data-informed

surveillance methods detected a greater proportion of outbreaks as the number of outbreaks per

month decreased; even the Network Connectivity method, which was not influenced by the

number of outbreaks in a month, showed this pattern. For example, when we used the data-

informed surveillance methods to survey 5% of epiunits, they detected over 60% of outbreaks

when outbreaks were rare (fewer than 15 outbreaks per month) (Fig. 3,4a).

We also found that restricting our analysis to individual serotypes did not improve our ability to

find outbreaks. For example, sentinel surveillance methods that searched for serotype O FMD

outbreaks detected slightly fewer outbreaks overall than serotype A specific surveillance

methods and the non-serotype specific surveillance methods (supplementary material – Table

S2, S3). This result implies that collecting serotype-specific information does not improve our

ability to allocate sentinel surveillance locations and that informed sentinel allocation decisions

could be made from non-serotype-specific information alone.

Additionally, we determined that collecting time-specific shipment network information improved

our ability to find outbreaks. When we used the cattle shipment network calculated from all of

the cattle shipment data (January 2007 to July 2012), the Network Connectivity surveillance

method detected fewer outbreaks than the average number detected by the Network

Connectivity surveillance method informed by only two months of data (supplementary material

– Fig. S3; also holds for the serotype-specific methods–supplementary material – Table S2, S3).

This is likely because the two-month cattle shipment Network Connectivity surveillance method

is only informed by cattle shipments that happened around the date of the potential outbreaks

and is not affected by additional shipments that happened far before or far after the outbreaks

being searched for. Operationally, this implies that countries with endemic FMD should prioritize

collecting shipment data from around the time of the outbreak start date and should be wary of

using shipment data that is from far before or far after an outbreak as it may be misleading at

worst, or exhibit diminishing returns at best.

The results of this study were calculated from data from the Republic of Türkiye, and it is

important to acknowledge that farming demography and animal movement networks may differ

in other countries with endemic FMD, but we can still extrapolate some general suggestions for

designing sentinel surveillance programs in countries with endemic FMD that have limited

resources. Since surveillance methods informed by network data detected more outbreaks at

high surveillance effort levels but roughly the same number of outbreaks as other methods at a

low surveillance effort level (on average), our results would suggest that countries with endemic

FMD and limited surveillance resources should prioritize collecting whichever of those data

sources would require the least additional resources, while countries with endemic FMD and

more resources available for surveillance might prioritize collecting network data (as, at higher

surveillance efforts, these methods detect 5-10% more outbreaks which could be quite

significant, practically speaking). Note that Stage 1 of the Progressive Control Pathway for FMD

developed by FAO and EuFMD (42) encourages building systems to describe and document

outbreaks and cattle movement, thus acknowledging the importance of these two types of data.

In the Republic of Türkiye, many of the cattle shipment network edges were fairly short (most

within the same district, median 28.3 km) which may be why the network-informed methods

performed similarly to the Spatial Proximity method (Fig. 2). If farms are further apart on

average, network information may become even more important as there would likely be less

transmission between neighbouring farms. Our results also indicate that dynamic sentinels

might be ideal; we determined that the best sentinel surveillance methods were sensitive to

time-specific network and outbreak information and thus the best sentinel surveillance locations

changed over time. Since the surveillance methods detected more outbreaks when outbreaks

were rare, it implies that outbreaks may be easier to predict and track using data-informed

methods in countries with endemic FMD when they are rare, which is promising as it is often

most important to find outbreaks when there are fewer of them in order to better control the

disease.

Past studies of sentinel surveillance allocation for FMD and other diseases that used network

data tended to focus on allocating sentinels through a Network Connectivity-type method, and

found similar results. Dawson et al. (28) used livestock transport network data from the UK to

determine that using a Network Connectivity-type method for surveillance allocation finds more

simulated FMD-like outbreaks than various random allocation methods. Interestingly, in Dawson

et al. (28)'s simulated non-endemic setting, the top 20% of nodes detected through the network

connectivity-type method are sufficient to predict epidemic size to within 90% confidence; in our

study, however, the top 20% of epiunits detected using the Network Connectivity method only

encompassed approximately 40% of the outbreaks. This implies that Network Connectivity-type

methods may perform better in simulated epidemic settings (which have perfect observation of

the network and the outbreaks) than it did for realized endemic FMD outbreaks. Frössling et al.

(36) combined serological survey results and cattle movement data to compare a Network

Connectivity-type method and a Network Proximity-type method for allocating herds to survey

for detecting BRSV (bovine respiratory syncytial virus) and BCV (bovine coronavirus) infections,

also finding that surveillance methods informed by network data from the relevant time period

found more infections than random methods (also finding that the Network Connectivity-type

method performed slightly better than the Network Proximity-type method). Vidondo & Voelkl

(37) simulated epidemic outbreaks in cattle farms in Switzerland and determined that dynamic

network measures are better at detecting simulated outbreaks than static network measures,

similar to our finding that the Network Connectivity method informed by the two-month networks

detected more outbreaks than the All-Month Network Connectivity method. Network

Connectivity-type methods have also been shown to be more effective than random methods for

finding outbreaks in endemic settings for other diseases (e.g. Ribeiro-Lima et al. (43) for bovine

tuberculosis).

The data that we have from the Republic of Türkiye contain very detailed information on the

reported outbreaks, and on shipments of cattle between January 2007 and July 2012. However,

unreported outbreaks or shipments could limit the potential for outbreak detection in our

analysis. We assume that we are missing either outbreaks or cattle movements or both because

there were some outbreaks that were not detected by all three surveillance methods

(supplementary material – Fig. S6), implying the existence of either un-reported outbreaks that

could bridge the gaps between reported outbreaks or missing connections between epiunits.

We also only had the exact locations of 40,746 out of 54,096 epiunits, which likely impeded the

effectiveness of the Spatial Proximity method. However, as one is unlikely to encounter a more

complete dataset from a country with endemic FMD it is important to work around these data

limitations to derive findings to inform surveillance allocations in other, more resource-limited,

countries with endemic FMD.

Further, we know that the Republic of Türkiye recommends certain control measures (19) but

we do not know to what extent those control measures were enacted around the country from

January 2007 to July 2012, so we did not factor this into our sentinel surveillance method

designs. It is possible that control measures in the Republic of Türkiye managed to stop the

spread of FMD outbreaks to certain epiunits between January 2007 and July 2012 that would

otherwise have experienced outbreaks, which would make all of our sentinel surveillance

methods appear less effective. This uncertainty may also restrict the applicability of our results

to other countries with endemic FMD that have fewer control measures in place, as FMD may

spread differently in them. However, it is likely that our data-informed sentinel surveillance

methods would perform better in countries with no FMD control measures, because the

methods we propose here do not account for the presence of control measures.

In this study, we focus on using each sentinel surveillance method independently so that we can

easily control the surveillance effort level and assess how well each method performs alone.

However, it would be possible to develop a composite measure of outbreak risk using multiple

methods at the same time (e.g. Dawson et al. (28) and Kendall et al. (44) suggest a hybrid of

the Network Connectivity and Network Proximity methods) or varying the method we use over

time (adaptive management framework, (45)). It is possible that combining surveillance methods

(either across time or at one time) might also help reduce the number of outbreaks detected by

no method (supplementary material – Fig. S6) and also allow us to survey more than 35% of

epiunits consistently. In this study, to ensure direct comparison of each surveillance method, we

could only explore surveillance efforts up to 35% as the Network Proximity method was unable

to survey 35% of the network in certain months (supplementary material – Fig. S4, S5).

Since all of the data-informed sentinel surveillance methods performed comparably and the

most data-informed method was only marginally better and was limited in the number of

candidate epiunits it detected, the results of this study suggest that sentinel surveillance sites in

countries with endemic FMD could usefully be allocated through Spatial Proximity or Network

Connectivity surveillance methods (whichever would require the least additional effort to collect

the requisite information). However, based on these results, if a country is able to allocate

sufficient resources to ensure a high surveillance effort, or is planning to increase the effort it

allocates to surveillance in the future, it would make sense to prioritize building a network-data-

motivated surveillance system. Significant effort is expended on surveillance efforts, so finding

the most efficient locations to survey is crucial to ensure that countries do not waste effort

surveying farms that are relatively unlikely to experience outbreaks, especially if such effort

could be better spent on other forms of management to help reduce future FMD outbreaks.

Funding

AG, KS, MF were supported by NSF-NIH-NIFA (National Science Foundation - National

Institutes of Health - National Institute of Food and Agriculture) Ecology and Evolution of

Infectious Disease (https://new.nsf.gov/funding/opportunities/ecology-evolution-infectious-

diseases-eeid) award DEB 1911962. AG was also supported by a Natural Sciences and

Engineering Research Council of Canada (NSERC) Postdoctoral Fellowship (https://www.nserc-

crsng.gc.ca/students-etudiants/pd-np/pdf-bp_eng.asp). MT was funded by a Biotechnology and

Biological Sciences Research Council (BBSRC, https://www.ukri.org/councils/bbsrc/) grant

BB/T004312/1. The funders had no role in study design, data collection and analysis, decision

to publish, or preparation of the manuscript.

Data Availability

The authors do not have permission to share the location data required to reproduce the results

of the Spatial Proximity method, so the authors provide randomized location data instead to

allow the code for that method to be run. The source code and data used to produce the results

and analyses presented in this manuscript are available at

https://github.com/ArielGreiner/FMDLimitedSurveillance/ and at 10.25446/oxford.26488138. All

data queries should be made by contacting the SAP Institute in the Republic of Türkiye

(abdulnaci.bulut@tarimorman.gov.tr).

Competing Interests

606 The authors have declared that no competing interests exist.

Author Contributions

609 AG led the design of the study, ran the model and wrote the first draft of the manuscript. JLH-D,

610 MF and KS contributed to the study concept and design, helped with the analyses and

611 commented on previous versions of the manuscript. MT contributed to the study concept and

design and acquired the agricultural data through agreement with the SAP Institute of the

Republic of Türkiye and EuFMD. All authors read and approved the final manuscript.

References

1. Alexandersen S, Zhang Z, Donaldson AI, Garland AJM. The Pathogenesis and Diagnosis of Foot-and-Mouth Disease. Journal of Comparative Pathology. 2003 Jul;129(1):1–36.

2. Veterinary Services UA. Foot-and-Mouth Disease Response Plan: The Red Book. 2020.

3. James AD, Rushton J. The economics of foot and mouth disease. Revue scientifique et technique (International Office of Epizootics). 2003 Jan 1;21:637–44.

4. Knight-Jones TJD, Rushton J. The economic impacts of foot and mouth disease – What are they, how big are they and where do they occur? Preventive Veterinary Medicine. 2013 Nov;112(3–4):161–73.

5. Knight-Jones TJD, McLaws M, Rushton J. Foot-and-Mouth Disease Impact on Smallholders - What Do We Know, What Don't We Know and How Can We Find Out More? Transboundary and Emerging Diseases. 2017;64(4):1079–94.

6. Affairs GBD for E Food and Rural, Anderson I. Foot and mouth disease 2007: a review and lessons learned. The Stationery Office; 2008. 144 p.

7. Sutmoller P, Barteling SS, Olascoaga RC, Sumption KJ. Control and eradication of foot-and-mouth disease. Virus Research. 2003 Jan 1;91(1):101–44.

8. Anonymous. Council Directive 2003/85/EC on Community measures for the control of foot-and-mouth disease repealing Directive 85/511/EEC and Decisions 89/531/EEC and 96/665/EEC and amending Directive 92/46/EEC. Article 21 & 45. Off J Eur Union. 2003;L306.

9. Schley D, Gubbins S, Paton DJ. Quantifying the Risk of Localised Animal Movement Bans for Foot-and-Mouth Disease. Gravenor MB, editor. PLoS ONE. 2009 May 8;4(5):e5481.

10. Tildesley MJ, Brand S, Brooks Pollock E, Bradbury NV, Werkman M, Keeling MJ. The role of movement restrictions in limiting the economic impact of livestock infections. Nat Sustain. 2019 Sep;2(9):834–40.

11. Ferguson NM, Donnelly CA, Anderson RM. The Foot-and-Mouth Epidemic in Great Britain: Pattern of Spread and Impact of Interventions. Science. 2001 May 11;292(5519):1155–60.

12. Ferguson NM, Donnelly CA, Anderson RM. Transmission intensity and impact of control policies on the foot and mouth epidemic in Great Britain. Nature. 2001 Oct;413(6855):542–8.

13. Tildesley MJ, Bessell PR, Keeling MJ, Woolhouse MEJ. The role of pre-emptive culling in the control of foot-and-mouth disease. Proceedings of the Royal Society B: Biological Sciences. 2009 Jul;276(1671):3239–48.

14.   Ringa N, Bauch CT. Dynamics and control of foot-and-mouth disease in endemic
countries: A pair approximation model. Journal of Theoretical Biology. 2014 Sep 21;357:150–
9.

15.   Schnell PM, Shao Y, Pomeroy LW, Tien JH, Moritz M, Garabed R. Modeling the role of
carrier and mobile herds on foot-and-mouth disease virus endemicity in the Far North Region
of Cameroon. Epidemics. 2019 Dec 1;29:100355.

16.   Zaheer MU, Salman MD, Steneroden KK, Magzamen SL, Weber SE, Case S, et al.
Challenges to the Application of Spatially Explicit Stochastic Simulation Models for Foot-and-
Mouth Disease Control in Endemic Settings: A Systematic Review. Computational and
Mathematical Methods in Medicine. 2020;2020(1):7841941.

17.   Herrera-Diestra JL, Tildesley M, Shea K, Ferrari MJ. Cattle transport network predicts
endemic and epidemic foot-and-mouth disease risk on farms in Turkey. Hill AL, editor. PLoS
Comput Biol. 2022 Aug 19;18(8):e1010354.

18.   Sumption K, Domenech J, Ferrari G. Progressive control of FMD on a global scale.
Veterinary Record. 2012 Jun;170(25):637–9.

19.   F.A.O. Foot-and-Mouth Disease: Quarterly Report. Rome January–March. 2023;2023.

20.   F.A.O., O.I.E. The global foot and mouth disease control strategy: strengthening animal
health systems through improved control of major diseases.

21.   Caporale V, Giovannini A, Zepeda C. Surveillance strategies for foot and mouth disease
to prove absence of disease and absence of viral circulation. Revue scientifique et technique
(International Office of Epizootics). 2012 Dec 1;31:747–59.

22.   Tildesley MJ, Savill NJ, Shaw DJ, Deardon R, Brooks SP, Woolhouse MEJ, et al.
Optimal reactive vaccination strategies for a foot-and-mouth outbreak in the UK. Nature.
2006 Mar;440(7080):83–6.

23.   Teutsch SM, Churchill RE. Principles and Practice of Public Health Surveillance. Oxford
University Press; 2000. 422 p.

24.   Bai Y, Yang B, Lin L, Herrera JL, Du Z, Holme P. Optimizing sentinel surveillance in
temporal network epidemiology. Sci Rep. 2017 Jul 6;7(1):4804.

25.   Colman E, Holme P, Sayama H, Gershenson C. Efficient sentinel surveillance strategies
for preventing epidemics on networks. Paolotti D, editor. PLoS Comput Biol. 2019 Nov
25;15(11):e1007517.

26.   Karsai M, Perra N. Control Strategies of Contagion Processes in Time-Varying
Networks. In: Masuda N, Holme P, editors. Temporal Network Epidemiology [Internet].
Singapore: Springer; 2017 [cited 2024 Jul 30]. p. 179–97. Available from:
https://doi.org/10.1007/978-981-10-5287-3_8

27.   Schirdewahn F, Lentz HHK, Colizza V, Koher A, Hövel P, Vidondo B. Early warning of
infectious disease outbreaks on cattle-transport networks. Clegg S, editor. PLoS ONE. 2021
Jan 6;16(1):e0244999.

28.   Dawson PM, Werkman M, Brooks-Pollock E, Tildesley MJ. Epidemic predictions in an
imperfect world: modelling disease spread with partial data. Proceedings of the Royal Society
B: Biological Sciences. 2015 Jun 7;282(1808):20150205.

29.   Newman MEJ. Spread of epidemic disease on networks. Phys Rev E. 2002 Jul
26;66(1):016128.

30.   Guyver‐Fletcher G, Gorsich EE, Tildesley MJ. A model exploration of carrier and
movement transmission as potential explanatory causes for the persistence of
foot‐and‐mouth disease in endemic regions. Transbounding Emerging Dis. 2022
Sep;69(5):2712–26.

31.   Dawson PM. On the analysis of livestock networks and the modelling of foot-and-mouth
disease [Internet] [phd]. University of Warwick; 2016 [cited 2024 Jul 30]. Available from:
http://webcat.warwick.ac.uk/record=b3062082~S15

32.   Guyver-Fletcher GD. Investigating the efficacy of vaccine strategies in Turkey using a
mathematical epidemiological model [Internet] [phd]. University of Warwick; 2022 [cited 2024
Jul 30]. Available from: http://webcat.warwick.ac.uk/record=b3877301

33.   Fisher SSA. Foot-and-Mouth Disease (FMD) Response Ready Reference Guide -
Etiology and Ecology. 2020;

34.   Yoon H, Jeong W, Choi J, Kang YM, Park HS. Epidemiology and investigation of foot-
and-mouth disease (FMD) in the Republic of Korea. Epidemiology of Communicable and
Non-Communicable Diseases Attributes of Lifestyle and Nature on Humankind. 2016;3–18.

35.   Klinkenberg D, Fraser C, Heesterbeek H. The Effectiveness of Contact Tracing in
Emerging Epidemics. PLOS ONE. 2006 Dec 20;1(1):e12.

36.   Frössling J, Ohlson A, Björkman C, Håkansson N, Nöremark M. Application of network
analysis parameters in risk-based surveillance – Examples based on cattle trade data and
bovine infections in Sweden. Preventive Veterinary Medicine. 2012 Jul 1;105(3):202–8.

37.   Vidondo B, Voelkl B. Dynamic network measures reveal the impact of cattle markets and
alpine summering on the risk of epidemic outbreaks in the Swiss cattle population. BMC Vet
Res. 2018 Mar 13;14(1):88.

38.   Rayfield B, Fortin MJ, Fall A. Connectivity for conservation: a framework to classify
network measures. Ecology. 2011;92(4):847–58.

39.   Dale MRT, Fortin MJ. Spatial Analysis: A Guide For Ecologists. Cambridge University
Press; 2014. 453 p.

40.   Csárdi G, Nepusz T, Traag V, Horvát S, Zanini F, Noom D, et al. igraph: Network
Analysis and Visualization in R. doi: 10.5281/zenodo. 7682609, R package version 1.5. 1.
2023.

41.   Gilbert M, Aktas S, Mohammed H, Roeder P, Sumption K, Tufan M, et al. Patterns of
spread and persistence of foot-and-mouth disease types A, O and Asia-1 in Turkey: a meta-
population approach. Epidemiology & Infection. 2005;133(3):537–45.

42.   de la Santé Animale OM. The progressive control pathway for foot-and-mouth disease
control (PCP-FMD): principles, stage descriptions and standards. 2018 [cited 2024 Jul 30];
Available from: https://www.sidalc.net/search/Record/unfao:840354/Description

43.   Ribeiro-Lima J, Enns EA, Thompson B, Craft ME, Wells SJ. From network analysis to
risk analysis—An approach to risk-based surveillance for bovine tuberculosis in Minnesota,
US. Preventive Veterinary Medicine. 2015 Mar 1;118(4):328–40.

44.   Kendall C, Kerr LRFS, Gondim RC, Werneck GL, Macena RHM, Pontes MK, et al. An
Empirical Comparison of Respondent-driven Sampling, Time Location Sampling, and
Snowball Sampling for Behavioral Surveillance in Men Who Have Sex with Men, Fortaleza,
Brazil. AIDS Behav. 2008 Jul 1;12(1):97–104.

45.   Shea K, Tildesley MJ, Runge MC, Fonnesbeck CJ, Ferrari MJ. Adaptive Management
and the Value of Information: Learning Via Intervention in Epidemiology. Dobson AP, editor.
PLoS Biol. 2014 Oct 21;12(10):e1001970.