E. coli phylogeny drives co-amoxiclav resistance through variable

expression of *bla*_{TEM-1}

William Matlock[1,2,*], Gillian Rodger[1,2,*], Emma Pritchard[1,2], Matthew Colpus[1], Natalia

Kapel[1], Lucinda Barrett[3], Marcus Morgan[3], Sarah Oakley[3], Katie L. Hopkins[4], Aysha

Roohi[1,2], Drosos Karageorgopoulos[3], Matthew B. Avison[5], A. Sarah Walker[1,2], Samuel

Lipworth[1,2,3,†], Nicole Stoesser[1,2,3,†]

[1] Nuffield Department of Medicine, University of Oxford, Oxford, UK

[2] The National Institute for Health Research Health Protection Research Unit in Healthcare Associated

Infections and Antimicrobial Resistance at the University of Oxford, Oxford, UK

[3] Oxford University Hospitals NHS Foundation Trust, Oxford, UK

[4] UK Health Security Agency, Colindale, UK

[5] School of Cellular and Molecular Medicine, University of Bristol, Bristol, UK

* Contributed equally

† Contributed equally

Correspondence: William Matlock (william.matlock@ndm.ox.ac.uk) and Nicole Stoesser

(nicole.stoesser@ndm.ox.ac.uk)

Abstract

Co-amoxiclav resistance in *E. coli* is a clinically important phenotype associated with increased mortality. The class A beta-lactamase $bla_{\text{TEM-1}}$ is often carried by co-amoxiclav-resistant pathogens, but exhibits high phenotypic heterogeneity, making genotype-phenotype predictions challenging. We present a curated dataset of n=377 *E. coli* isolates representing all 8 known phylogroups, where the only acquired beta-lactamase is $bla_{\text{TEM-1}}$. For all isolates, we generate hybrid assemblies and co-amoxiclav MICs, and for a subset (n=67/377), $bla_{\text{TEM-1}}$ qPCR expression data. First, we test whether certain *E. coli* lineages are intrinsically better or worse at expressing $bla_{\text{TEM-1}}$, for example, due to lineage differences in regulatory systems, which are challenging to directly quantify. Using genotypic features of the isolates ($bla_{\text{TEM-1}}$ promoter variants and copy number), we develop a hierarchical Bayesian model for $bla_{\text{TEM-1}}$ expression that controls for phylogeny. We establish that $bla_{\text{TEM-1}}$ expression intrinsically varies across the phylogeny, with some lineages (e.g. phylogroups B1 and C, ST12) better at expression than others (e.g. phylogroups E and F, ST372). Next, we test whether phylogenetic variation in expression influences the resistance of the isolates. With a second model, we use genotypic features ($bla_{\text{TEM-1}}$ promoter variants, copy number, duplications; *ampC* promoter variants; efflux pump AcrF presence) to predict isolate MIC, again controlling for phylogeny. Lastly, we use a third model to demonstrate that the phylogenetic influence on $bla_{\text{TEM-1}}$ expression causally drives the variation in co-amoxiclav MIC. This underscores the importance of incorporating phylogeny into genotype-phenotype predictions, and the study of resistance more generally.

Introduction

The class A beta-lactamase bla_{TEM-1} was first identified in 1965 in a clinical *Escherichia coli* isolate[1]. Originally, it was mobilised by two of the earliest named transposons, Tn*2* and Tn*3*, located on plasmids[2]. In the decades since the genetic context of bla_{TEM-1} has evolved[3], and other mobile genetic elements such as IS*26*[4] and a diverse array of plasmids[5] contribute to its dissemination. At the time of writing, NCBI contains over 170,000 unique isolates carrying bla_{TEM-1} distributed across 28 genera, including the common clinical pathogens *Escherichia coli*, *Klebsiella pneumoniae*, and *Acinetobacter baumannii*[6]. The emergence and dissemination of beta-lactam resistance has been a major healthcare challenge[7], and bla_{TEM-1} represents a key example.

In the UK, beta-lactam and beta-lactamase inhibitor combinations such as co-amoxiclav (amoxicillin and clavulanic acid), are commonly used as a first-line treatment for severe infections[8]. For Enterobacterales, the current EUCAST co-amoxiclav Minimum Inhibitory Concentration (MIC) breakpoint for resistance is 8/2 µg/mL[9] across all indications, with a recent study concluding that empiric co-amoxiclav treatment of *E. coli* bacteraemia with MICs $>$ 32/2 µg/mL was associated with significantly higher mortality in *E. coli* bacteraemia[10]. However, the carriage of bla_{TEM-1} is associated with high phenotypic heterogeneity, making genotype-phenotype predictions challenging.

Small-scale, experimental bla_{TEM-1} systems have demonstrated that the interplay of location (plasmid or chromosome) and copies in the genome, through varying dosage, contributes to variable resistance[11,12]. In addition, other determinants such as mutations in the promoter of bla_{TEM-1}[13] and the chromosomally intrinsic *ampC* gene[14], and efflux pumps[15], are associated with *E. coli* beta-lactam resistance. Moreover, different regulatory systems[16,17], epistasis

(interaction between genes)[18,19], and epigenetics (heritable phenotypic changes without

alterations to the underlying DNA sequence)[20], might also influence co-amoxiclav resistance.

For example, five different *E. coli* strains carrying the same pLL35 plasmid (which carries

$bla_{CTX-M-15}$ and $bla_{TEM-112}$) varied in cefotaxime resistance[21]. Likewise, the introduction of a

pOXA-48 plasmid to six different *E. coli* strains resulted in variable co-amoxiclav

resistance[22]. This indicates that strain background plays a role in resistance.

Successful genotype-to-phenotype prediction requires a comprehensive understanding of not

only individual resistant determinants but also their combined effects. Moreover, this

understanding must be translated to clinically relevant pathogens. Yet, to accurately model

resistance in these systems, a large sample of linked genomic and phenotypic data is required,

which until recently has been limited by sequencing technology and costs.

In this study, we curated and completely reconstructed the genomes of nearly 400 clinical *E.*

coli bacteraemia isolates to reflect a "real-world" but relatively simple genetic scenario where

the only known acquired beta-lactamase gene identified was bla_{TEM-1}, all identical at the

amino acid level. We quantified the co-amoxiclav MICs for these isolates and generated

bla_{TEM-1} qPCR expression data for a subset. We then modelled bla_{TEM-1} expression and co-

amoxiclav MIC whilst controlling confounding genetic mechanisms and chromosomal

phylogeny.

Materials and methods

Isolate selection

We considered $n=548$ candidate *E. coli* bacteraemia isolates cultured from patients presenting

to Oxford University Hospitals NHS Foundation Trust between 2013–2018, and selected

from a larger study of systematically sequenced isolates based on screening their short-read

only assemblies with NCBIAMRFinder (v. 3.11.2) for bla_{TEM-1}, and the absence of other

beta-lactamases[23].

DNA extraction and sequencing

Sub-cultures of isolate stocks, stored at -80°C in 10 % glycerol nutrient broth, were grown

on Columbia blood agar (CBA) overnight at 37°C. DNA was extracted using the EasyMag

system (bioMerieux) and quantified using the Broad Range DNA Qubit kit (Thermo Fisher

Scientific, UK). DNA extracts were multiplexed as 24 samples per sequencing run using the

Oxford Nanopore Technologies (ONT) Rapid Barcoding kit (SQK-RBK110.96) according to

the manufacturer's protocol. Sequencing was performed on a GridION using version FLO-

MIN106 R9.4.1 flow cells with MinKNOW software (v. 21.11.7) and basecalled using

Guppy (v. 3.84). Short-read sequencing was performed on the Illumina HiSeq 4000 pooling

192 isolates per lane, generating 150bp paired end-reads[24].

Dataset curation and genome assembly

Full details are given in Supplementary File 1. Briefly, short- and long-read quality control

used fastp (v. 0.23.4) and filtlong (v. 0.2.1), respectively[25,26]. We also used Rasusa (v. 0.7.1)

on n=3/548 long-read sets due to memory constraints[27]. Genome assembly used Flye (v.

2.9.2-b1786) with bwa (v. 0.7.17-r1188) and Polypolish (v. 0.5.0), and Unicycler (v. 0.5.0)

which uses SPAdes (v. 3.15.5), miniasm (v. 0.3-r179), and Racon (v. 1.5.0)[28–34]. Plasmid

contig validation used Mash screen (v. 2.3) with PLSDB (v. 2023_06_23_v2)[35,36]. All

assemblies were annotated with NCBIAMRFinder (v. 3.11.26 and database v. 2023-11-

15.1)[23]. Alongside, we validated the presence of bla_{TEM-1} using tblastn (v. 2.15.0+) with the

NCBI Reference Gene Catalog TEM-1 RefSeq protein WP_000027057.1 and 100% amino

acid identity[37]. Following genome assembly, we removed $n=171/548$ isolates, either because

(i) the chromosome did not circularise (116/171), (ii) it carried a non-bla_{TEM-1} bla_{TEM} variant

and/or an additional acquired beta-lactamase (54/171), or (iii) the chromosome was too short

consistent with misassembly (~3.5Mbp; 1/171). This left a final dataset of $n=377$ isolates.

Antibiotic susceptibility testing

125   Antibiotic susceptibility testing was performed using the BD Phoenix™ system in

126   accordance with the manufacturers' instructions, generating MICs for co-amoxiclav.

Generation of cDNA template

RNA extraction and DNase treatment were performed on replicates of each isolate ($n=3$

biological/$n=3$ technical) as described previously[38]. RNA was quantified post DNase

treatment using Broad Range RNA Qubit kit (Thermo Fisher Scientific, UK), normalised to 1

μg and reverse transcribed to cDNA using SuperScript IV VILO (Thermo Fisher Scientific,

UK) under the following conditions: 25°C for 10 minutes, 42°C for 60 minutes and 85°C for

5 minutes.

qPCR quantification of bla_{TEM-1} expression

bla_{TEM-1} expression was quantified in a selection of isolates: initially $n=35$ isolates in

triplicate, referred to as batch 1; then a further $n=48$ isolates in duplicate referred to as batch

2. Batch 1 were randomly selected stratified by MICs ($n=2$ MIC $<=2$, $n=5$ MIC 4/2, $n=9$ MIC

8/2, $n=10$ MIC 16/2, $n=4$ MIC 32/2, $n=5$ MIC $>32/2$). Batch 2 was enhanced for specific

bla_{TEM-1} promoter mutations, selecting all isolates with a single bla_{TEM-1} gene with C32T

(with or without a G146A mutation) that had not already been tested, and then randomly

selecting from other wildtype and G146A, single bla_{TEM-1} gene isolates. For all qPCR

reactions, *E. coli* cDNA was normalised to 1ng and amplified in a duplex qPCR reaction

targeting bla_{TEM-1} and 16S. qPCR standard curves were prepared for both bla_{TEM-1} (Genbank

Accession: DQ221255.1) and 16S (Genbank Accession: LC747145.1) sequences cloned into

pMX vectors (Thermo Fisher Scientific, UK). Tenfold dilutions of linearised plasmids (1-

1×10^7 copies/reaction) were used as a standard curve for each experiment. Both curves were

linear in the range tested (16S: $R^2 > 0.991$; TEM-1: $R^2 > 0.91$,). The slopes of the standard

curves for 16S and bla_{TEM-1} were -3.607 and -3.522 respectively. qPCR was performed using

a custom 20 µl TaqMan gene expression assay consisting of TaqMan™ Multiplex Master

Mix, TaqMan unlabelled primers and a TaqMan probe with dye label (FAM for TEM-1 and

VIC for 16S) carried out on the QuantStudio5™ real-time PCR system (Thermo Fisher

Scientific, UK). Cycling conditions were 95°C for 20 seconds, followed by 40 cycles of 95°C

for 3 □ seconds and 60°C for 30 seconds, with Mustang purple as the passive reference. For

batch 1, triplicate samples were analysed and standardized against 16S rRNA gene

expression. Triplicate reactions for each isolate demonstrated good reproducibility for batch 1

(Figure S1). Of note, for isolate OXEC-75, TEM-1 expression was very low-level, and 5

reactions (1 technical replicate for biological replicate 1, 1 technical replicate for biological

replicate 2, and all 3 technical replicates for biological replicate 3) failed to amplify any

product. Due to resource constraints, we reduced replicates for batch 2 ($n=1$ biological/$n=2$

technical; Figure S2). To reduce model complexity, we omitted some batch 1 isolates

($n=16/35$) which carried more than one copy of bla_{TEM-1} in the genome, leaving a total of

$n=67$ isolates. ΔCt values were calculated by subtracting mean 16S Ct from mean TEM-1 Ct.

Assembly annotations

We annotated the chromosomes using Prokka (v . 1.14.6) with default parameters except `--`

`centre X --compliant` (see `annotate.sh`)[39]. Abricate (v. 1.0.1) was used with

default parameters and the PlasmidFinder database (v. 2023-Nov-4) to annotate for plasmid

replicons[40,41]. Plasmid mobilities were predicted using MOB-suite's MOB-typer (v. 3.1.4)

with default parameters[41]. Briefly, a plasmid was labelled as putatively conjugative if it had

both a relaxase and mating pair formation (MPF) complex, mobilisable if it had either a

relaxase or an origin of transfer (oriT) but no MPF, and non-mobilisable if it had no relaxase

and oriT. Lastly, we assigned sequence types (STs) and phylogroups to our *E. coli*

chromosomes using mlst (v. 2.23.0) with default parameters and EzClermont (v. 0.7.0)

default parameters, respectively[42,43]. We used blastn (v. 2.15.0+) with a custom database of

known bla_{TEM-1} promoters[13,44,45]. Due to the high similarity between the P3, Pa/b, P4, and P5

reference sequences, we chose the top hit in each position.

SNV analysis

We first determined the sets of sequences we wanted to align: (i) bla_{TEM-1} (*n*=451; some

genomes carried multiple copies), (ii) bla_{TEM-1} promoters (*n*=409), (iii) *ampC* (*n*=377), and

(iv) *ampC* promoters (*n*=377). For bla_{TEM-1} and *ampC*, we extracted the relevant sequences

using the coordinate and strand information from the NCBIAMRFinder output (see

extractGene.py). For the bla_{TEM-1} promoters, we used coordinate and strand information

from the earlier blastn results. For the *ampC* promoters, we took the sequence 200bp

upstream of the *ampC* gene then manually excised the -42 to +37 region in AliView[46]. Sets of

sequences were aligned using MAFFT (v. 7.520) with default parameters except --auto[47].

Variable sites were examined using snp-sites (v. 2.5.1) with default parameters and in -v

mode[48].

Contig copy number

We used BWA (v. 0.7.17-r1188) to map the quality-controlled short-reads to each contig,

then SAMtools (v. 1.18) for subsequent processing (see `copyNumber.sh`)[29,49]. For each

contig, we calculated the mean depth over its length, then within each assembly, normalised

by the mean depth of the chromosome.

Chromosomal core gene phylogeny

Building the chromosomal phylogeny involved four main steps: annotating the chromosomes,

identifying the core genes, aligning them, and building a phylogeny. Initially, all the

chromosomes carried a copy of *ampC*, meaning it was a core gene and would be included in

the phylogeny. Since we wanted to manually verify EzClermont phylogroup classifications

with the phylogeny and then compare phylogroups to the distribution of *ampC* gene variants,

we excised the *ampC* sequence from all the chromosomes beforehand to avoid confounding

our analysis (see `removeGene.py`). To identify the core genes (those with ≥98% frequency

in the sample), we used Panaroo (v. 1.4.2) with default parameters except `--clean-mode`

`sensitive --aligner mafft -a core --core_threshold 0.98`[50]. Panaroo

also aligned our core genes using MAFFT (v. 7.520; see `runPanaroo.sh`)[47]. Lastly, we

built the core gene maximum-likelihood phylogeny using IQ-Tree (v. 2.3.0) with default

parameters except `-m GTR + F + I + R4 -keep-ident -B 1000 -mem 10G`

using `-s core_gene_alignment_filtered.aln` from Panaroo (see

`runIQTREE.sh`)[51]. The substitution model used was general time reversible (GTR) using

empirical base frequencies from the alignment (F), allowing for invariant sites (I) and

variable rates of substitution (R4).

Statistical analysis and visualisation

All statistical analysis was performed in R (v. 4.4.0) using RStudio (v.

2024.04.2+764)[52,53].We implemented MCMC generalised linear mixed models using the

MCMCglmm library in R[54]. Model specifications, convergence diagnostics, parameter

estimations, and outputs are reported in Supplementary File 2; see `modelExpression.R`,

`modelMIC.R`, and `modelCombined.R`, to reproduce the bla_{TEM-1} expression, co-

amoxiclav MIC, and causal models, respectively). Homogeneity and completeness are

defined in Rosenberg, A. and Hirschberg, J (2007) and were also implemented in R[55]. A 95%

highest posterior density (HPD) credible interval finds the closest points (a and b) for which

$F(b) - F(a) = 0.95$, where F is the empirical density of the posterior. Figures were plotted

with the ggplot2 library[56].

Computational reproducibility

See `buildResults.R` to reproduce all statistics and figures in the manuscript. All scripts

referenced in the Materials and methods can be found in https://github.com/wtmatlock/tem.

Data availability

Metadata for all n=377 genomes included in the final analysis is given in Supplementary

Table 1. Metadata for all n=451 bla_{TEM-1} annotations identified in these genomes is given in

Supplementary Table 2. qPCR expression data for all replicates is given in Supplementary

Table 3. NCBI accessions for short- and long-read sets and assemblies are given in

Supplementary Table 4.

Results

240   **A curated dataset of *E. coli* isolates with hybrid assemblies and co-amoxiclav MICs**

241 We began with $n=548$ candidate *E. coli* isolates, which following hybrid assembly, were

242 curated into a final dataset of $n=377/548$ (see Materials and methods and Supplementary File

243 1). In total, 77% (291/377) of assemblies were complete (all contigs were circularised), with

244 the remaining 27% (86/377) having at least a circularised chromosome to confidently

245 distinguish between chromosomal and plasmid-associated bla_{TEM-1}. Assemblies contained

246 median=3 (IQR=2-5) plasmid contigs.

248 We identified $n=451$ bla_{TEM-1} genes on 431 contigs (13% [58/431] chromosomal versus 87%

249 [373/431] plasmid). Isolates carried a median=1 copy of bla_{TEM-1} (range=1-6). Carrying more

250 than one copy of bla_{TEM-1} on a single contig was rare: of all bla_{TEM-1}-positive contigs, 97%

251 (400/412) versus 3% (12/412) had no duplications versus at least one. The bla_{TEM-1} genes had

252 synonymous single nucleotide polymorphisms (SNPs) in positions 18, 138, 228, 396, 474,

253 705, and 717, totalling $n=7$ single nucleotide variant (SNV) profiles across the replicons, yet

254 diversity was dominated by bla_{TEM-1b} at 73% (329/451; SNV profile TATTTCG; see

255 Materials and methods)[57]. Where bla_{TEM-1}-positive contigs had at least two copies of bla_{TEM-1},

256 they were almost always the same SNV duplicated (11/12).

258 By examining the "genomic arrangement" of bla_{TEM-1} (namely the replicons it was found on

259 as well as any copies), we found most isolates carried a single non-chromosomal copy

260 (73.5% [277/377]; Figure 1a-b). More generally, whilst the plasmid contigs totalled only

261 3.6% of the total sequence length (bp) across the assemblies

262 (71,126,646bp/1,969,804,202bp), they carried 85.8% of the bla_{TEM-1} genes (387/451). Such

263 bla_{TEM-1}-carrying plasmids were represented across the *E. coli* phylogeny (Figure 1c).

264 Overall, the dataset comprised 5.6% (21/377) phylogroup A, 8.0% (30/377) B1, 48.5%

265 (183/377) B2, 4.8% (18/377) C, 27.3% (103/377) D, 0.5% (2/377) E, 4.2% (16/377) F, and

1.1% (4/377) G. In total, we manually corrected $n=5$ EzClermont phylogroup classifications

using a core-gene phylogeny (see Materials and methods): OXEC-108 (G to D), OXEC-317

(B2 to D), OXEC-333 (U to B1), OXEC-344 (U to B1), and OXEC-406 (U to B1). The

EzClermont publication presented a 98.4% (123/125) true-positive rate on their validation

set, which is in line with our ~98.7% (372/377)[43].

Five known upstream promoters modulate the expression of bla_{TEM-1}: P3, Pa/Pb, P4, P5, and

Pc/d [13,44]. We linked 91% (409/451) bla_{TEM-1} genes to a promoter immediately upstream, of

which a majority, 64% (262/409), were identical to the P3 reference. More generally,

excluding $n=2$ different Pc/Pd-like promoters which have large deletions, we identified SNPs

in positions 32, 43, 65, 141, 162, and 175, totalling $n=8$ SNV profiles (by Sutcliffe

numbering[58]). Notably, 15% (39/262) of promoters had the Pa/Pb-associated C32T mutation

which produces two overlapping promoter sequences.

Isolates were associated with a diverse range of co-amoxiclav MICs for the 377 isolates

(μg/mL; $\leq$2/2 [4 (1.1%)], 4/2 [24 (6.4%)], 8/2 [144 (38.2%)], 16/2 [86 (22.8%)], 32/2 [44

(11.7%)], and >32/2 [75 (19.9%)]; Figure 1d; see Materials and methods).

Figure 1. A genotypically and phenotypically heterogenous population of *bla*_{TEM-1}-carrying *E. coli*. (a) The genomic arrangement of *bla*_{TEM-1} in the genomes ordered in descending **(b)** frequency. **(c)** Phylogroup and **(d)** co-amoxiclav MIC distribution for each genomic arrangement.

285 **_bla__{TEM-1} associated with conjugative plasmids**

286   Whilst chromosomal copies of *bla*_{TEM-1} can remain with a lineage over time, plasmidic copies

287   might come by conjugation and leave by loss. This could give the host cell access to a

288   transient boost in resistance without impeding long-term fitness (see Materials and methods).

289   Confining the analysis to circularised plasmids (1,036/1,512) and using a 2-sample test for

290   equality of proportions with continuity correction, the *bla*_{TEM-1}-positive plasmids (333/1,036)

291   were significantly more likely to be putatively conjugative (85.9% [286/333]) compared to

292   the *bla*_{TEM-1}-negative plasmids (28.4% [200/703]; prop=200/703; $\chi^2 = 297.02$, df = 1, *p*-value

< 2.2e-16). Moreover, for the most common genomic arrangement of bla_{TEM-1} (i.e.a single

plasmid [277/377]), amongst circularised plasmids (252/277), 90% were putatively

conjugative (227/252).

For a genome carrying bla_{TEM-1} on the chromosome, the gene's copy number and total

number of genes in the genome are equivalent. For a genome carrying bla_{TEM-1} on a plasmid,

this might not be the case. This is because plasmids can exist as multiple copies. The

calculated copy number of all plasmidic contigs (n=1,512) was median=3.13 (range=0.04-

57.00). Of these, n=19/1,512 contigs (with 6/19 circularised) had calculated copy numbers

less than one (see Materials and methods). This was potentially due to uneven short-read

coverage. However, none carried bla_{TEM-1} and so were not used in the later modelling. Taking

the circularised plasmids with copy number at least one (1,030/1,512), longer plasmids

(>10kbp; 668/1,030) were generally low copy number (median=2.36), whilst shorter

plasmids (≤10kbp; 362/1,030) were generally high copy number (median=11.01, see Figure

S3), consistent with previous studies[59].

***E. coli* phylogeny shapes *bla*_{TEM-1} expression**

Within different *E. coli* lineages, bla_{TEM-1} and its promoter are potentially subject to different

regulatory systems and epigenetic interactions, which may in turn affect bla_{TEM-1} expression.

To test this, we first selected a random subsample of n=67/377 isolates with a single copy of

bla_{TEM-1} in the genome, either on a chromosome (15/67) or a plasmid (52/67). Moreover, we

only selected isolates with zero, one, or two mutations in the bla_{TEM-1} promoter sequence:

C32T, which produces two overlapping promoters and is known to increase expression[13,] and

G175A (according to Sutcliffe numbering based on the PBR322 plasmid[58]; see Table 1).

Isolates were distributed across the entire *E. coli* phylogeny (phylogroup A [6/67], B1 [4/67],

B2 [40/67], C [4/67], D [8/67], E [1/67], and F ([4/67]). We then performed qPCR to evaluate

for $bla_{\text{TEM-1}}$ expression (see Materials and methods). Every isolate had at least two replicates

(2 [48/67], 4 [1/67], or 9 [18/67]), giving a total of $n=262$ $bla_{\text{TEM-1}}$ ΔCt observations (TEM-1

Ct – 16S Ct; see Materials and methods) for modelling.

Promoter SNV / Replicon	Chromosome	Plasmid	Total
CG (wildtype)	1	22	**23**
G175A	2	21	**23**
C32T	3	5	**7**
C32T, G175A	9	4	**13**
Total	**15**	**52**	**67**

Table 1. Replicon distribution of $n=67$ $bla_{\text{TEM-1}}$ promoter variants. Single nucleotide variants (SNVs) are for the 32nd and 175th positions by Sutcliffe numbering[58].

To test for the effects of *E. coli* lineage, we built a maximum likelihood core gene phylogeny

for all $n=377$ chromosomes (see Materials and methods). In total, we identified 17,836 gene

clusters, of which 18.7% (3,342/17,836) were core genes (those found in $\geq$98% of

chromosomes). The phylogeny (midpoint rooted and restricted to the $n=67/377$ isolates in

the expression analysis) is given in Figure 2a. Using $b=1000$ ultrafast bootstraps, all

phylogroup node supports were 100%, and more generally, 76.7% (287/374) of internal node

supports were 100%, and 87.4% (327/374) were at least 95% (see Materials and methods).

Moreover, the Robinson-Foulds distance between the ML tree and consensus tree was 4,

indicating nearly identical topology.

Briefly, the expression linear mixed model employed Markov Chain Monte Carlo (MCMC)

to estimate parameters. The response variable $bla_{\text{TEM-1}}$ ΔCt (normalised and 95th percentile

truncated) was related to the fixed effects (i) $bla_{\text{TEM-1}}$ cell copy number (normalised), (ii)

presence of the C32T promoter mutation, (iii) presence of the G175A mutation, and (iv) their

interaction. Random effects were incorporated to account for qPCR replicates and

Figure 2. Intrinsic expression of bla_{TEM-1} shapes co-amoxiclav MIC across the *E. coli* phylogeny. (a) A midpoint-rooted core gene phylogeny of *E. coli* chromosomes from the 67 isolates with expression quantified. Tips are coloured by phylogroup. ST12 (higher than average effect) and ST372 (lower than average effect) are highlighted by green and orange outlines, respectively. Tip shape distinguishes location of bla_{TEM-1} on chromosomes (circles) and plasmids (squares). **(b)** Posterior means (coloured circles) and 95% HPD intervals (horizontal lines) for phylogenetic effect on negative bla_{TEM-1} ΔCt for each tip (multiplied by -1 for ease of comparison). Red indicates above average expression and blue indicates below average expression. **(c)** Posterior means (coloured circles) and 95% HPD intervals (horizontal lines) for phylogenetic effect on co-amoxiclav MIC for each tip. Pink indicates above average MIC and green indicates below average MIC.

In decreasing order of effect size, C32T, G175A, and a one unit increase in contig copy

number all increased expression (decreased ΔCt; Table 2). There was no additional effect of

G175A if C32T was also present (-1.69 < -1.71). The posterior for contribution of variance

from phylogeny demonstrated a long right tail (mean=0.07; 95% highest posterior density,

HPD=[0.00, 0.21]; see Materials and methods), suggestive of high heterogeneity. For qPCR

replicates, the contribution of variance exhibited minimal skew (mean=0.15; 95%

HPD=[0.08, 0.23]). To investigate this further, we computed the posterior mean and 95%

HPD credible interval for each tip in the phylogeny (Figure 2b). Compared to the average

across the *E. coli* phylogeny, some phylogroups (B1, C) and STs (12) were associated with

increased bla_{TEM-1} expression, whilst some phylogroups (E, F) and STs (372) were associated

with decreased bla_{TEM-1} expression.

Variable	Beta coefficient posterior mean	95% HPD (l, u)	p_{MCMC}
Intercept	0.28	0.03, 0.59	0.0245
C32T*	-1.71	-2.08, -1.34	<1e-05
G175A*	-0.31	-0.57, -0.04	0.0202
C32T*:G175A* interaction	0.33	-0.14, 0.81	0.1779
bla_{TEM-1} cell copy number	-0.12	-0.24, 0.01	0.0679

Table 2. Parameter estimates for bla_{TEM-1} ΔCt genotype-phenotype model. All values are taken from chain 1. For Replicates and Phylogeny, estimates represent the average contribution to variance across the sample. Effect of both C32T* and G175A* is -1.71-0.31+0.33=-1.69.

* Numbering by Sutcliffe[58]

***ampC* gene variation is highly concordant with *E. coli* lineage**

In *E. coli*, the chromosomally intrinsic *ampC* encodes a class C beta-lactamase with

expression typically induced from an external stimulus, namely the presence of a penicillin

such as amoxicillin, or a beta-lactamase inhibitor such as clavulanic acid. At the time of

writing, the beta-lactamase database (BLDB) contains *n*=2,281 non-synonymous variants of

the gene.

To quantify how well *ampC* variants agree with phylogroup and ST, we calculated the

homogeneity (h) and completeness (c; both range from 0 to 1; see Materials and methods).

Briefly, $h = 1$ means that a phylogroup or ST contains a single *ampC* variant. Conversely, $c =$

1 means that all instances of an *ampC* variant fall within the same phylogroup or ST. For

phylogroups, we found $h = 0.489$ and $c = 0.964$, and for STs (excluding 38/377 which were

unassigned), $h = 0.938$ and $c = 0.877$. Overall, this suggests that phylogroups tend to contain

distinct *ampC* variants, which are generally ST-specific, and overall, that *E. coli* phylogeny is

a suitable proxy for *ampC* variation.

Whilst many *E. coli ampC* variants present a narrow spectrum of hydrolytic activity, some

can potentially hydrolyse third-generation cephalosporins following mutations in the

promoter sequence. To explore promoter variation, we aligned all $n=377$ *ampC* promoter

sequences. Mutations outside positions -42 to +37 (according to Jaurin numbering[14]) were

disregarded based on existing characterisations[60,61]. In total, $n=12$ *ampC* promoter SNVs

were identified, with variation dominated by the *E. coli* K12 wildtype at 47% (177/377).

Table 3 documents all $n=11$ mutations identified. A given *ampC* variant associated almost

uniquely with an *ampC* promoter variant, yet *ampC* promoter variants were associated with

multiple *ampC* variants ($h = 0.483$ and $c = 0.941$).

Region	Position	*E. coli* K12 (*n*)	Mutation (*n*)
Spacer	-28	G (243)	A (134)
	-18	G (329)	A (48)
Between -10 box and attenuator	-1	C (329)	T (48)
	+11	T (371)	- (6)
Attenuator	+17	C (316)	T (61)
	+22	C (367)	T (10)
	+26	T (367)	G (10)
	+27	A (367)	T (10)
	+30	G (376)	A (1)
	+32	G (364)	A (13)
	+37	G (376)	T (1)

Table 3. Variation in *n*=377 *ampC* promoters. Positions are according to Jaurin numbering[14].

Efflux pump AcrF is not encoded by all *E. coli*

Efflux pumps can confer resistance to antibiotics by transporting molecules outside the cell[15].

In particular, members of the resistance-nodulation-cell division superfamily (RND),

including AcrF (encoded by *acrF*), can export penams. We annotated our *n*=377 genomes for

efflux pump genes, totalling *n*=1,067 hits (at least 100% coverage and at least 95% identity;

see Materials and methods), invariably on the chromosome. Chromosomes had median=3

annotations (range=1-4). AcrF was encoded by 65% (244/377) of isolates, always as a single

copy. Moreover, it was widely distributed across the phylogeny: 8.2% (20/244) phylogroup

A, 11.1% (27/244) B1, 29.9% (73/244) B2, 7.0% (17/244) C, 38.9% (95/244) D, 0.8%

(2/244) E, 2.5% (6/244) F, and 1.6% (4/244) G. All other efflux pump annotations are given

in Supplementary Table 1.

***E. coli* phylogeny drives co-amoxiclav resistance through expression**

We next investigated whether the *E. coli* lineages with intrinsically higher bla_{TEM-1}

expression also had intrinsically higher co-amoxiclav MICs. This would be consistent with

lineage differences in regulatory regions and epigenetic interactions driving increased

resistance.

We employed an MCMC to estimate parameters in an ordinal mixed model. The response

Variable		Beta coefficient posterior mean	95% HPD (l, u)	p_{MCMC}
Intercept		3.89	2.89, 4.89	<1e-05
bla_{TEM-1} cell copy number		2.01	1.34, 2.68	<1e-05
bla_{TEM-1} genome copy number (>1 vs. 1)		0.99	0.04, 1.93	0.0399
bla_{TEM-1} promoter SNV vs. wildtype	G175A*	0.16	-0.37, 0.67	0.5435
	C32T*	6.06	4.14, 8.06	<1e-05

variable isolate co-amoxiclav MIC (μg/mL; levels ≤2/2, 4/22, 8/22, 16/2, 32/2, >32/2) was

predicted by the fixed effects (i) bla_{TEM-1} cell copy number (normalised and 95[th] percentile

truncated), (ii) bla_{TEM-1} genome copy number (>1 vs. 1), (iii) non-wildtype bla_{TEM-1} promoter

SNVs, (iv) non-wildtype $ampC$ promoter SNVs, and (v) presence of AcrF. For the model, we

only used isolates for which every bla_{TEM-1} gene was linked to a promoter, and all the

promoters were the same variant. We then filtered out isolates with bla_{TEM-1} promoter and

$ampC$ promoter variants that appeared less than 10 times. This left n=292/377 isolates. Full

model specification, convergence diagnostics, and outputs are given in Supplementary File 2.

	C32T* and G175A*	5.88	3.91, 7.94	<1e-05
ampC promoter SNV vs. wildtype	G-28A[†] and C17T[†]	0.44	-0.62, 1.51	0.4096
	G-28A[†]	0.87	-0.93, 2.60	0.3152
	G-18A[†] and C-1T[†]	0.83	-1.45, 3.07	0.4444
AcrF		0.02	-0.51, 0.54	0.9475

Table 4. Parameter estimates for co-amoxiclav MIC genotype-phenotype model. All values are taken from chain 1.

* Numbering by Sutcliffe[58]
[†] Numbering by Jaurin[14]

In decreasing order of effect size, the presence of C32T and G175A in the bla_{TEM-1} promoter,

the presence of just C32T, bla_{TEM-1} cell copy number, and bla_{TEM-1} genome copy number all

increased co-amoxiclav MIC (Table 4); the remaining effects were compatible with chance.

As with the expression model, the posterior distribution for contribution of variance from

phylogeny demonstrated a long right tail (mean=2.85; 95% HPD=[0.75, 5.25]). The

phylogeny for the n=292 isolates with co-amoxiclav MIC tip effects is given in Figure S4.

Lastly, we developed a combined model to test whether the phylogenetic influence on bla_{TEM-}

$_1$ expression casually varies co-amoxiclav MIC (see Supplementary File 2). Here, we only

used the predictors identified as significant from previous expression and MIC models

(bla_{TEM-1} cell and genome copy numbers, and bla_{TEM-1} promoter SNV). Briefly, the model

estimates a parameter that scales the phylogenetic and non-phylogenetic random effects from

expression to MIC. Under causality, the scaling parameter should be constant across all

random effect terms. We found the scaling parameter had a posterior mean=-1.13 (95%

HPD=[-0.72, -1.49]; p_{MCMC}=0.002; values taken from chain 1). This supports a direct and

substantial influence of expression on MIC, mediated by phylogenetic relationships.

Discussion

In our dataset of clinical *E. coli*, bla_{TEM-1} was overwhelmingly carried by conjugative

plasmids. These means it can spread between bacterial hosts and different genetic

backgrounds. We demonstrated that different bacterial hosts intrinsically vary in their ability

to express bla_{TEM-1} when accounting for variation in promoters (C32T and G175A mutations)

and contig copy number. Moreover, our findings suggest that some clinically successful

lineages (e.g. ST12) are better at expressing bla_{TEM-1} than less clinically successful lineages

(e.g. phylogroup F). With a second model, we also found that different *E. coli* lineages vary

intrinsically in co-amoxiclav MIC (accounting for bla_{TEM-1} genome and cell copies, bla_{TEM-1}

and *ampC* promoter variants, and AcrF presence). Again, we observed that some clinically

successful lineages (e.g. ST12) had higher resistance than less clinically successful lineages

(e.g. phylogroup F). A third model demonstrated that these two traits were casually linked: *E.*

coli phylogeny drives co-amoxiclav resistance through variable expression of bla_{TEM-1}. We

believe this finding should generalise to other AMR genes, and underscores the necessity of

fully resolving bacterial genomes to incorporate accurate genetic, genomic, and phylogenetic

information in resistance prediction models. Future work could include evaluations of single

amino acid substitutions in TEM-1 (that hydrolyse third-generation cephalosporins and

carbapenemases[62]) which are typically carried in more complex genetic backgrounds.

This study has limitations. Firstly, it is possible that, due to fragmented plasmid assemblies,

some isolates identified as having multiple copies of bla_{TEM-1} on multiple plasmids instead

had multiple copies on the same plasmid. Nonetheless, our expression analysis only

considered isolates with a single copy of bla_{TEM-1} in the genome, mitigating this concern.

Secondly, we only examined expression for a subsample of our isolates due to resource

limitations. Thirdly, whilst there is not an agreed upon standard reference gene for

quantifying beta-lactamase expression[63–65], previous work has shown 16S to be stable[64].

Crucially, our delta Ct values were consistent within isolates. Fourthly, whilst we found no

signal for the AcrF pump influencing co-amoxiclav MIC, recent work on ST11 suggests that

not all *E. coli* possess a functional copy of the gene[66]. However, it was out of scope of this

work to characterise all AcrF gene variants. We also only observed two potentially relevant

porin mutations (a premature stop codon in OmpC on OXEC-40's chromosome and in OmpF

on OXEC-423's chromosome; see Supplementary Table 1), limiting our ability to investigate

their effects on phenotype. Penultimately, automated susceptibility testing methods, like the

BD Phoenix™ used here, may not agree completely with reference methods; yet previous

work has shown strong agreement with the EUCAST agar dilution method[67]. Lastly, plasmid

copy number is not static. Moreover, in the presence of antibiotics, it has been demonstrated

that resistance gene-carrying plasmids can increase their copy number to increase the chance

of survival[68]. Our point estimates of plasmid copy number were derived from genome

assemblies sequenced in the absence of antibiotics, which likely represent a lower bound.

Nonetheless, we found strong signal to suggest the import of plasmid copy number on

resistance, even if under our sensitivity testing, plasmid copy number potentially increased

within isolates.

Future studies should examine specific regulatory pathways, epistatic interactions, and

epigenetic mechanisms such as DNA methylation, that link phylogenetic background with

bla_{TEM-1} expression and resistance. Additionally, examining other resistance genes and their

expression patterns in a similar phylogenetic framework could provide a broader

understanding and prediction of resistance mechanisms across different bacterial species and

antibiotics. Expression of a resistance gene like bla_{TEM-1} might carry a fitness penalty to the

host cell. In the absence of antibiotics, this can select against cells with resistance gene-

carrying plasmids. We speculate that in bacterial populations, conjugative plasmids could

escape purging by moving from bacterial lineages with high intrinsic expression, which bear

the fitness cost, to low intrinsic expression, where the fitness cost is reduced. This process

could help maintain the presence of resistance genes within the population even in the

absence of antibiotic pressure.

Funding information

This work was funded by the National Institute for Health Research (NIHR) Health

Protection Research Unit in Healthcare Associated Infections and Antimicrobial Resistance

(NIHR200915), a partnership between the UK Health Security Agency (UKHSA) and the

University of Oxford. It was also supported by the NIHR Oxford Biomedical Research

Centre (BRC). The computational aspects of this research were funded from the NIHR

Oxford BRC with additional support from the Wellcome Trust Core Award Grant Number

203141/Z/16/Z. The views expressed are those of the author(s) and not necessarily those of

the NIHR, UKHSA or the Department of Health and Social Care.

Acknowledgements

496   The authors thank Jarrod Hadfield for providing guidance on implementing the MCMCglmm

497   library.

Conflicts of interest

500   The authors declare that there are no conflicts of interest.

Bibliography

1.   Datta, N. & Kontomichalou, P. Penicillinase synthesis controlled by infectious R

factors in enterobacteriaceae. *Nature* **208**, (1965).

2.   Partridge, S. R. & Hall, R. M. Evolution of transposons containing blaTEM genes.

Antimicrobial Agents and Chemotherapy vol. 49 Preprint at

https://doi.org/10.1128/AAC.49.3.1267-1268.2005 (2005).

3.   Shaw, L. P. & Neher, R. A. Visualizing and quantifying structural diversity around

mobile resistance genes. *Microb Genom* **9**, (2023).

4.   Bailey, J. K., Pinyon, J. L., Anantham, S. & Hall, R. M. Distribution of the blaTEM

gene and blaTEM-containing transposons in commensal Escherichia coli. *Journal of*

Antimicrobial Chemotherapy **66**, (2011).

5.   Alcock, B. P. *et al.* CARD 2023: expanded curation, support for machine learning, and

resistome prediction at the Comprehensive Antibiotic Resistance Database. *Nucleic*

Acids Res **51**, (2023).

6.   Sayers, E. W. *et al.* Database resources of the national center for biotechnology

information. *Nucleic Acids Res* **50**, (2022).

7.   Murray, C. J. *et al.* Global burden of bacterial antimicrobial resistance in 2019: a

systematic analysis. *The Lancet* **399**, (2022).

8.   Sanderson, T. A web tool for exploring the usage of medicines in hospitals in England.

Wellcome Open Res **9**, 147 (2024).

9.   EUCAST: Clinical breakpoints and dosing of antibiotics.

https://www.eucast.org/clinical_breakpoints.

10.   Yoon, C. H. *et al.* Mortality risks associated with empirical antibiotic activity in

Escherichia coli bacteraemia: an analysis of electronic health records. *Journal of*

Antimicrobial Chemotherapy **77**, (2022).

11.   Martinez, J. L. *et al.* Resistance to beta-lactam/clavulanate. *Lancet* vol. 2 Preprint at

https://doi.org/10.1016/s0140-6736(87)91180-9 (1987).

12.   Reguera, J. A., Baquero, F., Perez-Diaz, J. C. & Martinez, J. L. Synergistic effect of

dosage and bacterial inoculum in TEM-1 mediated antibiotic resistance. *European*

Journal of Clinical Microbiology & Infectious Diseases **7**, (1988).

13.   Lartigue, M. F., Leflon-Guibout, V., Poirel, L., Nordmann, P. & Nicolas-Chanoine, M.

H. Promoters P3, Pa/Pb, P4, and P5 upstream from blaTEM genes and their

relationship to β-lactam resistance. *Antimicrob Agents Chemother* **46**, (2002).

14.   Jaurin, B., Grundström, T. & Normark, S. Sequence elements determining ampC

promoter strength in E. coli. *EMBO J* **1**, (1982).

15.   Siasat, P. A. & Blair, J. M. A. Microbial Primer: Multidrug efflux pumps.

Microbiology (United Kingdom) **169**, (2023).

16.   Vital, M. *et al.* Gene expression analysis of E. coli strains provides insights into the

role of gene regulation in diversification. *ISME Journal* **9**, (2015).

17.   McNally, A. *et al.* Combined Analysis of Variation in Core, Accessory and Regulatory

Genome Regions Provides a Super-Resolution View into the Evolution of Bacterial

Populations. *PLoS Genet* **12**, (2016).

18.   Card, K. J., Thomas, M. D., Graves, J. L., Barrick, J. E. & Lenski, R. E. Genomic

evolution of antibiotic resistance is contingent on genetic background following a

long-term experiment with Escherichia coli. *Proc Natl Acad Sci U S A* **118**, (2021).

19.   Wong, A. Epistasis and the evolution of antimicrobial resistance. *Frontiers in*

Microbiology vol. 8 Preprint at https://doi.org/10.3389/fmicb.2017.00246 (2017).

20.   Wang, X., Yu, D. & Chen, L. Antimicrobial resistance and mechanisms of epigenetic

regulation. *Frontiers in Cellular and Infection Microbiology* vol. 13 Preprint at

https://doi.org/10.3389/fcimb.2023.1199646 (2023).

21.   Dunn, S., Carrilero, L., Brockhurst, M. & McNally, A. Limited and Strain-Specific

553   Transcriptional and Growth Responses to Acquisition of a Multidrug Resistance

554   Plasmid in Genetically Diverse Escherichia coli Lineages. *mSystems* **6**, (2021).

555   22.   Valle, A. A. Del *et al.* Antimicrobial resistance level and conjugation permissiveness

556   shape plasmid distribution in clinical enterobacteria. *Proc Natl Acad Sci U S A* **120**,

557   (2023).

558   23.   Feldgarden, M. *et al.* AMRFinderPlus and the Reference Gene Catalog facilitate

559   examination of the genomic links among antimicrobial resistance, stress response, and

560   virulence. *Sci Rep* **11**, (2021).

561   24.   Lipworth, S. *et al.* Ten-year longitudinal molecular epidemiology study of Escherichia

562   coli and Klebsiella species bloodstream infections in Oxfordshire, UK. *Genome Med*

563   **13**, (2021).

564   25.   Chen, S., Zhou, Y., Chen, Y. & Gu, J. Fastp: An ultra-fast all-in-one FASTQ

565   preprocessor. in *Bioinformatics* vol. 34 (2018).

566   26.   Wick, R. R. Filtlong. *GitHub* (2019).

567   27.   Hall, M. Rasusa: Randomly subsample sequencing reads to a specified coverage. *J*

568   *Open Source Softw* **7**, (2022).

569   28.   Oxford Nanopore Technologies. Medaka: Sequence correction provided by ONT

570   Research. *GitHub* (2018).

571   29.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler

572   transform. *Bioinformatics* **25**, (2009).

573   30.   Wick, R. R. & Holt, K. E. Polypolish: Short-read polishing of long-read bacterial

574   genome assemblies. *PLoS Comput Biol* **18**, (2022).

575   31.   Wick, R. R., Judd, L. M., Gorrie, C. L. & Holt, K. E. Unicycler: Resolving bacterial

576   genome assemblies from short and long sequencing reads. *PLoS Comput Biol* **13**,

577   (2017).

578   32.   Bankevich, A. *et al.* SPAdes: A new genome assembly algorithm and its applications

579   to single-cell sequencing. *Journal of Computational Biology* **19**, (2012).

580   33.   Li, H. Minimap and miniasm: Fast mapping and de novo assembly for noisy long

581   sequences. *Bioinformatics* **32**, (2016).

582   34.   Vaser, R., Sović, I., Nagarajan, N. & Šikić, M. Fast and accurate de novo genome

583   assembly from long uncorrected reads. *Genome Res* **27**, (2017).

584   35.   Ondov, B. D. *et al.* Mash Screen: High-throughput sequence containment estimation

585   for genome discovery. *Genome Biol* **20**, (2019).

586   36.   Galata, V., Fehlmann, T., Backes, C. & Keller, A. PLSDB: A resource of complete

587   bacterial plasmids. *Nucleic Acids Res* **47**, (2019).

588   37.   Camacho, C. *et al.* BLAST+: Architecture and applications. *BMC Bioinformatics* **10**,

589   (2009).

590   38.   Rodger, G. *et al.* Comparison of direct cDNA and PCR-cDNA Nanopore sequencing

591   of Escherichia coli isolates. *bioRxiv* 2024.01.23.576853 (2024)

592   doi:10.1101/2024.01.23.576853.

593   39.   Seemann, T. Prokka: Rapid prokaryotic genome annotation. *Bioinformatics* **30**, (2014).

594   40.   Carattoli, A. *et al.* In Silico detection and typing of plasmids using plasmidfinder and

595   plasmid multilocus sequence typing. *Antimicrob Agents Chemother* **58**, (2014).

596   41.   Robertson, J. & Nash, J. H. E. MOB-suite: software tools for clustering, reconstruction

597   and typing of plasmids from draft assemblies. *Microb Genom* **4**, (2018).

42.   Seemann T. mlst. Preprint at (2017).

43.   Waters, N. R., Abram, F., Brennan, F., Holmes, A. & Pritchard, L. Easy phylotyping

of Escherichia coli via the EzClermont web app and command-line tool. *Access*

Microbiol **2**, (2020).

44.   Robin, F. *et al.* Evolution of TEM-type enzymes: Biochemical and genetic

characterization of two new complex mutant TEM enzymes, TEM-151 and TEM-152,

from a single patient. *Antimicrob Agents Chemother* **51**, (2007).

45.   Stoesser, N. *et al.* Predicting antimicrobial susceptibilities for Escherichia coli and

Klebsiella pneumoniae isolates using whole genomic sequence data. *Journal of*

Antimicrobial Chemotherapy **68**, (2013).

46.   Larsson, A. AliView: A fast and lightweight alignment viewer and editor for large

datasets. *Bioinformatics* **30**, (2014).

47.   Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:

Improvements in performance and usability. *Mol Biol Evol* **30**, (2013).

48.   Page, A. J. *et al.* SNP-sites: rapid efficient extraction of SNPs from multi-FASTA

alignments. *Microb Genom* **2**, (2016).

49.   Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, (2021).

50.   Tonkin-Hill, G. *et al.* Producing polished prokaryotic pangenomes with the Panaroo

pipeline. *Genome Biol* **21**, (2020).

51.   Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic

Inference in the Genomic Era. *Mol Biol Evol* **37**, (2020).

52.   R Core Team. R: A language and environment for statistical computing. *R Foundation*

for Statistical Computing Preprint at (2019).

53.   RStudio Team. RStudio: Integrated Development for R. *RStudio, Inc., Boston, MA*

Preprint at (2021).

54.   Hadfield, J. D. MCMC methods for multi-response generalized linear mixed models:

The MCMCglmm R package. *J Stat Softw* **33**, (2010).

55.   Rosenberg, A. & Hirschberg, J. V-Measure: A conditional entropy-based external

cluster evaluation measure. in *EMNLP-CoNLL 2007 - Proceedings of the 2007 Joint*

Conference on Empirical Methods in Natural Language Processing and

Computational Natural Language Learning (2007).

56.   Gómez-Rubio, V. ggplot2 - Elegant Graphics for Data Analysis (2nd Edition) . *J Stat*

Softw **77**, (2017).

57.   Goussard, S. & Courvalin, P. Updated sequence information for TEM β-lactamase

genes. *Antimicrob Agents Chemother* **43**, (1999).

58.   Sutcliffe, J. G. Nucleotide sequence of the ampicillin resistance gene of Escherichia

coli plasmid pBR322. *Proc Natl Acad Sci U S A* **75**, (1978).

59.   Shaw, L. P. *et al.* Niche and local geography shape the pangenome of wastewater-and

livestock-associated Enterobacteriaceae. *Sci Adv* **7**, (2021).

60.   Tracz, D. M. *et al.* ampC gene expression in promoter mutants of cefoxitin-resistant

Escherichia coli clinical isolates. *FEMS Microbiol Lett* **270**, (2007).

61.   Caroff, N., Espaze, E., Bérard, I., Richet, H. & Reynaud, A. Mutations in the ampC

promoter of Escherichia coli isolates resistant to oxyiminocephalosporins without

extended spectrum β-lactamase production. *FEMS Microbiol Lett* **173**, (1999).

62.   Naas, T. *et al.* Beta-lactamase database (BLDB)–structure and function. *J Enzyme*

Inhib Med Chem **32**, (2017).

63.   Rodríguez-Villodres, Á. *et al.* Survival of infection with TEM β-lactamase-producing

Escherichia coli with Pan-β-lactam resistance. *Journal of Infection* **89**, 106268 (2024).

64.   Singh, T. *et al.* Transcriptome analysis of beta-lactamase genes in diarrheagenic

Escherichia coli. *Sci Rep* **9**, 3626 (2019).

65.   Kjeldsen, T. S. B. *et al.* CTX-M-1 β-lactamase expression in Escherichia coli is

649   dependent on cefotaxime concentration, growth phase and gene location. *Journal of*

650   *Antimicrobial Chemotherapy* **70**, 62–70 (2015).

651   66.   Pugh, H. L., Connor, C., Siasat, P., McNally, A. & Blair, J. M. A. E. coli ST11

652   (O157:H7) does not encode a functional AcrF efflux pump. *Microbiology (United*

653   *Kingdom)* **169**, (2023).

654   67.   Davies, T. J. *et al.* Reconciling the potentially irreconcilable? Genotypic and

655   Phenotypic Amoxicillin-Clavulanate Resistance in Escherichia coli. *Antimicrob Agents*

656   *Chemother* **64**, (2020).

657   68.   Hernandez-Beltran, J. C. R. *et al.* Plasmid-mediated phenotypic noise leads to transient

658   antibiotic resistance in bacteria. *Nat Commun* **15**, (2024).

a

Phylogroup
- A
- B1
- B2
- C
- D
- E
- F
- G

Posterior mean
2
1
0
−1

0.01

b

Posterior 95% HDI
−2.5   0.0   2.5   5.0