Microglia are Required for Developmental Specification of AgRP Innervation in the Hypothalamus of Offspring Exposed to Maternal High-Fat Diet During Lactation

Haley N. Mendoza-Romero, Jessica E. Biddinger, Michelle N. Bedenbaugh and Richard B. Simerly[1]

Dept of Molecular Physiology & Biophysics, Vanderbilt University, Nashville, TN, 37232, USA

[1]Corresponding author: Richard B. Simerly
Email: richard.simerly@vanderbilt.edu

Competing Interest Statement: No competing interests

Classifications: Major category: Developmental Biology; Minor category: Neuroscience

Keywords:
Microglia
Neural Development
Hypothalamus
Agouti-related peptide
Paraventricular hypothalamic nucleus

Impact Statement

Microglia appear to play an essential role in specifying patterns of hypothalamic innervation during development in response to maternal HFD exposure, which may contribute to developmental programming of metabolic phenotype.

Abstract

Agouti-related peptide (AgRP) neurons in the arcuate nucleus of the hypothalamus respond to multiple metabolic signals and distribute neuroendocrine information to other brain regions such as the paraventricular hypothalamic nucleus (PVH), which plays a central role in metabolic homeostasis. Neural projections from AgRP neurons to the PVH form during the postnatal lactational period in mice and these projections are reduced in offspring of dams that consumed a high-fat diet (HFD) during lactation (MHFD-L). Here we used immunohistochemistry to visualize microglial morphology in MHFD-L offspring and identified changes that were regionally localized to the PVH and appeared temporally restricted to the period when AgRP neurons innervate this region. In addition, axon labeling experiments revealed that microglia engulf AgRP terminals in the PVH, and that the density of AgRP innervation to the PVH in MHFD-L offspring may be dependent on microglia, because microglial depletion blocked the decrease in PVH AgRP innervation observed in MHFD-L offspring, as well as prevented the increased body weight exhibited at weaning. Together, these findings suggest that microglia are activated by exposure to MHFD-L and interact directly with AgRP axons during postnatal development to permanently alter innervation of the PVH, with implications for developmental programming of metabolic phenotype.

1 Introduction

Maternal nutritional status has a profound effect on the metabolic phenotype of offspring. Children born to obese mothers experience higher rates of obesity later in life, with accompanying comorbidities that negatively impact health and longevity (Stettler et al., 2005; Prospective Studies Collaboration 2009; Tamashiro & Moran 2010; Andersen et al., 2012). Although this developmental programming of metabolic phenotype has been reproduced in several animal models (Samuelsson et al., 2008; Masuyama et al., 2014; García-Cáceres et al., 2019; Skowronski et al., 2023), the underlying mechanisms remain poorly defined. In mouse models, maternal obesity during lactation, a time when offspring are dependent on milk from their mothers for nutrition, appears to be particularly impactful. These changes to metabolic phenotype are thought to be mediated by changes in the milk (Gorski et al., 2006; Vogt et al., 2014; Calvo-Lerma et al., 2022) and occur without subsequent dietary challenge to the offspring themselves, suggesting that they are a consequence of developmental programming (Bolton et al., 2022; Skowronski et al., 2023). Because neural circuits known to control body weight develop during the lactational period, they are vulnerable to a variety of environmental signals that may affect their organization and function (Horvath et al., 2010; Bouret et al., 2015; Elson & Simerly 2015; Zeltser 2018; Skowronski et al., 2023).

AgRP neurons in the arcuate nucleus of the hypothalamus (ARH) function as "hunger neurons" that respond to key metabolic signals such as leptin, ghrelin, glucose and free fatty acids (Krashes et al., 2011; Betley et al., 2015; Chen et al., 2015; Sutton Hickey et al., 2023), and they distribute this information to other regions associated with energy balance regulation (Simerly 2008; Zagmutt, et al., 2018). Thus, the ability of AgRP neurons to influence other components of feeding circuitry is dependent on the formation of their neural connections,

which develop primarily during the first 2 weeks of postnatal life (Bouret et al., 2004a). During
development, AgRP axons extend from the ARH at postnatal day 4 (P4) and reach the
paraventricular hypothalamic nucleus (PVH) between P8-P10. Leptin is required for normal
targeting of AgRP axons to downstream regions, and in leptin-deficient mice both
neuroanatomical and related physiological defects persist into adulthood (Bouret et al., 2004b;
Bouyer & Simerly 2013). Maternal overnutrition affects formation of feeding circuits during
postnatal life with concomitant dysregulation of body weight (Plagemann et al., 1992; Lippert &
Bruning 2022; Skowronski et al., 2023). Limiting HFD exposure of dams to the first 3 weeks of
lactation (MHFD-L) causes suppression of neural projections from AgRP neurons to the PVH
in offspring and is associated with increased body weight later in life (Vogt et al., 2014). In fact,
MHFD-L was more effective than prenatal maternal HFD exposure in causing body weight
changes of adult offspring. MHFD-L did not alter cell number, peptidergic expression, or
cellular activity of AgRP neurons in the ARH, suggesting that maternal nutritional status during
lactation is particularly important for the establishment of neural connections related to the
control of body weight. Notably, the effects of both leptin (Kamitakahara et al., 2018) and
MHFD-L (Vogt et al., 2014) on targeting of AgRP projections display considerable regional
specificity.
110
Adult mice placed on HFD display a marked hypothalamic gliosis that reveals an acute
inflammatory response, which presages significant weight gain (Horvath et al., 2010;
Valdearcos et al., 2017; Spencer et al., 2019; Cansell et al., 2021). This hypothalamic
neuroinflammation is characterized by marked changes in the density and morphology of
microglia that are most pronounced in the ARH (Thaler et al., 2012; Valdearcos et al., 2014).
Microglia are the resident myeloid cells of the CNS and respond to a broad array of circulating
signals, including nutrients such as saturated fats and carbohydrates (Valdearcos et al., 2014;
Nadjar et al., 2017; Leyrolle et al., 2019; Butler et al., 2020). Moreover, activation of microglia
alone is sufficient to stimulate food intake and promote weight gain in adult mice, and
perturbations that block activation of microglia reduce the metabolic disruption associated with
neuroinflammation (Valdearcos et al., 2017; Rosin & Kurrasch 2019; Sun et al., 2024).
Because of their established role as nutrient-sensing sentinels of hypothalamic
neuroinflammation, and their documented participation in neural development (Stevens et al.,
2007; Schafer et al., 2012; Stephan et al., 2012), microglia have been proposed as possible
mediators of developmental programming caused by nutritional perturbations (Rosin &
Kurrasch 2019; Folick et al., 2021).
127
Evidence from several of lines of investigation indicate that microglia have multiple roles in
brain development (Bilbo & Schwarz 2009; Tremblay et al., 2011; Miyamoto et al., 2016; Li et
al., 2019). Although microglia were initially thought to remain quiescent until activation by
neuroinflammation, in vivo imaging experiments demonstrated continual activity of their cellular
processes, which actively survey their local environment (Nakajima & Kohsaka 2001; Town et
al., 2005; Wake et al., 2009; Stowell et al., 2018), including direct contact with axons and
dendrites (Schafer et al., 2013). In addition to impacting neuronal number and initial formation
of neural circuits through effects on axon targeting, microglia are thought to play an important
role in synaptic refinement through selective elimination of synapses, a process termed
synaptic pruning (Paolicelli et al., 2011; Schafer et al., 2012; Hong et al., 2016). Thus far, the
majority of developmental microglia studies have focused on their role in cortical or

hippocampal circuits. However, transcriptional profiling suggests a great deal of regional and
temporal variation in microglial cell type and activity (Hammond et al., 2019; Masuda et al.,
2020; Young et al., 2021), and the effects of dietary interventions have largely focused on the
ARH. Here we used Iba1 immunostaining to visualize and measure microglial morphology in
regions known to receive AgRP inputs. Morphological parameters of microglia were quantified
in the PVH and ARH nuclei of the hypothalamus, as well as the bed nuclei of the stria
terminalis (BST), a major limbic target of AgRP neurons, in offspring exposed to MHFD-L and
compared with offspring that were raised on a normal chow diet (NCD). We also used
genetically-targeted axonal labeling of AgRP neurons to directly visualize cellular interactions
between microglia and labeled AgRP terminals in the PVH and ARH to determine if MHFD-L
stimulates synaptic pruning in these regions. The results demonstrate regionally-specific
changes to microglia in the PVH of MHFD-L offspring, that are temporally restricted to the
period when AgRP neurons innervate the PVH. In addition, the axon labeling experiments
confirm a significant decrease in AgRP innervation of the PVH in MHFD-L offspring, and for
the first time provide direct evidence of microglial-mediated synaptic pruning of AgRP terminals
in the hypothalamus. Microglial depletion experiments determined that the significant decrease
in AgRP innervation of the PVH observed in MHFD-L offspring requires normal densities of
microglia, and that microglia are required for the weight gain seen in offspring of MHFD-L
dams at weaning. However, we did not detect a significant effect of MHFD-L on the degree of
synaptic pruning in the PVH, suggesting an alternative microglial signaling mechanism yet to
be identified.
160
161
162 **2 Results**
163
164 **2.1 Microglia exhibit morphological changes in the PVH in response to MHFD-L during**
165 **postnatal development.**
166
To assess the impact of MHFD-L on microglia in the brains of postnatal mice, we used Iba1
immunohistochemistry and confocal microscopy to visualize the distribution and morphology of
microglial cells in the PVH. Discrete regions of interest were imaged and a 3D modeling
analysis pipeline was used to measure structural changes in microglia (Figure1Ai-iii). We found
that in the PVH, the overall size of microglia was significantly enhanced in offspring exposed to
MHFD-L than those raised on NCD at P16 (Figure 1B,C). The increased size of microglia
observed in MHFD-L offspring is demonstrated through a 44% increase in the complexity of
microglial process branching, as determined by Sholl analysis (Figure 1F), as well as an 87%
increase in microglial process length (Figure 1G). Additionally, the volume of microglial cells
(volume measurements include cell body and processes; Figure 1I), and the spatial territory
they occupy (Figure 1H), was nearly doubled in MHFD-L offspring, while the density of
microglia between the dietary groups remained unaltered at P16 (Figure 1J). The changes in
microglia between NCD and MHFD-L mice at P16 (Figure 1B,C) appeared to be transient,
because by P30 (Figure 1D,E), we did not detect changes in microglial size (process
complexity, Figure 1F; and length, Figure 1G), microglia cell territory (Figure 1H), microglial
cell volume (Figure 1I), or density of microglial cells (Figure 1J) between MHFD-L and NCD
offspring. The density of microglia in the PVH was reduced from P16 to P30, although this was
independent of dietary treatment (Figure 1J). We also used Cre-dependent targeting of

synaptophysin-tdTomato to axons of AgRP neurons to assess the density of AgRP terminals in PVH regions of interest. The results confirmed that AgRP terminal density is significantly lower in the brains of MHFD-L offspring compared with NCD controls at both P16 and P30 (Figure 1K). Taken together, these results suggest that exposure to MHFD-L causes changes to the morphology of microglia that are consistent with enhanced activity and surveillance of their immediate microenvironment. Furthermore, the effects of MHFD-L on microglial morphology in the PVH correspond to an accompanying decrease in the density of AgRP inputs to the PVH.

2.2 Microglia do not exhibit morphological changes in the ARH or BST in response to MHFD-L exposure during postnatal development.

The ARH houses the cell bodies of AgRP neurons, and their number is established primarily during prenatal development (Ishii & Bouret 2012). We evaluated microglia morphology in the ARH of postnatal mice by using the same 3D modeling pipeline shown in Figure 1. In contrast to our findings in the PVH, microglial size and density in the ARH were not significantly different in NCD and MHFD-L offspring at P16 (Figure 2A,B; quantified in 2E-I). By P30 (Figure 2C,D) there was a 67% increase in process length (Figure 2F) of NCD mice compared to their P16 counterparts, but no significant differences in microglial morphology in the ARH were detected between the dietary treatment groups (Figure 2A-G). The volume of individual microglial cells (Figure 2H), as well as the spatial territory they occupy (Figure 2G), increased from P16 to P30, while the total number of microglial cells was reduced by nearly half (Figure 2I). We also measured numbers of AgRP neuronal cell bodies in the ARH in brains derived from NCD and MHFD-L offspring and confirmed that the number of AgRP neurons in the ARH is also resistant to MHFD-L exposure at both P16 and P30 (Figure 2J). In addition, we evaluated microglia and AgRP terminals in the anterolateral part of the BST (Figure 3), an extrahypothalamic target of AgRP neurons innervated during the lactational period (Bouret et al., 2004a; Cansell et al., 2012; Barbier et al., 2021). As was found for the ARH, neither the size nor number of microglia in the BST were significantly different between NCD and MHFD-L offspring at P16 or P30 (Figure 3A-D; quantified in 3E-I). Similarly, the density of AgRP terminals in the same region of interest was not affected by MHFD-L exposure (Figure 3J), although by P30 the number of AgRP terminals in the BST increased by 61% compared to their P16 counterparts (Figure 3J). Taken together, these data suggest that microglia in the ARH and BST are resistant to the morphological changes that occur in the PVH of MHFD-L offspring, suggesting a notable degree of spatial specificity in the role of hypothalamic microglia during postnatal development.

2.3 Microglia are required for changes in AgRP terminal density in PVH and body weight associated with MHFD-L exposure.

To determine if microglia are required for the observed changes in AgRP inputs to the PVH of MHFD-L offspring, the colony stimulating factor 1 receptor (CSF1R) inhibitor PLX5622 was administered from P4-P21 via daily intraperitoneal (i.p.) injection (Figure 4A). These postnatal treatments resulted in a significant decrease in microglia detected in the hypothalamus at P55 relative to age-matched controls (Figure 4B,C; quantified in 4H). Notably, the PLX5622

treatments blocked the reduction in AgRP fiber density observed in the medial dorsal
parvicellular part of the PVH (PVHmpd) of vehicle-treated MHFD-L offspring to a level that was
comparable to that of NCD offspring (Figure 4D-G; quantified in 4I). In contrast, MHFD-L
exposure did not affect the density of AgRP fibers in the lateral posterior magnocellular
compartment of the PVH (PVHpml) and no significant difference in AgRP fiber density was
detected between NCD and MHFD-L offspring treated with either PLX5622 or vehicle (Figure
4D-G; quantified in 4K), suggesting target specificity for the microglial-mediated effects on
development of AgRP inputs to the PVHmpd.
239
Depletion of microglia also appeared to protect against the increase in body weight normally
observed in MHFD-L mice. In keeping with previously published results (Vogt et al., 2014), the
body weight of vehicle-treated MHFD-L animals was significantly greater (24%), compared
with that of NCD animals at weaning (Figure 4J). However, MHFD-L mice treated with
PLX5622 during lactation exhibited significantly lower weights at weaning compared to those
of vehicle-treated MHFD-L animals (Figure 4J). Taken together, these findings suggest that
microglia mediate target-specific effects of MHFD-L exposure on the innervation of the PVH by
AgRP neurons, and that microglia play a role in mediating the effects of MHFD-L on body
weight.
249
250
251 **2.4 Engulfment of AgRP terminals by microglia in the PVH and the ARH.**
252
Microglia are thought to impact development of neuronal connections through an active
engulfment mechanism and the lysosomal associated membrane protein CD68 has
been implicated in this process. Here, we used immunohistochemistry to visualize the
presence of in Iba1-labeled microglia in mice with genetically targeted labeling of AgRP
terminals (Figure 5). Many apparent contacts between microglial processes and AgRP
terminals in the PVH and ARH were observed at P16 and P30, including internalized
AgRP terminals (Figure 5A-L). However, the extent of internalization did not appear to
be influenced by MHFD-L exposure; there were no significant differences between
internalized AgRP terminals in MHFD-L and NCD offspring at P16, in either the PVH or
ARH (Figure 5M,O). Similarly, we did not detect a statistically significant difference in
microglial CD68 levels in the PVH between diet groups at either P16 or P30 (Figure
5N). Consistent with previous reports (Wong et al., 2005; Hart et al., 2012), the density
of CD68 labeled profiles nearly doubled in the PVH between P16 and P30 (Figure 5N),
as microglia become more phagocytic with age. In the ARH, CD68 staining also
increased between P16 and P30 (Figure 5P), supporting the notion that microglia
increase their phagocytic capacity with age. Nevertheless, our analysis demonstrates
that microglia interact directly with AgRP terminals, with clear evidence of engulfment.
MHFD-L exposure does not appear to promote microglia-mediated engulfment, at least
not in the specific PVH and ARH domains examined.
272
273
274 **3 Discussion**
275

It is well established that microglia are responsive to HFD exposure in adult rodents (Thaler et al., 2012; Morari et al., 2014; Valdearcos et al., 2014; Baufeld et al., 2016; Valdearcos et al., 2017) and multiple lines of evidence support an important role for microglia in mediating key aspects of neural circuit development (Checchin et al., 2006; Hoshiko et al., 2012; Li et al., 2012; Hagemeyer et al., 2017). Here we demonstrate that microglia are required for significant elevations in body weight that emerge from postnatal exposure to HFD and are associated with a sustained decrease in the density of afferents from AgRP neurons to the PVH in offspring. Exposure to MHFD-L caused distinct morphological changes to microglia that may be consistent with enhanced activity, which were observed in the PVH, but not the ARH or BST. Moreover, the morphological changes to microglia observed appear to be primarily limited to the critical period for development of AgRP inputs to PVH neurons in MHFD-L offspring. Although our results demonstrate that microglia engage in engulfment of AgRP terminals in the PVH during development, synaptic pruning by microglia does not appear to represent the cellular mechanism mediating the effects of MHFD-L exposure on innervation of the PVH by AgRP neurons.

3.1 A Role for Microglia in Mediating Body Weight Changes Observed in MHFD-L Offspring.

Maternal HFD exposure during perinatal development leads to increased body weight, fat content, and susceptibility to diet-induced obesity in offspring (Samuelsson et al., 2008; Tamashiro et al., 2009; Masuyama & Hiramatsu 2014). The postnatal period, which corresponds to lactation, and when mice derive their nutrition primarily from milk, is especially sensitive to nutritional environment and exposure to HFD that is exclusive to this period plays a particularly dominant role in specifying metabolic phenotype later in life (Chen et al., 2008; Sun et al., 2012; Vogt et al., 2014). In the present study, global depletion of microglia with the CSF1R inhibitor PLX5622 blocked the ability of MHFD-L to increase body weight in mice by weaning at P21, indicating microglia may mediate metabolic changes caused by MHFD-L exposure. Consistent with this, microglial depletion with PLX5622 in adult mice mitigates the effects of HFD exposure (Rosin & Kurrasch 2019), and our results indicate that microglia may function similarly in offspring during postnatal life to effect changes in body weight, even if the maternal HFD exposure is restricted to the lactation period. Further studies are required to define the long-term metabolic profile resulting from developmental microglial manipulations. However, given the abundant literature on sustained impact of MHFD-L on metabolic phenotype, enduring disruptions are likely. It should be noted that PLX5622 treatment is not spatially limited to the PVH or ARH, leaving open the possibility that the effects of microglial depletion on body weight occur outside of these nuclei, or are due to collective activation of microglia in multiple components of feeding circuitry (Green et al., 2020). Localization of the specific site of action for microglial specification of mature body weight during development will require utilization of specific markers for hypothalamic microglia that account for regional and phenotypic heterogeneity, perhaps through intersectional genetic methods and combinatorial pharmacology (Hammond et al., 2019; Kim et al., 2021).

3.2 MHFD-L Induces Spatially Limited Changes in Microglial Morphology.

322
Morphological changes in microglia have been reported in response to a variety of
environmental exposures. In the hypothalamus, adult mice fed a HFD show both proliferation
and changes in microglial process length and complexity (Thaler et al., 2012; Valdearcos et al.,
2017). In our studies, MHFD-L exposure caused a marked increase in the overall size of
microglia in the PVH that is related to increases in both the length and branching complexity of
immunolabeled Iba1 cellular processes. This increase in the territory occupied by microglia in
the PVH was not accompanied by an increase in microglial number, nor were numbers of
microglia affected in the PVH by MHFD-L exposure. However, in contrast to the PVH, changes
in microglial morphology were not observed in the ARH in response to MHFD-L, although we
did observe an overall increase in process length in the ARH between P16 and P30 of both
MHFD-L and NCD mice. This finding is consistent with previously published reports on
microglial maturation (Sun et al., 2024). The spatially restricted enhancement of microglial
activation in the PVH resulting from MHFD-L exposure appears to contribute to an expansion
of parenchymal territory surveilled by PVH microglia, as reflected in the volume measurements
accomplished with geometrical modeling of process length and complexity. This interpretation
is supported by *in vitro* and *in vivo* observations of enhanced process extension and increased
neuronal interactions resulting from inflammatory activation of microglia (Wake et al., 2009;
Schafer et al., 2013; Dissing-Olesen et al., 2014; Stowell et al., 2018; Bolton et al., 2022).
341
As in the ARH, we did not observe comparable changes in microglial morphology in the BST,
an extrahypothalamic target of AgRP neurons innervated during the lactation period (Cansell
et al., 2012; Barbier et al., 2021). These observations underscore the remarkable molecular
heterogeneity of microglial phenotypes that appear to occupy various hypothalamic niches
during development, and may have equally diverse developmental roles and responses to
environmental signals (Bilbo & Schwarz, 2012; Frost & Schafer, 2016; Li & Barres, 2018;
Ngozi & Bolton, 2022). Moreover, the observed morphological changes in the PVH caused by
MHFD-L appear to be transient as there no significant differences in microglial processes by
P30, and the density of microglial cells in the PVH was significantly reduced in the older mice,
suggesting a decline in overall activity. That the observed changes in microglial morphology
occur within the critical period for development of AgRP projections to the PVH suggests a
possible role for microglia linking nutrition with specification of axonal targeting (Kamitakahara
et al., 2018).
355
356
357 **3.3 Microglia Mediate Impaired Innervation of the PVH by AgRP Neurons.**
358
AgRP neuronal projections develop primarily during the first two weeks of life, which
corresponds to a critical period for the neurotrophic action of leptin on axonal outgrowth and
targeting of AgRP inputs to distinct components of the PVH (Bouret et al., 2004a; Elson &
Simerly, 2015). Exposure to MHFD-L during this critical developmental window permanently
impairs AgRP projections to the PVH, DMH and LH (Vogt et al., 2014) that are associated with
increased body weight in adulthood. Here we confirm that MHFD-L impairs innervation of the
PVH, and this defect was apparent by P16. The PVH is innervated by AgRP neurons between
P8 and P10 (Bouret et al., 2004a). We found that depletion of microglia with PLX5622 between
P4 and P21, a period that aligns not only with AgRP innervation of the PVH but also with

maximum changes in microglial morphology, blocked the effects of MHFD-L on AgRP terminals in the PVH. However, this partial rescue of innervation appeared to be limited to the PVHmpd, suggesting that there is a regional specialization in the activity of microglia in the PVH. We did not observe a change in AgRP innervation of the BST in MHFD-L offspring that mirrored the changes seen in the PVH of the same animals, further supporting the conclusion that microglia display at least some degree of spatial heterogeneity in mediating site-specific alterations in AgRP axon targeting.

Microglia are known to impact a variety of developmental events, including alterations in cell number through programmed cell death or neurogenesis, as well as axonal targeting and remodeling of neural circuits (Frost & Schafer, 2016; Li & Barres 2018). Because PLX5622 causes a decrease in microglia in the ARH as well as in the PVH, our microglial depletion studies do not eliminate the possibility that microglia may act locally on AgRP neurons to affect axonal growth. It is unlikely that the impaired innervation of the PVH observed in MHFD-L offspring is due to a reduction in the number of AgRP neurons in the ARH (Vogt et al., 2014; Valdearcos et al., 2017). Sun and colleagues (2024) reported that microglial depletion during postnatal life increases numbers of NPY neurons, as well as enhances local densities of AgRP fibers in the ARH, possibly through enhanced formation of perineuronal nets. Depletion of microglia during gestation causes a significant decrease in the number of proopiomelanocortin (POMC) neurons in the ARH and leads to an acceleration of weight gain (Rosin & Kurrasch, 2019), consistent with neurogenesis of ARH neurons occurring in mid gestation and increased susceptibility to nutritional impacts during embryonic life (Ishii & Bouret, 2012; Elson & Simerly, 2015). Interestingly, genetic deletion of leptin receptors from myeloid cells reduced numbers of POMC neurons in the ARH, suppressed POMC innervation of the PVH, and decreased microglial process complexity (Gao et al., 2017). Taken together, these results suggest that leptin signaling in microglia may act at the level of the ARH to promote outgrowth of AgRP projections to the PVH, while MHFD-L activates microglia in the PVH to specify patterns of AgRP afferents that are not only regionally specific, but also target discrete subdomains of the PVH. However, whether microglia inhibit synaptogenesis or are involved in synaptic refinement through an alternative regressive mechanism will require further investigation.

3.4 Microglia in PVH Participate in Synaptic Pruning.

Microglia have been proposed as mediators of synaptic pruning, a process whereby synapses that form early in development are eliminated as others are strengthened and maintained (Katz & Shatz, 1996; Sanes & Lichtman, 1999; Frost & Schafer 2016; Li & Barres, 2018). Although this process has been studied most extensively in somatosensory cortex (Miyamoto et al., 2016), hippocampus (Paolicelli et al., 2011; Wang et al., 2020) and the visual system (Tremblay et al., 2010; Schafer et al., 2012), there is evidence for involvement of microglia in synaptic pruning of immunolabeled glutamatergic terminals associated with corticotropin releasing hormone neurons in the PVH (Bolton et al., 2022). In the present study, we used genetically targeted axonal labeling to provide evidence that microglia in the PVH participate in synaptic pruning of AgRP synapses during the critical period for PVH innervation, and when PVH microglia exhibit high levels of process extension. The lysosomal marker CD68 was colocalized with internalized AgRP terminals in PVH microglia, and although elevated at P30,

there were no differences between offspring of MHFD-L and NCD dams. Additionally, we did
not find a significant difference in the density of engulfed AgRP terminals in the PVH of MHFD-
L offspring at either P16 or P30. However, enhanced engulfment of AgRP terminals in MHFD-L
offspring may occur at a later point in development not assessed in this study. It is also
possible that the synaptophysin-tdTomato axonal label may have been lost from pruned
synapses during engulfment. A more probable interpretation is that PVH microglia participate
in synaptic refinement through other cellular mechanisms, including microglial release of
secreted factors such as the interleukin IL-6 (Kim et al., 2024) or microglial-derived BDNF
(Parkhurst et al., 2013). Future studies that include cell type-specific manipulations of
microglial signaling and live cell imaging may clarify these potential developmental
mechanisms.
425
426
427 **4 Conclusions**
428
MHFD-L causes elevated levels of saturated carbohydrates and fats in milk (Gorski et al.,
2006; Vogt et al., 2014; Calvo-Lerma et al., 2022). The resulting overnutrition resulting from
exposure to this enhanced diet is thought to underlie the propensity towards obesity observed
in offspring later in life (Skowronski et al., 2023). Although microglia are likely mediators of
multiple neurobiological events influencing how hypothalamic circuits function during regulation
of energy balance, the precise signaling mechanisms remain ill defined. There may be
common molecular mechanisms underlying the effects of HFD exposure on microglial
activation in adults and those occurring during postnatal development, but how these signaling
events exert a lasting impact on the organization and function of feeding circuitry has not been
defined. The results presented here demonstrate an important role for microglia on sculpting
the density of inputs from AgRP neurons to the PVH that is not only spatially restricted, but
also aligned temporally with synaptogenesis in the PVH. Furthermore, PVH microglia clearly
interact directly with AgRP afferent axons during this critical period and may be refined through
engulfment by microglia. However, synaptic pruning through engulfment does not appear to be
sufficient to affect the significant reduction in AgRP innervation of PVH neurons observed
following MHFD-L exposure, suggesting involvement of additional microglial signaling
mechanisms that are not only important for specifying patterns of innervation of the PVH by
AgRP neurons, but may also contribute more broadly to developmental programming of
metabolic phenotype.
448
449

Reagent type or resource	Designation	Source or reference	Identifiers	Additional information

Genetic reagent (*M. musculus*)	*Agrp*^{tm1(cre)Lowl}/J	Jackson Laboratory	Stock #: 012899 RRID: IMSR_JAX:012899	MGI ID: J:140858
Genetic reagent (*M. musculus*)	Ai34(RCL-Syp/tdT)-D (B6;129S-*Gt(ROSA)26Sor*^{tm34.1(CAG-Syp/tdTomato)/Hze}/J	Jackson Laboratory	Stock #: 012570 RRID: IMSR_JAX:012570	MGI ID: J:170755
Antibody	Rabbit polyclonal anti-Iba1	FUJIFILM Wako	Cat. #: 019-19741 RRID: AB_ 839504	IHC (1:2000)
Antibody	Rat monoclonal anti-CD68 [FA-11]	Abcam	Cat. #: ab53444 RRID: AB_869007	IHC (1:500)
Antibody	Donkey polyclonal anti-rabbit Alexa Fluor 488	ThermoFisher Scientific	Cat. #: A32790 RRID: AB_2762833	IHC (1:500)
Antibody	Donkey polyclonal anti-rat Alexa Fluor 647	ThermoFisher Scientific	Cat. #: A48272 AB_2893138	IHC (1:500)
Chemical compound	PLX5622 hemifumarate, CSF1R inhibitor	MedChemExpress	Cat. #: HY114153A	
Software, Algorithm	Imaris	Bitplane	V9.5	
Software, Algorithm	GraphPad Prism	Prism	Prism 10	

5 Materials and Methods

5.1 Key Resources Table

5.2 Animals

All animal care and experimental procedures were performed in accordance with the guidelines of the National Institutes of Health and the Institutional Care and Use Committee of Vanderbilt University. Mice were housed at 22°C on a 12:12 h light:dark cycle provided ad libitum access to NCD (PicoLab Rodent Diet 20 #5053) and water unless otherwise specified. AgRP-Cre mice *(Agrp*^{tm1(cre)Lowl}/J; stock number: 01289) and mice expressing the Cre-dependent fluorescent reporter synaptophysin-tdTomato (RCL-Syp/tdT)-D (B6;129S-

Gt(ROSA)26Sor$^{tm34.1(CAG-Syp/tdTomato)/Hze}$/J; stock number: 012570) were obtained from the
Jackson Laboratory (Bar Harbor, ME) and maintained in our colony at Vanderbilt University. To
visualize AgRP inputs, *Agrp*-Cre mice were crossed with Ai34D mice to generate Agrp-
Cre::Ai34D mice, as described previously (Biddinger et al., 2020).

To generate offspring of dams exposed to HFD during lactation (MHFD-L), mice had ad libitum
access to NCD (PicoLab Rodent Diet 20 #5053: 25% protein; 62% carbohydrates; 13% fat; 4
kcal/g energy density) prior to and during mating. On the first postnatal day (P1) all litters were
adjusted to 7 pups to normalize nutrition and dams were switched to either HFD (Research
Diets D12451: 20% protein; 35% carbohydrate; 45% fat; 4.7 kcal/g energy density) or kept on
the same NCD. The dams remained on either HFD or NCD throughout lactation, and offspring
were weaned onto the same normal chow diet that the dams had received prior to MHFD-L
treatment, regardless of lactation dietary condition, and remained on NCD until they were
processed for perfusion.

5.3 PLX5622 Microglia Depletion
To reduce microglia during postnatal development, mouse pups were treated daily from P4 to
P21 with either PLX5622, a colony stimulating factor 1 receptor (CSF1R) inhibitor or DMSO
vehicle via intraperitoneal injection (Riquier & Sollars 2020). Briefly, PLX5622 hemifurate solid
(Cat. #HY114153A MedChemExpress, Monmouth Junction, NJ, USA) was suspended in
DMSO at a concentration of 172 mg/ml. The injection working solution was prepared to include
20% Kolliphor RH40 diluted in PBS, which resulted in doses with a 6.5mg/ml PLX5622
concentration and injection concentration of 15 mg/kg.

5.4 Immunohistochemistry
Mice were perfused at P16 and P30 and processed for immunofluorescence by using primary
antibodies to Iba1 (1:2000; FUJIFILM Wako, Osaka, Japan) to visualize microglia and CD68
(1:500; Abcam, Cambridge, MA, USA) to assess phagocytic capacity of the microglia. Mice
were first anesthetized with tribromoethanol (TBE) and then perfused transcardially with cold
0.9% saline, followed by cold fixative (4% paraformaldehyde in borate buffer, pH 9.5) for 10
minutes. After fixative perfusion, brains were removed from the skull and postfixed in the same
fixative overnight. The next day, the tissue was transferred to 20% sucrose for cryoprotection
overnight. A freezing-stage sliding microtome was used to collect 30 μm-thick coronal sections
and free-floating tissue sections were stored in cryoprotectant solution at -20°C until further
processing. To prepare tissue for immunohistochemical processing, brain sections were
removed from cryoprotectant and rinsed several times in 0.02M KPBS. Free-floating sections
were incubated in blocking buffer containing 2% normal donkey serum and 0.3% Triton-X 100
in 0.02M KPBS overnight at 4°C.Tissue sections were incubated in the same blocking buffer
with primary antibodies for 48 hr at 4°C. Following primary antibody incubation, sections were
rinsed several times in KPBS, and then incubated in the appropriate species-specific
fluorophore-conjugated Alexa-Fluor secondary antibodies for 1 hr at RT. Tissue sections were
again rinsed several times with KPBS, and mounted onto charged microscope slides and
coverslipped using ProLong antifade mounting medium (Life Technologies, Carlsbad, CA,
USA).

5.5 Image Acquisition and Analysis

Sections through the PVH, ARH, and BST were identified and morphological features of the
nuclei were visualized with Hoescht 33342, which were then used to define matching regions
of interest (ROI) for quantitative analysis carried out by a user blind to treatment group.
Confocal image stacks were collected using a laser scanning confocal microscope (Zeiss LSM
800) for each ROI through the entire thickness of the region at a frequency of 0.1μm using the
40x objective. Imaris visualization software (Bitplane V9.5, Salisbury Cove, ME, USA) was
used to create 3D reconstructions of each multichannel set of images.
515
Profiles of Iba1-immunolabeled microglia were segmented and skeletonized by using the
Filaments tool in Imaris to quantify changes in microglia structure. Sholl analysis (Norris et al.,
2014), was performed on the skeletonized structures to determine the branching complexity of
microglial processes. Briefly, 3D concentric spheres are drawn around each selected microglial
cell and contact points between microglial processes and the spheres counted. 3D
reconstructions of the microglia were also evaluated for process length and cell volume. To
estimate the 3D space occupied by each microglial cell, a polyhedron was drawn around the
microglia by using the built-in Convex Hull function under Filaments to estimate the regional
volume occupied by each cell analyzed.
525
In order to assess cellular interactions between AgRP terminals and microglia, 3D renderings
of microglia and AgRP terminals were used to determine densities of contact points between
AgRP terminals and microglial processes by using an Imaris MATLAB script to automate the
analysis. AgRP terminals were reconstructed as "spots" of 0.8 mm diameter (corresponding to
the largest measured size) and their total number was calculated in each ROI. Briefly, the
automatic detection algorithm applies a 3D Mexican hat filter using the spot size and then
locates the spot centroid at the local maxima of the filtered image. The number of spots
located at no more than 1μm from the microglia surface was automatically determined as an
estimate of contact points between AgRP terminals and microglia. Next, spots that were
determined to be more than 0.5μm away from internal microglial surfaces were determined to
be internalized by the microglia and counted as engulfed AgRP terminals. Intracellular CD68
levels in microglia were estimated by segmentation of CD68 labeling profiles in image stacks
followed by creation of 3D renderings and their volume computed. Finally, to assess the effect
of MHFD-L on numbers of AgRP neurons in the ARH, AgRP neuronal cell bodies were
identified by aligning synaptophysin-tdTomato labeled somal profiles with Hoescht 33342-
stained nuclei and counting the number of visualized neurons manually.
542
543
544
5.6 Statistical Analyses
Data are presented as group mean values ± SEM, as well as individual data points.
Statistical analyses were performed using GraphPad Prism software (Version 10).
Unpaired t-tests were used to compare data between two groups. In the microglia
depletion experiment, two-way analysis of variance (ANOVA) was used to test for
differences in group means, followed by Tukey's multiple comparisons post hoc test to
identify specific group differences. Differences between groups were considered
statistically significant at $p<0.05$.
553

Acknowledgments

We thank the members of the Simerly Lab for comments and discussion on early versions of this manuscript and Nicholas Thomas-Low for assistance with the figures. This work was supported by NIH grants R01DK106476 (RBS) and T32DK07563 (HNM-R).

559
560

Author Contributions

HNM-R and RBS conceived of this research and designed experiments. HNM-R conducted experiments with technical support from JEB and MNB. HNM-R and JEB analyzed data and prepared figures. HNM-R and RBS wrote the manuscript and all authors participated in editing the manuscript.

Data Availability

All data collected in this study are included in the manuscript and have been deposited in the Open Science Framework.

570
571

References

Andersen, L. G., Holst, C., Michaelsen, K. F., Baker, J. L., & Sørensen, T. I. A. (2012). Weight and weight gain during early infancy predict childhood obesity: a case-cohort study. *International Journal of Obesity*, *36*(10), 1306–1311. https://doi.org/10.1038/ijo.2012.134

Barbier, M., González, J. A., Houdayer, C., Burdakov, D., Risold, P., & Croizier, S. (2021). Projections from the dorsomedial division of the bed nucleus of the stria terminalis to hypothalamic nuclei in the mouse. *Journal of Comparative Neurology*, *529*(5), 929–956. https://doi.org/10.1002/cne.24988

Baufeld, C., Osterloh, A., Prokop, S., Miller, K. R., & Heppner, F. L. (2016). High-fat diet-induced brain region-specific phenotypic spectrum of CNS resident microglia. *Acta Neuropathologica*, *132*(3), 361–375. https://doi.org/10.1007/s00401-016-1595-4

Betley, J. N., Xu, S., Cao, Z. F. H., Gong, R., Magnus, C. J., Yu, Y., & Sternson, S. M. (2015). Neurons for hunger and thirst transmit a negative-valence teaching signal. *Nature, 521*(7551), 180-185. https://doi.org/10.1038/nature14416

Biddinger, J. E., Lazarenko, R. M., Scott, M. M., & Simerly, R. (2020). Leptin suppresses development of GLP-1 inputs to the paraventricular nucleus of the hypothalamus. *ELife*, *9*, e59857. https://doi.org/10.7554/elife.59857

Bilbo, S. D., & Schwarz, J. M. (2009). Early-Life Programming of Later-Life Brain and Behavior: A Critical Role for the Immune System. *Frontiers in Behavioral Neuroscience*, *3*, 14. https://doi.org/10.3389/neuro.08.014.2009

Bilbo, S. D., & Schwarz, J. M. (2012). The immune system and developmental programming of brain and behavior. *Frontiers in Neuroendocrinology*, *33*(3), 267–286. https://doi.org/10.1016/j.yfrne.2012.08.006

Bolton, J. L., Short, A. K., Othy, S., Kooiker, C. L., Shao, M., Gunn, B. G., Beck, J., Bai, X., Law, S. M., Savage, J. C., Lambert, J. J., Belelli, D., Tremblay, M.-È., Cahalan, M. D., & Baram, T. Z. (2022). Early stress-induced impaired microglial pruning of excitatory synapses on immature CRH-expressing neurons provokes aberrant adult stress responses. *Cell Reports*, *38*(13), 110600. https://doi.org/10.1016/j.celrep.2022.110600

Bouret, S. G., Draper, S. J., & Simerly, R. B. (2004a). Formation of Projection Pathways from the Arcuate Nucleus of the Hypothalamus to Hypothalamic Regions Implicated in the Neural Control of Feeding Behavior in Mice. *The Journal of Neuroscience*, *24*(11), 2797–2805. https://doi.org/10.1523/jneurosci.5369-03.2004

Bouret, S. G., Draper, S. J., & Simerly, R. B. (2004b). Trophic Action of Leptin on Hypothalamic Neurons That Regulate Feeding. *Science*, *304*(5667), 108–110. https://doi.org/10.1126/science.1095004

Bouret, S., Levin, B. E., & Ozanne, S. E. (2015). Gene-Environment Interactions Controlling Energy and Glucose Homeostasis and the Developmental Origins of Obesity. *Physiological Reviews*, *95*(1), 47–82. https://doi.org/10.1152/physrev.00007.2014

Bouyer, K., & Simerly, R. B. (2013). Neonatal Leptin Exposure Specifies Innervation of Presympathetic Hypothalamic Neurons and Improves the Metabolic Status of Leptin-Deficient Mice. *The Journal of Neuroscience*, *33*(2), 840–851. https://doi.org/10.1523/jneurosci.3215-12.2013

Butler, M. J., Cole, R. M., Deems, N. P., Belury, M. A., & Barrientos, R. M. (2020). Fatty food, fatty acids, and microglial priming in the adult and aged hippocampus and amygdala. *Brain, Behavior, and Immunity*, *89*, 145–158. https://doi.org/10.1016/j.bbi.2020.06.010

Calvo-Lerma, J., Selma-Royo, M., Hervas, D., Yang, B., Intonen, L., González, S., Martínez-Costa, C., Linderborg, K. M., & Collado, M. C. (2022). Breast Milk Lipidome Is Associated With Maternal Diet and Infants' Growth. *Frontiers in Nutrition*, *9*, 854786. https://doi.org/10.3389/fnut.2022.854786

Cansell, C., Denis, R. G. P., Joly-Amado, A., Castel, J., & Luquet, S. (2012). Arcuate AgRP neurons and the regulation of energy balance. *Frontiers in Endocrinology*, *3*, 169. https://doi.org/10.3389/fendo.2012.00169

Cansell, C., Stobbe, K., Sanchez, C., Thuc, O. L., Mosser, C., Ben-Fradj, S., Leredde, J., Lebeaupin, C., Debayle, D., Fleuriot, L., Brau, F., Devaux, N., Benani, A., Audinat, E., Blondeau, N., Nahon, J., & Rovère, C. (2021). Dietary fat exacerbates postprandial hypothalamic inflammation involving glial fibrillary acidic protein-positive cells and microglia in male mice. *Glia*, *69*(1), 42–60. https://doi.org/10.1002/glia.23882

Checchin, D., Sennlaub, F., Levavasseur, E., Leduc, M., & Chemtob, S. (2006). Potential Role
of Microglia in Retinal Blood Vessel Formation. *Investigative Opthalmology & Visual*
Science, *47*(8), 3595. https://doi.org/10.1167/iovs.05-1522

Chen, H., Simar, D., Lambert, K., Mercier, J., and Morris, M.J. (2008). Maternal and postnatal
overnutrition differentially impact appetite regulators and fuel metabolism. *Endocrinology*
149, 5348–5356. https://doi-org/10.1210/en.2008-0582

Chen, Y., Lin, Y. C., Kuo, T. W., & Knight, Z. A. (2015). Sensory detection of food
rapidly modulates arcuate feeding circuits. *Cell*, *160*(5), 829-841.
https://doi.org/10.1016/j.cell.2015.01.033

Dissing-Olesen, L., LeDue, J. M., Rungta, R. L., Hefendehl, J. K., Choi, H. B., & MacVicar, B.
A. (2014). Activation of Neuronal NMDA Receptors Triggers Transient ATP-Mediated
Microglial Process Outgrowth. *The Journal of Neuroscience*, *34*(32), 10511–10527.
https://doi.org/10.1523/jneurosci.0405-14.2014

Elson, A. E., & Simerly, R. B. (2015). Developmental specification of metabolic circuitry.
Frontiers in Neuroendocrinology, *39*, 38–51. https://doi.org/10.1016/j.yfrne.2015.09.003

Folick, A., Koliwad, S. K., & Valdearcos, M. (2021). Microglial Lipid Biology in the
Hypothalamic Regulation of Metabolic Homeostasis. *Frontiers in Endocrinology*, *12*,
668396. https://doi.org/10.3389/fendo.2021.668396

Frost, J. L., & Schafer, D. P. (2016). Microglia: Architects of the Developing Nervous System.
Trends in Cell Biology, *26*(8), 587–597. https://doi.org/10.1016/j.tcb.2016.02.006

Gao, Y., Vidal-Itriago, A., Milanova, I., Korpel, N. L., Kalsbeek, M. J., Tom, R. Z., Kalsbeek, A.,
Hofmann, S. M., & Yi, C.-X. (2017). Deficiency of leptin receptor in myeloid cells disrupts
hypothalamic metabolic circuits and causes body weight increase. *Molecular Metabolism*, *7*,
155–160. https://doi.org/10.1016/j.molmet.2017.11.003

García-Cáceres, C., Balland, E., Prevot, V., Luquet, S., Woods, S. C., Koch, M., Horvath, T. L.,
Yi, C.-X., Chowen, J. A., Verkhratsky, A., Araque, A., Bechmann, I., & Tschöp, M. H. (2019).
Role of astrocytes, microglia, and tanycytes in brain control of systemic metabolism. *Nature*
Neuroscience, *22*(1), 7–14. https://doi.org/10.1038/s41593-018-0286-y

Norris, G., Derecki, N., & Kipnis, J. (2014). Microglial Scholl Analysis. *Protocol Exchange*.
https://doi.org/10.1038/protex.2014.029

Gorski, J. N., Dunn-Meynell, A. A., Hartman, T. G., & Levin, B. E. (2006). Postnatal
environment overrides genetic and prenatal factors influencing offspring obesity and insulin
resistance. *American Journal of Physiology-Regulatory, Integrative and Comparative*
Physiology, *291*(3), R768–R778. https://doi.org/10.1152/ajpregu.00138.2006

Green, K. N., Crapser, J. D., & Hohsfield, L. A. (2020). To Kill a Microglia: A Case for CSF1R Inhibitors. *Trends in Immunology*, *41*(9), 771–784. https://doi.org/10.1016/j.it.2020.07.001

Hagemeyer, N., Hanft, K.-M., Akriditou, M.-A., Unger, N., Park, E. S., Stanley, E. R., Staszewski, O., Dimou, L., & Prinz, M. (2017). Microglia contribute to normal myelinogenesis and to oligodendrocyte progenitor maintenance during adulthood. *Acta Neuropathologica*, *134*(3), 441–458. https://doi.org/10.1007/s00401-017-1747-1

Hammond, T. R., Dufort, C., Dissing-Olesen, L., Giera, S., Young, A., Wysoker, A., Walker, A. J., Gergits, F., Segel, M., Nemesh, J., Marsh, S. E., Saunders, A., Macosko, E., Ginhoux, F., Chen, J., Franklin, R. J. M., Piao, X., McCarroll, S. A., & Stevens, B. (2019). Single-Cell RNA Sequencing of Microglia throughout the Mouse Lifespan and in the Injured Brain Reveals Complex Cell-State Changes. *Immunity*, *50*(1), 253-271.e6. https://doi.org/10.1016/j.immuni.2018.11.004

Hart, A. D., Wyttenbach, A., Perry, V. H., & Teeling, J. L. (2012). Age related changes in microglial phenotype vary between CNS regions: Grey versus white matter differences. *Brain, Behavior, and Immunity*, *26*(5), 754–765. https://doi.org/10.1016/j.bbi.2011.11.006

Hong, S., Dissing-Olesen, L., & Stevens, B. (2016). New insights on the role of microglia in synaptic pruning in health and disease. *Current Opinion in Neurobiology*, *36*, 128–134. https://doi.org/10.1016/j.conb.2015.12.004

Horvath, T. L., Sarman, B., García-Cáceres, C., Enriori, P. J., Sotonyi, P., Shanabrough, M., Borok, E., Argente, J., Chowen, J. A., Perez-Tilve, D., Pfluger, P. T., Brönneke, H. S., Levin, B. E., Diano, S., Cowley, M. A., & Tschöp, M. H. (2010). Synaptic input organization of the melanocortin system predicts diet-induced hypothalamic reactive gliosis and obesity. *Proceedings of the National Academy of Sciences*, *107*(33), 14875–14880. https://doi.org/10.1073/pnas.1004282107

Hoshiko, M., Arnoux, I., Avignone, E., Yamamoto, N., & Audinat, E. (2012). Deficiency of the Microglial Receptor CX3CR1 Impairs Postnatal Functional Development of Thalamocortical Synapses in the Barrel Cortex. *The Journal of Neuroscience*, *32*(43), 15106–15111. https://doi.org/10.1523/jneurosci.1167-12.2012

Ishii, Y., & Bouret, S. G. (2012). Embryonic Birthdate of Hypothalamic Leptin-Activated Neurons in Mice. *Endocrinology*, *153*(8), 3657–3667. https://doi.org/10.1210/en.2012-1328

Kamitakahara, A., Bouyer, K., Wang, C., & Simerly, R. (2018). A critical period for the trophic actions of leptin on AgRP neurons in the arcuate nucleus of the hypothalamus. *Journal of Comparative Neurology*, *526*(1), 133–145. https://doi.org/10.1002/cne.24327

Katz, L. C., & Shatz, C. J. (1996). Synaptic Activity and the Construction of Cortical Circuits. *Science*, *274*(5290), 1133–1138. https://doi.org/10.1126/science.274.5290.1133

Kim, J. D., Copperi, F., & Diano, S. (2024). Microglia in Central Control of Metabolism. *Physiology*, *39*(1), 5–17. https://doi.org/10.1152/physiol.00021.2023

Kim, J.-S., Kolesnikov, M., Peled-Hajaj, S., Scheyltjens, I., Xia, Y., Trzebanski, S., Haimon, Z., Shemer, A., Lubart, A., Hove, H. V., Chappell-Maor, L., Boura-Halfon, S., Movahedi, K., Blinder, P., & Jung, S. (2021). A Binary Cre Transgenic Approach Dissects Microglia and CNS Border-Associated Macrophages. *Immunity*, *54*(1), 176-190.e7. https://doi.org/10.1016/j.immuni.2020.11.007

Krashes, M. J., Koda, S., Ye, C., Rogan, S. C., Adams, A. C., Cusher, D. S., ... & Lowell, B. B. (2011). Rapid, reversible activation of AgRP neurons drives feeding behavior in mice. The Journal of clinical investigation, 121(4), 1424-1428. https://www.jci.org/articles/view/46229

Leyrolle, Q., Layé, S., & Nadjar, A. (2019). Direct and indirect effects of lipids on microglia function. *Neuroscience Letters*, *708*, 134348. https://doi.org/10.1016/j.neulet.2019.134348

Li, Q., & Barres, B. A. (2018). Microglia and macrophages in brain homeostasis and disease. *Nature Reviews Immunology*, *18*(4), 225–242. https://doi.org/10.1038/nri.2017.125

Li, Q., Cheng, Z., Zhou, L., Darmanis, S., Neff, N. F., Okamoto, J., Gulati, G., Bennett, M. L., Sun, L. O., Clarke, L. E., Marschallinger, J., Yu, G., Quake, S. R., Wyss-Coray, T., & Barres, B. A. (2019). Developmental Heterogeneity of Microglia and Brain Myeloid Cells Revealed by Deep Single-Cell RNA Sequencing. *Neuron*, *101*(2), 207-223.e10. https://doi.org/10.1016/j.neuron.2018.12.006

Li, Y., Du, X., Liu, C., Wen, Z., & Du, J. (2012). Reciprocal Regulation between Resting Microglial Dynamics and Neuronal Activity In Vivo. *Developmental Cell*, *23*(6), 1189–1202. https://doi.org/10.1016/j.devcel.2012.10.027

Lippert, R. N., & Brüning, J. C. (2022). Maternal Metabolic Programming of the Developing Central Nervous System: Unified Pathways to Metabolic and Psychiatric Disorders. *Biological Psychiatry*, *91*(10), 898–906. https://doi.org/10.1016/j.biopsych.2021.06.002

Masuda, T., Sankowski, R., Staszewski, O., & Prinz, M. (2020). Microglia Heterogeneity in the Single-Cell Era. *Cell Reports*, *30*(5), 1271–1281. https://doi.org/10.1016/j.celrep.2020.01.010

Masuyama, H., & Hiramatsu, Y. (2014). Additive Effects of Maternal High-Fat Diet during Lactation on Mouse Offspring. *PLoS ONE*, *9*(3), e92805. https://doi.org/10.1371/journal.pone.0092805

Miyamoto, A., Wake, H., Ishikawa, A. W., Eto, K., Shibata, K., Murakoshi, H., Koizumi, S., Moorhouse, A. J., Yoshimura, Y., & Nabekura, J. (2016). Microglia contact induces synapse formation in developing somatosensory cortex. *Nature Communications*, *7*(1), 12540. https://doi.org/10.1038/ncomms12540

Morari, J., Anhe, G. F., Nascimento, L. F., Moura, R. F. de, Razolli, D., Solon, C., Guadagnini, D., Souza, G., Mattos, A. H., Tobar, N., Ramos, C. D., Pascoal, V. D., Saad, M. J., Lopes-Cendes, I., Moraes, J. C., & Velloso, L. A. (2014). Fractalkine (CX3CL1) is involved in the early activation of hypothalamic inflammation in experimental obesity. *Diabetes*, *63*(11), 3770–3784. https://doi.org/10.2337/db13-1495

Nadjar, A., Leyrolle, Q., Joffre, C., & Laye, S. (2017). Bioactive lipids as new class of microglial modulators: When nutrition meets neuroimmunology. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, *79*(Pt A), 19–26. https://doi.org/10.1016/j.pnpbp.2016.07.004

Nakajima, K., & Kohsaka, S. (2001). Microglia: Activation and Their Significance in the Central Nervous System. *The Journal of Biochemistry*, *130*(2), 169–175. https://doi.org/10.1093/oxfordjournals.jbchem.a002969

Ngozi, Z., & Bolton, J. L. (2022). Microglia Don't Treat All Neurons the Same: The Importance of Neuronal Subtype in Microglia-Neuron Interactions in the Developing Hypothalamus. *Frontiers in Cellular Neuroscience*, *16*, 867217. https://doi.org/10.3389/fncel.2022.867217

Norris, G., Derecki, N., & Kipnis, J. (2014). Microglial Scholl Analysis. *Protocol Exchange*. https://doi.org/10.1038/protex.2014.029

Paolicelli, R. C., Bolasco, G., Pagani, F., Maggi, L., Scianni, M., Panzanelli, P., Giustetto, M., Ferreira, T. A., Guiducci, E., Dumas, L., Ragozzino, D., & Gross, C. T. (2011). Synaptic Pruning by Microglia Is Necessary for Normal Brain Development. *Science*, *333*(6048), 1456–1458. https://doi.org/10.1126/science.1202529

Parkhurst, C. N., Yang, G., Ninan, I., Savas, J. N., Yates, J. R., Lafaille, J. J., Hempstead, B. L., Littman, D. R., & Gan, W.-B. (2013). Microglia Promote Learning-Dependent Synapse Formation through Brain-Derived Neurotrophic Factor. *Cell*, *155*(7), 1596–1609. https://doi.org/10.1016/j.cell.2013.11.030

Plagemann, A., Heidrich, I., Götz, F., Rohde, W., & Dörner, G. (1992). Obesity and Enhanced Diabetes and Cardiovascular Risk in Adult Rats due to Early Postnatal Overfeeding. *Experimental and Clinical Endocrinology*, *99*(03), 154–158. https://doi.org/10.1055/s-0029-1211159

Prospective Studies Collaboration; Whitlock, G., Lewington, S., Sherliker, P., Clarke, R., Emberson, J., Halsey, J., Qizilbash, N., Collins, R., & Peto, R. (2009). Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet*, *373*(9669), 1083–1096. https://doi.org/10.1016/s0140-6736(09)60318-4

Riquier, A. J., & Sollars, S. I. (2020). Astrocytic response to neural injury is larger during development than in adulthood and is not predicated upon the presence of microglia. *Brain, Behavior, & Immunity - Health*, *1*, 100010. https://doi.org/10.1016/j.bbih.2019.100010

Rosin, J. M., & Kurrasch, D. M. (2019). Emerging roles for hypothalamic microglia as
regulators of physiological homeostasis. *Frontiers in Neuroendocrinology*, 100748.
https://doi.org/10.1016/j.yfrne.2019.100748

Samuelsson, A.-M., Matthews, P. A., Argenton, M., Christie, M. R., McConnell, J. M., Jansen,
E. H. J. M., Piersma, A. H., Ozanne, S. E., Twinn, D. F., Remacle, C., Rowlerson, A.,
Poston, L., & Taylor, P. D. (2008). Diet-Induced Obesity in Female Mice Leads to Offspring
Hyperphagia, Adiposity, Hypertension, and Insulin Resistance: A Novel Murine Model of
Developmental Programming. *Hypertension*, *51*(2), 383–392.
https://doi.org/10.1161/hypertensionaha.107.101477

Sanes, J. R., & Lichtman, J. W. (1999). DEVELOPMENT OF THE VERTEBRATE
NEUROMUSCULAR JUNCTION. *Annual Review of Neuroscience*, *22*(1), 389–442.
https://doi.org/10.1146/annurev.neuro.22.1.389

Schafer, D. P., Lehrman, E. K., Kautzman, A. G., Koyama, R., Mardinly, A. R., Yamasaki, R.,
Ransohoff, R. M., Greenberg, M. E., Barres, B. A., & Stevens, B. (2012). Microglia Sculpt
Postnatal Neural Circuits in an Activity and Complement-Dependent Manner. *Neuron*, *74*(4),
691–705. https://doi.org/10.1016/j.neuron.2012.03.026

Schafer, D. P., Lehrman, E. K., & Stevens, B. (2013). The "quad-partite" synapse:
Microglia-synapse interactions in the developing and mature CNS. *Glia*, *61*(1), 24–36.
https://doi.org/10.1002/glia.22389

Simerly, R. B. (2008). Hypothalamic substrates of metabolic imprinting. *Physiology & Behavior*,
94(1), 79–89. https://doi.org/10.1016/j.physbeh.2007.11.023

Skowronski, A. A., Leibel, R. L., & LeDuc, C. A. (2023). Neurodevelopmental Programming of
Adiposity: Contributions to Obesity Risk. *Endocrine Reviews*, *45*(2), 253–280.
https://doi.org/10.1210/endrev/bnad031

Spencer, S. J., Basri, B., Sominsky, L., Soch, A., Ayala, M. T., Reineck, P., Gibson, B. C., &
Barrientos, R. M. (2019). High-fat diet worsens the impact of aging on microglial function
and morphology in a region-specific manner. *Neurobiology of Aging*, *74*, 121–134.
https://doi.org/10.1016/j.neurobiolaging.2018.10.018

Stephan, A. H., Barres, B. A., & Stevens, B. (2012). The Complement System: An Unexpected
Role in Synaptic Pruning During Development and Disease. *Neuroscience*, *35*(1), 369–389.
https://doi.org/10.1146/annurev-neuro-061010-113810

Stettler, N., Stallings, V. A., Troxel, A. B., Zhao, J., Schinnar, R., Nelson, S. E., Ziegler, E. E.,
& Strom, B. L. (2005). Weight gain in the first week of life and overweight in adulthood: a
cohort study of European American subjects fed infant formula. *Circulation*, *111*(15), 1897–
1903. https://doi.org/10.1161/01.cir.0000161797.67671.a7

Stevens, B., Allen, N. J., Vazquez, L. E., Howell, G. R., Christopherson, K. S., Nouri, N., Micheva, K. D., Mehalow, A. K., Huberman, A. D., Stafford, B., Sher, A., Litke, A. M., Lambris, J. D., Smith, S. J., John, S. W. M., & Barres, B. A. (2007). The Classical Complement Cascade Mediates CNS Synapse Elimination. *Cell*, *131*(6), 1164–1178. https://doi.org/10.1016/j.cell.2007.10.036

Stowell, R. D., Wong, E. L., Batchelor, H. N., Mendes, M. S., Lamantia, C. E., Whitelaw, B. S., & Majewska, A. K. (2018). Cerebellar microglia are dynamically unique and survey Purkinje neurons in vivo. *Developmental Neurobiology*, *78*(6), 627–644. https://doi.org/10.1002/dneu.22572

Sun, J., Wang, X., Sun, R., Xiao, X., Wang, Y., Peng, Y., & Gao, Y. (2024). Microglia shape AgRP neuron postnatal development via regulating perineuronal net plasticity. *Molecular Psychiatry*, 306-316. https://doi.org/10.1038/s41380-023-02326-2

Sutton Hickey, A. K., Duane, S. C., Mickelsen, L. E., Karolczak, E. O., Shamma, A. M., Skillings, A., Li, C., & Krashes, M. J. (2023). AgRP neurons coordinate the mitigation of activity-based anorexia. *Molecular Psychiatry*, 306-316. https://doi.org/10.1038/s41380-023-02326-2

Tamashiro, K. L. K., Terrillion, C. E., Hyun, J., Koenig, J.I., & Moran, T. H. (2009). Prenatal Stress or High-Fat Diet Increases Susceptibility to Diet-Induced Obesity in Rat Offspring. *Diabetes*, *58*(5), 1116-1125. https://doi-org/10.2337/db08-1129

Tamashiro, K. L. K., & Moran, T. H. (2010). Perinatal environment and its influences on metabolic programming of offspring. *Physiology & Behavior*, *100*(5), 560–566. https://doi.org/10.1016/j.physbeh.2010.04.008

Thaler, J. P., Yi, C.-X., Schur, E. A., Guyenet, S. J., Hwang, B. H., Dietrich, M. O., Zhao, X., Sarruf, D. A., Izgur, V., Maravilla, K. R., Nguyen, H. T., Fischer, J. D., Matsen, M. E., Wisse, B. E., Morton, G. J., Horvath, T. L., Baskin, D. G., Tschöp, M. H., & Schwartz, M. W. (2012). Obesity is associated with hypothalamic injury in rodents and humans. *Journal of Clinical Investigation*, *122*(1), 153–162. https://doi.org/10.1172/jci59660

Town, T., Nikolic, V., & Tan, J. (2005). The microglial "activation" continuum: from innate to adaptive responses. *Journal of Neuroinflammation*, *2*(1), 24. https://doi.org/10.1186/1742-2094-2-24

Tremblay, M.-È., Lowery, R. L., & Majewska, A. K. (2010). Microglial Interactions with Synapses Are Modulated by Visual Experience. *PLoS Biology*, *8*(11), e1000527. https://doi.org/10.1371/journal.pbio.1000527

Tremblay, M.-È., Stevens, B., Sierra, A., Wake, H., Bessis, A., & Nimmerjahn, A. (2011). The Role of Microglia in the Healthy Brain. *The Journal of Neuroscience*, *31*(45), 16064–16069. https://doi.org/10.1523/jneurosci.4158-11.2011

Valdearcos, M., Douglass, J. D., Robblee, M. M., Dorfman, M. D., Stifler, D. R., Bennett, M. L.,
Gerritse, I., Fasnacht, R., Barres, B. A., Thaler, J. P., & Koliwad, S. K. (2017). Microglial
Inflammatory Signaling Orchestrates the Hypothalamic Immune Response to Dietary
Excess and Mediates Obesity Susceptibility. *Cell Metabolism*, *26*(1), 185-197.e3.
https://doi.org/10.1016/j.cmet.2017.05.015

Valdearcos, M., Robblee, M. M., Benjamin, D. I., Nomura, D. K., Xu, A. W., & Koliwad, S. K.
(2014). Microglia Dictate the Impact of Saturated Fat Consumption on Hypothalamic
Inflammation and Neuronal Function. *Cell Reports*, *9*(6), 2124–2138.
https://doi.org/10.1016/j.celrep.2014.11.018

Vogt, M. C., Paeger, L., Hess, S., Steculorum, S. M., Awazawa, M., Hampel, B., Neupert, S.,
Nicholls, H. T., Mauer, J., Hausen, A. C., Predel, R., Kloppenburg, P., Horvath, T. L., &
Brüning, J. C. (2014). Neonatal Insulin Action Impairs Hypothalamic Neurocircuit Formation
in Response to Maternal High-Fat Feeding. *Cell*, *156*(3), 495–509.
https://doi.org/10.1016/j.cell.2014.01.008

Wake, H., Moorhouse, A. J., Jinno, S., Kohsaka, S., & Nabekura, J. (2009). Resting Microglia
Directly Monitor the Functional State of Synapses In Vivo and Determine the Fate of
Ischemic Terminals. *The Journal of Neuroscience*, *29*(13), 3974–3980.
https://doi.org/10.1523/jneurosci.4363-08.2009

Wang, C., Yue, H., Hu, Z., Shen, Y., Ma, J., Li, J., Wang, X.-D., Wang, L., Sun, B., Shi, P.,
Wang, L., & Gu, Y. (2020). Microglia mediate forgetting via complement-dependent synaptic
elimination. *Science*, *367*(6478), 688–694. https://doi.org/10.1126/science.aaz2288

Wong, A. M., Patel, N. V., Patel, N. K., Wei, M., Morgan, T. E., Beer, M. C. de, Villiers, W. J. S.
de, & Finch, C. E. (2005). Macrosialin increases during normal brain aging are attenuated
by caloric restriction. *Neuroscience Letters*, *390*(2), 76–80.
https://doi.org/10.1016/j.neulet.2005.07.058

Young, A. M., Kumasaka, N., Calvert, F., Hammond, T. R., Knights, A., Panousis, N., Park, J.
S., Schwartzentruber, J., Liu, J., Kundu, K., Segel, M., Murphy, N. A., McMurran, C. E.,
Bulstrode, H., Correia, J., Budohoski, K. P., Joannides, A., Guilfoyle, M. R., Trivedi, R., …
Gaffney, D. J. (2021). A map of transcriptional heterogeneity and regulatory variation in
human microglia. *Nature Genetics*, *53*(6), 861–868. https://doi.org/10.1038/s41588-021-
00875-2

Zagmutt, S., Mera, P., Soler-Vázquez, M. C., Herrero, L., & Serra, D. (2018). Targeting AgRP
neurons to maintain energy balance: Lessons from animal models. *Biochemical*
Pharmacology, *155*, 224–232. https://doi.org/10.1016/j.bcp.2018.07.008

Zeltser, L. M. (2018). Feeding circuit development and early-life influences on future feeding
behaviour. *Nature Reviews Neuroscience*, *19*(5), 302–316.
https://doi.org/10.1038/nrn.2018.23

877
878
879 **Figure Legends**
880
Figure 1. MHFD-L: Microglial morphology in the PVH. (A) Image analysis pipeline.
Fluorescence image of an Iba1-immunostained microglial cell in the PVH (Ai). Confocal
images through labeled cells were used to generate 3D reconstructions (Aii), which
were then used to create 3D models of microglial cells by using the Filaments tool in
Imaris. Polyhedrons were generated around each cell using the Convex Hull function of
Imaris to estimate the total tissue "territory" occupied by the microglial cell (Aiii). (B-E)
Images of microglial cells (green) and labeled AgRP terminals (red) in the PVH of mice
at P16 (B,C) or P30 (D,E) that were raised on NCD (B, D) or MHFD-L (C, E). Graphical
comparisons between groups to show that MHFD-L increased microglial ramification
complexity (F), process length (G), microglial cell territory (H), and cell volume (I) at
P16. The density of microglia in the PVH decreased between P16 and P30, irrespective
of diet (J). Density of AgRP terminals were decreased in MHFD-L offspring at both P16
and P30 (K). Bars represent the mean ± SEM and each point represents one animal.
*P<.05, **P<.005. Abbreviations: MHFD-L, maternal HFD during lactation; NCD, Normal
Chow Diet; PVH, paraventricular nucleus of the hypothalamus.
896
897
Figure 2. MHFD-L: Microglial morphology in the ARH. Microglial cells (green) and
labeled AgRP terminals (red) in the ARH of mice at P16 (A, B) or P30 (C,D) that were
exposed to NCD (A, C) or MHFD-L (B, D). Graphical comparisons between groups to
show that microglial ramification complexity (E) remained the same, regardless of age
or diet. Microglial process length (F), cell territory (G), and cell volume (H) increased
between P16 and P30, but were not changed as a result of diet. The density of
microglia in the ARH decreased between P16 and P30, irrespective of diet (I). There
were no apparent changes in numbers of AgRP neurons (J). Bars represent the mean ±
SEM and each point represents one animal. *P<.05, **P<.005. Abbreviations: ARH,
arcuate nucleus of the hypothalamus; MHFD-L, maternal HFD during lactation; NCD,
Normal Chow Diet.
909
910
Figure 3. MHFD-L: Microglial morphology in the BST. Microglial cells (green) and
labeled AgRP terminals (red) in the BST of mice at P16 (A, B) or P30 (C,D) that were
exposed to NCD (A, C) or MHFD-L (B, D). Graphical comparisons between groups to
show that microglial ramification complexity (E), process length (F), cell territory (G),
and cell volume (H) did not significantly change between P16 and P30, nor were they
changed as a result of diet. The density of microglia in the BST decreased between P16
and P30, irrespective of diet (I). The density of AgRP terminals increased between P16
and P30, but there was no effect of maternal diet (J). Bars represent the mean ± SEM
and each point represents one animal. *P<.05. Abbreviations: BST, bed nucleus of the
stria terminalis; MHFD-L, maternal HFD during lactation; NCD, Normal Chow Diet.
921
922

Figure 4. Microglial depletion during lactation period. Schematic of MHFD-L exposure and PLX5622 treatment experimental design (A). Microglial cells (green) in the PVH of adult mice (P55) treated during lactation with vehicle (B) or PLX5622 (C). Images of labeled AgRP terminals (red) to illustrate the density of labeling in distinct compartments of the PVH (white boxes denote locations of ROIs) of NCD offspring (D,E) and MHFD-L offspring (F,G). Graphical comparisons to illustrate the effects of postnatal PLX5622 treatments on microglia density in the PVH (H), body weight (J) and the density of AgRP terminals in the PVHmpd (I) and PVHpml (J). Bars represent the mean ± SEM and each point represents one animal. Unpaired t-test was used to compare cell number in 4H; two-way ANOVA was used to test for differences in group means, followed by Tukey's multiple comparisons posthoc test to identify specific group differences in 4I-K. P-values less than 0.05 were considered significant; *P<.05, **P<.005, ***P<.0005. Abbreviations: AgRP, agouti-related peptide; CSF1R, Colony Stimulating Factor 1 Receptor; MHFD-L, maternal HFD during lactation; PVH, paraventricular nucleus of the hypothalamus; mpd, medial parvocellular compartment of the PVH; pml, posterior magnocellular compartment of the PVH.

Figure 5. Microglial interaction with AgRP axon terminals in the PVH and ARH. (A-D) Representative images of microglial cells (green), labeled AgRP terminals (red), and CD68 (lysosomal associated membrane protein and phagocytic capacity marker, pink) that compare their cellular relationships in the PVH (A,B) and ARH (C,D) at P16 and P30. (E-H) Digital 3D reconstructions of cells shown in (A-D) after application of filaments tool to visualize internalized AgRP terminals. (I-L) Cells shown in E-H after application of digital zoom to more clearly illustrate engulfment of labeled AgRP terminals by microglia and location of CD-68 labeled profiles. (M-P) Graphical comparisons between groups to illustrate the effects of age and MHFD-L exposure on CD68 expression and AgRP terminal engulfment. Bars represent the mean ± SEM and each point represents one animal. *P<.05, **P<.005. Abbreviations: AgRP, agouti-related peptide; ARH, arcuate nucleus or the hypothalamus; CD68, Cluster of Differentiation 68; MHFD-L, maternal HFD during lactation; PVH, paraventricular nucleus of the hypothalamus.

M PVH Engulfed AgRP

Internalization (%) vs Age (Postnatal day)
□ NCD ■ MHFD-L

N PVH Microglial CD68

CD68 Labeling vs Age (Postnatal day)
**

O ARH Engulfed AgRP

Internalization (%) vs Age (Postnatal day)

P ARH Microglial CD68

CD68 Labeling vs Age (Postnatal day)
* *

P16 PVH
P30 PVH
P16 ARH
P30 ARH

Iba-1
CD68
AgRP-cre::SynTom

Iba-1
CD68
AgRP internalized
AgRP-cre::SynTom

15 μm
3 μm

A

PLX5622 or Vehicle (15mg/kg)

Agrp^cre::Ai34

P1   P4   P21   P55

← NCD or MHFD-L → | ← Weaned to NCD → | Analyze AgRP Terminal Density in PVH

3V — pml, mpd

B + Vehicle

C + PLX

Iba-1

40 µm

D NCD   **F** MHFD-L

3V — pml, mpd

E NCD   **G** MHFD-L

3V

AgRP

40 µm

H Iba1+ Cells

Cells (#/volume)

Vehicle   PLX5622

**

J Body Weight

Body Weight (g)

NCD/Veh   NCD/PLX   MHFD-L/Veh   MHFD-L/PLX

***   ****   *

I PVHmpd AgRP

Terminal Density (#/vol)

NCD/Veh   NCD/PLX   MHFD-L/Veh   MHFD-L/PLX

***   **

K PVHpml AgRP

Terminal Density (#/vol)

NCD/Veh   NCD/PLX   MHFD-L/Veh   MHFD-L/PLX

A
(i) Iba-1
10 μm

(ii) 3D Model

(iii) Filament model

B *AgRP* *Iba-1*
P16 PVH NCD

C P16 PVH MHFD-L

D P30 PVH NCD
20 μm

E P30 PVH MHFD-L

F Process Complexity

□ NCD ■ MHFD-L

Intersections / Age (Postnatal day)

**

G Process Length

Filament Length (μm) / Age (Postnatal day)

*

H Microglia Cell Territory

Area (μm²) / Age (Postnatal day)

*

I Cell Volume

Volume (μm³) / Age (Postnatal day)

*

J Cell Density

Cells (number/vol) / Age (Postnatal day)

***

K AgRP Terminals

Terminal Density (number/vol) / Age (Postnatal day)

*** ** *