Microglia are Required for Developmental Specification of AgRP Innervation in the Hypothalamus of Offspring Exposed to Maternal High Fat Diet During Lactation

Haley N. Mendoza-Romero, Jessica E. Biddinger, Michelle N. Bedenbaugh and Richard B. Simerly[1]

Dept of Molecular Physiology & Biophysics, Vanderbilt University, Nashville, TN, 37232, USA

[1]**Corresponding author:** Richard B. Simerly
Email: richard.simerly@vanderbilt.edu

Competing Interest Statement: No competing interests

Classifications: Major category: Developmental Biology; Minor category: Neuroscience

Keywords:
Microglia
Neural Development
Hypothalamus
Agouti-related peptide
Paraventricular hypothalamic nucleus

32
Abstract

Nutritional fluctuations that occur early in life dictate metabolic adaptations that will affect susceptibility to weight gain and obesity later in life. The postnatal period in mice represents a time of dynamic changes in hypothalamic development and maternal consumption of a high fat diet during the lactation period (MHFD) changes the composition of milk and leads to enhanced susceptibility to obesity in offspring. Agouti-related peptide (AgRP) neurons in the arcuate nucleus of the hypothalamus (ARH) react to changes in multiple metabolic signals and distribute neuroendocrine information to other brain regions, such as the paraventricular hypothalamic nucleus (PVH), which is known to integrate a variety of signals that regulate body weight. Development of neural projections from AgRP neurons to the PVH occurs during the lactation period and these projections are reduced in MHFD offspring, but underlying developmental mechanisms remain largely unknown. Microglia are the resident immune cells of the central nervous system and are involved in refinement of neural connections and modulation of synaptic transmission. Because high fat diet exposure causes activation of microglia in adults, a similar activation may occur in offspring exposed to MHFD and play a role in sculpting hypothalamic feeding circuitry. Genetically targeted axonal labeling and immunohistochemistry were used to visualize AgRP axons and microglia in postnatal mice derived from MHFD dams and morphological changes quantified. The results demonstrate regionally localized changes to microglial morphology in the PVH of MHFD offspring that suggest enhanced surveillance activity and are temporally restricted to the period when AgRP neurons innervate the PVH. In addition, axon labeling experiments confirm a significant decrease in AgRP innervation of the PVH in MHFD offspring and provide direct evidence of synaptic pruning of AgRP inputs to the PVH. Microglial depletion with the Colony-stimulating factor 1 receptor inhibitor PLX5622 determined that the decrease in AgRP innervation observed in MHFD offspring is dependent on microglia, and that microglia are required for weight gain that emerges as early as weaning in offspring of MHFD dams. However, these changes do not appear to be dependent on the degree of microglial mediated synaptic pruning. Together, these findings suggest that microglia are activated by exposure to MHFD and interact directly with AgRP axons during development to permanently alter their density, with implications for developmental programming of metabolic phenotype.

Significance Statement

Maternal high fat diet exposure results in enhanced risk for negative health outcomes in humans and multiple animal models. Here we demonstrate that microglia are required for changes in body weight and perturbations to hypothalamic circuits caused by maternal high fat diet exposure that is limited to the lactational period. We identified spatially and temporally limited morphological changes to microglia that reflect an enhancement of surveillance activity and align with a critical period of hypothalamic circuit formation. We also identify direct cellular interactions between microglia and developing axons, as well as evidence for synaptic engulfment, although this mechanism does not appear to be responsible for changes to neural patterning caused by maternal high fat diet exposure. Together these findings identify an essential role for microglia in specifying patterns of hypothalamic innervation during

77 development in response to maternal high fat diet exposure, which may contribute to
78 developmental programming of metabolic phenotype.
79
80
81 **1 Introduction**
82
Maternal nutritional status has a profound effect on the metabolic phenotype of offspring.
Children born to obese mothers experience higher rates of obesity later in life, with
accompanying comorbidities that negatively impact health and longevity (Stettler et al., 2005;
Prospective Studies Collaboration et al., 2009; Tamashiro and Moran 2010; Andersen et al.,
2012). Although this developmental programming of metabolic phenotype has been
reproduced in a number of animal models (Samuelsson et al., 2008; Masuyama et al., 2014;
Garcia-Caceres et al., 2019; Skowronski, Leibel and Leduc, 2023), the underlying mechanisms
remain poorly defined. In mouse models, maternal obesity during lactation (MHFD), a time
when offspring are dependent on milk from their mothers for nutrition, appears to be
particularly impactful. These changes to metabolic phenotype are thought to be mediated by
changes in the milk (Gorski et al., 2006; Vogt et al., 2014; Calvo-Lerma et al., 2022) and occur
without subsequent dietary challenge to the offspring themselves, suggesting that they are a
consequence of developmental programming (Bolton et al., 2022; Skowronski, Leibel and
LeDuc, 2023). Because neural circuits known to control body weight develop during the
lactational period, they are vulnerable to a variety of environmental signals that may affect their
organization and function (Horvath et al., 2010; Elson and Simerly 2015; Bouret, Levin and
Ozanne, 2015; Zeltser, 2018; Skowronski, Leibel and Leduc, 2023).
100
AgRP neurons in the arcuate nucleus of the hypothalamus (ARH) function as "hunger neurons"
that respond to key metabolic signals such as leptin, ghrelin, glucose and free fatty acids
(Krashes et al., 2011; Betley et al.,2015; Chen et al., 2015; Sutton Hickey et al., 2023), and
they distribute this information to other regions associated with energy balance regulation
(Simerly 2008; Zagmutt, et al., 2018). Thus, the ability of AgRP neurons to influence other
components of feeding circuitry is dependent on formation of their neural connections, which
form primarily during the first 2 weeks of life (Bouret, Draper and Simerly, 2004a). During
development, AgRP axons extend from the ARH at postnatal day 4 (P4) and reach the
paraventricular hypothalamic nucleus (PVH) between P8-P10. Leptin is required for normal
targeting of AgRP axons to their downstream targets, and in leptin deficient mice both
neuroanatomical and related physiological defects persist into adulthood (Bouret et al., 2004b;
Bouyer et al., 2013; Elson and Simerly 2015). Maternal overnutrition affects formation of
feeding circuits during postnatal life with concomitant dysregulation of body weight (Plagemann
et al., 1992; Elson and Simerly, 2015; Lippert and Bruning, 2022; Skowronski, Leibel and
LeDuc, 2023). Limiting high fat diet exposure of dams to the first 3 weeks of lactation (MHFD)
causes suppression of neural projections from AgRP neurons to the PVH in offspring and is
associated with increased body weight later in life (Vogt et al., 2014). In fact, MHFD was more
effective than prenatal maternal high fat diet exposure in causing body weight changes of adult
offspring. MHFD did not alter cell number, peptidergic expression, or cellular activity of AgRP
neurons in the ARH, suggesting that maternal nutritional status during lactation is particularly
important for the establishment of neural connections related to the control of body weight.

Notably, the effects of both leptin (Kamitakahara et al., 2018) and MHFD (Vogt et al., 2014) on
targeting of AgRP projections display considerable regional specificity.
124
Adult mice placed on HFD display a marked hypothalamic gliosis that reveals an acute
inflammatory response, which presages significant weight gain (Horvath et al., 2010;
Valdearcos et al., 2017; Spencer et al., 2019; Cansell et al., 2021). This hypothalamic
neuroinflammation is characterized by marked changes in the density and morphology of
microglia that are most pronounced in the ARH (Thaler et al., 2012; Valdearcos et al., 2014).
Microglia are the resident myeloid cells of the CNS and respond to a broad array of circulating
signals, including nutrients such as saturated fats and carbohydrates (Valdearcos et al., 2014;
Nadjar, et al., 2017; Leyroll, Laye & Nadjar, 2019; Butler et al., 2020). Moreover, activation of
microglia alone is sufficient to stimulate food intake and promote weight gain in adult mice, and
perturbations that block activation of microglia reduce the metabolic disruption associated with
neuroinflammation (Valdearcos et al., 2017; Rosin et al., 2018; Sun et al., 2023). Because of
their established role as nutrient-sensing sentinels of hypothalamic neuroinflammation, and
their documented participation in neural development (Stevens et al., 2007; Schafer et al.,
2012; Stephan et al., 2012), microglia have been proposed as possible mediators of
developmental programming caused by nutritional perturbations (Rosin and Kurrasch, 2019;
Folick et al., 2021).
141
Evidence from several of lines of investigation indicate that microglia have multiple roles in
brain development (Schwarz and Bilbo 2009; Tremblay et al., 2011; Miyamoto et al., 2016; Li
et al., 2019). Although microglia were initially thought to remain quiescent until activation by
neuroinflammation, (Nakajima & Kohsaka, 2001; Nikolic & Tan, 2005) in vivo imaging
experiments in the cerebral cortex demonstrated continual activity of their cellular processes,
which actively survey their local environment (Stowell et al., 2018), including direct contact with
axons and dendrites (Wake et al., 2009; Schafer et al., 2013). In addition to impacting neuronal
number and initial formation of neural circuits, microglia are thought to play an important role in
synaptic refinement through selective elimination of synapses, a process termed synaptic
pruning (Paolicelli et al., 2011; Schafer et al., 2012; Hong, Dissing-Olesen & Stevens, 2016).
Thus far, the majority of microglial developmental studies have focused on their role in cortical
or hippocampal circuits. However, transcriptional profiling suggests a great deal of regional
and temporal variation in microglial cell type and activity (Hammond et al., 2019; Masuda et al.,
2020; Young et al., 2021), and the effects of dietary interventions have largely focused on the
ARH. Here we used Iba1 immunostaining to visualize and measure morphological changes in
microglial morphology in the PVH of offspring exposed to MHFD and compared these effects
with those observed in the ARH and bed nuclei of the stria terminalis (BST), a limbic target of
AgRP neurons. We also used genetically targeted axonal labeling of AgRP neurons to directly
visualize cellular interactions between microglia and labeled AgRP terminals in the PVH and
ARH to determine if MHFD stimulates synaptic pruning in these regions. The results
demonstrate regionally specific changes to microglia in the PVH of MHFD offspring that are
temporally restricted to the period when AgRP neurons innervate the PVH. In addition, the
axon labeling experiments confirm a significant decrease in AgRP innervation of the PVH in
MHFD offspring, and for the first time provide direct evidence of microglial-mediated synaptic
pruning of AgRP terminals in the hypothalamus. Microglial depletion experiments determined
that the significant decrease in AgRP innervation of the PVH observed in MHFD offspring

requires normal densities of microglia, and that microglia are required for the weight gain seen
in offspring of MHFD dams at weaning. However, we did not detect a significant effect of
MHFD on the degree of synaptic pruning in the PVH, suggesting an alternative microglial
signaling mechanism yet to be identified.
172
173 **2 Results**
174
175 **2.1 Microglia exhibit morphological changes in the PVH in response to MHFD during**
176 **postnatal development.**
177
To assess the impact of MHFD on microglia in the brains of postnatal mice, we used Iba1
immunohistochemistry and confocal microscopy to visualize the distribution and morphology of
microglial cells in the PVH. Discrete regions of interest were imaged and a 3D modeling
analysis pipeline was used to measure structural changes in microglia (Figure1Ai-iii). Although
there was no apparent difference between the density of microglia in the PVH of MHFD and
NCD offspring at P16 (Figure 1J), the overall size of microglia in the PVH of MHFD offspring
was significantly greater, compared to that of NCD mice (Figure 1B, C). The enhanced size of
microglia in the MHFD offspring corresponded to an 87% increase in overall process length
and a 44% increase in branching complexity (Figure 1F, G). Additionally, the volume of
microglial cells (volume of cell body and processes) and the spatial territory they occupy was
nearly doubled in MHFD offspring (Figure 1H-I). These morphological changes appeared to be
transient because both the density and size of microglia (process length and complexity) were
reduced in postweaning mice perfused on P30 (Figure 1D-G). We also assessed the density of
AgRP terminals in the same regions of interest in the PVH and confirmed that their density is
significantly lower in the brains of MHFD offspring compared with that of NCD controls (Figure
1K), as suggested by data published previously (Vogt et al. 2014). Taken together, these
results suggest that exposure to MHFD has a profound effect on the morphology of microglia
that is consistent with enhanced activity and surveillance of their immediate microenvironment.
Furthermore, the effects of MHFD on microglial morphology in the PVH of offspring display
both temporal and regional specificity, which correspond to a decrease in the density of AgRP
inputs to the PVH.
199
200
201
202 **2.2 Microglia do not exhibit morphological changes in the ARH or BST in response to**
203 **MHFD exposure during postnatal development.**
204
The ARH houses the cell bodies of AgRP neurons, and their number is established primarily
during prenatal development (Ishii & Bouret 2012). We evaluated microglia morphology in the
ARH of postnatal mice by using the same 3D modeling pipeline shown in Figure 1. In sharp
contrast to our findings in the PVH, the number and size of microglia in the ARH were not
significantly different in NCD and MHFD offspring at P16 (Figure 2A, B; 2E, F). By P30, there
was a 67% increase in process length compared to their P16 counterparts, but no significant
difference in microglial morphology in the ARH detected between treatment groups (Figure A-
I). The volume of microglial cells, as well as the spatial territory they occupy, also increased
between P16 and P30 (Figure 2G-H), but the number of microglia visualized was reduced by

nearly half (Figure 2I). We also measured numbers of AgRP neuronal cell bodies in the ARH in
brains derived from NCD and MHFD offspring and confirmed that the number of AgRP
neurons in the ARH are also resistant to MHFD exposure (Figure 2J). In addition, we
evaluated microglia and AgRP terminals in the anterolateral part of the BST, an
extrahypothalamic target of AgRP neurons innervated during the lactational period (Bouret,
Draper and Simerly, 2004a; Cansell et al., 2012; Barbier et al., 2021). As was found for the
ARH, neither the number nor size of microglia in the BST were significantly different between
NCD and MHFD offspring at P16 or P30 (Figure 3A-D; 3E-I). Similarly, the density of AgRP
terminals in the same region of interest was not affected by MHFD exposure (Figure 3J). By
P30 the number of AgRP terminals in the BST increased by 61% compared to their P16
counterparts (Figure 3J). Taken together, these data suggest that microglia in the ARH and
BST are resistant to the increases in cellular complexity that occur in the PVH of MHFD
offspring, suggesting a notable degree of spatial specificity in the role of hypothalamic
microglia during postnatal development.
228
229
230 **2.3 Microglia are required for changes in AgRP terminal density in PVH and body weight**
231 **associated with MHFD exposure**
232
To determine if microglia are required for the observed changes in AgRP inputs to the PVH of
MHFD offspring, we depleted microglia during the lactation period. The colony-stimulating
factor-1 receptor inhibitor (CSF1R) PLX5622 was administered daily via intraperitoneal
injections between P4-P21. These postnatal treatments resulted in a significant decrease in
microglia in the PVH at the time of weaning (Figure 4A-B; Figure 4G) that appeared to be
widespread throughout the rostral forebrain. Notably, the PLX5622 treatments blocked the
reduction in AgRP fiber density observed in the medial dorsal parvicellular part of the PVH
(PVHmpd) of vehicle-treated MHFD offspring to a level that was comparable to that of NCD
offspring (Figure 4C-F; quantified in 4H). In contrast, MHFD exposure did not affect the density
of AgRP fibers in the lateral posterior magnocellular compartment of the PVH (PVHpml) and
no significant difference in AgRP fiber density was detected between MHFD offspring treated
with either PLX or vehicle (Figure 4C-F; 4J), suggesting target specificity for the microglia
mediated effects on development of AgRP inputs to the PVHmpd.
246
Depletion of microglia also appeared to protect against the increase in body weight normally
observed in MHFD mice. In keeping with previously published results (Vogt et al. 2014), the
weight of vehicle treated MHFD animals was significantly greater (24%), compared with that of
NCD animals (Figure 4I). However, MHFD mice treated with PLX5622 during lactation
exhibited significantly lower weights at P21 compared to those of vehicle-treated MHFD
animals. Taken together, these findings suggest that microglia mediate target-specific effects
of MHFD exposure on the innervation of the PVH by AgRP neurons, and that microglia play a
role in mediating the effects of MHFD on body weight.
255
256 **2.4 Engulfment of AgRP terminals by microglia in the PVH and the ARH**
257
Microglia are thought to impact development of neuronal connections through an active
engulfment mechanism and the lysosomal associated membrane protein CD68 has

been implicated in this process. Here, we used immunohistochemistry to visualize the
presence of in Iba1-labeled microglia in mice with genetically targeted labeling of AgRP
terminals. Many apparent contacts between microglial processes and AgRP terminals in
the PVH and ARH were observed at P16 and P30, including internalized AgRP
terminals (Figure 5A-L). However, the extent of internalization did not appear to be
influenced by MHFD exposure; there were no significant differences between
internalized AgRP terminals in MHFD and NCD offspring at P16, in either the PVH or
ARH (Figure 5M, O). Similarly, we did not detect a statistically significant difference in
microglial CD68 levels in the PVH between diet groups at P30 (Figure 5N). Consistent
with previous reports, (Wong et al. 2005; Hart et al. 2012) the density of CD68 labeled
profiles nearly doubled in the PVH between P16 and P30 (Figure 5N), as microglia
become more phagocytic with age. CD68 staining also increased in the ARH between
P16 and P30 (Figure 5P), supporting the notion that microglia increase their phagocytic
capacity with age. Nevertheless, our analysis demonstrates that microglia interact
directly with AgRP terminals, with clear evidence of engulfment. MHFD exposure does
not appear to promote microglia-mediated engulfment, at least not in the specific PVH
and ARH domains examined.
277
278 **3 Discussion**
279
It is well established that microglia are responsive to HFD exposure in adult rodents (Thaler et
al., 2012; Morari et al., 2014; Valdearcos et al., 2014; Baufeld et al., 2016; Valdearcos et al.,
2017) and multiple lines of evidence support an important role for microglia in mediating key
aspects of neural circuit development (Checchin et al., 2006; Hoshiko et al. 2012; Li et al.,
2012; Hagemeyer et al., 2017). Here we demonstrate that microglia are required for significant
elevations in body weight that emerge from postnatal exposure to HFD and are associated
with a sustained decrease in the density of afferents from AgRP neurons to the PVH in
offspring. Exposure to MHFD caused distinct morphological changes to microglia that are
consistent with enhanced activity, which were observed in the PVH, but not the ARH or BST.
Moreover, the morphological changes to microglia observed appear to be primarily limited to
the critical period for development of AgRP inputs to PVH neurons in MHFD offspring.
Although our results demonstrate that microglia engage in engulfment of AgRP terminals in the
PVH during development, synaptic pruning by microglia does not appear to represent the
cellular mechanism mediating the effects of MHFD exposure on innervation of the PVH by
AgRP neurons.
295
296 **3.1 A Role for Microglia in Mediating Body Weight Changes Observed in MHFD**
297 **Offspring**
298
Maternal high fat diet exposure during a period that extends across both gestation and
lactation causes offspring to have increased body weight, fat content and susceptibility to diet-
induced obesity (Samuelsson et al., 2008; Masuyama et al., 2014; Gomes et al., 2018). In
contrast to the more variable impact of maternal HFD exposure just during gestation, limiting
HFD exposure to the postnatal period, which corresponds to lactation, when mice derive their
nutrition primarily from milk, reliably predisposes adult mice to obesity. Feeding pregnant dams
HFD during the week prior to parturition leads to either increased body weight in offspring

(Akhaphong et al., 2021) or no significant effect (Sun et al, 2012; Vogt et al., 2014), suggesting
that MHFD plays a dominant role in specifying metabolic phenotype later in life (Sun et al.,
2012; Vogt et al. 2014). In the present study, global depletion of microglia with the CSF1R
inhibitor PLX5622 blocked the ability of MHFD to increase body weight in mice at P21,
indicating that microglia may mediate metabolic changes caused by MHFD exposure that are
apparent as early as weaning. A previous study that used intragastric administration of
PLX3397 to neonatal mice reported enhanced food intake in the offspring during lactation (Sun
et al., 2023). Notably, these mice were not derived from MHFD dams. Microglial depletion with
PLX5622 in adult mice was found to mitigate the effects of HFD exposure (Rosin et al., 2018)
and our results indicate that microglia may function similarly in offspring during postnatal life to
effect changes in body weight, even if the maternal HFD exposure is restricted to the lactation
period. Further studies are required to define the long-term metabolic profile resulting from
developmental microglial manipulations. However, given the abundant literature on sustained
impact of MHFD on metabolic phenotype, enduring disruptions are likely. It should be noted
that PLX5622 treatment is not spatially limited to the PVH or ARH, leaving open the possibility
that the effects of microglial depletion on body weight occur outside of these nuclei, or are due
to collective activation of microglia in multiple components of feeding circuitry (Green, Crasper
and Hohsfield, 2020). Localization of the specific site of action for microglial specification of
mature body weight during development will require utilization of specific markers for
hypothalamic microglia that account for regional and phenotypic heterogeneity, perhaps
through intersectional genetic methods and combinatorial pharmacology (Hammond et al.,
2019; Kim et al., 2021).
328
329 **3.2 MHFD Induces Spatially Limited Changes in Microglial Morphology**
330
Morphological changes in microglia have been reported in response to a variety of
environmental exposures. In the hypothalamus, adult mice fed a high fat diet show both
proliferation and changes in process length and complexity (Thaler et al., 2012; Valdearcos et
al, 2017). In our studies, MHFD exposure caused a marked increase in the overall size of
microglia in the PVH related to increases in both the length and branching complexity of Iba-1-
stained cellular processes. This increase in the territory occupied by microglia in the PVH was
not accompanied by an increase in microglial number, nor were numbers of microglia affected
in the PVH by MHFD exposure.   However, in contrast to the PVH, changes in microglial
morphology were not observed in the ARH in response to MHFD, although we did observe an
overall increase in process length in the ARH between P16 and P30 of both MHFD and NCD
mice. This finding is consistent with previously published reports on microglial maturation (Sun
et al., 2023). The spatially restricted enhancement of microglial activation in the PVH resulting
from MHFD exposure appears to contribute to an expansion of parenchymal territory surveiled
by PVH microglia, as reflected in the volume measurements accomplished with geometrical
modeling of process length and complexity. This interpretation is supported by *in vitro* and *in*
vivo observations of enhanced process extension and increased neuronal interactions
resulting from inflammatory activation of microglia (Wake et al., 2009; Schafer et al., 2013;
Dissing-Olesen et al., 2014; Stowell et al., 2018; Bolton et al., 2022).
349
350 As in the ARH, we did not observe comparable changes in microglial morphology in the BST,
351 an extrahypothalamic target of AgRP neurons innervated during the lactation period (Cansell

et al., 2012; Barbier et al., 2021). These observations underscore the remarkable molecular heterogeneity of microglial phenotypes that appear to occupy various hypothalamic niches during development, and which may have equally diverse developmental roles and responses to environmental signals (Bilbo and Schwarz, 2012; Frost and Schafer, 2016; Li and Barres, 2018; Ngozi and Bolton, 2022). Moreover, the observed morphological changes in the PVH caused by MHFD appear to be transient as there no significant differences in microglial processes by P30, and the density of microglial cells in the PVH was significantly reduced in the older mice, suggesting a decline in overall activity. Temporal alignment of these morphological events with the critical period for development of AgRP projections to the PVH suggests a possible role for microglia linking nutrition with specification of axonal targeting (Kamitakahara et al., 2018).

3.3 Microglia Mediate Impaired Innervation of the PVH by AgRP Neurons

AgRP neuronal projections develop primarily during the first two weeks of life, which corresponds to a critical period for the neurotrophic action of leptin on axonal outgrowth and targeting of AgRP inputs to distinct components of the PVH (Simerly, 2008; Elson and Simerly, 2015). Exposure to MHFD during this critical developmental window permanently impairs AgRP projections to the PVH, DMH and LH (Vogt et al., 2014) that are associated with increased body weight in adulthood. Here we confirm that MHFD impairs innervation of the PVH, and this defect was apparent as early as P16. The PVH is innervated by AgRP neurons between P8 and P10 (Bouret, Draper and Simerly, 2004a). We found that depletion of microglia with PLX5622 between P4 and P21, a period that aligns not only with AgRP innervation of the PVH but also with maximum changes in microglial morphology, blocked the effects of MHFD on AgRP terminals in the PVH. However, this partial rescue of innervation appeared to be limited to the PVHmpd, suggesting that there is a profound regional specialization in the activity of microglia in the PVH. We did not observe a change in AgRP innervation of the BST in MHFD offspring that mirrored the changes seen in the PVH of the same animals, further supporting the conclusion that microglia display significant spatial heterogeneity in mediating site-specific alterations in AgRP axon targeting.

Microglia are known to impact a variety of developmental events, including alterations in cell number through programmed cell death or neurogenesis, as well as axonal targeting and remodeling of neural circuits (Frost and Schafer, 2016; Li and Barres 2018). It is unlikely that the impaired innervation of the PVH observed in MHFD offspring is due to a reduction in the number of AgRP neurons in the ARH (Vogt et al, 2014; Valdearcos et al., 2017). Sun and colleagues reported that microglial depletion during postnatal life actually increases numbers of AgRP neurons, as well as enhances local densities of AgRP fibers in the ARH, possibly through enhanced formation of perineuronal nets (Sun et al., 2023). However, changes in AgRP targets outside of the ARH were not assessed. Depletion of microglia during gestation causes a significant decrease in the number of proopiomelanocortin (POMC) neurons in the ARH and leads to an acceleration of weight gain (Rosin et al., 2018), consistent with neurogenesis of ARH neurons occurring in mid gestation and being susceptible to nutritional impacts during embryonic life (Ishii and Bouret, 2012; Elson and Simerly, 2015). Interestingly, genetic deletion of leptin receptors from myeloid cells reduced numbers of POMC neurons in the ARH, suppressed POMC innervation of the PVH, and decreased microglial process

complexity (Gao et al., 2017). Taken together, these results suggest that leptin signaling in
microglia may act at the level of the ARH to promote growth of AgRP projections to the PVH,
while MHFD activates microglia in the PVH to specify patterns of AgRP afferents that are not
only regionally specific, but also target discrete domains of the PVH. However, whether
microglia inhibit synaptogenesis or are involved in synaptic refinement through an alternative
regressive mechanism will require further investigation.
404
405 **3.4 Microglia in PVH Participate in Synaptic Pruning**
406
Microglia play an important role in synaptic pruning, a process whereby synapses that form
early in development are eliminated as others are strengthened and maintained (Katz and
Shatz, 1996; Sanes and Lichtman, 1999; Frost and Schafer 2016; Li and Barres, 2018).
Although this process has been studied most extensively in somatosensory cortex (Miyamoto
et al., 2016), hippocampus (Paolicelli et al., 2011; Wang et al. 2020) and the visual system
(Tremblay et al., 2010; Schafer et al., 2012), there is clear evidence for involvement of
microglia in synaptic pruning of immunolabeled glutamatergic terminals associated with
corticotropin releasing hormone neurons in the PVH (Bolton et al., 2022). In the present study,
we used AgRP axonal labeling to provide evidence that microglia in the PVH participate in
synaptic pruning of AgRP/GABAergic synapses during the critical period for PVH innervation,
and when PVH microglia exhibit high levels of process extension. The lysosomal marker CD68
was colocalized with internalized AgRP terminals in PVH microglia, and although elevated at
P30, there were no differences between offspring of MHFD and NCD dams. Additionally, we
did not find a significant difference between the density of engulfed AgRP terminals in the PVH
of MHFD offspring at either P16 or P30. However, enhanced engulfment of AgRP terminals in
MHFD offspring may occur at a later point in development. It is also possible that the
synaptophysin-tdTomato axonal label may have been lost from pruned synapses during
engulfment. A more probable interpretation is that PVH microglia participate in synaptic
refinement through other cellular mechanisms, including microglial release of secreted factors
such as IL6 (Kim, Copperi, and Diano, 2024) or microglial derived BDNF (Parkhurst et al.,
2013). Additional investigation is required to resolve this question.
428
429 **4 Conclusions**
430
MHFD causes elevated levels of saturated carbohydrates and fats in milk (Gorski et al., 2006;
Vogt et al., 2014; Calvo-Lerma et al., 2022). The resulting overnutrition resulting from exposure
to this enhanced diet is thought to underly the propensity towards obesity observed in offspring
later in life (Skowronski, Leibel and Leduc, 2023). Although microglia are likely mediators of
multiple neurobiological events influencing how hypothalamic circuits function during regulation
of energy balance, the precise signaling mechanisms remain ill defined. There may be
common molecular mechanisms underlying the effects of HFD exposure on microglial
activation in adults and those occurring during postnatal development, but how these signaling
events exert a lasting impact on the organization and function of feeding circuitry has not been
defined. The results presented here clearly demonstrate an important role for microglia,
specifically in the PVH, on sculpting the density of AgRP inputs to the PVH that is not only
spatially restricted, but also aligned temporally with synaptogenesis in the PVH. Furthermore,
PVH microglia clearly interact directly with AgRP afferent axons during this critical period and

appear to be refined through engulfment by microglia. However, synaptic pruning does not
appear to be sufficient to affect the significant reduction in AgRP innervation of PVH neurons
observed following MHFD exposure, suggesting involvement of additional microglial signaling
mechanisms that are not only important for specifying patterns of innervation of the PVH by
AgRP neurons, but may also contribute more broadly to developmental programming of
metabolic phenotype.
450
451 **5 Materials and Methods**
452
453 **5.1 Key Resources Table**

Reagent type or resource	Designation	Source or reference	Identifiers	Additional information
Genetic reagent (*M. musculus*)	*AgRP*-IRES-Cre	Jackson Laboratory	Stock #: 012899 RRID: IMSR_JAX:012899	MGI ID: J:140858
Genetic reagent (*M. musculus*)	Synaptophysin-tdTomato	Jackson Laboratory	Stock #: 012570 RRID: IMSR_JAX:012570	MGI ID: J:170755
Antibody	Rabbit polyclonal anti-Iba1	FUJIFILM Wako Shibayagi	Cat. #: 019-19741 RRID: AB_ 839504	IHC (1:2000)
Antibody	Rat monoclonal anti-CD68 [FA-11]	Abcam	Cat. #: ab53444 RRID: AB_869007	IHC (1:500)
Antibody	Donkey polyclonal anti-rabbit Alexa Fluor 488	ThermoFisher Scientific	Cat. #: A32790 RRID: AB_2762833	IHC (1:500)
Antibody	Donkey polyclonal anti-rat Alexa Fluor 647	ThermoFisher Scientific	Cat. #: A48272 AB_2893138	IHC (1:500)
Pharmacological Inhibitor	PLX5622 hemifumarate, CSF1R inhibitor	MedChemExpress	Cat. # HY114153A	
Software, Algorithm	Imaris	Bitplane	V9.5	
Software, Algorithm	GraphPad Prism	Prism	Prism 7	

454
455
5.2 Animal Care
Transgenic mice expressing IRES-Cre recombinase under the control of the AgRP promoter,
AgRP-Cre mice (AgRP-IRES-Cre Stock no: 012899; MGI:J:140858) and mice expressing the
cre-dependent fluorescent reporter synaptophysin-tdTomato (SynTom mice; Ai34D-Rosa-
CAG-LSL-Synaptophysin-tdTomato-WPRE; stock number: 012570; MGI:J:170755) were
obtained from the Jackson Laboratory (Bar Harbor, ME) and maintained in our colony at

Vanderbilt University. Wild-type C57BL/6J mice (stock number: 000664) were also obtained
from The Jackson Laboratory. To visualize AgRP inputs, AgRP-Cre mice were crossed with
SynTom mice to generate AgRP-Cre::SynTom mice, as described previously (Biddinger, et.al.,
2020).
466
467 All animal care and experimental procedures were performed in accordance with the
468 guidelines of the National Institutes of Health and the Institutional Care and Use Committee of
469 Vanderbilt University. Mice were housed at 22°C on a 12:12 h light:dark cycle (lights on at 6:00
470 a.m.: lights off at 6:00 p.m.). Mice were provided ad libitum access to a standard chow diet
471 (PicoLab Rodent Diet 20 #5053), unless otherwise specified.
472
To generate offspring of dams exposed to HFD during lactation (MHFD) mice had ad libitum
access to normal chow (PicoLab Rodent Diet, 5053LabDiet: 25% protein; 62% carbohydrates;
13% fat; 4 kcal/g energy density) during mating. On the first postnatal day (P1) all litters were
adjusted to 7 pups to normalize nutrition and dams were switched to either high fat diet (HFD)
(Research Diets D12451: 20% protein; 35% carbohydrate; 45% fat; 4.7 kcal/g energy density)
or kept on normal chow diet (NCD). The dams remained on either HFD or NCD through
lactation, and at weaning the offspring were switched to a normal chow diet, regardless of
lactation diet condition.
481
5.3 PLX5622 Microglia Depletion
To reduce microglia during postnatal development (P4-P21) mice were treated with PLX5622,
a colony stimulating factor 1 receptor (CSF1R) inhibitor or with DMSO vehicle via
intraperitoneal injection (Riquier et al. 2020). Briefly, PLX5622 hemifurate solid (Cat.
#HY114153A MedChemExpress, Monmouth Junction, NJ, USA) was suspended in dimethyl
sulfoxide at a concentration of 172 mg/ml. The injection working solution was prepared to
include 20% Kolliphor RH40 diluted in PBS, which resulted in doses with a 6.5mg/ml PLX5622
concentration and injection concentration of 15 mg/kg. IP injections were given once daily from
P4 to P21 and injection dose was determined by each animal's body weight at time of
injection.
492
5.4 Immunohistochemistry
Mice were perfused at P16 and P30 and processed for immunofluorescence by using an
antibody to Iba-1 (1:2000; FUJIFILM Wako Pure Chemicals, Osaka, Japan) to visualize
microglia and an antibody to CD68 (1:500; Abcam, Cambridge, MA, USA) to assess
phagocytic capacity of the microglia. Mice were first anesthetized with tribromoethanol (TBE)
and then perfused transcardially with cold 0.9% saline, followed by cold fixative (4%
paraformaldehyde in borate buffer, pH 9.5) for 10 minutes. Brains were then removed from
skull and postfixed in the same fixative overnight. Brains were then cryoprotected overnight in
a 20% sucrose solution before being frozen in optimal temperature cutting compound (Sakura
Finetek Inc., Torrance, CA) and sectioned on a freezing stage sliding microtome at 30μm.
Free-floating brain sections were rinsed in KPBS and then incubated in blocking buffer
containing 2% normal donkey serum and 0.3% Triton-X overnight at 4°C. Sections were
transferred to primary antibody incubation for 48hr with rabbit anti-Iba-1 antibody and rat anti-
CD68 antibody. Following primary antibody incubation, sections were rinsed several times in
KPBS, incubated for an hour at room temperature in blocking buffer containing secondary

antibodies against rabbit and rat (raised in donkey) conjugated with Alexa-Fluor fluorochromes
(1:500; Life Technologies, Carlsbad, CA, USA), mounted onto charged microscope slides and
coverslipped using ProLong antifade mounting medium (Life Technologies, Carlsbad, CA,
USA).
512
5.5 Image Acquisition and Analysis
Sections through the PVH, ARH, and BST were examined on a laser scanning confocal
microscope (Zeiss 800) and cytoarchitectonic features of the nuclei were visualized with
Hoescht 33342, and used to define matching regions of interest (ROI) for quantitative analysis.
Confocal image stacks were collected for each ROI through the entire thickness of the region
at a frequency of $0.1\mu m$ using the 40x objective. Imaris visualization software (Bitplane V9.5,
Salisbury Cove, ME, USA) was used to create 3D reconstructions of each multichannel set of
images.
521
3D reconstructions of microglia based on Iba-1 immunofluorescence were made using made
using Imaris (Bitplane, v9.5). Profiles were skeletonized automatically by using the Filaments
tool in order to quantify changes in microglia structure. Sholl analysis (Geoffry et al., 2014),
was performed on the skeletonized structures to determine complexity of branching of
microglial processes. Briefly, 3D concentric spheres are drawn around the microglia skeleton
and each contact point between microglial process and sphere is counted as 1 level of
branching. The higher the number of contacts, the higher complexity of branching in the
microglia. 3D reconstructions of the microglia were also evaluated for length of processes as
well as cell volume and the area that is occupied. To evaluate the 3D space occupied by each
microglial cell, a polyhedron was drawn around the microglia by using the built-in Convex Hull
function under Filaments, which is used to calculate volume of the polyhedron automatically,
accounting for the 3D space around the microglia.
534
In order to assess cellular interactions between AgRP terminals and microglia, the 3D
renderings were used to determine level of contact by using a MATLAB script to automate the
analysis. Both channels were subjected to background subtraction and Gaussian filtering. The
automatic threshold calculated was based on k-means statistical methods and was used in the
majority of analyses. The AgRP terminals were reconstructed as "spots" of 0.8 mm diameter
(corresponding to the largest measured size) and their total number was automatically
calculated. Briefly, the automatic detection algorithm applies a 3D Mexican hat filter using the
spot size and then locates the spot centroid at the local maxima of the filtered image. The
number of spots located at no more than $1\mu m$ from the microglia surface was automatically
determined and indicated as a contact point. Next, spots that were determined to be less than
$0.5\mu m$ away from internal microglial surfaces were determined to be internalized by the
microglia and counted as engulfed. CD68 levels were determined using automated analysis to
create 3D renderings and their volume computed.
548
549 **5.6 Statistical Analyses**
550 Data are presented as the mean values ± SEM. Descriptive statistics and unpaired t-tests were
551 used to compute group differences using GraphPad Prism software (GraphPad Software, San
552 Diego, CA). P-value less than 0 .05 were considered significant.
553

Acknowledgments

We thank the members of the Simerly Lab for comments and discussion on early versions of this manuscript. This work was supported by NIH grants R01DK106476 (RBS) and T32DK07563 (HNM-R).

Author Contributions

HNM-R and RBS conceived of this research and designed experiments. HNM-R conducted experiments with technical support from JEB and MNB. HNM-R and JEB analyzed data and prepared figures. HNM-R and RBS wrote the manuscript and all authors participated in editing the manuscript.

References

Akhaphong, B., Gregg, B., Kumusoglu, D., Jo, S., Singer, K., Scheys, J., DelProposto, J., Lumeng, C., Bernal-Mizrachi, E., & Alejandro, E. U. (2022). Maternal High-Fat Diet During Pre-Conception and Gestation Predisposes Adult Female Offspring to Metabolic Dysfunction in Mice. *Frontiers in Endocrinology*, *12*, 780300. https://doi.org/10.3389/fendo.2021.780300

Andersen, L. G., Holst, C., Michaelsen, K. F., Baker, J. L., & Sørensen, T. I. A. (2012). Weight and weight gain during early infancy predict childhood obesity: a case-cohort study. *International Journal of Obesity*, *36*(10), 1306–1311. https://doi.org/10.1038/ijo.2012.134

Barbier, M., González, J. A., Houdayer, C., Burdakov, D., Risold, P., & Croizier, S. (2021). Projections from the dorsomedial division of the bed nucleus of the stria terminalis to hypothalamic nuclei in the mouse. *Journal of Comparative Neurology*, *529*(5), 929–956. https://doi.org/10.1002/cne.24988

Baufeld, C., Osterloh, A., Prokop, S., Miller, K. R., & Heppner, F. L. (2016). High-fat diet-induced brain region-specific phenotypic spectrum of CNS resident microglia. *Acta Neuropathologica*, *132*(3), 361–375. https://doi.org/10.1007/s00401-016-1595-4

Betley, J. N., Xu, S., Cao, Z. F. H., Gong, R., Magnus, C. J., Yu, Y., & Sternson, S. M. (2015). Neurons for hunger and thirst transmit a negative-valence teaching signal. *Nature, 521*(7551), 180-185. https://doi.org/10.1038/nature14416

Biddinger, J. E., Lazarenko, R. M., Scott, M. M., & Simerly, R. (2020). Leptin suppresses development of GLP-1 inputs to the paraventricular nucleus of the hypothalamus. *ELife, 9*, e59857. https://doi.org/10.7554/elife.59857

Bilbo, S. D., & Schwarz, J. M. (2009). Early-Life Programming of Later-Life Brain and Behavior: A Critical Role for the Immune System. *Frontiers in Behavioral Neuroscience*, *3*, 14. https://doi.org/10.3389/neuro.08.014.2009

Bilbo, S. D., & Schwarz, J. M. (2012). The immune system and developmental programming of brain and behavior. *Frontiers in Neuroendocrinology*, *33*(3), 267–286. https://doi.org/10.1016/j.yfrne.2012.08.006

Bolton, J. L., Short, A. K., Othy, S., Kooiker, C. L., Shao, M., Gunn, B. G., Beck, J., Bai, X., Law, S. M., Savage, J. C., Lambert, J. J., Belelli, D., Tremblay, M.-È., Cahalan, M. D., & Baram, T. Z. (2022). Early stress-induced impaired microglial pruning of excitatory synapses on immature CRH-expressing neurons provokes aberrant adult stress responses. *Cell Reports*, *38*(13), 110600. https://doi.org/10.1016/j.celrep.2022.110600

Bouret, S. G., Draper, S. J., & Simerly, R. B. (2004a). Formation of Projection Pathways from the Arcuate Nucleus of the Hypothalamus to Hypothalamic Regions Implicated in the Neural Control of Feeding Behavior in Mice. *The Journal of Neuroscience*, *24*(11), 2797–2805. https://doi.org/10.1523/jneurosci.5369-03.2004

Bouret, S. G., Draper, S. J., & Simerly, R. B. (2004b). Trophic Action of Leptin on Hypothalamic Neurons That Regulate Feeding. *Science*, *304*(5667), 108–110. https://doi.org/10.1126/science.1095004

Bouret, S., Levin, B. E., & Ozanne, S. E. (2015). Gene-Environment Interactions Controlling Energy and Glucose Homeostasis and the Developmental Origins of Obesity. *Physiological Reviews*, *95*(1), 47–82. https://doi.org/10.1152/physrev.00007.2014

Bouyer, K., & Simerly, R. B. (2013). Neonatal Leptin Exposure Specifies Innervation of Presympathetic Hypothalamic Neurons and Improves the Metabolic Status of Leptin-Deficient Mice. *The Journal of Neuroscience*, *33*(2), 840–851. https://doi.org/10.1523/jneurosci.3215-12.2013

Butler, M. J., Cole, R. M., Deems, N. P., Belury, M. A., & Barrientos, R. M. (2020). Fatty food, fatty acids, and microglial priming in the adult and aged hippocampus and amygdala. *Brain, Behavior, and Immunity*, *89*, 145–158. https://doi.org/10.1016/j.bbi.2020.06.010

Calvo-Lerma, J., Selma-Royo, M., Hervas, D., Yang, B., Intonen, L., González, S., Martínez-Costa, C., Linderborg, K. M., & Collado, M. C. (2022). Breast Milk Lipidome Is Associated With Maternal Diet and Infants' Growth. *Frontiers in Nutrition*, *9*, 854786. https://doi.org/10.3389/fnut.2022.854786

Cansell, C., Denis, R. G. P., Joly-Amado, A., Castel, J., & Luquet, S. (2012). Arcuate AgRP neurons and the regulation of energy balance. *Frontiers in Endocrinology*, *3*, 169. https://doi.org/10.3389/fendo.2012.00169

Cansell, C., Stobbe, K., Sanchez, C., Thuc, O. L., Mosser, C., Ben-Fradj, S., Leredde, J., Lebeaupin, C., Debayle, D., Fleuriot, L., Brau, F., Devaux, N., Benani, A., Audinat, E., Blondeau, N., Nahon, J., & Rovère, C. (2021). Dietary fat exacerbates postprandial hypothalamic inflammation involving glial fibrillary acidic protein-positive cells and microglia in male mice. *Glia*, *69*(1), 42–60. https://doi.org/10.1002/glia.23882

Checchin, D., Sennlaub, F., Levavasseur, E., Leduc, M., & Chemtob, S. (2006). Potential Role of Microglia in Retinal Blood Vessel Formation. *Investigative Opthalmology & Visual Science*, *47*(8), 3595. https://doi.org/10.1167/iovs.05-1522

Chen, Y., Lin, Y. C., Kuo, T. W., & Knight, Z. A. (2015). Sensory detection of food rapidly modulates arcuate feeding circuits. *Cell*, *160*(5), 829-841. https://doi.org/10.1016/j.cell.2015.01.033

Collaboration, P. S., Whitlock, G., Lewington, S., Sherliker, P., Clarke, R., Emberson, J., Halsey, J., Qizilbash, N., Collins, R., & Peto, R. (2009). Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet*, *373*(9669), 1083–1096. https://doi.org/10.1016/s0140-6736(09)60318-4

Dissing-Olesen, L., LeDue, J. M., Rungta, R. L., Hefendehl, J. K., Choi, H. B., & MacVicar, B. A. (2014). Activation of Neuronal NMDA Receptors Triggers Transient ATP-Mediated Microglial Process Outgrowth. *The Journal of Neuroscience*, *34*(32), 10511–10527. https://doi.org/10.1523/jneurosci.0405-14.2014

Elson, A. E., & Simerly, R. B. (2015). Developmental specification of metabolic circuitry. *Frontiers in Neuroendocrinology*, *39*, 38–51. https://doi.org/10.1016/j.yfrne.2015.09.003

Folick, A., Koliwad, S. K., & Valdearcos, M. (2021). Microglial Lipid Biology in the Hypothalamic Regulation of Metabolic Homeostasis. *Frontiers in Endocrinology*, *12*, 668396. https://doi.org/10.3389/fendo.2021.668396

Frost, J. L., & Schafer, D. P. (2016). Microglia: Architects of the Developing Nervous System. *Trends in Cell Biology*, *26*(8), 587–597. https://doi.org/10.1016/j.tcb.2016.02.006

Gao, Y., Vidal-Itriago, A., Milanova, I., Korpel, N. L., Kalsbeek, M. J., Tom, R. Z., Kalsbeek, A., Hofmann, S. M., & Yi, C.-X. (2017). Deficiency of leptin receptor in myeloid cells disrupts hypothalamic metabolic circuits and causes body weight increase. *Molecular Metabolism*, *7*, 155–160. https://doi.org/10.1016/j.molmet.2017.11.003

García-Cáceres, C., Balland, E., Prevot, V., Luquet, S., Woods, S. C., Koch, M., Horvath, T. L., Yi, C.-X., Chowen, J. A., Verkhratsky, A., Araque, A., Bechmann, I., & Tschöp, M. H. (2019). Role of astrocytes, microglia, and tanycytes in brain control of systemic metabolism. *Nature Neuroscience*, *22*(1), 7–14. https://doi.org/10.1038/s41593-018-0286-y

Geoffry, Derecki, N., & Kipnis, J. (2014). Microglial Scholl Analysis. *Protocol Exchange*.

Gomes, D., Kries, R. von, Delius, M., Mansmann, U., Nast, M., Stubert, M., Langhammer, L., Haas, N. A., Netz, H., Obermeier, V., Kuhle, S., Holdt, L. M., Teupser, D., Hasbargen, U., Roscher, A. A., & Ensenauer, R. (2018). Late-pregnancy dysglycemia in obese pregnancies after negative testing for gestational diabetes and risk of future childhood overweight: An interim analysis from a longitudinal mother–child cohort study. *PLoS Medicine*, *15*(10), e1002681. https://doi.org/10.1371/journal.pmed.1002681

Gorski, J. N., Dunn-Meynell, A. A., Hartman, T. G., & Levin, B. E. (2006). Postnatal environment overrides genetic and prenatal factors influencing offspring obesity and insulin resistance. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, *291*(3), R768–R778. https://doi.org/10.1152/ajpregu.00138.2006

Green, K. N., Crapser, J. D., & Hohsfield, L. A. (2020). To Kill a Microglia: A Case for CSF1R Inhibitors. *Trends in Immunology*, *41*(9), 771–784. https://doi.org/10.1016/j.it.2020.07.001

Hagemeyer, N., Hanft, K.-M., Akriditou, M.-A., Unger, N., Park, E. S., Stanley, E. R., Staszewski, O., Dimou, L., & Prinz, M. (2017). Microglia contribute to normal myelinogenesis and to oligodendrocyte progenitor maintenance during adulthood. *Acta Neuropathologica*, *134*(3), 441–458. https://doi.org/10.1007/s00401-017-1747-1

Hammond, T. R., Dufort, C., Dissing-Olesen, L., Giera, S., Young, A., Wysoker, A., Walker, A. J., Gergits, F., Segel, M., Nemesh, J., Marsh, S. E., Saunders, A., Macosko, E., Ginhoux, F., Chen, J., Franklin, R. J. M., Piao, X., McCarroll, S. A., & Stevens, B. (2019). Single-Cell RNA Sequencing of Microglia throughout the Mouse Lifespan and in the Injured Brain Reveals Complex Cell-State Changes. *Immunity*, *50*(1), 253-271.e6. https://doi.org/10.1016/j.immuni.2018.11.004

Hart, A. D., Wyttenbach, A., Perry, V. H., & Teeling, J. L. (2012). Age related changes in microglial phenotype vary between CNS regions: Grey versus white matter differences. *Brain, Behavior, and Immunity*, *26*(5), 754–765. https://doi.org/10.1016/j.bbi.2011.11.006

Hong, S., Dissing-Olesen, L., & Stevens, B. (2016). New insights on the role of microglia in synaptic pruning in health and disease. *Current Opinion in Neurobiology*, *36*, 128–134. https://doi.org/10.1016/j.conb.2015.12.004

Horvath, T. L., Sarman, B., García-Cáceres, C., Enriori, P. J., Sotonyi, P., Shanabrough, M., Borok, E., Argente, J., Chowen, J. A., Perez-Tilve, D., Pfluger, P. T., Brönneke, H. S., Levin, B. E., Diano, S., Cowley, M. A., & Tschöp, M. H. (2010). Synaptic input organization of the melanocortin system predicts diet-induced hypothalamic reactive gliosis and obesity. *Proceedings of the National Academy of Sciences*, *107*(33), 14875–14880. https://doi.org/10.1073/pnas.1004282107

Hoshiko, M., Arnoux, I., Avignone, E., Yamamoto, N., & Audinat, E. (2012). Deficiency of the Microglial Receptor CX3CR1 Impairs Postnatal Functional Development of Thalamocortical Synapses in the Barrel Cortex. *The Journal of Neuroscience*, *32*(43), 15106–15111. https://doi.org/10.1523/jneurosci.1167-12.2012

Ishii, Y., & Bouret, S. G. (2012). Embryonic Birthdate of Hypothalamic Leptin-Activated Neurons in Mice. *Endocrinology*, *153*(8), 3657–3667. https://doi.org/10.1210/en.2012-1328

Kamitakahara, A., Bouyer, K., Wang, C., & Simerly, R. (2018). A critical period for the trophic actions of leptin on AgRP neurons in the arcuate nucleus of the hypothalamus. *Journal of Comparative Neurology*, *526*(1), 133–145. https://doi.org/10.1002/cne.24327

Katz, L. C., & Shatz, C. J. (1996). Synaptic Activity and the Construction of Cortical Circuits.
Science, *274*(5290), 1133–1138. https://doi.org/10.1126/science.274.5290.1133

Kim, J. D., Copperi, F., & Diano, S. (2024). Microglia in Central Control of Metabolism.
Physiology, *39*(1), 5–17. https://doi.org/10.1152/physiol.00021.2023

Kim, J.-S, Kolesnikov, M., Peled-Hajaj, S., Scheyltjens, I., Xia, Y., Trzebanski, S., Haimon, Z.,
Shemer, A., Lubart, A., Hove, H. V., Chappell-Maor, L., Boura-Halfon, S., Movahedi, K.,
Blinder, P., & Jung, S. (2021). A Binary Cre Transgenic Approach Dissects Microglia and
CNS Border-Associated Macrophages. *Immunity*, *54*(1), 176-190.e7.
https://doi.org/10.1016/j.immuni.2020.11.007

Krashes, M. J., Koda, S., Ye, C., Rogan, S. C., Adams, A. C., Cusher, D. S., ... & Lowell, B. B.
(2011). Rapid, reversible activation of AgRP neurons drives feeding behavior in mice. The
Journal of clinical investigation, 121(4), 1424-1428. https:// www.jci.org/articles/view/46229

Leyrolle, Q., Layé, S., & Nadjar, A. (2019). Direct and indirect effects of lipids on microglia
function. *Neuroscience Letters*, *708*, 134348. https://doi.org/10.1016/j.neulet.2019.134348

Li, Q., & Barres, B. A. (2018). Microglia and macrophages in brain homeostasis and disease.
Nature Reviews Immunology, *18*(4), 225–242. https://doi.org/10.1038/nri.2017.125

Li, Q., Cheng, Z., Zhou, L., Darmanis, S., Neff, N. F., Okamoto, J., Gulati, G., Bennett, M. L.,
Sun, L. O., Clarke, L. E., Marschallinger, J., Yu, G., Quake, S. R., Wyss-Coray, T., &
Barres, B. A. (2019). Developmental Heterogeneity of Microglia and Brain Myeloid Cells
Revealed by Deep Single-Cell RNA Sequencing. *Neuron*, *101*(2), 207-223.e10.
https://doi.org/10.1016/j.neuron.2018.12.006

Li, Y., Du, X., Liu, C., Wen, Z., & Du, J. (2012). Reciprocal Regulation between Resting
Microglial Dynamics and Neuronal Activity In Vivo. *Developmental Cell*, *23*(6), 1189–1202.
https://doi.org/10.1016/j.devcel.2012.10.027

Lippert, R. N., & Brüning, J. C. (2022). Maternal Metabolic Programming of the Developing
Central Nervous System: Unified Pathways to Metabolic and Psychiatric Disorders.
Biological Psychiatry, *91*(10), 898–906. https://doi.org/10.1016/j.biopsych.2021.06.002

Masuda, T., Sankowski, R., Staszewski, O., & Prinz, M. (2020). Microglia Heterogeneity in the
Single-Cell Era. *Cell Reports*, *30*(5), 1271–1281.
https://doi.org/10.1016/j.celrep.2020.01.010

Masuyama, H., & Hiramatsu, Y. (2014). Additive Effects of Maternal High Fat Diet during
Lactation on Mouse Offspring. *PLoS ONE*, *9*(3), e92805.
https://doi.org/10.1371/journal.pone.0092805

Miyamoto, A., Wake, H., Ishikawa, A. W., Eto, K., Shibata, K., Murakoshi, H., Koizumi, S.,
Moorhouse, A. J., Yoshimura, Y., & Nabekura, J. (2016). Microglia contact induces synapse

formation in developing somatosensory cortex. *Nature Communications*, *7*(1), 12540. https://doi.org/10.1038/ncomms12540

Morari, J., Anhe, G. F., Nascimento, L. F., Moura, R. F. de, Razolli, D., Solon, C., Guadagnini, D., Souza, G., Mattos, A. H., Tobar, N., Ramos, C. D., Pascoal, V. D., Saad, M. J., Lopes-Cendes, I., Moraes, J. C., & Velloso, L. A. (2014). Fractalkine (CX3CL1) is involved in the early activation of hypothalamic inflammation in experimental obesity. *Diabetes*, *63*(11), 3770–3784. https://doi.org/10.2337/db13-1495

Nadjar, A., Leyrolle, Q., Joffre, C., & Laye, S. (2017). Bioactive lipids as new class of microglial modulators: When nutrition meets neuroimunology. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, *79*(Pt A), 19–26. https://doi.org/10.1016/j.pnpbp.2016.07.004

Nakajima, K., & Kohsaka, S. (2001). Microglia: Activation and Their Significance in the Central Nervous System. *The Journal of Biochemistry*, *130*(2), 169–175. https://doi.org/10.1093/oxfordjournals.jbchem.a002969

Ngozi, Z., & Bolton, J. L. (2022). Microglia Don't Treat All Neurons the Same: The Importance of Neuronal Subtype in Microglia-Neuron Interactions in the Developing Hypothalamus. *Frontiers in Cellular Neuroscience*, *16*, 867217. https://doi.org/10.3389/fncel.2022.867217

Ohsawa, K., Irino, Y., Sanagi, T., Nakamura, Y., Suzuki, E., Inoue, K., & Kohsaka, S. (2010). P2Y12 receptor-mediated integrin-β1 activation regulates microglial process extension induced by ATP. *Glia*, *58*(7), 790–801. https://doi.org/10.1002/glia.20963

Paolicelli, R. C., Bolasco, G., Pagani, F., Maggi, L., Scianni, M., Panzanelli, P., Giustetto, M., Ferreira, T. A., Giuducci, E., Dumas, L., Ragozzino, D., & Gross, C. T. (2011). Synaptic Pruning by Microglia Is Necessary for Normal Brain Development. *Science*, *333*(6048), 1456–1458. https://doi.org/10.1126/science.1202529

Parkhurst, C. N., Yang, G., Ninan, I., Savas, J. N., Yates, J. R., Lafaille, J. J., Hempstead, B. L., Littman, D. R., & Gan, W.-B. (2013). Microglia Promote Learning-Dependent Synapse Formation through Brain-Derived Neurotrophic Factor. *Cell*, *155*(7), 1596–1609. https://doi.org/10.1016/j.cell.2013.11.030

Plagemann, A., Heidrich, I., Götz, F., Rohde, W., & Dörner, G. (1992). Obesity and Enhanced Diabetes and Cardiovascular Risk in Adult Rats due to Early Postnatal Overfeeding. *Experimental and Clinical Endocrinology*, *99*(03), 154–158. https://doi.org/10.1055/s-0029-1211159

Riquier, A. J., & Sollars, S. I. (2020). Astrocytic response to neural injury is larger during development than in adulthood and is not predicated upon the presence of microglia. *Brain, Behavior, & Immunity - Health*, *1*, 100010. https://doi.org/10.1016/j.bbih.2019.100010

Rosin, J. M., & Kurrasch, D. M. (2019). Emerging roles for hypothalamic microglia as
regulators of physiological homeostasis. *Frontiers in Neuroendocrinology*, 100748.
https://doi.org/10.1016/j.yfrne.2019.100748

Samuelsson, A.-M., Matthews, P. A., Argenton, M., Christie, M. R., McConnell, J. M., Jansen,
E. H. J. M., Piersma, A. H., Ozanne, S. E., Twinn, D. F., Remacle, C., Rowlerson, A.,
Poston, L., & Taylor, P. D. (2008). Diet-Induced Obesity in Female Mice Leads to Offspring
Hyperphagia, Adiposity, Hypertension, and Insulin Resistance: A Novel Murine Model of
Developmental Programming. *Hypertension*, *51*(2), 383–392.
https://doi.org/10.1161/hypertensionaha.107.101477

Sanes, J. R., & Lichtman, J. W. (1999). DEVELOPMENT OF THE VERTEBRATE
NEUROMUSCULAR JUNCTION. *Annual Review of Neuroscience*, *22*(1), 389–442.
https://doi.org/10.1146/annurev.neuro.22.1.389

Schafer, D. P., Lehrman, E. K., Kautzman, A. G., Koyama, R., Mardinly, A. R., Yamasaki, R.,
Ransohoff, R. M., Greenberg, M. E., Barres, B. A., & Stevens, B. (2012). Microglia Sculpt
Postnatal Neural Circuits in an Activity and Complement-Dependent Manner. *Neuron*, *74*(4),
691–705. https://doi.org/10.1016/j.neuron.2012.03.026

Schafer, D. P., Lehrman, E. K., & Stevens, B. (2013). The "quad-partite" synapse: Microglia-
synapse interactions in the developing and mature CNS. *Glia*, *61*(1), 24–36.
https://doi.org/10.1002/glia.22389

Simerly, R. B. (2008). Hypothalamic substrates of metabolic imprinting. *Physiology & Behavior*,
94(1), 79–89. https://doi.org/10.1016/j.physbeh.2007.11.023

Skowronski, A. A., Leibel, R. L., & LeDuc, C. A. (2023). Neurodevelopmental Programming of
Adiposity: Contributions to Obesity Risk. *Endocrine Reviews*, *45*(2), 253–280.
https://doi.org/10.1210/endrev/bnad031

Spencer, S. J., Basri, B., Sominsky, L., Soch, A., Ayala, M. T., Reineck, P., Gibson, B. C., &
Barrientos, R. M. (2019). High-fat diet worsens the impact of aging on microglial function
and morphology in a region-specific manner. *Neurobiology of Aging*, *74*, 121–134.
https://doi.org/10.1016/j.neurobiolaging.2018.10.018

Stephan, A. H., Barres, B. A., & Stevens, B. (2012). The Complement System: An Unexpected
Role in Synaptic Pruning During Development and Disease. *Neuroscience*, *35*(1), 369–389.
https://doi.org/10.1146/annurev-neuro-061010-113810

Stettler, N., Stallings, V. A., Troxel, A. B., Zhao, J., Schinnar, R., Nelson, S. E., Ziegler, E. E.,
& Strom, B. L. (2005). Weight gain in the first week of life and overweight in adulthood: a
cohort study of European American subjects fed infant formula. *Circulation*, *111*(15), 1897–
1903. https://doi.org/10.1161/01.cir.0000161797.67671.a7

Stevens, B., Allen, N. J., Vazquez, L. E., Howell, G. R., Christopherson, K. S., Nouri, N., Micheva, K. D., Mehalow, A. K., Huberman, A. D., Stafford, B., Sher, A., Litke, A. M., Lambris, J. D., Smith, S. J., John, S. W. M., & Barres, B. A. (2007). The Classical Complement Cascade Mediates CNS Synapse Elimination. *Cell*, *131*(6), 1164–1178. https://doi.org/10.1016/j.cell.2007.10.036

Stowell, R. D., Wong, E. L., Batchelor, H. N., Mendes, M. S., Lamantia, C. E., Whitelaw, B. S., & Majewska, A. K. (2018). Cerebellar microglia are dynamically unique and survey Purkinje neurons in vivo. *Developmental Neurobiology*, *78*(6), 627–644. https://doi.org/10.1002/dneu.22572

Sun, J., Wang, X., Sun, R., Xiao, X., Wang, Y., Peng, Y., & Gao, Y. (2023). Microglia shape AgRP neuron postnatal development via regulating perineuronal net plasticity. *Molecular Psychiatry*, 1–11. https://doi.org/10.1038/s41380-023-02326-2

Sutton Hickey, A. K., Duane, S. C., Mickelsen, L. E., Karolczak, E. O., Shamma, A. M., Skillings, A., ... & Krashes, M. J. (2023). AgRP neurons coordinate the mitigation of activity-based anorexia. *Molecular psychiatry, 28*(4), 1622-1635. https://doi.org/10.1038/s41380-022-01932-w

Tamashiro, K. L. K., & Moran, T. H. (2010). Perinatal environment and its influences on metabolic programming of offspring. *Physiology & Behavior*, *100*(5), 560–566. https://doi.org/10.1016/j.physbeh.2010.04.008

Thaler, J. P., Yi, C.-X., Schur, E. A., Guyenet, S. J., Hwang, B. H., Dietrich, M. O., Zhao, X., Sarruf, D. A., Izgur, V., Maravilla, K. R., Nguyen, H. T., Fischer, J. D., Matsen, M. E., Wisse, B. E., Morton, G. J., Horvath, T. L., Baskin, D. G., Tschöp, M. H., & Schwartz, M. W. (2012). Obesity is associated with hypothalamic injury in rodents and humans. *Journal of Clinical Investigation*, *122*(1), 153–162. https://doi.org/10.1172/jci59660

Town, T., Nikolic, V., & Tan, J. (2005). The microglial "activation" continuum: from innate to adaptive responses. *Journal of Neuroinflammation*, *2*(1), 24. https://doi.org/10.1186/1742-2094-2-24

Tremblay, M.-È., Lowery, R. L., & Majewska, A. K. (2010). Microglial Interactions with Synapses Are Modulated by Visual Experience. *PLoS Biology*, *8*(11), e1000527. https://doi.org/10.1371/journal.pbio.1000527

Tremblay, M.-È., Stevens, B., Sierra, A., Wake, H., Bessis, A., & Nimmerjahn, A. (2011). The Role of Microglia in the Healthy Brain. *The Journal of Neuroscience*, *31*(45), 16064–16069. https://doi.org/10.1523/jneurosci.4158-11.2011

Valdearcos, M., Douglass, J. D., Robblee, M. M., Dorfman, M. D., Stifler, D. R., Bennett, M. L., Gerritse, I., Fasnacht, R., Barres, B. A., Thaler, J. P., & Koliwad, S. K. (2017). Microglial Inflammatory Signaling Orchestrates the Hypothalamic Immune Response to Dietary

Excess and Mediates Obesity Susceptibility. *Cell Metabolism*, *26*(1), 185-197.e3. https://doi.org/10.1016/j.cmet.2017.05.015

Valdearcos, M., Robblee, M. M., Benjamin, D. I., Nomura, D. K., Xu, A. W., & Koliwad, S. K. (2014). Microglia Dictate the Impact of Saturated Fat Consumption on Hypothalamic Inflammation and Neuronal Function. *Cell Reports*, *9*(6), 2124–2138. https://doi.org/10.1016/j.celrep.2014.11.018

Vogt, M. C., Paeger, L., Hess, S., Steculorum, S. M., Awazawa, M., Hampel, B., Neupert, S., Nicholls, H. T., Mauer, J., Hausen, A. C., Predel, R., Kloppenburg, P., Horvath, T. L., & Brüning, J. C. (2014). Neonatal Insulin Action Impairs Hypothalamic Neurocircuit Formation in Response to Maternal High-Fat Feeding. *Cell*, *156*(3), 495–509. https://doi.org/10.1016/j.cell.2014.01.008

Wake, H., Moorhouse, A. J., Jinno, S., Kohsaka, S., & Nabekura, J. (2009). Resting Microglia Directly Monitor the Functional State of Synapses In Vivo and Determine the Fate of Ischemic Terminals. *The Journal of Neuroscience*, *29*(13), 3974–3980. https://doi.org/10.1523/jneurosci.4363-08.2009

Wang, C., Yue, H., Hu, Z., Shen, Y., Ma, J., Li, J., Wang, X.-D., Wang, L., Sun, B., Shi, P., Wang, L., & Gu, Y. (2020). Microglia mediate forgetting via complement-dependent synaptic elimination. *Science*, *367*(6478), 688–694. https://doi.org/10.1126/science.aaz2288

Wong, A. M., Patel, N. V., Patel, N. K., Wei, M., Morgan, T. E., Beer, M. C. de, Villiers, W. J. S. de, & Finch, C. E. (2005). Macrosialin increases during normal brain aging are attenuated by caloric restriction. *Neuroscience Letters*, *390*(2), 76–80. https://doi.org/10.1016/j.neulet.2005.07.058

Young, A. M., Kumasaka, N., Calvert, F., Hammond, T. R., Knights, A., Panousis, N., Park, J. S., Schwartzentruber, J., Liu, J., Kundu, K., Segel, M., Murphy, N. A., McMurran, C. E., Bulstrode, H., Correia, J., Budohoski, K. P., Joannides, A., Guilfoyle, M. R., Trivedi, R., … Gaffney, D. J. (2021). A map of transcriptional heterogeneity and regulatory variation in human microglia. *Nature Genetics*, *53*(6), 861–868. https://doi.org/10.1038/s41588-021-00875-2

Zagmutt, S., Mera, P., Soler-Vázquez, M. C., Herrero, L., & Serra, D. (2018). Targeting AgRP neurons to maintain energy balance: Lessons from animal models. *Biochemical Pharmacology*, *155*, 224–232. https://doi.org/10.1016/j.bcp.2018.07.008

Zeltser, L. M. (2018). Feeding circuit development and early-life influences on future feeding behaviour. *Nature Reviews Neuroscience*, *19*(5), 302–316. https://doi.org/10.1038/nrn.2018.23

 Figures

Figure 1. MHFD: Microglial morphology in the PVH. (A) Image analysis pipeline. Maximum projection
of an Iba1 stained microglial cell in the PVH (Ai). Confocal images through labeled cells were used to

generate 3D reconstructions (Aii). Each 3D rendering was then used to create a skeletonized model by
using the Filaments tool in Imaris. Polyhedrons were generated around each cell using the Convex Hull
function of Imaris to estimate the total tissue "territory" occupied by the microglial cell (Aiii). (B-E)
Maximum projection images of microglial cells (green) and labeled AgRP terminals (red) in the PVH of
mice at P16 (B,C) or P30 (D,E) that were either exposed to MHFD (C, E) or NCD (B, D). (F-J) Graphical
comparisons between groups to show that MHFD increased microglial ramification complexity (F),
microglial cell territory (G), cell volume (H) and process length (I) and AgRP terminals (K). The density of
microglia in the PVH decreased between P16 and P30, irrespective of diet (J). Bars represent the mean $\pm$
SEM and each point represents one animal. *P<.05, **P<.005. Abbreviations: MHFD, maternal high fat
diet during lactation; NCD, Normal Chow Diet; PVH, paraventricular nucleus of the hypothalamus.
896

897
898
Figure 2. MHFD: Microglial morphology in the ARH. Maximum projection images of microglial cells
(green) in the ARH of mice at P16 (A, B) or P30 (C,D) that were either exposed to MHFD (B, D) or NCD
(A, C). Graphical comparisons between groups to show that microglial ramification complexity (E)
remained the same, regardless of age or diet. Microglial cell territory (F), cell volume (G) and process
length (H) increased between P16 and P30, but were not changed as a result of diet. The density of

microglia in the ARH decreased between P16 and P30, irrespective of diet (I). There were no changes in
numbers of AgRP neurons (J). Bars represent the mean ± SEM and each point represents one animal.
*P<.05, **P<.005. Abbreviations: ARH, arcuate nucleus of the hypothalamus; MHFD, maternal high fat
diet during lactation; NCD, Normal Chow Diet.
908

909
Figure 3. MHFD: Microglial morphology in the BST. Maximum projection images of microglial cells
(green) and labeled AgRP terminals (red) in the BST of mice at P16 (A, B) or P30 (C,D) that were either
exposed to MHFD (B, D) or NCD (A, C). Graphical comparisons between groups to show that microglial

ramification complexity (E) remained the same, regardless of age or diet. Microglial cell territory (F), cell
volume (G) and process length (H) did not significantly change between P16 and P30 and were not
changed as a result of diet. The density of microglia in the BST decreased between P16 and P30,
irrespective of diet (I). The density of AgRP terminals increased between P16 and P30, but there was no
effect of maternal diet (J). Bars represent the mean ± SEM and each point represents one animal.
*P<.05. Abbreviations: BST, bed nucleus of the stria terminalis; MHFD, maternal high fat diet during
lactation; NCD, Normal Chow Diet.
920
921
922

923
924
Figure 4. Microglial depletion during lactation period. Images of microglial cells (green) to assess
microglia number (A,B). Maximum projection images of labeled AgRP terminals (red) to assess AgRP
terminal density in distinct compartments of the PVH at P55 long after the period of daily injections (C-F).
Graphical comparison between groups to show that daily postnatal PLX5622 injections caused a 33%
decrease in microglia in the PVH (G). Maximum projections of confocal images to illustrate the density of
AgRP labeling in the PVH of NCD offspring (C,D) and MHFD offspring (E,F). (G-J) Graphical comparison
to illustrate the effects of postnatal PLX5622 treatments on microglia in the PVH (G), body weight (I) and
the density of AgRP terminals in the PVHmpd (H) and PVHpml (J). Bars represent the mean ± SEM and
each point represents one animal. *P<.05, **P<.005, ***P<.0005. Abbreviations: AgRP, agouti-related
peptide; CSF1R, Colony-Stimulating Factor 1 Receptor; MHFD, maternal high fat diet during lactation;
MPD, medial parvocellular compartment of the PVH; PML, posterior magnocellular compartment of the
PVH; PVH, paraventricular nucleus of the hypothalamus.
937

939
Figure 5. Microglial interaction with AgRP axon terminals in the PVH and ARH. (A-D) Maximum
projection images of microglial cells (green) labeled AgRP terminals (red), and CD68 (lysosomal
associated membrane protein and phagocytic capacity marker, pink). (E-H) Digital 3D reconstructions
shown in (A-D) after application of a digital zoom to more clearly illustrate engulfment of labeled AgRP
terminals by microglia (I-L). (M-P) Graphical comparisons between groups to illustrate the effects of age
and MHFD exposure on CD68 expression and AgRP terminal engulfment. Bars represent the mean ±
SEM and each point represents one animal. *P<.05, **P<.005. Abbreviations: AgRP, agouti-related
peptide; ARH, arcuate nucleus or the hypothalamus; CD68, Cluster of Differentiation 68; MHFD, maternal
high fat diet during lactation; PVH, paraventricular nucleus of the hypothalamus.
949