Title: Test-retest reliability of TMS motor evoked responses and silent periods during

explosive voluntary isometric contractions

Castelli[1] F., Mian[1] O.S., Bruton[1,2] A., Chembila Valappil[1] A.C., Tillin[1] N.A.

[1] University of Roehampton, Life Sciences, Whitelands College, Holybourne Avenue, London,

SW15 4JD, United Kingdom

[2] Brunel University London, Centre for Cognitive and Clinical Neuroscience, Sport

Health and Exercise Sciences, Uxbridge, UB8 3PH, United Kingdom

Castellf1@roehampton.ac.uk

9   **Abstract**

Purpose

This study assessed the intraclass correlation coefficient (ICC) and coefficient of variation (CV) test-
retest reliability of TMS responses MEPs and silent periods at early, middle, and late phases of the
rising time-torque curve during explosive voluntary contractions. We also investigated how the
number of consecutively averaged measurements over 3 to 15 separate contractions influenced
reliability.

Methods

On two separate occasions 3-7 days apart, 14 adults completed 48 isometric explosives (1-s)
contractions of the knee extensors, superimposed with TMS. The TMS elicited an MEP and silent
period in the quadriceps muscles at 45 (early), 115 (middle), or 190 ms (late) during each
contraction. TMS was also superimposed at the plateau of 15 separate MVCs. Test-retest ICC and
CV were calculated for consecutively averaged MEPs and silent periods.

Results

No one condition was more reliable than another. For MEP amplitude, in all conditions except the
explosive late phase, ICCs generally increased, and CV decreased, with an increase in the number of
averaged contractions, and were >0.50 ICC and <15% CV within 7 contractions. For silent period,
ICCs and CVs were unaffected by the number of consecutively averaged contractions and remained
>0.50 ICC and <10% CV.

Conclusion

Test-retest reliability of TMS responses is comparable between phases of explosive contraction and
at the plateau of MVC. To enhance MEP reliability, with the increases he number of averaged
contractions of except the explosive late phase, we recommend studies average data across > the 3-
5 contractions typically reported in the literature for studies using TMS during MVCs.

Key words: Reliability, TMS, explosive contractions, test-retest

Abbreviations

AMT = Active motor threshold

CV = coefficient of variance

EMG = Electromyography

ICC = Inter-class correlation

MEP = Motor evoked potential

MVC = Maximal voluntary contraction

MU = Motor unit

RF = Rectus femoris

RMS = Root mean squared

RTD = Rate of torque development

SD = Standard deviation

SP = Silent period

TMS = Transcranial magnetic stimulation

VM = Vastus medialis

VL = Vastus lateralis

Introduction

The rate of torque development (RTD) measures the ability of muscles to rapidly increase

force around a joint (Tillin and Folland, 2014), and is functionally important where time to develop

force is limited, such as sprinting (Tillin, Pain and Folland, 2013) or balance recovery (Sundstrup *et*

al., 2010). RTD is often measured in early (0-50 ms), middle (50-100 ms) and late (>100 ms) phases

of explosive contraction performed from rest (Maffiuletti *et al.*, 2016), and different physiological

factors limit the RTD in these separate phases. One limiting factor is the neural drive to muscle,

which positively correlates with early- and middle-phase RTD (Folland, Buckthorpe and Hannah,

2014; Del Vecchio *et al.*, 2019). Despite the relevance of neural drive to RTD, the corticospinal

mechanisms affecting RTD are not well established.

Transcranial magnetic stimulation (TMS), superimposed during voluntary contractions, elicits

a motor-evoked potential (MEP) in the surface EMG signal of the target contracting muscle. The

MEP amplitude is thought to reflect corticospinal excitability (Rossini *et al.*, 2015), whilst a period of

electrical inactivity immediately after the MEP, referred to as the silent period (Damron *et al.*, 2008),

is thought to reflect corticospinal inhibitory mechanisms (Säisänen *et al.*, 2008). Despite extensive

use of TMS to explore corticospinal excitability/inhibition at the plateau of a maximal voluntary

contraction (MVC; Todd, Taylor and Gandevia, 2016), TMS has not been commonly used during the

rising torque of explosive voluntary contractions, where RTD is typically measured.

Before using TMS to assess corticospinal excitability/inhibition during explosive contractions,

it is important to establish the test-retest absolute and relative reliability of TMS responses (MEP

amplitude and silent period) during such conditions; however, this has not been done. In contrast,

the test-retest reliability for TMS responses at the plateau of MVCs has been investigated in various

muscles. For absolute and normalised MEP amplitude at the MVC plateau in the lower limbs, studies

have reported moderate inter-class correlation (ICC; 0.52-0.79) (Mileva, Sumners and Bowtell, 2012;

Souron *et al.*, 2016) and coefficient of variation (CV) of 10-11% (Souron *et al.*, 2016). For the silent

period at the MVC plateau, the test-retest ICC has ranged from moderate in the vastus lateralis

(0.61-0.70; (Di Virgilio *et al.*, 2022)) to excellent in the soleus and tibialis anterior (0.93-0.95; Mileva,

Sumners and Bowtell, 2012; Souron *et al.*, 2016) and one study has reported a CV of 8.6% (Souron *et*

al., 2016) in the tibialis anterior. It is unclear whether the reliability of MEP amplitudes and silent

periods during explosive contractions will be comparable to that observed at the MVC plateau.

Typically, the reliability of torque and EMG amplitude measurements is lower in early compared to

later phases of explosive contraction, and generally lower for explosive contractions compared to

the MVC plateau (De Ruiter *et al.*, 2004; Buckthorpe *et al.*, 2012), so a similar pattern may be

observed for the reliability of MEP amplitudes and silent periods.

TMS responses are typically averaged across multiple separate stimuli (in separate

contractions) to minimise the influence of random variation and improve reliability. Thus, when

assessing the test-retest reliability of MEP amplitudes and silent periods, it is important to consider

how many stimulations might be required to achieve adequately reliable averages. A relatively high

number of stimuli (≥20) are needed to obtain moderate to good test-retest reliability (ICC = 0.50 -

0.75; Goldsworthy, Hordacre and Ridding, 2016; Biabani *et al.*, 2018) of MEP amplitudes recorded

when the participant is passive. In contrast, as few as four stimuli have been shown to produce good

(≥0.75) test-retest ICC for MEP amplitude during submaximal contractions held at a constant force

(Wheaton *et al.*, 2009; Lewis, Signal and Taylor, 2013; Temesi, Ly and Millet, 2017). The effect of the

number of contractions on test-retest reliability of average TMS responses has not been determined

during either MVC or explosive contractions. Previous studies of TMS responses during MVCs have

typically only collected and averaged 3-5 responses (Luc *et al.*, 2014; Tallent *et al.*, 2017; Škarabot *et*

al., 2019). This could be due to time constraints or concerns over fatigue with multiple contractions;

however, it may be at the expense of poor reliability. Thus, there is a need to assess the effect of

the number of contractions on the test-retest reliability of average TMS responses during MVCs and

explosive contractions.

In studies of TMS responses, measures are typically averaged across all stimulations

obtained within a testing session. However, a subset of contractions are typically used for studies of

explosive contractions, with the best three out of ten contractions being recommended for

averaging RTD measurements (Maffiuletti *et al.*, 2016). Therefore, for studies of MEPs recorded

during explosive contractions, it may be desirable to investigate MEPs obtained from the best three

contractions rather than from all contractions, and investigation into the reliability of this approach

is necessary.

This study aims to assess the absolute (ICC) and relative (CV) test-retest reliability of TMS

responses (MEP amplitude and silent period) recorded at different time points (early, middle, and

late phases) during explosive voluntary contractions and at the plateau of MVCs. As part of this aim,

we intend to (i) document how the number of consecutive contractions (between 3-15) over which

TMS responses are averaged, influences test-retest reliability; and (ii) document test-retest reliability

of TMS responses averaged across the best 3, out of 10, contractions. We determine the best three

as those with the highest torque (Torque averaging method) or EMG RMS (EMG averaging method)

prior to MEP on-set.

Methods

Participants

Fourteen participants, nine males (age 31 ± 5 years, height, 178.7 ± 6.6cm, and mass 79.2 ±

4.7kg) and five females (age 29 ± 6 years, height, 165.5 ± 5.5cm, and mass 59.3 ± 6.2kg) were

recruited to take part in this study. All participants habitually performed 120-180 minutes of

moderate to high-intensity activity per week, and were deemed recreationally active (McKay *et al.*,

2022). Participants were also free from injury and disease (screened via a questionnaire adapted

from Balady *et al.*, 1998) and free from contraindications to TMS (screened via a questionnaire

adapted from Rossi *et al.*, 2011). Due to changes in endogenous hormones throughout the

menstrual cycle potentially affecting neuromuscular responses to TMS (Ansdell *et al.*, 2019), female

participants undertook experimental trials exclusively during their self-reported early to mid-
follicular phase (first ten days from the first day of menstruation), when endogenous hormone
concentration is low and relatively stable (de Jonge, Thompson and Han, 2019). The University of
Roehampton ethics committee approved the study, and all participants provided written informed
consent before participating.

Overview

Participants visited the laboratory on three separate occasions and were asked to avoid strenuous
exercise and alcohol consumption for 24 hours before each visit. Each session lasted approximately
120-150 minutes, with consecutive sessions separated by 3-7 days.   The first visit was a
familiarisation session, and the second and third were measurement sessions. The measurement
sessions were performed at a consistent time of day and involved an identical protocol, with
measurements used to assess the test-retest reliability of the variables of interest. Each session
involved knee extensor torque and EMG measurement during maximal voluntary and explosive
contractions, using TMS to obtain superimposed MEP and femoral nerve stimulation to obtain
compound muscle action potentials at rest.

Torque measurements and surface electromyography (EMG)

Participants were tightly secured in a custom-built strength testing chair (Fig. 6b in Maffiuletti *et al.*,
2016) with a waist belt and shoulder straps. The hip and knee angles were set at 100° and 105°,
respectively (full extension being 180°). All contractions were isometric knee extensions performed
with the right leg. An ankle strap joined to a calibrated S-shaped load cell (FSB-1.5kN, Force Logic,
Reading, UK) was secured 4 cm proximal to the medial malleolus. The force signal was amplified
(x375) and then sampled at 2000 Hz (Mirco3 1401 and Spike2 v.8; CED., Cambridge, UK). Offline, the
force was filtered (fourth-order low-pass Butterworth, 250 Hz cut-off), corrected for limb weight,
and multiplied by the external moment arm to calculate joint torque.

The skin was prepared by shaving, cleaning (70% ethanol), and lightly abrading the area

where EMG electrodes were placed.   A single, bipolar silver-silver-chloride gel-electrode

configuration (2-cm diameter and 2-cm inter-electrode distance; Dual Electrode, Noraxon, Arizona,

USA) was placed over the belly of each of the rectus femoris (RF), vastus lateralis (VL) and vastus

medialis (VM), based on SENIAM guidelines (Hermens *et al.*, 1999).   The EMG system has an

inherent 312-ms delay, so whilst suitable for measurements involved in the study, EMG signals

sampled by it could not be used to detect activation on-set and trigger the TMS in real-time during

the explosive contractions (explained below in Experimental procedures).   Thus, a second wired

bipolar EMG electrode (2 cm diameter and 2 cm inter-electrode distance; Dual Electrode, Biometrics

Ltd, Gwent, UK) was placed on the belly of the VM to trigger the TMS during contraction. Electrode

locations were marked with a permanent marker pen, and participants were asked to maintain these

marks throughout the study by re-applying them if necessary. The three wireless EMG signals were

transmitted to a desktop receiver (TeleMYO D.T.S., Noraxon, Arizona, USA) and sampled, along with

the single wired EMG signal, at 2000 Hz via the same A/D convertor and software as the force signal.

Once offline, wireless EMG signals were filtered (fourth-order Butterworth, band-pass, 6-500 Hz) and

time-corrected for the 312-ms delay inherent in the Noraxon system.

Transcranial magnetic stimulation (TMS)

TMS with a 1-ms pulse width was delivered via a double cone coil (110-mm Magstim 200,

Whitland, UK) over the scalp in an optimal position to elicit MEPs in the right quadriceps muscles.

The following procedures were completed in every session. Participants wore a swim cap, and the

vertex of the head – identified as 50% of the distance between nasion and inion and between right

and left perpendicular point – was marked on the swim cap. A 5-by-5-cm grid with 1-cm spacing

between grid lines was drawn on the swim cap, lateral (left hemisphere) and posterior from the

vertex. The coil was moved posteriorly and laterally from the vertex in ~0.5-cm steps, and in each

position, the participant completed four submaximal voluntary contractions at 20% MVC torque

(established in the warm-up; see below) with superimposed TMS on each contraction at a

submaximal (range 50-60%) stimulator output. The position determined the optimal coil position for

the rest of the session, which was subjectively the highest consistent MEP amplitudes (peak-to-peak)

over the four superimposed contractions for all three muscles (RF, VL and VM). The optimal coil

position was marked by drawing the edge of the coil over the swim cap, ensuring accurate relocation

throughout the session. The active motor threshold (AMT) was then determined via a series of 20%

MVC torque contractions superimposed with TMS stimulator output starting at 39%. The AMT was

defined as the minimum TMS intensity required to elicit five visible MEPs amongst the background

EMG activity of the VM and RF, out of ten consecutive superimposed contractions. If the muscles

had less or more than five visible MEPs, the machine intensity was reduced or increased by 2% of

the machine output. We prioritised the VM and RF (not the VL) as these muscles produced more

consistent and visible MEPs in pilot testing. Where it was impossible to match AMT for VM and RF,

we settled on being one visible MEP away from 5/10 (e.g., 6/10 VM and 4/10 RF). The same

investigator held the coil by hand throughout the measurement sessions, continuously monitoring

its position and orientation. For TMS delivered during maximal and explosive contractions (see

experimental protocol), the intensity was set at 140% of the stimulator output at AMT (Groppa *et*

al., 2012; Rossini *et al.*, 2015; Rossi *et al.*, 2020). These procedures were repeated for all sessions.

Femoral nerve electrical stimulation

Single, square-wave pulses (200 μs duration) were delivered (DS7AH, Digitimer,

Hertfordshire, UK) over the femoral nerve in the inguinal triangle to evoke twitch contractions and

obtain compound muscle action potentials (M-waves) at rest. The anode (5 x 8 cm carbon rubber;

EMS Physio Ltd., Oxfordshire, UK) was placed over the head of the greater trochanter. The optimal

location of the cathode (1 cm diameter tip; S1 Compex Motor PointPen, Digitimer, UK) was

determined as that which evoked the greatest peak twitch torque for a given submaximal

stimulation intensity (80-120 mA). The cathode was taped down and held in position by the same

investigator as a series of twitches at incremental intensities were evoked until there was a plateau

in the peak-to-peak M-wave amplitude (M_{max}) of all three muscles (RF, VL, VM). The stimulator

intensity was then increased to 150% of the intensity at M_{max}, ensuring supramaximal intensity.

Three supramaximal twitch contractions were then evoked, each separated by 15 s, and M_{max} was

averaged across the three contractions for each muscle. The procedures were repeated for all

sessions.

Experimental protocol

Participants first completed a warm-up involving a series of explosive, submaximal, and maximal

voluntary contractions (the latter being used to establish MVC torque), followed by procedures for

obtaining optimal TMS coil position, ATM, and M_{max}. Participants then completed a series of MVCs

and explosive voluntary contractions with and without superimposed TMS. The instruction for MVCs

was to "push as hard as possible" for 3-5 s and explosive contractions to "push as fast and hard as

possible", emphasising fast for 1 s. The protocol was organised into three blocks of contractions (Fig.

1). Each block involved 24 explosive contractions (8 without and 16 with superimposed TMS) and 8

MVCs (3 without and 5 with superimposed TMS), distributed across four sets. Each set involved six

explosive contractions (the final four with TMS stimulation) and two MVCs. In sets 1-3, one of the

two MVCs had superimposed TMS (randomly ordered), and in set four, both MVCs had

superimposed TMS (Fig. 1). Participants rested 10 s between explosive contractions, 30 s between

MVCs, 120 s between sets, and 300 s between blocks. Each block was identical except for the timing

of TMS application during explosive contractions, with each block using a different TMS timing

condition for these contractions (see next section). The order of the blocks was randomised across

participants but held constant across sessions for the same participant. Overall, the 3-block protocol

yielded 24 explosive contractions and 9 MVCs without TMS, and 48 explosive contractions (16 per

stimulus time condition) and 15 MVCs with TMS.

223   [INSERT FIGURE 1 HERE]

Fig 1 *The order of experimental contractions in a block. There were 3 blocks per experimental session and each block*
involved a different condition for explosive contractions performed with superimposed TMS. In one condition/block TMS
occurred in the early phase, in another the middle phase, and in another the late phase, of explosive contraction. The order
that blocks/conditions were completed was randomised. Each block consisted of 4 sets of contractions as shown. The thin
blue bar represents explosive contractions, while the wide blue bar represents MVCs. The red triangle on the top of the blue
bars represents the superimposed TMS

230   [INSERT FIGURE 2 HERE]

Fig 2 Representative traces of motor evoked potentials (full black line) in the vastus medialis and knee extensor torque
(empty dotted line) in each of the four conditions: Early (45 ms), middle (115 ms), and Late (190 ms) from EMG on-set
during explosive contractions, and at MVC plateau. The vertical black line across all graphs represents the time of
stimulation. The blue line in each graph represents when the TMS was triggered in each condition. The red dotted arrows
represent the peak-to-peak amplitude of the MEPs used for analysis

For superimposed MVCs, the TMS was triggered manually by the same experienced

investigator during the torque plateau of the MVC. For explosive contractions, the TMS was

triggered automatically when the VM EMG signal of the wired system exceeded a set threshold. The

threshold was set in the Spike2 software as the lowest amplitude for that session above the highest

peaks and troughs of the baseline noise. Commencing from manually detected EMG on-set (0 ms;

determined via the methods of Tillin *et al.*, 2010), the threshold was exceeded after ~5 ms, and

triggered the TMS at either 8, 78, or 153 ms (accounting for a 3 ms system delay), for the early,

middle, and late-phase conditions, respectively (Fig 2). The centre of the evoked MEPs would then

occur at ~45 (early), 115 (middle), or 190 ms (late) from manually detected EMG on-set (Fig. 2).

Data screening and analysis

Out of 15 superimposed MVCs, only those where the TMS was delivered >90% of MVC

torque were used for further analysis. Out of the 16 superimposed explosive contractions per

condition, only those that met the following criteria were used for further analysis: (i) average

baseline force did not change by >2 Nm during the 200 ms preceding manually detected force on-set

(detected as in Tillin *et al.*, 2010); and (ii) there was a genuine attempt at an explosive contraction. A

genuine attempt at an explosive contraction was defined as the instantaneous slope of torque-time,

just prior to MEP on-set, being within 3 standard deviations (SD) of mean instantaneous slope at the

same time point for contractions without TMS. The time points for measuring instantaneous slope

were 30, 105, and 180 ms after manually detected torque on-set, for early-, middle- and late-phase

conditions, respectively. Given the criteria above, the maximum number of MEPs analysed further

in each condition was limited to the number of usable contractions in the participant with the lowest

number of usable contractions.　This was 9 (early), 10 (middle), and 15 (late) for the explosive

contractions, and 15 for the MVCs.

For useable contractions, MEP amplitude was defined as the peak-to-trough of the

superimposed response (Fig. 2) and is reported in absolute terms and normalised to M_{max}. The silent

period was measured from the point of stimulation to the resumption of EMG activity after the MEP

off set.　To determine the resumption of EMG activity, the second derivative of EMG amplitude over

time was established (1-ms time constant) and then the signal was rectified.　EMG activity

resumption was defined when the amplitude of the second derivative increased above 5 SD of the

mean baseline (calculated in a 500 ms period prior to the contraction) for 70% of the next 10 ms

(adapted from Damron *et al.*, 2008; Fig.3).　This automated process was confirmed via manual

inspection of the signal as recommended by Damron et al., (2008).　MEP amplitude (absolute and

normalised to M_{max} separately) and silent periods were extracted on a muscle level before being

averaged across the three muscles to obtain a quadriceps mean.　This is standard practice in studies

looking to relate general quadriceps EMG data to net knee extensor torque and RTD (Tillin *et al.*,

2010; Tillin, Pain and Folland, 2012, 2013, 2018; Folland, Buckthorpe and Hannah, 2014; Behrens *et*

al., 2015; Morales-Artacho *et al.*, 2018; Cossich and Maffiuletti, 2020).

Averaging methods

For each condition (early, middle, and late explosive phases, and at MVC plateau), mean

quadriceps consecutive running averages of the dependent variables were calculated from the first

three stimuli/contractions, up to the maximum number of usable contractions: 9 (early), 10 (middle),

and 15 (late) for the explosive contractions, and 15 for the MVCs.　In the torque and EMG averaging

methods, dependent variables were averaged over the best 3 out of the first 10 contractions (9 for

early-phase explosive) in each condition. For the torque averaging method, the best 3 explosive

contractions were those with the greatest torque just prior to MEP (i.e., at 30, 105, and 180 ms from

torque on-set, in early, middle, and late conditions, respectively), whilst the best 3 MVCs were those

with the highest average torque over 1 s prior to the MEP. For the EMG averaging method, the best

explosive contractions were those with the highest EMG RMS amplitude (averaged of all three

muscles) from on-set to the start of the MEP (i.e., 0-23, 0-98, 0-173 ms, in early, middle, and late,

respectively), whilst the best three MVCs were those with the greatest EMG RMS in the 1 s period

before MEP.

287   [INSERT FIGURE 3 HERE]

Fig 1 Example of how the silent period off-set was detected in the VM of one participant. The data shows the second
derivative of the EMG-time trace after rectification. Circled data are an example of SP off-set detection. The orange line
represents the detection threshold (5 SD of the mean baseline), whilst the red dot, which is the detected off-set, represents
the last time point before the EMG signal crosses threshold, and remains above the threshold for ≥70% of the data points
recorded in the next 10 ms

Statistical analysis

Paired t-tests were used to compare the differences between the test and re-test sessions

within each condition for each dependent variable. The relative reliability of TMS responses was

assessed via the relative test-retest interclass correlation coefficient (ICC) determined via a two-way

mixed-effects model, with the absolute agreement, as suggested for a test-retest design (Koo and Li,

2016). ICCs were interpreted as <0.5 poor, ≥0.5 but <0.75 moderate, ≥0.75 and <0.9 good, and ≥0.9

excellent (Koo and Li, 2016). The coefficient of variation (CV) determined the absolute reliability

(degree of fluctuation of repeated measurements within individuals) of dependent variables and was

calculated for each participant (100 * test-retest standard deviation (SD) / test-retest mean) before

being averaged across participants.

Separate ICCs and CVs were produced for each averaging approach; consecutive running

averages and the best three based on both torque and EMG averaging methods. Logarithmic

functions $(y = a \ln(x) + k)$ were fit to data describing the relationship between ICC or CV and the

number of consecutive contractions/stimuli over which the dependent variables were averaged to

establish trends in the data and enable extrapolation of the ICC or CV beyond the maximum number

of useable contractions. Only statistically significant (p<0.05) logarithmic functions are reported in

the results. Logarithmic functions were chosen as they provided excellent fits with significant

(p<0.05) relationships (r^2=0.97-0.75 and SD=0.08) and did not cross zero values of CV (as with, e.g.,

exponential fits), which are physiologically impossible. Best-fit logarithmic functions were made in

SPSS v.26 (IBM Inc., Armonk, NY, USA). For ICCs, a confidence interval of 95% was used for the lower

bound (Lb) and upper bound (Ub), and CVs were calculated using MATLAB (R2021a Update 5,

MathWorks Inc., Natick, MA, USA). The significance level was set at $p \leq 0.05$. Data are presented in

the results as mean ± SD.

Results

There was no significant difference (p≥0.168) between the test and retest sessions for any of

the dependent variables in each condition (Table 1).

	MEP amplitude (mV)		MEP amplitude (% M_{max})		Silent Period (s)	
	Test	Re-test	Test	Re-test	Test	Re-test
Early	0.96 ± 0.40	0.92 ± 0.38	30.2 ±8.3	30.2 ± 9.4	0.078 ± 0.002	0.078 ± 0.002
Middle	0.97 ± 0.47	0.95 ± 0.40	30.0 ± 6.9	31.7 ± 7.8	0.083 ± 0.002	0.083 ± 0.001
Late	1.01 ± 0.45	0.96 ± 0.41	31.8 ± 7.7	31.3 ± 6.9	0.088 ± 0.002	0.089 ± 0.001
MVC	1.10 ± 0.46	1.02 ± 0.39	35.4 ± 7.3	35.5 ± 10.3	0.091 ± 0.005	0.093 ± 0.004

Table 1 shows the absolute and normalised MEP amplitude, and silent period for each condition in each session (test and re-
test). Normalised MEP amplitude is a % of maximal M-wave (M_{max}). Conditions are early (45), middle (115), and late (190)
phases of explosive contractions, and at the MVC plateau. Data are mean ±SD of the first 10 (9 for early) contractions
performed in each condition.

Reliability as a function of the number of MEPs

The test-retest ICCs for absolute and normalised MEP amplitude generally increased with

the number of averaged MEPs (Fig.3 A, B), whilst the test-retest CV for the same variables generally

decreased (Fig.3 C, D). These trends indicate increasing test-retest MEP amplitude reliability with

increasing numbers of averaged MEPs. The exception to these general trends was the late phase. In

this condition, although the CV of normalised MEP amplitude was related to the number of averaged

MEPs (Fig 3D), the CV of absolute MEP amplitude, the ICC of absolute MEP amplitude, and the ICC of

normalised MEP amplitude were not related to the number of averaged MEPs (Figs 3 A, B, C). These

trends, apparent upon visual inspection, were supported by logarithmic fits to the data, which were

significant in all cases aside from the late phase cases highlighted in the previous sentence (Table 2).

ICCs were generally higher for absolute MEP amplitude (Fig 3A) than normalised MEP

amplitude (Fig 3B). For absolute MEP amplitude, good ICC (ICC>0.75) was found within just three

averaged MEPs for all conditions except middle (where it was attained within nine averaged MEPs).

For normalised MEP amplitude, good ICC was only achieved in the early condition (attained within

seven contractions), whilst other conditions only reached moderate ICCs (ICC>0.50). CV tended to

be similar for absolute and normalised MEP amplitudes, reaching 10-15% values for all conditions

within seven contractions (Figs 3C, 3D). We were interested in identifying the number of averaged

MEPs needed to attain good ICC (ICC>0.75) and CV<10% for all variables and conditions. As is

noticeable from Fig 3 A-D, this was not consistently reached using the number of MEPs measured in

this study (9-15, depending on the condition). Therefore, the fitted logarithmic functions provided

estimates by extrapolating beyond fitted data where necessary (see Table 2 "n of" columns).

In contrast to the general trends for MEP amplitude reliability, the silent period reliability did

not vary as a function of number of averaged MEPs (Fig.3E, F). In support, attempted logarithmic fits

were all non-significant (p>0.05; fits not shown). Moderate ICCs (0.5≤ICC≤0.75) were generally

observed for the silent period in all conditions (Fig 3E). CVs for the silent period (Fig 3F) were lower

than for MEP amplitude, particularly during MVC.

                                     [INSERT FIGURE 4 HERE]

Fig 4 *test-retest ICC (A , B, E) and CV (C, D, F) for dependent variables as a function of the number of consecutive*
contractions over which data are averaged. Dependent variables are absolute (A and C) and normalised (B and D) MEP
amplitude, and silent period (E and F). For normalised data, MEP amplitude is normalised to maximal M-wave. MEP
amplitudes and silent periods were recorded at the plateau of MVCs, and at 3 different time points from activation on-set
during explosive contractions: early (45 ms), middle (115 ms), and Late (190 ms). Logarithmic functions (y=a ln(x)+k) were
fitted to data and are plotted as a dotted line for conditions where the fit was significant (p<0.05)

ICC

MEP absolute	a	k	R^2	p	n of	MEP normalised	a	k	R^2	p	n of
Early	0.116	0.704	0.96	0.001	2	Early	0.115	0.537	0.95	0.001	7
Middle	0.231	0.265	0.92	0.001	9	Middle	0.392	-0.395	0.84	0.002	19*
Late	0.002	0.819	-0.20	0.950	-	Late	-0.093	0.660	0.13	0.220	-
MVC	0.427	0.200	0.93	0.001	4	MVC	0.154	0.273	0.85	0.002	23*

CV

MEP absolute	a	k	R^2	p	n of	MEP normalised	a	k	R^2	p	n of
Early	-5.17	23.38	0.90	0.001	14*	Early	-4.77	24.28	0.87	0.001	20*
Middle	-7.23	29.55	0.94	0.001	15*	Middle	-5.41	22.61	0.93	0.001	11*
Late	-1.76	15.82	0.16	0.210	-	Late	-3.04	18.71	0.95	0.001	18*
MVC	-2.16	15.19	0.68	0.013	11	MVC	-2.90	19.04	0.79	0.005	23*

Table 2. Coefficients (a and k), the goodness of fit (R^2) of the logarithmic functions $y = a\,ln(x) + k$. Functions were fit to
describe the relationship between test-retest ICC or CV and the number of consecutive contractions over which MEP
*amplitudes were averaged. The 'n of' is the number of MEP amplitudes averaged to reach ICC≥0.75 or CV≤10%. The ***
denotes where 'n of' has been extrapolated beyond the measured data. MEP amplitudes were recorded at the plateau of
MVCs, and at three different time points from EMG on-set during explosive contractions: Early (45 ms), middle (115 ms),
and Late (190 ms).

Reliability for best three contractions using EMG or torque methods.

Compared to averaging the first 10 consecutive contractions, the ICC tended to be lower (-0.07±0.01) and CV slightly higher (+1.7±0.2%) for either of the best 3 averaging methods, across all conditions and dependent variables (Table 4). There was no consistency for the torque (best 3) averaging method to be better than the EMG averaging method, nor vice versa, with the higher ICC and lower CV between methods being dependent on condition and variable.

Dependent Variables	Averaging Method		
Absolute MEP amplitude	First 10	Best 3	
		EMG	Torque
Early	0.95 (0.84-0.98)	0.93 (0.79-0.98)	0.88 (0.65-0.96)
Middle	0.79 (0.47-0.93)	0.74 (0.26-0.89)	0.69 (0.64-0.97)
Late	0.85 (0.61-0.95)	0.83 (0.56-0.94)	0.84 (0.57-0.94)
MVC	0.85 (0.59-0.95)	0.60 (0.14-0.85)	0.77 (0.43-0.92)
Normalised MEP amplitude	First 10	Best 3	
		EMG	Torque
Early	0.78 (0.39-0.92)	0.72 (0.34-0.90)	0.65 (0.21-0.87)
Middle	0.55 (0.06-0.83)	0.37 (-0.1-0.74)	0.28 (-0.3-0.69)
Late	0.49 (-0.05-0.81)	0.49 (-0.07-0.80)	0.40 (-0.17-0.76)
MVC	0.63 (0.14-0.86)	0.53 (0.02-0.82)	0.72 (0.32-0.90)
Silent Period	First 10	Best 3	
		EMG	Torque
Early	0.62 (0.14-0.88)	0.49 (0.15-0.88)	0.64 (-0.2-0.92)
Middle	0.71 (0.28-0.90)	0.63 (0.21-0.89)	0.68 (0.16-0.93)
Late	0.68 (0.21-0.89)	0.63 (0.13-0.87)	0.63 (0.13-0.87)
MVC	0.71 (0.23-0.89)	0.66 (0.20-0.88)	0.77 (0.42-0.92)

Table 3. test-retest ICC, with 95% confidence interval in parenthesis, of dependent variables determined via 3 different averaging methods: average of the first ten consecutive contractions, and average of the best 3 based on those contractions with the highest EMG or torque prior to the MEP. Dependent variables are absolute MEPs, normalised MEPs (normalised to maximal M-wave), and silent period. MEP data were recorded at the plateau of MVCs, and at 3 different time points from EMG on-set during explosive contractions: early (45 ms), middle (115 ms), and late (190 ms)

Dependent Variables	Averaging Method		
Absolute MEP amplitude	First 10	Best 3	
		EMG	Torque
Early	11.7±10	11.9±9.6	16.5±14.6
Middle	12.6±8.9	15.5±11	14.3±10.1
Late	10.7±7.8	12.6±8.5	12.6±8.2
MVC	10.7±6	14.7±11.9	10.0±10.1
Normalised MEP amplitude	First 10	Best 3	
		EMG	Torque
Early	14.0±9.8	11.9±10.6	16.9±14.3
Middle	10.0±7.8	12.3±9.9	13.4±11
Late	11.0±5.1	12.1±7.6	13.1±7.8
MVC	10.0±7.2	17.1±10.3	13.1±9.3
Silent Period	First 10	Best 3	
		EMG	Torque
Early	9.7±9.2	10.3±9.3	8.2±7.3
Middle	7.9±7.1	9.4±7.3	9.0±7.2
Late	7.8±5.7	8.5±5.2	8.9±5.3
MVC	4.1±3.4	4.7±4.1	4.3±2.4

Table 4. test-retest CV mean ± SD of dependent variables determined via 3 different averaging methods: average of the first ten consecutive contractions, and average of the best 3 based on those contractions with the highest EMG or torque prior to the MEP. Dependent variables are absolute MEPs, normalised MEPs (normalised to maximal M-wave), and silent period. MEP data were recorded at the plateau of MVCs, and at 3 different time points from EMG on-set during explosive contractions: early (45 ms), middle (115 ms), and late (190 ms)

Discussion

This is the first study to investigate the absolute (ICC) and relative (CV) test-retest reliability of TMS responses (MEP amplitude and silent period duration) recorded at different time points

(early, middle, and late) during explosive voluntary contractions, and at the plateau of MVCs. In

general, no one condition (explosive early, middle, late, or MVC plateau) produced consistently more

reliable (absolute or relative) TMS responses across the different averaging methods than another.

In all conditions except the explosive late phase, both absolute and relative reliability of MEP

amplitude (absolute and normalised) generally increased with an increase in the number of

consecutive stimuli (separate contractions) averaged. In contrast, increasing the number of

averaged stimuli did not improve silent period reliability in any condition. This study also found

averaging the best 3 out of the first 10 contractions based on EMG amplitude or torque output,

produced comparable, albeit slightly poorer, reliability than averaging the same 10 contractions for

all TMS responses and conditions.

Reliability during MVCs

To date, only a handful of studies have investigated the test-retest reliability of MEPs evoked

at the MVC plateau, and these have all been based on measures averaged across 3-5 MVCs (Kamen,

2004; Mileva, Sumners and Bowtell, 2012; Souron *et al.*, 2016). For comparison within this section,

we will focus on our findings within this range of 3-5 contractions, with the influence of higher

numbers of contractions being discussed later.

Within 3-5 MVCs, we observed moderate ICC (0.50-0.75) for absolute and normalised MEP

amplitudes, which is within the range (0.47-0.79) of what others have observed (Kamen, 2004;

Sidhu, Bentley and Carroll, 2009; Mileva, Sumners and Bowtell, 2012; Souron *et al.*, 2016; Table 5 in

supplementary material). We also observed moderate ICC for silent periods at MVC plateau (0.66-

0.70) which was comparable to what has previously been reported for the rectus femoris (0.70;

Mileva, Sumners and Bowtell, 2012; 0.61-70; Di Virgilio *et al.*, 2022) and lower than reported for the

tibialis anterior (0.93-0.95; Souron *et al.*, 2016; Table 5 in supplementary material). For test-retest

CV, within 3-5 MVCs, we observed 11-16% values for absolute and normalised MEP amplitude and

below 5% for the silent period. One previous study has reported test-retest CV for TMS responses at

MVC plateau in the tibialis anterior (Souron *et al.*, 2016) and reported slightly lower CVs for MEP

amplitude (10-11%) and a slightly higher CV for silent period (9%), compared to our results for the

quadriceps. A more detailed comparison of our results and those of previous studies is difficult

because of the disparity in methods for factors such as the type of ICC used, muscles involved, TMS

intensity used, and method of calculating silent period. Nevertheless, based on our results and

those of previous studies, averaging data across only 3-5 MVCs should provide moderate ICCs and

CVs<16% for MEP amplitudes and silent periods.

Independently of the number of averaged MEPs, absolute MEP amplitude (V) at MVC

plateau showed consistently better ICC values than normalised MEP amplitude (% M_{max}), and this

pattern was consistent for other conditions also. In contrast, a previous study of the tibialis anterior

(Souron *et al.*, 2016) found ICC for MEP amplitude at the MVC plateau to be similar between

absolute and normalised data. This difference between our study and Souron et al. (2016) may be

due to the different muscle groups used, although there were other methodological differences

(e.g., process for determining coil position and intensity), again making a direct comparison difficult.

The greater ICC in absolute compared to normalised MEP amplitude that we observed might

originate from the ICC calculation itself. ICC scores will be influenced by inter-participant variability,

with greater variability likely contributing to greater ICCs (Koo and Li, 2016). Normalising the MEP

amplitudes to M_{max} will control for some of the factors that cause inter-participant variability, such

as muscle size and adiposity (Besomi *et al.*, 2020), which in turn might reduce ICCs. In contrast to

ICC, CV measurements are not influenced by inter-participant variability which may explain why CVs

were generally similar for absolute and normalised MEP amplitudes.

Reliability during explosive contractions

To our knowledge, no prior study has investigated the reliability of TMS responses during

explosive contractions. We had suspected the reliability of TMS responses to be lowest in early-

phase explosive contractions and highest at the MVC plateau, as this is commonly observed for

torque and EMG amplitudes in these different conditions (Folland, Buckthorpe and Hannah, 2014;

Tillin and Folland, 2014). However, our findings indicated the test-retest reliability of TMS responses

was similar for the different conditions, an observation that appeared unrelated to the number of

contractions averaged. Just prior to the onset of rapid contractions, in a period of time that overlaps

with the early phase of EMG measurements in the current study, there is an increase in subthreshold

cortical excitability and increased probability in evoking a detectable MEP (Mackinnon and Rothwell,

2000). Early subthreshold cortical excitability changes consistently during explosive contractions,

but the probability of conversion to detectable voluntary motor output in surface EMG is less

reliable (Folland, Buckthorpe and Hannah, 2014; Tillin and Folland, 2014). At later phases in the

contraction, corticospinal excitability and voluntary motor output variability align more closely,

explaining why early phase EMG and torque measurements are less reliable compared to MEP

amplitudes, which remain consistent across phases.

The only exception to there being no clear differences in reliability between conditions was

the observation that CV for the silent period was generally highest for the early phase and lowest for

the MVC plateau (Fig. 2F). This may be associated with the lower overall length of the silent period

in the early phase compared to the MVC plateau (Table 1). Given the CV is a measure of variability

relative to the mean, a condition with a lower mean silent period (as is the case for the explosive

early phase), will be susceptible to higher CV values where variability is constant. The shorter silent

period in the early phase may be due to neural drive in this phase being predominantly feedforward

(Škarabot *et al.*, 2021) and less affected by afferent feedback or intracortical circuitry causing

inhibition.

Number of averaged MEPs

For absolute and normalised MEPs, ICCs generally increased, and CVs decreased, with an

increasing number of contractions, in all conditions except the late phase. This is in agreement with

studies on MEP amplitudes recorded at rest or at submaximal force, which also observed improved

reliability by increasing the number of averaged stimuli (Goldsworthy, Hordacre and Ridding, 2016;

Biabani et al., 2018; Brownstein et al., 2018). Our results show that the conditions in our study,

except for the late-explosive phase, require 11-15 contractions (absolute MEP amplitude) and 18-23

contractions (normalised MEP amplitude) to ensure both good ICCs (≥0.75) and CVs≤10%. Further,

although caution is required when extrapolating the logarithmic functions reported in this study well

beyond the number of contractions measured, the shape of the logarithmic functions indicates little

benefit to reliability will be gained by sampling beyond the abovementioned ranges. Variability in

the MEP responses in our study are likely affected by the TMS intensity we used (140% AMT), which

evokes MEPs in the quadriceps half-way up the impulse-response relationship for the quadriceps

(Temesi, Ly and Millet, 2017). As this is the steepest section of the impulse-response relationship,

small changes in corticospinal excitability can greatly affect MEP amplitude (Temesi, Ly and Millet,

2017). Combined with the intrinsic fluctuations in ongoing oscillatory activity within the cortical area

(Hordacre et al., 2017), this likely explains the inherent variability in MEP amplitudes, which seems

to limit test-retest reliability, even with consecutive averaging. Nevertheless, based on our results,

we recommend future investigators consider collecting and averaging MEP amplitudes across more

than the standard 3-5 MVCs (or explosive contractions) used in previous studies, if this can be

managed within their protocol, with an optimal range likely falls between 10-20.

Unlike the other conditions, MEP amplitude reliability in the late-explosive phase (190 ms)

was not improved by increasing the number of averaged contractions. This implies that variability in

MEP amplitude during the late phase of explosive contraction remains constant and within a narrow

range from contraction to contraction, unlike during the earlier phases. We can only speculate

about the physiological factors explaining this observation. However, potentially afferent feedback,

which will be greater in the later than earlier phases of explosive contraction (Škarabot *et al.*, 2021),

inhibits the neural drive to muscle and so maybe limits the range of possible MEP amplitudes in the

late phase of explosive contraction.

In the silent period, ICC and CV remained constant despite increases in the number of

averaged stimuli. Thus, variability in the silent period appears to be consistent and remain within a

narrow range of possibilities from contraction to contraction. Factors that are likely contributing to

the variability in silent period include the intensity of stimulation – with lower intensities resulting in

higher variability (Damron *et al.*, 2008) – and differences in motor threshold compared to MEP as

changes in response curve reduce variability (Wassermann *et al.*, 1993). Our results, obtained at

140% AMT, show that collecting data or more than 3 explosive contractions or MVCs will nt provide

meaningful improvements in the reliability of silent period measurament.

In studies on explosive contractions, RFD is often averaged from the best 3 out of as many as

10 or more contractions (Maffiuletti *et al.*, 2016). Thus, if intending to relate TMS responses

measured during explosive contractions to RFD, it may be pertinent to also average the TMS

responses in the best 3 contractions. Of the two methods we used for determining the best 3 (out

of the first 10) contractions (torque and EMG methods), neither showed consistently better

reliability than the other across all conditions and variables. However, both methods for all

conditions provided comparable ICCs and CVs to the average of the first 10 contractions (Tables 3

and 4). It therefore appears that selecting the best 3 contractions to average MEP amplitudes

minimises the influence of MEP variability to a similar extent as increasing the number of averaged

stimuli over consecutive contractions from 3 to 10. Thus, averaging the best three contractions may

be a viable option for investigators in the future, particularly if looking to relate MEP amplitude

measurements to RTD.

Conclusion

Regardless of the averaging method or reliability metric (ICC and CV), no specific condition

produced consistently more reliable MEP amplitudes (absolute or normalised), and silent periods

than another. Generally, increasing the number of averaged consecutive contractions increased the

reliability for MEPs amplitude (absolute and normalised) in all conditions except for late explosive

phase, but did not improve the reliability of silent period duration. Thus, it is advisable to collect

more than the typical 3-5 contractions, as used by previous studies involving TMS during MVC, to

maximise reliability for MEP amplitude; however, 3-5 contractions will suffice for silent period

measurements. Alternatively, investigators could opt to average the best 3 contractions based on

those with the highest torque or EMG prior to MEP, which provides comparable reliability to

averaging the first 10 contractions.

Bibliography

Ansdell, P. *et al.* (2019) 'Menstrual cycle-associated modulations in neuromuscular function and

fatigability of the knee extensors in eumenorrheic women', *Journal of Applied Physiology*, 126(6), pp.

1701–1712. doi: 10.1152/japplphysiol.01041.2018.

Balady, G. J. *et al.* (1998) 'Recommendations for cardiovascular screening, staffing, writing group',

Medicine and Science in Sports and Exercise1, 30(6), pp. 1009–1018. Available at:

http://journals.lww.com/acsm-

msse/Fulltext/1998/06000/AHA_ACSM_Joint_Position_Statement__Recommendations.34.aspx.

Behrens, M. *et al.* (2015) 'Caffeine-induced increase in voluntary activation and strength of the

quadriceps muscle during isometric, concentric and eccentric contractions', *Scientific Reports*,

5(October 2014), pp. 1–10. doi: 10.1038/srep10209.

Besomi, M. *et al.* (2020) 'Consensus for experimental design in electromyography (CEDE) project:

Amplitude normalisation matrix', *Journal of Electromyography and Kinesiology*, 53(June), p. 102438.

doi: 10.1016/j.jelekin.2020.102438.

Biabani, M. *et al.* (2018) 'The minimal number of TMS trials required for the reliable assessment of

corticospinal excitability, short interval intracortical inhibition, and intracortical facilitation',

Neuroscience Letters, 674(March), pp. 94–100. doi: 10.1016/j.neulet.2018.03.026.

Brownstein, C. G. *et al.* (2018) 'An optimal protocol for measurement of corticospinal excitability,

short intracortical inhibition and intracortical facilitation in the rectus femoris', *Journal of the*

Neurological Sciences, 394(June), pp. 45–56. doi: 10.1016/j.jns.2018.09.001.

Buckthorpe, M. W. *et al.* (2012) 'Reliability of neuromuscular measurements during explosive

isometric contractions, with special reference to electromyography normalisation techniques',

Muscle and Nerve, 46(4), pp. 566–576. doi: 10.1002/mus.23322.

Cossich, V. and Maffiuletti, N. A. (2020) 'Early vs. late rate of torque development: Relation with

maximal strength and influencing factors', *Journal of Electromyography and Kinesiology*,

55(October), p. 102486. doi: 10.1016/j.jelekin.2020.102486.

Damron, L. A. *et al.* (2008) 'Quantification of the corticospinal silent period evoked via transcranial

magnetic stimulation', *Journal of Neuroscience Methods*, 173(1), pp. 121–128. doi:

10.1016/j.jneumeth.2008.06.001.

Folland, J. P., Buckthorpe, M. W. and Hannah, R. (2014) 'Human capacity for explosive force

production: Neural and contractile determinants', *Scandinavian Journal of Medicine & Science in*

Sports, 24(6), pp. 894–906. doi: 10.1111/sms.12131.

Goldsworthy, M. R., Hordacre, B. and Ridding, M. C. (2016) 'Minimum number of trials required for

within- and between-session reliability of TMS measures of corticospinal excitability', *Neuroscience*,

320, pp. 205–209. doi: 10.1016/j.neuroscience.2016.02.012.

Groppa, S. *et al.* (2012) 'A practical guide to diagnostic transcranial magnetic stimulation: Report of

an IFCN committee', *Clinical Neurophysiology*, 123(5), pp. 858–882. doi:

10.1016/j.clinph.2012.01.010.

Hermens, H. J. *et al.* (1999) 'European Recommendations for Surface ElectroMyoGraphy', *Roessingh*

Research and Development, pp. 8–11.

Hordacre, B. *et al.* (2017) 'Variability in neural excitability and plasticity induction in the human

cortex: A brain stimulation study', *Brain Stimulation*, 10(3), pp. 588–595. doi:

10.1016/j.brs.2016.12.001.

de Jonge, X. J., Thompson, B. and Han, A. (2019) *Methodological Recommendations for Menstrual*

Cycle Research in Sports and Exercise, *Medicine & Science in Sports & Exercise*. doi:

10.1249/MSS.0000000000002073.

Kamen, G. (2004) 'Reliability of motor-evoked during resting and active contraction conditions',

Medicine and Science in Sports and Exercise, 36(9), pp. 1574–1579. doi:

10.1249/01.MSS.0000139804.02576.6A.

Koo, T. K. and Li, M. Y. (2016) 'A Guideline of Selecting and Reporting Intraclass Correlation

Coefficients for Reliability Research', *Journal of Chiropractic Medicine*, 15(2), pp. 155–163. doi:

10.1016/j.jcm.2016.02.012.

Lewis, G. N., Signal, N. and Taylor, D. (2013) 'Reliability of lower limb motor evoked potentials in

stroke and healthy populations: How many responses are needed?', *Clinical Neurophysiology*,

125(4), pp. 748–754. doi: 10.1016/j.clinph.2013.07.029.

Luc, B. A. *et al.* (2014) 'Reliability of Corticomotor Excitability in Leg and Thigh Musculature at 14 and

28 Days', *Journal of Sport Rehabilitation*, 23(4), pp. 330–338. doi: 10.1123/JSR.2013-0069.

Mackinnon, C. D. and Rothwell, J. C. (2000) 'Padrão Trifásico - Time_varying changes in corticospinal

excitability', pp. 633–645.

Maffiuletti, N. A. *et al.* (2016) 'Rate of force development: physiological and methodological

considerations', *European Journal of Applied Physiology*, 116(6), pp. 1091–1116. doi:

10.1007/s00421-016-3346-6.

McKay, A. K. A. *et al.* (2022) 'Defining Training and Performance Caliber: A Participant Classification

Framework', *International Journal of Sports Physiology and Performance*, 17(2), pp. 317–331. doi:

10.1123/ijspp.2021-0451.

Mileva, K. N., Sumners, D. P. and Bowtell, J. L. (2012) 'Decline in voluntary activation contributes to

reduced maximal performance of fatigued human lower limb muscles', *European Journal of Applied

Physiology*, 112(12), pp. 3959–3970. doi: 10.1007/s00421-012-2381-1.

Morales-Artacho, A. J. *et al.* (2018) 'Associations of the Force-velocity Profile with Isometric Strength

and Neuromuscular Factors', *International Journal of Sports Medicine*, 39(13), pp. 984–994. doi:

10.1055/a-0644-3742.

Rossi, S. *et al.* (2011) 'Screening questionnaire before TMS: An update', *Clinical Neurophysiology*,

122(8), p. 1686. doi: 10.1016/j.clinph.2010.12.037.

Rossi, S. *et al.* (2020) 'Safety and recommendations for TMS use in healthy subjects and patient

populations, with updates on training, ethical and regulatory issues: Expert Guidelines', *Clinical

Neurophysiology*, 132(1), pp. 269–306. doi: 10.1016/j.clinph.2020.10.003.

Rossini, P. M. *et al.* (2015) 'Non-invasive electrical and magnetic stimulation of the brain, spinal cord,

roots and peripheral nerves: Basic principles and procedures for routine clinical and research

application: An updated report from an I.F.C.N. Committee', *Clinical Neurophysiology*, 126(6), pp.

1071–1107. doi: 10.1016/j.clinph.2015.02.001.

De Ruiter, C. J. *et al.* (2004) 'Initial phase of maximal voluntary and electrically stimulated knee

extension torque development at different knee angles', *Journal of Applied Physiology*, 97(5), pp.

1693–1701. doi: 10.1152/japplphysiol.00230.2004.

Säisänen, L. *et al.* (2008) 'Factors influencing cortical silent period: Optimised stimulus location,

intensity and muscle contraction', *Journal of Neuroscience Methods*, 169(1), pp. 231–238. doi:

10.1016/j.jneumeth.2007.12.005.

Sidhu, S. K., Bentley, D. J. and Carroll, T. J. (2009) 'Cortical voluntary activation of the human knee

extensors can be reliably estimated using transcranial magnetic stimulation', *Muscle and Nerve*,

39(2), pp. 186–196. doi: 10.1002/mus.21064.

Škarabot, J. *et al.* (2019) 'Electrical stimulation of human corticospinal axons at the level of the

lumbar spinal segments', *European Journal of Neuroscience*, 49(10), pp. 1254–1267. doi:

10.1111/ejn.14321.

Škarabot, J. *et al.* (2021) 'The knowns and unknowns of neural adaptations to resistance training',

European Journal of Applied Physiology, pp. 675–685. doi: 10.1007/s00421-020-04567-3.

Škarabot, J. *et al.* (2023) 'Decoding firings of a large population of human motor units from

high-density surface electromyogram in response to transcranial magnetic stimulation', *The Journal*

of Physiology, 601(10), pp. 1719–1744. doi: 10.1113/JP284043.

Souron, R. *et al.* (2016) 'Reliability of the functional measures of the corticospinal pathways to

dorsiflexor muscles during maximal voluntary contractions', *Journal of the Neurological Sciences*,

369(October), pp. 368–374. doi: 10.1016/j.jns.2016.09.003.

Sundstrup, E. *et al.* (2010) 'Muscle function and postural balance in lifelong trained male footballers

compared with sedentary elderly men and youngsters.', *Scandinavian journal of medicine & science*

in sports, 20 Suppl 1, pp. 90–97. doi: 10.1111/j.1600-0838.2010.01092.x.

Tallent, J. *et al.* (2017) 'Enhanced corticospinal excitability and volitional drive in response to

shortening and lengthening strength training and changes following detraining', *Frontiers in*

Physiology, 8(FEB). doi: 10.3389/fphys.2017.00057.

Temesi, J., Ly, S. N. and Millet, G. Y. (2017) 'Reliability of single- and paired-pulse transcranial

magnetic stimulation for the assessment of knee extensor muscle function', *Journal of the*

Neurological Sciences, 375, pp. 442–449. doi: 10.1016/j.jns.2017.02.037.

Tillin, N. A. *et al.* (2010) 'Neuromuscular performance of explosive power athletes versus untrained

individuals', *Medicine and Science in Sports and Exercise*, 42(4), pp. 781–790. doi:

10.1249/MSS.0b013e3181be9c7e.

Tillin, N. A. and Folland, J. P. (2014) 'Maximal and explosive strength training elicit distinct

neuromuscular adaptations, specific to the training stimulus', *European Journal of Applied*

Physiology, 114(2), pp. 365–374. doi: 10.1007/s00421-013-2781-x.

Tillin, N. A., Pain, M. T. G. and Folland, J. (2013) 'Explosive force production during isometric squats

correlates with athletic performance in rugby union players', *Journal of Sports Sciences*, 31(1), pp.

66–76. doi: 10.1080/02640414.2012.720704.

Tillin, N. A., Pain, M. T. G. and Folland, J. P. (2012) 'Short-term training for explosive strength causes

neural and mechanical adaptations', *Experimental Physiology*, 97(5), pp. 630–641. doi:

10.1113/expphysiol.2011.063040.

Tillin, N. A., Pain, M. T. G. and Folland, J. P. (2018) 'Contraction speed and type influences rapid

utilisation of available muscle force: Neural and contractile mechanisms', *Journal of Experimental*

Biology, 221(24). doi: 10.1242/jeb.193367.

Todd, G., Taylor, J. L. and Gandevia, S. C. (2016) 'Measurement of voluntary activation based on

transcranial magnetic stimulation over the motor cortex', *Journal of Applied Physiology*, 121(3), pp.

678–686. doi: 10.1152/japplphysiol.00293.2016.

Del Vecchio, A. *et al.* (2019) 'You are as fast as your motor neurons: speed of recruitment and

maximal discharge of motor neurons determine the maximal rate of force development in humans',

Journal of Physiology, 597(9), pp. 2445–2456. doi: 10.1113/JP277396.

Di Virgilio, T. G. *et al.* (2022) 'The Reliability of Transcranial Magnetic Stimulation-Derived

Corticomotor Inhibition as a Brain Health Evaluation Tool in Soccer Players', *Sports Medicine - Open*,

8(1). doi: 10.1186/s40798-021-00399-3.

Wassermann, E. M. *et al.* (1993) 'Topography of the inhibitory and excitatory responses to

transcranial magnetic stimulation in a hand muscle', *Electroencephalography and Clinical*

Neurophysiology/ Evoked Potentials, 89(6), pp. 424–433. doi: 10.1016/0168-5597(93)90116-7.

Wheaton, L. A. *et al.* (2009) 'Reliability of TMS motor evoked potentials in quadriceps of subjects

with chronic hemiparesis after stroke', *Journal of the Neurological Sciences*, 276(1–2), pp. 115–117.

doi: 10.1016/j.jns.2008.09.012.

Fig 1

Fig 2

Fig 3

Fig 4