Arabidopsis natural variation induces complex transcriptomic heterochronies at the floral transition and perturbs the coupling between leaf and flower development by an unexpected genetic control

Dieudonné Sana[1], Kristianingsih Ruth[3], Laine Stéphanie[1], Jesson Béline[2], Vidal Véronique[2], Wells Rachel[4], Morris Richard[3] and Besnard Fabrice[1]

[1]*Laboratoire Reproduction et Développement des Plantes, Univ Lyon, ENS de Lyon, CNRS, INRAe, Lyon, France;* [2]*Helixio, Clermont-Ferrand, France;* [3]*Department of Computational and Systems Biology, John Innes Centre, Norwich, United Kingdom and* [4]*Department of Crop Genetics, John Innes Centre, Norwich, United Kingdom.*

correspondance: fabrice.besnard@ens-lyon.fr

This manuscript contains 6 main figures, 14 supporting figures and 11 supporting tables.

ABSTRACT

Plant development may be viewed as a sequence of tightly orchestrated events in space and time. The coordination of developmental stages gives rise to various organs, such as leaves or flowers. For instance, in *Arabidopsis thaliana*, leaf/bract development ceases once the first flower emerges. Unravelling the mechanisms of this regulation is, therefore, key for understanding how developmental programs are coordinated during the floral transition. In this study, we take advantage of a previously overlooked natural variation that desynchronizes bract repression from the initiation of the first flowers. We found that prolonged bract formation after the floral transition depends on complex genetic interactions, involving at least four loci. Surprisingly, none of these loci contained the floral identity genes previously implicated in bract repression, suggesting that other genes are involved in the synchronisation of bract and flower programs. Furthermore, although ectopic bracts in inflorescences have been interpreted as a prolonged vegetative state, time-series transcriptomics and curve registration revealed a more complex scenario whereby many genes desynchronize in terms of expression between these developmental programs. As a consequence, transcriptional differences peak at the floral transition when bract development is prolonged, affecting a wide variety of biological processes not necessarily associated with vegetative state. A transient increase in transcriptome divergence has been proposed to account for morphological variation between species in animals and plants under the "inverse hourglass" model. Our results suggest that such a model could also explain the sensitivity of certain developmental transitions to phenotypic variation within species, as reported here for the floral transition.

5–8 key words

Introduction

The development of living organisms generally proceeds through distinct temporal phases. As

illustrated by insects developing from embryo to adult through several larval stages and

metamorphosis, development is often described as stepwise ontogenetic sequences (Baressi

and Gilbert 2023). Some transitions are sharp, requiring the precise coordination of different

organs' developmental programs to generate new structures and ultimately form the organism

(Furlow and Neff 2006; Truman 2019; Yamaguchi 2021). At the molecular level, these

transitions involve highly dynamic changes in gene expression. Changes in the relative timing of

these developmental events, or heterochrony, are recognized as an important drivers of

morphological evolution (Alberch et al. 1979; Keyte and Smith 2014). Hence, understanding

how developmental timing is controlled from gene expression to organ formation is key to

understanding both phenotypic robustness and evolution.

Flowering is a major developmental and morphological transition, transforming plant

architecture. In most species, flowers emerge along stems from previous flowerless parts,

without any intermediate floral forms. Very often, the start of flower production is also phased

with other phenotypic changes. The floral transition thus provides an excellent developmental

system to investigate the mechanisms of phenotypic robustness, involving precise spatio-

temporal coordination events.

In several species, flower production coincides with the cessation of leaf production, resulting

in flower-bearing stems devoid of leaves. In botany, a leaf associated with a single flower is

called a bract (Penin 2008; Prenner et al. 2009). A bractless (i.e. leafless) inflorescence is a

typical trait of the *Brassicaceae* family, which includes the model plant *Arabidopsis thaliana*

(Penin 2008). The presence of bracts in angiosperm inflorescences varies widely among species,

showing that the coordination of the leaf program with reproductive development has been

independently and recurrently modified during evolution (Mach 2010).

In this study, we use the floral transition in *A. thaliana* as a model system to study how two

developmental programs, leaf and flower, are coordinated in a short time window. For

instance, several studies have explored how leaf development is halted precisely when the first

flower is formed in this species.

First, morphological deformations in the inflorescence meristems (Kwiatkowska 2006) and
localised gene repression (e.g., SHOOT MERISTEMLESS [STM]) or activation (e.g.,
AINTEGUMENTA [ANT] or *FILAMENTOUS FLOWER [FIL]*) indicate that a bract domain forms
before each flower but fails to outgrow (Long and Barton 2000; Dinneny et al. 2004). This
domain is later incorporated in floral tissues, remaining cryptic (Heisler et al. 2005; Goldshmidt
et al. 2008). Hence, the absence of a bract is due to its early repression when flower
development is initiated.

Second, several genes have been implicated in bract repression through identification of
mutants that restore bract formation. These genes, mostly transcriptional regulators, are all
functionally associated with the floral meristem identity (FMI) pathway (Siriwardana and Lamb
2012), including the central flower regulator *LEAFY* (*LFY*) (Schultz and Haughn 1991; Weigel et
al. 1992). The FMI pathway specifies floral rather than shoot identity (Weigel 1995) and its
impairment leads to flower-to-shoot conversions by restoring vegetative characteristics into
flowers (altered floral organ identity and phyllotaxis, internode elongation, indeterminacy),
including bract outgrowth (see **Table S1**). Furthermore, the genetic ablation of flowers induces
bract outgrowth (Nilsson et al. 1998). This suggests that many genes within the FMI regulatory
network redundantly act as bract repressors. In all these studies, bract de-repression is
consistently accompanied by pleiotropic phenotypes affecting flowers, branching or flowering
time.

Conversely, only few genes have been identified as promoting bract outgrowth, such as *JAGGED*
(Dinneny et al. 2004; Ohno et al. 2004), *AINTEGUMENTA* (*ANT*) and *AINTEGUMENTA-LIKE6*
(*AIL6*) (Manuela and Xu 2024 Mar 28) and their misexpression also produce pleiotropic
phenotypes in addition to bract development. Bracts can also be induced in wild-type *A.*
thaliana under light treatments enriched in far red light, together with flower-shoot chimeras
(Hempel and Feldman 1995; Hempel et al. 1998). The authors proposed a "conversion" model,
where a young branch meristem (already subtended by a leaf) can be converted (here upon
photo-induction) into a flower that inherits this pre-formed leaf, interpreted then as a bract.
However, the underlying genetic mechanisms also remain unclear.

In summary, the current literature suggests a straightforward explanation for the precise

coupling of bract repression with formation of the first flower in *A. thaliana:* this arises from the

bract-repressing activity of the FMI pathway acting in the flower. The molecular mechanisms

involved from flower to bract remain unclear but they are likely indirect and/or non-cell

autonomous, as FMI genes are mostly expressed in the developing flower and not in the bract

domain. Since bracts are almost identical to leaves produced during the preceding vegetative

phase (Dinneny et al. 2004), de-repressed bracts have sometimes been interpreted as the

persistence of a vegetative program (Hempel et al. 1998). Bract de-repression and improper

establishment of flower identity are therefore considered as two sides of the same coin.

In this study, we provide new data showing that, in natural populations, bract repression and

flower development can be uncoupled. We report the transient presence of bracts with wild-

type flowers precisely at the floral transition in wild *A. thaliana* accessions. By combining

precise phenotype characterization, quantitative genetics, genomics, and time-series

transcriptomics specifically in meristem tissues, we investigated this change in the timing of

flower and leaf formation at developmental, genetic and transcriptomic levels. Altogether, our

results provide two new main contributions to our understanding of the mechanisms

coordinating leaf and flower development. First, bract development can be prolonged without

affecting the main FMI pathway genes. Second, this prolonged bract formation at the floral

transition does not correspond to a global "juvenile" shift in the transcriptome and correlates

with more complex transcriptomic heterochronies. In line with the 'inverse hourglass model,'

we discuss how transcriptional divergence, resulting from the ongoing adaptation of flowering

timing, may lead to phenotypic variation at different evolutionary scales.

Results

Bract repression is not strictly synchronised with the floral transition in the natural

***accession* Tsu-0**

Like in *A. thaliana* and its reference accession (*Col-0*) (**Fig. 1A**), the typical ebracteate flower (i.e.

without bracts) of *Brassicaceae* generally appears as soon as the first flower is produced,

indicating synchronization of bract repression with the onset of the flower program. However,

in a number of *Brassicaceae* species, bract repression is delayed for several rounds of flower

formation, generating bracteate flowers (i.e. with bracts) at the base of their flowering branch

(a raceme) (Endress 2006; German et al. 2023): some species display bracts up to the first half

of the raceme (e.g. **Fig. S1A,C**), while others only have one or two bracts at the base (**Fig. S1B,D**

or e.g. in Al-Shehbaz 2015).

Likewise, we discovered that the natural *Tsu-0* accession in *A. thaliana* displays bracts on the

first one to five flowers of the raceme (**Fig. 1A,B**), in the main stem and cauline branches at the

same frequency (**Fig. 1C**). A gradient of bract outgrowth can be observed in *Tsu-0*, from fully

developed bracts resembling the cauline leaves associated to branches, up to filamentous,

short rudimentary structures (**Fig. 1D,E**). Scanning electron microscopy (SEM) showed that a

mild swelling of the peduncle basis could sometimes occur in *Tsu-0*, unlike in *Col-0* (**Fig. 1F**).

Both bracteate and ebracteate *Tsu-0* flowers were normal, containing the invariant organ

number typical of each whorls. Thus, developmental variation is restricted to the bract (**Fig.**

1G). To exclude that these leaves could be secondary outgrowth from mature floral peduncles,

we imaged the emergence of the first flower by SEM. In *Col-0*, the last lateral meristem

subtended by a cauline leaf was immediately followed by ebracteate flowers (**Fig. 1H**). In *Tsu-0*,

the first flowers produced after the last branch displayed bracts or rudimentary bracts (**Fig. 1I**),

recapitulating the gradient of bract forms observed in mature stems. Their abaxial position and

their precocious development also suggest that these structures developed from the cryptic

bract, that developed despite the onset of flower formation.

Since *Tsu-0* is a wild accession, we wondered whether a 'conversion' model as proposed by

Hempel and Feldman (1995; 1998) for other wild accessions could explain this phenotype.

However, phenotyping hundreds of plants did not identify any shoot-flowers chimaeras (**Fig.**

1D,F,H). Also, *Tsu-0* plants did not exhibit the negative correlation between the numbers of

bracts and cauline branches (**Fig. 2A**) that would be expected under the "conversion" model.

This suggests that *Tsu-0* bracts are not produced by shoot-to-flower conversions.

To summarize, in the natural accession *Tsu-0*, bract repression is no longer synchronized with

the onset of flower development. Instead, it is transiently altered to varying and generally

decreasing degrees in the first flowers of a branch, resulting in what we have termed 'basal

bracts'.

Desynchronisation of bract repression with the floral transition is common among
Arabidopsis natural accessions

To measure the ability of different genotypes to generate basal bracts, we defined a bract score

(**Fig. 2B**) that takes into account several branches (hence several floral transitions) per plant,

normalizing by the number of scored branches. Similar to *Tsu-0*, other genetically diverse

natural accessions produce basal bracts, albeit at highly variable rates (**Fig. 2C**). We could not

identify any correlation between basal bracts and geographic or genetic origins (**Fig. S2**). The

bract score shows a high level of intra-genotype variability. Despite this variability, there are

clear differences between the basal bracts of different genotypes (**Fig. 2C**). The production of

basal bracts is thus a common phenotypic trait among *A. thaliana* natural accessions from

different geographical origins and varying genetic backgrounds.

Basal bract frequency shows complex correlations with flowering time but is not
affected by variation in photo-induction

We then explored whether variation in the frequency of basal bracts in natural accessions could

be linked to variation in flowering time. Indeed, flowering time is known to be highly variable

among natural accessions (Salomé et al. 2011) and depends on both genetic and environmental

factors (Maple et al. 2024). We measured both bract score and flowering time in a panel of five

accessions and observed a global positive correlation across genotypes (**Fig. 2D**). For instance,

Tsu-0 flowers later than *Col-0* in either absolute time (ie., the number of days from

germination) or developmental time (measured by the number of leaves produced),

consistently across culture conditions (**Fig. S3**). This suggests that a longer flowering time could

promote the alteration of synchrony between bract repression and the onset of flower

development. However, within an accession, this correlation can be negative (e.g. in *Ler* and

Tsu-0), suggesting that the relationships between basal bracts and flowering time could be

more complex depending on the genetic backgrounds.

To explore further the effects of varying flowering time in a genetic background, *Col-0* and *Tsu-0*

plants were cultured in different photo-induction regimes that vary the flowering time (see

methods, see **Fig. S3**). The bract score remains remarkably constant across all tested conditions

for both accessions (**Fig. 2E**). This indicates that basal bract formation does not depend either

on photoperiod or on the absolute flowering time and rather suggests the importance of

intrinsic genetic determinism specific to each accession.

Delayed bract repression is controlled by multiple QTLs that do not contain genes previously associated to bract development

To analyze the genetics of basal bract formation, we conducted quantitative genetics using

crosses between *Tsu-0* and *Col-0* since these two accessions ranked respectively as high and

low bract producers (**Fig. 2C**). F_1 progeny showed intermediate bract scores while F_2 bract

scores spread within the parental range, with a distribution that does not indicate a simple

monogenic trait (**Fig. 3A and Fig. S4A**). Bulk Segregant Analysis (BSA) with mapping-by-

sequencing identified four quantitative trait loci (QTL): two on chromosome 1 and two on

chromosome 5 (**Fig. S4B,C**). The large spread of F_1 phenotypic values constrained F_2 plant

selection with few plants in the high-score bulk and possibly spurious homozygous in the low-

score bulk. These two limits may explain the large intervals obtained.

To better estimate the QTLs, we also used Recombinant Inbred Lines (RIL), which allowed

averaging of bract scores across isogenic plants. Using an existing RIL set (Simon et al. 2008), we

measured the average bract scores of several lines (**Fig. S5A**) for which we checked the absence

of recombination bias (**Fig. S5B**). The heritability of the bract score was high among RIL (65.45%

+/- 1.7) and a single QTL scanning mapped four peaks: two most significant on chromosome 1

(LOD score > 4 with a p-value < 0.1% named 1a and 1b) and two less significant on chromosome

5 (LOD score > 3, named 5a and 5b; **Fig. 3B**). Consistent with the BSA result, this analysis

provided slightly different QTL positions and shorter intervals. A two-QTL scan suggested

additive effects between QTLs 1a and 1b (**Fig. S5C**). To confirm and reduce the intervals of these

two major QTLs, we used two RILs identified as heterozygous in a region overlapping with these

QTLs at F6 generation (**Fig. 3C-H**). F_8 plants from these RIL were re-genotyped at a higher

density with new genetic markers (**Fig. S6A,B**) to select recombining intervals homozygous for

either allele of the two parents (**Fig. 3D, G** and **Fig. S6F**). Selected plants were used to generate

new lines forming a heterologous inbred family (HIF) with identical genotypes except in a small

interval, which allows testing the effect of alleles in that region only (see **Fig. 3D**). Scoring bract

among HIF confirmed both QTLs and mapped them to smaller intervals named 1a* and 1b*.

Additive effects of QTL 1a and 1b appeared when *Tsu-0* allele in 1b* was combined with *Col-0*

allele in 1a*, the bract score reaching about half of the *Tsu-0* parent bract score (see in 488

families, **Fig. 3G, H**). However, *Tsu-0* parent score is tied only when both QTLs bear *Tsu-0* alleles

(see in 334 families, **Fig. 3D, E**). As a control, we generated a HIF to test the region between 1a

and 1b and demonstrated that it did not influence the bract score (**Fig. S6C-E**).

The final mapped intervals remained large (4.1 to 11.8 cM), containing many genes (332 to 931,

2324 annotated genes for the four QTL, **Fig. 3I** and **Table S2**). The parental accessions that we

re-sequenced differ by over 770,000 small genomic variations (SNPs and indels), with more

than one variation every 175 bp on average, leaving few invariant genes to exclude (**Table S2**).

However, known bract-related genes were absent from the QTLs (**Fig. 3I**), whether those genes

were reported as promoting bract development (e.g. *JAGGED*) or as being involved in the FMI

pathway and acting as bract repressor genes (**Table S1**). Only *LSH1* was found to lie at the

border outside QTL 5a. This absence of major FMI genes in our QTL is consistent with the wild-

type phenotype of flowers associated with bracts in *Tsu-0* (**Fig. 1G**).

To decipher the causal variation responsible for basal bracts, we took advantage of pleiotropic

phenotypes co-segregating with basal bracts in the RIL. The number of cauline branches

positively correlated with the bract score and QTL1b overlaps with a QTL controlling cauline

branch number (**Fig. S7**). This suggests that the gene controlling basal bracts in QTL 1b* might

also promote more cauline branches. Second, "branched siliques" could sometimes be

observed among RILs and HIF at low penetrance (**Fig. S8A,B**). We interpreted this transgressive

phenotype as resulting from an incomplete floral determinism with reiteration of floral bud

formation in whorls, as reported for mutants such as *ap1* (Irish and Sussex 1990). In HIF 488,

this phenotype occurred only when *Tsu-0* QTL 1b* was associated with *Col-0* haplotypes in

other mapped QTLs (**Fig. S8C**). In HIF 334 however, this phenotype was not correlated to a

consistent genetic combination of mapped QTLs (**Fig. S8C**). This transgression suggests that the

maintenance of a wild-type phenotype in *Tsu-0* bracteate flowers relies on genetic interactions

and may interfere with the FMI pathway.

To conclude, our genetic analysis demonstrates that the genetic control of basal bracts in *Tsu-0*

is complex, involving several loci with additive effects on bract development and epistatic

interactions with other traits. It also suggests that the causal genes differ from those previously

identified in mutants with bracts.

Time-series transcriptomics reveal a peak of gene expression divergence at the floral
transition, while FMI genes remain synchronized

Developmental programs are governed by gene regulatory networks, often leading to

substantial transcriptional changes. We therefore sought to identify genes whose expression

changes with bract development. We profiled meristem transcriptomes before, during and

after floral transition in both accessions. Microdissected meristems were matched to the same

four developmental stages: vegetative, late vegetative, transition and floral (V, L, T and F,

respectively, **Fig. 4A**). We performed RNAseq and used principal component analysis (PCA) to

assess the consistency between replicates (**Fig. S9B**).

To monitor the progression from vegetative growth to flowering at the transcript level, we

analysed the expression dynamics of several key regulators of floral transition, such as

FLOWERING LOCUS T (FT) (Wigge et al. 2005), and of flower identity (e.g. *LFY* or *AP1)*. The

synchronized expression of these genes in the two accessions validated the precision of our

morphological staging (**Fig. 4B**): despite different absolute flowering times (**Fig. 4A**; **Fig. S3**; **Fig.**

S9A), our experimental design realigned the two accessions' developmental progression using

the floral transition as a common timer. Hence, gene expressions that desynchronize from

flower formation can easily be tracked. To ensure that our bulk RNAseq data does not obscure

changes in gene expression patterns related to flower identity, we introduced an LFY

transcriptional reporter in both accessions to precisely monitor the activation of this master

floral identity gene in space and time during the floral transition. Live confocal imaging revealed

similar dynamics in both genotypes, with no expression of *LFY* in branches and high expression

levels in all flowers from the first flower onwards (**Fig. 4C**), regardless of bract presence in *Tsu-0*

(**Fig. 4D,E**). These results, in line with our genetic experiments, confirm that alteration of *LFY*

dynamics during floral transition is likely not involved in the delayed repression of bracts. As

previously reported (Chahtane et al. 2018), bracts in *lfy* mutants are not precisely located at the

base of branches (**Fig. S10**), as opposed to *Tsu-0* basal bracts (**Fig. 1B**). This indicates that the

developmental contexts leading to the formation of bract in mutants versus basal bracts in

natural accessions may be really distinct, in line with the involvement of distinct genetic

pathways shown above (**Fig. 3**) .

In the time-course, the stage T should be the most relevant to study bract formation even if no

bract is visible at this stage. Indeed, bract and flower initiations start before any morphological

event (Heisler et al. 2005). The delay between stages T and F (a median of 1 to 2 days, **Fig. S9A**)

is compatible with the initiation of the first flowers and their bracts at stage T. Hence,

transcriptional changes associated with the delay of bract repression should transiently appear

at stage T and progressively fade by stage F (since only the 1 to 5 first flowers present a bract,

Fig. 1). At the global transcriptome scale, the stage T stood out as a critical transcriptomic

transition in both accessions, with the highest number of gene expression changes upon

reaching this stage (**Fig. 4F**). Interestingly, the divergence between the transcriptomes of the

two accessions (as measured by the number of differentially expressed genes -DEG) also peaks

at stage T, with up to 4,759 genes DEG (**Fig. 4G, table S3**), including genes potentially

responsible for bract formation. In conclusion, the relevant time window during which bract

repression is uncoupled from flower development corresponds to a period of highly dynamic

gene expression, both throughout a plant's development and among plants with different

genetic backgrounds.

Transcriptomic data identify candidate genes responsible for the desynchronization of
bract repression
To identify putative genes impacting bract formation among DEGs at stage T, we first cross-

referenced our transcriptomic (4,759 DE genes) and QTL-mapping data (2324 genes in

intervals). We further filtered out genes that had no genetic polymorphisms (including

variations with no putative functional impact, see methods) between *Col-0* and *Tsu-0*. This

selection yielded a list of 362 candidate genes spread over the four QTLs (**Table S4**), including

24 transcription factors, and none of them had previously been associated with bract

formation. This suggests that basal bract formation in a natural accession can be genetically

determined by a novel set of genes likely present in this list.

To determine whether these causal genetic variations influence the expression of genes known

for their role in bract development (**Table S1**) or act through an independent mechanism, we

systematically analyzed the expression dynamics of these genes. Only *PUCHI*, *SOC1*, *TFL1* and

LSH1 significantly differed at stage T between *Tsu-0* and *Col-0* (**Fig. 4H** and **Fig. S9D**). Single

puchi, *soc1* or *lsh1* null mutations alone do not induce basal bracts (Rieu et al. 2024; **Fig.**

S10C,D), so the isolated effect of their reduced expression in *Tsu-0* at stage T (54%, 52% and

41%, respectively) does not likely impact bract development. By transforming lateral branches

into flowers, *tfl1* mutants display basal bracts (**Fig. S10C,D**). But *tfl1* phenotype is recessive: no

basal bract exists on heterozygous TFL1//tfl1 plants (data not shown), so the effect of the

reduced *TFL1* expression at stage T in *Tsu-0* (71%) is again uncertain. To our knowledge

however, combinations of these mutations have not been reported. Moreover, these gene

expression differences persisted in stage F while bracts disappeared with older flowers: hence,

the reduced expression of *PUCHI*, *SOC1*, *LSH1* and *TFL1* is not correlated to bract formation at

all developmental stages. Altogether, this suggests that both the natural genetic determinism of

delayed bract repression and the underlying transcriptional program acting in the meristem are

largely independent of the genetic pathway identified in bract mutants.

Using time-series to identify a transcriptomic signature of leaf/bract development in
shoot meristem
To identify biological functions specific to the critical stage T in *Tsu-0* without prior

assumptions, we performed a GO term analysis on all DEGs compared with *Col-0*. However, the

results were not very informative, notably lacking terms related to leaf development (**Fig.**

S11A,C). Since *Tsu-0* and *Col-0* are distant backgrounds, it is not surprising that many gene

expression differences at stage T are not related to bract formation or even to development

(see the second axis of PCA, **Fig. S9B)**. To specifically isolate the gene regulatory network

responsible for delayed bract repression in *Tsu-0*, and thanks to the information provided by

the time-series, we designed a clustering method to isolate a transcriptomic signature common

to all leaf-producing stages versus non-leaf producing stages. Briefly, expression data were

gathered in two groups corresponding to either bract-less (*Col-0* T, F, and *Tsu-0* F) or leaf/bract-

producing meristems (the other samples, including *Tsu-0* T) and we selected genes whose

expression profile clusters these two groups apart (**Fig. 5C**). These genes can be considered

"bract signature genes", with a positive or inverse correlation with the presence of bract if their

expression is up- or down-regulated (respectively) at stage T in *Tsu-0*.

Only 124 genes met these criteria (**Fig. 5D, Table S5**), including *SOC1*, whose expression anti-

correlates with the bract presence. Interestingly, two enriched GO-terms emerged from this list

of bract signature genes: anthocyanin biosynthesis and salicylic acid (SA), associated with up-

and down-regulated genes, respectively (**Fig. 5E**). Backtracking these ontology terms for all the

4,759 DEGs at stage T (**Table S6** and **S7**) retrieved even more genes associated with these

pathways, supporting that their activity levels differ between *Tsu-0* and *Col-0* at the critical

stage T (**Fig. S11D**). High anthocyanin production in *Tsu-0* at stages L and T was evident from

the frequent purple coloration just below the meristems, contrasting with the green tissues in

Col-0 (**Fig. 5G**). Sometimes, this purple color extended to young organs where bracts were

initiated (**Fig. S11E**). As the stem grew, the pigment receded to the rosette junction in both

accessions (**Fig. 5G**), vanishing from the apical meristem. This trait specific to *Tsu-0* validates

our clustering strategy: it is able to capture genes whose expression differ between *Tsu-0* and

Col-0 at stage T and with overall contrasted levels between vegetative and flower stages, as

expected for a gene regulating bract repression in synchronization with flower development.

Interestingly, among these 124 bract signature genes, 12 also lie in the mapped QTLs (**Fig. 5H**),

including one anthocyanin biosynthetic enzyme (the dihydroflavonol reductase, DFR) and five

SA-responding genes (see **Table S5**). For example, **Figure 5I** shows the expression profiles of

two such genes. However, further work is required to test whether these candidates are

responsible for basal bract formation in *Tsu-0* and whether the anthocyanin and/or SA

pathways are involved in this natural variation.

Curve registration revealed massive gene desynchronization and challenged the
***hypothesis of a "prolonged vegetative state" during floral transition in* Tsu-0**
In the previous clustering approach, both up- and down-regulated genes display similar

temporal dynamics: they tend to be delayed in *Tsu-0* compared to *Col-0* (**Fig. 5I**). This result

suggests that a transcriptomic leaf signature is maintained at the stage T in *Tsu-0* compared to

Col-0. This desynchronization of the expression of some genes with the developmental clock

imposed in our experiment (from V to F stage) identifies transcriptomic heterochronies, i.e.

changes in the timing of gene expression.

To assess the importance of heterochrony in the divergence of *Tsu-0* and *Col-0* transcriptomes,

we first tested whether its effects could be detected in a global unsupervised analysis of gene

expression levels over the time-course. We therefore performed a PCA and plotted the two

main axes of variance after grouping samples by genotype and time points (**Fig. 6A**). While the

second axis of variance (~24.4%) differentiates genotypes, the main axis (47.9% of variance)

orders the sampling time points chronologically in both genotypes, representing a

transcriptomic age. Surprisingly, *Tsu-0* stage T does not align with *Col-0* stage T on this axis: it is

closer to a stage F, explaining the limited expression changes when progressing to the next

stage F (**Fig. 4F**).

As suggested by Calderwood et al. (2021), the transcriptomes of two genotypes during

flowering cannot be aligned to a single developmental time; each gene may desynchronize

differently, sometimes in opposite directions. To quantify each gene desynchronization

between *Col-0* and *Tsu-0*, we used curve registration (Calderwood et al. 2021) that can predict

subtle temporal shifts of gene expression (Kristianingsih, 2024). We analyzed the shift relative

to the floral transition, used as a common reference clock between the two accessions. Thus,

genes with no shift, like *AP1* (**Fig. 4B and Fig. 6B**), can be nevertheless shifted in absolute time,

because *Tsu-0* plants flower later with an older absolute age (**Fig. S3 and S9A**). In our approach,

positive and negative shifts reflect desynchronization relative to the floral transition and gene

expression dynamics may shift earlier or later. Unlike our previous clustering approach (**Fig. 5C**),

this registration method uses a common scaled expression level and computes only temporal

shifts, regardless of absolute expression levels (**Fig. 6B** and **Fig. S12**).

Only a few genes could not be registered (**Fig. 6B,** N = 43), suggesting that most genes follow

similar temporal dynamics in both accessions (see examples **Fig. S9C, D** or **Fig. 4B, H**). Most

genes could be classified into three categories: null, negative and positive shifts (**Fig. 6B**). A null

shift indicates that the gene expression in *Tsu-0* stays "in phase" with flowering, like *AP1*

(compared with **Fig. 4B)**. A positive or negative shift indicates that gene dynamics in *Tsu-0* occur

earlier (*e.g. AG*) or later (*e.g. DFR*), relative to the floral transition, revealing heterochronies

between transcriptomic and phenotypic processes. Transcriptome-wide, the shifts were broadly

distributed, showing massive and complex desynchronisation of transcriptome dynamics (**Fig.**

6C). Hence, the floral transition in *Tsu-0* did not impose a synchronization to the entire

meristematic transcriptome: only ~18% of genes (n=3879) stayed in phase with this process

while the majority showed delayed or advanced expression dynamics. Even more genes shifted

earlier relative to the floral transition, supporting the PCA interpretation (**Fig. 6A**). Since curve

registration computes temporal shifts in gene expression regardless of absolute expression

levels, we investigated whether these shifts could affect differences in gene expression levels.

Indeed, there is a strong correlation (p-value < 2e-16) between the absolute shift value and the

proportion of DEGs between *Tsu-0* and *Col-0* at all developmental stages (**Fig. 6D**). This suggests

that the desynchronization of gene dynamics between the two accessions contributes to

differences in gene expression levels. Furthermore, this effect is significantly higher at the T

stage for positive shift values ranging from 0.25 to 0.75 (**Fig. 6D** and **Fig. S12**): this suggests that

genes with "earlier" dynamics in *Tsu-0* explain at least partially the peak of transcriptome

divergence observed at the T stage (**Fig. 4G**).

In conclusion, the differences in the transcriptomes of *Tsu-0* and *Col-0* are impacted by a

significant level of heterochrony, with complex combinations of temporal shifts. Although the

bract is often considered a 'juvenile' vegetative structure, its formation at the floral transition

does not indicate a global juvenilization at the transcriptomic level.

Investigating which biological processes are affected by desynchronization of gene
expressions in meristems during flowering
As expected, genes in phase were primarily linked to flowering and developmental phase

change as shown by key regulators of these pathways (**Fig. S12B, C**), which also emerges from

GO term enrichment analysis (**Fig. 6D**). In contrast, genes previously associated with bract

mutants exhibit a wide range of shifts, from very early to very late. This shows that they are

variably affected by these desynchronizations (**Fig. S12A, C**), reinforcing the idea that they are

not involved in basal bract formation. GO term analysis for each shift category highlighted the

particular processes desynchronized from flowering in *Tsu-0* compared to *Col-0* (**Fig. 6D**). For

instance, vascular differentiation (two GO terms mentioning tracheary element and secondary

cell wall) emerges as a process shifted earlier in *Tsu-0* relative to the floral transition (**Fig. 6D**).

Vascular differentiation dynamics may be driven by other progression factors like absolute age,

suggesting a loose connection with the flowering pathway. Conversely, processes shifted later

in *Tsu-0* revealed terms related to cell division (spindle, cell cycle, mitosis) and ribosomal

biogenesis (six GO terms mentioning ribosome, rRNA, ribonucleoprotein and protein-RNA

complexes; **Fig. 6D**), which suggests that core meristematic functions are prolonged in *Tsu-0*.

Further work is yet needed to explore whether and how the desynchronization of these

processes relative to floral transition impacts bract de-repression.

Discussion

423   In this study, we investigated the transient formation of 'basal bracts' at the base of

424   inflorescences in natural accessions of the model plant in *A. thaliana*. We characterised basal

425   bract formation at developmental, genetic and transcriptomic levels. We found that this

426   phenotype is associated with a desynchronization of bract repression from flower formation

427   during the floral transition.

429   ***The mechanisms desynchronizing bract repression and flower development***

Evidence from the literature suggests that flower development, through redundant actions of

the genes from the FMI pathway, represses bract development, hence naturally coupling flower

establishment and bract repression. However, natural basal bract demonstrate that these two

processes can be uncoupled in a short time window during the floral transition when both

flowers and bracts develop together.

The wild-type phenotype of basal bract associated flowers (**Fig. 1**) and the proper dynamics of

many FMI genes (e.g. *LFY*, *AP1*, **Fig. 4**) suggest that the transient bract de-repression at floral

transition is not a consequence of an impaired FMI establishment, contrary to previously

reported mutants. In line with this, bracts are rarely produced with the first flowers in mutants

of the FMI pathway (**Fig. S10**). Our quantitative genetic (**Fig. 3**) and transcriptomic data (**Fig. 5**)

further show that the genes causing this transient bract de-repression are different from the

handful of FMI genes whose mutants display bracts.

However, the precise mechanisms involved remain unresolved. Finer mapping or alternative

quantitative genetic approach such as GWAS could help pinpoint causal genes for their definite

validation. However, the four identified QTLs, including two with additive effects (**Fig. 4 and Fig.**

S5), indicate that several complex genetic mechanisms may be at work. Our transcriptomic

analysis established correlations of differentially expressed genes and pathways in the

meristem at the time when bract de-repression occurs: up-regulation of anthocyanin pathway,

down-regulation of salicylic acid pathway (**Fig. 5**) or prolonged activity of ribosome biogenesis

or cell cycle (**Fig. 6**). This however, does not demonstrate their role in promoting bract

development: a transcriptomic signature of leaf development in stage T meristems could simply

be a consequence of the developing bracts rather than its cause.

Interestingly, several plant species evolved leaf-less inflorescences, such as grasses (Poaceae),

through apparently convergent and unrelated mechanisms (Mach 2010). Indeed, *Arabidopsis*

and *Poaceae* mutant bracts are not strictly homologous and so far, different genetic

mechanisms were reported (Whipple et al. 2010; Whipple 2017; Xiao et al. 2022). However,

some of these mechanisms are still partially unresolved (Kawakatsu et al. 2009; Wang et al.

2021). Since in both groups, bract exclusion from wild-type inflorescences involve a leaf

repression mechanism synchronized with a developmental phase change, further research on

these two groups may reveal whether they share more similarities than what is currently

described.

Linking floral transition, transcriptome divergence and developmental robustness

We found that gene expression differs the most between the two accessions at the floral

transition, at the time when bracts are de-repressed in *Tsu-0* (**Fig. 4**). Furthermore, curve

registration identified an important level of gene desynchronization, which likely contributes to

the divergence in gene expression levels, especially at the stage T (**Fig. 6**). Specifically, these

desynchronizations alter the expression timing of many genes relative to a minor gene pool

driving the floral transition, which we used as a reference in our experiments.

As flowering is a key adaptive trait that involves transcriptional reprogramming, the floral

transition may be a developmental stage that leads to large transcriptome differences among

natural populations. Indeed, massive and complex gene desynchronisation during floral

transition were reported at both intra- and inter-species level in *Brassica rapa* cultivars and *A.*

thaliana (Calderwood et al. 2021). This phenomenon may arise from the strong selection

pressure adapting flowering time to the environment (Bloomer and Dean 2017). As a result, the

flowering-related genes could shift and desynchronize from numerous others not functionally

related to flowering, creating a peak of transcriptome divergence in the short time window

when flowering-related genes vary quickly (**Fig. S13**). In line with this, a peak of gene expression

divergence has also been reported in meristems transitioning from vegetative to floral stages

between different species of Solanaceae (Lemmon et al. 2016).

In different unrelated phyla like animals, plants, brown algae or fungi, the "hourglass" (Kalinka

et al. 2010; Quint et al. 2012; Cheng et al. 2015; Lotharukpong et al. 2024) and "inverse

hourglass" (Lemmon et al. 2016; Levin et al. 2016) models propose that reduced, or

respectively increased, transcriptome divergence at specific stages of development drive lower

(respectively higher) phenotypic variations in the final body plans. For instance, the high

transcriptome divergence between *Solanaceae* species during floral transition correlates with

the diversification of their inflorescence architectures (Lemmon et al. 2016). Here, we report a

similar correlation within natural populations of *A. thaliana*: the transcriptome divergence

peaks at floral transition, which correlates with a developmental variation, the formation of

basal bracts. Interestingly, several observations suggest that developmental robustness (i.e. the

ability to produce an invariant phenotype) is reduced during floral transition in *Brassicaceae*, at

both inter- and intra-species levels. In addition to basal bracts (**Fig. S1**), botanists have also

documented examples of frequent flower-to-shoot transformations or flower dimorphisms

(Arber 1931a; Arber 1931b). A possible explanation would be that, as a consequence of the

constant adaptation of flowering time, the many changes in gene expression lead to

desynchronisation of genetic programs occurring during the short floral transition, creating

new, transient gene expression states that fuel developmental variation (**Fig. S13**). More

generally, this questions whether rapid developmental transitions of adaptive traits that change

the timings of events, may result in more gene desynchronization and developmental variation.

To test this hypothesis for the case of bract development, further investigation into the

identification of the genetic and developmental mechanisms that uncouple bract repression

from flower formation in natural *A. thaliana* accessions will be important. Such follow-up

studies may also shed light on how the presence of bracts has frequently evolved within and

between species, both at the base of branches in link with floral transition and in entire

inflorescences as well.

Materials and Methods

Plant growth conditions

Seeds were sown on peaty-clay soil, stratified at 4 °C for at least two days, and watered with

fertiliser (18-10-18 N-P-K) under LED lighting (sunlight spectrum NS12, 150 µmol.m-2.s-1).

Three different day/night regimes were used in the experiments: short-days (SD) with 8 h light

and 16 h dark (non-inductive photo-conditions for flowering); long-days (LD) with 16 h light and

8 h dark and continuous light (CL) with 24 h light (both photo-inductive flowering conditions).

Temperature and humidity are controlled as follows: 22 °C and 60% humidity during light for LD

and CL conditions, and 18 °C and 70% humidity constantly in SD and during night time in LD. For

the Bulk Segregant Analysis and the RNAseq time course, plants were grown 20 days in SD

before switching to LD. For the RIL phenotyping assays, plants have been directly cultured in LD.

Plant materials

Arabidopsis thaliana natural accessions and associated RILs were obtained from the Versailles

Arabidopsis Stock Centre (VASC), *Col-0*: 186AV, *Tsu-0*: 91AV, RIL set name: 3RV. F_1 and F_2 plants

used for the BSA were generated by crossing the parents *Tsu-0* and *Col-0* in both directions. The

following strains were obtained from the NASC and are in *Col-0* background if not otherwise

mentioned: *tfl1-13*: N6237; *tfl1-14*: N6238. *puchi-1, bop1-4* x *bop2-11* and *puchi-1 x bop1-4* x

bop2-11 (Karim et al. 2009), *lfy-12* (Maizel and Weigel 2004) and *soc1-2* (Michaels et al. 2003)

were described previously. Plants expressing pLFY::2xmCherry-N7 ; 35S::Lti6b-YFP (in *Col-0* and

Tsu-0) were generated for this study. *Lobularia maritima* and *Allaria petiolata* plants were

natural specimens collected outside the laboratory, in France. The phylogeny of *Brassicaceae*

tribes used in Fig. S1 is extracted from the Brassibase website (Kiefer et al. 2014).

Plasmid constructions and plant transformation

pLFY::2xmCherry-N7: since no polymorphism was sequenced in the *LFY* promoter between *Col-*

0 and *Tsu-0* accessions, a sequence starting at -2277pb upstream of the ATG was amplified by

PCR   from   *Col-0*   genomic   DNA   using

5'GGGGACAACTTTGTATAGAAAAGTTGATCCATTTTTCGCAAAGG   and

5'GGGGACTGCTTTTTTGTACAAACTTGAATCTATTTTTCTCTCTCTCTATC primers. PCR fragments

were purified and inserted into pDONR P4-P1R with a gateway BP reaction. This plasmid was

then recombined in a three-fragment gateway reaction with 2xmCherry pDONR211 and N7-tag

pDONR P2R-P3 (containing the nuclear tag N7 and a stop codon) and the destination vector

pK7m34GW. 35S::Lti6b-YFP: a pDONR P2R-P3 plasmid containing the sequence of the plasma-

membrane protein Lti6b (Cutler et al. 2000) was recombined in a three-fragment gateway

reaction with a 35S pDONR P4-P1R and the YFP pDONR P2R-P3 into the destination vector

pB7m34GW. Both constructs were then transformed into *Agrobacterium tumefaciens*

C58pMP90 strain by electroporation and transformed into both *Col-0* and *Tsu-0* plants by floral

dipping (Clough and Bent 1998). For each construct-by-genotype combination, several

independent transgenic lines were selected in T_2 for a single insertion event based on 3:1

resistant:sensitive mendelian segregation of the resistance provided by the transgene. The

expression patterns of pLFY were reproducible between selected lines and matched published

data for *Col-0*. We were unable to get a 35S::Lti6b-YFP line with a membrane signal as strong in

Tsu-0 as in *Col-0* (Fig. 2 and S2). However, despite the weak YFP signal, the morphology of the

tissue could still be captured.

Microscopic meristem imaging and image analysis

Meristems were imaged using a Scanning Electron Microscope (Hirox 3000 SEM), or with a

confocal microscope Zeiss 700 LSM, according to the manufacturer's instruction and without

prior fixation. Multitrack sequential acquisitions were performed as follows: YFP, excitation

wavelength (ex): 488 nm, emission wavelength (em): 300–590 nm; mCherry, ex: 555 nm, em:

300–620 nm. Detection wavelengths and laser power were identical for *Col-0* and *Tsu-0*, PMT

voltage was identical for mCherry to allow comparisons of pLFY signal intensity in the two

genotypes. The YFP PMT voltage was optimised on each plant. Confocal images were processed

using imageJ (Schneider et al. 2012): maximum projections of mCherry channel and 16-bit-

transformed standard deviation projections of the YFP channel were merged in a composite

image. mCherry intensities were unchanged while brightness and contrast of the YFP channel

were optimised on each plant to provide the best morphological outlines of the tissues.

Macroscopic plant phenotyping

561   Macroscopic pictures were taken using different devices, according to the manufacturer's

562   instructions: Keyence VHX-900F, Canon EOS 450D camera, camera device of a Samsung Galaxy

563   A10 and a Ulefone Armor X5 pro.

Phenotypic measurements

Basal bract score was determined by counting all bracts in the main stem and cauline branches,

normalising by the total number of branches (excluding rosette branches, **Fig. 1E**); the

inspection was limited to the first five flowers, especially for mutants. The number of bracts

was counted after internodes elongation on the last upper cauline branch to ensure bracts

were visible to the naked eye. Flowering time was measured with different methods

(mentioned in the main text): the number of days from the start of transfer to in growth

chambers to the day of bolting, the day when the first flower blooms (open petals), or as the

cumulated number of leaves (rosette only or rosette and cauline). For plastochron

measurements, several plants of both genotypes were synchronised and grown in the same

condition. Each sampling day after transfer to a long day, 5 to 10 plants were randomly

dissected under a binocular dissecting scope, removing and counting all organs (leaves or

flowers) from the first leaf to the youngest organ visible on the main meristem. The youngest

organs were counted as soon as they were separated by a boundary (corresponding to stage-2

flowers as defined by (Smyth et al. 1990).

DNA extraction and sequencing for Bulk Segregant Analysis (BSA)

For the BSA, an F_2 mapping population of 684 plants was generated by crossing (*Col-0* x *Tsu-0*)

in both directions. The bulk segregant analysis was split into four replicates. A 1cm² leaf sample

was sampled for each F_2 plant, kept at -20°C and genomic DNA was extracted individually if the

plant was selected in one of the bulk. Genomic DNA of 56 and 17 plants were selected and

pooled in the bulks of low and high bract-score plants, respectively. The genome of the parental

lines was re-sequenced using genomic DNA from bulk *Tsu-0* and *Col-0* seedlings, respectively.

All genomic DNAs were extracted and purified using a CTAB-based protocol (cetyl

trimethylammonium bromide), following instructions as in (Healey et al. 2014).

Each DNA bulk was prepared by pooling the genomic DNA of selected plants in equal quantities,

to reach a final concentration between 13 and 25 ng/µl. Pooled DNA bulks and parental DNA

were used to prepare libraries and sequenced on BGISEQ-500WGS according to the

manufacturer's instructions, yielding 5 Gb data of 100bp paired-end reads per library (target

coverage of 40X).

DNA sequencing analysis and genomic variant analysis

Sequencing data consists of two parental plus two bulks of sequencing data. A genomic variant

analysis was performed on each dataset following the workflow of short variant discovery

previously described in (Besnard et al. 2020). This resulted in four gVCF files (one per sample)

generated by the HaplotypeCaller tool of GATK (v3.8, McKenna et al., 2010). TAIR10 was used

as the *A. thaliana* reference genome. Then, the two parental gVCF were first joint-genotyped

using GATK's tool GenotypeGVCFs to emit a common vcf file for the two samples. This file was

used to select a list of specific SNPs and small indels of *Tsu-0* (91AV stock) versus *Col-0* (186AV

stock), filtering for positions with coverage metric DP>10. This reference list of *Tsu-0*

polymorphisms was then used as the --dbsnp option in a second pass of joint genotyping using

all four gVCF as input (two parental samples plus the two bulks) to emit a common vcf file.

Finally, relevant polymorphic positions from the reference list in the two bulks were selected

after filtering for a depth ≥3.

QTL mapping from Bulk Segregant Analysis (BSA)

QTL identification was carried out using the QTLseqr package (Mansfeld and Grumet 2018)

according to user instructions. After filtering data with the following parameters (refAlleleFreq

= 0, minTotalDepth= 10, maxTotalDepth = 90, minSampleDepth = 10, minGQ= 99,

depthDifference = 15), the deltaSNP index (Takagi et al. 2013) was generated and loci that

reached the approximately 95% confidence interval were retained.

QTL mapping using Recombinant Inbred Lines

Genotyped RILs from *Col-0 x Tsu-0* (3RV) are publicly available in the VASC. Based on BSA

results, a panel of 55 RIL were selected from their known genotypes on chromosomes 1 and 5

using GGT 2.0 software (van Berloo 2008). The detailed genotypes of each line used in this

study are available in **Table S9**. The presence of basal bracts was quantified in each line using

the bract score. QTL mapping was performed with R/qtl software according to user instructions

(Broman et al. 2019): we used the scanone function with the mean value of the bract score for

the 55 tested RILs as a trait and significant thresholds were computed by setting a permutation

number to 2000. For the bract score, the "non-parametric" (np) statistical model was used

while for paraclade number, we used a normal model implemented with the hk method. To

look for QTL interactions, the scantwo function was used with the normal model and the hk

method since the np model is not implemented for this function. MQM was performed with

default parameters. The (broad-sense) heritability for the bract score was defined as $H^2 = (var_T -$

$var_E) / var_T$, with var_T and var_E being the total and environmental variance, respectively. var_T

was computed as the total variance of the bract score of each plant over the entire QTL

mapping dataset (1830 RIL plants and 780 control parents split over 12 experiments) while var_E

was computed as the pooled variance of the line variances (computed over all plants of a line

across all experiments), which averages the different line variances as is computed as: $(\sum_{i=1,N}((n_i -$

$1) \times s_i^2))/(\sum_{i=1,N}(n_i - N))$ where N is the number of line; n_i and s_i the number of plants in the line i.

Confidence intervals of H^2 were provided by bootstrapping 70% of the strain 1000 times as in

(Chuffart et al. 2016).

KASP genotyping

We selected 19 new genotyping markers from the 140SNPvCol marker set (Lutz U. et al. 2017)

to cover the two large QTLs mapped in chr1, 1a and 1b (**Fig. S6B**) and corresponding

kompetitive allele-specific PCR (KASP) oligos were ordered to Biosearch Technologies™ LGC Ltd.

To further reduce the intervals, we designed two new SNP markers targeting *Tsu-0*

polymorphisms identified in our whole-genome re-sequencing data. Specific KASP oligos (three

per SNP) were designed by LGC from the 100-bp sequence surrounding the SNP. All 21 new

markers (**Table S10**) were validated on parental genomic DNA (*Col-0* and *Tsu-0*) in a KASP assay

before their use with recombinant inbred lines. Clean genomic DNA from CTAB-based

extraction was used for all samples. For each marker, 2.5 μL of DNA (diluted to a final

concentration of 5 ng/μL), 2.5 μL of KASP-TF V4.0 2X Master Mix low ROX and 0.07 μL KASP

assay mix were mixed in 384-well plates and the genotyping assays were run in a QuantStudio 6

Flex (Applied Biosystems), using standard user guidelines for thermal cycling and final

fluorescence analysis.

Fine mapping using Heterologous inbred families

Using F_6 genotyping data for the 3RV RIL set from the VASC, we selected 3 RILs heterozygous in

a region overlapping QTL1a, QTL1b and the interval in-between: RIL 334, 488 and 478,

respectively (**Fig. 4** and **Fig. S6**). For RIL 334 and 488, we genotyped by KASP 160 segregating F8

progenies and selected 5 and 8 plants, respectively, with homozygous recombination inside the

QTL region under study. For RIL 478, we genotyped 20 plants and selected only one; genotyping

reactions were performed using PCR primers designed to amplify the published marker of the

VASC (Simon et al. 2008)(**Table S11**) and the results were read by Sanger sequencing. All

selected F_8 plants were selfed to generate an inbred line of a fixed genotyped (F_9), the collection

of F_9 inbred lines forming a Heterologous Inbred Family (HIF). HIF lines (with at least 20 plants

per line) were phenotyped for the bract score with parental control in the same experiment.

Biological sample preparation for the transcriptome time-series profiling

After 20 days in SD, *Tsu-0* and *Col-0* meristems were dissected every day in LD conditions, to

capture the precise developmental stages (especially stages T and F, see below). Three

independent biological experiments were performed with 5 to 11 meristems per replicate.

Stages are defined as follows: V, the day when plants were transferred to LD from SD; L, after 4

to 5 days of LD (identical for both *Col-0* and *Tsu-0*, which have the same meristem shape at this

time, see **Fig. 4A**); T, the main meristem enlarges and domes, (Kwiatkowska 2008; Kinoshita et

al. 2020), and lateral meristems starts being visible at the axils of young leaf primordia; F, all

young organs in the meristems were identified as flowers (note that this stage occurred several

days before bolting). Stage F in mutants was defined when several rounded primordia become

visible at the SAM (that will become branch-like flowers or only branches). Dissections were

performed from 9:00 a.m. to noon, by alternating between *Tsu-0* and *Col-0*. Micro-dissected

meristems were snap frozen in liquid nitrogen and stored at -80 °C until RNA extraction. To

avoid the induction of stress-related gene expression, meristem dissection did not exceed 3 min

between the first organ removal and freezing. For each replicate, pooled meristems were

ground with a RockyII tissue lyser according to the manufacturer's instructions. RNA was

extracted using the PicoPure RNA Isolation Kit (Arcturus, Catalog KIT0202) according to the

standard protocol. RNA concentration ranged from 3 to 64 ng/µl (average 23 ng/µl), with a RIN

value between 5.6 and 7.6 (average 6.8).

RNA-sequencing

Library preparation was made using NEBNext® Ultra™ II Directional RNA Library Prep Kit for

Illumina (New England Biolabs); NEBNext® Poly(A) mRNA Magnetic Isolation Module (New

England Biolabs); and NEBNext® Multiplex Oligos for Illumina (Index Primers Sets 1, 2 et 3),

from 40ng of RNA and sequenced using the NextSeq500 (Illumina). Sequence quality was

controlled using the Sequencing Analysis Viewer, sequences that did not pass the quality filter

were removed. Following QC an average of 43 million sequences per sample was achieved.

RNA-seq analysis (pre-processing procedures)

Quality filtering using Trimmomatic (Bolger et al. 2014) retained for all samples 96% of the

reads, which were aligned to the TAIR10 genome using STAR (Dobin et al. 2013) with the

following parameters: non-canonical splice junctions were removed, multi-mapping reads were

limited to 10, and only reads mapped once were considered to determine splicing junctions.

Between 88 and 97% of the reads were mapped to a simple locus. Normalisation of read counts

was performed using the R Bioconductor packages DESeq (Love et al. 2014), with the following

parameters: genes for which the total number of reads was below 10 were discarded, and data

were transformed with Variant Stabilising Transformation (VST) function (Anders and Huber

2010), to harmonise the variance. Consistency between biological replicates was verified using

a Principal Component Analysis on all samples with VST-transformed data.

RNA-seq analysis: Differential Gene Expression (DGE) analysis

Differential analysis was performed using the R Bioconductor package edgeR (McCarthy et al.

2012). Reads were first normalised using TMM (Trimmed mean of M-values) to reduce library-

specific biases. Normalisation factors were between 0.94 and 1.049. A generalised linear model

was applied for the analysis of differentially expressed genes (DEG). Three types of DEG analysis

were considered: DEG at each stage between the different genotypes (**table S3**), DEG between

the stage within the same genotype, and DEG across all conditions (stages and genotypes).

Multiple DEG analyses were corrected using Benjamin-Hochberg correction, and genes with a p-
value < 0.05 were retained.

RNA-seq analysis: clustering gene expressions correlating with bract presence

To identify bract signature genes (see **Figure 5**), we first calculated the minimum and maximum
expression levels among stages with and without bracts. If the minimum expression level in the
stages with bracts was greater than or equal to the maximum expression level in the stages
without bracts, the genes were classified as bract signature genes whose expression levels
positively correlated with the presence of bracts. If the maximum expression level in the stages
with bracts was lower than or equal to the minimum expression level in the stages without
bracts, these genes were classified as bract signature genes whose expression levels anti- anti-
correlated with the presence of bracts. If neither of these conditions were satisfied, genes were
discarded.

Temporal registration of gene expression dynamics

To align gene expression profiles of *A. thaliana Tsu-0* (query data) with *Col-0* (reference data),
we utilised the curve registration method in the R package greatR (Kristianingsih, 2024). This
approach involved shifting the gene expression profiles of *Tsu-0* across developmental stages
(V, L, T, F) using shift factors ranging from -1 to 1; a stretch factor was not applied due to the
two datasets being over similar times. The optimal shift parameter for each gene pair was
identified by maximising the log-likelihood. The fit with the best registration factors for each
gene was then compared to the fit with a non-registered model (without transformation) using
the Bayesian Information Criterion (BIC) statistic. A lower BIC score for the registered model
(with transformation) indicated that the gene expression dynamics of *Tsu-0* and *Col-0* were
similar, and the expression profile differences could be resolved through registration.
Conversely, a higher BIC score suggested that the profiles were best described by two individual
curves, i.e. not aligned. Prior to transformation, gene expression levels in both datasets were
centred and scaled using the z-score scaling method. The standard deviation for the replicates
at each time point was set to 0.01.

GO term analysis

GO term enrichment analyses were performed using either the 'Single Enrichment Analysis'

tool from AgriGOv2 (Tian et al. 2017) with default parameters (Fisher's test with Yekutieli

adjustment method) or clusterProfiler 4.0 (Wu et al. 2021) with default parameters (enrichGO

and simplify functions with "BH" adjustment method). The Rich factor was computed by

dividing the number of genes associated with a GOterm ("Count") by the numerator of the

"BgRatio" computed by the enrichoGO function.

Gene selection inside genetic mapping intervals

The R library 'GenomicFeatures' (Lawrence et al. 2013) was used to import gene information

from the most recent annotation file of *A. thaliana* at the gff format (Cheng et al. 2017). Custom

R scripts were used to intersect all gene loci falling within genetic mapping intervals and

provide for each gene information from RNA-seq analysis and genomic variant analysis (see

Table S2 and **Table S4**). We used snpEff v5.0d (Cingolani et al. 2012) and its putative functional

categories (HIGH/MODERATE/LOW/MODIFIER) to predict the functional impacts of the genomic

variants over regions covering each annotated gene of the interval, including promoter (2 kbp

upstream of the transcriptional start site) as well as 500-bp downstream region in the case of

miRNA. In **Table S2** and **S4**, additional gene information was retrieved from ThaleMine (Pasha

et al. 2020) and GO terms from the TAIR bulk data retrieval tool.

Data handling, data visualization and descriptive statistics

The R software was used (R Core Team 2018), especially the following packages: Bioconductor

packages (Huber et al. 2015) for the analysis of omics data, especially GenomicFeatures, IRange

(Lawrence et al. 2013), rtracklayer (Lawrence et al. 2009), clusterProfiler (Wu et al. 2021),

limma (Ritchie et al. 2015), DESseq2 (Love et al. 2014), edgeR (Chen et al. 2016) and

org.At.tair.db (Carlson 2024); dplyr (Wickham et al. 2023), tidyr (Wickham et al. 2024) and

reshape2 (Wickham 2007) for data manipulation and ggplot2 for almost all plots (Wickham

2016), with viridis and plasma (Garnier et al. 2024) for some colour optimization.

Text and English

Artificial intelligence-based language correction software was used throughout the text to

ensure accurate spelling and grammar, as well as to achieve the most comprehensible style

possible, without influencing the content.

Acknowledgements

We thank Teva Vernoux and Yoan Coudert for critical reading of the manuscript. We thank

Mathieu Hanemian, Gaël Yvert and Bjorn Pieper for their advice in quantitative genetics. We

thank Thomas Widiez and Nathanaël Jacquier for their advice in Kasp genotyping. We thank the

following researchers for providing *A. thaliana* mutant seed stocks: Mitsuhiro Aida for *bop1-4* x

bop2-11, and *puchi-1 x bop1-4 x bop2-11* ; Hicham Chathane for *lfy-12* ; Georges Coupland and

Enric Bertran Garcia de Olalla for *soc1-2*. We acknowledge the PSMN (Pôle Scientifique de

Modélisation Numérique) computing centre of 'ENS de Lyon' for providing support in the

genomic variant analysis. We thank INRAE for a grant from the BAP department (call IB2019),

the Embassy of France in the United Kingdom for a short-term travel fellowship awarded to R.

K. R.M. and R.W. gratefully acknowledge support from the Biotechnology and Biological Science

Research Council Institute Strategic Programme 'Building Resilience in Crops' (BB/X01102X/1).

SD was supported by a Ph.D contract of the french ministry of Research and Higher Education.

Competing interests,

The authors declare that they have no competing interests

Author contributions

Sana Dieudonné: Conceptualization, Investigation, Software (implementation), Formal analysis,

Visualization, Methodology, Writing -review and editing; Ruth Kristianingsih: Formal Analysis,

Methodology, Software (designing, programming, testing, implementing), Visualization, Writing

-review and editing; Stephanie Laine, Investigation; Béline Jesson: Formal analysis, Véronique

Vidal: Supervision, Rachel Wells: Supervision, Writing -review and editing; Richard Morris:

Supervision, Writing -review and editing; Fabrice Besnard: Conceptualization, Project

administration, Supervision, Investigation, Formal analysis, Methodology, Software

(programming, implementing), Funding acquisition, Visualization, Writing - original draft, review

and editing.

Data availability

Whole-genome DNA-seq data of the parental lines (*Col-0* and *Tsu-0*) and of the two F2 pools

used for bulk segregant analysis are deposited under this identifier: <u>doi:10.57745/Z80SIM</u>.

Time-course RNA-seq data of *Col-0* and *Tsu-0* micro-dissected meristems during flowering are

available with these doi: <u>doi:10.57745/DKMQ06</u>, <u>doi:10.57745/7JI3E7</u>, respectively.

References

Alberch P, Gould SJ, Oster GF, Wake DB. 1979. Size and Shape in Ontogeny and
Phylogeny. Paleobiology. 5(3):296–317.

Al-Shehbaz IA. 2015. Cardamine xinfenii (Brassicaceae), a New Species from Sichuan
(China). Harvard Papers in Botany. 20(2):145–146.

Anders S, Huber W. 2010. Differential expression analysis for sequence count data.
Genome Biol. 11(10):R106. doi:10.1186/gb-2010-11-10-r106.

Arber A. 1931a. Studies in Floral Morphology. I. On Some Structural Features of the
Cruciferous Flower. The New Phytologist. 30(1):11–41.

Arber A. 1931b. Studies in Floral Morphology. II. Some Normal and Abnormal Crucifers:
with a Discussion on teratology and Atavism. The New Phytologist. 30(3):172–203.

Baressi M, Gilbert SF. 2023. Developmental Biology. 13th ed. Oxford University Press.

van Berloo R. 2008. GGT 2.0: Versatile Software for Visualization and Analysis of Genetic
Data. Journal of Heredity. 99(2):232–236. doi:10.1093/jhered/esm109.

Besnard F, Picao-Osorio J, Dubois C, Félix M-A. 2020. A broad mutational target explains
a fast rate of phenotypic evolution. Weigel D, Landry CR, Matus D, editors. eLife.
9:e54928. doi:10.7554/eLife.54928.

Bloomer RH, Dean C. 2017. Fine-tuning timing: natural variation informs the mechanistic
basis of the switch to flowering in Arabidopsis thaliana. Journal of Experimental Botany.
68(20):5439–5452. doi:10.1093/jxb/erx270.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics. 30(15):2114–2120. doi:10.1093/bioinformatics/btu170.

Broman KW, Gatti DM, Simecek P, Furlotte NA, Prins P, Sen Ś, Yandell BS, Churchill GA.
2019. R/qtl2: Software for Mapping Quantitative Trait Loci with High-Dimensional Data
and Multiparent Populations. Genetics. 211(2):495–502.
doi:10.1534/genetics.118.301595.

Calderwood A, Hepworth J, Woodhouse S, Bilham L, Jones DM, Tudor E, Ali M, Dean C,
Wells R, Irwin JA, et al. 2021. Comparative transcriptomics reveals desynchronisation of
gene expression during the floral transition between Arabidopsis and Brassica rapa
cultivars. Quant Plant Biol. 2:e4. doi:10.1017/qpb.2021.6.

Carlson M. 2024. org.At.tair.db: Genome wide annotation for Arabidopsis.

Chahtane H, Zhang B, Norberg M, LeMasson M, Thévenon E, Bakó L, Benlloch R,
Holmlund M, Parcy F, Nilsson O, et al. 2018. LEAFY activity is post-transcriptionally
regulated by BLADE ON PETIOLE2 and CULLIN3 in Arabidopsis. New Phytol. 220(2):579–
592. doi:10.1111/nph.15329.

Chen Y, Lun ATL, Smyth GK. 2016. From reads to genes to pathways: differential
expression analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-
likelihood pipeline. doi:10.12688/f1000research.8987.2. [accessed 2024 Jul 16].
https://f1000research.com/articles/5-1438.

Cheng C-Y, Krishnakumar V, Chan AP, Thibaud-Nissen F, Schobel S, Town CD. 2017.
Araport11: a complete reannotation of the Arabidopsis thaliana reference genome.
Plant J. 89(4):789–804. doi:10.1111/tpj.13415.

Cheng X, Hui JHL, Lee YY, Wan Law PT, Kwan HS. 2015. A "developmental hourglass" in
fungi. Mol Biol Evol. 32(6):1556–1566. doi:10.1093/molbev/msv047.

Chuffart F, Richard M, Jost D, Burny C, Duplus-Bottin H, Ohya Y, Yvert G. 2016. Exploiting
Single-Cell Quantitative Data to Map Genetic Variants Having Probabilistic Effects. PLOS
Genetics. 12(8):e1006213. doi:10.1371/journal.pgen.1006213.

Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM.
2012. A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118;
iso-2; iso-3. Fly (Austin). 6(2):80–92. doi:10.4161/fly.19695.

Clough SJ, Bent AF. 1998. Floral dip: a simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. Plant J. 16(6):735–743.

Cutler SR, Ehrhardt DW, Griffitts JS, Somerville CR. 2000. Random GFP::cDNA fusions
enable visualization of subcellular structures in cells of Arabidopsis at a high frequency.
Proc Natl Acad Sci U S A. 97(7):3718–3723. doi:10.1073/pnas.97.7.3718.

Dinneny JR, Yadegari R, Fischer RL, Yanofsky MF, Weigel D. 2004. The role of JAGGED in
shaping lateral organs. Development. 131(5):1101–1110. doi:10.1242/dev.00949.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 29(1):15–
21. doi:10.1093/bioinformatics/bts635.

Endress PK. 2006. Angiosperm Floral Evolution: Morphological Developmental
Framework. In: Advances in Botanical Research. Vol. 44. Academic Press.
(Developmental Genetics of the Flower). p. 1–61. [accessed 2023 Dec 1].
https://www.sciencedirect.com/science/article/pii/S0065229606440015.

Furlow JD, Neff ES. 2006. A developmental switch induced by thyroid hormone: *Xenopus*
laevis metamorphosis. Trends in Endocrinology & Metabolism. 17(2):40–47.
doi:10.1016/j.tem.2006.01.007.

Garnier, Simon, Ross, Noam, Rudis, Robert, Camargo, Pedro A, Sciaini, Marco, et al.
2024. viridis(Lite) - Colorblind-Friendly Color Maps for R.
https://sjmgarnier.github.io/viridis/.

German DA, Hendriks KP, Koch MA, Lens F, Lysak MA, Bailey CD, Mummenhoff K, Al-
Shehbaz IA. 2023. An updated classification of the Brassicaceae (Cruciferae). PhytoKeys.
220:127–144. doi:10.3897/phytokeys.220.97724.

Goldshmidt A, Alvarez JP, Bowman JL, Eshed Y. 2008. Signals derived from YABBY gene
activities in organ primordia regulate growth and partitioning of Arabidopsis shoot
apical meristems. Plant Cell. 20(5):1217–1230. doi:10.1105/tpc.107.057877.

Healey A, Furtado A, Cooper T, Henry RJ. 2014. Protocol: a simple method for extracting
next-generation sequencing quality genomic DNA from recalcitrant plant species. Plant
Methods. 10:21. doi:10.1186/1746-4811-10-21.

Heisler MG, Ohno C, Das P, Sieber P, Reddy GV, Long JA, Meyerowitz EM. 2005. Patterns
of auxin transport and gene expression during primordium development revealed by
live imaging of the Arabidopsis inflorescence meristem. Curr Biol. 15(21):1899–1911.
doi:10.1016/j.cub.2005.09.052.

Hempel FD, Feldman LJ. 1995. Specification of chimeric flowering shoots in wild-type
Arabidopsis. Plant J. 8(5):725–731. doi:10.1046/j.1365-313x.1995.08050725.x.

Hempel FD, Zambryski PC, Feldman LJ. 1998. Photoinduction of flower identity in
vegetatively biased primordia. Plant Cell. 10(10):1663–1676.
doi:10.1105/tpc.10.10.1663.

Huber W, Carey VJ, Gentleman R, Anders S, Carlson M, Carvalho BS, Bravo HC, Davis S,
Gatto L, Girke T, et al. 2015. Orchestrating high-throughput genomic analysis with
Bioconductor. Nat Methods. 12(2):115–121. doi:10.1038/nmeth.3252.

Irish VF, Sussex IM. 1990. Function of the apetala-1 gene during Arabidopsis floral
development. Plant Cell. 2(8):741–753. doi:10.1105/tpc.2.8.741.

Kalinka AT, Varga KM, Gerrard DT, Preibisch S, Corcoran DL, Jarrells J, Ohler U, Bergman
CM, Tomancak P. 2010. Gene expression divergence recapitulates the developmental
hourglass model. Nature. 468(7325):811–814. doi:10.1038/nature09634.

Karim MR, Hirota A, Kwiatkowska D, Tasaka M, Aida M. 2009. A role for Arabidopsis
PUCHI in floral meristem identity and bract suppression. Plant Cell. 21(5):1360–1372.
doi:10.1105/tpc.109.067025.

Kawakatsu T, Taramino G, Itoh J-I, Allen J, Sato Y, Hong S-K, Yule R, Nagasawa N, Kojima
M, Kusaba M, et al. 2009. PLASTOCHRON3/GOLIATH encodes a glutamate
carboxypeptidase required for proper development in rice. Plant J. 58(6):1028–1040.
doi:10.1111/j.1365-313X.2009.03841.x.

Keyte AL, Smith KK. 2014. Heterochrony and developmental timing mechanisms:
changing ontogenies in evolution. Semin Cell Dev Biol. 34:99–107.
doi:10.1016/j.semcdb.2014.06.015.

Kiefer M, Schmickl R, German DA, Mandáková T, Lysak MA, Al-Shehbaz IA, Franzke A,
Mummenhoff K, Stamatakis A, Koch MA. 2014. BrassiBase: introduction to a novel
knowledge database on Brassicaceae evolution. Plant Cell Physiol. 55(1):e3.
doi:10.1093/pcp/pct158.

Kinoshita A, Vayssières A, Richter R, Sang Q, Roggen A, van Driel AD, Smith RS, Coupland
G. 2020. Regulation of shoot meristem shape by photoperiodic signaling and
phytohormones during floral induction of Arabidopsis. Elife. 9:e60661.
doi:10.7554/eLife.60661.

Kristianingsih R. 2024. greatR: Gene Registration from Expression and Time-Courses in R.
https://ruthkr.github.io/greatR/.

Kwiatkowska D. 2006. Flower primordium formation at the Arabidopsis shoot apex:
quantitative analysis of surface geometry and growth. J Exp Bot. 57(3):571–580.
doi:10.1093/jxb/erj042.

Kwiatkowska D. 2008. Flowering and apical meristem growth dynamics. J Exp Bot.
59(2):187–201. doi:10.1093/jxb/erm290.

Lawrence M, Gentleman R, Carey V. 2009. rtracklayer: an R package for interfacing with
genome browsers. Bioinformatics. 25(14):1841–1842.
doi:10.1093/bioinformatics/btp328.

Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R, Morgan MT,
Carey VJ. 2013. Software for computing and annotating genomic ranges. PLoS Comput
Biol. 9(8):e1003118. doi:10.1371/journal.pcbi.1003118.

Lemmon ZH, Park SJ, Jiang K, Van Eck J, Schatz MC, Lippman ZB. 2016. The evolution of
inflorescence diversity in the nightshades and heterochrony during meristem
maturation. Genome Res. 26(12):1676–1686. doi:10.1101/gr.207837.116.

Levin M, Anavy L, Cole AG, Winter E, Mostov N, Khair S, Senderovich N, Kovalev E, Silver
DH, Feder M, et al. 2016. The mid-developmental transition and the evolution of animal
body plans. Nature. 531(7596):637–641. doi:10.1038/nature16994.

Long J, Barton MK. 2000. Initiation of axillary and floral meristems in Arabidopsis. Dev
Biol. 218(2):341–353. doi:10.1006/dbio.1999.9572.

Lotharukpong JS, Zheng M, Luthringer R, Liesner D, Drost H-G, Coelho SM. 2024. A
transcriptomic hourglass in brown algae. Nature. 635(8037):129–135.
doi:10.1038/s41586-024-08059-8.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol. 15(12):550. doi:10.1186/s13059-014-
0550-8.

Mach J. 2010. Different Words, Same Message: How Grasses and Arabidopsis Say "Hold
the Bract." The Plant Cell. 22(3):538. doi:10.1105/tpc.110.220311.

Maizel A, Weigel D. 2004. Temporally and spatially controlled induction of gene

expression in Arabidopsis thaliana. The Plant Journal. 38(1):164–171.
doi:10.1111/j.1365-313X.2004.02027.x.

Mansfeld BN, Grumet R. 2018. QTLseqr: An R Package for Bulk Segregant Analysis with
Next-Generation Sequencing. Plant Genome. 11(2).
doi:10.3835/plantgenome2018.01.0006.

Manuela D, Xu M. 2024 Mar 28. Aintegumenta And Redundant Aintegumenta-Like6 Are
Required For Bract Outgrowth In Arabidopsis. J Exp Bot.:erae138.
doi:10.1093/jxb/erae138.

Maple R, Zhu P, Hepworth J, Wang J-W, Dean C. 2024. Flowering time: From physiology,
through genetics to mechanism. Plant Physiol. 195(1):190–212.
doi:10.1093/plphys/kiae109.

McCarthy DJ, Chen Y, Smyth GK. 2012. Differential expression analysis of multifactor
RNA-Seq experiments with respect to biological variation. Nucleic Acids Res.
40(10):4288–4297. doi:10.1093/nar/gks042.

Michaels SD, Ditta G, Gustafson-Brown C, Pelaz S, Yanofsky M, Amasino RM. 2003.
AGL24 acts as a promoter of flowering in Arabidopsis and is positively regulated by
vernalization. The Plant Journal. 33(5):867–874. doi:10.1046/j.1365-313X.2003.01671.x.

Nilsson O, Wu E, Wolfe DS, Weigel D. 1998. Genetic ablation of flowers in transgenic
Arabidopsis. Plant J. 15(6):799–804. doi:10.1046/j.1365-313x.1998.00260.x.

Ohno CK, Reddy GV, Heisler MGB, Meyerowitz EM. 2004. The Arabidopsis JAGGED gene
encodes a zinc finger protein that promotes leaf tissue development. Development.
131(5):1111–1122. doi:10.1242/dev.00991.

Pasha A, Subramaniam S, Cleary A, Chen X, Berardini T, Farmer A, Town C, Provart N.
2020. Araport Lives: An Updated Framework for Arabidopsis Bioinformatics. Plant Cell.
32(9):2683–2686. doi:10.1105/tpc.20.00358.

Penin AA. 2008. Bract reduction in Cruciferae: possible genetic mechanisms and
evolution. Wulfenia. 15:63–73.

Prenner G, Vergara-Silva F, Rudall PJ. 2009. The key role of morphology in modelling
inflorescence architecture. Trends Plant Sci. 14(6):302–309.
doi:10.1016/j.tplants.2009.03.004.

Quint M, Drost H-G, Gabel A, Ullrich KK, Bönn M, Grosse I. 2012. A transcriptomic
hourglass in plant embryogenesis. Nature. 490(7418):98–101.
doi:10.1038/nature11394.

R Core Team. 2018. R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing. Vienna, Austria. https://www.R-project.org/.

Rieu P, Beretta VM, Caselli F, Thévénon E, Lucas J, Rizk M, Franchini E, Caporali E, Paleni
C, Nanao MH, et al. 2024. The ALOG domain defines a family of plant-specific
transcription factors acting during Arabidopsis flower development. Proceedings of the
National Academy of Sciences. 121(10):e2310464121. doi:10.1073/pnas.2310464121.

Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. 2015. limma powers
differential expression analyses for RNA-sequencing and microarray studies. Nucleic
Acids Research. 43(7):e47. doi:10.1093/nar/gkv007.

Salomé PA, Bomblies K, Laitinen RAE, Yant L, Mott R, Weigel D. 2011. Genetic
architecture of flowering-time variation in Arabidopsis thaliana. Genetics. 188(2):421–
433. doi:10.1534/genetics.111.126607.

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image
analysis. Nat Methods. 9(7):671–675. doi:10.1038/nmeth.2089.

Schultz EA, Haughn GW. 1991. LEAFY, a Homeotic Gene That Regulates Inflorescence
Development in Arabidopsis. Plant Cell. 3(8):771–781. doi:10.1105/tpc.3.8.771.

Simon M, Loudet O, Durand S, Bérard A, Brunel D, Sennesal F-X, Durand-Tardif M,
Pelletier G, Camilleri C. 2008. Quantitative trait loci mapping in five new large
recombinant inbred line populations of Arabidopsis thaliana genotyped with consensus
single-nucleotide polymorphism markers. Genetics. 178(4):2253–2264.
doi:10.1534/genetics.107.083899.

Siriwardana NS, Lamb RS. 2012. The poetry of reproduction: the role of LEAFY in
Arabidopsis thaliana flower formation. Int J Dev Biol. 56(4):207–221.
doi:10.1387/ijdb.113450ns.

Smyth DR, Bowman JL, Meyerowitz EM. 1990. Early flower development in Arabidopsis.
Plant Cell. 2(8):755–767. doi:10.1105/tpc.2.8.755.

Takagi H, Abe A, Yoshida K, Kosugi S, Natsume S, Mitsuoka C, Uemura A, Utsushi H,
Tamiru M, Takuno S, et al. 2013. QTL-seq: rapid mapping of quantitative trait loci in rice

by whole genome resequencing of DNA from two bulked populations. Plant J.
74(1):174–183. doi:10.1111/tpj.12105.

Tian T, Liu Y, Yan H, You Q, Yi X, Du Z, Xu W, Su Z. 2017. agriGO v2.0: a GO analysis
toolkit for the agricultural community, 2017 update. Nucleic Acids Res. 45(W1):W122–
W129. doi:10.1093/nar/gkx382.

Truman JW. 2019. The Evolution of Insect Metamorphosis. Current Biology.
29(23):R1252–R1268. doi:10.1016/j.cub.2019.10.009.

Wang L, Ming L, Liao K, Xia C, Sun S, Chang Y, Wang H, Fu D, Xu C, Wang Z, et al. 2021.
Bract suppression regulated by the miR156/529-SPLs-NL1-PLA1 module is required for
the transition from vegetative to reproductive branching in rice. Mol Plant. 14(7):1168–
1184. doi:10.1016/j.molp.2021.04.013.

Weigel D. 1995. The genetics of flower development: from floral induction to ovule
morphogenesis. Annu Rev Genet. 29:19–39. doi:10.1146/annurev.ge.29.120195.000315.

Weigel D, Alvarez J, Smyth DR, Yanofsky MF, Meyerowitz EM. 1992. LEAFY controls floral
meristem identity in Arabidopsis. Cell. 69(5):843–859. doi:10.1016/0092-
8674(92)90295-n.

Whipple CJ. 2017. Grass inflorescence architecture and evolution: the origin of novel
signaling centers. New Phytol. 216(2):367–372. doi:10.1111/nph.14538.

Whipple CJ, Hall DH, DeBlasio S, Taguchi-Shiobara F, Schmidt RJ, Jackson DP. 2010. A
conserved mechanism of bract suppression in the grass family. Plant Cell. 22(3):565–
578. doi:10.1105/tpc.109.073536.

Wickham H. 2007. Reshaping Data with the reshape Package. Journal of Statistical
Software. 21(12):1–20.

Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New
York. https://ggplot2.tidyverse.org.

Wickham H, François R, Henry L, Müller K, Vaughan D. 2023. dplyr: A Grammar of Data
Manipulation. https://dplyr.tidyverse.org.

Wickham H, Vaughan D, Girlich M. 2024. tidyr: Tidy Messy Data.
https://tidyr.tidyverse.org.

Wigge PA, Kim MC, Jaeger KE, Busch W, Schmid M, Lohmann JU, Weigel D. 2005.

Integration of Spatial and Temporal Information During Floral Induction in Arabidopsis.
Science. 309(5737):1056–1059. doi:10.1126/science.1114358.

Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, Feng T, Zhou L, Tang W, Zhan L, et al. 2021.
clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. The
Innovation. 2(3):100141. doi:10.1016/j.xinn.2021.100141.

Xiao Y, Guo J, Dong Z, Richardson A, Patterson E, Mangrum S, Bybee S, Bertolini E,
Bartlett M, Chuck G, et al. 2022. Boundary domain genes were recruited to suppress
bract growth and promote branching in maize. Sci Adv. 8(24):eabm6835.
doi:10.1126/sciadv.abm6835.

Yamaguchi N. 2021. LEAFY, a Pioneer Transcription Factor in Plants: A Mini-Review.
Front Plant Sci. 12. doi:10.3389/fpls.2021.701406. [accessed 2024 Dec 19].
https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2021.701406/
full.

1040   Supporting Information

Figures S1 to S13

Figure S1: Example of natural transient bract formation at the base of branches among
Brassicaceæ tribes.

Figure S2: The presence of basal bracts in *A. thaliana* accessions does not correlate with their
geographic or genetic origin.

Figure S3: *Tsu-0* plants flower later than *Col-0* plants in different culture conditions.

Figure S4: Bulk segregant analysis of basal bract formation (*Tsu-0* x *Col-0*) identifies four
putative QTLs.

Figure S5: Quantitative genetics of basal bracts formation in *Tsu-0* using a set of RILs with the
reference accession *Col-0*.

Figure S6: Fine mapping of the QTLs controlling basal bract formation in *Tsu-0*.

Figure S7: QTL1b* identified for basal bract correlates with a higher cauline branch number.

Figure S8: Transgressive phenotypes in HIF reveal epistasis controlling floral determinacy
between alleles of *Col-0* and *Tsu-0*.

Figure S9: Gene expression dynamics during the floral transition in micro-dissected meristems
of *Col-0* and *Tsu-0* plants.

Figure S10: Phenotypic differences between the bracts of the mutants affected in the FMI

pathway and the natural basal bracts observed in *Tsu-0*.

Figure S11: Functional analysis of the genes differentially expressed in *Tsu-0* meristem at the

stage T when bract repression is delayed.

Figure S13: Evolution of the proportions of DEGs between *Tsu-0* and *Col-0* depending on the

heterochronic shift and on the developmental stage

Figure S13: Temporal registration of expression dynamics in *Tsu-0* over the floral transition for

genes related to bract development, floral identity and floral transition.

Figure S14: A working model for natural basal bract formation in *Arabidopsis thaliana.*

1066

Table S1 to S11

Table S1: Literature presenting mutants in genes functionally associated with the FMI pathway

in and showing derepressed bract formation.

Note that tables S2 to S11 are large spreadsheets available for download:

Table S2: List of annotated genes lying in the four mapped QTLs controlling bracts in *Tsu-0*, with

additional information from RNAseq and genomic variant analysis.

Table S3: Differential gene expression analysis and log2CPM for the genes expressed in

microdissected meristems of *Tsu-0* and *Col-0* during the floral transition.

Table S4: Intersection of the genetic mapping, the genomic variant analysis and the RNAseq

data.

Table S5: Bract signature genes in *Tsu-0* identified by clustering apart expression from bract and

non-bract making stages in *Tsu-0* and *Col-0* time-series.

Table S6: Details for all genes associated with "response to salicylic acid" about their expression

at stage in *Tsu-0* and their identification as bract signature genes.

Table S7: Details for all genes associated with anthocyanin metabolism about their expression

at stage in *Tsu-0* and their identification as bract signature genes.

Table S8: Results of curve registration

Table S9: Genotypes of all RIL and HIF lines used in this study.

Table S10: SNP information relative to new KASP genotyping markers used in this study.

Table S11: SNP information relative to new sanger genotyping markers used in this study and

associated primers.

1088

Figure legends

Figure 1: **Desynchronization of leaf repression from flower formation in the natural *Tsu-0***
accession of *Arabidopsis thaliana*.
A, lateral views of the main stem of *Col-0* (left) and *Tsu-0* (right) accessions showing the
transition from branches (at the bottom) to flowers (to the top). The red arrow points to a bract (a
leaf) subtending the first flower in *Tsu-0*. Cartoons on the side summarize the structures
produced at each node. **B**, Distribution (as percent) of the position of bracts in *Tsu-0* (N plants =
50, n bracts = 337) in the main stem and cauline branches, where n°1 is the first flower after
floral transition. **C,** Number of bracts on the different branches of a *Tsu-0* inflorescence (dot is
the mean, vertical bars are sd). Branch positions are numbered with increasing numbers from
the base to the top of the main stem, which is noted as 0 by convention (N plants = 90). A linear
regression (horizontal red dotted line with standard error shaded in red) indicates a stable
average value of 1 bract on each branch. **D,** *Col-0* floral pedicels do not show any sign of bract
development (black arrowhead). **E**, Variable bract development (red arrows) observable in *Tsu-*
0, from true leaves (left) to filamentous or small rudimentary structures at the base of the floral
pedicel. **F**, Scanning electron microscopy of floral pedicels at the floral transition in *Col-0* (left)
and *Tsu-0* (right). In *Col-0*, the pedicel is a regular, naked cylinder at the base (black arrow)
while in *Tsu-0*, there is a bract on the first flower (red arrow) and swollen base of pedicels on the
two following flowers (red arrowhead). **G**, Number of floral organs in flowers with (dark red) or
without (light red) bract in *Tsu-0* plants. (**H,I**) Scanning electron microscopy of the main
meristem of plants captured right at the transition to flowering in *Col-0* (**F**) and *Tsu-0* (**G**). The
first flowers after the transition are pointed by white arrows and ordered with increasing
numbers in their order of production inferred from size and phyllotactic positions. Green
arrowheads: cauline branch (axillary meristem and a cauline leaf), -1 indicates the last branch
before floral transition. Red arrows: bracts. In *Col-0,* no bract is visible on the adaxial side of the
developing flowers.

1115

Figure 2: **Desynchronization of leaf repression from flower formation in *Tsu-0* and other**
natural accessions exhibits complex relationships with flowering time and photo-
induction conditions.
A, Correlation study between the number of lateral (cauline) branches and the number of bracts
on the main stem in *Tsu-0* (Pearson test, p-value > 0.1), which do not show a negative
correlation expected by the "conversion" model (see main text). **B**, Definition of a quantitative
bract score for a single plant. **C**, Quantification of basal bracts in different accessions using the
bract score. **D**, Correlation study between the bract score and the flowering time (assayed when
the first flower opens) in different accessions, labelled with different colours. Each dot is a plant
and a linear regression with standard deviation (color shade) is computed for each accession.
E, Effect of different light regimes on bract scores in *Col-0* and *Tsu-0* plants (green and red,
respectively). LD and SD stand for long- and short-day conditions, respectively and CL stands
for continuous light (see methods). Numbers (e.g. 20SD > LD) indicate the number of days in
the first condition before transfer to the second. The number of plants scored is indicated below
each bar.

1131

Figure 3: **Basal bract formation in *Tsu-0* is controlled by four major QTL that do not**

contain the FMI genes associated with bract repression. **A**, Genetic transmission of bract
score in F1 and F2 hybrids from a cross between *Col-0* and *Tsu-0* parent plants. **B**, QTL
mapping for bract score in a set of RILs identifies four putative QTLs, two in chromosome 1 (1a
and 1b) and two in chromosome 5 (5a and 5b). Dotted horizontal red lines indicate three
different significance thresholds computed from 2000 permutations. On top of the graph, H^2
indicates the (broad-sense) heritability of the bract score computed among the RIL set. **C**,
Genotype of the F6 generation of the line RIL334 in chromosome 1, showing a heterozygotic
region overlapping with QTL1a. **D**, Chromosome 1 genotypes of selected lines forming a
heterologous inbred family (HIF) obtained from RIL334 after two more generations of selfing
(F8). **E**, Box plots of bract scores of the different lines from the "334 family" (blue boxes) with
parental controls (*Col-0*: green, *Tsu-0*: red, N > 20 plants per line). Segregating bract scores
within the family map a narrower region, QTL1a*, spanning 541 annotated genes. **F**, Genotype
of the F6 generation of the line RIL488 in chromosome 1, showing a heterozygotic region
overlapping with QTL1b. **G**, Chromosome 1 genotypes of selected HIF lines obtained from
RIL488 at F8 and (**H**) a box plot of their bract scores (blue boxes), with parental controls (*Col-0*:
green, *Tsu-0*: red, N > 21 plants per line). This maps a narrower region, QTL1b*, spanning 332
annotated genes. In E and H, lines not sharing the same letter(s) are statistically different (*post*
hoc Tukey analysis with 0.05 sign. level from a glm of bract scores fitted with a quasi-poisson
distribution). **I**, Final genetic mapping of the 'basal bract' trait in *Tsu-0* and the location of genes
reported to impact bract development in previous mutant studies.
1153
Figure 4: **Transcriptomic divergence between *Tsu-0* and *Col-0* meristems peaks at the**
floral transition despite synchronization of the FMI pathway genes
A, Scanning electron microscopy showing the evolution of the main meristem in both *Col-0* and
Tsu-0 at the four different stages (V, L, T, F) used for RNAseq. Plants were first synchronized
by 21 days of a non-inductive short-day light regime before a transfer to inductive long days.
The date after the transfer is indicated in the top-right corner of each picture. Green arrowheads
point to branches (with a leaf) while magenta arrowheads point to the first flowers produced
after floral transition. In *Tsu-0*, red arrows show bracts. Scale bars: 50 μm. **B,** Dynamics of the
expression levels of FT, LFY and AP1 (three important regulators of floral transition and identity)
in both *Col-0* (green) and *Tsu-0* (red) over the four developmental stages. Differential
expression analysis reveals no difference at any stage between the two accessions. **C, D**.
Confocal live imaging of the main meristems of *Col-0* (**C**) and *Tsu-0* (**D**) plants at floral
transition, expressing a pLFY transcriptional reporter (magenta) and a membrane marker
(green) for morphology (top-view projections of stack acquisitions). Green arrowheads:
branches, ordered with decreasing numbers from the floral transition; white arrows: flowers
ordered with increasing numbers from the floral transition. In both genotypes, pLFY expression
is absent from branch meristems and suddenly appears in the first flower onwards.
Representative pictures of at least 6 plants per genotype captured at floral transition. **E**, side-
view of the first flower from image D (*Tsu-0*): a bract (red arrow) is visible on the abaxial side of
this young flower. Scale bars: 50 μm. **F**, Bar plots of the number of genes differentially
expressed between two consecutive stages in *Col-0* (green) and *Tsu-0* (red). The number of
changes peaks at the V-to-T stage transition, especially in *Tsu-0*. **G**, Bar plots of the number of
differentially expressed genes (DEG) between *Col-0* and *Tsu-0* at each stage of the time
course. T stage is when the highest number of DEG is measured (on top of the bar, different
letters indicate the statistical difference with a chi-square test of homogeneity after *post hoc*
analysis). **H**, Dynamics of the expression levels of four genes reported to be involved in bract
development, showing a significant difference at least at the T stage between *Col-0* (green) and
Tsu-0 (red). One star: genes are differentially expressed; two stars: the absolute fold change is
superior to 1.
1183

Figure 5: Identifying a transcriptomic signature of leaf development in shoot meristems
at the floral transition stage using time-series RNAseq.
A, Intersecting genetic mapping and transcriptomic data: if a genetic variant causing *Tsu-0*
bracts also induces the differential expression of the gene in *cis*, this gene should verify the
conditions defined by A, B, C and D. **B,** evolution of the number of candidate genes mapped in
the four QTL (condition A) by the successive addition of other conditions (B to D). **C**, Strategy to
cluster genes based on the presence (circle with the plain line) or the absence (dotted circle) of
leaf and/or bract in the different combinations of stages and genotypes (see text for details). **D**,
Volcano plot of gene expressions at the stage T between *Tsu-0* and *Col-0*. All genes expressed
in the SAM are plotted (n=21,652 grey dots) but only the genes fulfilling the clustering condition
defined in C ("bract signature genes") are highlighted in orange and blue for up- and down-
regulation, respectively (n=124). Vertical dashed lines: absolute fold change superior to 1,
horizontal dashed line: significance threshold at 5.10^{-2} (adjusted p.value with fdr method). **E**,
Seven 'biological process' GO terms are statistically enriched among the up-regulated genes
identified in D (bar plot providing the number of related genes). They are all related to the
anthocyanin biosynthesis pathway, which is cited in the two most significant terms (stars). **F**,
Only three 'biological process' GO terms are statistically enriched among the down-regulated
genes identified in D (bar plot providing the number of related genes); they collectively point to
defense response and more specifically the acid salicylic pathway (star). In E,F, color gradients
indicate the statistical significance of the GO term enrichment (Fisher's test with Yekutieli
adjustment method). **G**, Representative pictures of micro-dissected meristems in *Col-0* (upper
row) and *Tsu-0* (lower row) just at or before (left) or after (middle) stage T (N>15 for each
genotype) and a close-up of the base of the bolted main stem (right). At stage T, *Tsu-0*
meristems display a typical anthocyanin red coloration just below the meristem, which is not
observed in *Col-0*. After bolting, both genotypes show anthocyanin coloration at the base of the
stem. Scale bars: 100 µm (left and middle), 1 cm (right). **H**, Intersection of the gene set in A with
the 'bract signature genes'. **I**, Examples of the expression profile of two candidate genes,
showing an up- (left) and a down- (right) regulation at the T stage. *DFR* is an enzyme involved
in anthocyanin biosynthesis while *FMOGS-OX7* is annotated as a salicylic acid responding
enzyme. Two stars mean that the genes are differentially expressed and the absolute fold
change is superior to 1. The bract and bract-less clusters are outlined with a solid and dashed
circle, respectively, while the horizontal dotted line highlights their separation.
1216	
Figure 6: **Massive and complex transcriptional desynchronisation coexist in *Tsu-0* across**
the floral transition.
A, PCA of RNAseq data from microdissected meristems of *Col-0* and *Tsu-0* over the four
sampled stages (biological replicates averaged per time points). The two main axes of the PCA
can be interpreted as developmental time and genotype, respectively. **B**, Examples of temporal
registration of gene dynamics (right panels) between *Col-0* (green) and *Tsu-0* (red) from scaled
expression levels (left panel). Dots represent the expression levels of independent biological
replicates, while lines indicate the mean expression level at each time point. In the right panel,
the green (or red) dotted curves represent the fitted models for *Col-0* (or *Tsu-0*) independently,
while the grey dotted curve represents the joint model for both *Col-0* and *Tsu-0*. If the green and
red dotted curves are used, it means the two time series are best explained by two independent
models, indicating they are not similar. Conversely, if the grey dotted curve is used, it suggests
that a single model best explains both time series, indicating they are similar. The name of the
gene plotted is indicated on the left of each row. These four genes exemplify the four possible
categories (from top to bottom, respectively): genes that cannot be registered (N = 43, *e.g.*
CYP705A9), genes with identical temporal dynamics (shift = 0, *e.g. AP1,* see also Fig. 4B) and
genes whose expression dynamics in *Tsu-0* must be shifted negatively (*e.g. DFR,* see also Fig.
5I) or positively (*e.g*. *AG*) to align with *Col-0*. The last column provides a biological interpretation

of the computed shift (see main text): a null shift indicates that the expression dynamics in *Tsu-*
0 stay "*in phase*" with floral transition while negative or positive shifts indicate that the
expression dynamics in *Tsu-0* are desynchronized later or earlier, respectively, than the
phenotypic progression of floral transition. **C**, Distribution of heterochronic shifts between *Tsu-0*
and *Col-0* on the entire meristematic transcriptome, computed by the registration method from
B. The shift value is colour-coded in a red-to-blue gradient from -1 to 1. **D**, Correlation between
the heterochronic shifts in gene expression and the fraction of genes presenting a differential
expression between *Tsu-0* and *Col-0* for each stage, colour-coded from green to orange. The
blue dotted lines are two piecewise linear regressions modeling the effect of the shift with a
binomial model for positive and negative values (p-value < 2e-16) surrounded by a 5%
confidence interval (gray shade). Stars highlight the significantly higher proportion of DEGs at
the T stage (p-value code: *** < 0.001, * < 0.05 ; see also figure S12 for the complete statistical
analysis). **E**, GO term enrichment analysis associated with the three categories of heterochronic
shifts. The list of all significant 'biological process' GO terms (BH-adjusted p.value < 0.05) was
simplified using semantic similarity (cutoff = 0.7) and the Rich Factor was computed for the
remaining terms, revealing the proportion of genes involved among all the genes associated
with this GO term. Dot size indicates the count of genes and the color scale is the statistical
significance (BH-adjusted p.value) of the enrichment in the shift category. Stars indicate GO
terms referring to developmental processes.
1254
1255

1256   **FIGURE 1**

1257

1258   **FIGURE2**

1259

1260   **FIGURE3**

1262   **FIGURE4**

1263

1264   **FIGURE5**

1265

1266   **FIGURE6**

1267