Natural variation suggests new mechanisms for bract development in

Arabidopsis, desynchronising bract suppression from the floral transition

Dieudonné Sana[1], Kristianingsih Ruth[3], Laine Stéphanie[1], Jesson Béline[2], Vidal Véronique[2], Wells Rachel[4], Morris Richard[3] and Besnard Fabrice[1]

[1]Laboratoire RDP, (Univ. Lyon, ENS de Lyon, UCB Lyon 1, CNRS, INRAe, Inria), Lyon, France; [2]Helixio, Clermont-Ferrand, France; [3]Department of Computational and Systems Biology, John Innes Centre, Norwich, United Kingdom and [4]Department of Crop Genetics, John Innes Centre, Norwich, United Kingdom.

correspondance: fabrice.besnard@ens-lyon.fr

Word count indication:

section	Word count
Introduction	1301
Materials and Methods	3316
Results	4003
Discussion	1465

This manuscript contains 6 main figures, 16 supporting figures and 8 supporting tables

Content

Running title : bract natural variation

1 SUMMARY

• Bracts, the leaves subtending flowers, were lost multiple times in angiosperms, including
in *Brassicaceae*, where their development is blocked early. *Arabidopsis* mutants that
restore bract formation suggest that flower identity genes suppress the vegetative
program of bract development, but the exact mechanisms and the evolutionary origin of
bract loss remain unclear.

• We exploited natural variation in bracts that form only at the base of flowering branches
in *Arabidopsis*, to study bract development and its connection to floral transition. We
combined multiscale phenotyping, quantitative genetics, meristem imaging, time-series
transcriptomics and curve registration to capture the genetic and developmental
mechanisms unlocking bract development during floral transition.

• We mapped four Quantitative Trait Loci controlling bracts, excluding known bract mutant
genes. We demonstrated LEAFY and other flower regulators were not involved and
identified lists of new candidate genes and pathways, such as the anthocyanin pathway.
We found that bract develops when gene expression is desynchronised from the floral
transition, either later or earlier, revealing a more complex landscape than the previously
proposed prolonged vegetative state.

• We identified new mechanisms unlocking bract development. This natural variation
sheds a new light on development canalisation during floral transition and on bract loss
evolution.

21 5–8 key words

22 *Arabidopsis* natural variation, bract, evolutionary loss, floral transition, heterochrony,

23 quantitative genetics, time registration, time-series transcriptomics.

Introduction

Morphological evolution sometimes proceeds by losing structures or by regaining lost traits.

The genetic and developmental mechanisms leading to a trait loss or regain are being

elucidated (Cronk 2009; Sadier et al. 2022). Trait loss can simply result from gene(s) loss (Xu et

al. 2019). However, pleiotropy (one gene having different functions) can limit gene loss if it

causes detrimental effects and reduces fitness (Helsen et al. 2020). Also, a full genetic knock-

out of a trait may hinder its regain (Sadier et al. 2022). Hence, more studies are needed to

unravel the mechanisms behind the evolution of trait loss or regain.

In flowering plants, bracts provide an interesting model to address this loss and regain question

as they have been repeatedly lost and sometimes regained in different groups. Bracts refer to

any leaves developing in the inflorescence of angiosperms such as the edible leaves of an

artichoke. More accurately in botany, bracts are the leaves subtending a single flower, at the

junction of the floral peduncle with the stem (Dinneny et al. 2004; Endress 2006; Prenner et al.

2009). Bracts have evolved a wide diversity aligned to various adaptive functions:

photosynthesis, mechanical protection, pollinator attraction or seed dispersal. Some of these

adaptive changes have resulted in bract reduction or loss, like in *Poaceae* (grasses) or

Brassicaceae (cabbages). This loss may be an evolutionary adaptation to specific pollinators or

anemogamy, privileged resource allocation into reproductive versus vegetative organs or

developmental constraints (Whipple et al. 2010). To investigate the underlying genetic and

developmental mechanisms, several studies have been led in *Poaceae* (using rice, maize, wheat

and barley) and in *Arabidopsis thaliana* for *Brassicaceae*.

In these two groups, the reduced bracts are not strictly homologous. In *Poaceae* inflorescences

(ear, panicles or tassels), bracts are suppressed at major branching points while the bracts

closely associated with the floral unit (florets) are maintained (Whipple 2017; Xiao et al. 2022).

In *Brassicaceae*, the suppressed bract directly subtends the flower. Despite these differences,

two common points are shared. First, bract reduction is not total. Morphological deformations

(Kwiatkowska 2006) and localised gene expressions, such as *A. thaliana FILAMENTOUS FLOWER*

(*FIL*) or its maize ortholog *Zea mays yabby15* (*Zyb15*), indicate that a bract domain forms but

fails to outgrow (Long and Barton 2000; Dinneny et al. 2004; Whipple et al. 2010). In *A.*

thaliana, this domain is incorporated in floral tissues, remaining cryptic (Heisler et al. 2005;

Goldshmidt et al. 2008). Second, in both families, mutants can restore a bract from the cryptic

domain. This suggests that a gene regulatory network (GRN) actively suppresses bract

outgrowth in this domain. In *Poaceae* model plants, bract regain in mutants is often interpreted

as a heterochrony, or a change in developmental timing (Alberch et al. 1979). Bracts are

compared to leaves produced during juvenile phases, so the persistence of this developmental

program in the inflorescence suggests a 'prolonged vegetative phase' (Itoh et al. 1998;

Kawakatsu et al. 2006; Chuck et al. 2007; Kawakatsu et al. 2009; Chuck et al. 2010; Wang et al.

2021). This hypothesis, however, does not offer clear insights into the precise mechanisms

controlling bract regain in mutants, nor into the evolutionary scenario of bract loss in *Poaceae*.

Studies on bract mutants in *Poaceae* and *A. thaliana* have identified two GRNs controlling bract

suppression. In *Poaceae*, the GRN is organised around *NECK LEAF1 / TASSEL SHEATH1 / THIRD*

OUTER GLUME1 (NL1/TSH1/TRD1, or NTT) in rice, maize and barley respectively (Wang et al.

2009; Whipple et al. 2010; Houston et al. 2012). Loss-of-function mutants in grass species

produce bracts (Whipple et al. 2010; Wang et al. 2021), but not in *A. thaliana* mutants of the

orthologous gene *HANABA TARANU* (*HAN*) (Zhao et al. 2004). Upstream, NTT is directly

regulated by *SQUAMOSA PROMOTER BINDING PROTEIN LIKE* (*SPL*) transcriptional regulators

(Wang et al. 2009; Chuck et al. 2010; Chuck et al. 2014), themselves targeted by

microRNA156/529 (miR156/529) (Chuck et al. 2007; Wang et al. 2021). The miR156/SPL balance

regulates phase transitions from vegetative to reproductive stages in angiosperms (Wu and

Poethig 2006; Wang and Wang 2015), providing a genetic module controlling heterochrony in

plants (Buendía-Monreal and Gillmor 2018). This supports the interpretation that bract regain

in mutants is a heterochronic phenotype. Downstream, NTT influences genes involved in

hormone signalling, meristem and boundary identities, or leaf development (Xiao et al. 2022).

Yet, few direct NTT-target genes have been identified, such as the *PLASTOCHRON1* (*PLA1*) and

PLA2 genes in rice (Wang et al. 2021). These genes encode two enzymes whose function has

not been clearly identified (Miyoshi et al. 2004; Kawakatsu et al. 2006). Hence, NTT-

downstream mechanisms of bract repression need clarification.

In *A. thaliana*, bract mutants revealed different genes from the *Poaceae* orthologs, all

connected to the floral meristem identity (FMI) pathway. This genetic program specifies a floral

rather than a shoot identity (Weigel 1995). Impairing FMI leads to flower-to-shoot conversions

by restoring vegetative hallmarks into flowers: perturbed floral organ identity and phyllotaxis,

internode elongation, indeterminacy and also, bract outgrowth. In *A. thaliana*, FMI relies on a

complex gene regulatory network orchestrated by the transcription factor *LEAFY* (*LFY*)

(Siriwardana and Lamb 2012): *lfy* mutants fail to produce proper flowers and show diverse

bract outgrowths (Schultz and Haughn 1991; Weigel et al. 1992). Other FMI-related

transcription factors also cause bract de-repression when mutated alone or in combination,

suggesting they also act as bract-repressors: *UNUSUAL FLORAL ORGANS (UFO)* (Levin and

Meyerowitz 1995; Hepworth et al. 2006), *APETALA1* (*AP1*)(Irish and Sussex 1990; Bowman et al.

1993), *BLADE ON PETIOLE1 (BOP1)* and *BOP2* (Hepworth et al. 2005; Norberg et al. 2005; Xu et

al. 2010; Chahtane et al. 2018), *PUCHI* (Karim et al. 2009), *LATE MERISTEM IDENTITY 1 (LMI1)*,

AGAMOUS-LIKE 24 (AGL24), *SHORT VEGETATIVE PHASE (SVP)*(Xu et al. 2010; Grandi et al. 2012),

SUPPRESSOR OF CONSTANS 1 (SOC1), *SEPALLATA3 (SEP3)* (Liu et al. 2009), *FRUITFUL (FUL)*

(Melzer et al. 2008; Balanzà et al. 2014), *TERMINAL FLOWER 1 (TFL1)* (Shannon and Meeks-

Wagner 1991; Penin 2008) and *FILAMENTOUS FLOWER (FIL)* (Levin and Meyerowitz 1995; Sawa

et al. 1999; Siegfried et al. 1999). Fewer genes have been identified as promoting bract

outgrowth, such as *JAGGED* (Dinneny et al. 2004; Ohno et al. 2004), *AINTEGUMENTA* (*ANT*) and

AINTEGUMENTA-LIKE6 (*AIL6*) (Manuela and Xu 2024 Mar 28). In all these studies, bract de-

repression is accompanied by pleiotropic phenotypes affecting flowers, branching or flowering

time. Bracts, together with chimeric shoot-flowers, can also be induced in wild-type *A. thaliana*

under specific light treatments (Hempel and Feldman 1995; Hempel et al. 1998). Finally, the

genetic ablation of flowers induces bract outgrowths (Nilsson et al. 1998). Altogether, this

suggests that FMI establishment is responsible for bract repression, although the precise

mechanisms remain unclear.

To summarise, thanks to bract mutant studies, current models propose different GRNs to

acquire the ability to suppress bract, either from boundaries in *Poaceae* or from flowers in

Brassicaceae. In this latter case, it is unclear how FMI genes mostly expressed in flowers act

non-cell autonomously on the bract domain. Also, how this network acquired this new function

remains to be determined.

This study revisits bract inhibition in *A. thaliana* with natural variation. Important differences

suggest that the bracts naturally forming at the base of inflorescences involve developmental

mechanisms distinct from mutant bracts and that may be more similar to bract-bearing species.

Using two accessions, we mapped this heritable trait to four Quantitative Trait Loci (QTL).

Transcriptomic profiling of bract-producing meristems indicates that during floral transition, the

stage at which bract develops, the meristematic transcriptomes of the two accessions diverge

the most, with numerous desynchronisation of gene expression relative to the flowering event.

Challenging the model that FMI genes inhibit bracts, our results question whether changes in

the timing of gene expression may play a role in canalising phenotypes during floral transition

and in the evolution of bract loss.

Materials and Methods

Materials and Methods

Plant growth conditions

Seeds were sown on peaty-clay soil, stratified at 4°C for at least two days, and watered with

fertiliser (18-10-18 N-P-K) under LED lighting (sunlight spectrum NS12, 150 μmol.m-2.s-1).

Three different day/night regimes were used in the experiments: short-days (SD) with 8h light

and 16h dark; long-days (LD) with 16h light and 8h dark and continuous light (CL) with 24h light.

Temperature and humidity are controlled as follows: 22°C and 60% humidity during light for LD

and CL conditions, and 18°C and 70% humidity constantly in SD and during night time in LD. For

the Bulk Segregant Analysis and the RNAseq time course, plants were grown 20 days in SD

before switching to LD. For the RIL phenotyping assays, plants have been directly cultured in LD.

Plant materials

A. thaliana natural accessions and associated RILs were obtained from the Versailles

Arabidopsis Stock Centre (VASC), *Col-0*: 186AV, *Tsu-0*: 91AV, RIL set name: 3RV. F1 and F2

plants used for the BSA were generated by crossing the parents *Tsu-0* and *Col-0* in both

directions. The following strains were obtained from the NASC and are in *Col-0* background if

not otherwise mentioned: *ufo-1*: N16361 (*Col-2* background); jag-5d: N9506 (background: *Col-0*

gl1 pop1); *tfl1-13*: N6237; *tfl1-14*: N6238. *puchi-1*, *bop1-4* x *bop2-11* and *puchi-1* x *bop1-4* x

bop2-11 (Karim et al. 2009), *lfy-12* (Maizel and Weigel 2004), *soc1-2* (Michaels et al. 2003) and

svp x agl24 x soc1 (Liu et al. 2009) were described previously. Plants expressing

pLFY::2xmCherry-N7 ; 35S::Lti6b-YFP (in *Col-0* and *Tsu-0*) were generated for this study.

Tarenaya hassleriana and *Gynandropsis gynandra* seeds were kindly provided by Pr Eric Schranz

and Frank Becker, from Wageningen University (WUR, Holland). *Lunaria annua* and *Allaria*

petiolata plants are spontaneous specimens found outside the laboratory, in France. The

phylogeny of *Brassicaceae tribes* used in Fig. S1 is extracted from the Brassibase website

(Kiefer et al. 2014).

Plasmid constructions and plant transformation

<u>pLFY::2xmCherry-N7</u>: since no polymorphism was sequenced in the *LFY* promoter between *Col-*

0 and *Tsu-0* accessions, a sequence starting at -2277pb upstream of the ATG was amplified by

PCR   from   *Col-0*   genomic   DNA   using

5'GGGGACAACTTTGTATAGAAAAGTTGATCCATTTTTCGCAAAGG   and

5'GGGGACTGCTTTTTTGTACAAACTTGAATCTATTTTTCTCTCTCTCTATC primers. PCR fragments

were purified and inserted into pDONR P4-P1R with a gateway BP reaction. This plasmid was

then recombined in a three-fragment gateway reaction with 2xmCherry pDONR211 and N7-tag

pDONR P2R-P3 (containing the nuclear tag N7 and a stop codon) and the destination vector

pK7m34GW. <u>35S::Lti6b-YFP</u>: a pDONR P2R-P3 plasmid containing the sequence of the plasma-

membrane protein Lti6b (Cutler et al. 2000) was recombined in a three-fragment gateway

reaction with a 35S pDONR P4-P1R and the YFP pDONR P2R-P3 into the destination vector

pB7m34GW. Both constructs were then transformed into *Agrobacterium tumefaciens*

C58pMP90 strain by electroporation and transformed into both *Col-0* and *Tsu-0* plants by floral

dipping (Clough and Bent 1998). For each construct-by-genotype combination, several

independent transgenic lines were selected in T2 for a single insertion event based on 3:1

resistant:sensitive mendelian segregation of the resistance provided by the transgene. The

expression patterns of pLFY were reproducible between selected lines and matched published

data for *Col-0*. We were unable to get a 35S::Lti6b-YFP line with a membrane signal as strong in

Tsu-0 as in *Col-0* (Fig. 2 and S2). However, despite the weak YFP signal, the morphology of the

tissue could still be captured.

Microscopic meristem imaging and image analysis

Meristems were imaged using a Scanning Electron Microscope (Hirox 3000 SEM), or with a

confocal microscope Zeiss 700 LSM, according to the manufacturer's instruction and without

prior fixation. Multitrack sequential acquisitions were performed as follows: YFP, excitation

wavelength (ex): 488 nm, emission wavelength (em): 300–590 nm; mCherry, ex: 555 nm, em:

300–620 nm. Detection wavelengths and laser power were identical for *Col-0* and *Tsu-0*, PMT

voltage was identical for mCherry to allow comparisons of pLFY signal intensity in the two

genotypes. The YFP PMT voltage was optimised on each plant. Confocal images were processed

using imageJ (Schneider et al. 2012): maximum projections of mCherry channel and 16-bit-

transformed standard deviation projections of the YFP channel were merged in a composite

image. mCherry intensities were unchanged while brightness and contrast of the YFP channel

were optimised on each plant to provide the best morphological outlines of the tissues.

Macroscopic plant phenotyping

Macroscopic pictures were taken using different devices, according to the manufacturer's

instructions: Keyence VHX-900F, Canon EOS 450D camera, camera device of a Samsung Galaxy

A10 and a Ulefone Armor X5 pro.

Phenotypic measurements

Basal bract score was determined by counting all bracts in the main stem and cauline branches,

normalising by the total number of branches (excluding rosette branches, **Fig. 1E**); the

inspection was limited to the first five flowers, especially for mutants. The number of bracts

was counted after internodes elongation on the last upper cauline branch to ensure bracts

were visible to the naked eye. Flowering time was measured with different methods

(mentioned in the main text): the number of days from the start of transfer to in growth

chambers to the day of bolting, the day when the first flower blooms (open petals), or as the

cumulated number of leaves (rosette only or rosette and cauline). For plastochron

measurements, several plants of both genotypes were synchronised and grown in the same

condition. Each sampling day after transfer to a long day, 5 to 10 plants were randomly

dissected under a binocular dissecting scope, removing and counting all organs (leaves or

flowers) from the first leaf to the youngest organ visible on the main meristem. The youngest

organs were counted as soon as they were separated by a boundary (corresponding to stage-2

flowers as defined by (Smyth et al. 1990).

200 DNA extraction and sequencing for Bulk Segregant Analysis (BSA)

For the BSA, an F2 mapping population of 684 plants was generated by crossing (*Col-0* x *Tsu-0*)

in both directions. The bulk segregant analysis was split into four replicates. A 1cm² leaf sample

was sampled for each F2 plant, kept at -20°C and genomic DNA was extracted individually if the

plant was selected in one of the bulk. Genomic DNA of 56 and 17 plants were selected and

pooled in the bulks of low and high bract-score plants, respectively. The genome of the parental

lines was re-sequenced using genomic DNA from bulk *Tsu-0* and *Col-0* seedlings, respectively.

All genomic DNAs were extracted and purified using a CTAB-based protocol (cetyl

trimethylammonium bromide), following instructions as in (Healey et al. 2014).

Each DNA bulk was prepared by pooling the genomic DNA of selected plants in equal quantities,

to reach a final concentration between 13 and 25 ng/µl. Pooled DNA bulks and parental DNA

were used to prepare libraries and sequenced on BGISEQ-500WGS according to the

manufacturer's instructions, yielding 5 Gb data of 100bp paired-end reads per library (target

coverage of 40X).

214 DNA sequencing analysis and genomic variant analysis

Sequencing data consists of two parental plus two bulks of sequencing data. A genomic variant

analysis was performed on each dataset following the workflow of short variant discovery

previously described in (Besnard et al. 2020). This resulted in four gVCF files (one per sample)

generated by the HaplotypeCaller tool of GATK (v3.8, McKenna et al., 2010). TAIR10 was used

as the *A. thaliana* reference genome. Then, the two parental gVCF were first joint-genotyped

using GATK's tool GenotypeGVCFs to emit a common vcf file for the two samples. This file was

used to select a list of specific SNPs and small indels of *Tsu-0* (91AV stock) versus *Col-0* (186AV

stock), filtering for positions with coverage metric DP>10. This reference list of *Tsu-0*

polymorphisms was then used as the --dbsnp option in a second pass of joint genotyping using

all four gVCF as input (two parental samples plus the two bulks) to emit a common vcf file.

Finally, relevant polymorphic positions from the reference list in the two bulks were selected

after filtering for a depth ≥3.

227   **QTL mapping from Bulk Segregant Analysis**

228   QTL identification was carried out using the QTLseqr package (Mansfeld and Grumet 2018)

229   according to user instructions. After filtering data with the following parameters (refAlleleFreq

230   = 0, minTotalDepth= 10, maxTotalDepth = 90, minSampleDepth = 10, minGQ= 99,

231   depthDifference = 15), the deltaSNP index (Takagi et al. 2013) was generated and loci that

232   reached the ~95\% confidence interval were retained.

233   **QTL mapping using Recombinant Inbred Lines**

234   Genotyped RILs from *Col-0 x Tsu-0* (3RV) are publicly available in the VASC. Based on BSA

235   results, a panel of 55 RIL were selected from their known genotypes on chromosomes 1 and 5

236   using GGT 2.0 software (van Berloo 2008). The detailed genotypes of each line used in this

237   study are available in **Table S6**. The presence of basal bracts was quantified in each line using

238   the bract score. QTL mapping was performed with R/qtl software according to user instructions

239   (Broman et al. 2019): we used the scanone function with the mean value of the bract score for

240   the 55 tested RILs as a trait and significant thresholds were computed by setting a permutation

241   number to 2000. For the bract score, the "non-parametric" (np) statistical model was used

242   while for paraclade number, we used a normal model implemented with the hk method. To

243   look for QTL interactions, the scantwo function was used with the normal model and the hk

244   method since the np model is not implemented for this function. MQM was performed with

245   default parameters. The (broad-sense) heritability for the bract score was defined as $H^2 = (var_T -$

246 $var_E) / var_T$, with var_T and var_E being the total and environmental variance, respectively. var_T

247   was computed as the total variance of the bract score of each plant over the entire QTL

248   mapping dataset (1830 RIL plants and 780 control parents split over 12 experiments) while var_E

249   was computed as the pooled variance of the line variances (computed over all plants of a line

250   across experiments). Confidence intervals of H^2 were provided by bootstrapping 70% of the

251   strain 1000 times.

252   **KASP genotyping**

253   We selected 19 new genotyping markers from the 140SNPvCol marker set (Lutz U. et al. 2017)

254   to cover the two large QTLs mapped in chr1, 1a and 1b (**Fig. S5B**) and corresponding

255   kompetitive allele-specific PCR (KASP) oligos were ordered to Biosearch Technologies™ LGC Ltd.

To further reduce the intervals, we designed two new SNP markers targeting *Tsu-0*

polymorphisms identified in our whole-genome re-sequencing data. Specific KASP oligos (three

per SNP) were designed by LGC from the 100-bp sequence surrounding the SNP. All 21 new

markers (**Table S7**) were validated on parental genomic DNA (*Col-0* and *Tsu-0*) in a KASP assay

before their use with recombinant inbred lines. Clean genomic DNA from CTAB-based

extraction was used for all samples. For each marker, 2.5 µL of DNA (diluted to a final

concentration of 5 ng/µL), 2.5 µL of KASP-TF V4.0 2X Master Mix low ROX and 0.07 µL KASP

assay mix were mixed in 384-well plates and the genotyping assays were run in a QuantStudio 6

Flex (Applied Biosystems), using standard user guidelines for thermal cycling and final

fluorescence analysis.

Fine mapping using Heterologous inbred families

Using F6 genotyping data for the 3RV RIL set from the VASC, we selected 3 RILs heterozygous in

a region overlapping QTL1a, QTL1b and the interval in-between: RIL 334, 488 and 478,

respectively (**Fig. 3** and **Fig. S5**). For RIL 334 and 488, we genotyped by KASP 160 segregating F8

progenies and selected 5 and 8 plants, respectively, with homozygous recombination inside the

QTL region under study. For RIL 478, we genotyped 20 plants and selected only one; genotyping

reactions were performed using PCR primers designed to amplify the published marker of the

VASC (Simon et al. 2008)(**Table S8**) and the results were read by Sanger sequencing. All selected

F8 plants were selfed to generate an inbred line of a fixed genotyped (F9), the collection of F9

inbred lines forming a Heterologous Inbred Family (HIF). HIF lines (with at least 20 plants per

line) were phenotyped for the bract score with parental control in the same experiment.

Biological sample preparation for the transcriptome time-series profiling

After 20 days in SD, *Tsu-0* and *Col-0* meristems were dissected every day in LD conditions, in

order to capture the precise developmental stages (especially stages T and F, see below). The

mutants *lfy-12*; *puchi1 x bop1 x bop2*; and *jag5-D* meristems were dissected every day starting

from 1 week after LD transfer to target the stage F. Three independent biological experiments

were performed with 5 to 11 meristems per replicate. Stages are defined as follows: V, the day

when plants were transferred to LD from SD; L, after 4 to 5 days of LD (identical for both *Col-0*

and *Tsu-0,* which have the same meristem shape at this time); T, the main meristem enlarges

and domes, (Kwiatkowska 2008; Kinoshita et al. 2020), and lateral meristems starts being visible

at the axils of young leaf primordia; F, all young organs in the meristems were identified as

flowers (note that this stage occurred several days before bolting). Stage F in mutants was

defined when several rounded primordia become visible at the SAM (that will become branch-

like flowers or only branches). Dissections were performed from 9:00 a.m. to noon, by

alternating between *Tsu-0* and *Col-0*. Micro-dissected meristems were snap frozen in liquid

nitrogen and stored at -80°C until RNA extraction. To avoid the induction of stress-related gene

expression, meristem dissection did not exceed 3 min between the first organ removal and

freezing. For each replicate, pooled meristems were ground with a RockyII tissue lyser

according to the manufacturer's instructions. RNA was extracted using the PicoPure RNA

Isolation Kit (Arcturus, Catalog KIT0202) according to the standard protocol. RNA concentration

ranged from 3 to 64 ng/µl (average 23ng/µl), with a RIN value between 5.6 and 7.6 (average

6.8).

RNA-sequencing

Library preparation was made using NEBNext® Ultra™ II Directional RNA Library Prep Kit for

Illumina (New England Biolabs); NEBNext® Poly(A) mRNA Magnetic Isolation Module (New

England Biolabs); and NEBNext® Multiplex Oligos for Illumina (Index Primers Sets 1, 2 et 3),

from 40ng of RNA and sequenced using the NextSeq500 (Illumina). Sequence quality was

controlled using the Sequencing Analysis Viewer, sequences that did not pass the quality filter

were removed. Following QC an average of 43 million sequences per sample was achieved.

RNA-seq analysis (pre-processing procedures)

Quality filtering using Trimmomatic (Bolger et al. 2014) retained for all samples 96% of the

reads, which were aligned to TAIR10 genome using STAR (Dobin et al. 2013) with the following

parameters: non canonical splice junctions were removed, multi-mapping reads were limited to

10, and only reads mapped once were considered to determine splicing junctions. Between 88

and 97% of the reads were mapped to a simple locus. Normalisation of read counting was

performed using the R Bioconductor packages DESeq (Love et al. 2014), with the following

parameters: genes for which the total number of reads was below 10 were discarded, and data

were transformed with Variant Stabilising Transformation (VST) function (Anders and Huber

2010), to harmonise the variance. Consistency between biological replicates was verified using

a Principal Component Analysis on all samples with VST-transformed data.

RNA-seq analysis: Differential Gene Expression (DGE) analysis

Differential analysis was performed using the R Bioconductor package edgeR (McCarthy et al.

2012). Reads were first normalised using TMM (Trimmed mean of M-values) to reduce library-

specific biases. Normalisation factors were between 0.94 and 1.049. A generalised linear model

was applied for the analysis of differentially expressed genes (DEG). Three types of DEG analysis

were considered: DEG at each stage between the different genotypes, DEG between the stage

within the same genotype, and DEG across all conditions (stages and genotypes). Multiple DEG

analyses were corrected using Benjamin-Hochberg correction, and genes with a p-value < 0.05

were retained.

RNA-seq analysis: clustering gene expressions correlating with bract presence

To identify candidate genes that may act as positive or negative bract regulators (see **Figure**

5G), we first calculated the minimum and maximum expression levels among stages with and

without bracts. If the minimum expression level in the stages with bracts was greater than or

equal to the maximum expression level in the stages without bracts, the genes were considered

positive bract regulators. If the maximum expression level in the stages with bracts was lower

than or equal to the minimum expression level in the stages without bracts, the genes were

considered positive bract regulators. If neither of these conditions were satisfied, genes were

discarded.

Temporal registration of gene expression dynamics

To align gene expression profiles of *A. thaliana Tsu-0* (query data) with *Col-0* (reference data),

we utilised the curve registration method in the R package greatR (Kristianingsih 2024). This

approach involved shifting the gene expression profiles of *Tsu-0* across developmental stages

(V, L, T, F) using shift factors ranging from -1 to 1; a stretch factor was not applied due to the

two datasets being over similar times. The optimal shift parameter for each gene pair was

identified by maximising the log-likelihood. The fit with the best registration factors for each

gene was then compared to the fit with a non-registered model (without transformation) using

the Bayesian Information Criterion (BIC) statistic. A lower BIC score for the registered   model

(with transformation) indicated that the gene expression dynamics of *Tsu-0* and *Col-0* were

similar, and the expression profile differences could be resolved through registration.

Conversely, a higher BIC score suggested that the profiles were best described by two individual

curves, i.e. not aligned. Prior to transformation, gene expression levels in both datasets were

centred and scaled using the z-score scaling method. The standard deviation value for the

replicates at each time point was set to 0.01.

GO terms analysis

GO term enrichment analyses were performed using either the 'Single Enrichment Analysis'

tool from AgriGOv2 (Tian et al. 2017) with default parameters (Fisher's test with Yekutieli

adjustment method) or clusterProfiler 4.0 (Wu et al. 2021) with default parameters (enrichGO

and simplify functions with "BH" adjustment method). The Rich factor was computed by

dividing the number of genes associated with a GOterm ("Count") by the numerator of the

"BgRatio" computed by the enrichoGO function.

Gene selection inside genetic mapping intervals

The R library 'GenomicFeatures' (Lawrence et al. 2013) was used to import gene information

from the most recent annotation file of *A. thaliana* at the gff format (Cheng et al. 2017). Custom

R scripts were used to intersect all gene loci falling within genetic mapping data and provide for

each gene information from RNA-seq analysis and genomic variant analysis (see Table S1 and

Table S2). We used snpEff v5.0d (Cingolani et al. 2012) and its putative functional categories

(HIGH/MODERATE/LOW/MODIFIER) to predict the functional impacts of the genomic variants

over regions covering each annotated gene of the interval, including promoter (2 kbp upstream

of the transcriptional start site) as well as 500-bp downstream region in the case of miRNA. In

Table S1 and S2, additional gene information was retrieved from ThaleMine (Pasha et al. 2020)

and GO terms from the TAIR bulk data retrieval tool.

Data handling, data visualization and descriptive statistics

The R software was used (R Core Team 2018), especially the following packages: Bioconductor

packages (Huber et al. 2015) for the analysis of omics data, especially GenomicFeatures, IRange

(Lawrence et al. 2013), rtracklayer (Lawrence et al. 2009), clusterProfiler (Wu et al. 2021),

limma (Ritchie et al. 2015), DESseq2 (Love et al. 2014), edgeR (Chen et al. 2016) and

org.At.tair.db (Carlson 2024); dplyr (Wickham et al. 2023), tidyr (Wickham et al. 2024) and

reshape2 (Wickham 2007) for data manipulation and ggplot2 for almost all plots (Wickham

2016), with viridis and plasma (Garnier et al. 2024) for some colour optimization.

Text and english

Language correction software with artificial intelligence was occasionally used throughout the

text to ensure correct spelling and grammar as well as the most comprehensible style possible.

Results

Transient bract formation during the floral transition is common among A. thaliana

accessions

The typical *Brassicaceae* flower has no bract. Yet, a number of *Brassicaceae* species produce

some bracteate flowers (i.e. with bracts) at the base of their flowering branch (a raceme) (**Fig.**

1A and **Fig. S1 A-D**), a common botanical trait in this family (Endress 2006; German et al. 2023).

Hence, bract loss in *Brassicaceae* is incomplete, showing a reduction trend compared to its

sister clade *Cleomaceae* (**Fig. 1A**, **Fig. S1E**). Mapping the presence of bracts among the

Brassicaceae phylogeny does not reveal an evolutionary scenario (**Fig. S1A**). Some species

display bracts up to the first half of the raceme (e.g. **Fig. 1A**), while others only have one or two

bracts at the base (**Fig. S1 C, D**), e.g. in (Al-Shehbaz 2015). While the reference *Col-0* accession

of *A. thaliana* bears no bracts, we discovered that the natural *Tsu-0* accession displays bracts on

the first one to five flowers of the raceme (**Fig. 1A**). Under a stereomicroscope, a gradient of

bract outgrowth is visible, from fully developed bracts resembling the cauline leaves associated

to branches, up to filamentous, short rudimentary structures (**Fig. 1B**). Using scanning electron

microscopy, a mild swelling of the peduncle basis could sometimes be observed in *Tsu-0*, unlike

Col-0 smooth peduncle (**Fig. 1C**). In *Tsu-0*, bract development can thus be released to varying

degrees in the first flowers of a branch, which we have termed "basal bracts". Other genetically

diverse natural accessions produce basal bracts, albeit at a highly variable rate (**Fig. 1D**) and

without any clear correlation to geographic or genetic origins (**Fig. 1D, Fig. S1 F, G**).

Interestingly, *Col-0* also produces basal bracts at a low frequency, typically on lateral branches.

In general, we observed that basal bracts can develop at every flowering branch but with a

variable frequency, even within an accession and in fixed environmental conditions. To quantify

basal bracts, we

Figure 1: **The presence of basal bracts is a common, natural trait in *Arabidopsis thaliana* with quantitative variations among genetic backgrounds**. **A**, Examples of bracts (red arrows) in different angiosperms. *Brassicaceae* are mostly bractless but some species retain bracts at the base of inflorescence branches (e.g. in *L. maritima*). In *A. thaliana*, some natural accessions display basal bracts (e.g. *Tsu-0*) while others do not (e.g. *Col-0*). Schematic phylogenetic relationships are indicated with a cladogram below the pictures. White stars: first ebracteate flowers following previous bracteate flowers **B**, (from left to right) *A. thaliana*'s basal bracts can be true leaves or just small rudimentary filamentous structures at the base of the floral pedicel (red arrows). These structures are absent in younger flowers, as in the reference *Col-0* accession (rightmost panel, black arrowhead). **C**, Details of basal

bracts by scanning electron microscopy in a *Tsu-0* inflorescence tip (left panel), showing a bract on the first flower (red arrow) and swollen base of pedicels on the two following flowers (red arrowhead). Right panel: pedicels of *Col-0* plants do not show such structures (black arrowhead). **D**, Occurrence of basal bracts in different accessions, assessed by the percentage of plants with at least one basal bract in the inflorescence. Each dot is the average value of several plants (number indicated by dot size) of a scoring assay. A box plot indicates several assays per line and thicker horizontal black lines are the median value of all scoring assays (range 1-3) in the accession. *Col-0* and *Tsu* are highlighted in green and red, respectively. The geographical origin of each strain is located in the world map below. **E**, Definition of a quantitative bract score for a single plant (see main text for detail). **G**, Quantification of basal bracts in different accessions using the bract score.

defined a bract score (**Fig. 1E**) which captures both intra- and inter-genotype variations (**Fig.**

1F). *Tsu-0* and *Col-0* ranked respectively as high and low bract producers and were kept for

further investigation.

Basal bracts develop with wild-type flowers and differ from known mutant bracts

To investigate basal bract development, we imaged the emergence of the first flower. Scanning

electron microscopy revealed a sharp floral transition in *Col-0*: the last lateral meristem

subtended by a cauline leaf is immediately followed by ebracteate (i.e. bract-less) flowers (**Fig.**

S2A-B). In *Tsu-0*, the first flowers produced after the last branch bear bracts or rudimentary

bracts (**Fig. S2C-D)**. Their association with young flowers indicates that these bracts are not

secondary outgrowth from mature floral peduncles. In bract-making species like the sister clade

Cleomaceae, we observed that bract emerges before the flower (**Fig. S2E**). In *A. thaliana*, the

abaxial position of bracts and their precocious development suggest that they are true de-

repressed bracts. Given LFY's central role in suppressing bract development in *A. thaliana*, we

question whether bracts form due to LFY perturbation at floral transition. Using a p*LFY* reporter

line, we observed in both genotypes the same sharp activation of *LFY* transcription from the

first flower onwards (**Fig. 2A-B**), regardless of bract presence in *Tsu-0* (**Fig. 2B-C**). Accordingly,

Tsu-0 bracteate flowers were wild-type, with a normal number of floral organs (**Fig. 2D**). This

contrasts with reported bract mutants which presented severe phenotypes, such as loss of

floral organs or lack of determinacy with branched flowers (**Fig. S3**). Interestingly, *pLFY* was also

activated in the bract margins, albeit at lower

Figure 2: **Flowers bearing bracts express high levels of LFY and do not display mutant phenotypes**

A, B. Confocal live imaging of the main meristems of *Col-0* (**A**) and *Tsu-0* (**B**) plants at floral transition, expressing a pLFY transcriptional reporter (magenta) and a membrane marker (green) for morphology (top-view projections of stack acquisitions). Green arrowheads: branches, ordered with decreasing numbers from the floral transition; white arrows: flowers ordered with increasing numbers from the floral transition. In both genotypes, pLFY expression is absent from branch meristems and suddenly appears in the first flower onwards. Representative pictures of at least 6 plants per genotype captured at floral transition.

C, side-view of the first flower from image B (*Tsu-0*): a bract (red arrow) is visible on the abaxial side of this young flower.

D, Number of floral organs in flowers with (dark red) or without (light red) bract in *Tsu-0* plants.

E, F. Side views by confocal live imaging of *Tsu-0* flowers with bracts (red arrow) expressing the pLFY reporter (magenta) and the green morphological marker. Flowers show two developmental stages older than C and with increasing age from E to F, as shown by the growing abaxial sepal on top of the flower. pLFY is expressed at low levels in the bract margins (magenta arrow).

Scale bars: 50 µm

levels than in the floral meristems (**Fig. 2E-F**), contrary to the proposed role of LFY as a bract

inhibitor. To better understand LFY contribution to basal bracts, we then re-examined bracts in

mutants with lowered *LFY* expression: *lfy*, *ufo* (Hepworth et al. 2006) and *puchi x bop1 x bop2*

(Karim et al. 2009). As previously reported (Chahtane et al. 2018), bracts in these mutants were

at the tip of old branches and more frequently on secondary than on main shoots (**Fig. S4 A-G**).

In *Tsu-0*, however, basal bracts were limited to the first flowers at floral transition (**Fig. S4H**),

with the same frequency on each shoot (**Fig. S4I**). In addition, at floral transition, these mutants

produced branches typically lacking cauline leaves, called I* branches in the literature (Ratcliffe

et al. 1998), which have been interpreted as flower-to-shoot transformations. The first

determinate flowers coming after I* branches were also ebracteate (**Fig. S4 B, C**). Hence, at

floral transition, loss of FMI gene function is sufficient to lose floral fate but not to restore leaf

development, unlike at branch tips, suggesting *LFY* effect on bract formation is context-

dependent. We further tested the effect of intermediate LFY expression levels by scoring basal

bracts in *LFY//lfy-null* heterozygous plants (**Fig. S4J**). The meristem was sensitive to halving the

genetic dose of *LFY*, as shown by the slight increase in the number of branches. However, all

basal flowers were wild type and ebracteate, confirming that this range of variation in LFY levels

does not affect basal bract formation. Other bract mutants rarely produce bracts at the base of

inflorescence but either on each node (e.g. in *JAGGED* gain-of-function mutant) or at the tip of

branches (**Fig. S5**). Although *tfl1* mutants produce frequent basal bracts, these are likely cauline

leaves that initially subtended branches, which were later transformed into flowers, as

suggested by the presence of a few lateral branches (**Fig. S5**). These phenotypic differences

suggest the genetics of bract development in mutants differ from those in *Tsu-0*, which may be

influenced by the particular context of floral transition. This first report of bracts associated

with wild-type flowers in *A. thaliana* illustrates that flower and bract formation are not

necessarily incompatible in this species, questioning the underlying developmental

mechanisms.

Basal bract frequency is unaffected by variation in photo-induction or plastochron

length, but shows complex correlations with flowering time

To characterise factors governing basal bract formation, we evaluated the effects of photo-

inductions as reported by (Hempel and Feldman 1995; Hempel et al. 1998). In these seminal

experiments, basal bracts were induced together with graded phenotypes of chimeric

Figure 3: **Variation in natural basal bract frequency is not correlated with branch position, plastochron length or light conditions, but with variation in flowering time**

A, Correlation study between the number of lateral (cauline) branches and the number of bracts on the main stem only in *Col-0* and *Tsu-0* plants (green and red, respectively). (Pearson test, p-value > 0.1).

B, Effect of different light regimes on bract scores in *Col-0* and *Tsu-0* plants (green and red, respectively). LD and SD stand for long- and short-day conditions, respectively and CL stands for continuous light (see methods). Numbers (e.g. 20SD > LD) indicate the number of days in the first condition before transfer to the second. The number of plants scored is indicated below each bar.

C, Cumulated production of organs over time for *Col-0* and *Tsu-0* plants counted in microdissected shoot apices (green and red, respectively, each time point averaging 5 to 10 plants): the time window when plants make the floral transition is indicated by a horizontal arrow and vertical dashed lines. The rate of organ production (inverse to the plastochron) in this period is computed from the local slope of the curve. Plants were cultured for 21 days in short-day conditions before transfer in long-day (LD) conditions (see methods).

D, Result of three independent experiments of plastochron measurements at flowering

transition (see also fig. S5).

E, Differences in flowering time between *Col-0* and *Tsu-0* accession measured as bolting time in days (left) or as the number of vegetative nodes (rosette leaves and cauline leaves/branches) before the first flower on the main stem. Plants grown in LD conditions, N = 58 plants per genotype in two independent replicates.

F, Correlation study between the bract score and the flowering time (assayed when the first flower opens) in different accessions, labelled with different colours. Each dot is a plant and a linear regression standard deviation is computed for each accession.

shoot-flowers in natural accessions. These results lead to the "conversion" model, where a

strong photo-induction converts a young branch meristem (already subtended by a leaf) into a

bracteate flower. However, in *Tsu-0*, we never observed such shoot-flowers chimaeras (**Fig. 1, 2**

and S2) nor a negative correlation between the numbers of bracts and cauline branches (**Fig.**

3A), supporting that *Tsu-0* bracts are not linked to shoot-to-flower conversions. While Hempel

and Feldman observed flower formation in hours after their strong photo-induction, our floral

transition occurred at least one week after transfer to long days (**Fig. 3C**), suggesting milder

photo-inductive signals in our growth conditions. However, further varying photo-inductive

conditions did not affect the bract score in *Tsu-0* nor in *Col-0* (**Fig. 3B**). In the conversion

hypothesis, bract formation is promoted by shorter plastochrones, the time between two

lateral meristem initiations. Indeed, young meristems can be converted into flowers during a

short time window: with short plastochrones, young branch meristems sensitive to floral

conversion are frequently produced. In addition, the *plastochron1/2/3* mutants in rice have a

shorter plastochron (Miyoshi et al. 2004; Kawakatsu et al. 2006; Kawakatsu et al. 2009) and

make bracts. We evaluated precisely the plastochrones of *Col-0* and *Tsu-0* plants during the

floral transition by micro-dissecting shoot apices in time series. Despite the variability, we found

no consistent shorter plastochron in *Tsu-0* compared to *Col-0* (**Fig. 3C, D; Fig. S6A**). A clear

difference between *Tsu-0* and *Col-0* is their flowering time, which happens in older *Tsu-0* plants

in absolute time or in developmental time, and consistently across conditions (**Fig. 3E** and **Fig.**

S6B). A panel of five accessions showed a global positive correlation between flowering time

and bract score when comparing genotypes (**Fig. 3F**). However, within an accession, the

correlation can be negative (e.g. in *Ler* and *Tsu-0*), suggesting more complex relationships with

flowering time. Altogether, our results indicate that basal bract formation in *Tsu-0* is not linked

to variation in light or productivity (plastochrone) but may be influenced by flowering time.

Basal bract development is controlled by multiple QTLs suggesting new genetic

mechanisms

To identify the genetic determinism of basal bracts, we conducted quantitative genetics using

crosses between *Tsu-0* and *Col-0*. F1 progeny showed intermediate bract scores while F2 bract

scores spread within the parental range, with a distribution that does not indicate a simple

monogenic trait (**Fig. 4A and Fig. S7A**). Bulk Segregant Analysis (BSA) with mapping-by-

sequencing identified four quantitative trait loci (QTL): two on chromosome 1 and two on

chromosome 5 (**Fig. S7B, C**). The large spread of F1 phenotypic values constrained F2 plant

selection with few plants in the high-score bulk and possibly spurious homozygous in the low-

score bulk. These two limits may explain the large intervals obtained. To overcome the

uncertainty on the phenotypic value attributed to an F2 genotype, we then used Recombinant

Inbred Lines (RIL), which allowed averaging of bract scores across isogenic plants. Using an

existing RIL set (Simon et al. 2008), we measured the average bract scores of several lines (**Fig.**

S8A) for which we checked the absence of recombination bias (**Fig. S8B**). The heritability of

bract score was high among RIL (65.45% +/- 1.7) and single QTL scanning mapped four peaks:

two most significant on chromosome 1 (named 1a and 1b) and two less significant on

chromosome 5 (named 5a and 5b; **Fig. 4B**). Consistent with the BSA result, this analysis

provided slightly different QTL positions and shorter intervals. A two-QTL scan suggested

additive effects between QTLs 1a and 1b (**Fig. S8C**). To confirm and reduce the intervals of these

two major QTLs, we used two RILs identified as heterozygous in a region overlapping with these

QTLs at F6 generation (**Fig. 4C-H**). F8 plants from these RIL were re-genotyped at a higher

density with new genetic markers (**Fig. S9A, B**) to select recombining intervals homozygous for

either allele of the two parents (**Fig. 4D, G**). Selected plants founded new lines forming a

heterologous inbred family (HIF) with identical genotypes except in a small interval, which

allows testing the effect of alleles in that region only. Scoring bract scores among HIF confirmed

Figure 4: **Identification of 4 major QTLs controlling basal bract formation in *Tsu-0* suggests unknown genetic pathways**. **A**, Genetic transmission of bract score in F1 and F2 hybrids from a cross between *Col-0* and *Tsu-0* parent plants. **B**, QTL mapping for bract score in a set of RILs identifies four putative QTLs, two in chromosome 1 (1a and 1b) and two in chromosome 5 (5a and 5b). Dotted horizontal red lines indicate three different significance thresholds computed from 2000 permutations. On top of the graph, H^2 indicates the (broad-sense) heritability of the bract score computed among the RIL set. **C**, Genotype of the F6 generation of the line RIL334 in chromosome 1, showing a heterozygotic region overlapping with QTL1a. **D**, Chromosome 1 genotypes of selected lines forming a heterologous inbred family (HIF) obtained from RIL334 after two more generations of selfing (F8). **E**, Box plots of bract scores of the different lines from the "334 family" (blue boxes) with parental controls (*Col-0*: green, *Tsu-0*: red, N > 20 plants per line). Segregating bract scores within the family map a narrower region, QTL1a*, spanning 541 annotated genes. **F**, Genotype of the F6 generation of the line RIL488 in chromosome 1, showing a heterozygotic region overlapping with QTL1b. **G**, Chromosome 1 genotypes of selected HIF lines obtained from RIL488 at F8

and (**H**) a box plot of their bract scores (blue boxes), with parental controls (*Col-0*: green, *Tsu-0*: red, N > 21 plants per line). This maps a narrower region, QTL1b*, spanning 332 annotated genes. In E and H, lines not sharing the same letter(s) are statistically different (posthoc Tukey analysis with 0.05 sign. level from a glm of bract scores fitted with a quasi-poisson distribution). **I**, Final genetic mapping of the 'basal bract' trait in *Tsu-0* and the location of genes reported to impact bract development in previous mutant studies.

both QTLs and mapped them to smaller intervals named 1a* and 1b*. Additive effects of QTL 1a

and 1b appeared when *Tsu-0* allele in 1b* was combined with *Col-0* allele in 1a*, the bract

score reaching about half of the *Tsu-0* parent bract score (see in 488 families, **Fig. 4G, H**).

However, *Tsu-0* parent score is tied only when both QTLs bear *Tsu-0* alleles (see in 334 families,

Fig. 4D, E). As a control, we generated a HIF to test the region between 1a and 1b and

demonstrated that it did not influence the bract score (**Fig. S9C-E**). However, mapped intervals

remained large (4.1 to 11.8 cM), containing many genes (332 to 931, 2324 annotated genes for

the four QTL, **Fig. 3I** and **Table S1**). The parental accessions that we re-sequenced differ by over

770,000 small genomic variations (SNPs and indels), with more than one variation every 175 bp

on average, leaving few invariant genes to exclude (**Table S1**). Interestingly, known bract-

related genes were absent from the QTLs (**Fig. 4I**).. However, a careful RIL phenotype inspection

provided additional hints into the developmental pathways possibly involved in *Tsu-0* basal

bract formation. First, we observed that the cauline branch number positively correlates with

the bract score and QTL1b overlaps with a QTL controlling cauline branch number (**Fig. S10**).

This suggests that the gene controlling basal bracts in QTL 1b* might also promote more cauline

branches. Second, two transgressive, partially penetrant phenotypes appeared in the RILs: the

bract position often shifted to a mid-peduncle position (**Fig. S11, A-F**) and signs of incomplete

floral determinism (mostly branched flowers) were observed (**Fig. S11, N, O**). The shifted bract

position resembles flower-preceding prophylls in some natural species (Endress 2006; Prenner

et al. 2009). Interestingly, this phenotype is frequent in certain bract mutants like *soc1*, *tfl1* and

even systematic in *bop1 x bop2* (**Fig. S11 H-M**). However, this trait did not correlate with any

specific allelic combinations among the HIF families (**Fig. S11G**). Indeterminism in HIF 488 was

associated with *Tsu-0* QTL 1b* and *Col-0* haplotypes in other mapped QTLs (**Fig. S11P**). Other

indeterminism cases occurred in HIF 334, but without consistent genetic combinations of

mapped QTLs. Such transgressions suggest that complex genetic interactions may be required

in *Tsu-0* to ensure proper bract development and positioning with a wild-type flower. To

conclude, our data demonstrate that the genetic control of basal bracts in *Tsu-0* is complex and

relies on several genes having additive effects on bract development and epistasis on other

traits.

Transcriptomics of the floral transition suggests new pathways associated with bract

development

To identify genes whose expression changes with bract development, we profiled meristem

transcriptomes over the floral transition in both accessions. Microdissected meristems were

matched to the same four developmental stages: vegetative, late vegetative, transition and

floral (V, L, T and F, respectively, **Fig. 5A**). This ensured that, despite different absolute

flowering times (**Fig. S12A**), the transcriptomes of the two accessions were realigned using the

floral transition as a common developmental clock. After RNAseq, principal component analysis

(PCA) validated replicate consistency (**Fig. S12B**) and the precision of the staging was confirmed

by the synchronised expression between accessions of genes such as *FT*, *LFY* or *AP1*, key

regulators of floral transition and flower identity (**Fig. 5B,** see also **Fig. S12C** for other reference

genes). For LFY, this RNAseq data supported our results from the pLFY transcriptional reporter

lines (**Fig. 2**) and genetic studies (**Fig. S4**), indicating that *LFY* is not involved in *Tsu-0* basal bract

development. The stage T stood out as a critical transition, with the highest number of gene

expression changes in both accessions (**Fig. 5C**). At stage T, *Col-0* and *Tsu-0* meristem

transcriptomes also diverged the most, with up to 4,759 genes differentially expressed (DE)

(**Fig. 5D**). Bract and flower initiations start before any morphological event (Heisler et al. 2005).

Given the delay between stages T and F (a median of 1 to 2 days, **Fig. S12A**) and the

plastochron (**Fig. 3B, C**), the first flowers and their bracts must be initiated at stage T. Thus,

transcriptional changes associated with bract formation in *Tsu-0* should transiently appear at

stage T and progressively fade by stage F since only the 1 to 5 first flowers present a bract (**Fig.**

1C, **Fig. S4H**).

Figure 5: **Transcriptomic divergence between *Tsu-0* and *Col-0* meristems peaks at the**

Since GO terms yielded no informative clues from the high number of DE genes at stage T (**Fig.**

S14A), we first examined whether known bract-related genes could be affected, although we

knew from our mapping that they cannot be at the origin of basal bracts in *Tsu-0* (**Fig. 4I**). Only

PUCHI, *SOC1* and *TFL1* significantly differed at stage T between *Tsu-0* and *Col-0* (**Fig. 5E** and **Fig.**

S12D) but these differences persisted until stage F even though bracts disappeared with older

flowers, suggesting that the variation of these gene expressions is not essential in bract

formation.

To discover genes without *a priori*, we then performed a comparative transcriptomic approach

with additional mutant data. Since *lfy* and *puchi x bop1 x bop2* mutants stop making leaves at

floral transitions (I* branches, **Fig. S4B, D**) while *jagged-5d* plants always produce a bract,

transcriptomic cues for bract development may only be present in *jagged-5d* meristems and

not in the other two backgrounds. As expected from their genetic background and

developmental stage, these mutants clustered with *Col-0* stage F in a PCA (**Fig. S12B**). By

comparing DE genes of each mutant with *Col-0*, DE genes specific to jagged-5d were isolated

(**Fig. S13A**). This set revealed an enrichment for biological processes related to shoot, phyllome

development and photosynthesis, possibly linked to persistent bract formation (**Fig. S13B**). Only

413 of these genes were shared with DE genes of *Tsu-0* at stage T, with Go terms pointing again

to chloroplast functions and less expectedly, to metal ion transport and homeostasis (**Fig. S13C,**

D). If one of the causative genes mapped in *Tsu-0* (**Fig. 4, Table S1**) was differentially expressed

(so not an allele acting post-transcriptionally) and shared with *jagged-5d*, it should be contained

in this short list. Cross referencing transcriptomic and QTL-mapping datasets yielded 33

candidate genes (**Fig. S13E, Table S2**). None of them is known to be linked with bract

development or with flowering in general. Although these particular genes should be validated,

this mutant comparison suggests that new uncharacterized genetic pathways may be involved

in the development of bracts.

In a third approach to capture genes involved in bract formation, we specifically took advantage

of the time-series information. Briefly, expression data were gathered in two groups

corresponding to either bract-less (*Col-0* T, F, and *Tsu-0* F) or leaf/bract-producing meristems

(the other samples, including *Tsu-0* T) and we selected genes whose expression profile clusters

these two groups apart (**Fig. 5G**). Only 124 genes met the criteria of such "putative bract

regulators" (**Fig. 5H, Table S3**), including *SOC1*, as a putative negative regulator. Just 19 genes

overlapped with the genes also DE in *jagged-5d* (**Fig. S13C, Table S3**), without pointing to

particular biological functions. Instead, two enriched GO-terms emerged from these putative

bract regulators: anthocyanin biosynthesis and salicylic acid (SA), associated with up- and

down-regulated genes, respectively (**Fig. S14D-F**). Backtracking these ontology terms for all DE

genes at stage T (**Table S4** and **S5**) retrieved more genes associated with these pathways,

supporting that their activity levels differ between *Tsu-0* and *Col-0* at the critical stage T (**Fig.**

S14I). High anthocyanin production in *Tsu-0* at stages L and T was evident from the frequent

purple coloration just below the meristems, contrasting with the green tissues in *Col-0* (**Fig. 5I**).

Sometimes, this purple extended to young organs where bracts are initiated (**Fig. 14J**). As the

stem grew, the pigment receded to the rosette junction in both accessions (**Fig. 5I**). This

transient anthocyanin presence specific to *Tsu-0* supported our clustering strategy for capturing

genes involved in transient bract development during flowering transition. Interestingly, among

the 124 expression-based putative bract regulators (**Fig. 5H**), 12 also lied in the mapped QTLs

(**Fig. 4I**). They all differed from the 33 genes selected before via specific overlap with *jagged-5d*

(**Table S2**). Interestingly, we found one anthocyanin biosynthetic enzyme (the dihydroflavonol

reductase, DFR) and five SA-responding genes (see **Table S3**). For example, **Figure 5J** shows the

expression profiles of two such genes. Further work is required to test whether these

candidates contribute to basal bract formation in *Tsu-0* and if the anthocyanin and/or SA

pathways are involved in this natural variation. However, our transcriptomics exclude most of

the genes previously associated with bract development in mutants, and suggest instead new

candidate pathways promoting bract outgrowth with a wild-type flower during floral transition.

606   ***Bract development occurs in a time window when many genes are desynchronized***

Our clustering approach (**Fig. 5G**) tended to select genes whose transcriptional dynamics were

delayed in *Tsu-0* compared to *Col-0* (**Figure 5J**), with changes of RNA levels occurring later in

Tsu-0. This defines a transcriptional heterochrony. Since bracts have been sometimes

considered as a heterochrony because the juvenile leaf trait is maintained at older stages, we

decided to characterise the extent of changes in the timing of gene expression between the two

accessions. We first plotted the two main axes of variance from a PCA after grouping samples

by genotype and time points (**Fig. 6A**). While the second axis of variance (~24.4%) differentiates

genotypes, the main axis (47.9% of variance) orders the sampling time points chronologically in

both genotypes, representing a transcriptomic age. Surprisingly, *Tsu-0* stage T does not align

with *Col-0* stage T on this axis, clustering instead with the stages F: this transcriptome is closer

to a stage F, explaining the limited expression changes when progressing to the next stage F

(**Fig. 5C**).

As suggested by (Calderwood et al. 2021), the transcriptomes of two genotypes during

flowering cannot be aligned to a single developmental time; each gene may desynchronize

differently, sometimes in opposite directions. To quantify gene desynchronization between *Col-*

0 and *Tsu-0*, we leveraged our previous approach of curve registration (Calderwood et al. 2021)

to predict subtle temporal shifts of gene expression in our dataset (Kristianingsih 2024). This

shift is relative to the floral transition, used as the common reference clock between the two

accessions. Thus, genes with no shift, like *AP1* (**Fig. 5B and Fig. 6B**), may still be shifted in

absolute time, because *Tsu-0* plants flower later with an older absolute age. Positive and

negative shifts reflect desynchronization relative to the event of floral transition: gene

expression dynamics may shift earlier or later. Unlike our previous clustering approach, this

registration method uses a common scaled expression level and computes only temporal shifts,

regardless of absolute expression levels (**Fig. 6B** and **Fig. S15A**). Few genes could not be

registered (**Fig. 6B,** N = 43), suggesting that most genes follow similar temporal dynamics in

both accessions (see examples **Fig. S12C, D** or **Fig. 5B, E**). Most genes were classified into three

categories: null, negative and positive shifts (**Fig. 6B)**. A null shift indicates that the gene

expression in *Tsu-0* stays "in phase" with flowering, like *AP1* (compared with **Fig. 5B)**. A positive

Figure 6: **Massive and complex transcriptional desynchronisation coexist in *Tsu-0* across the floral transition.**

A, PCA of RNAseq data from microdissected meristems of *Col-0* and *Tsu-0* over the four sampled stages (biological replicates averaged per time points). The two main axes of the PCA can be interpreted as developmental time and genotype, respectively.

B, Examples of temporal registration of gene dynamics (right panels) between *Col-0* (green) and *Tsu-0* (red) from scaled expression levels (left panel). Dots represent the expression levels of independent biological replicates, while lines indicate the mean expression level at each time point. In the right panel, the green (or red) dotted curves represent the fitted models for *Col-0* (or *Tsu-0*) independently, while the grey dotted curve represents the joint model for both *Col-0* and *Tsu-0*. If the green and red dotted curves are used, it means the two time series are best explained by two independent models, indicating they are not similar. Conversely, if the grey dotted curve is used, it suggests that a single model best explains both time series, indicating they are similar. The name of the gene plotted is indicated on the left of each row. These four genes exemplify the four possible categories (from top to bottom, respectively): genes that cannot be registered (N = 43, *e.g. CYP705A9*), genes with identical temporal dynamics (shift = 0, *e.g. AP1,* see also Fig. 5B) and genes whose expression dynamics in *Tsu-0* must be shifted negatively (*e.g. DFR,* see also Fig. 5J) or positively (*e.g. AG*) to align with *Col-0*. The last column provides a biological interpretation of the computed shift (see main text): a null shift indicates that the expression dynamics in *Tsu-0* stay "*in phase*" with floral transition while negative or positive shifts indicate that the expression dynamics in *Tsu-0* are desynchronized later or earlier, respectively, than the phenotypic progression of floral transition.

C, Distribution of heterochronic shifts between *Tsu-0* and *Col-0* on the entire meristematic transcriptome, computed by the registration method from B. The shift value is colour-coded in a red-to-blue gradient from -1 to 1.

D, GO term enrichment analysis associated with the three categories of heterochronic shifts. The list of all significant 'biological process' GO terms (BH-adjusted p.value < 0.05) was simplified using semantic similarity (cutoff = 0.7) and the Rich Factor was computed for the remaining terms, revealing the proportion of genes involved among all the genes associated with this GO term. Dot size indicates the count of genes and the color scale is the statistical significance (BH-adjusted p.value) of the enrichment in the shift category. Stars indicate GO terms referring to developmental processes.

or negative shift indicates that gene dynamics in *Tsu-0* occur earlier *(e.g. AG)* or later *(e.g., DFR,*

see also **Fig. 5J**), relative to the floral transition, revealing heterochronies between

transcriptomic and phenotypic processes. Transcriptome-wide, the shifts were broadly

distributed, showing massive and complex desynchronisation of gene dynamics (**Fig. 6C**). The

floral transition in Tsu-0 did not impose its clock to the entire meristematic transcriptome: only

~18% of genes (n=3879) stayed in phase with this process while the majority showed delayed or

advanced expression dynamics. Even more genes shifted earlier relative to the floral transition,

supporting the PCA interpretation (**Fig. 6A**) and confirming that the bract, if considered as

"juvenile", did not represent the majority of heterochronies at the transcriptomic level.

As expected, genes in phase were primarily linked to flowering and developmental phase

change as shown by key regulators of these pathways (**Fig. S15B, C**), which also emerges from

GO term enrichment analysis (**Fig. 6D**). In contrast, known bract genes exhibit a wide range of

shifts, from very early to very late, suggesting that they are not collectively involved in *Tsu-0*

bract development (**Fig. S15A, C**). GO term analysis for each shift category highlighted the

particular processes desynchronized from flowering in *Tsu-0* compared to *Col-0* (**Fig. 6D**). For

instance, vascular differentiation (two GO terms mentioning tracheary element and secondary

cell wall) emerges as a process shifted earlier in *Tsu-0* relative to the floral transition (**Fig. 6D**).

Its dynamics may be driven by other progression factors like absolute age, suggesting its loose

connection with the flowering pathway. Conversely, processes shifted later in *Tsu-0* revealed

terms related to cell division (spindle, cell cycle, mitosis) and ribosomal biogenesis (six GO

terms mentioning ribosome, rRNA, ribonucleoprotein and protein-RNA complexes; **Fig. 6D**),

which suggests that core meristematic functions are prolonged in *Tsu-0*. Further work is yet

needed to explore whether the desynchronization of these processes relative to floral transition

impacts bract development.

Finally, this focus on gene desynchronisation provided an explanation to the fact that gene

expression diverged most at the stage T (**Fig. 5D**). The genes controlling floral transition change

swiftly their expression levels upon stage T when many other genes have different levels

between both accessions, because they are no longer synchronised with flowering in *Tsu-0* (**Fig.**

S16). A transient process such as flowering is likely to occur in varying gene expression states,

especially if it is fast and depends on a small number of genes. Massive desynchronization of

gene dynamics creates transcriptional variation, potentially leading to developmental variation

at the time of flowering, such as bract development. In conclusion, our work provides the first

analysis of transcriptome-wide meristematic heterochronies between two *A. thaliana*

accessions, revealing that natural bract development during floral transition is more complex

than a prolonged vegetative phase.

671   **Discussion**

672   This study documents a natural variation for the presence of bracts at the base of

673   flowering branches in *A. thaliana*, which was used to investigate their genetic and

674   developmental basis. Comprehensive phenotypic characterizations indicate strong differences

675   with known mutant bracts and similarities with bracts of bracteate species. Combining

676   quantitative genetics, genomics and meristem transcriptomics on the accession *Tsu-0*, our

677   results suggest new mechanisms controlling bract outgrowth and highlight the phenomenon of

678   massive gene desynchronizations at floral transition, raising new questions about their

679   implications in development and evolution.

680   ***A new mechanism unlocking bract development without affecting floral identity***

681   ***during floral transition***

682   We used two quantitative genetics approaches (BSA and RIL) and a transcriptomic approach to

683   identify the genetic mechanisms controlling basal bract formation in *Tsu-0*. We identified at

684   least four major QTLs, the two main on chromosome 1 with additive effects (**Fig. 4** and **Fig. S7**).

685   However, the density of polymorphisms between the two accessions and the high number of

686   differentially expressed genes resulted in many potential candidates within the mapped

687   intervals. Further fine-mapping or a GWAS analysis on a larger accession panel could provide a

688   higher resolution and even reveal more causative variations.

689   Our work also proposes new pathways for bract development: genetic interactions between

690   *SOC1*, *TFL1* and *PUCHI* (**Fig. 5E**), chloroplast or metal ion transport and homeostasis (**Fig. S13D**),

691   anthocyanin biosynthesis and response to salicylic acid (**Fig. 5I, J**; **Fig. S14E-J**), ribosome

692   biogenesis (**Fig. 6D**). The low number of transcription factors (**tables S1, S2, S3**) compared to

693   enzymes and genes related to metabolic pathways (**Fig. 5 and 6**; **Fig. S13 and S14**) aligns with

694   studies reporting that genes with basic metabolic and cellular functions control specific

695   developmental processes in plants (Tsukaya et al. 2013). Likewise, the *PLA1/2/3* genes In

696   *Poaceae* are three unrelated metabolic enzymes (a cytochrome P450, MEI2-like RNA-binding

697   protein and a glutamate carboxypeptidase II, respectively) that partially suppress bract

698   outgrowth in a redundant manner through a still unknown mechanism (Kawakatsu et al. 2009;

Wang et al. 2021). Our data may thus be helpful to investigate the mechanisms that control

bract development.

Despite the remaining uncertainty about the causal genes and pathways, our data show the

existence of a new bract developmental process different from regained bracts produced in

mutants. *Tsu-0* basal bracts display specific features: association with wild-type flowers (**Fig. 2**

and **Fig. S2, S3**); wild-type bract shape and position (**Fig. 1**, **Fig. S11**); presence restricted at the

base of the raceme (**Fig. S4, S5**); no modification of plastochron rate (**Fig. 3B, Fig. S6A**);

independence from light regime (**Fig. 3D**). Not only such phenotypes differ from those reported

in bract mutants, but none of the known "bract mutant genes" were found in QTL intervals,

indicating that the causal polymorphisms involve other genes and that bract development could

be de-repressed without affecting floral meristem identity. However, because of the

transgressive indeterminism observed in some RILs (**Fig. S11**), it cannot be ruled out that some

of the bract causal genes could also affect flower development, but genetic interactions would

suppress these floral phenotypes while maintaining the basal bracts.

Transcriptomic heterochronies at floral transition may challenge developmental

canalization

The transient formation of bracts at the base of each raceme raises many biological questions.

This trait is widespread within the *Brassicaceae* phylogeny (**Fig. 1S**). It is also known for decades

that several species within their natural context display developmental variations at the base of

the raceme, including bracts, flower-to-shoot transformations, or flower dimorphisms (Arber

1931a; Arber 1931b). Since these basal nodes are produced just at the floral transition, this

suggests that developmental canalization (ensuring an invariant phenotypic output) is less

effective at this stage. The variability of bract formation from branch to branch and plant to

plant (**Fig. 1 D-F, Fig. 4A**) also suggests limited developmental canalization. Our data show that

natural genetic variation is sufficient to reveal a higher frequency of phenotypic variations at

the base of branches. To our knowledge, no specific hypothesis has been proposed to explain

the lower developmental canalization at the floral transition.

We propose that transcriptional heterochronies may account for some of this phenomenon. In

bract-less species, bract development is often described as a prolonged vegetative state,

characterised by the persistence of a juvenile trait (a leaf) with an adult trait (a flower)(Alberch

et al. 1979; Buendía-Monreal and Gillmor 2018). Using microdissected meristems at different

stages (**Fig. 5**), we expected to find this signature in the bract-making *Tsu-0* transcriptome.

Instead, we discovered that the floral transition occurs in an older transcriptome rather than a

younger one (**Fig. 6A, C**). In absolute time, *Tsu-0* plants flower later (**Fig. 3 E, F; Fig. S12 A**), so

the flowering-related genes are actually shifted later. However, a large majority of genes get

out of synchronisation with flowering. In absolute time, some may keep their dynamics or even

shift earlier, both categorised as "earlier" in our analysis using flowering as the reference, while

other genes can shift even later than flowering (**Fig. 6 B, C**). Confirming our previous results

using *Brassica rapa* cultivars and *A. thaliana* (Calderwood et al. 2021), such massive and

complex gene desynchronisation during floral transition seems to be a general rule at both

intra- and inter-species level. Flowering time is under strong selection pressure in *A. thaliana*

(Bloomer and Dean 2017). If selective adaptation of flowering time constantly shifts and

desynchronizes the flowering-related genes from numerous others, this can create new global

gene expression states, especially at the transition when gene expressions vary quickly (**Fig.**

S16). This could promote transient developmental variations, like basal bracts.

The consequence of gene desynchronization is to create a peak of transcriptome divergence

upon floral transition. Such a peak has been reported in *Solanaceae* where it was associated

with the phenotypic evolution of inflorescence complexity (Lemmon et al. 2016). Mirroring the

"inverse hourglass" model for animal embryogenesis, morphological variations would be

promoted by transcriptomic divergence during intermediate developmental steps. Hence, the

sensitivity of floral transition to heterochronies at the transcriptional level could have larger

implications for phenotypic evolution from populations to species.

Evolution of bract loss in Brassicaceae and Angiosperms and the possible role of

heterochronies

Heterochrony is often considered as a powerful mechanism for evolutionary change (Buendía-

Monreal and Gillmor 2018; Petrone-Mendoza et al. 2023). Here, we propose that

heterochronies at the floral transition may lower developmental canalization, contributing to

bract derepression in some accessions. Conversely, heterochronies might have been involved in

bract loss in the *Brassicaceae* ancestor. Current evolutionary scenarios for bract loss are

deduced from mutants and do not involve heterochrony. In *A. thaliana*, flowers would have

evolved the ability to repress bract development (Whipple et al. 2010), while In *Poaceae*,

boundary regions may have acquired this function through genes like NTT (Whipple et al. 2010;

Xiao et al. 2022). However, direct regulation of NTT by SPL genes in maize (Xiao et al. 2022) and

rice (Wang et al. 2021) provide a link with a major regulator of heterochrony in plants, the

SPL/miR156 balance (Buendía-Monreal and Gillmor 2018). SPL genes have not been found to

control bract development in *A. thaliana*. Recently, *SOC1*, *FUL* and *AGL24* were shown to be

more effective than *LFY* or *BOP1/2* in repressing bracts (Manuela and Xu 2024 Mar 28). These

genes are special because they act both as FMI genes and as flowering time genes. Our data did

not correlate bract outgrowth with any FMI genes (**Fig. 2, Fig. 5**) while flowering time was

clearly impacted at the phenotypic (**Fig. 3E, F**) and transcriptional level (*e.g.*, *FLC* in **Fig. S12C or**

Fig. S15C). Since flowering time is prone to heterochrony, this could connect bract with

heterochrony in *A. thaliana*. Hence, despite differences in molecular players, bract outgrowth in

Brassicaceae and *Poaceae* may share more similarities than previously described.

Deducing evolutionary events from mutants relies on the assumption that natural evolution

proceeds in the opposite direction to artificial mutants, which should be taken with caution.

Indeed, actual ebracteate plants have accumulated many evolutionary changes since they

diverged from the last bracteate ancestor: genetic interactions revealed by mutants in current

species may be totally irrelevant in the ancestor. It has also been proposed as the "Dollo's law"

(Gould 1970) that lost traits cannot be regained and "*must be constructed afresh in some*

different mode" (Arber 1918). However, heterochronies have been proposed as a mechanism to

break Dollo's law (Cronk 2009). Since ebracteate plants still produce leaves elsewhere,

activating or repressing this functional developmental program by heterochronies could

effectively coordinate bract and flower development, explaining both bract loss and their

"flickering presence" (Marshall et al. 1994) within and between species in the entire

Brassicaceae family. Further work, notably including different bracteate and ebracteate species,

will be necessary to test these hypotheses and clarify the genetic and developmental

mechanisms that led to bract loss in *Brassicaceae*.

Acknowledgements

We thank Teva Vernoux for the critical reading of the manuscript. We thank Mathieu

Hanemian, Gaël Yvert and Bjorn Pieper for their advice in quantitative genetics. We thank

Thomas Widiez and Nathanaël Jacquier for their advice in Kasp genotyping. We thank Michiel

Vandenbussche for providing the picture of a Petunia plant (Fig. 1A). We thank the following

researchers for providing *A. thaliana* mutant seed stocks: Mitsuhiro Aida for *puchi-1*, *bop1-4* x

bop2-11, and *puchi-1 x bop1-4 x bop2-11* ; Jiawei Wang for *soc1-2 x agl24-1 x svp-41* ; Hicham

Chathane for *lfy-12* ; Georges Coupland and Enric Bertran Garcia de Olalla for *soc1-2*. We thank

Pr Eric Schranz and Frank Becker, from Wageningen University (WUR, Holland) for providing

seeds of *Tarenaya hassleriana* and *Gynandropis gynandra*. We acknowledge the PSMN (Pôle

Scientifique de Modélisation Numérique) computing centre of 'ENS de Lyon' for providing

support in the genomic variant analysis. We thank the Embassy of France in the United

Kingdom for a short-term travel fellowship awarded to R. K. R.M. and R.W. gratefully

acknowledge support from the Biotechnology and Biological Science Research Council Institute

Strategic Programme 'Building Resilience in Crops' (BB/X01102X/1).

Competing interests,

The authors declare that they have no competing interests

Author contributions

Sana Dieudonné: Conceptualization, Investigation, Software (implementation), Formal analysis,

Visualization, Methodology, Writing -review and editing; Ruth Kristianingsih: Formal Analysis,

Methodology, Software (designing, programming, testing, implementing), Visualization, Writing

-review and editing; Stephanie Laine, Investigation; Béline Jesson: Formal analysis, Véronique

Vidal: Supervision, Rachel Wells: Supervision, Writing -review and editing; Richard Morris:

Supervision, Writing -review and editing; Fabrice Besnard: Conceptualization, Project

administration, Supervision, Investigation, Formal analysis, Methodology, Software

(programming, implementing), Funding acquisition, Visualization, Writing - original draft, review

and editing.

Data availability

Whole-genome DNA-seq data of the parental lines (*Col-0* and *Tsu-0*) and of the two F2 pools

used for bulk segregant analysis are deposited under this identifier: doi:10.57745/Z80SIM.

Time-course RNA-seq data of *Col-0* and *Tsu-0* micro-dissected meristems during flowering are

available with these doi: doi:10.57745/DKMQ06, doi:10.57745/7JI3E7, respectively. RNA-seq

data of micro-dissected shoot meristems of the mutant lines (*puchi-1x bop1-4* x *bop2-11*, *lfy-12*

and *jagged-5d)* at early flowering stage are available at doi:10.57745/HAGJJH.

References

Alberch P, Gould SJ, Oster GF, Wake DB. 1979. Size and Shape in Ontogeny and Phylogeny.

Paleobiology. 5(3):296–317.

Al-Shehbaz IA. 2015. Cardamine xinfenii (Brassicaceae), a New Species from Sichuan (China).

Harvard Papers in Botany. 20(2):145–146.

Anders S, Huber W. 2010. Differential expression analysis for sequence count data. Genome

Biol. 11(10):R106. doi:10.1186/gb-2010-11-10-r106.

Arber A. 1918. The "Law of Loss" in Evolution. Proc Linn Soc Lond. 131:70–78.

Arber A. 1931a. Studies in Floral Morphology. I. On Some Structural Features of the Cruciferous

Flower. The New Phytologist. 30(1):11–41.

Arber A. 1931b. Studies in Floral Morphology. II. Some Normal and Abnormal Crucifers: with a

Discussion on teratology and Atavism. The New Phytologist. 30(3):172–203.

Balanzà V, Martínez-Fernández I, Ferrándiz C. 2014. Sequential action of FRUITFULL as a

modulator of the activity of the floral regulators SVP and SOC1. J Exp Bot. 65(4):1193–

1203. doi:10.1093/jxb/ert482.

van Berloo R. 2008. GGT 2.0: Versatile Software for Visualization and Analysis of Genetic Data.

Journal of Heredity. 99(2):232–236. doi:10.1093/jhered/esm109.

Besnard F, Picao-Osorio J, Dubois C, Félix M-A. 2020. A broad mutational target explains a fast

rate of phenotypic evolution. Weigel D, Landry CR, Matus D, editors. eLife. 9:e54928.

doi:10.7554/eLife.54928.

Bloomer RH, Dean C. 2017. Fine-tuning timing: natural variation informs the mechanistic basis

of the switch to flowering in Arabidopsis thaliana. Journal of Experimental Botany.

68(20):5439–5452. doi:10.1093/jxb/erx270.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence

data. Bioinformatics. 30(15):2114–2120. doi:10.1093/bioinformatics/btu170.

Bowman JL, Alvarez J, Weigel D, Meyerowitz EM, Smyth DR. 1993. Control of flower

development in Arabidopsis thaliana by APETALA1 and interacting genes. Development.

119(3):721–743. doi:10.1242/dev.119.3.721.

Broman KW, Gatti DM, Simecek P, Furlotte NA, Prins P, Sen Ś, Yandell BS, Churchill GA. 2019.

R/qtl2: Software for Mapping Quantitative Trait Loci with High-Dimensional Data and

Multiparent Populations. Genetics. 211(2):495–502. doi:10.1534/genetics.118.301595.

Buendía-Monreal M, Gillmor CS. 2018. The times they are a-changin': heterochrony in plant

development and evolution. Frontiers in Plant Science. 9. doi:10.3389/fpls.2018.01349.

Calderwood A, Hepworth J, Woodhouse S, Bilham L, Jones DM, Tudor E, Ali M, Dean C, Wells R,

Irwin JA, et al. 2021. Comparative transcriptomics reveals desynchronisation of gene

expression during the floral transition between Arabidopsis and Brassica rapa cultivars.

Quant Plant Biol. 2:e4. doi:10.1017/qpb.2021.6.

Carlson M. 2024. org.At.tair.db: Genome wide annotation for Arabidopsis.

Chahtane H, Zhang B, Norberg M, LeMasson M, Thévenon E, Bakó L, Benlloch R, Holmlund M,

Parcy F, Nilsson O, et al. 2018. LEAFY activity is post-transcriptionally regulated by BLADE

ON PETIOLE2 and CULLIN3 in Arabidopsis. New Phytol. 220(2):579–592.

doi:10.1111/nph.15329.

Chen Y, Lun ATL, Smyth GK. 2016. From reads to genes to pathways: differential expression

analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-likelihood

pipeline. doi:10.12688/f1000research.8987.2. [accessed 2024 Jul 16].

https://f1000research.com/articles/5-1438.

Cheng C-Y, Krishnakumar V, Chan AP, Thibaud-Nissen F, Schobel S, Town CD. 2017. Araport11: a

complete reannotation of the Arabidopsis thaliana reference genome. Plant J.

89(4):789–804. doi:10.1111/tpj.13415.

Chuck G, Cigan AM, Saeteurn K, Hake S. 2007. The heterochronic maize mutant Corngrass1

results from overexpression of a tandem microRNA. Nat Genet. 39(4):544–549.

doi:10.1038/ng2001.

Chuck G, Whipple C, Jackson D, Hake S. 2010. The maize SBP-box transcription factor encoded

by tasselsheath4 regulates bract development and the establishment of meristem

boundaries. Development. 137(8):1243–1250. doi:10.1242/dev.048348.

Chuck GS, Brown PJ, Meeley R, Hake S. 2014. Maize SBP-box transcription factors unbranched2

and unbranched3 affect yield traits by regulating the rate of lateral primordia initiation.

Proc Natl Acad Sci U S A. 111(52):18775–18780. doi:10.1073/pnas.1407401112.

Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM. 2012. A

program for annotating and predicting the effects of single nucleotide polymorphisms,

SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly

(Austin). 6(2):80–92. doi:10.4161/fly.19695.

Clough SJ, Bent AF. 1998. Floral dip: a simplified method for Agrobacterium-mediated

transformation of Arabidopsis thaliana. Plant J. 16(6):735–743.

Cronk QCB. 2009. Evolution in reverse gear: the molecular basis of loss and reversal. Cold Spring

Harb Symp Quant Biol. 74:259–266. doi:10.1101/sqb.2009.74.034.

Cutler SR, Ehrhardt DW, Griffitts JS, Somerville CR. 2000. Random GFP::cDNA fusions enable

visualization of subcellular structures in cells of Arabidopsis at a high frequency. Proc

Natl Acad Sci U S A. 97(7):3718–3723. doi:10.1073/pnas.97.7.3718.

Dinneny JR, Yadegari R, Fischer RL, Yanofsky MF, Weigel D. 2004. The role of JAGGED in shaping

lateral organs. Development. 131(5):1101–1110. doi:10.1242/dev.00949.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR.

2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 29(1):15–21.

doi:10.1093/bioinformatics/bts635.

Endress PK. 2006. Angiosperm Floral Evolution: Morphological Developmental Framework. In:

Advances in Botanical Research. Vol. 44. Academic Press. (Developmental Genetics of

the Flower). p. 1–61. [accessed 2023 Dec 1].

https://www.sciencedirect.com/science/article/pii/S0065229606440015.

Garnier, Simon, Ross, Noam, Rudis, Robert, Camargo, Pedro A, Sciaini, Marco, et al. 2024.

viridis(Lite) - Colorblind-Friendly Color Maps for R. https://sjmgarnier.github.io/viridis/.

German DA, Hendriks KP, Koch MA, Lens F, Lysak MA, Bailey CD, Mummenhoff K, Al-Shehbaz IA.
2023. An updated classification of the Brassicaceae (Cruciferae). PhytoKeys. 220:127–
144. doi:10.3897/phytokeys.220.97724.

Goldshmidt A, Alvarez JP, Bowman JL, Eshed Y. 2008. Signals derived from YABBY gene activities
in organ primordia regulate growth and partitioning of Arabidopsis shoot apical
meristems. Plant Cell. 20(5):1217–1230. doi:10.1105/tpc.107.057877.

Gould SJ. 1970. Dollo on Dollo's Law: Irreversibility and the Status of Evolutionary Laws. Journal
of the History of Biology. 3(2):189–212.

Grandi V, Gregis V, Kater MM. 2012. Uncovering genetic and molecular interactions among
floral meristem identity genes in Arabidopsis thaliana. Plant J. 69(5):881–893.
doi:10.1111/j.1365-313X.2011.04840.x.

Healey A, Furtado A, Cooper T, Henry RJ. 2014. Protocol: a simple method for extracting next-
generation sequencing quality genomic DNA from recalcitrant plant species. Plant
Methods. 10:21. doi:10.1186/1746-4811-10-21.

Heisler MG, Ohno C, Das P, Sieber P, Reddy GV, Long JA, Meyerowitz EM. 2005. Patterns of
auxin transport and gene expression during primordium development revealed by live
imaging of the Arabidopsis inflorescence meristem. Curr Biol. 15(21):1899–1911.
doi:10.1016/j.cub.2005.09.052.

Helsen J, Voordeckers K, Vanderwaeren L, Santermans T, Tsontaki M, Verstrepen KJ, Jelier R.
2020. Gene Loss Predictably Drives Evolutionary Adaptation. Mol Biol Evol. 37(10):2989–
3002. doi:10.1093/molbev/msaa172.

Hempel FD, Feldman LJ. 1995. Specification of chimeric flowering shoots in wild-type
Arabidopsis. Plant J. 8(5):725–731. doi:10.1046/j.1365-313x.1995.08050725.x.

Hempel FD, Zambryski PC, Feldman LJ. 1998. Photoinduction of flower identity in vegetatively
biased primordia. Plant Cell. 10(10):1663–1676. doi:10.1105/tpc.10.10.1663.

Hepworth SR, Klenz JE, Haughn GW. 2006. UFO in the Arabidopsis inflorescence apex is
required for floral-meristem identity and bract suppression. Planta. 223(4):769–778.
doi:10.1007/s00425-005-0138-3.

Hepworth SR, Zhang Y, McKim S, Li X, Haughn GW. 2005. BLADE-ON-PETIOLE-dependent

signaling controls leaf and floral patterning in Arabidopsis. Plant Cell. 17(5):1434–1448.

doi:10.1105/tpc.104.030536.

Houston K, Druka A, Bonar N, Macaulay M, Lundqvist U, Franckowiak J, Morgante M, Stein N,

Waugh R. 2012. Analysis of the barley bract suppression gene Trd1. Theor Appl Genet.

125(1):33–45. doi:10.1007/s00122-012-1814-x.

Huber W, Carey VJ, Gentleman R, Anders S, Carlson M, Carvalho BS, Bravo HC, Davis S, Gatto L,

Girke T, et al. 2015. Orchestrating high-throughput genomic analysis with Bioconductor.

Nat Methods. 12(2):115–121. doi:10.1038/nmeth.3252.

Irish VF, Sussex IM. 1990. Function of the apetala-1 gene during Arabidopsis floral

development. Plant Cell. 2(8):741–753. doi:10.1105/tpc.2.8.741.

Itoh JI, Hasegawa A, Kitano H, Nagato Y. 1998. A recessive heterochronic mutation,

plastochron1, shortens the plastochron and elongates the vegetative phase in rice. Plant

Cell. 10(9):1511–1522. doi:10.1105/tpc.10.9.1511.

Karim MR, Hirota A, Kwiatkowska D, Tasaka M, Aida M. 2009. A role for Arabidopsis PUCHI in

floral meristem identity and bract suppression. Plant Cell. 21(5):1360–1372.

doi:10.1105/tpc.109.067025.

Kawakatsu T, Itoh J-I, Miyoshi K, Kurata N, Alvarez N, Veit B, Nagato Y. 2006. PLASTOCHRON2

regulates leaf initiation and maturation in rice. Plant Cell. 18(3):612–625.

doi:10.1105/tpc.105.037622.

Kawakatsu T, Taramino G, Itoh J-I, Allen J, Sato Y, Hong S-K, Yule R, Nagasawa N, Kojima M,

Kusaba M, et al. 2009. PLASTOCHRON3/GOLIATH encodes a glutamate

carboxypeptidase required for proper development in rice. Plant J. 58(6):1028–1040.

doi:10.1111/j.1365-313X.2009.03841.x.

Kiefer M, Schmickl R, German DA, Mandáková T, Lysak MA, Al-Shehbaz IA, Franzke A,

Mummenhoff K, Stamatakis A, Koch MA. 2014. BrassiBase: introduction to a novel

knowledge database on Brassicaceae evolution. Plant Cell Physiol. 55(1):e3.

doi:10.1093/pcp/pct158.

Kinoshita A, Vayssières A, Richter R, Sang Q, Roggen A, van Driel AD, Smith RS, Coupland G.

2020. Regulation of shoot meristem shape by photoperiodic signaling and

phytohormones during floral induction of Arabidopsis. Elife. 9:e60661.

doi:10.7554/eLife.60661.

Kristianingsih R. 2024. greatR: Gene Registration from Expression and Time-Courses in R.

https://ruthkr.github.io/greatR/.

Kwiatkowska D. 2006. Flower primordium formation at the Arabidopsis shoot apex:

quantitative analysis of surface geometry and growth. J Exp Bot. 57(3):571–580.

doi:10.1093/jxb/erj042.

Kwiatkowska D. 2008. Flowering and apical meristem growth dynamics. J Exp Bot. 59(2):187–

201. doi:10.1093/jxb/erm290.

Lawrence M, Gentleman R, Carey V. 2009. rtracklayer: an R package for interfacing with

genome browsers. Bioinformatics. 25(14):1841–1842.

doi:10.1093/bioinformatics/btp328.

Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R, Morgan MT, Carey VJ.

2013. Software for computing and annotating genomic ranges. PLoS Comput Biol.

9(8):e1003118. doi:10.1371/journal.pcbi.1003118.

Lemmon ZH, Park SJ, Jiang K, Van Eck J, Schatz MC, Lippman ZB. 2016. The evolution of

inflorescence diversity in the nightshades and heterochrony during meristem

maturation. Genome Res. 26(12):1676–1686. doi:10.1101/gr.207837.116.

Levin JZ, Meyerowitz EM. 1995. UFO: an Arabidopsis gene involved in both floral meristem and

floral organ development. Plant Cell. 7(5):529–548. doi:10.1105/tpc.7.5.529.

Liu C, Xi W, Shen L, Tan C, Yu H. 2009. Regulation of floral patterning by flowering time genes.

Dev Cell. 16(5):711–722. doi:10.1016/j.devcel.2009.03.011.

Long J, Barton MK. 2000. Initiation of axillary and floral meristems in Arabidopsis. Dev Biol.

218(2):341–353. doi:10.1006/dbio.1999.9572.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol. 15(12):550. doi:10.1186/s13059-014-0550-8.

Maizel A, Weigel D. 2004. Temporally and spatially controlled induction of gene expression in

Arabidopsis thaliana. The Plant Journal. 38(1):164–171. doi:10.1111/j.1365-

313X.2004.02027.x.

Mansfeld BN, Grumet R. 2018. QTLseqr: An R Package for Bulk Segregant Analysis with Next-

Generation Sequencing. Plant Genome. 11(2). doi:10.3835/plantgenome2018.01.0006.

Manuela D, Xu M. 2024 Mar 28. Aintegumenta And Redundant Aintegumenta-Like6 Are

Required For Bract Outgrowth In Arabidopsis. J Exp Bot.:erae138.

doi:10.1093/jxb/erae138.

Marshall CR, Raff EC, Raff RA. 1994. Dollo's law and the death and resurrection of genes. Proc

Natl Acad Sci U S A. 91(25):12283–12287. doi:10.1073/pnas.91.25.12283.

McCarthy DJ, Chen Y, Smyth GK. 2012. Differential expression analysis of multifactor RNA-Seq

experiments with respect to biological variation. Nucleic Acids Res. 40(10):4288–4297.

doi:10.1093/nar/gks042.

Melzer S, Lens F, Gennen J, Vanneste S, Rohde A, Beeckman T. 2008. Flowering-time genes

modulate meristem determinacy and growth form in Arabidopsis thaliana. Nat Genet.

40(12):1489–1492. doi:10.1038/ng.253.

Michaels SD, Ditta G, Gustafson-Brown C, Pelaz S, Yanofsky M, Amasino RM. 2003. AGL24 acts

as a promoter of flowering in Arabidopsis and is positively regulated by vernalization.

The Plant Journal. 33(5):867–874. doi:10.1046/j.1365-313X.2003.01671.x.

Miyoshi K, Ahn B-O, Kawakatsu T, Ito Y, Itoh J-I, Nagato Y, Kurata N. 2004. PLASTOCHRON1, a

timekeeper of leaf initiation in rice, encodes cytochrome P450. Proc Natl Acad Sci U S A.

101(3):875–880. doi:10.1073/pnas.2636936100.

Nilsson O, Wu E, Wolfe DS, Weigel D. 1998. Genetic ablation of flowers in transgenic

Arabidopsis. Plant J. 15(6):799–804. doi:10.1046/j.1365-313x.1998.00260.x.

Norberg M, Holmlund M, Nilsson O. 2005. The BLADE ON PETIOLE genes act redundantly to

control the growth and development of lateral organs. Development. 132(9):2203–

2213. doi:10.1242/dev.01815.

Ohno CK, Reddy GV, Heisler MGB, Meyerowitz EM. 2004. The Arabidopsis JAGGED gene

encodes a zinc finger protein that promotes leaf tissue development. Development.

131(5):1111–1122. doi:10.1242/dev.00991.

Pasha A, Subramaniam S, Cleary A, Chen X, Berardini T, Farmer A, Town C, Provart N. 2020.

Araport Lives: An Updated Framework for Arabidopsis Bioinformatics. Plant Cell.

32(9):2683–2686. doi:10.1105/tpc.20.00358.

Penin AA. 2008. Bract reduction in Cruciferae: possible genetic mechanisms and evolution.

Wulfenia. 15:63–73.

Petrone-Mendoza E, Vergara-Silva F, Olson ME. 2023. Plant morpho evo-devo. Trends Plant Sci.

28(11):1257–1276. doi:10.1016/j.tplants.2023.06.007.

Prenner G, Vergara-Silva F, Rudall PJ. 2009. The key role of morphology in modelling

inflorescence architecture. Trends Plant Sci. 14(6):302–309.

doi:10.1016/j.tplants.2009.03.004.

R Core Team. 2018. R: A Language and Environment for Statistical Computing. R Foundation for

Statistical Computing. Vienna, Austria. https://www.R-project.org/.

Ratcliffe OJ, Amaya I, Vincent CA, Rothstein S, Carpenter R, Coen ES, Bradley DJ. 1998. A

common mechanism controls the life cycle and architecture of plants. Development.

125(9):1609–1615. doi:10.1242/dev.125.9.1609.

Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. 2015. limma powers differential

expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Research.

43(7):e47. doi:10.1093/nar/gkv007.

Sadier A, Sears KE, Womack M. 2022. Unraveling the heritage of lost traits. J Exp Zool B Mol Dev

Evol. 338(1–2):107–118. doi:10.1002/jez.b.23030.

Sawa S, Ito T, Shimura Y, Okada K. 1999. FILAMENTOUS FLOWER controls the formation and

development of arabidopsis inflorescences and floral meristems. Plant Cell. 11(1):69–86.

doi:10.1105/tpc.11.1.69.

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis.

Nat Methods. 9(7):671–675. doi:10.1038/nmeth.2089.

Schultz EA, Haughn GW. 1991. LEAFY, a Homeotic Gene That Regulates Inflorescence

Development in Arabidopsis. Plant Cell. 3(8):771–781. doi:10.1105/tpc.3.8.771.

Shannon S, Meeks-Wagner DR. 1991. A Mutation in the Arabidopsis TFL1 Gene Affects

Inflorescence Meristem Development. Plant Cell. 3(9):877–892.

doi:10.1105/tpc.3.9.877.

Siegfried KR, Eshed Y, Baum SF, Otsuga D, Drews GN, Bowman JL. 1999. Members of the YABBY

gene family specify abaxial cell fate in Arabidopsis. Development. 126(18):4117–4128.

doi:10.1242/dev.126.18.4117.

Simon M, Loudet O, Durand S, Bérard A, Brunel D, Sennesal F-X, Durand-Tardif M, Pelletier G,

Camilleri C. 2008. Quantitative trait loci mapping in five new large recombinant inbred

line populations of Arabidopsis thaliana genotyped with consensus single-nucleotide

polymorphism markers. Genetics. 178(4):2253–2264. doi:10.1534/genetics.107.083899.

Siriwardana NS, Lamb RS. 2012. The poetry of reproduction: the role of LEAFY in Arabidopsis

thaliana flower formation. Int J Dev Biol. 56(4):207–221. doi:10.1387/ijdb.113450ns.

Smyth DR, Bowman JL, Meyerowitz EM. 1990. Early flower development in Arabidopsis. Plant

Cell. 2(8):755–767. doi:10.1105/tpc.2.8.755.

Takagi H, Abe A, Yoshida K, Kosugi S, Natsume S, Mitsuoka C, Uemura A, Utsushi H, Tamiru M,

Takuno S, et al. 2013. QTL-seq: rapid mapping of quantitative trait loci in rice by whole

genome resequencing of DNA from two bulked populations. Plant J. 74(1):174–183.

doi:10.1111/tpj.12105.

Tian T, Liu Y, Yan H, You Q, Yi X, Du Z, Xu W, Su Z. 2017. agriGO v2.0: a GO analysis toolkit for

the agricultural community, 2017 update. Nucleic Acids Res. 45(W1):W122–W129.

doi:10.1093/nar/gkx382.

Tsukaya H, Byrne ME, Horiguchi G, Sugiyama M, Van Lijsebettens M, Lenhard M. 2013. How do

"housekeeping" genes control organogenesis?--Unexpected new findings on the role of

housekeeping genes in cell and organ differentiation. J Plant Res. 126(1):3–15.

doi:10.1007/s10265-012-0518-2.

Wang Hai, Wang Haiyang. 2015. The miR156/SPL Module, a Regulatory Hub and Versatile

Toolbox, Gears up Crops for Enhanced Agronomic Traits. Mol Plant. 8(5):677–688.

doi:10.1016/j.molp.2015.01.008.

Wang L, Ming L, Liao K, Xia C, Sun S, Chang Y, Wang H, Fu D, Xu C, Wang Z, et al. 2021. Bract

suppression regulated by the miR156/529-SPLs-NL1-PLA1 module is required for the

transition from vegetative to reproductive branching in rice. Mol Plant. 14(7):1168–

1184. doi:10.1016/j.molp.2021.04.013.

Wang L, Yin H, Qian Q, Yang J, Huang C, Hu X, Luo D. 2009. NECK LEAF 1, a GATA type

transcription factor, modulates organogenesis by regulating the expression of multiple

regulatory genes during reproductive development in rice. Cell Res. 19(5):598–611.

doi:10.1038/cr.2009.36.

Weigel D. 1995. The genetics of flower development: from floral induction to ovule

morphogenesis. Annu Rev Genet. 29:19–39. doi:10.1146/annurev.ge.29.120195.000315.

Weigel D, Alvarez J, Smyth DR, Yanofsky MF, Meyerowitz EM. 1992. LEAFY controls floral

meristem identity in Arabidopsis. Cell. 69(5):843–859. doi:10.1016/0092-

8674(92)90295-n.

Whipple CJ. 2017. Grass inflorescence architecture and evolution: the origin of novel signaling

centers. New Phytol. 216(2):367–372. doi:10.1111/nph.14538.

Whipple CJ, Hall DH, DeBlasio S, Taguchi-Shiobara F, Schmidt RJ, Jackson DP. 2010. A conserved

mechanism of bract suppression in the grass family. Plant Cell. 22(3):565–578.

doi:10.1105/tpc.109.073536.

Wickham H. 2007. Reshaping Data with the reshape Package. Journal of Statistical Software.

21(12):1–20.

Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

https://ggplot2.tidyverse.org.

Wickham H, François R, Henry L, Müller K, Vaughan D. 2023. dplyr: A Grammar of Data

Manipulation. https://dplyr.tidyverse.org.

Wickham H, Vaughan D, Girlich M. 2024. tidyr: Tidy Messy Data. https://tidyr.tidyverse.org.

Wu G, Poethig RS. 2006. Temporal regulation of shoot development in Arabidopsis thaliana by

miR156 and its target SPL3. Development. 133(18):3539–3547. doi:10.1242/dev.02521.

Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, Feng T, Zhou L, Tang W, Zhan L, et al. 2021.

clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. The

Innovation. 2(3):100141. doi:10.1016/j.xinn.2021.100141.

Xiao Y, Guo J, Dong Z, Richardson A, Patterson E, Mangrum S, Bybee S, Bertolini E, Bartlett M,

Chuck G, et al. 2022. Boundary domain genes were recruited to suppress bract growth

and promote branching in maize. Sci Adv. 8(24):eabm6835.

doi:10.1126/sciadv.abm6835.

Xu M, Hu T, McKim SM, Murmu J, Haughn GW, Hepworth SR. 2010. Arabidopsis BLADE-ON-

PETIOLE1 and 2 promote floral meristem fate and determinacy in a previously undefined

pathway targeting APETALA1 and AGAMOUS-LIKE24. Plant J. 63(6):974–989.

doi:10.1111/j.1365-313X.2010.04299.x.

Xu Y-C, Niu X-M, Li X-X, He W, Chen J-F, Zou Y-P, Wu Q, Zhang YE, Busch W, Guo Y-L. 2019.

Adaptation and Phenotypic Diversification in Arabidopsis through Loss-of-Function

Mutations in Protein-Coding Genes[OPEN]. Plant Cell. 31(5):1012–1025.

doi:10.1105/tpc.18.00791.

Zhao Y, Medrano L, Ohashi K, Fletcher JC, Yu H, Sakai H, Meyerowitz EM. 2004. HANABA

TARANU is a GATA transcription factor that regulates shoot apical meristem and flower

development in Arabidopsis. Plant Cell. 16(10):2586–2600. doi:10.1105/tpc.104.024869.

Supporting Information

Figures S1 to S16

Figure S1: Basal bracts are common among *Brassicaceæ* tribes and their presence is not

correlated to broad geographic nor genetic origin in *A. thaliana.*

Figure S2: Comparing basal bract formation in *Tsu-0* with bracts forming in a bract-making sister

species.

Figure S3: Known bract mutants in *A. thaliana* display abnormal floral phenotypes and

indeterminacy.

Figure S4: Bracts of mutants impaired in floral meristem identity genes are not located at the

floral transition like *Tsu-0* natural basal bracts.

Figure S5: Known bract mutants in *A. thaliana* do not display bracts specifically at the floral

transition.

Figure S6: Plastochron variation and flowering heterochronies in *Tsu-0* versus *Col-0* accessions

Figure S7: Bulk segregant analysis of basal bract formation (*Tsu-0* x *Col-0*) identifies four

putative QTLs.

Figure S8: Quantitative genetics of basal bracts formation in *Tsu-0* using a set of RILs with the

reference accession *Col-0*.

Figure S9: Finer mapping of QTLs.

Figure S10: QTL1b* identified for basal bract correlates with a higher cauline branch number.

Figure S11: Transgressive phenotypes in RILs and their genetic determinism provide further

information of the genetic pathways involved.

Figure S12: Transcriptomic profile associated with basal bract formation

Figure S13: Transcriptomic cross-comparisons between bract mutants and *Tsu-0* at T stage

capture an enrichment for processes related to photosynthesis and provide a short list of 33

causal candidate genes.

Figure S14: GO term analysis at T stage and the identification of anthocyanin biosynthesis and

SA-responding pathway enrichment among putative bract regulators.

Figure S15: Temporal Registration of expression dynamics in *Tsu-0* over the floral transition for

genes related to bract development, floral identity and floral transition.

Figure S16: A working model for natural basal bract formation in *Arabidopsis thaliana.*

1146

Tables S1 to S8 are large spreadsheets available for download:

Table S1: List of annotated genes lying in mapped QTLs controlling bracts in *Tsu-0,* with

additional information from RNAseq and genomic variant analysis.

Table S2: Intersection of genes differentially expressed in *Tsu-0* at stage T and specifically in the

jagged-5D mutant meristem and mapped within Tsu-0 bract QTLs.

Table S3: Putative bract regulators in *Tsu-0* identified by clustering apart expression from bract

and non-bract making stages.

Table S4: Details for all genes associated with "response to salicylic acid" about their expression

at stage in *Tsu-0* and their identification as putative bract regulators.

Table S5: Details for all genes associated with anthocyanin metabolism about their expression

at stage in *Tsu-0* and their identification as putative bract regulators.

Table S6: Genotypes of all RIL and HIF lines used in this study.

Table S7: SNP information relative to new KASP genotyping markers used in this study.

Table S8: SNP information relative to new sanger genotyping markers used in this study and

associated primers.

1162

Figure legends

Figure 1: **The presence of basal bracts is a common, natural trait in *Arabidopsis thaliana***

with quantitative variations among genetic backgrounds. **A**, Examples of bracts (red

arrows) in different angiosperms. *Brassicaceae* are mostly bractless but some species retain

bracts at the base of inflorescence branches (e.g. in *L. maritima*). In *A. thaliana*, some natural

accessions display basal bracts (e.g. *Tsu-0*) while others do not (e.g. *Col-0*). Schematic

phylogenetic relationships are indicated with a cladogram below the pictures. White stars: first

ebracteate flowers following previous bracteate flowers **B**, (from left to right) *A. thaliana*'s basal

bracts can be true leaves or just small rudimentary filamentous structures at the base of the

floral pedicel (red arrows). These structures are absent in younger flowers, as in the reference

Col-0 accession (rightmost panel, black arrowhead). **C**, Details of basal bracts by scanning

electron microscopy in a *Tsu-0* inflorescence tip (left panel), showing a bract on the first flower

(red arrow) and swollen base of pedicels on the two following flowers (red arrowhead). Right

panel: pedicels of *Col-0* plants do not show such structures (black arrowhead). **D**, Occurrence of

basal bracts in different accessions, assessed by the percentage of plants with at least one

basal bract in the inflorescence. Each dot is the average value of several plants (number

indicated by dot size) of a scoring assay. A box plot indicates several assays per line and

thicker horizontal black lines are the median value of all scoring assays (range 1-3) in the

accession. *Col-0* and *Tsu* are highlighted in green and red, respectively. The geographical origin

of each strain is located in the world map below. **E**, Definition of a quantitative bract score for a

single plant (see main text for detail). **G**, Quantification of basal bracts in different accessions

using the bract score.

1185

Figure 2: **Flowers bearing bracts express high levels of LFY and do not display mutant**

phenotypes

A, B. Confocal live imaging of the main meristems of *Col-0* (**A**) and *Tsu-0* (**B**) plants at floral

transition, expressing a pLFY transcriptional reporter (magenta) and a membrane marker

(green) for morphology (top-view projections of stack acquisitions). Green arrowheads:

branches, ordered with decreasing numbers from the floral transition; white arrows: flowers

ordered with increasing numbers from the floral transition. In both genotypes, pLFY expression

is absent from branch meristems and suddenly appears in the first flower onwards.

Representative pictures of at least 6 plants per genotype captured at floral transition.

C, side-view of the first flower from image B (*Tsu-0*): a bract (red arrow) is visible on the abaxial

side of this young flower.

D, Number of floral organs in flowers with (dark red) or without (light red) bract in *Tsu-0* plants.

E, F. Side views by confocal live imaging of *Tsu-0* flowers with bracts (red arrow) expressing the

pLFY reporter (magenta) and the green morphological marker. Flowers show two

developmental stages older than C and with increasing age from E to F, as shown by the

growing abaxial sepal on top of the flower. pLFY is expressed at low levels in the bract margins

(magenta arrow).

Scale bars: 50 µm

1204

Figure 3: **Variation in natural basal bract frequency is not correlated with branch position,**

plastochron length or light conditions, but with variation in flowering time

A, Correlation study between the number of lateral (cauline) branches and the number of bracts

on the main stem only in *Col-0* and *Tsu-0* plants (green and red, respectively). (Pearson test, p-

value > 0.1).

B, Effect of different light regimes on bract scores in *Col-0* and *Tsu-0* plants (green and red,

respectively). LD and SD stand for long- and short-day conditions, respectively and CL stands

for continuous light (see methods). Numbers (e.g. 20SD > LD) indicate the number of days in

the first condition before transfer to the second. The number of plants scored is indicated below

each bar.

C, Cumulated production of organs over time for *Col-0* and *Tsu-0* plants counted in

microdissected shoot apices (green and red, respectively, each time point averaging 5 to 10

plants): the time window when plants make the floral transition is indicated by a horizontal arrow

and vertical dashed lines. The rate of organ production (inverse to the plastochron) in this period

is computed from the local slope of the curve. Plants were cultured for 21 days in short-day

conditions before transfer in long-day (LD) conditions (see methods).

D, Result of three independent experiments of plastochron measurements at flowering transition

(see also fig. S5).

E, Differences in flowering time between *Col-0* and *Tsu-0* accession measured as bolting time in

days (left) or as the number of vegetative nodes (rosette leaves and cauline leaves/branches)

before the first flower on the main stem. Plants grown in LD conditions, N = 58 plants per

genotype in two independent replicates.

F, Correlation study between the bract score and the flowering time (assayed when the first

flower opens) in different accessions, labelled with different colours. Each dot is a plant and a

linear regression standard deviation is computed for each accession.

1230

Figure 4: **Identification of 4 major QTLs controlling basal bract formation in *Tsu-0***

suggests unknown genetic pathways. **A**, Genetic transmission of bract score in F1 and F2

hybrids from a cross between *Col-0* and *Tsu-0* parent plants. **B**, QTL mapping for bract score in

a set of RILs identifies four putative QTLs, two in chromosome 1 (1a and 1b) and two in

chromosome 5 (5a and 5b). Dotted horizontal red lines indicate three different significance

thresholds computed from 2000 permutations. On top of the graph, H^2 indicates the (broad-

sense) heritability of the bract score computed among the RIL set. **C**, Genotype of the F6

generation of the line RIL334 in chromosome 1, showing a heterozygotic region overlapping

with QTL1a. **D**, Chromosome 1 genotypes of selected lines forming a heterologous inbred

family (HIF) obtained from RIL334 after two more generations of selfing (F8). **E**, Box plots of

bract scores of the different lines from the "334 family" (blue boxes) with parental controls (*Col-*

0: green, *Tsu-0*: red, N > 20 plants per line). Segregating bract scores within the family map a

narrower region, QTL1a*, spanning 541 annotated genes. **F**, Genotype of the F6 generation of

the line RIL488 in chromosome 1, showing a heterozygotic region overlapping with QTL1b. **G**,

Chromosome 1 genotypes of selected HIF lines obtained from RIL488 at F8 and (**H**) a box plot

of their bract scores (blue boxes), with parental controls (*Col-0*: green, *Tsu-0*: red, N > 21 plants

per line). This maps a narrower region, QTL1b*, spanning 332 annotated genes. In E and H,

lines not sharing the same letter(s) are statistically different (posthoc Tukey analysis with 0.05

sign. level from a glm of bract scores fitted with a quasi-poisson distribution). **I**, Final genetic

mapping of the 'basal bract' trait in *Tsu-0* and the location of genes reported to impact bract

development in previous mutant studies.

1252

Figure 5: **Transcriptomic divergence between *Tsu-0* and *Col-0* meristems peaks at the**

floral transition and suggests an unknown developmental control for natural basal bract

formation

A, Scanning electron microscopy showing the evolution of the main meristem in both *Col-0* and

Tsu-0 at the four different stages (V, L, T, F) used for RNAseq. Plants were first synchronized

by 21 days of a non-inductive short-day light regime before a transfer to inductive long days.

The date after the transfer is indicated in the top-right corner of each picture. Green arrowheads

point to branches (with a leaf) while magenta arrowheads point to the first flowers produced

after floral transition. In *Tsu-0*, red arrows show bracts.

B, Dynamics of the expression levels of FT, LFY and AP1 (three important regulators of floral

transition and identity) in both *Col-0* (green) and *Tsu-0* (red) over the four developmental

stages. Differential expression analysis reveals no difference at any stage between the two

accessions.

C, Bar plots of the number of genes differentially expressed between two consecutive stages in

Col-0 (green) and *Tsu-0* (red). The number of changes peaks at the V-to-T stage transition,

especially in *Tsu-0*.

D, Bar plots of the number of differentially expressed genes (DEG) between *Col-0* and *Tsu-0* at

each stage of the time course. T stage is when the highest number of DEG is measured (on top

of the bar, different letters indicate the statistical difference with a chi-square test of

homogeneity after posthoc analysis).

E, Dynamics of the expression levels of three previously known "bract" genes, showing a

significant difference at least at the T stage between *Col-0* (green) and *Tsu-0* (red). Two stars

mean that the genes are differentially expressed and the fold change is superior to 1.

G, Strategy to cluster genes based on the presence/absence of leaf and/or bract in the different

combinations of stages and genotypes (see text for details).

H, Volcano plot of gene expressions at the T stage between the two accessions. All genes

expressed in the SAM are plotted (n=21,652 grey dots) but only the genes fulfilling the

clustering condition defined in G ("putative bract regulators") are highlighted in orange and blue

for up- and down-regulation, respectively (n=124). Vertical dashed lines: absolute fold change

superior to 1, horizontal dashed line: significance threshold at 5.10^{-2} (adjusted p.value with fdr

method).

I, Representative pictures of micro-dissected meristems in *Col-0* (upper row) and *Tsu-0* (lower

row) just at or before (left) or after (middle) stage T (N>15 for each genotype) and a close-up of

the base of the bolted main stem (right). At stage T, *Tsu-0* meristems display a typical

anthocyanin red coloration just below the meristem, which is not observed in *Col-0*. After

bolting, both genotypes show anthocyanin coloration at the base of the stem. Scale bars: 100

µm (left and middle), 1 cm (right).

J, Examples of the expression profile of two candidate genes, showing an up- (left) and a down-

(right) regulation at the T stage. *DFR* is an enzyme involved in anthocyanin biosynthesis while

FMOGS-OX7 is annotated as a salicylic acid responding enzyme. Two stars mean that the

genes are differentially expressed and the fold change is superior to 1. The bract and bract-less

clusters are outlined with a solid and dashed circle, respectively, while the horizontal dotted line

highlights their separation.

1296

Figure 6: **Massive and complex transcriptional desynchronisation coexist in *Tsu-0* across**

the floral transition.

A, PCA of RNAseq data from microdissected meristems of *Col-0* and *Tsu-0* over the four

sampled stages (biological replicates averaged per time points). The two main axes of the PCA

can be interpreted as developmental time and genotype, respectively.

B, Examples of temporal registration of gene dynamics (right panels) between *Col-0* (green)

and *Tsu-0* (red) from scaled expression levels (left panel). Dots represent the expression levels

of independent biological replicates, while lines indicate the mean expression level at each time

point. In the right panel, the green (or red) dotted curves represent the fitted models for *Col-0* (or

Tsu-0) independently, while the grey dotted curve represents the joint model for both *Col-0* and

Tsu-0. If the green and red dotted curves are used, it means the two time series are best

explained by two independent models, indicating they are not similar. Conversely, if the grey

dotted curve is used, it suggests that a single model best explains both time series, indicating

they are similar. The name of the gene plotted is indicated on the left of each row. These four

genes exemplify the four possible categories (from top to bottom, respectively): genes that

cannot be registered (N = 43, *e.g. CYP705A9*), genes with identical temporal dynamics (shift =

0, *e.g. AP1,* see also Fig. 5B) and genes whose expression dynamics in *Tsu-0* must be shifted

negatively (*e.g. DFR,* see also Fig. 5J) or positively (*e.g. AG*) to align with *Col-0*. The last

column provides a biological interpretation of the computed shift (see main text): a null shift

indicates that the expression dynamics in *Tsu-0* stay "*in phase*" with floral transition while

negative or positive shifts indicate that the expression dynamics in *Tsu-0* are desynchronized

later or earlier, respectively, than the phenotypic progression of floral transition.

C, Distribution of heterochronic shifts between *Tsu-0* and *Col-0* on the entire meristematic

transcriptome, computed by the registration method from B. The shift value is colour-coded in a

red-to-blue gradient from -1 to 1.

D, GO term enrichment analysis associated with the three categories of heterochronic shifts.

The list of all significant 'biological process' GO terms (BH-adjusted p.value < 0.05) was

simplified using semantic similarity (cutoff = 0.7) and the Rich Factor was computed for the

remaining terms, revealing the proportion of genes involved among all the genes associated

with this GO term. Dot size indicates the count of genes and the color scale is the statistical

significance (BH-adjusted p.value) of the enrichment in the shift category. Stars indicate GO

terms referring to developmental processes.