Preclinical testing of RRx-001 in mouse models of experimental

endometriosis reveals promising therapeutic impacts

Iona McIntyre[1, 2], Vadim Vasilyev[1], Chiara Lia Perrone[1], Priya Dhami[1, 2], Kavita Panir[1, 2], Matthew

Rosser[1, 2], Erin Greaves[1, 2]

1. Division of Biomedical Sciences, Warwick Medical School, University of Warwick, Coventry,

United Kingdom (UK)

2. Centre for Early Life, University of Warwick, Coventry, UK

Corresponding author: Dr Erin Greaves

Email: Erin.Greaves@Warwick.ac.uk

Tel: 024 7696 8640

Key words: Endometriosis, therapeutic, macrophage

Abstract

Endometriosis is a chronic inflammatory condition characterised by the presence of ectopic endometrial-like tissue (lesions), associated reduced fertility and chronic pain. Impacting both the health and psycho-social functioning of millions of women worldwide, there is an urgent need for innovative non-hormonal, non-invasive treatments for the disorder. Both peritoneal and lesion-resident macrophages have been strongly implicated in the pathogenesis of endometriosis; key roles include promotion of lesion growth, neuroangiogenesis and nerve sensitization. With such a central role in the disease, macrophages represent a novel therapeutic target. In the current preclinical study, we sought to repurpose the macrophage targeting anti-cancer drug RRx-001 for the treatment of endometriosis. We utilised mouse models of induced endometriosis to demonstrate that RRx-001 acts to reduce endometriosis lesions and attenuate associated pain-like behaviours, without negatively impacting fertility. Using single nuclei multiome analyses, we identified a modification of macrophage subpopulations in the peritoneal cavity, specifically a reduced acquisition of a pro-disease phenotype and an accumulation of a pro-resolving phenotype. These observations signify the potential of RRx-001 as a novel therapeutic for endometriosis management.

Introduction

Endometriosis is a common, chronic, neuro-inflammatory condition that affects 1 in 10 women (and those assigned female sex at birth) during their reproductive years, although adolescent and postmenopausal cases are well documented(*1*). Endometriosis is defined by ectopic endometrial-like tissue which forms non-neoplastic 'lesions', commonly within the peritoneal cavity and rarely (<1%), in extra-pelvic locations(*2, 3*). Endometriosis has a complex and diverse symptomology which includes severe dysmenorrhea, chronic pelvic pain, dyspareunia, fatigue, and infertility. These symptoms have a cumulative impact on health, well-being, and psychosocial functioning. Beyond the personal impact on those that suffer from the condition and that of their partners and family, endometriosis also represents a substantial socioeconomic burden, due not only to the cost of treatment, but also as a result of significant reductions in productivity(*4*).

Despite the high prevalence of endometriosis, there remains limited etiological understanding of the disease(*5*). Sampsons' theory of retrograde menstruation is the most commonly accepted hypothesis for the origin of endometrial-like tissue within the peritoneal cavity, referring to the reflux of sloughed menstrual debris through the fallopian tubes during menstruation(*6*). Although retrograde menstruation is common, occurring in 90% of those with patent fallopian tubes, only a proportion develop endometriosis, indicative of a multifactorial pathogenesis, with immune dysregulation prominent amongst many suspected etiological factors(*7-9*). Following an immunological inability to eliminate refluxed ectopic endometrial fragments, adherence and cell proliferation under hypoxic stress occurs, thus initiating the development of endometriotic lesions(*10, 11*). The current medical management of endometriosis is non-curative and includes analgesic management, hormonal suppression, and surgical debulking of lesions(*12*). This current strategy is insufficient due to common post-surgical recurrence of lesions, unwanted side-effects from hormonal medications (anovulation and those associated with hypoestrogenism), and a high prevalence of endometriosis-associated pain that is refractory to medical treatment(*13-15*). Thus, there is an urgent, unmet need for novel non-hormonal therapeutics for endometriosis.

Macrophages have been strongly implicated in the promotion of endometriosis, emerging as a potential target for a non-hormonal endometriosis therapy(*16*). These dynamic mononuclear phagocytes are present in all tissues, playing integral roles in innate immunity, tissue homeostasis, and the induction and resolution of sterile inflammation. Macrophages demonstrate considerable transcriptional and phenotypical plasticity, existing on a broad spectrum between two highly polarised states: pro-inflammatory and pro-repair, with a varied and dynamic functionality(*17*). Macrophage function is dependent on both the perception of extracellular signals within the tissue niche and

cellular ontogeny: seeded pre-birth from the embryonic yolk sac/fetal liver, or monocyte-derived and

recruited from the circulation(*18, 19*).

In endometriosis, macrophages (and their precursors, monocytes) are an important

component of the intra- and extra-lesion microenvironment. They are continuously recruited and play

pathophysiological roles in lesion establishment, angiogenesis, and nociception(*20, 21*). In both

patients and animal models, pro-endometriosis macrophages have been demonstrated to trend

predominantly towards a pro-repair phenotype(*22, 23*). However, broad heterogeneity in peritoneal

and lesion-resident populations has been revealed in detailed analyses using techniques such as single

cell RNA-sequencing (scRNA-seq) and mass cytometry by time-of-flight (mass CyTOF), demonstrating

subpopulations across the spectrum, playing subtype-specific roles in the progression or regression of

endometriosis(*24-27*). An inability of macrophages to sufficiently recognise and clear refluxed ectopic

endometrial-like tissue, alongside deficient phagocytic capabilities, have been identified in several

endometriosis studies(*28*). In the peritoneal cavity of mice, the predominant macrophage populations

are small peritoneal macrophages (SpM) and large peritoneal macrophages (LpM), with monocyte-

derived LpM demonstrated to be protective against endometriosis development, possibly due to

enhanced lipid metabolism and cholesterol efflux, highlighting an intrinsic link between macrophage

ontogeny and functionality(*26, 29*). Conversely, endometrial-derived lesion-resident macrophages

enhance lesion growth(*26, 29*) and monocytes that are recruited directly to lesions from the

vasculature differentiate *in situ* into macrophages that exhibit a pro-disease phenotype(*26*).

Macrophages and monocyte subpopulations are also altered in the endometrium and peripheral

blood of women with endometriosis, suggesting systemic myeloid dysfunction(*30*).

Parallels in phenotype and function can be drawn between macrophages in endometriosis and solid

cancers. As with lesions, mononuclear cells are continuously recruited into tumour tissues where they

differentiate into macrophages. Niche specific factors, including hypoxic stress, lead to phenotype

specification and pro-disease functions and an attenuation of phagocytic capabilities(*31*). Tumour

associated macrophages (TAM) are broadly described as having a pro-repair phenotype, however,

similar to endometriosis macrophages, deep immunophenotyping has identified diverse TAM

subpopulations across the phenotypic continuum, including macrophages in highly phagocytic,

cytotoxic, and potentially anti-tumorigenic states(*32-34*). Paradoxically, when appropriately polarised,

tumour-resident macrophages can mediate cancer cell phagocytosis, and enhance tumour

clearance(*35*). Multiple anti-cancer immunotherapies that reprogram TAM have emerged in recent

years, aiming to alter pro-tumorigenic phenotypes and enhance tumour-clearing activity(*36*). Due to

the inherent similarities between the immunopathophysiology, and recent single cell analysis

revealing the presence of pro-disease TAM-like macrophages within endometriotic lesions(*26*), we

have sought to repurpose the therapeutic strategy of macrophage repolarisation for endometriosis.

RRx-001 is a first-in-class small molecule inhibitor currently in several phase I-III clinical trials

for a variety of solid cancers, with properties linked to multiple mechanisms of activity including

macrophage modulation, phagocytic enhancement, vascular normalisation, inflammasome inhibition

and epigenetic modifications, amongst others(*37*). In the current study, we have evaluated the impact

of RRx-001 in preclinical mouse models of experimental endometriosis to determine the effect on

endometriotic lesion growth and associated pain-like behaviours, and to identify any impacts on

oestrous cyclicity and reproductive outcomes. As RRx-001 is known to modulate macrophage

phenotype, we utilised 10x multiomic sequencing to investigate the transcriptomic and epigenetic

landscape of peritoneal macrophages isolated from mice with induced endometriosis, dosed with or

without RRx-001. Collectively, we describe vital preclinical data that indicates RRx-001 could be a

promising new therapeutic for endometriosis

Results

RRx-001 reduced endometriotic lesion growth and attenuated pain-like behaviours in a 'menses'

mouse model of experimental endometriosis. The impact of RRx-001 was evaluated in a syngeneic

bioluminescent 'menses' mouse model of induced endometriosis. In brief, ovariectomised recipient

mice received an intraperitoneal injection of 'menses'-like endometrium and oestradiol

supplementation(*38*). Mice were randomly assigned to a treatment group and received either

intravenous injection of RRx-001 or vehicle control, 6 hours prior to donor tissue transfer, and then

twice weekly for 2 weeks (Fig.1A). *In vivo* bioluminescent imaging at days 3, 7, 10 and 14 was

performed for continuous monitoring of lesions (Fig.1B). Spontaneous resolution and/or lesion

progression was observed in the 'menses' model of endometriosis (as previously documented),

leading to a high intragroup variability of bioluminescent signal(*38*). Despite the occurrence of some

spontaneous resolution, there was no significant difference in bioluminescent signal across the vehicle

group at different timepoints. Conversely, mice dosed with RRx-001 exhibited a reduction between

days 3 and 7, and days 3 and 14 (p=0.0079 and p=0.020, respectively; Fig.1C). Lesions were recovered

at day 14 and histologically verified (Fig.S1); mice dosed with RRx-001 exhibited decreased lesion

number (p=0.013; Fig.1D) and a reduced cross-sectional lesion area (median = 1.48 mm^2), compared

to vehicle control (median = 4 mm^2, p=0.04; Fig.1E). To determine if RRx-001 attenuated evoked pain-

like behaviours, mechanical hyperalgesia was assessed using von Frey filaments on days 11, 12 and 13

post-lesion induction. Mice that received RRx-001 exhibited an increased abdominal retraction

threshold (p=0.0004; Fig.1F), implying reduced abdominal sensitivity, possibly due to the resolution of

lesions within the peritoneal cavity. No differences were observed in hind paw withdrawal

measurements (Fig.1G).

We performed flow cytometry analysis on the peritoneal lavage of endometriosis mice

treated with or without RRx-001 and found no alteration in the abundance of LpM or SpM populations

(Fig.S2A and C-D). As RRx-001 has been reported to modulate macrophage phenotype, and was

previously found to attenuate expression of Sirpα(*39*) (receptor Sirpα-Cd47 ligand interaction

protects 'self' cells from phagocytosis), we investigated the expression of Sirpα on peritoneal

macrophages; expression was increased on LpM isolated from mice with induced endometriosis (p =

0.0046, Fig.S2B and E) and RRx-001 treatment attenuated Sirpα levels (p=0.0053, Fig.S2B and E).

RRx-001 reduced endometriotic lesion growth and attenuated evoked *and* non-evoked pain-like

behaviours in a minimally invasive mouse model of experimental endometriosis. To extend our pre-

clinical testing of RRx-001, we utilised an additional, complementary mouse model of endometriosis.

In this minimally invasive bioluminescent model, full thickness uterine fragments were injected

intraperitoneally into intact recipient mice. Lesions established in this model exhibit less spontaneous

regression and improved longevity(*38*), and since recipients are intact, post-incision pain sensitivity is

avoided allowing improved pain behaviour testing. Mice were randomly assigned to a treatment group

and received a full thickness tissue transfer (day 0). One week following lesion induction, mice were

dosed (day 8) with either RRx-001 or vehicle, and then twice weekly for 2 weeks (Fig.2A). Mice exposed

to RRx-001 exhibited a reduction in lesion bioluminescence from day 7 (pre-treatment) to day 21

(p=0.25; Fig2B, C). No reduction in bioluminescence was observed in the vehicle group. Lesions were

recovered at day 21 and histologically verified. The RRx-001 group had a reduced lesion number

(p=0.0056; Fig.2D) and a reduced cross-sectional area (median = 1.6 mm^2) compared to the vehicle

treated mice (median = 3.1 mm^2, p=0.020; Fig.2E).

To perform a comprehensive investigation into the impact of RRx-001 on endometriosis-

associated pain, we used multiple evoked and non-evoked behavioural tests at i) baseline (pre-

endometriosis induction), ii) after endometriosis induction but prior to treatment (days 7/8) and iii)

after RRx-001/vehicle treatment (days 17/18; Fig.2A). To determine if RRx-001 treatment led to an

attenuation of evoked pain-like behaviours, mechanical and thermal allodynia were assessed using

von Frey filaments and the hot plate test, respectively. We did not detect any significant differences

between groups prior to endometriosis induction (day -1/-2) or pre-treatment (day 7/8) across evoked

behavioural tests (Fig.3A-C). Post treatment, RRx-001 treatment led to a significant decrease in

mechanical allodynia, with increased withdrawal thresholds in both abdominal (p=0.002; Fig.3A) and

hind paw measurements (p=0.013; Fig.3B). A significant increase in hot plate latency at day 17 in RRx-

001 treated mice (p=0.0064; Fig.3C) was also detected. Next, we assessed the impact of RRx-001 on

non-evoked pain-like behaviour using the ethologically relevant measures, thermal plate preference,

burrowing and nesting. Burrowing and nesting are known to be reduced or eliminated with decreased

well-being and can therefore be used as a surrogate marker of visceral pain evoked by

endometriosis(*40*). Prior to endometriosis induction or pre-treatment at days 7/8, we did not detect

any statistical differences between RRx-001 and vehicle groups in non-evoked behavioural tests

(Fig.3D-F). Post treatment, thermal plate preference testing (TPPT) revealed a significant difference

between groups at day 18 (p=0.005; Fig.3D) which may be indicative of a potential alleviation of

hypothermal sensitivities following RRx-001 treatment. Of note, mice exhibited only exploratory

behaviour on the cold plate, whereas the warm plate was also used for grooming by all mice,

suggesting a thermal preference for warmth, as opposed to cold allodynia. No post-treatment

differences were identified in burrowing behaviour between experimental groups (Fig.3E). RRx-001

treated mice did exhibit significantly increased nest complexity score (p< 0.0001, Fig.3F,

representative nest images shown in Fig.3G) indicating improved well-being due to an attenuation of

endometriosis-associated pain.

RRx-001 impacts the transcriptomic and epigenomic landscape of peritoneal macrophages in mice

with induced endometriosis. 10x single nuclei multiomic profiling was used to generate paired RNA

and ATAC libraries for sequencing to investigate the transcriptomic and epigenomic impacts of RRx-

001 on peritoneal macrophages (Fig.4A). For this experiment, we used the 'menses' model of induced

endometriosis, as the intraperitoneal injection of decidualised endometrial tissue most closely

recapitulates the refluxed menstrual tissue thought to initiate endometriotic lesions. At day 7 post-

lesion induction, fluorescent activated cell sorting (FACs) was utilised to isolate Lineage-, Cd45+

leukocytes from the peritoneal lavage of RRx-001 and vehicle treated endometriosis mice, as well as

sham controls (n=9, n=5, n=3, respectively, Fig.S3A), followed by multiomic processing using the 10X

pipeline. We performed an integrative analysis of peritoneal macrophages combining both modalities

into a 'weighted-nearest-neighbour' (WNN) framework, with any cells identified as non-macrophages

removed (Fig.S3B-E)(*41*). UMAP projection revealed 7 macrophage clusters; inset shows the samples

split by library ID (Fig.4B). We found consistent weighting of RNA and ATAC across clusters (although

an increased RNA modality weight was seen in the proliferative cluster), uniformity of high-quality

nucleosome signal (ATAC) and a consistently high log10GenesPerUMI (RNA) (Fig.S3C). All peritoneal

macrophage clusters were *Cd45+* and *Cd11b+*, and our feature plots of *H2-Ab1* and *Adgre1* enable

visualisation of SpM by low *Adgre1* (F4/80) and high *H2-Ab1* (MHCII) expression, prototypical LpM

with high *Adgre1* and low *H2-Ab1*, and intermediate *Adgre1* in other LpM populations, as previously characterised in mice with induced endometriosis at the single cell level(*26, 42*) (Fig.4C).

An analysis of sample composition by cluster revealed macrophage subpopulations were shared between experimental groups. Prototypical LpM (defined by *Timd4* and characteristic DEGs *Prg4*, *Ltbp1*, *Alox15*) exhibited comparable proportions in each sample, as did proliferative LpM and SpM. The predominant population in the vehicle treated endometriosis mice, termed pro-disease LpM, expressed the DEGs *F13a1, Gas6, Lyve1* and *Nrp1*, previously linked to endometriosis(*30, 42*), this population was markedly reduced in both RRx-001 treated and sham mice (Fig.4D,E). In RRx-001 treated mice, the predominant population was termed pro-resolving LpM due to the associated lesion resolution observed and the expression of genes such as *Ms4a4*, *Ccr2*, *Calhm6*, *Fcrla* and *Saa3* associated with inflammatory monocyte-derived infiltration, response to pathogens / antigens and lipid-associated phenotypes, previously linked to lesion resolution (Fig.4D,E)(*26, 29*). We included sham mice in this analysis, which exhibit a post-surgical loss and subsequent replenishment of prototypical LpM and therefore model a resolution of transient inflammation. As expected, transitory LpM were predominant in sham mice; both populations expressed intermediate levels of *Adgre1* and *Timd4* and were separated by enhanced *Vsig4* expression (in transitory LpM 1; Fig.4D,E), a gene known to represent longer time-in-residency, suggesting transitory LpM1 have been present in the peritoneal niche longer than transitory LpM2(*43*).

Abundant expression of *Ccr2* and low *Timd4* on SpM and pro-resolving LpM indicated recent recruitment/monocytic-derivation of these populations, an ontogeny previously shown to be protective against endometriosis development(*29*); paired ATAC data also demonstrated increased chromatin accessibility of the *Ccr2* gene in these populations and significant peak-gene links (Fig.4F). Top DEGs separating pro-resolving and pro-disease LpM were those linked with monocytic ontogeny and complement protein C1q as well as *Saa3* (Fig.4G). Enhanced mRNA levels of the transcription factor, *Cebpb* (associated with regulating macrophage lipid metabolism and phagocytosis(*44*)) were evident in the pro-resolving population (Fig.4G), suggesting enhanced Cebpb regulation of this phenotype. Conversely, transcription factor footprinting did not support this; with comparable enrichment observed in pro-resolving, pro-disease and prototypical populations. Expected footprinting differences, such as the characteristic Gata6 control of prototypical LpM, were detectable in the ATAC data, with most enrichment observed in prototypical LpM and least enrichment observed in pro-disease LpM (Fig.4H)(*45*). We also performed DNA sequence motif enrichment analysis of the ATAC data to identify overrepresented motifs in the differentially accessible peaks of RRx-001 treated mice (all macrophages), with hypergeometric probability testing of enrichment against a set of background peaks matched for GC content. We identified a strong enrichment of Kruppel-like factor

motifs (Klf15, Klf14, Klf2), with Klf family members having broadly reported roles in the activation and
polarisation of macrophages, recent research has identified highly specific and unexpected roles in
macrophage differentiation for these broadly expressed transcription factors(*46*). The greatest
overrepresentation was seen in the Klf14 (4.03-fold enriched, p < 0.0001) and Klf15 (2.91-fold
enriched, p < 0.0001) motifs, which alongside Klf2, have been linked to modulation of inflammatory
signalling pathways. Sp4 and 9 motifs were also enriched (3.06-fold and 2.96-fold enrichment
respectively). Of particular interest was the enrichment of Nrf1 motifs (5.29-fold enrichment,
p<0.0001); RRx-001 has previous been shown to induce Nrf2 which mediates an antioxidant
response(*47*). Nrf1 works in concert with other transcription factors such as Nrf2 to modulate cellular
responses to oxidative stress and maintain cellular homeostasis.

Next, slingshot was used to perform cell lineage and pseudotime inference analysis on
reduced dimensional data using previously identified WNN clusters. SpM was specified as the initial
cluster as the infiltration and transformation of monocytes into SpM with eventual differentiation into
LpM is known to occur under inflammatory conditions, such as endometriosis. Lineage inference
identified three branching trajectories from SpM (Fig.5A)(*23*); pseudo-temporal trajectories to the
terminal populations of pro-disease LpM, prototypical LpM and proliferative LpM were identified.
Interestingly, pro-resolving LpM appeared to be an intermediate in all trajectories, further supportive
of a recent monocyte-derived ontogeny for this population (Fig.5B). Multinomial logistic regression
was used to assess the pseudo-temporal sample imbalance in individual lineages, while adjusting for
baseline sample probabilities. Macrophages from sham mice were more likely to be enriched further
along in the pseudotime trajectories of both prototypical and proliferative lineages, as expected due
to a greater replenishment post transient inflammation and were less likely to be present in the pro-
disease LpM lineage. Macrophages from vehicle treated mice were most abundant in the pro-disease
lineage and predominant across all pseudotime points indicating increased pro-disease differentiation
(Fig.5B). Conversely, in RRx-001 treated mice, macrophages were present at earlier pseudotime points
in the prototypical and proliferative lineages and were also less present in the pro-disease LpM
lineage, suggesting a reduction in pro-disease phenotypical differentiation. These data led us to
conclude that RRx-001 appears to prevent the acquisition of pro-disease phenotype. As our analysis
revealed pseudo temporal differences at early pseudotime points, we performed an aggregate
analysis of the LpM and SpM populations from RRx-001 and vehicle mice which also indicated
differences in SpM gene expression (Fig.5C). To further probe these differences, we performed gene
set enrichment analysis with ClusterProfiler using GO terms (RRx-001 vs Vehicle), and identified terms
related to activation of macrophages, antigen processing and defence responses (Fig.5D).
Furthermore, KEGG pathways suggested an enhancement of phagosome and lysosomal activity,

alongside antigen processing, with the majority activated in both the LpM and SpM populations (Fig. 5E), supporting the concept that treatment of induced endometriosis with RRx-001 may impact peritoneal myeloid populations early in the differentiation trajectory. In line with RRx-001 enhancing phagocytic activity in macrophage populations we demonstrated reduced average expression level of Sirpα in transcriptomic and gene activity data (inferred from ATAC) compared to vehicle treated mice, corresponding with our flow cytometric analysis at day 14 (Figs.5F, S1C) and indicating that RRx-001 acts to disrupt the Sirpα-CD47 immune evasion signal that is enhanced in endometriosis.

RRx-001 does not impact reproductive function in mice. The therapeutic activity of RRx-001 has been safely evaluated in over 300 patients with no increase in infections or anaemia detected, indicating limited side-effects. Impact on reproductive function has not yet been evaluated(*37*). Currently, gold-standard treatment of endometriosis focuses on surgical intervention or therapeutic ovarian suppression (oral contraceptives/GnRH agonists). A non-hormonal medical alternative that does not interrupt reproductive function is urgently needed. To this end, we sought to determine if pre-pregnancy treatment with RRx-001 led to any aberrant changes to fertility or developmental impacts in mice. Pre-pregnancy intravenous dosing was chosen for these initial experiments as an RRx-001-bound haemoglobin adduct persists for the lifetime of the red blood cell, reported to be 40.7 ± 1.9 (S.D) days in mice, potentially leading to long-term impacts(*48*). Tissue-resident endometrial macrophages dynamically regulate tissue homeostasis and remodelling during the oestrous cycle in mice(*49*), thus we initially investigated the impact of RRx-001 on oestrous cycle length, and within individual stages, to identify any perturbations. Oestrous cycles were synchronised using the Whitten effect(*50*), followed by daily staging by wet vaginal cytology preparations for > 4 complete cycles prior to treatment with either RRx-001 or vehicle for 1-week. Daily staging continued for > 4 further cycles. No significant changes in overall cycle length or individual cycle stage were detected in RRx-001 treated mice pre- or post-treatment (Fig.6A, B).

Next, mice were dosed with RRx-001 or vehicle for 1-week, with harem breeding set up 24-hours post final drug dose for investigations into reproductive function. Within these experiments, pregnancy success rates following copulatory plug detection were as expected within breeding colonies: 13/16 RRx-001 treated mice became pregnant (1 did not plug, 2 not pregnant post-plug) and 14/15 vehicle treated mice became pregnant (1 not pregnant post-plug). Sexual receptivity was assessed by daily seminal plug checks, detected between 0.5 - 4.5 days post-mating. No significant differences in time to seminal plug were observed (Fig.6C). At embryonic day (E)10.5, no significant differences in average embryo number or embryo weight were determined between experimental groups (Fig.6D, E). At E15.5, Hofbauer cells, a heterogenous population of placental macrophages

which broadly exhibit a pro-repair phenotype, were isolated from the chorionic villus of the foetal

placenta and assessed using quantitative spectral flow cytometry. We sought to assess Hofbauer

phenotype; dysregulation of these cells is associated with placental pathologies including inadequate

development, inflammation and infection(*51*). No significant differences were identified in the MFIs

of SIRPα, CD206 and CD40 between RRx-001 and vehicle groups (Fig.6F). As expected for a

predominantly pro-repair population, CD206 was expressed across both experimental groups, with no

expression of pro-inflammatory CD40.

Finally, the remaining pregnant mice and pups were monitored throughout pregnancy and

during the postnatal period. We found no impact of RRx-001 on gestational length (Fig.6G).

Furthermore, mice treated with RRx-001 did not have any alterations to number of pups (ranging from

2-11 in the vehicle group, and 3-11 in the RRx-001 treated group; Fig.6H), or sex ratio of litter (Fig.6I).

Post parturition, mothers remained healthy and omental milk spots were observed in all pups on day

1 indicative of successful lactation and nursing. Pups were weighed at postnatal day 7, with weekly

weighing up to 28 days of age, with no significant differences detected at any timepoint (Fig.6J).

Collectively, these data indicate that RRx-001 does not impact reproductive function.

Discussion

In the current study we have revealed that RRx-001 has a robust therapeutic impact in mouse models

of induced endometriosis; treatment decreased lesion number and size, and attenuated pain

associated behaviours. Using multiomic single nuclei sequencing we determined an evident impact on

macrophage populations whereby the abundance of pro-disease macrophages was reduced, and the

proportion of pro-resolving macrophages was enhanced. Trajectory analysis inferred that RRx-001

tips the balance in favour of pro-resolving macrophage by preventing differentiation to a pro-disease

phenotype. Finally, preliminary investigations into reproductive function suggest that RRx-001 does

not impede fertility or aberrantly affect embryonic development *in utero.* Taken together, we present

promising preclinical data highlighting RRx-001 as novel macrophage directed therapy for the

treatment of endometriosis.

In the 'menses' model of induced endometriosis we found that when treatment with RRx-001

was initiated prior to establishment of lesions, there was no difference in bioluminescent signal

between vehicle control and drug treated mice on day 3, indicating that RRx-001 does not impact the

initial stages of ectopic endometrial tissue adhesion within the peritoneal cavity. This aligns with the

proposed erythrophagoimmunotherapeutic activity of RRx-001; in which the covalent encapsulation

of RRx-001 in red blood cells vectors RRx-001 to hypoxic (tumour) vasculature – leading to vasculature

occlusion, haemolysis and erythrophagocytosis by tumour associated macrophages(*39*). This

phenomenon may also be occurring in lesions, which are also exhibit hypoxic areas, following the
initial stages of vascularisation that occur in the first week(*52*). Interestingly, in the menses model,
RRx-001 treatment led to a reduction in bioluminescent signal between days 3 to 7 and days 3 to 14,
but not day 3 to 10, this may be related to cases seen in the treatment of lung cancer with RRx-001,
when tumour pseudo-progression is detected early in treatment due to tumoral lymphocyte
infiltration, before regression occurs later, leading us to hypothesise that this may also be occurring
within the endometriotic lesions(*52*). Future studies will establish the dynamics of pseudo-progression
if present, and determine the regime required for elimination of lesions.

We also investigated the impact of RRx-001 in a minimally invasive model of induced
endometriosis; we previously determined that endometriotic lesions established using this model
exhibit greater longevity(*38*). In this model, full thickness uterine fragments are transferred containing
myometrial tissue, and resulting lesions are usually larger than in the menses model and have a cystic
appearance. We used two alternate models in this study to model alternate 'phenotypes' of
endometriosis, with the minimally invasive model representing a more persistent form of
endometriosis. Here, we also demonstrated a significant reduction in lesion growth when treatment
was initiated 1-week post-lesion induction, rather that prior to lesion induction, again indicative that
RRx-001 resolves established lesions. In both models, treatment with RRx-001 led to an attenuation
of abdominal mechanical hyperalgesia, indicative of reduced pain in this region and perhaps due to
the accompanying resolution of lesions with treatment. We also included non-evoked behavioural
tests as they better recapitulate impacts to well-being that are associated with endometriosis and are
used as surrogate measures of visceral pain. The inclusion of non-evoked behavioural tests were
described recently as an absolute requirement in the replication and translation of the complex pain
experience of endometriosis patients(*40*). In our studies, nest building(*53*) was the most sensitive
assay for identifying behavioural phenotypes; impaired nest building was seen in mice at day 7 post-
lesion induction. Following treatment with RRx-001 nest building was restored to a quality beyond
that observed in naïve, pre-endometriosis mice.

Our multiomic analysis revealed that RRx-001 leads to a reduction in pro-disease LpM and an
accumulation of pro-resolving LpM (prototypical LpM remained constant in each group (sham, RRx-
001, vehicle). Since our trajectory analysis suggested that the pro-resolving LpM phenotype is a
transitory cell state we predict that RRx-001 prevents the acquisition of a pro-disease phenotype by
newly recruited monocyte-derived LpM. The transcriptional phenotype of pro-resolving LpM
exhibited some similarities to that of monocyte-derived LpM observed in our previous studies(*26*)
which were attributed to protective functions in the mouse model of endometriosis. DEGs include
lipid associated genes and those encoding C1q; suggesting roles in the reduction of available lipid,

hormone synthesis and activation of membrane lytic complexes. Treatment with RRx-001 also
appeared to increase genes associated with response to pathogens and antigens suggesting improved
visibility of ectopic tissue to peritoneal macrophages.

Together these may reduce endometriosis drivers and promote destruction of ectopic tissue. RRx-001
has previously been shown to have significant impacts on the epigenome of cancer cells; effects vary
depending on the dose and timepoints evaluated - sites that were hypo- and hyper-methylated as well
as globally increased acetylation were reported(*54*). We predict that in the peritoneal cavity RRx-001
prevents acquisition of pro-disease phenotype via epigenetic modification. Of particular interest, we
found that Klf (including Klf14) binding motifs were enriched in peritoneal macrophages derived from
mice exposed to RRx-00. Previous studies have identified a role for Klf14 in reducing glucose
metabolism and immune function in macrophages; this seems to exert a protective function in
inflammatory conditions by limiting inflammatory cytokine secretion(*46*). These modifications also
highlight the paradoxical nature of RRx-001 mechanisms reported in the literature, whereby
antigenicity is enhanced whilst simultaneously promoting an antioxidative and anti-inflammatory
environment.

Much of the research conducted on the mechanism of action of RRx-001 has occurred in the
oncology field, where studies describe targeting of RRx-001 to hypoxic and aberrant vasculature
prevalent within tumours via binding to haemoglobin within red blood cells. Drug bound RBCs travel
to hypoxic vasculature where they accumulate because of their RRx-001 induced rigidity(*55*). During
this process, nitric oxide (NO) is released which is thought to confer cytotoxic activity within
tumours(*56*). Macrophages also engulf RRx-001 bound RBCs, inducing polarisation from a pro-repair
to pro-inflammatory state and down-regulating the CD47-SIRPα pathway, which reduces tumour cell
immune evasion(*57, 58*). Immunogenicity of cancer cells is also induced via increased expression of
interferon (IFN)-responsive genes(*59*). Given the similarity that exist between cancer and
endometriosis (including pro-disease macrophage polarisation and pathological immunotolerance)
and that endometriosis lesions also exhibit a hypoxic environment, we predict that significant RRx-001
impacts are also occurring within lesions. Future work will interrogate the intra-lesional environment
in response to RRx-001. The mediation of indirect effects within associated environments has not yet
been characterized, however from our results it is plausible to suggest that the modulation of
macrophage populations within the peritoneal cavity occurs as a secondary outcome resulting from
endometriotic lesion resolution by lesion-resident macrophages subsequently reducing the
inflammatory burden within the peritoneal cavity, modified cytokine signalling from intra-lesional
immune cells, the release of reactive oxygen and nitrogen species, or a combination of these impacts.

Thus, further work is required to elucidate the impacts of RRx-001 on lesion-resident macrophage
populations, including interrogation of off-target effects in other tissues.

Our initial results evaluating fertility in mice exposed to RRx-001, indicate that pre-pregnancy
treatment with RRx-001 does not impair the normal reproductive function of female mice or embryo
development indicating that it has potential suitability as a non-hormonal and fertility sparing
treatment for endometriosis. Further work will identify whether RRx-001 can rescue endometriosis-
associated infertility.

To summarize, we have revealed a novel indication for the anti-cancer drug RRx-001 in the
treatment of endometriosis. It acts to reduce endometriosis lesions, associated pain-like behaviours
and modulates macrophage populations in the peritoneal cavity without negatively impacting fertility.
RRx-001 is currently in phase I - III clinical trials for different solid cancers, is well tolerated with
available safety data. In this first phase of preclinical testing, we highlight a robust effect in
experimental endometriosis and plan to proceed to a feasibility trial in women with the disorder. RRx-
001 and its macrophage modifying actions has the potential to improve the lives of millions of women
suffering from endometriosis.

Acknowledgements. The authors thank Dr Laura Baxter at the Bioinformatics RTP, University of
Warwick, for pre-processing of multiomic data. We would also like to thank Deborah McIntyre for the
independent quantitative scoring of nests.

Funding. This work was supported by an MRC iCASE (Ferring Pharmaceuticals were the industrial
partner and provided additional funding for consumables and travel) studentship to I.M and an MRC
Project Grant (MR/S002456) to E.G. The authors declare no conflict of interest.

Author contributions: E.G conceptualised and supervised the study and acquired funding. I.M
conducted animal experiments, analysed data, and performed multiomic analyses. V.V performed
multiomic analyses. C.L.P and M.R performed experiments and analysed data. P.D and K.P performed
animal experiments. I.M and E.G wrote the manuscript.

Data and materials availability: All data associated with this study is present within the manuscript
and the included supplementary materials. Analysis script for both multiomic and cross-species
analyses are publicly available on github.com under project x. Multiomic sequencing data are
deposited in Gene Expression Omnibus under accession number x.

Materials and Methods

Study design. These experiments were designed to determine the impact of RRx-001 treatment in a mouse model of experimental endometriosis and to identify any potential impact on fertility outcomes. For the purpose of determining the impact of RRx-001 on induced endometriosis, we implemented two complementary mouse models ('menses'(*60*) and 'minimally-invasive'(*38*)) in which to robustly examine endometriotic lesion growth and pain-like behaviours. Additionally, we utilised multiomic sequencing to determine transcriptomic and epigenomic effects of RRx-001 on peritoneal macrophages isolated from the 'menses' mouse model. To assess fertility outcomes, we examined the effect of pre-pregnancy dosing of RRx-001 on the oestrous cycle and reproductive functions. Animals were randomised to either RRx-001 treatment or control group using the GraphPad QuickCalcs online tool (https://www.graphpad.com/quickcalcs/randomize1/). Behavioural experiments were performed under blinded conditions. Details of mouse models, sample size and statistical methods are described in the appropriate sections below.

Animals. Wild-type FVB/N mice were purchased from either Envigo (Cambridge, UK) or Charles River (UK) and maintained under standard conditions at the University of Warwick, UK. FVB-Tg(CAG-luc,-GFP)L2G85Chco/J (stock number 008450|L2G85) were originally purchased from The Jackson Laboratory (Bar Harbor, ME, USA), and a breeding colony maintained under standard conditions at the University of Warwick. Mice were housed in standard conditions with a 12:12 light-dark cycle, an ambient temperature of 21°C, humidity at 50%, and *ad libitum* access to food (standard rodent chow) and water. The diet of weanlings was supplemented by apple sauce. Following arrival to the facility, mice acclimatised for one week prior to commencement of experimental procedures. All procedures performed were permitted under license by the United Kingdom Home Office Animal Experimentation (Scientific Procedures) Act; PPL PP0568394, and in accordance with the guidelines of the Committee for Research and Ethical Issues of the International Association for the Study of Pain.

Mouse models of experimental endometriosis. We implemented two variations of the bioluminescent syngeneic mouse models of induced endometriosis; a 'menses' model and a minimally invasive model, described in detail previously(*38, 60*). In brief, for the 'menses' model, recipient FVB/N mice (n=24) were ovariectomised and received an intraperitoneal injection of decidualised and progesterone withdrawn donor endometrium recovered from FVB/Cag-Luc mice (~40mg, minced, resuspended in saline and passed through an 18 g needle) and exogenous oestradiol supplementation (oestradiol valerate 500 ng, twice weekly). Surgical controls, referred to as 'sham' mice, were ovariectomised FVB/N mice that received intraperitoneal injection of saline instead of tissue. In the

minimally invasive model, intact recipient FVB/N mice (n = 32) received an intraperitoneal injection of

FVB/Cag-Luc donor full-thickness uterus (equivalent to 1 uterine horn per recipient, minced and

passed through an 18 g needle). Oestrus stage was determined in donors and recipients by vaginal

smear and retrospective cytological analysis(*61*)

Experimental groups. Mice with induced endometriosis were randomly assigned to either (1) RRx-001

treatment or (2) vehicle control group. In the bioluminescent experiments, groups included were (1)

induced endometriosis and RRx-001 treatment ('menses' model n=13, minimally invasive model

n=16), or (2) induced endometriosis and vehicle control ('menses' model n=11, minimally invasive

model n=16), with dosing schedules as indicated in experimental schematics. In the 10x multiome

experiment, groups included were (1) induced endometriosis ('menses') and RRx-001 treatment (n=9),

(2) induced endometriosis and vehicle control (n=5), and (3) sham (n=3). In the fertility experiments,

mice were randomly assigned to either the (1) RRx-001 treatment (n=20), or (2) vehicle control group

(n=18), with sample size and dosing schedules as described within the results for individual

experiments. In all our studies, RRx-001 (Selleck Chemicals, USA) was dosed intravenously at 10mg/kg

in DMA:PEG-400 1:2, diluted 1:4 in sterile saline immediately prior to injection, 2x weekly (Monday

and Thursday, 12pm), with control mice receiving intravenous vehicle only.

Non-invasive *in vivo* bioluminescence quantification. As described previously, mice were injected

subcutaneously with 1.5 mg D-Luciferin Potassium Salt (75mg/kg, reconstituted in sterile PBS and

sterilised by 0.22 μm filtration), following a 5-minute period to allow a maximal luciferase signal

plateau, mice were imaged under maintained anaesthesia using a PhotonIMAGER (Biospace Lab,

Paris)(*38*). Lesion bioluminescence was measured using a standard region of interest across

experiments, on the front and back of each mouse for 7 minutes per side. Using M3 Vision software

(Biospace Lab), the background reading was subtracted, front and back readings averaged, and

photons/s/cm2/sr calculated for bioluminescent signal quantification.

Histological and ImageJ analyses. Post tissue fixation and embedding in paraffin, 3 μM sections of

recovered lesions were stained using a standard haematoxylin/eosin protocol. Images were taken on

a LeicaDMI microscope (Leica, Germany), with white balance and histological verification of

endometriotic lesion hallmarks (stroma +/- glandular epithelium) conducted in FIJI (ImageJ2, Version

2.3.0). In FIJI, analysis to determine cross-sectional area was performed on lesion images captured

using 4x magnification, image scale was set to a known size and the 'select' tool was implemented to

outline the lesion while excluding peritoneal wall/adipose tissue, and the 'measure' function was then

used to calculate cross-sectional lesion area.

Behavioural assessments.

von Frey. We determined mechanical allodynia as an average of measurements taken on days 11/12/13 post lesion induction ('menses' model) or day -1 (baseline), 8 (1 week; initiation of treatment) and 18 (10 days following start of treatment; minimally invasive model) using calibrated Semmes-Weinstein von Frey filaments (Stoelting, Wood Vale, IL, USA), as described previously[38, 62]. Briefly, mice were allowed to adapt to the apparatus until exploratory responses ceased. Filaments were then applied 10 times in ascending order to the abdomen or hind paw. The filament force (grams) was recorded when withdrawal responses were observed in 50% of applications

Hot plate. Thermal hyperalgesia was determined prior to lesion induction, and on day 7 and day 17 post-lesion induction using a hot plate (Ugo Basile, Italy). Mice were positioned onto a preheated plate set at 52°C and their responses were observed. A time-to response (seconds) was noted when mice exhibited a reaction to the thermal stimulus. If no response was observed within 25 seconds, the mice were removed from the plate.

Thermal plate preference. Thermal hyperalgesia was determined prior to lesion induction, and on day 7 and day 17 post-lesion induction using TPP thermal place preference plates (Ugo Basile, Italy). The thermal plates were set at distinct temperatures, one at 12.5°C and the other at 30°C. Mice were allowed a 15-minute-period for exploration of both plates, during which the duration spent at 12.5°C was recorded.

Burrowing. The ethological rodent behaviour of burrowing was determined prior to lesion induction and on day 8 and day 18 post-lesion induction. Mice were individually supplied with a burrowing tube (diameter 6.8 cm, length 20 cm), made of clear acrylic, open and elevated by 3 cm at one end to prevent the accidental displacement of burrowing substrate (tubes manufactured at Life Sciences, University of Warwick). The mouse was placed in a plastic cage with the burrowing tube filled with 110g of wooden shredding; following a one-hour burrowing assay, the material remaining was weighed and recorded, and the percentage of material displaced within an hour calculated.

Nesting. The ethological rodent behaviour of nesting was determined prior to lesion induction and on day 8 and day 18 post-lesion induction as described[53]. Briefly, 1 hour prior to dark phase, mice were individually housed with wooden shredding and a Nestlet (VetTech, UK). The following morning, the nest is imaged, and the quality of the nest created from the shredded nestlet is recorded with a rating scale of 1-5. (e.g. 1 no nest; 5 fully shredded material arranged into a neat nest). Mice had *ad libitum* access to food and water during the test and were re-grouped immediately following the nesting assay (Table S1).

Flow cytometry. Peritoneal lavage was collected into 5 ml of ice-cold DMEM, red blood cells were lysed (name) and cells filtered through a 40 μM filter and then blocked with monocyte block (name) prior to resuspension and staining of 1 million cells per sample with the antibody panel shown. Samples were analysed using a FACS Aria Fusion with FACS Chorus software (BD Biosciences) and FlowJo v.9 software. Hofbauer cells were isolated from the foetal placenta of mouse embryos at E15.5. Dissection of the foetal placenta from embryonic yolk sacs and subsequent placental digestion into a single cell suspension were performed as described previously, with three foetal placentas pooled from each mouse(*63, 64*). Samples were filtered, RBCs lysed, blocked, and resuspended as above, prior to staining of 1 million cells per sample with the antibody panel. Samples were analysed on a Sony spectral ID7000 spectral cytometer (Sony Biotechnology, USA).

Marker	Fluorophore	Clone	Supplier	Product Number
CD45	Pacific Blue	30-F11	Biolegend	103125
Sirpα	BV510	P84	Biolegend	144005
Ly6C	BV711	HK1.4	Biolegend	128037
CD40	FITC	3/23	Biolegend	124607
c-kit	PerCP Cy5.5	ACK2	Biolegend	135133
CD163	PE	S15049F	Biolegend	156703
CD34	PE-Cy5	MEC14.7	Biolegend	119311
CD206	PE-Cy7	C068C2	Biolegend	141719
Lineage exclusion CD19 CD3 CD335 Ly6G SiglecF	APC	 1D3/CD19 17A2 29A1.4 1A8 S17007L	Biolegend	 152410 100235 137607 127613 155507
MHCII	APC-Cy7	M5/114.15.2	Biolegend	107627
F4/80	APC Fire750	BM8	Biolegend	123117
Live/Dead	Zombie UV	-	Biolegend	423107

Preparation of peritoneal immune cells for 10X multiomic sequencing. Peritoneal lavage was pooled within experimental groups (endometriosis/RRx-001 n=9, endometriosis/vehicle n=5, and sham n=3) prior to processing and staining as described above, with fluorescence activated cell sorting based on surface marker expression (Lin-/Cd45+) performed to enrich samples for macrophage populations.

100,000 cells per sample were sorted into 1ml PBS + 2% FBS using a FACs Aria. Gentle nuclei isolation

was performed, and nuclei were then processed and barcoded according to the established pipelines

to generate individual paired RNA and ATAC libraries using a 10x Nuclei Isolation Kit, a 10x Multiome

Kit and 10x Chromium Controller™ (10x Genomics). Libraries were sequenced individually by the

Genomics Lab at the University of Warwick on high-output 150 cycle cartridges using a NextSeq

500/550 (Illumina®, San Diego, USA).

Bioinformatic analysis. We followed recent guideline best practices for single cell analysis across

modalities as described by Heumos et al for our paired analysis of snRNA-seq and snATAC-seq

libraries(*65*). BCL files for each flow cell directory from the ATAC and GEX sequencing runs were

demultiplexed using Cell Ranger ARC analysis pipelines to generate FASTQ files. Seurat (5.0.3) and

Signac (1.13.0) in R studio (4.3.2) were used to create Seurat objects (v3) based on GEX data, with

ATAC-seq data added as a second assay. Using ATAC-seq data, a common peak set was created and

peaks were called in individual samples using MACS2 v2, with peaks on non-standard chromosomes

and in genomic blacklist regions removed(*66*). Independent quality filtering of each dataset was

performed on both modalities to remove low-quality cells based on number of detected molecules for

each modality, the number of genes detected per UMI, nucleosome signal and TSS enrichment. As

described in datasets generated by Yoshimura et al and 10x Genomics, the gentle nuclei isolation

protocol allows mitochondria to attach to the nuclei prior to GEM generation resulting in a high

mitochondrial percentage(*67*). Post-matrix generation we filtered reads aligned to mitochondrial or

ribosomal regions, and matrices were then subset on common genes. Finally, the scDblFinder and

SoupX were used to identify and remove doublets and assess ambient contamination, respectively(*68,*

69). Pre-processing of RNA datasets was performed using the SCTransform v1 workflow (Seurat),

followed by conservative anchor-based RPCA-integration using the default k.anchor parameter (5) to

align populations (Seurat). We then computed the LSI of individual peak datasets using standard

functionalities (Signac), followed by Harmony integration of datasets(*70*). The Seurat v4 weighted

nearest neighbour (WNN) method of 'FindMultiModalNeighbors' was utilised to compute a joint

neighbour graph to represent both the GEX and ATAC modalities, followed by the use of standard

workflows for WNN UMAP visualisation and clustering (Seurat). Clusters were identified using marker

genes identified in Wilcoxon rank-sum tests in a one-vs-all fashion for manual annotation according

to our previously published annotations of mouse peritoneal macrophages(*26*). Clusters

corresponding to non-macrophage populations or expressing < 1% of total cells were removed in order

to isolate 11,877 peritoneal macrophages (Fig.S3B, C). To account for within-sample correlation we

implemented MAST, a generalised mixed effect model, for differential gene expression testing(*71*).

Differentially accessible peaks were identified utilising logistic regression within the Signac
'FindMarkers' function. DNA sequence motif information (JASPAR2020) was added using the
'AddMotifs' function, with enriched motifs identified using 'FindMotifs'.

For data visualisation purposes the Seurat and Signac functionalities 'DimPlot', 'FeaturePlot',
'DotPlot', 'CoveragePlot', 'CellScatter', 'VlnPlot', 'MotifPlot', and the EnhancedVolcano package were
implemented(*72*) (*73, 74*). The clusterProfiler package was used for statistical analysis, output
simplification and visualisation of GO and KEGG terms(*75*). For trajectory analysis, DynGuidelines was
utilised to select the most applicable package and the indicated Slingshot package were
implemented(*76, 77*).

Statistics
Statistical analyses were performed using GraphPad Prism 10.1.1 for macOS (San Diego, CA). All
graphical data plotted as mean with standard deviation. For data meeting the necessary assumptions
for parametric analyses (Shapiro-Wilk), Student's t-tests or two-way analysis of variance (ANOVA) with
post hoc testing conducted with the Bonferroni test. If non-normality of data was detected, non-
parametric analyses were performed using a Mann-Whitney U test. Statistical significance for all tests
was established at $P < 0.05$. Significance levels were denoted as follows: * $P < 0.05$, ** $P < 0.01$, *** P
< 0.001, **** $P < 0.0001$.

Figures and legends

Figure 1. RRx-001 reduces lesion growth and mechanical allodynia in a 'menses' model of induced endometriosis. (A) Schematic of experimental design. (B) Representative frontal images from days 3, 7, 10 and 14 from RRx-001 and vehicle treated mice. (C) Lesion bioluminescent signal of RRx-001 and vehicle treated mice with endometriosis; statistical analysis performed using a two-way ANOVA with Bonferroni post hoc. (D) Number of lesions and (E) size of lesions recovered from RRx-001 and vehicle mice; statistical analyses performed using a Mann-Whitney test. (F) Mechanical hyperalgesia measured using von Frey filaments applied to the abdomen and (G) hind paw (F and G plotted as an average of 3 measurements taken over days 11-13 post-endometriosis induction; statistical analyses performed using a Mann-Whitney test). * p < 0.05, ** p < 0.01, **** p < 0.001.

Figure 2. The impact of RRx-001 on lesion growth and evoked and non-evoked behavioural tests in

a minimally invasive model of induced endometriosis. (A) Schematic of experimental design. (B)

Representative frontal images from days 3, 7, 14 and 21 from RRx-001 and vehicle treated mice. (C)

Bioluminescent signal of RRx-001 and vehicle treated mice; statistical analysis performed using a two-

way ANOVA with Bonferroni post hoc. (D) Number of lesions and (E) size of lesions recovered from

RRx-001 and vehicle mice; statistical analyses performed using a Mann-Whitney test. * $p < 0.05$, ** p

< 0.01.

Figure 3. RRx-001 reduces evoked and non-evoked pain-associated behaviors in a minimally invasive

model of induced endometriosis. (A) Mechanical allodynia measured using von Frey filaments applied

to the abdomen and (B) hind paw; statistical analyses performed using multiple Mann-Whitney tests.

(C) Hot plate latency assessed at 52°C; statistical analyses performed using multiple unpaired t-tests.

(D) Percentage of material displaced in a burrowing assay; statistical analyses performed using

multiple unpaired t-tests. (E) Thermal plate preference test analysis of % time spent on the 12.5 °C

plate; statistical analyses performed using multiple unpaired t-tests. (F) Nest complexity scores;

statistical analyses performed using multiple unpaired t-tests. (G) Example nests corresponding to

scores 1-5. * p < 0.05, ** p < 0.01, **** p < 0.0001.

Figure 4. Multiomic analyses of RNA and ATAC modalities of peritoneal macrophages. (A) Schematic

of experimental design. (B) UMAP visualisation based on a weighted combination of RNA and ATAC

data from peritoneal macrophages isolated from vehicle treated endometriosis mice, RRx-001 treated

endometriosis mice and sham mice. Inset is UMAP projection split by sample ID. (C) Feature plots of

Ptprc (Cd45), *Itgam* (Cd11b), *Adgre1* (F4/80) and *H2-Ab1* (MhcII) expression levels. (D) Proportion of

each cluster occupied by each sample. (E) Dot plot of featured genes. (F) Chromatin accessibility, gene

expression data and peak-gene links for Ccr2 expression in selected macrophage populations. (G)

Volcano plot of up- and down-regulated DEGs of pro-resolving LpM compared to pro-disease LpM. (H)

Transcription factor footprinting of Cebpb and Gata6 in pro-resolving LpM, pro-disease LpM and

prototypical LpM populations. (I) DNA sequence motif enrichment analysis of ATAC data in peritoneal

macrophages derived from RRx-001 treated endometriosis mice.

Figure 5. RRx-001 impacts differentiation trajectories and phenotypes of LpM and SpM populations in mice with induced endometriosis. (A) Slingshot fitted trajectories and principal curves. (B) Pseudotime ordering of inferred lineages (left) with multinomial logistic regression analysis of pseudo-temporal sample imbalance (right). (C) Scatter plot of aggregate (pseudobulk) expression of LpM and SpM populations from RRx-001 and vehicle treated mice (Pearson correlation reported). (D) ClusterProfiler gene set enrichment analysis using GO terms showing the top 5 terms in the LpM and SpM of RRx-001 treated mice with induced endometriosis. (E) ClusterProfiler gene set enrichment analysis using KEGG pathways showing the top 5 pathways in the LpM and SpM of RRx-001 treated

730 mice with induced endometriosis. (F) Violin plots of *Sirpα* expression in RNA-seq data and the gene

731 activity matrix derived from ATAC-seq data.

Figure 6. Pre-pregnancy treatment with RRx-001 does not cause oestrous cycle perturbations or impact reproductive function. (A) Length of oestrous cycle and (B) days spent within each stage; n=3 vehicle, n=4 RRx-001; statistical analyses performed using Student's t test and two-way ANOVA with Bonferroni post hoc. (C) Time to detection of seminal plug; n=15 RRx-001, n=15 vehicle; statistical analysis performed using a Student's t test. (D) Embryonic number and (E) average weight at E10.5; n=5 RRx-001, n=8 vehicle; statistical analyses performed using a Student's t test. (F) Median fluorescence intensity of SIRPα, CD206 (pro-repair) and CD40 (pro-inflammatory) in Hofbauer cells

769   pooled from 3 placentas per mouse at embryonic day 15.5; n=3 RRx-001, n=4 vehicle, statistical

770   analysis performed using Student's t test. (G) Gestational length of pregnancy, (H) number of pups

771   born to each mouse, (I) sex ratio of litter and (J) average litter weights from postnatal day 7 to

772   postnatal day 28; n=4 RRx-001, n=3 vehicle; statistical analyses performed using Student's t tests. No

773   statistical significance reached $p < 0.05$.

References

1. A. L. Shafrir, L. Farland, D. Shah, H. Harris, M. Kvaskoff, K. Zondervan, S. Missmer, Risk for and consequences of endometriosis: a critical epidemiologic review. *Best practice & research Clinical obstetrics & gynaecology* **51**, 1-15 (2018).
2. K. T. Zondervan, C. M. Becker, K. Koga, S. A. Missmer, R. N. Taylor, P. Viganò, Endometriosis. *Nature Reviews Disease Primers* **4**, 9 (2018).
3. T. Hirata, K. Koga, Y. Osuga, Extra-pelvic endometriosis: a review. *Reproductive medicine and biology* **19**, 323-333 (2020).
4. K. T. Zondervan, C. M. Becker, S. A. Missmer, Endometriosis. *N Engl J Med* **382**, 1244-1256 (2020).
5. K. Ellis, D. Munro, J. Clarke, Endometriosis is undervalued: a call to action. *Frontiers in Global Women's Health* **3**, 54 (2022).
6. J. A. Sampson, Metastatic or embolic endometriosis, due to the menstrual dissemination of endometrial tissue into the venous circulation. *The American journal of pathology* **3**, 93 (1927).
7. J. Halme, M. G. Hammond, J. F. Hulka, S. G. Raj, L. M. Talbert, Retrograde menstruation in healthy women and in patients with endometriosis. *Obstetrics and gynecology* **64**, 151-154 (1984).
8. L. K. Symons, J. E. Miller, V. R. Kay, R. M. Marks, K. Liblik, M. Koti, C. Tayade, The immunopathophysiology of endometriosis. *Trends in molecular medicine* **24**, 748-762 (2018).
9. A. W. Horne, S. A. Missmer, Pathophysiology, diagnosis, and management of endometriosis. *bmj* **379**, (2022).
10. R. O. Burney, L. C. Giudice, Pathogenesis and pathophysiology of endometriosis. *Fertility and sterility* **98**, 511-519 (2012).
11. M. H. Wu, K. Y. Hsiao, S. J. Tsai, Hypoxia: The force of endometriosis. *Journal of Obstetrics and Gynaecology Research* **45**, 532-541 (2019).
12. C. Chapron, L. Marcellin, B. Borghese, P. Santulli, Rethinking mechanisms, diagnosis and management of endometriosis. *Nature Reviews Endocrinology* **15**, 666-682 (2019).
13. K. Nirgianakis, L. Ma, B. McKinnon, M. D. Mueller, Recurrence patterns after surgery in patients with different endometriosis subtypes: a long-term hospital-based cohort study. *Journal of clinical medicine* **9**, 496 (2020).
14. G. Dunselman, N. Vermeulen, C. Becker, C. Calhaz-Jorge, T. D'Hooghe, B. De Bie, O. Heikinheimo, A. Horne, L. Kiesel, A. Nap, ESHRE guideline: management of women with endometriosis. *Human reproduction* **29**, 400-412 (2014).
15. C. M. Becker, W. T. Gattrell, K. Gude, S. S. Singh, Reevaluating response and failure of medical treatment of endometriosis: a systematic review. *Fertility and sterility* **108**, 125-136 (2017).
16. C. Hogg, A. W. Horne, E. Greaves, Endometriosis-associated macrophages: origin, phenotype, and function. *Frontiers in endocrinology* **11**, 7 (2020).
17. A. Shapouri-Moghaddam, S. Mohammadian, H. Vazini, M. Taghadosi, S. A. Esmaeili, F. Mardani, B. Seifi, A. Mohammadi, J. T. Afshari, A. Sahebkar, Macrophage plasticity, polarization, and function in health and disease. *J Cell Physiol* **233**, 6425-6440 (2018).
18. F. Ginhoux, M. Guilliams, Tissue-resident macrophage ontogeny and homeostasis. *Immunity* **44**, 439-449 (2016).
19. P. Italiani, D. Boraschi, From monocytes to M1/M2 macrophages: phenotypical vs. functional differentiation. *Frontiers in immunology* **5**, 514 (2014).

850   20.   M. Bacci, A. Capobianco, A. Monno, L. Cottone, F. Di Puppo, B. Camisa, M. Mariani, C.
851   Brignole, M. Ponzoni, S. Ferrari, P. Panina-Bordignon, A. A. Manfredi, P. Rovere-Querini,
852   Macrophages are alternatively activated in patients with endometriosis and required for
853   growth and vascularization of lesions in a mouse model of disease. *Am J Pathol* **175**, 547-
854   556 (2009).
855   21.   R. Forster, A. Sarginson, A. Velichkova, C. Hogg, A. Dorning, A. W. Horne, P. T. K.
856   Saunders, E. Greaves, Macrophage-derived insulin-like growth factor-1 is a key
857   neurotrophic and nerve-sensitizing factor in pain associated with endometriosis. *Faseb j* **33**,
858   11210-11222 (2019).
859   22.   T. N. Ramírez-Pavez, M. Martínez-Esparza, A. J. Ruiz-Alcaraz, P. Marín-Sánchez, F.
860   Machado-Linde, P. García-Peñarrubia, The role of peritoneal macrophages in
861   endometriosis. *International Journal of Molecular Sciences* **22**, 10792 (2021).
862   23.   C. C. Bain, C. A. Hawley, H. Garner, C. L. Scott, A. Schridde, N. J. Steers, M. Mack, A.
863   Joshi, M. Guilliams, A. M. I. Mowat, Long-lived self-renewing bone marrow-derived
864   macrophages displace embryo-derived cells to inhabit adult serous cavities. *Nature
865   communications* **7**, ncomms11852 (2016).
866   24.   Y. Tan, W. F. Flynn, S. Sivajothi, D. Luo, S. B. Bozal, M. Davé, A. A. Luciano, P. Robson,
867   D. E. Luciano, E. T. Courtois, Single-cell analysis of endometriosis reveals a coordinated
868   transcriptional programme driving immunotolerance and angiogenesis across eutopic and
869   ectopic tissues. *Nat Cell Biol* **24**, 1306-1318 (2022).
870   25.   M. Guo, C. Bafligil, T. Tapmeier, C. Hubbard, S. Manek, C. Shang, F. O. Martinez, N.
871   Schmidt, M. Obendorf, H. Hess-Stumpp, T. M. Zollner, S. Kennedy, C. M. Becker, K. T.
872   Zondervan, A. P. Cribbs, U. Oppermann, Mass cytometry analysis reveals a distinct immune
873   environment in peritoneal fluid in endometriosis: a characterisation study. *BMC Med* **18**, 3
874   (2020).
875   26.   Y. Henlon, K. Panir, I. McIntyre, C. Hogg, P. Dhami, A. O. Cuff, A. Senior, N.
876   Moolchandani-Adwani, E. T. Courtois, A. W. Horne, M. Rosser, S. Ott, E. Greaves, Single-cell
877   analysis identifies distinct macrophage phenotypes associated with pro-disease and pro-
878   resolving functions in the endometriotic niche. *bioRxiv*, 2024.2003.2007.583861 (2024).
879   27.   G. Zou, J. Wang, X. Xu, P. Xu, L. Zhu, Q. Yu, Y. Peng, X. Guo, T. Li, X. Zhang, Cell subtypes
880   and immune dysfunction in peritoneal fluid of endometriosis revealed by single-cell RNA-
881   sequencing. *Cell Biosci* **11**, 98 (2021).
882   28.   J. Li, S. Yan, Q. Li, Y. Huang, M. Ji, X. Jiao, M. Yuan, G. Wang, Macrophage-associated
883   immune checkpoint CD47 blocking ameliorates endometriosis. *Mol Hum Reprod* **28**,
884   (2022).
885   29.   C. Hogg, K. Panir, P. Dhami, M. Rosser, M. Mack, D. Soong, J. W. Pollard, S. J. Jenkins,
886   A. W. Horne, E. Greaves, Macrophages inhibit and enhance endometriosis depending on
887   their origin. *Proc Natl Acad Sci U S A* **118**, (2021).
888   30.   J. Vallvé-Juanico, A. F. George, S. Sen, R. Thomas, M.-G. Shin, D. Kushnoor, J. J.
889   Vásquez, K. C. Vo, J. C. Irwin, N. R. Roan, Deep immunophenotyping reveals endometriosis
890   is marked by dysregulation of the mononuclear phagocytic system in endometrium and
891   peripheral blood. *BMC medicine* **20**, 1-19 (2022).
892   31.   A. Labani-Motlagh, M. Ashja-Mahdavi, A. Loskog, The tumor microenvironment: a
893   milieu hindering and obstructing antitumor immune responses. *Frontiers in immunology*
894   **11**, 940 (2020).
895   32.   K. Movahedi, D. Laoui, C. Gysemans, M. Baeten, G. Stangé, J. Van den Bossche, M.
896   Mack, D. Pipeleers, P. In't Veld, P. De Baetselier, Different tumor microenvironments

contain functionally distinct subsets of macrophages derived from Ly6C (high) monocytes.
Cancer research **70**, 5728-5739 (2010).

33. E. Azizi, A. J. Carr, G. Plitas, A. E. Cornish, C. Konopacki, S. Prabhakaran, J. Nainys, K.
Wu, V. Kiseliovas, M. Setty, Single-cell map of diverse immune phenotypes in the breast
tumor microenvironment. *Cell* **174**, 1293-1308. e1236 (2018).

34. Y. Pan, Y. Yu, X. Wang, T. Zhang, Tumor-associated macrophages in tumor immunity.
Frontiers in immunology **11**, 583084 (2020).

35. A. Mantovani, F. Marchesi, A. Malesci, L. Laghi, P. Allavena, Tumour-associated
macrophages as treatment targets in oncology. *Nature reviews Clinical oncology* **14**, 399-
416 (2017).

36. A. Mantovani, P. Allavena, F. Marchesi, C. Garlanda, Macrophages as tools and targets
in cancer therapy. *Nature Reviews Drug Discovery* **21**, 799-820 (2022).

37. N. Jayabalan, B. Oronsky, P. Cabrales, T. Reid, S. Caroen, A. M. Johnson, N. A. Birch, J.
D. O'Sullivan, R. Gordon, A review of RRx-001: a late-stage multi-indication inhibitor of
NLRP3 activation and chronic inflammation. *Drugs* **83**, 389-402 (2023).

38. A. Dorning, P. Dhami, K. Panir, C. Hogg, E. Park, G. D. Ferguson, D. Hargrove, J. Karras,
A. W. Horne, E. Greaves, Bioluminescent imaging in induced mouse models of
endometriosis reveals differences in four model variations. *Dis Model Mech* **14**, (2021).

39. B. Oronsky, C. A. Carter, S. Caroen, C. Scribner, A. Oronsky, T. R. Reid, RRx-001, a first-
in-class small molecule inhibitor of MYC and a downregulator of CD47, is an"
erythrophagoimmunotherapeutic". *OncoImmunology* **9**, 1746172 (2020).

40. M. A. Tejada, C. Antunez, P. Nunez-Badinez, B. De Leo, P. T. Saunders, K. Vincent, A.
Cano, J. Nagel, R. Gomez, Rodent Animal Models of Endometriosis-Associated Pain: Unmet
Needs and Resources Available for Improving Translational Research in Endometriosis.
International Journal of Molecular Sciences **24**, 2422 (2023).

41. Y. Hao, S. Hao, E. Andersen-Nissen, W. M. Mauck, S. Zheng, A. Butler, M. J. Lee, A. J.
Wilk, C. Darby, M. Zager, Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-
3587. e3529 (2021).

42. L. Zhao, M. Shi, S. Winuthayanon, J. A. MacLean, K. Hayashi, Niclosamide targets the
dynamic progression of macrophages for the resolution of endometriosis in a mouse model.
Communications Biology **5**, 1225 (2022).

43. P. A. Louwe, L. Badiola Gomez, H. Webster, G. Perona-Wright, C. C. Bain, S. J. Forbes,
S. J. Jenkins, Recruited macrophages that colonize the post-inflammatory peritoneal niche
convert into functionally divergent resident cells. *Nat Commun* **12**, 1770 (2021).

44. D. Dörr, B. Obermayer, J. M. Weiner, K. Zimmermann, C. Anania, L. K. Wagner, E. M.
Lyras, V. Sapozhnikova, D. Lara-Astiaso, F. Prósper, C/EBPβ regulates lipid metabolism and
Pparg isoform 2 expression in alveolar macrophages. *Science Immunology* **7**, eabj0140
(2022).

45. M. Rosas, L. C. Davies, P. J. Giles, C.-T. Liao, B. Kharfan, T. C. Stone, V. B. O'Donnell, D.
J. Fraser, S. A. Jones, P. R. Taylor, The transcription factor Gata6 links tissue macrophage
phenotype and proliferative renewal. *Science* **344**, 645-648 (2014).

46. Y. Yuan, G. Fan, Y. Liu, L. Liu, T. Zhang, P. Liu, Q. Tu, X. Zhang, S. Luo, L. Yao, F. Chen, J.
Li, The transcription factor KLF14 regulates macrophage glycolysis and immune function by
inhibiting HK2 in sepsis. *Cellular & Molecular Immunology* **19**, 504-515 (2022).

47. B. Oronsky, L. Takahashi, R. Gordon, P. Cabrales, S. Caroen, T. Reid, RRx-001: a
chimeric triple action NLRP3 inhibitor, Nrf2 inducer, and nitric oxide superagonist. *Front
Oncol* **13**, 1204143 (2023).

48.   L. Van Putten, F. Croon, The life span of red cells in the rat and the mouse as
determined by labeling with DFP32 in vivo. *Blood* **13**, 789-794 (1958).

49.   L. Zhang, W. Long, W. Xu, X. Chen, X. Zhao, B. Wu, Digital cell atlas of mouse uterus:
from regenerative stage to maturational stage. *Frontiers in Genetics* **13**, 847646 (2022).

50.   W. K. Whitten, Modification of the oestrous cycle of the mouse by external stimuli
associated with the male. *Journal of Endocrinology* **13**, 399-404 (1956).

51.   L. Reyes, B. Wolfe, T. Golos, Hofbauer cells: placental macrophages of fetal origin.
Macrophages: origin, functions and biointervention, 45-60 (2017).

52.   C. A. Carter, B. Schmitz, P. G. Peterson, M. Quinn, A. Degesys, J. Jenkins, B. Oronsky, J.
Scicinski, S. Caroen, T. R. Reid, Immune reactivity and pseudoprogression or tumor flare in
a serially biopsied neuroendocrine patient treated with the epigenetic agent RRx-001. *Case
reports in oncology* **9**, 164-170 (2016).

53.   R. M. Deacon, Assessing nest building in mice. *Nat Protoc* **1**, 1117-1119 (2006).

54.   H. Zhao, S. Ning, J. Scicinski, B. Oronsky, S. J. Knox, D. M. Peehl, Epigenetic effects of
RRx-001: a possible unifying mechanism of anticancer activity. *Oncotarget* **6**, 43172-43181
(2015).

55.   T. Reid, B. Oronsky, J. Scicinski, C. L. Scribner, S. J. Knox, S. Ning, D. M. Peehl, R. Korn,
M. Stirn, C. A. Carter, A. Oronsky, M. J. Taylor, W. L. Fitch, P. Cabrales, M. M. Kim, H. A.
Burris, C. D. Lao, N. E. D. Abrouk, G. R. Fanger, J. R. Infante, Safety and activity of RRx-001
in patients with advanced cancer: a first-in-human, open-label, dose-escalation phase 1
study. *The Lancet Oncology* **16**, 1133-1142 (2015).

56.   M. H. Fens, P. Cabrales, J. Scicinski, S. K. Larkin, J. H. Suh, F. A. Kuypers, N. Oronsky, M.
Lybeck, A. Oronsky, B. Oronsky, Targeting tumor hypoxia with the epigenetic anticancer
agent, RRx-001: a superagonist of nitric oxide generation. *Med Oncol* **33**, 85 (2016).

57.   B. Oronsky, P. Cabrales, S. Caroen, X. Guo, C. Scribner, A. Oronsky, T. R. Reid, RRx-001,
a downregulator of the CD47- SIRPα checkpoint pathway, does not cause anemia or
thrombocytopenia. *Expert Opin Drug Metab Toxicol* **17**, 355-357 (2021).

58.   P. Cabrales, RRx-001 Acts as a Dual Small Molecule Checkpoint Inhibitor by
Downregulating CD47 on Cancer Cells and SIRP-α on Monocytes/Macrophages. *Transl
Oncol* **12**, 626-632 (2019).

59.   H. Zhao, S. Ning, R. Nolley, J. Scicinski, B. Oronsky, S. J. Knox, D. M. Peehl, The
immunomodulatory anticancer agent, RRx-001, induces an interferon response through
epigenetic induction of viral mimicry. *Clin Epigenetics* **9**, 4 (2017).

60.   E. Greaves, F. L. Cousins, A. Murray, A. Esnal-Zufiaurre, A. Fassbender, A. W. Horne, P.
T. Saunders, A novel mouse model of endometriosis mimics human phenotype and reveals
insights into the inflammatory contribution of shed endometrium. *Am J Pathol* **184**, 1930-
1939 (2014).

61.   A. C. McLean, N. Valenzuela, S. Fai, S. A. Bennett, Performing vaginal lavage, crystal
violet staining, and vaginal cytological evaluation for mouse estrous cycle staging
identification. *J Vis Exp*, e4389 (2012).

62.   E. Greaves, A. W. Horne, H. Jerina, M. Mikolajczak, L. Hilferty, R. Mitchell, S. M.
Fleetwood-Walker, P. T. Saunders, EP2 receptor antagonism reduces peripheral and central
hyperalgesia in a preclinical mouse model of endometriosis. *Sci Rep* **7**, 44169 (2017).

63.   K. Ottersbach, E. Dzierzak, in *Leukemia: Methods and Protocols,* C. W. Eric So, Ed.
(Humana Press, Totowa, NJ, 2009), pp. 335-346.

64.   G. Liang, B. Huang, F. Wang, F. Liu, Protocols for isolation and characterization of
mouse placental hemogenic endothelial cells. *STAR protocols* **2**, 100884 (2021).

65.   L. Heumos, A. C. Schaar, C. Lance, A. Litinetskaya, F. Drost, L. Zappia, M. D. Lücken, D.
C. Strobl, J. Henao, F. Curion, Best practices for single-cell analysis across modalities. *Nature*
Reviews Genetics **24**, 550-572 (2023).
66.   Y. Zhang, T. Liu, C. A. Meyer, J. Eeckhoute, D. S. Johnson, B. E. Bernstein, C. Nusbaum,
R. M. Myers, M. Brown, W. Li, Model-based analysis of ChIP-Seq (MACS). *Genome biology*
9, 1-9 (2008).
67.   Y. Yoshimura, Y. Muto, N. Ledru, H. Wu, K. Omachi, J. H. Miner, B. D. Humphreys, A
single-cell multiomic analysis of kidney organoid differentiation. *Proceedings of the*
National Academy of Sciences **120**, e2219699120 (2023).
68.   P.-L. Germain, A. Lun, C. G. Meixide, W. Macnair, M. D. Robinson, Doublet
identification in single-cell sequencing data using scDblFinder. *F1000Research* **10**, (2021).
69.   M. D. Young, S. Behjati, SoupX removes ambient RNA contamination from droplet-
based single-cell RNA sequencing data. *Gigascience* **9**, giaa151 (2020).
70.   I. Korsunsky, N. Millard, J. Fan, K. Slowikowski, F. Zhang, K. Wei, Y. Baglaenko, M.
Brenner, P.-r. Loh, S. Raychaudhuri, Fast, sensitive and accurate integration of single-cell
data with Harmony. *Nature methods* **16**, 1289-1296 (2019).
71.   G. Finak, A. McDavid, M. Yajima, J. Deng, V. Gersuk, A. K. Shalek, C. K. Slichter, H. W.
Miller, M. J. McElrath, M. Prlic, MAST: a flexible statistical framework for assessing
transcriptional changes and characterizing heterogeneity in single-cell RNA sequencing
data. *Genome biology* **16**, 1-13 (2015).
72.   T. Stuart, A. Butler, P. Hoffman, C. Hafemeister, E. Papalexi, W. M. Mauck, Y. Hao, M.
Stoeckius, P. Smibert, R. Satija, Comprehensive integration of single-cell data. *cell* **177**,
1888-1902. e1821 (2019).
73.   K. Blighe, S. Rana, M. Lewis, EnhancedVolcano: Publication-ready volcano plots with
enhanced colouring and labeling. *R package version* **1**, 10.18129 (2019).
74.   T. Stuart, A. Srivastava, S. Madad, C. A. Lareau, R. Satija, Single-cell chromatin state
analysis with Signac. *Nature Methods* **18**, 1333-1341 (2021).
75.   G. Yu, L.-G. Wang, Y. Han, Q.-Y. He, clusterProfiler: an R package for comparing
biological themes among gene clusters. *Omics: a journal of integrative biology* **16**, 284-287
(2012).
76.   W. Saelens, R. Cannoodt, H. Todorov, Y. Saeys, A comparison of single-cell trajectory
inference methods. *Nature biotechnology* **37**, 547-554 (2019).
77.   K. Street, D. Risso, R. B. Fletcher, D. Das, J. Ngai, N. Yosef, E. Purdom, S. Dudoit,
Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics. *BMC*
genomics **19**, 1-16 (2018).
1026