Diversity in Chemical Subunits and Linkages: A Key Molecular Determinant of Microbial Richness, Microbiota Interactions, and Substrate Utilization

Hugh C. McCullough[a,b], Hyun-Seob Song[a,b,c#] and Jennifer M. Auchtung[a,b#]

[a]Nebraska Foods for Health Center, University of Nebraska-Lincoln, Lincoln, NE USA

[b]Department of Food Science and Technology, University of Nebraska-Lincoln, Lincoln, NE USA

[c]Department of Biological Systems Engineering, University of Nebraska-Lincoln, Lincoln, NE USA

Running title: Chemical complexity impacts microbial community structure

#Address correspondence to Hyun-Seob Song, hsong5@unl.edu and Jennifer M. Auchtung, jauchtung2@unl.edu

ABSTRACT

Dietary fibers play a significant role in shaping the composition and function of microbial

communities in the human colon. Our understanding of the specific chemical traits of dietary

fibers that influence microbial diversity, interactions, and function remains limited. Towards

filling this knowledge gap, we developed a novel measure, termed Chemical Subunits and

Linkages (CheSL) Shannon diversity, to characterize the effects of carbohydrate complexity on

human fecal bacteria cultured *in vitro* under controlled, continuous flow conditions using media

that systematically varied in carbohydrate composition. Our analysis revealed that CheSL

Shannon diversity demonstrated a strong Pearson correlation with microbial richness across

multiple fecal samples and study designs. Additionally, we observed that microbial communities

in media with higher CheSL Shannon diversity scores exhibited greater peptide utilization, and

more connected, reproducible structures in computationally inferred microbial interaction

networks. Together, these findings demonstrate that CheSL Shannon diversity can be a useful

tool to quantify the effects of carbohydrate complexity on microbial diversity, metabolic

potential, and interactions. Furthermore, our work highlights how robust and stable community

data can be generated by engineering media composition and structure. These studies provide a

valuable framework for future research on microbial community interactions and their potential

impacts on host health.

IMPORTANCE

For the human adult gut microbiota, higher microbial diversity strongly correlates with positive

health outcomes. This correlation is likely due to increased community resilience that results

from functional redundancy that can occur within diverse communities. While previous studies

have shown that dietary fibers influence microbiota composition and function, we lack a

complete mechanistic understanding of how differences in the composition of fibers are likely to

functionally impact microbiota diversity. To address this need, we developed Chemical Subunits

and Linkages (CheSL) Shannon diversity, a novel measure that describes carbohydrate

complexity. Using this measure, we were able to correlate changes in carbohydrate complexity

with alterations in microbial diversity and interspecies interactions. Overall, these analyses

provide new perspectives on dietary optimization strategies to improve human health.

INTRODUCTION

Members of the gut microbiota are known to contribute to different aspects of human health,

including metabolizing otherwise inaccessible dietary substrates, educating the immune system,

and providing protection from pathogen infection (1–4). Ongoing research efforts focus on

developing treatments and interventions to engineer the gut microbiota to improve host health

(5–8); approaches have included addition of new dietary substrates (e.g., prebiotics and dietary

fibers (9–11)), new microbes (e.g., probiotics and live microbial therapeutics (1, 12, 13)), or

combinations of substrates and microbes (e.g., synbiotics (2, 14)). Previous studies have shown

diets are a major driver of microbial composition (15). In mice, consumption of a westernized

diet caused large changes in microbiota composition compared to consumption of a low fat, low

sugar diet enriched in complex polysaccharides (16, 17). Similarly, long-term dietary patterns

and microbiota composition are correlated in humans (18). Dietary fibers are of particular

interest for gut microbiota modulation because they cannot be metabolized by human enzymes

and gut microbiota metabolism of these substrates can produce products (e.g., butyrate)

metabolized by the host (19–22).

The "nutrient-niche theory" proposed by Freter et al. (23, 24) can serve as a basic

framework to understand the impacts of diets on microbiota composition. It states that the

metabolic potential of microbial community members and the availability of nutrients dictate

how niches are occupied (23, 24). Particularly, increasing the number of unique utilizable

substrates expands the fundamental niche space potentially facilitating increased microbial

richness. However, competition for limiting nutrients and environmental conditions that limit

growth may restrict the ability of microbes to colonize (25). Recent work has begun to

investigate how carbohydrate structural diversity impacts microbial community composition.

Chung et al. demonstrated that more complex fibers and/or mixtures of fibers increased

microbial community richness and diversity compared to simple fibers for communities of

human fecal microbes, although there was some variation between the two fecal communities

tested (26). Yao et al observed similar effects culturing human fecal microbiotas, with the more

complex fiber sorghum arabinoxylan supporting higher microbial community richness and

diversity than the simple fiber inulin or mixtures of the primary monosaccharides found in either

fiber (27). However, a follow-up study observed that the subtle differences in carbohydrate

structure could have significant impacts on microbial richness of fecal communities cultured

under high dilution pressure (28). Using a defined community of ten fecal microbes, Loss et al

demonstrated that higher carbohydrate complexity and low sugar concentrations led to higher

community diversity by increasing the ratio of positive to negative interactions between strains

(29).

In this study, we analyzed the response of human fecal microbial communities to changes

in the diversity and concentration of carbohydrates through a combination of *in vitro*

experiments and computational modeling using a previously described continuous flow model of

the lumen of the distal colon (30). Our workflow was composed of three parts (Fig. 1): (1) data

generation and analysis to identify key traits of carbohydrates determining microbial richness, (2)

post-analysis of metabolic potential and activity, and (3) network inference to predict community

structure - microbial interactions and central microbiota. Inspired by the nutrient-niche theory,

we developed a molecular-level diversity metric of carbohydrates termed Chemical Subunit and

Linkage (CheSL) Shannon diversity, which we used to quantify carbohydrate diversity for

mixtures of carbohydrates based on their internal linkages and subunits. Communities cultured in

media with high CheSL Shannon diversity demonstrated strong positive correlations with

microbial richness, peptide utilization (measured and predicted through Stickland fermentation),

and microbial interactions in computationally inferred networks. Through these combined data

analyses and model predictions, we also determined that microbial communities assembled under

conditions with higher CheSL Shannon diversity led to more reproducible interactions with

conserved central taxa (*Bacteroides*, *Lachnospiraceae* and *Ruminococcaceae*). Finally, we were

able to draw parallels between enrichment of specific microbes in the presence of carbohydrate

mixtures that were consistent with predicted abundances of metabolic pathways and enrichment

previously observed *in vivo*.

RESULTS

Developing Metrics to Quantify Carbohydrate Complexity

To investigate the potential contribution of carbohydrate complexity to unique niches

within a medium, we developed a new metric termed CheSL, which enables quantifying

carbohydrate complexity based on the number of unique free monosaccharides and linkages

present in each carbohydrate. Accounting for internal linkages in evaluating carbohydrate

complexity is critical because polysaccharides may provide one or more unique niches

depending on their complexity and the likelihood that enzymes required for degradation are

encoded within a single microbe may vary based upon linkage complexity. For example,

previous studies have demonstrated that while *Bifidobacterium longum* (31) and *Bacteroides*

thetaiotaomicron, two common members of the gut microbiota, possess enzymes necessary for

complete degradation of the simple fiber starch (32), neither *B. longum* nor *B. caccae* can fully

metabolize arabinogalactan, a more complex polysaccharide composed of five different types of

glycosidic bonds that connect primarily arabinose and galactose monomers (Table 1) (33).

The primary inputs for CheSL include the fractional composition of linkages in each

carbohydrate and the fractional composition of carbohydrates in each medium. This information

can be effectively represented by defining a linkage matrix (**L**) and media matrix (**M**),

respectively.

The matrix **L**, representing the fractional composition of the linkage arrangements present

across the carbohydrates contained in all media, can be given as follows:

125
$$\mathbf{L} = \begin{bmatrix} l_{1,1} & \cdots & l_{1,n_C} \\ \vdots & \ddots & \vdots \\ l_{n_L,1} & \cdots & l_{n_L,n_C} \end{bmatrix} \tag{1}$$

Where the rows and columns in **L** represent linkage arrangements and carbohydrates,

respectively; n_L denotes the total number of unique linkages associated with subunits contained

in all carbohydrates; the $(i,j)^{th}$ element of **L**, i.e., $l_{i,j}$, denotes the proportion of linkage

arrangement *i* within carbohydrate *j*.

The matrix **M** representing media-specific carbohydrate composition can be given as

follows:

$$\mathbf{M} = \begin{bmatrix} m_{1,1} & \cdots & m_{1,n_M} \\ \vdots & \ddots & \vdots \\ m_{n_C,1} & \cdots & m_{n_C,n_M} \end{bmatrix} \quad (2)$$

133 Where the rows and columns in $\mathbf{M}$ represent carbohydrates and media, respectively; n_c and n_M

134 denote the total numbers of unique carbohydrates and media used in experiments, respectively;

135 the $(j,k)^{th}$ element of $\mathbf{M}$, i.e., $m_{j,k}$, denotes the concentration of a specific carbohydrate j in a

136 medium k.

$$\mathbf{L} = \begin{bmatrix} l_{1,1} & \cdots & l_{1,n_C} \\ \vdots & \ddots & \vdots \\ l_{n_L,1} & \cdots & l_{n_L,n_C} \end{bmatrix} \quad (3)$$

138 We can also define a $(n_L \times n_M)$ matrix $\mathbf{S}$ to represent the concentrations of linkage

139 arrangements (or subunits) in media, which are readily obtainable by multiplying $\mathbf{L}$ and $\mathbf{M}$, i.e.,

140
$$\mathbf{S} = \mathbf{LM} \quad (4)$$

141 where the rows and columns in $\mathbf{S}$ represent linkage arrangements and media, respectively; the

142 $(i,k)^{th}$ element of $\mathbf{S}$, i.e., $s_{i,k}$, denotes the i^{th} linkage arrangement (or free subunit)

143 concentration for the k^{th} medium. The fractional abundance of the i^{th} linkage arrangement (or

144 free subunit) within the k^{th} medium, denoted by $f_{i,k}$, is simply calculated from $s_{i,k}$ ($i =$

145 $1,2,\dots,n_L$) as follows:

146
$$f_{i,k} = \frac{s_{i,k}}{\sum_{i'=1}^{n_L} s_{i',k}} \quad (5)$$

147 We can finally calculate CheSL Shannon diversity and CheSL Shannon evenness for the

148 k^{th} medium, denoted CSI_k and CSE_k, respectively, using the equations below:

149
$$CSI_k = -\sum_{i \in I_{NZ,k}} f_{i,k} \ln(f_{i,k}) \quad (6)$$

150
$$CSE_k = \frac{CSI_k}{\ln(|I_{NZ,k}|)} \quad (7)$$

$I_{NZ,k}$ denotes the indices of non-zero elements of $\mathbf{s_k}$. For, CheSL Shannon evenness, we divide

CSI_k by the number of non-zero elements in $\mathbf{s_k}$ (or the cardinality of $I_{NZ,k}$), which represents the

CheSL richness.

In the metric we developed, monomeric glucose was assigned a CheSL value of 0,

because pathways to import and metabolize glucose are conserved across many species of

bacteria (34–38) and as a result glucose metabolism was unlikely to provide a unique niche for

microbes within most communities. Other unique monosaccharides (e.g., arabinose, N-

acetylglucosamine, D-glucuronic acid) were assigned CheSL richness values of one, as pathways

for their uptake and utilization were not expected to be as widely conserved as glucose and could

represent unique niches. Other CheSL values for polysaccharides were primarily assigned based

upon the number of unique linkage arrangements, as the capability to metabolize

monosaccharides liberated from polysaccharides was expected to be part of the same unique

niche. With the exception of soluble starch, carbohydrate composition and linkage data were

estimated based upon previously published literature (Table 1).

Using the two hypothetical media exemplified in the bottom of Fig. 2, we illustrate how

to calculate CheSL richness, CheSL Shannon diversity, and CheSL evenness metrics. Across

these two media, there are three carbohydrates. Carbohydrate one has two linkage arrangements,

of which 66% are arrangement one and 33% are arrangement two. Carbohydrate two was a

polysaccharide, with three linkage arrangements (three, four, and five) in equal proportions.

Carbohydrate three contains three linkage arrangements, 50% are linkage arrangement six, 33.33%

are linkage arrangement seven, and 16.66% are linkage arrangement eight. The linkage matrix $\mathbf{L}$

is below:

$$\mathbf{L} = \begin{bmatrix} .6666 & 0 & 0 \\ .3333 & 0 & 0 \\ 0 & .3333 & 0 \\ 0 & .3333 & 0 \\ 0 & .3333 & 0 \\ 0 & 0 & .5 \\ 0 & 0 & .3333 \\ 0 & 0 & .1666 \end{bmatrix} \quad (8)$$

174   The first medium in the study, $\mathbf{m_1}$, contains 1 g/L carbohydrate one and 2 g/L

175   carbohydrate 2. The second medium in the study, $\mathbf{m_2}$, contains 1 g/L carbohydrate one, 2 g/L

176   carbohydrate two and 2 g/L carbohydrate three. The media matrix ($\mathbf{M}$) is shown below:

177
$$\mathbf{M} = \begin{bmatrix} 1 & 1 \\ 2 & 2 \\ 0 & 2 \end{bmatrix} \quad (9)$$

178   The concentration of linkages, $\mathbf{S}$, can be calculated from L and M.

$$\mathbf{S} = \mathbf{LM} = \begin{bmatrix} 1 & 2 & 0 \\ 1 & 2 & 2 \end{bmatrix} \begin{bmatrix} .6666 & .3333 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & .3333 & .3333 & .3333 & 0 & 0 & 0 \\ 0 & 0 & .5 & 0 & 0 & .5 & .3333 & .1666 \end{bmatrix}$$

179
$$= \begin{bmatrix} .6666 & .6666 \\ .3333 & .3333 \\ .6666 & .6666 \\ .6666 & .6666 \\ .6666 & .6666 \\ 0 & 1 \\ 0 & .6666 \\ 0 & .3333 \end{bmatrix} \quad (10)$$

These calculations demonstrate that the medium, $\mathbf{m_1}$ has 0.3333 g/L worth of

carbohydrate associated with linkage two and 0.6666 g/L associated with linkage arrangements

one, three, four, and five. $\mathbf{m_2}$ has 0.666 g/L worth of carbohydrate associated with linkage

arrangements one, three, four, five, and seven, 0.3333 g/L worth of carbohydrate associated with

linkage arrangements two and eight, and 1 g/L worth of carbohydrate associated with linkage

arrangement six.

From this, we calculated the fractional contributions of each linkage pool (f_j and f_k). For f_j,

this was 0.222 for linkage arrangement one, three, four, and five, and 0.111 for linkage

arrangement two. For f_k, this was 0.1333 for linkage one, three, four, five, seven, 0.066 for

linkage arrangements two and eight, and 0.2 for linkage arrangement six.

190
$$f_k = \begin{bmatrix} .2222 & .1333 \\ .1111 & .0666 \\ .2222 & .1333 \\ .2222 & .1333 \\ .2222 & .1333 \\ 0 & .2 \\ 0 & .1333 \\ 0 & .0666 \end{bmatrix} \tag{10}$$

Using this fractional matrix and the formulas described above, we determined that for

medium m_1, the CheSL richness was 5, CheSL Shannon diversity was 1.58, and CheSL Shannon

evenness was 0.982, whereas for medium m_2, the CheSL richness was 8, CheSL Shannon

diversity was 2.03, and CheSL Shannon evenness was 0.97.

CheSL Shannon Diversity Connects Carbohydrate Complexity to Microbial Richness

We hypothesized that increasing the number of unique utilizable substrates would expand

the fundamental niche space and facilitate increased microbial richness. To test this, microbial

communities from the fecal samples of two healthy humans (Fecal Samples A and B (FSA,

FSB)) were cultured in continuous-flow minibioreactor arrays (MBRAs) with media that differed

in composition and abundance of carbohydrates. Previously, we demonstrated that community

assembly in MBRAs was a result of environmental filtering that selected for core, cultivatable

taxa as well as stochastic processes that contributed to variation between replicate communities

(30). We expected that unique utilizable carbohydrate linkages would contribute to community

assembly primarily through environmental filtering but that stochastic processes would also

impact assembly.

Replicate cultures were inoculated into four media (NS, MM, 5MM, and IL) that varied

in composition and abundance of carbohydrates but were composed of the same base of proteins,

lipids, and trace minerals (**Table S1**). All media contained low levels of inulin and

arabinogalactan. NS also contained low levels of mono (glucose) and disaccharides (cellobiose

and maltose). MM contained low levels of mucosal monosaccharides and soluble starch. 5MM

contained higher levels of mucosal monosaccharides and soluble starch compared to MM. IL

was supplemented with ileostomy effluent, which likely contains a complex mixture of

carbohydrates and other factors, requiring approximation of substrate concentrations from

available literature (39, 40). Samples were collected from communities daily and changes in

microbial community composition were determined through sequencing the V4 region of the 16S

rRNA gene. Contamination limited collection of data from 5MM to FSB only.

Contrary to our original expectation, we found that microbiota richness (number of

observed ASVs) was most strongly correlated with CheSL Shannon diversity rather than CheSL

richness (Fig. 3). Specifically, we observed a Pearson correlation coefficient between ASV

richness and CheSL richness of R=0.64 (p=0.0024, Fig. 3A), between ASV richness and CheSL

Shannon evenness of 0.21 (p=0.38, Fig. 3B), and between ASV richness and CheSL Shannon

diversity of R=0.74 (p=0.00019, Fig. 3C). Individual analysis of each fecal sample (Fig. S1A-F),

indicated that the strong correlation was primarily a result of observations with FSB, as

correlations for FSA were lower, which may have been a result of the relatively narrow range of

CheSL Shannon diversity and CheSL Shannon evenness tested for FSA.

We evaluated the effects on microbial richness when FSA was cultured in additional

media that varied across a broader range of CheSL Shannon Diversity and CheSL Evenness

(Table 2). Unfortunately, FSB samples were no longer available for further testing. Medium B

was similar to NS, but also had low levels of soluble starch. ILC included a mixture of

carbohydrates inferred to be present in IL present at low concentrations (39, 40). The remaining

four media had a single carbohydrate polymer or its monomers present at higher levels, but this

carbohydrate varied in complexity. GP medium had higher concentrations of glucose

polysaccharide (starch), GM medium had higher concentrations of glucose monosaccharide, AP

medium had higher concentrations of arabinogalactan polysaccharide, and AM medium had high

concentrations of the arabinogalactan monomers galactose and arabinose. We cultured replicate

communities from FSA in these different media in continuous-flow MBRAs and analyzed

changes in microbial community composition through sequencing of the V4 region of the 16S

rRNA gene.

Adding this new data to our existing FSA data, we observed higher correlations between

ASV richness and CheSL richness (Fig. 3D; R=0.71, p=4.4X10^{-6}), CheSL Shannon evenness

(Fig. 3E, R=0.63, p=0.00011), and CheSL Shannon diversity (Fig. 3F, R=0.8, p=3.5X10^{-8})

across FSA samples. Analysis of FSA samples from this second experiment alone yielded higher

correlations (Fig. S1G-S1I), more consistent with what was observed for FSB cultured across a

range of media (Fig. S1A-S1C). As the two FSA experiments were performed on a

cryopreserved fecal sample stored for years between experiments, the high correlation across the

two experiments indicated the robustness of the observations.

We also evaluated two previously published data sets examining effects of carbohydrate

diversity on fecal microbial community assembly (26, 27). For these data sets, carbohydrate

linkage information was estimated from the available literature (Supplementary Table S2), and

microbial richness data was obtained from the text and/or supplementary information. (26–28,

41–43). We observed similar correlations between microbiota richness and CheSL richness,

CheSL Shannon evenness, and CheSL Shannon diversity (Fig. S2), across these studies, although

substrate complexity and the needed approximations made for certain polysaccharides used (i.e.,

pectin), may have reduced statistical significance of data analyzed from Chung et al (26).

Overall, these analyses demonstrated that CheSL Shannon diversity can be broadly applied, but

also highlighted limitations for use with certain fecal communities and carbohydrate structures.

CheSL Shannon diversity predicts peptide utilization

We used PICRUSt2 to infer genome content from 16S rRNA gene sequences and STAMP to

identify differentially abundant functional pathways between communities cultured in different

media. We focused on data from FSA cultured in the media described in Table 2 because we also

had access to cell-free culture supernatants for further testing. One observation of interest, given

our recent studies indicating Stickland fermentation contributes to persistence of *C. difficile* in

fecal communities cultured in NS medium (44), was enrichment for taxa potentially encoding

genes for L-arginine degradation (Stickland Fermentation; Fig. 4A) in microbial communities

with higher shared richness (cultured in B, GM, AP, and ILC). We also observed predicted

pathways for fermentation of pyruvate to butanoate (butyrate; Fig. 4B) were enriched in these

same microbial communities. We hypothesized that differential abundance of Stickland

fermentation pathways could indicate differences in peptide fermentation between communities.

We found that peptide utilization varied between communities cultured in different types of

media (Fig. 4C). When we performed a correlation between peptide utilization and CheSL

Shannon diversity, we observed a strong correlation (R = -0.77, P = 9.5×10^{-6}, Fig. 4D), similar

to that observed between peptide utilization and microbiota richness alone (Fig. S3A) and higher

than that observed for CheSL Shannon evenness (Fig. S3B). From this data, we infer that for

communities cultured in the presence of more complex mixtures of carbohydrates, peptide

utilization may serve as an additional substrate niche dependent on community composition;

alternatively, these communities may have higher needs for peptide utilization to facilitate

carbohydrate degradation capabilities.

Bacteroides and _Lachnospiraceae_ are central members in conserved, rich communities

As microbial interactions can have significant impacts on microbial community

assembly, stability and function (25), we investigated how variations in carbohydrate

composition affected microbial interactions. We used LIMITS, a computational algorithm that

infers microbial interactions by fitting a discrete Lotka-Volterra model against the temporal

profiles of ASV relative abundance data (45, 46), to analyze our sequence data. The LIMITS

equation was derived by implicitly assuming that total community biomass does not significantly

vary in time. Thus, performance is improved when LIMITS is conducted with data collected

from communities with minimal variations of the total biomass, such as those found when

culturing microbial communities under continuous-flow conditions approaching steady state.

However, LIMITS ignores temporal variations of interspecies interactions by assuming

interaction coefficients to be constant in time. While relevant in certain cases, this assumption is

invalid in general (47, 48). To address this limitation, we employed a moving window approach

by: (1) dividing the data into multiple subsets, each representing a shorter time period; (2) using

LIMITS to identify (constant) microbial interactions within these shorter periods; and (3)

iteratively inferring the network by shifting the time window to subsequent timepoints. By

analyzing the goodness of fit of these interaction networks (mean cross-correlation between

inferred networks and observed networks) across time windows, we observed that the mean

cross-correlation across media types increased after the early phase of _in vitro_ culture (Fig. S4).

We initially compared ensemble networks at the ASV level across replicate communities

cultured in the same media. However, very few interactions were conserved, which is consistent

with lower levels of ASV conservation observed between replicate communities. We posited that

neutral processes acting during initial community assembly likely contributed to variations in

ASV composition but that environmental filtering due to carbohydrate composition could select

for ASVs with similar functions. We assessed conserved interactions at the genus level, as

conserved functions have been previously observed to occur at the genus level in human fecal

communities (49), and identified interactions conserved between replicates within a medium.

Our analysis of the ensemble models primarily focused on FSA samples cultured in B,

AP, AM, GP, GM and ILC (Fig. 5), because the frequency of sample collection in this study

facilitated higher quality ensemble models. (Data from FSA and FSB in NS, MM, and IL can be

found in Fig. S5). We saw a distinct trend in the number of nodes, representing interacting

taxonomic groups, and the shared richness of the cultures. (Fig. 5A). These networks were not

driven solely by correlations between genera, but rather population dynamics fit to generalized

Lotka Volterra models by LIMITS (45), as was demonstrated by different network features

between media (e.g., AP, ILC, and B). However, it is important to note that not all taxonomic

groups were represented in the networks, as not all groups had conserved (or any) interactions

detected in some conditions.

We found three consistently central taxonomic groups, *Bacteroides*, *Lachnospiraceae*,

and *Ruminococcaceae* of unclassified genera (Fig. 5B). Particularly *Lachnospiraceae* and

Ruminococcacecae of unclassified genera exhibit conserved centrality in media with high CheSL

Shannon Diversity which produce complex ensemble networks. While the loss of

Lachnospiraceae spp. interactions in low CheSL Shannon Diversity networks could possibly be

explained by differences in abundance between media types (e.g., the negative $\log_2$ fold change

in GP compared to B), differences in abundance of *Ruminococcaeceae* spp. could not explain the

absence of interactions between communities cultured in different media, as these species were

not differentially enriched (Fig. 6). We also found greater connectivity in conserved networks

among media types which supported the highest microbiota richness. To support future uses of

ILC medium, we found communities cultured in this medium exhibited more conserved, positive

interactions between the three central taxonomic groups mentioned earlier (Fig. 5F). Overall, we

observed more conserved relationships across replicate cultures with higher CheSL Shannon

diversity, suggesting that this metric could be used to not only increase microbiota richness, but

also the reproducibility of microbial interactions in cultures derived from complex sources.

For media with low conservation of interactions between replicate networks (i.e., GP and

AP), we gained some insights into the effects of media on community composition through

differential enrichment analysis (Fig. 6). We observed that most taxa decreased in abundance in

GP compared to B (Fig 6), except for *Clostridium* spp. and *Bifidobacterium* spp (Supplementary

Table 3). For AM, we saw a similar selection against many genera, but enrichment of *Collinsella*

spp. (Fig 6). Based upon the ensemble network, this enrichment may be driven by *Bacteroides*

spp. (Fig. 5B, 6). Alternatively, enrichment may be due to the comparatively low ratio of

monosaccharides to polysaccharides, as was previously found in human fecal samples (50).

DISCUSSION

In this work, we developed novel metrics to describe carbohydrate complexity based on

Chemical Substrate and Linkage (CheSL) richness, CheSL Shannon diversity, and CheSL

Shannon evenness and demonstrated that mixtures of carbohydrates with higher levels of CheSL

Shannon diversity supported growth of human fecal microbial communities with higher richness

and conserved microbial interactions. This was contrary to our initial hypothesis that CheSL

richness would be the primary driver of microbial richness, and points to a role for CheSL

Shannon evenness (which also contributes to CheSL Shannon diversity) in microbial community

assembly. We hypothesize that CheSL Shannon evenness is most likely to contribute to

increased microbiota richness when carbohydrate mixtures are highly uneven. Specifically, when

species who occupy different carbohydrate niches compete for shared resources, generation of

higher levels of biomass by the species who occupy the carbohydrate niche with higher

concentrations may deplete other limiting nutrients that will lead to the extinction of other

species unable to compete for this limiting nutrient. Thus, these observations do not contradict,

but rather build upon nutrient-niche theory (23, 24) and integrate the concepts of competitive

fitness in the context of community stability and carbohydrate diversity.

This work builds on the body of emerging literature linking carbohydrate complexity to

microbial community composition and function. Similar to what was observed by Chung et al.

and Yao et al., media with higher levels of CheSL Shannon diversity support higher levels of

microbiota richness in communities cultured from fecal samples across the samples tested (26,

27). As Loss et al used a defined community of 10 microbes selected to provide representation of

microbiota diversity, they did not see effects of carbohydrate complexity on richness (29).

However, reanalysis of their data demonstrated correlations between microbiota diversity and

carbohydrate complexity (Fig. S6). We also observe correlation between CheSL Shannon

Diversity and microbial diversity (Fig. S6), but the correlations observed with microbiota

richness were of primary interest for this study. Of note, the CheSL Shannon Evenness metric

that we describe is similar to the carbohydrate complexity measure determined by Loss (29). Our

work expands on their studies through development of CheSL Shannon diversity, which for the

data in this study, exhibits stronger correlation with microbiota richness than CheSL Shannon

evenness.

Other studies have found more complex relationships between carbohydrate complexity

and microbiota richness that are likely due to differences in ability of microbial communities to

cleave linkages identified (28, 51, 52). Degradation of complex polysaccharides is dependent

upon the presence of specific carbohydrate active enzymes (CAZymes) within microbial

community members (53). In the absence of enzymes able to break unique linkages within

complex polysaccharides, monosaccharides remain inaccessible and cannot provide unique

niches. The presence of specific CAZymes which allow degradation of linkages are the gateway

to energy for the microbes, and in turn, linkages serve as a representation of the latent energy

granting niche space. Thus, it was not entirely surprising when we observed that when more

carbohydrate mixtures and microbial communities were analyzed, correlations between CheSL

Shannon diversity and microbiota richness were lower (Fig. S2). However, we suggest that

predictive accuracy of CheSL can be improved with prior knowledge of the CAZyme diversity

within the complex community being targeted. Inherent assumptions used in this study were that

the complex fecal microbiota would be capable of utilizing the selected carbohydrates as they are

abundant in diet or are host derived. One should be careful in use of CheSL for predictions of in

cases of low initial functional diversity or addition of uncommon substrates to communities

which likely lack the enzymatic potential for degradation (e.g., cellulose to human fecal sample

derived community). Further, the CheSL metrics should be applied across media which are

controlled with respect to other substrates. Finally, it is also important to note that CheSL

measures are estimates based upon published literature and will improve as additional

carbohydrate composition and linkage information is published

From our observations, among samples cultured in media with the greatest CheSL

Shannon diversity (B, AP, and ILC), there is higher similarity between microbial interaction

network structure and shared microbiota richness. This may suggest the diversity of nutrient

niches inferred by the abundance and evenness of linkages may control the reproducibility of

community assembly from complex microbial samples in continuous flow bioreactors, thereby

impacting the shared microbiota richness of the samples. Loss and colleagues similarly observed

that higher carbohydrate complexity led to higher reproducibility in community assembly (29).

From our findings, the initial assembly of a community around *Bacteroides*, *Lachnospriaceae,*

and *Ruminococcaceae* resulted in a network of interactions supporting reproducible cultures with

greater microbiota diversity under these culture conditions.

We also found enrichment and microbial associations previously observed in *in vivo*

systems exposed to similar treatments– another indicator of our *in vitro* system's capability to

support representative communities. Particularly the relationship between *Bacteroides* and

Lachnospiraeceae were supported through studies demonstrating cross feeding between

Bacteroides thetaiotaomicron and *Anaerostipes caccae* (54). *Clostridium sensu stricto* and

Bifidobacterium has previously been observed in humans consuming diets high in starch aligning

with our GP medium - which over-represents soluble starch for which these cultures continued to

support of these taxa despite significant reduction in diversity (31, 55, 56).

Communities formed in conditions with high CheSL Shannon Diversity also exhibited

enrichment of genes for potential Stickland fermentation of amino acids. Consistent with these

observations, we observed higher levels of peptide depletion in these media. Future studies will

be needed to determine whether utilization of peptides in media with high CheSL Shannon

Diversity is conserved across other microbial communities and media types. Typically,

proteolytic fermentation is associated with negative health outcomes, with a tradeoff between

saccharolytic and proteolytic microbes (57). In our studies, peptide consumption occurred in the

presence of dietary fibers and in communities enriched with genes for butyrate production,

suggesting that potential beneficial functions of the microbiota were also maintained.

Like any ecological set of interactions, interactions between GI microbiota are directly

determined by the biotic and abiotic properties of the environment to which the community is

exposed. Using approaches that adhere to limitations in data analysis and modeling as described

in this work, we gained additional resolution into the relationships between fecal bacteria and the

potential interaction landscapes which can be realized from the availability of specific substrates.

From progression in process-based approaches to interaction prediction, we further informed and

connected population dynamics with metabolic potential and ecological theories of nutrient niche

partitioning (23, 24). Using CheSL Shannon diversity as a guide when designing *in vitro* culture

media has the potential to improve reproducibility and increase the similarity of *in vitro*

communities to those found in the fecal samples without introducing unknown dietary

components that would be found in media supplements, such as ileostomy effluents or foods that

have undergone simulated digestion. Our initial test at an improved carbohydrate medium

supports this potential, with ILC granting a similar network structure to IL and the greatest

shared richness of media which had a known substrate concentration (IL being of complex

origin). By improving the reproducibility of community assembly in continuous flow conditions,

we can better detect differences between samples and reactors driven by other interventions.

Additionally, the experimental conditions we used, the improved quality of fit realized using a

continuous flow bioreactor, strict filtering, and division of our time series into windows which

represent transitory states in the community, all together may aid in inferring interactions for

future mechanistic investigation of microbial metabolic interactions.

MATERIALS AND METHODS

Samples used in this study

Fecal samples were previously collected from two adult volunteers who were self-described as

healthy and had not consumed antibiotics within the previous two months (30). Samples were

homogenized under anaerobic conditions, aliquoted into anaerobic tubes and stored at -80°C

until cultivation. Ileostomy effluents were collected from participants who had ostomies placed

due to previous illness or injury as previously described (58). Samples were stored at -20°C until

use. Informed consent was obtained prior to sample collection according to a protocol approved

by the Institutional Review Board of Michigan State University (10-736SM).

Media used in this study

Microbes from fecal samples were cultured in bioreactor basal medium that varied in

carbohydrate composition. 1 L of basal medium contained 1 g tryptone, 2 g proteose peptone #3,

2 g yeast extract, 0.4 g sodium chloride, 0.5 g bovine bile, 0.01 g magnesium sulfate, 0.01 g

calcium chloride, and 2 mL Tween 80, which was sterilized by autoclaving. 2 g sodium

bicarbonate, 0.2 g inulin from chicory, 0.1 g arabinogalactan from larch wood, 0.2 g soluble

starch, 0.04 g potassium phosphate dibasic, 0.04 g potassium phosphate monobasic, and 0.001 g

vitamin K3 were filter sterilized and added to each liter of basal medium after autoclaving.

Different variations of carbohydrates added to basal medium were used to test the impact of

carbohydrate composition and concentration on microbial communies (Table S1). NS (No

starch) lacked soluble starch and contained 0.15 g/L D-cellobiose, 0.15 g/L maltose, and 0.04

g/L D-glucose in addition to basal medium. MM (mucosal monosaccharides) contained an

additional 0.2 g/L of soluble starch, 0.05 g/L N-acetylglucosamine, 0.05 g/L N-acetylneuraminic

acid, 0.05 g/L D-glucuronic acid, and 0.05 g/L L-fucose. 5MM (5X higher mucosal

monosaccharides) contained an additional 1.8 g/L soluble starch (2g/L final concentration), 0.2

g/L N-acetylglucosamine, 0.2 g/L N-acetylneuraminic acid, 0.2 g/L D-glucuronic acid, and 0.2

g/L L-fucose (0.25 g/L final concentration each). GP (glucose polysaccharide) contained an

additional 1.8 g/L starch (2 g/L final concentration). GM (glucose monomer) contained an

additional 1.96 g/L D-glucose (2 g/L final concentration). AP (arabinogalactan polymer)

contained an additional 1.9 g/L of arabinogalactan (2 g/L final concentration). AM

(arabinogalactan monomers) lacked arabinogalactan and instead contained the two primary

monosaccharides present in arabinogalactan (59, 60)- 1.79 g/L galactose and 0.21g/L arabinose.

IL (ileostomy effluent) contained an additional 0.2 g/L soluble starch, 0.05 g/L N-

acetylglucosamine, 0.05 g/L N-acetylneuraminic acid, 0.05 g/L D-glucuronic acid, 0.05 g/L L-

fucose, and 100 mL of ileostomy effluent. Ileostomy effluent was thawed, pooled from multiple

donors, and filtered through a coffee filter to remove particulates prior to sterilization along with

base medium by autoclaving. Carbohydrate diversity of ileostomy effluent was estimated from

references (39, 40). Finally, ILC (ileostomy carbohydrates) reduced the concentrations of

arabinogalactan, inulin, and soluble starch to 0.05 g/L for each of these carbohydrates and also

contained 0.05 g/L cellobiose, 0.05g/L fructose, 0.05g/L galactose, 0.05 g/L N-acetylneuraminic

acid, 0.05 g/L N-acetylglucosamine, 0.05 g/L L-fucose, and 0.05 g/L D-glucuronic acid.

Culturing human GI microbes in continuous flow bioreactors

We used MBRAs (30), continuous-flow culture vessels that simulate environmental parameters

of the lumen of the distal human colon and maintain communities in steady-state based on the

carrying capacity of the reactors, to culture communities of human GI microbes from fecal

samples. MBRAs are parallelized, therefore, we could culture replicate communities of human

GI microbes in medium that varied by carbohydrate composition and concentration and

determine how these changes impacted microbial interactions. Prior to inoculation into MBRAs,

fecal samples were thawed and homogenized in anaerobic phosphate buffered saline (PBS) as a

20% w/v slurry as previously described (30). In the first experiment, Fecal Sample A (FSA) was

cultured in NS (n=2 replicates), MM (n=3 replicates), or IL (n=3 replicates) and Fecal sample B

(FSB) was cultured in NS (n=3 replicates), MM (n=3 replicates), 5MM (n=3 replicates), or IL

(n=3 replicates). Media contamination prevented cultivation of FSA in 5MM, whereas technical

difficulties led to loss of third replicate of FSA in NS. Both FSA and FSB formerly referred to as

Fecal Sample B and Fecal Sample C respectively, were previously characterized when cultured

in BRM2, a medium highly similar to NS, with an 8 hour retention time (30). For these

experiments, 16 hours after inoculation, continuous flow was initiated at 0.94 mL/hr (16 hr

retention time for 15 ml minibioreactor). One mL samples were collected from each culture daily

for 10 days as previously described (30).

In the second experiment, FSA was selected for studies because of fecal sample

availability. FSA was cultured in B, GP, GM, AP, AM, and ILC media, with n=4

replicates/medium. As before, continuous flow was initiated 16 hr after inoculation at 0.94

mL/hr. 1 mL samples were collected every 12 hours for 8 days. The increased frequency of

sampling and number of replicates was to improve our ability to subsequently infer microbial

interactions from data. In both experiments, MBRAs were maintained in an anaerobic chamber

(Coy Laboratories) heated to 37°C with a gas atmosphere of 90% N_2, 5% CO_2, and 5% H_2.

16S rRNA gene amplicon sequencing

The V4 region of the 16S rRNA gene was amplified directly from mechanically disrupted cells

with broad range 16S rRNA primers 515F/805R as previously described (30). Purified amplicons

were pooled in equimolar concentration and sequenced using Illumina MiSeq v2 according to

Manufacturer's protocol. Sequence data were processed using mothur V1.35.0. Sequences were

aligned using the Silva r132 16S rRNA gene sequence database (61). Any chimeric sequences

were removed following identification by UCHIME, and any contaminating sequences identified

as chloroplasts, mitochondria, Archaea or Eukarya are removed as well (62). Sequences were

denoised using mothur's pre.cluster function, giving ASVs, and taxonomy was assigned using

the Silva r138 16S rRNA gene sequence database. Downstream analysis of the taxonomic and

count data was handled in R, using the phyloseq package (63, 64); code will be published on

Github upon acceptance. An ASV table is provided in Supplementary Table S3, which is posted

at https://github.com/AuchtungLab/CheSL.

Data pre-processing for computational network inference

Since Lotka-Volterra equations model niche-based effects determining population dynamics,

multiple filtering steps were required to mitigate spurious interactions. During the early phases of

culture, two factors could influence community stability: the loss of ASV presence as non-viable

cells were removed by dilution and partnership switching as communities adapted to the culture

conditions. To improve LIMITS inferred model quality of fit, we tested whether a sliding

window approach, sampling time regions across the time series, could improve consistency of

522 interactions over time. We tested windows of varying sizes from 6 to 15 time points, across the

523 entire time series. To inhibit performance loss from transient ASVs, we filtered ASVs by

524 minimum presence across all time points within a window, requiring presence of a single read

525 from an ASV across all time points. This pre-processing avoided issues associated with taxa

526 falling and rising above the threshold of detection, which would influence the model similarly to

527 migration and extinction events, and disturb inferences of a fixed ecosystem's interactions (65).

528 From this filtering we obtained what we call a "shared microbiota richness" which represents the

529 count of ASVs found consistently across time, excluding ASVs that may be alternating above

530 and below threshold of detection.

531 **Inference of interspecies interactions using LIMITS**

532 LIMITS analysis was performed using the seqtime package in R by Faust et al. (46). Prior to

533 running LIMITS, we preprocessed the data as described above. From there, we applied the

534 LIMITS algorithm using a sliding window approach as described above. Inferred interactions

535 were then used to simulate time series for the given window of time and the resulting simulated

536 data were compared to the experimental data. Mean cross correlations were used to quantify the

537 quality of fit of the simulated data to the experimental data. Specifically, mean cross correlations

538 were calculated by determining the correlation between each taxon's simulated and experimental

539 data over the time series; the mean correlation of all taxa was then calculated; this method of

540 quality of fit analysis was also implemented through the seqtime package (46). This sliding

541 window method granted greater quality of fit than applying LIMITS to the entire experiment's

542 time series. In addition to improved quality of fit, a sliding window approach allowed us to select

543 the time windows which most likely represent interactions driven by deterministic effects, rather

than including data from the first day(s) of cultivation which likely included influences of

founder effects, partner selection, and adjustment to the bioreactor system.

From each replicate and window, a network was inferred using LIMITS. To identify

conserved interactions, we calculated consensus across networks from replicates of the same

fecal sample and media culture conditions. using the same fecal sample inoculum. After

calculating average interaction values and filtering by a consensus threshold of 2, meaning at

least 2 networks identified the interaction with the same directionality, we obtained ensemble

networks for each media type.

Characterization of chemical subunit and linkage diversity

As described in the results, we developed CheSL Richness, CheSL Shannon diversity and CheSL

Shannon evenness measures to calculate the complexity of carbohydrate mixtures based upon

distinct linkages within di- and polysaccharides or the presence of unique monosaccharides. For

example, the polysaccharide arabinogalactan is primarily composed of arabinose (80%) and

galactose (15%) monomers (59) that are linked in eight unique types linkages within the

polysaccharide (Table 1), with the relative abundances of these linkages are also known. We

additionally informed the ratios of our linkages for amylopectin from literature, which indicated

predominantly α-1–4 linkages, with 5% of linkages being α-1–6 linkages (66). For disaccharides,

which have a single linkage type, there is no effective change in diversity when applying CheSL

if it is the sole carbohydrate, but it is important to decompose into linkage types in cases where

there is overlap. For example, we find that maltose, with α-1–4 linkages, contributes to the same

pool as amylopectin (67). While cellobiose and inulin, have their own unique linkage pools -

β(1—4) glucose for cellobiose and β(2—1) fructose and α-(l—2) glucose for inulin (27, 68). We

estimated fructans linkage diversity from a study containing glycosidic linkage proportions from

wheat kernels, yielding six distinct linkage types (69). For estimations of linkage diversity for

CheSL calculations to apply to Chung et al.'s data, we referenced literature for pectin, beta-

glucan, and glucomannan respectively (41–43). Composition and linkage data not described here

can be found in Table 1 and in Supplementary Tables S1 and S2. Fig. 2 provides a representation

of how CheSL diversity can be calculated across a medium containing multiple carbohydrates.

Statistical analysis – comparison of means and correlations

To make comparisons for media shared richness, we used the R package MASS's

implementation of a negative binomial GLM, glm.nb, as the richness did not fit a normal

distribution determined by the Shapiro-Wilk test, and the data was over-dispersed for a Poisson

distribution (70). Pairwise comparisons were made with the R package emmeans and Tukey's

HSD was used to adjust for family-wise error rate (71). For correlations, the R Package ggpubr

was used alongside tidyverse and ggplot2 to visualize and calculate the Pearson correlation

between diversity metrics (72–74).

Inferred metabolic functional potential with PICRUSt

After characterization of conserved interactions and visualization using Cytoscape, we inferred

Enzyme Commission Pathway abundance for our cultures from fully sequenced isolates with

similar 16S rRNA gene sequences using PICRUSt2 (75). The potential significance of Pfam

pathway abundance was assessed using STAMP (76).

Measurement of amylose/amylopectin ratio

To measure the ratio of amylose/amylopectin for this study we use a NEOGEN Megazyme

amylose/amylopectin Assay kit, product code K-AMYL. The assay was performed according to

the manufacturer's instructions. This kit was used to determine the proportions of amylose and

amylopectin comprising the soluble starch used in various media formulation. This ratio was

then used to calculate CheSL diversity measures.

Differential analysis of microbial abundances

We performed differential abundance analysis with DESeq2 (77) between communities cultured

in basal bioreactor medium (B) and media with differing carbohydrate composition. The DESeq2

model we used to explain composition as a function of media type, with time as a covariate.

Additionally, we performed DESeq2 analysis between communities cultured in media composed

of the polymer form (starch (GP), arabinogalactan (AP)) and respective monomer forms (glucose

(GM), galactose and arabinose (AM)), as they represent potential differences between

monosaccharide-based niches and linkage-based niches at the genus level. Changes in abundance

that were identified by DESeq2 as significant and had an absolute fold-change in abundance

greater than two were plotted.

Acknowledgements

The authors thank Dr. Robert A. Britton (Baylor College of Medicine, Houston, Texas USA) for

access to sequence abundance data collected in his laboratory by JMA and shown in Fig. 3A-3C

as well as access to the FSA fecal sample used in follow up experiments.

H.C.M and J.M.A. were responsible for conceptualization and design of the studies. Data

collection was performed by H.C.M. and J.M.A. Analysis was performed by H.C.M. with

guidance provided by H.-S.S. and J.M.A. J.M.A. acquired funding for the studies. The initial

draft of the manuscript was written by H.C.M. and J.M.A. and all authors revised and edited the

final manuscript. Bioinformatics was done using the Holland Computing Center, supported by

the UNL Office of Research and Economic Development and the Nebraska Research Initiative.

Data Availability

16S rRNA gene data is available in the NCBI Sequence Read Archive under BioProject ID

PRJNA1146550.

Ethics Approval

Protocols for collection and use of fecal samples were reviewed and approved by Institutional

Review Boards at Michigan State University (10-736SM) and University of Nebraska-Lincoln

(protocol number 18585).

Funding

H.C.M. was supported by a graduate fellowship from the Nebraska Food for Health Center. This

work was partially supported by NSF grant OIA-2044049 to JMA, and by Nebraska Tobacco

Settlement Biomedical Research Enhancement Funds to H.-S.S and J.M.A. The content is solely

the responsibility of the authors and does not necessarily represent the official views of the

funders, who had no role in study design, data collection and interpretation, or the decision to

submit the work for publication.

Conflicts of Interest

J.M.A. has a significant financial interest in Synbiotic Health. There are no other conflicts of

interest to declare.

REFERENCES

1. Crovesy L, Ostrowski M, Ferreira DMTP, Rosado EL, Soares-Mota M. 2017. Effect of

Lactobacillus on body weight and body fat in overweight subjects: A systematic review of

randomized controlled clinical trials. Int J Obes 41:1607–1614.

2. Kok CR, Quintero DFG, Niyirora C, Rose D, Li A, Hutkins R. 2019. An in vitro

enrichment strategy for formulating synergistic synbiotics. Appl Environ Microbiol 85:1–

15.

3. Tofalo R, Cocchi S, Suzzi G. 2019. Polyamines and gut microbiota. Front Nutr 6:1–5.

4. O'Toole PW, Cooney JC. 2008. Probiotic Bacteria Influence the Composition and Function

of the Intestinal Microbiota. Interdiscip Perspect Infect Dis 2008:1–9.

5. Harcombe WR, Betts A, Shapiro JW, Marx CJ. 2016. Adding biotic complexity alters the

metabolic benefits of mutualism. Evol Int J Org Evol 70:1871–1881.

6. Litvak Y, Mon KKZ, Nguyen H, Chanthavixay G, Liou M, Velazquez EM, Kutter L,

Alcantara MA, Byndloss MX, Tiffany CR, Walker GT, Faber F, Zhu Y, Bronner DN,

Byndloss AJ, Tsolis RM, Zhou H, Bäumler AJ. 2019. Commensal Enterobacteriaceae

Protect against Salmonella Colonization through Oxygen Competition. Cell Host Microbe

25:128-139.e5.

7. Simon JC, Marchesi JR, Mougel C, Selosse MA. 2019. Host-microbiota interactions: From

holobiont theory to analysis. Microbiome 7:1–5.

8.   Van Herreweghen F, De Paepe K, Marzorati M, Van de Wiele T. 2020. Mucin as a

Functional Niche Is a More Important Driver of In Vitro Gut Microbiota Composition and

Functionality than Akkermansia muciniphila Supplementation. Appl Environ Microbiol

AEM.02647-20.

9.   Prosky L, Hoebregs H. 1999. Nutritional and Health Benefits of Inulin and Oligofructose

Methods to Determine Food Inulin and Oligofructose 1. J Nutr 129:1418–1423.

10.   Ahmadi S, Mainali R, Nagpal R, Sheikh-Zeinoddin M, Soleimanian-Zad S, Wang S, Deep

G, Kumar Mishra S, Yadav H. 2017. Dietary Polysaccharides in the Amelioration of Gut

Microbiome Dysbiosis and Metabolic Diseases. Obes Control Ther Open Access

4:10.15226/2374-8354/4/2/00140.

11.   Chleilat F, Klancic T, Ma K, Schick A, Nettleton JE, Reimer RA. 2020. Human Milk

Oligosaccharide Supplementation Affects Intestinal Barrier Function and Microbial

Composition in the Gastrointestinal Tract of Young Sprague Dawley Rats. Nutrients

12:1532.

12.   Lee NK, Kim WS, Paik HD. 2019. Bacillus strains as human probiotics: characterization,

safety, microbiome, and probiotic carrier. Food Sci Biotechnol 28:1297–1305.

13.   Seekatz AM, Theriot CM, Molloy CT, Wozniak KL, Bergin IL, Young VB. 2015. Fecal

microbiota transplantation eliminates Clostridium difficile in a murine model of relapsing

disease. Infect Immun 83:3838–3846.

14.   Krumbeck JA, Rasmussen HE, Hutkins RW, Clarke J, Shawron K, Keshavarzian A, Walter

J. 2018. Probiotic Bifidobacterium strains and galactooligosaccharides improve intestinal

barrier function in obese adults but show no synergism when used together as synbiotics.

Microbiome 6:121.

15. Friedman J, Alm EJ. 2012. Inferring Correlation Networks from Genomic Survey Data.

PLoS Comput Biol 8:1–11.

16. Yan S, Shi R, Li L, Ma S, Zhang H, Ye J, Wang J, Pan J, Wang Q, Jin X, Liu X, Liu Z.

2019. Mannan Oligosaccharide Suppresses Lipid Accumulation and Appetite in Western-

Diet-Induced Obese Mice Via Reshaping Gut Microbiome and Enhancing Short-Chain

Fatty Acids Production. Mol Nutr Food Res 63:1–13.

17. Sonnenburg ED, Sonnenburg JL. 2014. Starving our microbial self: The deleterious

consequences of a diet deficient in microbiota-accessible carbohydrates. Cell Metab

20:779–786.

18. Wu GD, Chen J, Hoffmann C, Bittinger K, Chen YY, Keilbaugh SA, Bewtra M, Knights D,

Walters WA, Knight R, Sinha R, Gilroy E, Gupta K, Baldassano R, Nessel L, Li H,

Bushman FD, Lewis JD. 2011. Linking long-term dietary patterns with gut microbial

enterotypes. Science 334:105–108.

19. Makki K, Deehan EC, Walter J, Bäckhed F. 2018. The Impact of Dietary Fiber on Gut

Microbiota in Host Health and Disease. Cell Host Microbe 23:705–715.

20. Van Hul M, Cani PD. 2019. Targeting Carbohydrates and Polyphenols for a Healthy

Microbiome and Healthy Weight. Curr Nutr Rep 8:307–316.

21. Hryckowian AJ, Van Treuren W, Smits SA, Davis NM, Gardner JO, Bouley DM,

Sonnenburg JL. 2018. Microbiota-Accessible carbohydrates suppress Clostridium difficile

infection in a murine model. Nat Microbiol 3:662–669.

22. Chen J, Vitetta L. 2020. The Role of Butyrate in Attenuating Pathobiont-Induced

Hyperinflammation 20:1–18.

23. Freter R, Brickner H, Botney M, Cleven D, Aranki A. 1983. Mechanisms that control

bacterial populations in continuous-flow culture models of mouse large intestinal flora.

Infect Immun 39:676–685.

24. Freter R, Brickner H, Fekete J, Vickerman MM, Carey KE. 1983. Survival and

implantation of Escherichia coli in the intestinal tract. Infect Immun 39:686–703.

25. Pereira FC, Berry D. 2017. Microbial nutrient niches in the gut. Environ Microbiol

19:1366–1378.

26. Chung WSF, Walker AW, Vermeiren J, Sheridan PO, Bosscher D, Garcia-Campayo V,

Parkhill J, Flint HJ, Duncan SH. 2018. Impact of carbohydrate substrate complexity on the

diversity of the human colonic microbiota. FEMS Microbiol Ecol 95:1–13.

27. Yao T, Chen MH, Lindemann SR. 2020. Structurally complex carbohydrates maintain

diversity in gut-derived microbial consortia under high dilution pressure. FEMS Microbiol

Ecol 96:1–16.

28.   Yao T, Deemer DG, Chen M-H, Reuhs BL, Hamaker BR, Lindemann SR. 2023.

706   Differences in fine arabinoxylan structures govern microbial selection and competition

707   among human gut microbiota. Carbohydr Polym 316:121039.

708   29.   Ostrem Loss E, Thompson J, Cheung PLK, Qian Y, Venturelli OS. 2023. Carbohydrate

709   complexity limits microbial growth and reduces the sensitivity of human gut communities

710   to perturbations. Nat Ecol Evol 7:127–142.

711   30.   Auchtung JM, Robinson CD, Britton RA. 2015. Cultivation of stable, reproducible

712   microbial communities from different fecal donors using minibioreactor arrays (MBRAs).

713   Microbiome 3:1–15.

714   31.   Liu S, Ren F, Zhao L, Jiang L, Hao Y, Jin J, Zhang M, Guo H, Lei X, Sun E, Liu H. 2015.

715   Starch and starch hydrolysates are favorable carbon sources for Bifidobacteria in the human

716   gut. BMC Microbiol 15:1–9.

717   32.   Anderson KL, Salyers AA. 1989. Biochemical evidence that starch breakdown by

718   Bacteroides thetaiotaomicron involves outer membrane starch-binding sites and periplasmic

719   starch-degrading enzymes. J Bacteriol 171:3192–3198.

720   33.   Wang Y, Lapointe G. 2020. Arabinogalactan utilization by bifidobacterium longum subsp.

721   Longum ncc 2705 and bacteroides caccae atcc 43185 in monoculture and coculture.

722   Microorganisms 8:1–16.

723   34.   Gil R, Silva FJ, Peretó J, Moya A. 2004. Determination of the Core of a Minimal Bacterial

724   Gene Set. Microbiol Mol Biol Rev 68:518–537.

35.   Edirisinghe JN, Weisenhorn P, Conrad N, Xia F, Overbeek R, Stevens RL, Henry CS.

2016. Modeling central metabolism and energy biosynthesis across microbial life. BMC

Genomics 17:568.

36.   Morrissey EM, Mau RL, Schwartz E, Caporaso JG, Dijkstra P, Van Gestel N, Koch BJ, Liu

CM, Hayer M, McHugh TA, Marks JC, Price LB, Hungate BA. 2016. Phylogenetic

organization of bacterial activity. ISME J 10:2336–2340.

37.   Martiny AC, Treseder K, Pusch G. 2013. Phylogenetic conservatism of functional traits in

microorganisms. ISME J 7:830–838.

38.   Reimer LC, Sardà Carbasse J, Koblitz J, Ebeling C, Podstawka A, Overmann J. 2022. Bac

Dive in 2022: the knowledge base for standardized bacterial and archaeal data. Nucleic

Acids Res 50:D741–D746.

39.   Barrett JS, Gearry RB, Muir JG, Irving PM, Rose R, Rosella O, Haines ML, Shepherd SJ,

Gibson PR. 2010. Dietary poorly absorbed, short-chain carbohydrates increase delivery of

water and fermentable substrates to the proximal colon. Aliment Pharmacol Ther 31:874–

882.

40.   Berghouse L, Hori S, Hill M, Hudson M, Lennard-Jones JE, Rogers E. 1984. Comparison

between the bacterial and oligosaccharide content of ileostomy effluent in subjects taking

diets rich in refined or unrefined carbohydrate. Gut 25:1071–1077.

41.   Hu S, Wang J, Nie S, Wang Q, Xu X. 2022. Chain conformations and steady-shear

viscosity properties of pectic polysaccharides from apple and tomato. Food Chem X

14:100296.

42.   Chang S-C, Saldivar RK, Liang P-H, Hsieh YSY. 2021. Structures, Biosynthesis, and

Physiological Functions of (1,3;1,4)-β-d-Glucans. Cells 10:510.

43.   Abdel-raouf ME-S, Sayed A, Mostafa M. 2022. Application of Guar Gum and Its

Derivatives in Agriculture, p. 117–132. *In* Murthy, HN (ed.), Gums, Resins and Latexes of

Plant Origin: Chemistry, Biological Activities and Uses. Springer International Publishing,

Cham.

44.   Huang X, Johnson AE, Auchtung TA, McCullough HC, Haidacher SJ, Hoch KM, Horvath

TD, Auchtung JM. Clostridioides difficile colonization is not mediated by bile salts and

requires Stickland fermentation of proline in an in vitro model of infection.

45.   Fisher CK, Mehta P. 2014. Identifying keystone species in the human gut microbiome from

metagenomic timeseries using sparse linear regression. PLoS ONE 9:1–10.

46.   Faust K, Bauchinger F, Laroche B, de Buyl S, Lahti L, Washburne AD, Gonze D, Widder

S. 2018. Signatures of ecological processes in microbial community time series.

Microbiome 6:1–13.

47.   Song H-S, Lee N-R, Kessell AK, McCullough HC, Park S-Y, Zhou K, Lee D-Y. 2024.

Kinetics-based inference of environment-dependent microbial interactions and their

dynamic variation. mSystems 9:e01305-23.

48.   Song H-S, Lee J-Y, Haruta S, Nelson WC, Lee D-Y, Lindemann SR, Fredrickson JK,

Bernstein HC. 2019. Minimal Interspecies Interaction Adjustment (MIIA): Inference of

Neighbor-Dependent Interactions in Microbial Communities. Front Microbiol 10:1264.

49.   Zhang J, Guo Z, Xue Z, Sun Z, Zhang M, Wang L, Wang G, Wang F, Xu J, Cao H, Xu H,

Lv Q, Zhong Z, Chen Y, Qimuge S, Menghe B, Zheng Y, Zhao L, Chen W, Zhang H. 2015.

A phylo-functional core of gut microbiota in healthy young Chinese cohorts across

lifestyles, geography and ethnicities. ISME J 9:1979–1990.

50.   Gomez-Arango LF, Barrett HL, Wilkinson SA, Callaway LK, McIntyre HD, Morrison M,

Dekker Nitert M. 2018. Low dietary fiber intake increases Collinsella abundance in the gut

microbiota of overweight and obese pregnant women. Gut Microbes 9:189–201.

51.   Cantu-Jungles TM, Bulut N, Chambry E, Ruthes A, Iacomini M, Keshavarzian A, Johnson

TA, Hamaker BR. 2021. Dietary Fiber Hierarchical Specificity: the Missing Link for

Predictable and Strong Shifts in Gut Bacterial Communities. mBio 12:e01028-21.

52.   Kok CR, Rose DJ, Cui J, Whisenhunt L, Hutkins R. 2024. Identification of carbohydrate

gene clusters obtained from in vitro fermentations as predictive biomarkers of prebiotic

responses. BMC Microbiol 24:183.

53.   Wardman JF, Bains RK, Rahfeld P, Withers SG. 2022. Carbohydrate-active enzymes

(CAZymes) in the gut microbiome. Nat Rev Microbiol 20:542–556.

54.   Chia LW, Mank M, Blijenberg B, Aalvink S, Bongers RS, Stahl B, Knol J, Belzer C. 2020.

Bacteroides thetaiotaomicron fosters the growth of butyrate-producing anaerostipes caccae

in the presence of lactose and total human milk carbohydrates. Microorganisms 8:1–13.

55.   Wang Y, Leong LEX, Keating RL, Kanno T, Abell GCJ, Mobegi FM, Choo JM,

Wesselingh SL, Mason AJ, Burr LD, Rogers GB. 2019. Opportunistic bacteria confer the

ability to ferment prebiotic starch in the adult cystic fibrosis gut. Gut Microbes 10:367–381.

56.   Jung DH, Seo DH, Kim YJ, Chung WH, Nam YD, Park CS. 2020. The presence of

788   resistant starch-degrading amylases in Bifidobacterium adolescentis of the human gut. Int J

789   Biol Macromol 161:389–397.

790   57.   Korpela K. 2018. Diet, Microbiota, and Metabolic Health: Trade-Off Between

791   Saccharolytic and Proteolytic Fermentation. Annu Rev Food Sci Technol 9:65–84.

792   58.   Collins J, Robinson C, Danhof H, Knetsch CW, Van Leeuwen HC, Lawley TD, Auchtung

793   JM, Britton RA. 2018. Dietary trehalose enhances virulence of epidemic Clostridium

794   difficile. Nature 553:291–294.

795   59.   Goellner EM, Utermoehlen J, Kramer R, Classen B. 2011. Structure of arabinogalactan

796   from Larix laricina and its reactivity with antibodies directed against type-II-

797   arabinogalactans. Carbohydr Polym 86:1739–1744.

798   60.   Chandrasekaran R, Janaswamy S. 2002. Morphology of Western larch arabinogalactan.

799   Carbohydr Res 337:2211–2222.

800   61.   Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA,

801   Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ,

802   Weber CF. 2009. Introducing mothur: Open-source, platform-independent, community-

803   supported software for describing and comparing microbial communities. Appl Environ

804   Microbiol 75:7537–7541.

805   62.   Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. 2011. UCHIME improves

806   sensitivity and speed of chimera detection 27:2194–2200.

63. McMurdie PJ, Holmes S. 2013. phyloseq: An R Package for Reproducible Interactive

Analysis and Graphics of Microbiome Census Data. PLoS ONE 8:e61217.

64. R Core Team. 2022. R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna, Austria.

65. Poos MS, Jackson DA. 2012. Addressing the removal of rare species in multivariate

bioassessments: The impact of methodological choices. Ecol Indic 18:82–90.

66. Ai Y, Jane J l. 2016. Starch: Structure, Property, and Determination, p. 165–174. *In*

Caballero, B, Finglas, PM, Toldrá, F (eds.), Encyclopedia of Food and Health. Academic

Press, Oxford.

67. Nawaz MA, Pervez S, Jamal M, Jan T, Khan W, Rauf A, Aman A, Qader SAU. 2019.

Maltose deterioration approach: Catalytic behavior optimization and stability profile of

maltase from Bacillus licheniformis KIBGE-IB4. Biotechnol Rep 24:e00400.

68. French AD. 2017. Glucose, not cellobiose, is the repeating unit of cellulose and why that is

important. Cellulose 24:4605–4609.

69. King ML, Xing X, Reintjes G, Klassen L, Low KE, Alexander TW, Waldner M, Patel TR,

Wade Abbott D. 2024. In vitro and ex vivo metabolism of chemically diverse fructans by

bovine rumen Bifidobacterium and Lactobacillus species. Anim Microbiome 6:50.

70. Venables W. 2002. Statistics complements to modern applied statistics with S Fourth

edition. No Title.

71. Lenth R. 2024. emmeans: Estimated Marginal Means, aka Least-Squares Means (1.10.5).

72. Kassambara A. 2023. ggpubr: "ggplot2" Based Publication Ready Plots (0.6.0).

73. Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, Grolemund G,

Hayes A, Henry L, Hester J, Kuhn M, Pedersen T, Miller E, Bache S, Müller K, Ooms J,

Robinson D, Seidel D, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K, Yutani H.

2019. Welcome to the Tidyverse. J Open Source Softw 4:1686.

74. Villanueva RAM, Chen ZJ. 2019. ggplot2: Elegant Graphics for Data Analysis (2nd ed.).

Meas Interdiscip Res Perspect 17:160–167.

75. Douglas GM, Maffei VJ, Zaneveld JR, Yurgel SN, Brown JR, Taylor CM, Huttenhower C,

Langille MGI. 2020. PICRUSt2 for prediction of metagenome functions. Nat Biotechnol

38:685–688.

76. Parks DH, Tyson GW, Hugenholtz P, Beiko RG. 2014. STAMP: Statistical analysis of

taxonomic and functional profiles. Bioinformatics 30:3123–3124.

77. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. Genome Biol 15:1–21.

Tables

Table 1: Linkages of Disaccharides and Polysaccharides tested. Linkages from the different

polysaccharides used in this study, with estimated linkage abundances from literature, and in the

case of starch, fractions of amylose and amylopectin determined as described in methods. For

linkages, t indicates terminal, 3,4,6, refers to which carbon the bond is on, f indicates it is in the

furanose form and p, the pyranose form.

Polymeric Carbohydrate	Linkage	Linkage Abundance	Monosaccharide	Reference
Arabinogalactan	Gal tp	0.257	Galactose	Goellner et al. (59)
	Gal 3p	0.049	Galactose	
	Gal 6p	0.258	Galactose	
	Gal 3,4p	0.01	Galactose	
	Gal 3,6p	0.289	Galactose	
	Ara tf	0.045	Arabinose	
	Ara tp	0.048	Arabinose	
	Ara 3f	0.044	Arabinose	
Starch - Amylose (25%)	α-(l-4)	0.25	Glucose	Ai and Jane (66)
Starch - Amylopectin (75%)	α-(l-4)	0.7125	Glucose	Ai and Jane (66)
	α-(l-6)	0.0375	Glucose	
Inulin	β(2—1)	.862	Fructose	Yao et al. (27)
	α-(l-2)	.138	Glucose	
Cellobiose	β(1—4)	1	Glucose	French (68)
Maltose	α-(l-4)	1	Glucose	Nawaz et al. (67)
Fructans	Fru tf	0.005	Fructose	King et al. (69)
	Fru β(2—1)f	0.045	Fructose	
	Fru β(2—6)f	0.34	Fructose	
	Fru β(2—6,1)f	0.22	Fructose	
	Glc tp	0.05	Glucose	
	Glc (l-6)p	0.34	Glucose	

852   **Table 2: Variations in Bioreactor Medium**

Carbohydrate (g/L)	B[1]	ILC[1]	GP[1]	GM[1]	AP[1]	AM[1]
Arabinogalactan	0.1	0.05	0.1	0.1	2.0	-
Inulin	0.2	0.05	0.2	0.2	0.2	0.2
Soluble starch	0.2	0.05	2.0	0.2	0.2	0.2
Cellobiose	0.15	0.05	0.15	0.15	0.15	0.15
Maltose	0.15	-	0.15	0.15	0.15	0.15
Glucose	0.04	-	0.04	2.0	0.04	0.04
Fructose	-	0.05	-	-	-	-
Galactose	-	0.05	-	-	-	1.71
Arabinose	-	-	-	-	-	0.29
N-acetylneuraminic acid	-	0.05	-	-	-	-
N-acetylglucosamine	-	0.05	-	-	-	-
L-fucose	-	0.05	-	-	-	-
D-glucuronic acid	-	0.05	-	-	-	-
Total carbohydrate (g/L)	0.84	0.5	2.64	2.8	2.74	2.74

[1]Abbreviations for basal media variations: B: basal medium; ILC: ileostomy carbohydrates; GP:

glucose polymer (soluble starch); GM: glucose monomer; AP: arabinogalactan polymer; AM:

arabinogalactan monomers.

FIGURE LEGENDS

Figure 1: Graphical Abstract A conceptual overview of this study, centered on applicability of

CheSL metrics to different community culture traits. Empirical data is shown in dark green, post-

analysis traits are shown in light green, inferred characteristics are show in blue, and the

interpretations are shown in red.

Figure 2: Graphical representation of Carbohydrate Chemical Subunits and Linkages

(CheSL) metrics. Graphical representations of approaches to calculate microbial and CheSL

richness, Shannon evenness and Shannon diversity. In these examples, different microbes are

represented by different shapes and colors and different linkages are color-coded.

Figure 3: Effects of CheSL richness, CheSL Shannon evenness, and CheSL Shannon

diversity on microbiota richness. Correlations of shared microbiota richness (observed ASVs

shared across a time window as described in Methods) with (A) CheSL richness, (B) CheSL

Shannon evenness and (C) CheSL Shannon diversity for FSA (circles) and FSB (diamonds)

Correlations of shared microbiota richness with (D) CheSL richness, (E) CheSL Shannon

evenness, and (F) CheSL Shannon diversity for FSA communities cultured in additional media

along with those described in panels (A)-(C). In all plots, Pearson correlations coefficients and p-

values are reported. Correlation analysis for FSA and FSB alone in the media shown in panels

(A)-(C) and for FSA in the new media shown in panels (D)-(F) can be found in Supplementary

Figure S1.

Figure 4. Differentially abundant functional pathways suggest divergent metabolism and

production of beneficial compounds (A-B) STAMP generated plots of pathways differentially

abundant in FSA communities cultured in media with high CheSL Shannon diversity complexity

and low CheSL Shannon diversity. (C) Total peptide concentrations measured in steady state

culture of each media type exhibiting differences in peptide utilization by communities compared

to fresh media control (D) Correlation between CheSL Shannon diversity and the peptide

concentration in spent medium. Pearson correlation coefficient and p-value are reported.

Additional correlations can be found in Supplementary Figure S3.

Figure 5: Ensemble Networks of Conserved Interactions demonstrate conserved structure

of communities cultured in media that support high shared microbiota richness and

within-group similarity (A) Shared microbiota abundance measured across three late culture

windows for each FSA replicate culture (n=4), cultured in the media indicated. Unless otherwise

noted (ns, for non-significant) comparisons between medium were highly significant with an

adjusted p-value $< 5\text{x}10^{-5}$ from Tukey's HSD (B) Conserved ensemble networks were built for

each media from the networks inferred from the window with the highest quality of fit for each

replicate cultured in AM, GM, AP, B and ILC as marked below each network. A key describing

taxa for each numbered node is provided. Positive interactions are indicated by red lines and

negative interactions are indicated by blue lines. An ensemble network for GP could not be

determined due to lack of conserved interactions.

Figure 6: Differential Abundance of Taxa Between FSA Communities Cultured in

Different Media ASVs were binned into shared taxa at the genus level (or lowest level of

taxonomic resolution available for taxa whose classification could not be determined at the genus

level). Differential abundance of taxa between communities cultured in different media types

was determined using DESeq2. Color indicates the $\log_2$-fold change in abundance between

cultures and the size of the points indicate the statistical significance of the abundance

difference.   Unless otherwise indicated, media were compared to medium B as baseline.

Comparisons were also made between communities cultured in overrepresented polysaccharide

media (GP and AP) and their respective overrepresented monosaccharide media (GM and AM).

SUPPLEMENTAL MATERIAL

Supplementary File 1: Figures S1 to S6

Figure Legends:

Supplementary Figure S1. Effects of CheSL richness, CheSL Shannon evenness, and

CheSL Shannon diversity on microbiota richness within fecal sample communities and

between experiments. Correlations between shared ASV richness and (A,D,G) CheSL

Richness, (B,E,H) CheSL Shannon evenness, and (C,F,I) CheSL Shannon diversity are shown

FSB (A-C) and FSA (D-F) cultured in NS, IL, MM and 5MM (A-C only), and for FSA cultured

in B, AM, AP, GM, GP, and ILC (G-I). Pearson correlation coefficients and p-values are

reported.

Supplementary Figure S2. Effects of CheSL richness, CheSL Shannon evenness, and

CheSL Shannon diversity on microbiota richness from previously collected data. (A-C)

Correlations between shared OTU Richness and (A) CheSL richness, (B) CheSL Shannon

evenness, and (C) CheSL Shannon diversity from data reported by Yao and colleagues (27). (D-

F) Correlations between shared OTU richness on days 15, 18, and 20 for Donor 1 (calculated

from data published as Supplementary Table S1) and (D) CheSL richness, (E) CheSL Shannon

evenness, and (F) CheSL Shannon diversity from data reported by Chung and colleagues (26).

Supplementary Table S2 provide details for CheSL calculations. Pearson correlation coefficients

and p-values are reported.

Supplementary Figure S3. Reduction in peptide levels strongly correlates with microbiota

richness. As described in Figure 4, we measured peptide levels in supernatants from stable FSA

communities cultured in AM, AP, GM, GP, B, and ILC (n=4/medium). Here we report

correlations with (A) shared ASV richness and (B) CheSL Shannon evenness. Pearson

correlation coefficients and p-values are reported.

Supplementary Figure S4. Shared richness and mean cross correlation values of FSA

communities cultured in AM, AP, B, GM, GP, and ILC across different time points in

culture. As described in Methods, we calculated (A) shared richness and (B) mean cross

correlation values for three groupings of time windows during early culture (including the first 3

timepoints A-B), mid (excluding the first three and last three timepoints C-D), and late

(excluding the first six timepoints E-F) to identify best time windows for building conserved

ensemble networks.

Supplementary Figure S5. Conserved ensemble networks for FSA and FSB cultured in NS,

MM, and IL. Conserved ensemble networks were built for each media/fecal sample

combination from the networks inferred from the window with the highest quality of fit for each

replicate. A key describing taxa for each numbered node is provided. Positive interactions are

indicated by red lines and negative interactions are indicated by blue lines. (A) Networks for

FSA communities grown in NS, MM, or IL as marked below each network. (B) Networks for

FSB communities grown in NS and IL as marked below each network. Ensemble networks for

FSB grown in MM and 5MM could not be determined due to lack of conserved interactions.

Supplementary Figure S6. Correlation between carbohydrate diversity and Inverse

Simpson diversity for data described by Loss and colleagues. (A) Fractional data for microbe

abundance provided in previously published Supplementary Table 2 was converted to whole

numbers and used to calculate Inverse Simpson microbial diversity with mothur v.1.48. Inverse

Simpson microbial diversity values were plotted as a function of the carbohydrate diversity

measure reported by Loss. Pearson correlation coefficient and p-value are reported. Panels (B)

and (C) repeat data from Figure S1 for reference and show correlations between ASV abundance

for FSA communities cultured in the indicated media and (B) CheSL Shannon evennes and (C)

CheSL Shannon diversity.

Supplementary File 2: Supplementary Tables S1 and S2

Supplementary Table S1: CheSL Richness, CheSL Shannon Evenness, and CheSL

Shannon Diversity calculations for media used in this study.

Supplementary Table S2: CheSL Richness, CheSL Shannon evenness, and CheSL Shannon

diversity calculations for analysis of previously published media described in

Supplementary Figure S2.

Supplementary File 3: Supplementary Table 3: Table of ASVs generated and analyzed in

this study.

Figure 1

Diversity

Richness: 3
Evenness: 0.657
Diversity: 0.721

Richness: 4
Evenness: 1.0
Diversity: 1.39

CheSL Diversity

Richness: 5
Evenness: 0.982
Diversity: 1.58

Evenness: 0.97
Diversity: 2.03

Medium m_1

Medium m_2

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6