Lessons learned while evaluating how pairing aversive experiences with same or different sex social stimulus mice affects subsequent social engagement

Abbreviated title: **Sex effects on social conditioning and behavior**

Jasmin N. Beaver[a,b,c], Marissa M. Nicodemus[a,b], Isabella R. Spalding[a], Lauren R. Scrimshaw[a,b], Sohini Dutta[b,d], Aaron M. Jasnow[e], Lee M. Gilman[a,b,c]*

[a]Department of Psychological Sciences, Kent State University, Kent, OH, USA 44242
[b]Brain Health Research Institute, Kent State University, Kent, OH, USA 44242
[c]Healthy Communities Research Institute, Kent State University, Kent, OH, USA 44242
[d]School of Biomedical Sciences, Kent State University, Kent, OH, USA 44242
[e]Department of Pharmacology, Physiology, and Neuroscience, University of South Carolina School of Medicine, Columbia, SC, USA 29209

Author contributions: JNB, SD, AMJ, LMG designed research. JNB, MMN, IRS, LRS, and LMG performed research. LMG contributed analytic tools. JNB and LMG analyzed data and wrote the manuscript. All authors reviewed and approved the manuscript.

Correspondence should be addressed to:
  Lee M. Gilman, Ph.D.
  lgilman1@kent.edu
  600 Hilltop Dr.
  209 Kent Hall
  Kent State University
  Kent, OH, USA 44242

Acknowledgements: We gratefully acknowledge the mice used in this study, the unrivaled veterinary care by Stan Dannemiller, DVM, MS, DACLAM, and the dedicated work of our vivarium caretakers. Figure 1 was created with BioRender.com (Toronto, ON).

Conflict of Interest: The authors declare no competing financial interests.

Funding sources: This work was supported by R15 MH118705 to AMJ and LMG, and by Kent State University.

Abstract

Mice offer a wealth of opportunities for investigating brain circuits regulating multiple behaviors, largely due to their genetic tractability. Social behaviors are translationally relevant, considering both mice and humans are highly social mammals, and human social behavior disruptions are key symptoms of myriad neuropsychiatric disorders. Stresses related to social experiences are particularly influential in the severity and maintenance of neuropsychiatric disorders like anxiety disorders, and trauma and stressor-related disorders. Yet, induction and study of social stress in mice has disproportionately focused on males, influenced heavily by their inherent territorial nature. Social target-instigated stress (i.e., defeat), while ethologically relevant, is quite variable and predominantly specific to males, making rigorous and sex-inclusive studies challenging. In pursuit of a controllable, consistent, high throughput, and sex-inclusive method for social stress elicitation, we modified a paradigm to train male and female F1 129S1/SvlmJ × C57BL/6J mice to associate (via classical conditioning) same or different sex C57BL/6J targets with a mild, aversive stimulus. While further paradigm optimization is required, social interaction testing 24 h after conditioning indicates males socially conditioned better to male targets by exhibiting reduced social interaction, whereas females appeared not to form social stimulus associations. Serum corticosterone levels inversely corresponded to social avoidance after different sex, but not same sex, conditioning, suggesting corticosterone-mediated arousal influences cross-sex interactions. These rigorously controlled null outcomes align with past pursuits' limited success in creating a sex-inclusive social stress paradigm.

Significance Statement

Validated paradigms to study social stress in female mice, and across sexes, are needed. We modified a published male mouse protocol by using classical conditioning to pair an aversive stressor with a target. Our goal was to create a uniform, cross-sex, high-throughput social stress technique to advance future research. Though our modified paradigm requires future improvements, we did acquire evidence that males can be socially conditioned in this way, and female same sex social engagement can be attenuated by a preceding non-social aversive experience. These null findings, while not achieving our goal, provide useful information to advance future sex-inclusive social stress investigations.

Introduction

Social interaction behavior is a cross-mammalian phenomenon seen in humans, non-human primates, rats, hamsters, and mice, among other animals (Bloomberg et al., 1994; Jasnow and Huhman, 2001; Kamps et al., 2001; Leung, 2011; Monfils and Agee, 2019; Lee et al., 2020). Social interaction behavior shifts can be informative of the physical and/or emotional state of the mammal (Ago et al., 2014; Wilson and Koenig, 2014; Venniro et al., 2018). In humans, typical social interaction behavior disruptions are characteristic of numerous neuropsychiatric conditions including trauma and stress-related disorders, mood and anxiety disorders, and autism spectrum disorder. (American Psychiatric Association, 2022).

Rodent studies examining social behavior and socially associated stressors within various contexts have advanced identification of regulating neural pathways, and assisted development of new therapeutic approaches (Jasnow and Huhman, 2001; Coccurello et al., 2009; Smith and Wang, 2012; Gilman et al., 2015; Caldwell et al., 2017; Bergamini et al., 2018; Donovan et al., 2020; Sahani et al., 2022; Venniro et al., 2022). However, social behavior research in the genetically tractable mouse species (*Mus musculus*) has been relatively limited in studying behavioral and neurophysiological consequences of aversive social interactions in *female* mice (Kondrakiewicz et al., 2019; Kuske and Trainor, 2021). This is largely due to capitalization of male-specific territorial aggression in most rodent social conditioning paradigms (Blanchard et al., 2001; Gilman et al., 2015; Latsko et al., 2016; Peña et al., 2019; Lopez and Bagot, 2021; Furman et al., 2022; Lyons et al., 2023; Pantoja-Urbán et al., 2024). A breadth of social stress techniques have been employed in hamsters, mice, and rats (Miyashita et al., 2006; McCormick et al., 2007; Bernberg et al., 2008; Kercmar et al., 2011; McCormick et al., 2013; Iñiguez et al., 2018; Sterley et al., 2018; Lee et al., 2020; Furman et al., 2022; Pan et al., 2023). Of these, the most prevalent paradigm in mice is that of social defeat (both acute and chronic) (Jasnow and

Huhman, 2001; Jasnow et al., 2015; Golden et al., 2011; Dulka et al., 2015; Gilman et al., 2015;

Latsko et al., 2016; Bonnefil et al., 2019; Wang et al., 2021). Aside from the male-centric nature

of social defeat, additional concerns of reproducibility arise from the inherently variable range of

aversive social experiences each 'defeated' experimental mouse encounters. Such variability,

mostly beyond experimenter control, plus the single sex bias of social defeat, left us wanting to

explore an improved paradigm. For this, we paired controllable, uniform, aversive

unconditioned stimuli with the presence of a social stimulus (conditioned stimulus; target) to

study aversive social conditioning *across sexes* in mice using a higher throughput approach

(requiring one conditioning day, rather than multiple days-weeks).

To accomplish this, we modified a paradigm previously utilized in male mice, involving manual

administration of an aversive stimulus (mild foot shock) selectively when a male mouse actively

investigated a target, with the goal of attenuating subsequent social engagement (Toth et al.,

2012; Zoicas et al., 2014; de la Zerda et al., 2022; Grossmann et al., 2024). Rather than

continuing this operant-style approach, where a mouse's behavior dictates an outcome, we

shifted to a classical conditioning approach in which shock delivery is standardized across

individual mice. We anticipated experimental mice would associate the presence of the target

with the aversive unconditioned stimulus. Additionally, all mice would receive the same number

of shocks, instead of variable numbers based upon behavior.

Here, we evaluated how this paradigm affected social engagement and fear behaviors *across*

sexes after mice were exposed to a novel target. Specifically, mice were socially conditioned

with an aversive stimulus (mild foot shock; stress) when in the presence of their assigned target,

independent of investigative behavior (Social Stimulus + Stress group). Then, mice were tested

for social engagement and fear behavior in a separate environment with their assigned target

present, followed by testing of freezing in the conditioning environment in the absence of any

target. The former test enabled evaluation of both social and fear behaviors, while the latter test

assessed contextual memory sans social stimuli. These behaviors were tested under

circumstances when the experimental and target mice were the same sex, and when they were

different sexes. We hypothesized Social Stimulus + Stress mice would exhibit reduced social

engagement (indexed as time spent directly adjacent to the target enclosure) when tested with

their assigned target in a novel environment, regardless of if the Social Stimulus + Stress and

target mice were the same or different sexes. Our broad goal was to develop a paradigm that

would facilitate rigorous future investigations of social behavior across mouse sexes, which are

currently underrepresented in literature (Kuske and Trainor, 2021).

Methods

Mice

Hybrid F1 offspring (i.e., 129SB6F1/J) of both sexes resulting from pairing female 129S1/SvlmJ

(RRID:IMSR_JAX:002448) and male C57BL/6J (RRID:IMSR_JAX:000664) mice (hereafter

experimental mice), and male and female C57BL/6J mice (hereafter targets), all ≥9 weeks old or

older, were group-housed (two to five per cage) within sex. Experimental mice were hybrid F1

offspring of 129S1/SvlmJ and C57BL/6J mice because these two strains are the most prevalent

among transgenic mice, and because hybrids provide broader generalizability (Kumar et al.,

2021; Keady et al., 2022). Such hybrids are used for a variety of physiology and neuroscience

research purposes (e.g., (Curley et al., 2010; Kelley et al., 2011; Agrimson et al., 2017; Niibori

et al., 2020; Correia et al., 2024)). Targets were C57BL/6J mice housed in a separate room

from experimental mice. C57BL/6J mice were used as targets to ensure they were entirely

novel to experimental mice through separate housing; because C57BL/6Js are often used as

the primary or background strain for testing social behaviors in mice (see reviews (Laman-

Maharg and Trainor, 2017; Varholick et al., 2021; Takahashi, 2025)); and because our F1

breeding was limited to generation of experimental mice. All mice had *ad libitum* access to food

and water in rooms maintained on a 12:12 light/dark cycle with lights on at 07:00 local standard

time (i.e., Zeitgeber time 0), and temperature maintained at 22 ± 2°C. All mice were fed LabDiet

5001 rodent laboratory chow (LabDiet, Brentwood, MO) and were kept on 7090 Teklad Sani-

chip bedding (Envigo, East Millstone, NJ) in cages containing Nestlets (Ancare, Bellmore, NY)

and huts (Bio-Serv, Flemington, NJ) for enrichment. Experiments were approved by Kent State

University's Institutional Animal Care and Use Committee, and adhered to the National

Research Council's Guide for the Care and Use of Laboratory Animals, 8th Ed. (National

Research Council, 2011).

Social Conditioning Paradigm

Timeline

The entire paradigm spans four consecutive days. Procedures for each individual day are

outlined below in order. On Days 0 and 1, experimental mice underwent different experimental

manipulations according to their specific treatment group. On Days 2 and 3, all experimental

mice experienced identical testing conditions regardless of their treatment group (Fig. 1).

Apparatus and Software

On Days 0, 1, and 3, experimental mice were placed in Coulbourn Instruments chambers (17.8

cm D × 17.8 cm W × 30.5 cm H; Allentown, PA). Chambers comprised two opposing aluminum

walls each adjoining two clear acrylic walls. Targets were put inside a transparent enclosure

placed within a corner of the social conditioning chamber. The transparent enclosure included

23 holes (each 0.32 cm diameter) on each of the two sides facing the experimental mouse to

enable olfactory cue exchanges. Previous work demonstrates that olfaction is the most

important modality when recognizing and interacting with novel same-species targets (see

review (Sterley and Bains, 2021)); tactile (whiskers) and auditory modalities appear to only be

recruited when recognizing cage mates (de la Zerda et al., 2022) in male mice (females were

not studied). Chambers and enclosures were cleaned with 70% ethanol before and after each

session; chambers had visible illumination, and one clear acrylic wall was marked with a blue

dotted pattern. These components contributed olfactory and visual cues to the context, in

addition to the tactile cue of the stainless steel grid floor for the experimental mouse. Cameras

mounted above the chambers were used to record movements, and freezing behavior (≥0.25 s)

on Days 0, 1, and 3 was quantified using FreezeFrame software (v. 5.201, Actimetrics,

Wilmette, IL). Freezing – ceasing all movement save breathing – is an innate behavior

expressed by mice in response to a real or perceived threat, enabling threat assessment while

minimizing detection (see review (Blanchard and Blanchard, 1988)). All videos were manually

reviewed to ensure the software satisfactorily distinguished between freezing behavior and

mere absence of locomotion.

On Day 2, experimental mice were placed in a novel room (distinct from the room used for Days

0, 1, and 3) containing open arenas (41.9 cm W × 41.9 cm D × 39.6 cm H; Coulbourn

Instruments). Each open arena had one empty PVC tube (8.9 cm outer diameter) in a single

corner of the square chamber with wire mesh (0.25 × 0.25 cm openings) covering a rectangular

portion (11.4 cm W × 4.1 cm H) cut out of the tube bottom. Because of the mesh at the bottom

of the PVC tubes containing targets, exchanges of visual, auditory, and olfactory cues between

experimental mice and targets were possible, but tactile interactions were minimized. Social

interaction behavior was quantified as post-test duration when at least 80% of an experimental

mouse's body was in the interaction zone, a 7 cm radius zone (~223 cm^2 – 240 cm^2, i.e., 14-

15% of the open arena) around the base of the PVC tube. Freezing behavior, regardless of

arena location, lasting at least 250 ms during the post-test was also measured, then converted

into a percentage of the post-test time. Social and freezing behaviors were detected using ANY-

maze software (v. 7.09 Stoelting Co., Wood Dale, IL).

Pre-exposure (Day 0)

Pre-exposure was used to help reduce the novelty of the transparent enclosure and/or the

target to the experimental mouse, plus minimize any potential sex differences in acquisition

(Siegfried et al., 1987; Wiltgen et al., 2001; Guzmán et al., 2009; Brown et al., 2011). On Day 0,

experimental mice were placed in chambers with (Shock Control and Social Stimulus + Stress

mice) or without (Social, Pre-Contingency, and Post-Contingency Controls) a target (age-

matched; either same sex or different sex relative to experimental mouse, depending on if

assigned to same sex or different sex experiment) for pre-exposure (Fig. 1). During pre-

exposure, experimental mice were allowed to explore the chamber and target enclosure for five

min, then both mice were returned to their respective home cages. No aversive stimulus was

presented.

Social Conditioning (Day 1)

On Day 1, experimental mice were placed in the same social conditioning chamber as Day 0

and received five, 1 s mild foot shocks (1.0 mA) in the presence of the same target that they

encountered on Day 0 (Fig. 1). This was to have experimental mice form associations between

aversive foot shocks and their respective target (Fig. 1). Conditioning lasted for nine min; after

a two min baseline, shocks were then administered at 120, 210, 300, 390, and 480 s. Targets

did not receive foot shocks.

Social Engagement Testing (Day 2)

On Day 2, experimental mice intentionally underwent social engagement testing in a novel

environment with a novel target enclosure to ensure sensitivity of our social interaction measure

while minimizing potential contextual confounds; in other words, to isolate effects on social

engagement from those of context fear learning (the latter assessed on Day 3). Mice

acclimated to the room in their home cage for 30 min prior to behavior testing commencing.

Experimental mice were placed in the corner of the arena opposite the PVC tube and could

investigate freely for 2.5 min (pre-test). Immediately after these 2.5 min, the target that each

experimental mouse had previously been conditioned with on Day 1 was then placed in the PVC

tube in the corner. Experimental mice were allowed to continue investigating the arena for an

additional five min (post-test; Fig. 1). Proprietary incompatibilities and video file encoding

precluded analyses of freezing behavior during social interaction testing from being evaluated

with FreezeFrame software; this is why ANY-maze software was used instead here. Post-test

social interaction time was log-transformed ($Y=log(Y+0.001)$ to account for three mice with zero

s post-test social interaction) so that data would be normally distributed for statistical analyses.

Females were intentionally not assessed for estrus cycle stage for five reasons: 1) to minimize

mouse usage (Russell and Burch, 1959), we did not power our studies for assessment of

estrus; 2) with the goal of developing a social stress paradigm, we are seeking effects robust

enough to not depend upon estrus in intact, randomly cycling females; 3) we intentionally

focused here on fear and social behaviors, and did not measure sexual behaviors (e.g.,

lordosis); 4) evidence that overall mouse behaviors are not affected by estrus stage

(Prendergast et al., 2014; Levy et al., 2023), (but see (Chari et al., 2020)); 5) cross-species

evidence indicates vaginal lavage to determine estrus cycle is stressful (Becegato et al., 2021;

Bahadır-Varol et al., 2022), and we sought to minimize stress confounds here.

Context Testing (Day 3)

On the last day of the paradigm, experimental mice were placed in the social conditioning

chamber for behavior testing in the *absence* of any target (Fig. 1). The target enclosure was still

present to keep the context consistent. All other tactile, visual, and olfactory cues from Days 0

and 1 were present. Testing lasted 10 min and did not involve any foot shocks.

Treatment Groups

The Social Stimulus + Stress group involved mice that underwent social conditioning in the

presence of a non-shocked target (either same sex or different sex relative to Social Stimulus +

Stress mouse, depending upon experiment), with the hypothesis that Social Stimulus + Stress

mice would associate the target with this aversive experience. We planned four Control groups

for this one Social Stimulus + Stress group; each experiment (same sex or different sex) had its

own respective set of four Control groups for its respective Social Stimulus + Stress group. The

goals of these were to control for: 1) foot shock exposure, i.e., Shock Controls; 2) presence of

target during pre-exposure & foot shock, i.e., Social Controls; 3) exposure to target temporally

distal to foot shock exposure, such that social encounters still occurred in a manner not

contingent with foot shock nor the conditioning context, i.e., Pre- and Post-Contingency Controls

(Fig. 1). Shock Controls experienced procedures identical to those of the Social Stimulus +

Stress group, except a foot shock was never administered on Day 1 (Fig. 1). Social Controls

were treated the same as the Social Stimulus + Stress group, except experimental mice never

encountered their assigned target until testing on Day 2 (Fig. 1). Pre- and Post-Contingency

Controls involved experimental mice encountering targets either four h before (Pre-Contingency

Controls) or four h after (Post-Contingency Controls) pre-exposure (Day 0) and social

conditioning (Day 1); in other words, targets were not present in the conditioning chamber

during these two periods. Instead, experimental mice were exposed to their target in a separate

room for five min or nine min (the same amount of time as pre-exposure or social conditioning,

respectively) four h before (Pre-Contingency Controls) or four h after (Post-Contingency

Controls) the experimental mice were exposed to the conditioning chamber. This exposure

involved experimental mice and their respective target being placed in a clean mouse cage

separated with an acrylic divider to allow for visual, auditory, and olfactory exchange, but

minimizing tactile interaction, mirroring the experiences of Social Stimulus + Stress mice and

Shock Controls within the social conditioning context on Days 0-1. Mice in Pre- or Post-

Contingency Control conditions still experienced foot shocks on Day 1, and encountered their

assigned target for testing on Day 2 (Fig. 1). This temporal separation of four h was to minimize

consolidation of social encounters from interfering/intermingling with consolidation of the

aversive foot shock encounter in the social conditioning chamber, while still making execution of

these experiments feasible within a 12 h lights-on period (Wanisch et al., 2008; Cai et al., 2016;

Lynch et al., 2017) (see reviews (Abel and Lattal, 2001; Sheppard et al., 2018)).

Figure 1. Study timeline and treatment groups. Five groups were used in this study for each experiment (same sex experiment and different sex experiment): one socially conditioned (Social Stimulus + Stress) group and four Control groups. Day 0 was for pre-exposure; Day 1 was when social conditioning occurred; Day 2 tested for social engagement; Day 3 tested for context-specific freezing behavior. On Day 0, Social Stimulus + Stress mice (A; above horizontal lines) encountered their assigned target in the social conditioning context, where olfactory, visual, and auditory cues could be exchanged while tactile interactions were minimized. No aversive stimulus was applied on Day 0, but on Day 1, experimental mice and their assigned target were returned to this context. Social Stimulus + Stress (but not target) mice then experienced five mild foot shocks to associate their target with that aversive experience. Social Stimulus + Stress mice were then tested for social engagement with that same target on Day 2, followed by testing for context fear behavior on Day 3 in the absence of their assigned target. Each Control group (below horizontal lines) was composed of different mice. Shock Control (B) mice underwent the same exact procedure as detailed for the Social Stimulus + Stress group, except no foot shock was administered on Day 1. Social Control (C) mice similarly experienced the same procedure as mice in the Social Stimulus + Stress group, save that Social Control mice did not encounter any target until Day 2 for testing of social engagement. Pre-Contingency Control (D) mice encountered their target four h before pre-exposure and social conditioning, meaning no target was present for Pre-Contingency Control mice when they were in the social conditioning chamber for pre-exposure or social conditioning. To encounter their assigned target, Pre-Contingency Control mice were placed into a clean cage with a clear acrylic divider separating them from their target. This divider allowed visual, olfactory, and auditory exchanges with the target but minimized tactile interactions, the same as for Social Stimulus + Stress mice and Shock Controls when in the social conditioning context. On Day 0, Pre-Contingency Control mice encountered their target for five min; on Day 1, for

nine min; these were timed to match the length of target exposure that mice in the Social

Stimulus + Stress group experienced. Post-Contingency Control (E) mice were treated the

same as Pre-Contingency Control mice, except the former's encounters with their assigned

targets occurred four h *after*, rather than before, pre-exposure and social conditioning.

Serum Corticosterone

Thirty min after all experimental mice underwent contextual fear testing on Day 3, they were

briefly anesthetized with isoflurane then rapidly decapitated for trunk blood collection. Blood

clotted at room temperature for 10 min, then was spun at 3500 rpm for one h at 4°C. Serum was

collected and stored at -80°C until corticosterone analyses could be performed. Serum

corticosterone was measured using Enzo Life Sciences corticosterone enzyme-linked

immunosorbent assay kits (Farmingdale, NY). Assays were run according to the manufacturer's

instructions using their small volume protocol. Plates were read at 405 nm with correction at 580

nm. The sensitivity of the assay was 26.99 pg/mL. After serum corticosterone levels were

interpolated from each plate's standard curve, they were log-transformed to account for the

typical skewness of serum corticosterone (Teilmann et al., 2014; Uarquin et al., 2016; Weber et

al., 2023).

Statistical Analyses

Data were graphed with GraphPad Prism 10.3.0 (442), Beta (GraphPad Software, San Diego,

CA), and analyzed using IBM SPSS Statistics 28.0.0.0 (IBM, Armonk, NY), with the significance

threshold set *a priori* at $p<0.05$. Non-significant trends ($p<0.10$) were mentioned only when the

associated partial η^2 was ≥0.060. Data were graphed as the mean (μ) ± standard error of the

mean. Details of identified outliers (greater than the mean ± four standard deviations (σ)) are

provided in the Supplemental Dataset. The same sex experiment and different sex experiment

each included their own Social Stimulus + Stress group plus accompanying four Control groups.

Experiments were analyzed separately. Social conditioning acquisition was analyzed using a

three-way repeated measures general linear model (time × treatment group × sex of

experimental mouse). Greenhouse Geisser corrections were utilized for within-subjects

analyses. Measurements of contextual fear expression average, log-transformed post-test

social interaction time, percent time freezing during social interaction testing, and log-

transformed corticosterone were each analyzed with a two-way general linear model (treatment

group × sex of experimental mouse). Our primary outcomes of interest for this study were

freezing during social conditioning acquisition, social engagement behavior, and freezing during

context testing. Accordingly, our *a priori* hypotheses and planned contrasts were:

- **Hypothesis 1**: Freezing during the final time point of social conditioning acquisition would be minimal in Shock Control mice compared to all other groups; and female experimental mice would exhibit higher levels of freezing than males.

- **Hypothesis 2**: In female more than male experimental mice, log-transformed social engagement would be decreased in Social Stimulus + Stress mice compared to all four Control groups.

- **Hypothesis 3**: Freezing during testing in the social conditioning chamber would be minimal in Shock Control mice compared to all other groups; and female experimental mice would exhibit higher levels of freezing than males.

These planned contrasts were selectively analyzed without correction alongside our omnibus

analyses. For our secondary outcomes that either were used to assess the possibility of

confounds (percent time freezing during social interaction testing) or were exploratory (log-

transformed serum corticosterone), and as follow-ups to our planned contrasts on our primary

outcomes, we performed *post hoc* pairwise comparisons with Bonferroni correction. The results

of these *post hoc* tests for our primary outcomes are reported in Supplemental Figures S1-S3.

Results

Social Conditioning Acquisition

Social conditioning acquisition is illustrated in Fig. 2A, 2B for mice in the same sex target

experiment, and in Fig. 2C, 2D for mice in the different sex target experiment. All mice receiving

a foot shock expressed >40% freezing during at least one post-shock period during social

conditioning.

For acquisition in the same sex experiment, no three-way interaction of time × sex × group

occurred (Table 1). We observed a significant two-way interaction between time × group, and a

non-significant trend for group × sex was noted (p=0.070; Table 1). A significant main effect of

sex was also found for mice in the same sex experiment (Table 1). Our *a priori* planned

contrast for **Hypothesis 1** within the same sex experiment ($F_{(1,69)}$=1.140 p=0.289, partial

η^2=0.016) did not support our anticipated observation (Fig. 2A, B). Graphs of these data

showing the outcomes of pairwise comparisons are located in Supplemental Figure S1A, B.

Table 1. Two-way repeated measures general linear model on learning of social conditioning

for mice of both sexes with same sex targets.

Same Sex Experiment – Social Conditioning Acquisition			
	F Statistic	**p value**	**partial η^2**
Time	$F_{(4.18,288)}$=135.9	<0.001	0.663
Group	$F_{(4,69)}$=45.09	<0.001	0.723

Sex	$F_{(1,69)}$=13.54	<0.001	0.164
Time × Group	$F_{(16.7,288)}$=9.518	<0.001	0.356
Time × Sex	$F_{(4.18,288)}$=1.714	0.144	0.024
Group × Sex	$F_{(4,69)}$=2.275	0.070	0.117
Time × Group × Sex	$F_{(16.7,288)}$=0.787	0.706	0.044

Social conditioning acquisition for mice in the different sex experiment is shown in Fig. 2C, 2D.

A significant three-way interaction of time × sex × group was observed (Table 2). Similar to the

same sex experiment, however, our planned contrast did not reach significance for **Hypothesis**

1 ($F_{(1,70)}$=0.537 p=0.466, partial η^2=0.008). Indeed, no significant pairwise comparisons

indicated within-group significant sex differences at the final social conditioning acquisition time

point, either for the same sex or different sex experiments (Supplemental Figure S1).

Combined with same sex experiment acquisition (Supplemental Figure S1), these findings

indicate that social exposure – whether concurrent with or temporally distal from an aversive

unconditioned stimulus – can have transient impacts on the freezing behavior exhibited by

experimental mice across sexes. All conditioning procedures here, though, culminated in similar

final freezing levels across sexes of both experimental mice and their targets. This facilitates

comparisons of the different conditioning manipulations on subsequent social and freezing

behaviors by minimizing potential acquisition confounds.

Table 2. Two-way repeated measures general linear model on learning of social conditioning

for mice of both sexes with different sex targets.

Different Sex Experiment – Learning Social Conditioning			
	F Statistic	**p value**	**partial η^2**

Time	$F_{(4.35,304)}=184.2$	<0.001	0.725
Group	$F_{(4,70)}=38.88$	<0.001	0.690
Sex	$F_{(1,70)}=6.694$	0.012	0.087
Time × Group	$F_{(17.4,304)}=12.67$	<0.001	0.420
Time × Sex	$F_{(4.35,304)}=3.317$	0.009	0.045
Group × Sex	$F_{(4,70)}=0.892$	0.474	0.048
Time × Group × Sex	$F_{(17.4,304)}=1.658$	0.048	0.087

Figure 2. *Day 1 acquisition during social conditioning procedure*. Percent time freezing during social conditioning acquisition for mice in same sex (Panels A, B) and different sex (Panels C, D) experiments for male (A, C) and female (B, D) mice. Numbers of mice graphed within Panels A-D in order: Social Stimulus + Stress (n=8, 9, 9, 8); Shock Control (8, 8, 7, 7); Social Control (n=8 for all); Pre-Contingency Control (n=7, 8, 8, 9); Post-Contingency Control (n=7, 8, 8, 8). Average freezing for the first two min, prior to commencement of acquisition, is

plotted on the x-axis as baseline. The average percent freezing for each 30 second period

following each of the five mild foot shocks are thereafter plotted along x-axis (Post-shock

Periods 1-5). Data graphed as mean ± standard error of the mean.

Social Interaction Testing

Social conditioning in the same sex experiment resulted in a sex × group interaction on social

engagement (Table 3). Moreover, our planned contrast supported **Hypothesis 2** for the same

sex experiment (Fig. 3A; $F_{(1,68)}$=7.037 p=0.010, partial η^2=0.094), though our contrast estimate

was negative. This informed us that our proposed directionality of the sex × group interaction

was not supported by the data. In evaluating means for our planned contrast, we found that log-

transformed social engagement values for Social Stimulus + Stress males (μ=1.008, σ=1.698,

n=8) were indeed decreased compared to all male Controls (μ=2.036, σ=0.269, n=29).

However, the mean for Social Stimulus + Stress females (μ=1.813, σ=0.991, n=32) was actually

higher than that for all pooled female Controls (μ=1.968, σ=0.417, n=9), likely due to some

Social Control females with low social engagement values (Fig. 3A). Pairwise comparisons

(Supplemental Figure S2A, B) corroborate the low mean for Social Control females, plus

indicate that Social Stimulus + Stress males displayed less social engagement than Social

Stimulus + Stress females, reiterating that our paradigm was not successful in its goal.

Table 3. Two-way general linear model on social interaction for mice of both sexes with same

sex targets.

Same Sex Experiment – Log-transformed Social Interaction			
	F Statistic	**p value**	**partial η^2**
Group	$F_{(4,68)}$=2.967	0.025	0.149

Sex	$F_{(1,68)}=0.004$	0.951	0.000
Group × Sex	$F_{(4,68)}=3.097$	0.021	0.154

Fear behavior, indexed as percent time freezing during the social interaction post-test, was also

evaluated. While the goal of our modified paradigm was to ultimately decrease social

engagement following conditioning, we also assessed the level of freezing expressed in the

novel social testing arena. In the same sex experiment, no sex × group interaction occurred,

but main effects of both group and sex were observed (Table 4). Pairwise comparisons

indicated that freezing was higher in Social Stimulus + Stress females versus Shock Control

females, while the same comparison in males revealed no differences (Fig. 3C). In a somewhat

inverted pattern compared to Controls during same sex social interaction, Social Control

females exhibited more freezing during social interaction testing than Social Control males,

Shock Control females, and Pre- and Post-Contingency Control females (Fig. 3C). Overall,

though, freezing levels during social interaction testing were minimal as compared to those

occurring during acquisition (Fig. 2) and context testing (Fig. 4A-B).

Table 4. Two-way general linear model on percent time freezing during social interaction for

mice of both sexes with same sex targets.

Same Sex Experiment – % Freezing During Social Interaction			
	F Statistic	**p value**	**partial η^2**
Group	$F_{(4,68)}=11.45$	<0.001	0.402
Sex	$F_{(1,68)}=6.252$	0.015	0.084
Group × Sex	$F_{(4,68)}=1.207$	0.316	0.066

Contrasting with the same sex experiment, there was no significant interaction nor any main

effects in the different sex experiment for social engagement (Table 5). Further distinguishing

the different sex experiment with the same sex experiment, our **Hypothesis 2** planned contrast

for the different sex experiment did not achieve significance (Fig. 3B; $F_{(1,67)}=0.033$ p=0.856,

partial η^2=0.000). Pairwise comparisons demonstrated that Social Stimulus + Stress did not

generate any sex differences in social engagement (Supplemental Figure S2B). The only sex

effect observed was in Shock Controls, with females demonstrating less social engagement

than males, though this appears driven by a single Shock Control female (Supplemental Figure

S2B). Combined, these findings indicate that the social engagement of female mice with other

females is more affected by prior aversive experiences in the *absence* of any social stimuli,

whereas males are more socially affected by temporally concurrent aversive associations with

their own sex.

Table 5. Two-way general linear model on social interaction for mice of both sexes with

different sex targets.

Different Sex Experiment – Log-transformed Social Interaction			
	F Statistic	**p value**	**partial η^2**
Group	$F_{(4,67)}=0.726$	0.577	0.042
Sex	$F_{(1,67)}=1.756$	0.190	0.026
Group × Sex	$F_{(4,67)}=1.172$	0.331	0.065

Unlike the same sex experiment, neither an interaction nor main effects of sex and group were

observed with freezing behavior during social interaction testing for the different sex experiment

(Table 6). Moreover, no significant pairwise comparisons were observed (Fig. 3D). The

predominant consistency across experiments for freezing was that expression of fear was

minimal during the post-test of social interaction, as compared to freezing quantified in the

conditioning context (Figs. 2, 4A-B).

Table 6. Two-way general linear model on percent time freezing during social interaction for

mice of both sexes with different sex targets.

Different Sex Experiment – % Freezing During Social Interaction			
	F Statistic	**p value**	**partial η^2**
Group	$F_{(4,67)}=0.698$	0.596	0.040
Sex	$F_{(1,67)}=3.414$	0.069	0.048
Group × Sex	$F_{(4,67)}=0.491$	0.743	0.028

Figure 3. *Day 2 social interaction and fear behaviors following social conditioning*. Log-transformed social interaction (Panels A, B) and percent time freezing during post-test social interaction (Panels C, D) data are shown for mice in the same (A, C) and different (B, D) sex experiments. Numbers of mice graphed within each panel, left to right: A) n=8, 7, 8, 7, 7, 9, 8, 8, 8, 8; B) n=9, 6, 8, 8, 8, 8, 7, 7, 9, 7; C) n=8, 7, 8, 7, 7, 9, 8, 8, 8, 8; D) n=9, 6, 8, 8, 8, 8, 7, 7, 9, 7. Y axes for A, B were split to visualize data points for three total mice across three separate groups that exhibited no social interaction. Y axes for C, D were split to facilitate clearer visualization of the low freezing levels exhibited during social engagement testing. An *a priori* planned contrast evaluating if log-transformed social interaction was reduced in Social Stimulus + Stress mice across sexes compared to all four Control groups within the same

experiment was significant for mice in the same sex experiment (Panel A; $F_{(1,68)}$=7.037 p=0.010,

partial η^2=0.094). Results of pairwise comparisons with Bonferroni correction, left to right in

Panel C: [B]indicates (Social Controls male vs. female) p=0.004; [a]indicates (Shock Control vs.

Social Stimulus + Stress) p=0.008; [e]indicates (Shock Control vs. Social Control) p<0.001;

[h]indicates (Social Control vs. Pre-Contingency Control) p<0.001; [i]indicates (Social Control vs.

Pre-Contingency Control) p=0.001. Data graphed as mean ± standard error of the mean.

Social Conditioning Context Fear Testing

Because of the natural extinction process that can occur during testing, we specifically

examined behavioral expression of social conditioning context fear during minutes two through

six of testing, to best capture fear expression with minimal confounds from extinction processes

(Lynch et al., 2017; Beaver et al., 2023; Weber et al., 2023). This testing further allowed us to

determine if social engagement testing on Day 2 might have any carry-over effects on

contextual fear expression on Day 3 (Tables 7-8, Fig. 4A-B).

For the same sex experiment, no two-way interaction between group × sex was found, but a

significant main effect of group emerged (Table 7). Given omnibus statistics did not report an

interaction, it is not surprising that our planned contrast for **Hypothesis 3** in the same sex

experiment likewise was not significant (Fig. 4A; $F_{(1,69)}$=0.733 p=0.395, partial η^2=0.011). This

was further supported by pairwise comparisons (Supplemental Figure S3A), which indicated

that male Social Stimulus + Stress mice exhibited significantly less average freezing in the

conditioning context compared to female Social Stimulus + Stress mice (Fig. 4A). Shock

Control mice of both sexes exhibited reduced fear behavior compared to all other groups, as

expected (Fig. 4A).

Table 7. Two-way general linear model on both sexes' context fear testing average with same

sex targets.

Same Sex Experiment – Fear Expression Average			
	F Statistic	**p value**	**partial η²**
Group	$F_{(4,69)}=108.5$	<0.001	0.863
Sex	$F_{(1,69)}=2.859$	0.095	0.040
Group × Sex	$F_{(4,69)}=0.676$	0.611	0.038

In the different sex experiment, no significant group × sex interaction was found, nor was a

significant effect of sex (Table 8). A significant group effect was similarly observed for the

different sex experiment (Fig. 4B). As with the same sex experiment, the lack of a significant

omnibus interaction was reflected in the non-significant outcome of our planned contrast for

Hypothesis 3 ($F_{(1,67)}=0.037$ p=0.848, partial $η^2$=0.001). Pairwise comparisons once more

supported low freezing in Shock Controls across sex relative to all other groups, plus indicated

that male Social Stimulus + Stress mice froze significantly less than Post-Contingency Control

males (Supplemental Figure S3B).

Integrated with the results from social interaction testing, these findings indicate that conditioned

shifts in social engagement are distinct from conditioned fear to the social conditioning context.

In parallel, freezing behavior was relatively minimal during social interaction testing compared to

testing in the conditioning context, suggesting both that freezing behavior did not confound

social interaction outcomes and that mice were able to sufficiently discriminate between

contexts. Curiously, interaction of male Social Stimulus + Stress mice with familiar targets in a

novel environment on Day 2 might lead to mild reductions in contextual fear expression on Day

3.

Table 8. Two-way general linear model on both sexes' context fear testing average with

different sex targets.

Different Sex Experiment – Fear Expression Average			
	F Statistic	**p value**	**partial η^2**
Group	$F_{(4,67)}$=155.9	<0.001	0.903
Sex	$F_{(1,67)}$=0.008	0.929	0.000
Group × Sex	$F_{(4,67)}$=1.220	0.311	0.068

Serum Corticosterone

Log-transformed corticosterone levels of mice in the same sex experiment exhibited no

significant interaction of group × sex, but a main effect of group was observed (Supplemental

Table S1, Supplemental Figure S3C). Pairwise comparisons showed that, across sex, Social

Stimulus + Stress mice in the same sex experiment exhibited elevated log-transformed

corticosterone levels compared to their respective same sex Shock Control mice (Supplemental

Figure S3C). While this was not surprising, male Pre- and Post-Contingency Control mice also

exhibited higher log-transformed corticosterone levels than Shock Control mice, whereas Social

Control mice did not (Supplemental Figure S3C). Conversely, female Social Control mice had

higher log-transformed corticosterone levels than Shock Control mice, but female Pre- and

Post-Contingency Control mice did not (Supplemental Figure S3C). Given no target was present

during Day 3 testing, these findings suggest enduring and sex-specific physiological differences

following social conditioning, despite minimal behavioral shifts. Specifically, any social

interaction with their assigned target on Day 1, whether concurrent or temporally separate with

foot shock exposure, elevated male log-transformed corticosterone levels but attenuated female

log-transformed corticosterone levels after context testing on Day 3 *in the absence of any*

targets. Only Social Stimulus + Stress females did not align with this pattern, exhibiting

heightened log-transformed corticosterone levels on Day 3 similar to those of Social Control

females (Supplemental Figure S3C).

Log-transformed corticosterone levels of mice in the different sex experiment, while not

exhibiting a significant sex × group interaction, did display significant main effects of both group

and sex (Supplemental Table S2). In contrast to the same sex experiment, pairwise

comparisons indicated that within each sex, levels of log-transformed corticosterone did not

differ across groups, including relative to Shock Control mice (Supplemental Figure S3D). The

only difference that reached significance was male Social Stimulus + Stress mice having lower

log-transformed corticosterone levels compared to female Social Stimulus + Stress mice

(Supplemental Figure S3D). Combined with log-transformed corticosterone data from the same

sex experiment (Supplemental Figure S3C), these findings illustrate how same sex target

interactions on the same day as an aversive learning experience can elicit sex-specific and

enduring hormone changes. Moreover, these lasting physiological shifts are not mirrored after

different sex target interactions. This indicates the combination of target and experimental sex,

when temporally concurrent or adjacent to a fear learning event, influences endocrine

responses to context days later.

Figure 4. Day 3 social conditioning context fear expression averages. Average context fear behavior (Panels A, B) data are shown for mice in all groups in the same sex (A) and different sex (B) experiments. Numbers of mice graphed within Panels A-B in order, left to right: A) n=8, 8, 8, 7, 7, 9, 8, 8, 8, 8; B) n=8, 7, 8, 7, 7, 8, 7, 8, 9, 8. Left to right: [S]p=0.049, [d]indicates (Pre-Contingency Control vs. Social Stimulus + Stress) p=0.004. *p<0.001 Social, Pre-Contingency, and Post-Contingency Controls and Social Stimulus + Stress mice vs. Shock Control. Data graphed as mean ± standard error of the mean.

Discussion

Our goal in executing these experiments was to begin establishing a paradigm for inducing consistent, reproducible socially paired stress in mice across sexes. While our results indicate this specific paradigm does not yet accomplish our goal, we want to share our approach and findings so that others can utilize this information to efficiently prioritize future efforts in this domain. Moreover, despite a mostly negative outcome towards our overarching goal, we

nevertheless uncovered sex-specific patterns of behavioral and physiological stress responsivity

that are useful to fields incorporating sex as a biological variable, and/or seeking to employ

different sex social stressors in mice.

All mice that underwent social conditioning with mild foot shocks (Social, Pre-Contingency, and

Post-Contingency Controls plus Social Stimulus + Stress mice) ultimately ended their

acquisition stage at similar freezing levels. Subsequent outcomes in behavioral testing for

social engagement and contextual fear expression, as well as for serum corticosterone, could,

therefore, be interpreted with minimal concern for acquisition confounds. Conversely, this

consistency might have contributed to our negative findings. Alternatively, they might indicate

that our social conditioning protocol resulted in a ceiling effect. Additional studies would need to

parse this out.

Our social interaction findings were not as robust as anticipated. For our same sex experiment,

we found that male Social Stimulus + Stress mice engaged less during social testing compared

to Shock Control mice, and compared to Social Control mice that never encountered a target

during training. Additionally, male Social Stimulus + Stress mice exhibited less social

engagement than female Social Stimulus + Stress mice in same sex experiment, whereas

social engagement was unaffected across sexes in the different sex experiment. This reveals

that male mice likely form stronger associations when aversive experiences occur in conjunction

with the presence of another male mouse, whereas female mice appear to reduce social

engagement after experiencing non-social aversive stimuli. Such a sex difference could help

explain some of the challenges in developing social stress paradigms using female laboratory

mice (*Mus musculus*). Relatedly, this could indicate that, in the presence of a potential sexual

partner, threat assessment is suppressed. The inverse has been demonstrated in female rats;

that is, fear attenuates sexual behaviors, at least in part through amygdala and hypothalamus

estrogen receptor signaling (Moëne et al., 2019). Indirect evidence in male rats housed with

females post-context fear conditioning suggests that encounters with different sexes may

suppress threat assessment. These male rats subsequently exhibited attenuated fear

expression, involving dopamine receptor signaling in the hippocampus (Bai et al., 2009).

However, a different group studying mice reported that ejaculation by males was required for

retention of extinguished social fear conditioning (Grossmann et al., 2024). Continued

evaluations are needed to determine whether our observations – under conditions where

copulation is impossible – that male exposure to a female can diminish threat assessment

behaviors are reproducible. Evidence for social transmission of stress in rodents suggests an

alternative interpretation (Brandl et al., 2022). Instead, it could be that because the target for

each Social Stimulus + Stress mouse did not experience the same stress (mild foot shock), this

in turn affects the Social Stimulus + Stress mouse's perception of their own aversive experience

(Guzmán et al., 2009), likely in a sex-specific manner.

These sex-specific behavior patterns correspond to evidence that accumbal dopamine signaling

dynamics in mice are sex-dependent, both upon the sex of the studied mouse and the sex of

their assigned target (Dai et al., 2022). Our findings align with the success of using social defeat

to stress male mice. Additionally, they indicate that future studies seeking to suppress female

mice's social interaction behavior through a socially associated stimulus will meet with more

success if using an approach distinct from ours, which produced null results. Indeed, others

have encountered challenges when trying to elicit behavioral shifts in female mice, even after

four weeks of stress that involved only female targets (Díez-Solinska et al., 2023). One group

reported that female mice consistently prefer a socially paired food reinforcer, even if the pairing

was with a same sex target that had just previously undergone acute stress (mild foot shock)

(Kietzman et al., 2022). Because males' food preference was not studied by the researchers,

and neither did their females encounter stressed different sex targets, it remains unclear if these

effects are sex-specific, and if they would generalize to different sex interactions. Future studies

will also need to determine how socially conditioning female mice to male targets subsequently

affects interactions with female targets, and vice versa.

One day after social engagement testing, mice were tested for fear expression in the social

conditioning context in the absence of any targets. A sex difference in context fear expression

was observed between Social Stimulus + Stress mice in the same sex experiment, but not in

the different sex experiment. This is the only directionally comparable outcome with social

engagement, in that male Social Stimulus + Stress mice in the same sex experiment both

exhibited less context fear and less social engagement versus female Social Stimulus + Stress

mice. This could be interpreted in at least three ways. Male Social Stimulus + Stress mice

might have found their male target more salient than the context, thereby leading them to form

stronger associations between the mild foot shocks and their targets as compared to female

Social Stimulus + Stress mice. Another possibility is that when mice are in the vicinity of a

possible sexual partner, they could experience comparably impaired encoding of context cues,

though as discussed in the preceding three paragraphs, literature investigating such target-sex

based encoding effects is limited. Alternatively, the presence of the male target may have led to

social buffering of the mild foot shocks for male Social Stimulus + Stress mice (Guzmán et al.,

2009; Brandl et al., 2022), whereas this process did not occur for female Social Stimulus +

Stress mice in the same sex experiment. Because our goal was for mice to associate the

aversive mild foot shock with their assigned target, any social buffering of social conditioning

context fear could counteract our goal. This is why our use of targets of a different strain, that

were never housed in the same room as experimental mice, ensured they had never previously

been encountered (even via olfaction) by the mice tested here and therefore at baseline there

was no familiarity. Given the critical contribution of attachment to social buffering (see review

(Kiyokawa and Hennessy, 2018)), we can be reasonably confident that there were no social

buffering effects upon social conditioning context fear in same sex experiment mice. Thus, we

are inclined toward the first interpretation.

Serum was collected 30 minutes after context fear testing, 24 h after experimental mice's last

encounter with their assigned target. We were interested to discover that sex-specific group

differences in log-transformed corticosterone levels in the same sex experiment appeared to be

best explained by Day 1 group conditions, rather than behavior on Days 1-3. For example,

male Shock Control mice's log-transformed corticosterone levels were lower than Pre- and

Post-Contingency Controls and Social Stimulus + Stress groups – the three conditions where

both target exposure and foot shocks occurred within an eight h window on Day 1. Social

Control male mice only experienced foot shocks, and did not encounter a target, on Day 1.

Shock Control males similarly only encountered targets, and did not experience foot shocks on

Day 1; and male Social Control log-transformed corticosterone levels did not differ from Shock

Control males. Like males in the same sex experiment, female Social Stimulus + Stress mice

had elevated log-transformed corticosterone relative to Shock Control females. In contrast,

however, with this experiment's females, only Social Control females had higher log-

transformed corticosterone levels than Shock Control females. Pre- and Post-Contingency

Control females' log-transformed corticosterone was not different from Shock Controls, and the

former groups encountered targets four h before or after, respectively, experiencing foot shocks.

Collectively, this suggests social encounters temporally distal from the foot shock, but still

occurring within the same day, might augment males' and mitigate females' log-transformed

corticosterone responses to a stressful experience *two days later*. In contrast, coinciding foot

shocks and target exposure in Social Stimulus + Stress mice obscured this sex difference in

endocrine response.

Unlike the same sex experiment, the different sex experiment mice's log-transformed

corticosterone levels did not map onto groups in either sex, despite a significant group effect

being detected. Also distinct from the same sex experiment is that log-transformed

corticosterone levels of Social Stimulus + Stress mice in the different sex experiment had a sex

difference. Specifically, female Social Stimulus + Stress mice had greater log-transformed

corticosterone levels than male Social Stimulus + Stress mice. This does not map onto their

social engagement on Day 2, nor their average context fear expression on Day 3. This log-

transformed corticosterone level difference could, therefore, be an enduring endocrine effect of

encountering their assigned targets on Day 2.

To ensure we rigorously evaluated this modified paradigm's hypothesized utility as a higher

throughput approach for more consistent induction of social stress across sexes, we included

four separate controls. Had we only included one Control group, e.g., Shock Controls, we might

have incorrectly concluded that our modified paradigm successfully reduced social engagement

in females, if not across sexes. Through the inclusion of four Control groups, however, we have

concluded that our findings do not support our hypothesis, and that we cannot endorse this

paradigm in its current state. Instead, our negative findings indicate that this protocol requires

additional optimization to achieve its overarching goal. Nevertheless, we are sharing our

approach and findings so that others with similar procedure goals can learn from our

observations, and improve upon our study's limitations. If we had been able to continue our

investigations, we would have next reduced the mild foot shock amplitude, the number of foot

shocks, or some combination thereof. Given the group differences in freezing that emerged

early during our social conditioning, milder and/or less aversive stimuli could uncover enduring

effects of group conditions. To begin though, it was important here that all our groups attained

similar starting points to minimize confounds for testing interpretations. We would also advise

other researchers to consider optimizing their apparatus and software, if possible, to facilitate

consistent analyses of interaction and freezing, if their conditioning/stress component generates

freezing. We used one software to administer foot shocks while simultaneously recording and

quantifying freezing, and another software to track animal location and identify time spent within

specific arena locations, each in separate apparatus. Both software programs excel at their

purposes, but each have unique – and unfortunately incompatible – video formats plus

requirements regarding lighting, contrast, and background for optimal behavior/location

detection.

Another limitation is that we could not evaluate log-transformed corticosterone levels over time

without introducing additional confounding stressors to repeatedly sample blood. Daily blood

sampling would reveal if our log-transformed corticosterone observations were attributable to

specific days, how log-transformed corticosterone levels mapped on to each day's behavior

measure, and/or if log-transformed corticosterone differences are transient or persistent. Our

experiments were all performed in sexually naïve mice. We recognize that sexually

experienced mice (for example (Newman et al., 2019)) of both sexes might respond,

behaviorally and physiologically, in a manner distinct from what we report here. Earlier

practices of dividing mice into 'susceptible' and 'resilient' groups following social stress have

more recently fallen out of favor, given categorization of those responses are context-specific,

require substantial rodent numbers for optimal data confidence, and neither phenotype can be

universally deemed 'better' or 'worse' (Koolhaas et al., 2017; Lyons et al., 2023). We therefore

evaluated all Social Stimulus + Stress mice as a single group. Still, bimodal distribution of

social engagement in Social Stimulus + Stress mice looks possible in all but the different sex

Social Stimulus + Stress males. Increasing numbers to establish definitive thresholds with

sufficient power for phenotypic grouping could prove informative, particularly for the same sex

experiment. The same limitations as listed above would still apply, however. Finally, our

studies used a brief protocol spanning only four days, intended for high throughput and maximal

comparison to other social stress literature. We therefore cannot speak to whether these

outcomes persist for weeks or months, timeframes that are ethologically relevant for social

learning and memory. Once a short-term protocol has been optimized, duration of effects would

be a critical next step in determining validity of the paradigm.

Here, we have reported a novel, albeit unsuccessful, foray into sex-inclusive efforts to develop a

controllable, less labor-intensive, and higher throughput social stress paradigm in mice. These

experiments have provided utilizable insights regarding how mice of both sexes associate a

same- or different-sex target with an aversive experience. We report that these associations

seem strongest in males after same sex encounters, whereas females appeared not to form

social stimulus associations. Such distinctive responses are probably evolutionarily favorable.

While our investigation has generated new information, our modified paradigm did not

accomplish what we anticipated. We share our negative findings here to encourage and assist

other labs to achieve their own positive outcomes, and to spare them (and mice) the time and

effort of duplicating the present work. Development of a sex-inclusive social stress paradigm for

mice remains a worthwhile pursuit, as such would be a boon across multiple disciplines.

References

References

Abel T, Lattal KM (2001) Molecular mechanisms of memory acquisition, consolidation and

retrieval. Curr Opin Neurobiol 11:180–187.

Ago Y, Tanaka T, Ota Y, Kitamoto M, Imoto E, Takuma K, Matsuda T (2014) Social crowding in

the night-time reduces an anxiety-like behavior and increases social interaction in

adolescent mice. Behav Brain Res 270:37–46.

Agrimson KS, Oatley MJ, Mitchell D, Oatley JM, Griswold MD, Hogarth CA (2017) Retinoic acid

deficiency leads to an increase in spermatogonial stem number in the neonatal mouse

testis, but excess retinoic acid results in no change. Dev Biol 432:229–236.

American Psychiatric Association (2022) Diagnostic and statistical manual of mental disorders

(5th ed., text rev.). https://doi.org/10.1176/appi.books.9780890425787

Bahadır-Varol A, Esen EC, Eren-Koçak E (2022) Repeated collection of vaginal smear causes

stress in mice. Arch Neuropsychiatry 59:325–329.

Bai H, Cao J, Liu N, Xu L, Luo J (2009) Sexual behavior modulates contextual fear memory

through dopamine D1/D5 receptors. Hippocampus 19:289–298.

Beaver JN, Weber BL, Ford MT, Anello AE, Ruffin KM, Kassis SK, Gilman TL (2023)

Generalization of contextual fear is sex-specifically affected by high salt intake. PLOS ONE

18:e0286221.

Becegato M, Meurer YSR, Paiva-Santos MA, Lima AC, Marinho GF, Bioni VS, Soares MBL,

Leão AHFF, Suchecki D, Silva RH (2021) Impaired discriminative avoidance and increased

plasma corticosterone levels induced by vaginal lavage procedure in rats. Physiol Behav

232:113343.

Bergamini G, Mechtersheimer J, Azzinnari D, Sigrist H, Buerge M, Dallmann R, Freije R,

Kouraki A, Opacka-Juffry J, Seifritz E, Ferger B, Suter T, Pryce CR (2018) Chronic social

stress induces peripheral and central immune activation, blunted mesolimbic dopamine

function, and reduced reward-directed behaviour in mice. Neurobiol Stress 8:42–56.

Bernberg E, Andersson IJ, Gan L, Naylor AS, Johansson ME, Bergström G (2008) Effects of

social isolation and environmental enrichment on atherosclerosis in ApoE−/− mice. Stress

11:381–389.

Blanchard DC, Blanchard RJ (1988) Ethoexperimental Approaches to the Biology of Emotion.

Annu Rev Psychol 39:43–68.

Blanchard RJ, McKittrick CR, Blanchard DC (2001) Animal models of social stress: effects on

behavior and brain neurochemical systems. Physiol Behav 73:261–271.

Bloomberg L, Meyers J, Braverman MT (1994) The Importance of Social Interaction: A New

Perspective on Social Epidemiology, Social Risk Factors, and Health. Heal Educ Behav

21:447–463.

Bonnefil V, Dietz K, Amatruda M, Wentling M, Aubry AV, Dupree JL, Temple G, Park H-J,

Burghardt NS, Casaccia P, Liu J (2019) Region-specific myelin differences define

behavioral consequences of chronic social defeat stress in mice. eLife 8:e40855.

Brandl HB, Pruessner JC, Farine DR (2022) The social transmission of stress in animal

collectives. Proc Royal Soc B 289:20212158.

Brown KL, Kennard JA, Sherer DJ, Comalli DM, Woodruff-Pak DS (2011) The context

preexposure facilitation effect in mice: A dose–response analysis of pretraining scopolamine

administration. Behav Brain Res 225:290–296.

Cai DJ et al. (2016) A shared neural ensemble links distinct contextual memories encoded close

in time. Nature 534:115–118.

Caldwell HK, Aulino EA, Rodriguez KM, Witchey SK, Yaw AM (2017) Social Context, Stress,

Neuropsychiatric Disorders, and the Vasopressin 1b Receptor. Front Neurosci 11:567.

Chari T, Griswold S, Andrews NA, Fagiolini M (2020) The Stage of the Estrus Cycle Is Critical

for Interpretation of Female Mouse Social Interaction Behavior. Front Behav Neurosci

14:113.

Coccurello R, D'Amato FR, Moles A (2009) Chronic social stress, hedonism and vulnerability to

obesity: Lessons from Rodents. Neurosci Biobehav Rev 33:537–550.

Correia A da S, Schmitz M, Fischer A, Correia S da S, Simonetti FL, Saher G, Goya-Maldonado

R, Arora AS, Fischer A, Outeiro TF, Zerr I (2024) Cellular prion protein acts as mediator of

amyloid beta uptake by caveolin-1 causing cellular dysfunctions in vitro and in vivo.

Alzheimer's Dement 20:6776–6792.

Curley JP, Rock V, Moynihan AM, Bateson P, Keverne EB, Champagne FA (2010)

Developmental Shifts in the Behavioral Phenotypes of Inbred Mice: The Role of Postnatal

and Juvenile Social Experiences. Behav Genet 40:220–232.

Dai B, Sun F, Tong X, Ding Y, Kuang A, Osakada T, Li Y, Lin D (2022) Responses and

functions of dopamine in nucleus accumbens core during social behaviors. Cell Rep

40:111246.

de la Zerda SH, Netser S, Magalnik H, Briller M, Marzan D, Glatt S, Abergel Y, Wagner S

(2022) Social recognition in laboratory mice requires integration of behaviorally-induced

somatosensory, auditory and olfactory cues. Psychoneuroendocrinology 143:105859.

Díez-Solinska A, Azkona G, Muñoz-Culla M, Beitia-Oyarzabal G, Goñi-Balentziaga O, Gómez-

Lazaro E, Vegas O (2023) The role of sociability in social instability stress: Behavioral,

neuroendocrine and monoaminergic effects. Physiol Behav 270:114306.

Donovan M, Mackey CS, Platt GN, Rounds J, Brown AN, Trickey DJ, Liu Y, Jones KM, Wang Z

(2020) Social isolation alters behavior, the gut-immune-brain axis, and neurochemical

circuits in male and female prairie voles. Neurobiol Stress 13:100278.

Dulka BN, Lynch JF, Latsko MS, Mulvany JL, Jasnow AM (2015) Phenotypic responses to

social defeat are associated with differences in cued and contextual fear discrimination.

Behav Process 118:115–122.

Furman O, Tsoory M, Chen A (2022) Differential chronic social stress models in male and

female mice. Eur J Neurosci 55:2777–2793.

Gilman TL, DaMert JP, Meduri JD, Jasnow AM (2015) Grin1 deletion in CRF neurons sex-

dependently enhances fear, sociability, and social stress responsivity.

Psychoneuroendocrino 58:33–45.

Golden SA, Covington HE, Berton O, Russo SJ (2011) A standardized protocol for repeated

social defeat stress in mice. Nat Protoc 6:1183–1191.

Grossmann CP, Sommer C, Fahliogullari IB, Neumann ID, Menon R (2024) Mating-induced

release of oxytocin in the mouse lateral septum: Implications for social fear extinction.

Psychoneuroendocrinology 166:107083.

Guzmán YF, Tronson NC, Guedea A, Huh KH, Gao C, Radulovic J (2009) Social modeling of

conditioned fear in mice by non-fearful conspecifics. Behav Brain Res 201:173–178.

Iñiguez SD, Flores-Ramirez FJ, Riggs LM, Alipio JB, Garcia-Carachure I, Hernandez MA,

Sanchez DO, Lobo MK, Serrano PA, Braren SH, Castillo SA (2018) Vicarious Social Defeat

Stress Induces Depression-Related Outcomes in Female Mice. Biol Psychiatry 83:9–17.

Jasnow AM, Huhman KL (2001) Activation of GABAA receptors in the amygdala blocks the

acquisition and expression of conditioned defeat in Syrian hamsters. Brain Res 920:142–

150.

Jasnow AM, Shi C, Israel JE, Davis M, Huhman KL (2005) Memory of social defeat is facilitated

by cAMP response element-binding protein overexpression in the amygdala. Behav

Neurosci 119:1125.

Kamps D, Royer J, Dugan E, Kravits T, Gonzalez-Lopez A, Garcia J, Carnazzo K, Morrison L,

Kane LG (2001) Peer Training to Facilitate Social Interaction for Elementary Students with

Autism and Their Peers. Except Child 68:173–187.

Keady J, Fisher M, Anderson E, LeMalefant R, Turner J (2022) Age-specific impacts of nicotine

and withdrawal on hippocampal neuregulin signalling. Eur J Neurosci 56:4705–4719.

Kelley JB, Anderson KL, Altmann SL, Itzhak Y (2011) Long-term memory of visually cued fear

conditioning: roles of the neuronal nitric oxide synthase gene and cyclic AMP response

element-binding protein. Neuroscience 174:91–103.

Kercmar J, Büdefeld T, Grgurevic N, Tobet SA, Majdic G (2011) Adolescent social isolation

changes social recognition in adult mice. Behav Brain Res 216:647–651.

Kietzman HW, Trinoskey-Rice G, Blumenthal SA, Guo JD, Gourley SL (2022) Social

incentivization of instrumental choice in mice requires amygdala-prelimbic cortex-nucleus

accumbens connectivity. Nat Commun 13:4768.

Kiyokawa Y, Hennessy MB (2018) Comparative studies of social buffering: A consideration of

approaches, terminology, and pitfalls. Neurosci Biobehav Rev 86:131–141.

Kondrakiewicz K, Kostecki M, Szadzińska W, Knapska E (2019) Ecological validity of social

interaction tests in rats and mice. Genes, Brain Behav 18:e12525.

Koolhaas JM, Boer SF de, Buwalda B, Meerlo P (2017) Social stress models in rodents:

Towards enhanced validity. Neurobiology Stress 6:104–112.

Kumar M, Rainville JR, Williams K, Lile JA, Hodes GE, Vassoler FM, Turner JR (2021) Sexually

dimorphic neuroimmune response to chronic opioid treatment and withdrawal.

Neuropharmacology 186:108469.

Kuske JX, Trainor BC (2021) Neuroscience of Social Stress. Curr Top Behav Neurosci 54:95–

124.

Laman-Maharg A, Trainor BC (2017) Stress, sex, and motivated behaviors. J Neurosci Res

95:83–92.

Latsko MS, Farnbauch LA, Gilman TL, Lynch JF, Jasnow AM (2016) Corticosterone may

interact with peripubertal development to shape adult resistance to social defeat. Horm

Behav 82:38–45.

Lee J, Raycraft L, Johnson AW (2020) The dynamic regulation of appetitive behavior through

lateral hypothalamic Orexin and Melanin Concentrating Hormone expressing cells. Physiol

Behav:113234.

Leung L (2011) Loneliness, social support, and preference for online social interaction: the

mediating effects of identity experimentation online among children and adolescents. Chin J

Commun 4:381–399.

Levy DR, Hunter N, Lin S, Robinson EM, Gillis W, Conlin EB, Anyoha R, Shansky RM, Datta SR

(2023) Mouse spontaneous behavior reflects individual variation rather than estrous state.

Curr Biol 33:1358-1364.e4.

Lopez J, Bagot RC (2021) Defining Valid Chronic Stress Models for Depression With Female

Rodents. Biol Psychiatry 90:226–235.

Lynch JF, Winiecki P, Gilman TL, Adkins JM, Jasnow AM (2017) Hippocampal GABAB(1a)

Receptors Constrain Generalized Contextual Fear. Neuropsychopharmacol 42:914–924.

Lyons DM, Ayash S, Schatzberg AF, Müller MB (2023) Ecological validity of social defeat

stressors in mouse models of vulnerability and resilience. Neurosci Biobehav Rev

145:105032.

McCormick CM, Green MR, Cameron NM, Nixon F, Levy MJ, Clark RA (2013) Deficits in male

sexual behavior in adulthood after social instability stress in adolescence in rats. Horm

Behav 63:5–12.

McCormick CM, Merrick A, Secen J, Helmreich DL (2007) Social Instability in Adolescence

Alters the Central and Peripheral Hypothalamic-Pituitary-Adrenal Responses to a Repeated

Homotypic Stressor in Male and Female Rats. J Neuroendocr 19:116–126.

Miyashita T, Yamaguchi T, Motoyama K, Unno K, Nakano Y, Shimoi K (2006) Social stress

increases biopyrrins, oxidative metabolites of bilirubin, in mouse urine. Biochem Biophys

Res Commun 349:775–780.

Moëne OL, Stavarache M, Ogawa S, Musatov S, Ågmo A (2019) Estrogen receptors α and β in

the central amygdala and the ventromedial nucleus of the hypothalamus: Sociosexual

behaviors, fear and arousal in female rats during emotionally challenging events. Behav

Brain Res 367:128–142.

Monfils MH, Agee LA (2019) Insights from social transmission of information in rodents. Genes,

Brain Behav 18:e12534.

National Research Council (2011) Guide for the Care and Use of Laboratory Animals (8th ed.).

National Academies Press, Washington, DC. DOI: https://doi.org/10.17226/12910

Newman EL, Covington HE, Suh J, Bicakci MB, Ressler KJ, DeBold JF, Miczek KA (2019)

Fighting Females: Neural and Behavioral Consequences of Social Defeat Stress in Female

Mice. Biol Psychiatry 86:657–668.

Niibori Y, Lee SJ, Minassian BA, Hampson DR (2020) Sexually Divergent Mortality and Partial

Phenotypic Rescue After Gene Therapy in a Mouse Model of Dravet Syndrome. Hum Gene

Ther 31:339–351.

Pan Y, Mou Q, Huang Z, Chen S, Shi Y, Ye M, Shao M, Wang Z (2023) Chronic social defeat

alters behaviors and neuronal activation in the brain of female Mongolian gerbils. Behav

Brain Res 448:114456.

Pantoja-Urbán AH, Richer S, Mittermaier A, Giroux M, Nouel D, Hernandez G, Flores C (2024)

Gains and Losses: Resilience to Social Defeat Stress in Adolescent Female Mice. Biol

Psychiatry 95:37–47.

Peña CJ, Nestler EJ, Bagot RC (2019) Environmental Programming of Susceptibility and

Resilience to Stress in Adulthood in Male Mice. Front Behav Neurosci 13:40.

Pizarro JM, Lumley LA, Medina W, Robison CL, Chang WE, Alagappan A, Bah MJ, Dawood

MY, Shah JD, Mark B, Kendall N, Smith MA, Saviolakis GA, Meyerhoff JL (2004) Acute

social defeat reduces neurotrophin expression in brain cortical and subcortical areas in

mice. Brain Res 1025:10–20.

Prendergast BJ, Onishi KG, Zucker I (2014) Female mice liberated for inclusion in neuroscience

and biomedical research. Neurosci Biobehav Rev 40:1–5.

Russell W, Burch R (1959) The Principles of Humane Experimental Technique. Nature

184:1675–1676 Available at: https://doi.org/10.1038/1841675b0.

Sahani V, Hurd YL, Bachi K (2022) Neuroscience of Social Stress. Curr Top Behav Neurosci

54:483–515.

Sheppard PAS, Koss WA, Frick KM, Choleris E (2018) Rapid actions of oestrogens and their

receptors on memory acquisition and consolidation in females. J Neuroendocr 30.

Siegfried B, Frischknecht H-R, Riggio G, Waser PG (1987) Preexposure to a Nonaggressive

Opponent Prevents Low-Intensity, Social-Conflict Analgesia in Mice. Behav Neurosci

101:423–426.

Smith AS, Wang Z (2012) Salubrious effects of oxytocin on social stress-induced deficits. Horm

Behav 61:320–330.

Sterley T-L, Baimoukhametova D, Füzesi T, Zurek AA, Daviu N, Rasiah NP, Rosenegger D,

Bains JS (2018) Social transmission and buffering of synaptic changes after stress. Nat

Neurosci 21:393–403.

Sterley T-L, Bains JS (2021) Social communication of affective states. Curr Opin Neurobiol

68:44–51.

Takahashi A (2025) The role of social isolation stress in escalated aggression in rodent models.

Neurosci Res 211:75–84.

Teilmann AC, Kalliokoski O, Sørensen DB, Hau J, Abelson KSP (2014) Manual versus

automated blood sampling: impact of repeated blood sampling on stress parameters and

behavior in male NMRI mice. Lab Anim 48:278–291.

Toth I, Neumann ID, Slattery DA (2012) Social Fear Conditioning: A Novel and Specific Animal

Model to Study Social Anxiety Disorder. Neuropsychopharmacology 37:1433–1443.

Uarquin DG, Meyer JS, Cardenas FP, Rojas MJ (2016) Effect of Overcrowding on Hair

Corticosterone Concentrations in Juvenile Male Wistar Rats. J Am Assoc Laboratory Animal

Sci Jaalas 55:749–755 Available at: https://pubmed.ncbi.nlm.nih.gov/27931312/.

Varholick JA, Bailoo JD, Jenkins A, Voelkl B, Würbel H (2021) A Systematic Review and Meta-

Analysis of the Relationship Between Social Dominance Status and Common Behavioral

Phenotypes in Male Laboratory Mice. Front Behav Neurosci 14:624036.

Venniro M, Marino RAM, Chow JJ, Caprioli D, Epstein DH, Ramsey LA, Shaham Y (2022) The

Protective Effect of Social Reward on Opioid and Psychostimulant Reward and Relapse:

Behavior, Pharmacology, and Brain Regions. J Neurosci 42:9298–9314.

Venniro M, Zhang M, Caprioli D, Hoots JK, Golden SA, Heins C, Morales M, Epstein DH,

Shaham Y (2018) Volitional social interaction prevents drug addiction in rat models. Nat

Neurosci 21:1520–1529.

Wang W, Liu W, Duan D, Bai H, Wang Z, Xing Y (2021) Chronic Social Defeat Stress Mouse

Model: Current View on Its Behavioral Deficits and Modifications. Behav Neurosci 135:326–

335.

Wanisch K, Wotjak CT, Engelmann M (2008) Long-lasting second stage of recognition memory

consolidation in mice. Behav Brain Res 186:191–196.

Weber BL, Nicodemus MM, Hite AK, Spalding IR, Beaver JN, Scrimshaw LR, Kassis SK,

Reichert JM, Ford MT, Russell CN, Hallal EM, Gilman TL (2023) Heterotypic Stressors

Unmask Behavioral Influences of PMAT Deficiency in Mice. Int J Mol Sci 24:16494.

Wilson CA, Koenig JI (2014) Social interaction and social withdrawal in rodents as readouts for

investigating the negative symptoms of schizophrenia. Eur Neuropsychopharmacol 24:759–

773.

Wiltgen BJ, Sanders MJ, Behne NS, Fanselow MS (2001) Sex Differences, Context

Preexposure, and the Immediate Shock Deficit in Pavlovian Context Conditioning With Mice.

Behav Neurosci 115:26–32.

Zoicas I, Slattery DA, Neumann ID (2014) Brain Oxytocin in Social Fear Conditioning and Its

Extinction: Involvement of the Lateral Septum. Neuropsychopharmacology 39:3027–3035.