Male and female mice respectively form stronger social aversive memories with same and different sex conspecifics

Jasmin N. Beaver[a,b,c], Marissa M. Nicodemus[a,b], Isabella R. Spalding[a], Sohini Dutta[b,d], Aaron M. Jasnow[e], T. Lee Gilman[a,b,c]*

[a]Department of Psychological Sciences, [b]Brain Health Research Institute, [c]Healthy Communities Research Institute, [d]School of Biomedical Sciences, Kent State University, Kent, OH, USA 44242, [e]Department of Pharmacology, Physiology, and Neuroscience, University of South Carolina School of Medicine, Columbia, SC, USA 29209

Present addresses:

Marissa M. Nicodemus, B.S.
Department of Psychology, West Virginia University
1124 Life Sciences Building
P.O. Box 6040
Morgantown, WV, USA 26506-6040

Sohini Dutta, Ph.D.
Department of Neuroscience
Mount Sinai Medical Hospital, Main
1460 Madison Avenue
New York, NY, USA 10029

*Corresponding author
lgilman1@kent.edu
600 Hilltop Dr.
209 Kent Hall
Kent State University
Kent, OH, USA 44242

Abstract

Mice offer a wealth of opportunities for investigating brain circuits regulating multiple behaviors, largely due to their genetic tractability. Social behaviors are of translational relevance, considering both mice and humans are highly social mammals, and disruptions in human social behavior are key symptoms of myriad neuropsychiatric disorders. Stresses related to social experiences are particularly influential in the severity and maintenance of neuropsychiatric disorders like anxiety disorders, and trauma and stressor-related disorders. Yet, induction and study of social stress in mice is disproportionately focused on males, influenced heavily by their natural territorial nature. Conspecific-elicited stress (i.e., defeat), while ethologically relevant, is quite variable and predominantly specific to males, making rigorous and sex-inclusive studies challenging. In pursuit of a controllable, consistent, high throughput, and sex-inclusive paradigm for eliciting social stress, we have discovered intriguing sex-specific social aversions that are dependent upon the sex of both experimental and conspecific mice. Specifically, we trained male and female F1 129S1/SvlmJ × C57BL/6J mice to associate (via classical conditioning) same or different sex C57BL/6J conspecifics with a mild, aversive stimulus. Upon subsequent testing for social interaction 24 h later, we found that males socially conditioned better to male conspecifics by exhibiting reduced social interaction, whereas females socially conditioned better to male conspecifics. Serum corticosterone levels inversely corresponded to social avoidance after different sex, but not same sex, conditioning, suggesting corticosterone-mediated arousal could influence cross sex interactions. While our paradigm has further optimization ahead, these current findings reveal why past pursuits to develop same sex female social stress paradigms may have met with limited success. Future research should expand investigation of utilizing male mouse conspecifics to instigate social stress across sexes.

Introduction

Social interaction behavior is a cross-mammalian phenomenon seen in humans, non-human primates, rats, hamsters, and mice, among other animals [1–6]. Shifts in social interaction behavior can be informative of the physical and/or mental state of the mammal [7–9]. In humans, disruptions in typical social interaction behavior are characteristic of numerous neuropsychiatric conditions including trauma and stress-related disorders, mood and anxiety disorders, schizophrenia, autism spectrum disorder, and substance use disorders [10].

Rodent studies examining social behavior and socially associated stressors within various contexts have advanced identification of neural pathways important for stress responsivity and social (dis)engagement, as well as assisted development of new therapeutic approaches and targets for neuropsychiatric disorders [6,11–20]. However, social behavior research in the genetically tractable mouse species (*Mus musculus*) has been relatively limited in studying behavioral consequences (and thereby the underlying neurocircuitry and neurophysiological shifts) of negative, aversive, or otherwise stressful social interactions in *female* mice [21,22]. This is largely due to most current social conditioning paradigms employing only male rodents [23–28] - something we ourselves are guilty of [14,29] - by capitalizing upon male-specific territorial aggression. A breadth of social stress techniques have been employed in hamsters, mice, and rats, including social instability [30,31], social transmission of stress [2,32], social isolation [33,34], social defeat [6,35,36], and social crowding [26,37]. Of these, the most prevalent paradigm in mice is that of social defeat (both acute and chronic) [14,29,38–43]. Aside from the inherent male-centric nature of social defeat stress, additional concerns of consistency and reproducibility arise from the inherently variable range of aversive social behaviors, from defensive postures to aggressive bites, that each experimental mouse perceives and experiences. Such variability - much beyond experimenter control - plus the

single sex bias of social defeat stress left us seeking an improved paradigm. In particular, we

sought to pair controllable, uniform, aversive stimuli with the presence of a social stimulus to

study aversive social conditioning *across sexes* in mice using a high throughput approach

(requiring 1 day, as opposed to multiple days or weeks).

To accomplish this, we modified a paradigm previously utilized in males (44–47). In the original

paradigm, manual administration of an aversive stimulus (mild foot shock) occurs selectively

when a male mouse actively investigates a conspecific, with the goal of attenuating subsequent

social engagement (44–47). Rather than using this operant-style approach, where a behavior

exhibited by the experimental mouse directly affects the outcome, we instead wanted to develop

a more classical conditioning approach that was less labor-intensive (i.e., did not require real-

time monitoring for manual shock administration). In this way, experimental mice would not

necessarily need to approach the social stimulus (conspecific) to develop an association

between it and the aversive stimulus. Additionally, all mice would receive the same number of

shocks, instead of variable numbers based upon behavior (44–47).

Using our modified version of this novel approach, we evaluated how employing this paradigm

affected multiple measures across *both sexes* after being exposed to an assigned, previously

never encountered, conspecific. Specifically, mice were socially conditioned (SC) with an

aversive stimulus (mild foot shock) when in the presence of a novel conspecific, independent of

investigative behavior. Then, mice were tested for social engagement in a novel environment in

the presence of the same conspecific, followed by testing of behavior in the conditioning

environment in the absence of any social stimulus. These behaviors were tested both under

circumstances when the experimental and conspecific mice were the same sex and when they

were different sexes. We hypothesized that SC mice would exhibit reduced social behavior

when tested with their assigned conspecific in a novel environment, regardless of if the SC and

conspecific mice were the same or different sexes. Our broad goal was to develop a paradigm

that would provide a useful first step for future investigations of social behavior across mouse

sexes, which are currently underrepresented in literature (22).

Methods

Mice

Hybrid F1 offspring of both sexes resulting from pairing female 129S1/SvlmJ and male

C57BL/6J mice (hereafter experimental mice), and male and female C57BL/6J mice (hereafter

conspecifics), all ≥9 weeks old or older, were group-housed (2-5 per cage) within sex. All mice

had *ad libitum* access to food and water in rooms maintained on a 12:12 light/dark cycle with

lights on at 07:00 local standard time, and temperature maintained at 22 ± 2°C. All mice were

fed LabDiet 5001 rodent laboratory chow (LabDiet, Brentwood, MO) and were kept on 7090

Teklad Sani-chip bedding (Envigo, East Millstone, NJ) in cages containing Nestlets (Ancare,

Bellmore, NY) and huts (Bio-Serv, Flemington, NJ) for enrichment. Experiments were approved

by the Kent State University Institutional Animal Care and Use Committee, and adhered to the

National Research Council's Guide for the Care and Use of Laboratory Animals, 8th Ed.

undefined.

Social Conditioning Paradigm

The entire paradigm spans four consecutive days. Procedures for each individual day are

outlined below in order.

Pre-exposure (Day 0)

Pre-exposure was used to help reduce the novelty of the transparent enclosure and/or the conspecific to the experimental mouse, plus minimize any potential sex differences in acquisition (48–51). On Day 0, experimental mice were placed in Coulbourn Instruments chambers (17.8 cm D × 17.8 cm W × 30.5 cm H; Allentown, PA) with (Control A and SC mice) or without (Control B-D mice) a conspecific (age-matched; either same sex or different sex relative to experimental mouse, depending on if assigned to same sex or different sex experiment) for pre-exposure (Fig. 1). Chambers comprised two opposing aluminum walls each adjoining two clear acrylic walls. Conspecifics were put inside a transparent enclosure placed within a corner of the social conditioning chamber. The transparent enclosure included 23 holes (each 0.32 cm diameter) on each of the 2 sides facing the SC mouse to enable olfactory cue exchanges. Previous work demonstrates that olfaction is the most important modality when recognizing and interacting with novel conspecifics (see review (52)); tactile (whiskers) and auditory modalities appear to only be recruited when recognizing cage mates (46) in male mice (females were not studied). Chambers and enclosures were cleaned with 70% ethanol before and after each session; chambers had visible illumination and one clear acrylic wall was marked with a blue dotted pattern. These components contributed olfactory and visual cues to the context, in addition to the tactile cue of the stainless steel grid floor for the experimental mouse. During pre-exposure, experimental mice were allowed to explore the chamber and conspecific enclosure for 5 min, then both mice were returned to their respective home cages.

Social Conditioning (Day 1)

On Day 1, experimental mice were placed in the same social conditioning chamber as Day 0 and received five, 1 s mild foot shocks (1.0 mA) in the presence of the same conspecific that they encountered on Day 0 (Figure 1A). This was to have experimental mice form associations between aversive foot shocks and their respective conspecific (Fig. 1). Conditioning lasted for 9

minutes; after a 2 minute baseline, shocks were then administered at 120, 210, 300, 390, and

480 s. Conspecifics did not receive foot shocks. Cameras mounted above the chambers were

used to record movements, and freezing behavior during conditioning was quantified using

FreezeFrame software (v. 5.201, Actimetrics, Wilmette, IL). Freezing behavior is indicative of

fear/aversive conditioning in mice; freezing was defined as the absence of movement except

breathing.

Social Interaction Testing (Day 2)

On Day 2, experimental mice were placed in a novel room containing open arenas (41.9 cm W

× 41.9 cm D × 39.6 cm H). Mice acclimated to the room in their home cage for 30 min prior to

behavior testing commencing. Each open arena had one empty PVC tube (8.9 cm outer

diameter) in a single corner of the square chamber with wire mesh (6.35 mm^2 openings)

covering a rectangular portion (11.4 cm W x 4.1 cm H) cut out of the tube bottom. Experimental

mice were placed in the corner of the arena opposite the PVC tube and could investigate freely

for 2.5 minutes (pre-test). Immediately after these 2.5 min, the conspecific that each

experimental mouse had previously been conditioned with on Day 1 was then placed in the PVC

tube in the corner. Experimental mice were allowed to continue investigating the arena for an

additional 5 min (post-test; Fig. 1). Because of the mesh at the bottom of the PVC tubes

containing conspecifics, exchanges of visual, auditory, and olfactory cues between experimental

mice and conspecifics were possible, but tactile interactions were minimized. Interaction

behavior was quantified as duration when at least 80% of an experimental mouse's body was in

the interaction zone, a 7 cm radius zone (~223 cm^2 – 240 cm^2, i.e., 14-15% of the open arena)

around the base of the PVC tube. This was quantified using ANY-maze software (v. 7.09

Stoelting Co., Wood Dale, IL). Post-test social interaction time was log-transformed

(Y=log(Y+0.001) to account for 3 mice with 0 s post-test social interaction) so that data would be

normally distributed for statistical analyses. Females were intentionally not assessed for estrus

cycle stage for five reasons: 1) to minimize mouse usage (53), we did not power our studies for

assessment of estrus; 2) with the goal of developing a social stress paradigm, we are seeking

effects robust enough to not depend upon estrus in intact, randomly cycling females; 3) we

intentionally focused here on fear and social behaviors, and did not measure sexual behaviors

(e.g., lordosis); 4) evidence that overall mouse behaviors are not affected by estrus stage

(54,55), (but see (56)); 5) cross-species evidence indicates vaginal lavage to determine estrus

cycle is stressful (57,58), and we sought to minimize stress confounds here.

Context Testing (Day 3)

On the last day of the social conditioning paradigm, experimental mice were placed in the social

conditioning chamber for behavior testing in the *absence* of any conspecific (Fig. 1). The

conspecific enclosure was still present to keep the context consistent with conditioning. All

other tactile, visual, and olfactory cues from Day 1 were present. Testing lasted 10 min and did

not involve any foot shocks. Freezing behavior was again quantified using FreezeFrame

software.

Treatment Groups

The SC group involved mice that underwent social conditioning in the presence of a non-

shocked conspecific (either same sex or different sex relative to SC mouse, depending upon

experiment), with the hypothesis that SC mice would associate the conspecific with this aversive

experience. We planned four Control groups for this one SC group; each experiment (same sex

or different sex) had its own respective set of four Control groups for its respective SC group.

The goals of these were to control for: A) foot shock exposure; B) presence of conspecific

during pre-exposure & foot shock; C & D) exposure to conspecific temporally distal to foot shock

exposure. Control A involved procedures identical to those of the SC group, except a foot shock

was never administered on Day 1 (Fig. 1). Control B was the same as the SC group procedure,

except experimental mice never encountered the conspecific until testing on Day 2 (Fig. 1).

Controls C and D had experimental mice encountering conspecifics either 4 h before (Control C)

or 4 h after (Control D) pre-exposure and social conditioning; in other words, conspecifics were

not present in the conditioning chamber during these two periods. Instead, experimental mice

were exposed to their conspecific in a separate room for 5 minutes or 9 minutes (the same

amount of time as pre-exposure or social conditioning, respectively) 4 h before (Control C) or 4

h after (Control D) the experimental mice were exposed to the conditioning chamber. This

exposure involved experimental mice and their respective conspecific being placed in a clean

mouse cage separated with an acrylic divider to allow for visual, auditory, and olfactory

exchange, but no tactile interaction. Mice in Control C or D conditions still experienced foot

shock on Day 1, and encountered their assigned conspecific for testing on Day 2 (Fig. 1). This

temporal separation of 4 h was to minimize consolidation of social encounters from

interfering/intermingling with consolidation of the aversive foot shock encounter in the social

conditioning chamber, while still making execution of these experiments feasible within a 12 h

lights-on period (59–61) (see reviews (62,63)).

Figure 1. Study timeline and treatment groups. Five groups were used in this study for each experiment (same sex experiment, and different sex experiment): one socially conditioned (SC) group and four Control groups. Day 0 was for pre-exposure; Day 1 was when social conditioning occurred; Day 2 tested for social engagement; Day 3 tested for context-specific freezing behavior. Each Control group was composed of different mice. Control A mice underwent the same exact procedure as detailed for the SC group (above horizontal line),

258	except no foot shock was administered on Day 1. Control B mice similarly experienced the
259	same procedure as mice in the SC group (above horizontal line), save that Control B mice did
260	not encounter any conspecific until Day 2 for testing of social engagement. Control C involved
261	mice encountering their conspecific 4 h before pre-exposure and social conditioning, meaning
262	no conspecific was present for Control C mice when they were in the social conditioning
263	chamber for pre-exposure or social conditioning. To encounter their assigned conspecific,
264	Control C mice were placed into a clean cage with a clear acrylic divider separating them from
265	their conspecific. This divider allowed visual, olfactory, and auditory exchanges but no tactile
266	interactions. On Day 0, Control C mice encountered their conspecific for 5 min; on Day 1, for 9
267	min; these were timed to match the length of conspecific exposure that mice in the SC group
268	experienced. Control D mice were treated the same as Control C mice, except their encounters
269	with their assigned conspecifics occurred 4 h *after*, rather than before, pre-exposure and social
270	conditioning.

Serum Corticosterone

Thirty minutes after all experimental mice underwent contextual fear testing on Day 3, they were

briefly anesthetized with isoflurane then rapidly decapitated for trunk blood collection. Blood

clotted at room temperature for 10 minutes, then was spun at 3500 rpm for 1 h at 4°C. Serum

was collected and stored at -80°C until corticosterone analyses could be performed. Serum

corticosterone was measured using Enzo Life Sciences corticosterone enzyme-linked

immunosorbent assay (ELISA) kits (Farmingdale, NY). Assays were run according to the

manufacturer's instructions using their small volume protocol. Plates were read at 405 nm with

correction at 580 nm. The sensitivity of the assay was 26.99 pg/mL.

Statistical Analyses

Data were graphed with GraphPad Prism 10.3.0 (442), Beta (GraphPad Software, San Diego,

CA), and analyzed using IBM SPSS Statistics 28.0.0.0 (IBM, Armonk, NY), with the significance

threshold set *a priori* at $p < 0.05$. Non-significant trends ($p < 0.10$) were mentioned only when the

associated partial η^2 was > 0.060. Data were graphed as the mean ± 95% confidence interval

(CI). Details of identified outliers (greater than the mean ± 4 standard deviations) are provided

in the Supplemental Dataset. The same sex experiment and different sex experiment each

included their own SC group plus accompanying four Control groups. Control A mice were

analyzed across experiments. Within each experiment, the SC group and other three Control

groups (all of which experienced foot shocks, i.e., shocked mice) were analyzed. Social

conditioning and context fear expression time course were analyzed using 3-way repeated

measures (RM) ANOVAs (shocked mice: time × treatment group × sex of experimental mouse)

(Control A mice: time × experiment × sex of experimental mouse) and pairwise comparisons

with Bonferroni correction. Greenhouse Geisser corrections were utilized for within-subjects

analyses. Measurements of contextual fear expression average, social interaction, and serum

corticosterone were analyzed with a 2-way ANOVA (SC and Control B-D mice: treatment group

× sex of experimental mouse; Control A mice: experiment × sex of experimental mouse) and

pairwise comparisons with Bonferroni correction.

Results

Social Conditioning Acquisition

Social conditioning acquisition was examined in experimental mice in either the presence or

absence of a conspecific. One experiment used same-sex conspecifics, and the other

experiment used different sex conspecifics; each of these experiments included its own SC

group and corresponding four Control groups.

Control A (no foot shock) mice

Control A mice were utilized to distinguish the effects of receiving mild foot shocks on

subsequent social behavior. Figure 2A and 2C show social conditioning 'acquisition' on Day 1

for Control A mice with either a same sex conspecific (Figure 2A) or a different sex conspecific

(Figure 2C). For acquisition, there was a significant three-way interaction of time × conspecific

sex × experiment (Table 1). This appears to be driven primarily by inconsistent time course

patterns across sexes, and mice in the different sex experiment exhibiting lower freezing levels

at several timepoints compared to same sex experiment mice (Figure 2A, 2C).

Table 1. Three-way RM ANOVAs on Control A mice's social conditioning acquisition across

experiments and sexes.

Control A Mice - Fear Acquisition			
	F Statistic	**p value**	**Partial η^2**
Time	$F_{(3.325, 86.45)} = 0.334$	0.821	0.013
Sex	$F_{(1, 26)} = 1.265$	0.271	0.046
Experiment	$F_{(1, 26)} = 14.520$	<0.001	0.358
Time × Sex	$F_{(3.325, 86.45)} = 1.836$	0.141	0.066
Time × Experiment	$F_{(3.325, 86.45)} = 1.128$	0.345	0.042
Sex × Experiment	$F_{(1, 26)} = 0.639$	0.431	0.024
Time × Sex × Experiment	**$F_{(3.325, 86.45)} = 2.642$**	**0.049**	**0.092**

Bold indicates highest interaction or main effect with p<0.05

Two days after social 'conditioning' (see Figure 1 for timeline), Control A mice were tested for

their fear to the context in which they were 'conditioned'. Testing lasted for 10 minutes, and the

time course data for testing Control A mice are in Figure 2B (same sex conspecific) and 2D

(different sex conspecific). We observed no significant three-way interaction of time × sex ×

experiment (Table 2). There were likewise no significant two-way interactions of time × sex, time

× experiment, or sex × experiment (Table 2). Main effects of sex and time were not significant,

though a non-significant trend for time was noted (p=0.099; Table 2). A significant main effect

of experiment was detected (Table 2), and pairwise comparisons support a pattern of female

and male mice in the different sex experiment freezing less than mice in the same sex

experiment (Figure 2B, 2D), as observed with 'training'.

Table 2. Three-way RM ANOVA on Control A mice's context testing time course across

experiments and sexes.

Control A Mice - Context Testing Time Course			
	F Statistic	**p value**	**Partial η^2**
Time	*F(7.884,205.0)=1.710*	*0.099*	*0.062*
Sex	F(1,26)=0.391	0.537	0.015
Experiment	**F(1,26)=16.16**	**<0.001**	**0.383**
Time × Sex	F(7.884,205.0)=0.599	0.776	0.023
Time × Experiment	F(7.884,205.0)=1.063	0.391	0.039
Sex × Experiment	F(1,26)=0.004	0.949	0.000
Time × Sex × Experiment	F(7.884,205.0)=0.963	0.465	0.036

Bold indicates highest interaction or main effect with p<0.05; *italics* indicates p<0.10 and partial η^2 >0.060

Consistent with 'training' and time course testing of Control A mice, averaged freezing during

minutes two through six of testing revealed no sex × experiment interaction, but a significant

main effect of experiment (Table 3). Once more, different sex experiment mice exhibited less

freezing than their counterparts in the same sex experiment (Figure 2E).

Table 3. Two-way ANOVA on Control A mice's context testing average across experiments and

sexes.

Control A Mice - Context Testing Average			
	F Statistic	**p value**	**Partial η^2**
Sex	F(1,26)=0.050	0.825	0.002

Experiment	F(1,26)=15.00	<0.001	0.366
Sex × Experiment	F(1,26)=0.045	0.833	0.002

349   **Bold** indicates highest interaction or main effect with $p < 0.05$

Figure 2. *Day 1 social conditioning 'acquisition' and Day 3 context fear testing in Control A mice.* Control A (no foot shock) mice underwent 'acquisition' or 'training' for social conditioning after a two min baseline period on Day 1 (Panels A,C). Percent time freezing for the same 30 s periods that are graphed for SC and Control B-D mice (Figure 3) are indicated at time points 1-5 in Panels A,C. On Day 3, freezing behavior was quantified for 10 min when mice were re-exposed to the social conditioning chamber (Panels B,D). Average freezing behavior was evaluated between minutes two through six (yellow shading in Panels B,D) of the 10 min test and graphed in Panels C,F. Mice either went through the Control A social conditioning process with same sex conspecifics (Panels A,B,E) or different sex conspecifics (Panels C,D,E). Same sex n=8, different sex n=7. *indicates p=0.017 (A), p=0.026 (Panel C) compared to other sex within the same experiment. [a]indicates p=0.026, p=0.026, p=0.007, left to right, comparing females in same sex (Panel A) versus different sex (Panel C) experiment 'training' at specified timepoints. [b]indicates p=0.016, p=0.002, left to right, comparing males in same sex (Panel A) versus different sex (Panel C) experiment 'training' at specified timepoints. [c]indicates p=0.008, p=0.020, p=0.018, p<0.001, p=0.027, p=0.001, p=0.037, p=0.033, p=0.020, p=0.027, left to right, comparing females in same sex (Panel B) versus different sex (Panel D) experiment testing at specified timepoints. [d]indicates p=0.024, p=0.006, p=0.030, p=0.011, p=0.030, p=0.046, left to right, comparing males in same sex (Panel B) versus different sex (Panel D) experiment testing at specified timepoints. [e]indicates p=0.016, [f]indicates p=0.008, comparing across experiments within sex for average testing freezing (Panel E). Data graphed as mean ± 95% CI.

Socially Conditioned and Control B-D mice

Social conditioning acquisition for mice that received mild foot shocks are illustrated in Figure 3A, 3B for mice in the same sex conspecific experiment, and in Figure 3C, 3D for mice in the

different sex conspecific experiment. All mice expressed >40% freezing during at least one

post-shock period during social conditioning. For acquisition in the same sex experiment, no

three-way interaction of time × sex × group (of which there are four: socially conditioned mice or

Control B-D mice) occurred (Table 4). We did not observe any significant two-way interactions

either (Table 4). Nonetheless, significant main effects of time, sex, and group were found for

mice in the same sex experiment (Table 4).

Table 4. Three-way RM ANOVAs on fear acquisition for mice of both sexes with same sex

conspecifics.

Same Sex Experiment – Fear Acquisition			
	F Statistic	**p value**	**Partial η^2**
Time	**F(4.05,223.0)=145.923**	**<0.001**	**0.726**
Sex	**F(1,55)=14.90**	**<0.001**	**0.213**
Group	**F(3,55)=5.322**	**0.003**	**0.225**
Time × Group	F(12.16,223.0)=1.014	0.437	0.052
Time × Sex	F(4.05,223.0)=1.766	0.136	0.031
Group × Sex	F(3,55)=1.833	0.143	0.093
Time × Group × Sex	F(12.16,223.0)=0.550	0.882	0.029

Bold indicates highest interaction or main effect with p<0.05

Pairwise comparisons indicated that socially conditioned males exhibited less freezing at two

training timepoints relative to Control B mice (Figure 3A) in the same sex experiment. For all

but the last post-shock period in the same sex experiment, female socially conditioned mice

froze significantly more than male socially conditioned mice (Figure 3A, 3B).

Social conditioning acquisition for mice in the different sex experiment is shown in Figure 3C,

3D. No significant three-way interaction of time × sex × group was observed (Table 5), though a

non-significant trend (p=0.076) was noted. While no sex × group interaction was detected, we

did find significant time × group and time × sex interactions (Table 5).

Table 5. Three-way RM ANOVAs on fear acquisition for mice of both sexes with different sex

conspecifics.

Different Sex Experiment – Fear Acquisition			
	F Statistic	**p value**	**Partial η²**
Time	$F(4.342, 251.8) = 203.7$	<0.001	0.778
Sex	$F(1,58) = 5.464$	0.023	0.086
Group	$F(3,58) = 1.098$	0.357	0.054
Time × Group	**$F(13.03, 251.8) = 3.548$**	**<0.001**	**0.155**
Time × Sex	**$F(4.342, 251.8) = 3.321$**	**0.009**	**0.054**
Group × Sex	$F(3,58) = 0.950$	0.423	0.047
Time × Group × Sex	*$F(13.03, 251.8) = 1.636$*	*0.076*	*0.078*

Bold indicates highest interaction or main effect with p<0.05; *italics* indicates p<0.10 and partial η² >0.060

Only the second post-shock period was different across sexes in SC mice in the different sex

experiment (Figure 3C, 3D). For Control C mice, females displayed greater freezing for the first

three post-shock periods than males in the different sex experiment (Figure 3C, 3D). All other

between-group comparisons within each sex did not exhibit consistent patterns, and as with

same sex experiment mice, different sex experiment mice concluded training at similar freezing

levels. Combined with same sex experiment acquisition, these findings indicate that social

exposure – whether concurrent or 4 h prior – can transiently augment social conditioning

acquisition in female mice, depending upon the sex of the conspecific.

Figure 3. *Day 1 acquisition during social conditioning procedure.* Male (Panels A, C) and

female (Panels B, D) mouse numbers are, respectively: Socially conditioned (SC; n=8, 9, 9, 8);

Control B (n=8 for all); Control C (n=7, 8, 8, 9); Control D (n=7, 8, 8, 8). Same sex experiment

data are in Panels A,B; different sex experiment data in Panels C,D. Average freezing for the

first two minutes, prior to commencement of acquisition, is plotted on the x-axis as baseline. The

average percent freezing for each 30 second period following each of the five mild foot shocks

are thereafter plotted along x-axis (Post-shock Periods 1-5). *Same sex experiment*: [a]indicates

p=0.037, p=0.006 (left to right, Panel A) when comparing within males and between SC and

Control B groups. [b]indicates p=0.022 (Panel B) when comparing between sexes and within

Control D group. [c]indicates p=0.011, p=0.047, p=0.012, p=0.016 (left to right, Panel B) when

comparing between sexes within the SC group. *Different sex experiment*: [m]indicates p=0.042

(Panel C) when comparing within males and between Control C and SC groups. [n]indicates

p=0.014 (Panel C) or p=0.046 (Panel D) when comparing within sex and between Control B and

SC groups. [o]indicates p=0.033 (Panel D) when comparing within females and between Control

C and D groups. [p]indicates p=0.010 (Panel D) when comparing between sexes and within

Control B group. [q]indicates p=0.032, p=0.012, p=0.019 (left to right, Panel D) when comparing

between sexes and within Control C group. [r]indicates p=0.020 (Panel D) when comparing

between sexes and within SC group. Data graphed as mean ± 95% CI.

Social Interaction

Twenty four hours after social conditioning, mice were tested for social engagement using the

social interaction test. Three mice did not interact at all with their conspecific, and thus were

excluded from social interaction analyses (1 different sex Control A female; 1 same sex Control

B female; 1 same sex SC male; see Supplemental Dataset). Figure 4A illustrates social

interaction data for Control A mice across experiments. There was no significant sex ×

experiment interaction, but there was a significant main effect of experiment (Table 6). This was

driven by males in the different sex experiment interacting more with female than male

conspecifics (Figure 4A).

Table 6. Two-way ANOVA on social interaction for Control A mice across experiments and

sexes.

Control A Mice – Log-transformed Social Interaction			
	F Statistic	**p value**	**Partial η^2**
Sex	F(1,23)=0.429	0.519	0.018
Experiment	**F(1,23)=8.529**	**0.008**	**0.271**
Sex × Experiment	F(1,23)=2.773	0.109	0.108

Bold indicates highest interaction or main effect with p<0.05

Social conditioning in the same sex experiment did not result in a sex × group interaction, but a

main effect of group was found (Table 7). Pairwise comparisons indicated that SC males

exhibited less social interaction than SC females (Figure 4B).

Table 7. Two-way ANOVA on social interaction for mice of both sexes with same sex

conspecifics.

Same Sex Experiment – Log-transformed Social Interaction			
	F Statistic	**p value**	**Partial η^2**
Sex	F(1,52)=0.438	0.511	0.008
Group	**F(3,52)=2.923**	**0.042**	**0.144**
Sex × Group	F(3,52)=1.870	0.146	0.097

Bold indicates highest interaction or main effect with p<0.05

In contrast to the same sex experiment, there were no significant interactions nor main effects in

the different sex experiment (Table 8). A non-significant trend for group was noted (p=0.066,

Table 8). Pairwise comparisons demonstrated that females exhibited significantly reduced

social interaction, both compared to female Control B mice and to SC male mice (Figure 4C).

Combined, these findings indicate that the social behavior of females is more affected by

aversive associations with males, whereas males are more socially affected by aversive

associations with their own sex.

Table 8. Two-way ANOVA on social interaction for mice of both sexes with different sex

conspecifics.

Different Sex Experiment – Log-transformed Social Interaction			
	F Statistic	**p value**	**Partial η^2**
Sex	F(1,56)=0.405	0.527	0.007
Group	*F(3,56)=2.538*	*0.066*	*0.120*
Sex × Group	F(3,56)=1.678	0.182	0.082

464 *Italics* indicates p<0.10 and partial η^2 >0.060

Control A

Same Sex

Different Sex

 Figure 4. *Day 2 social interaction behavior following social conditioning.* Log-

transformed social interaction data are shown for mice in Control A groups (Panel A), and in the

same (Panel B) and different (Panel C) sex experiments. Numbers of mice graphed within each

panel, left to right: A) n=7, 8, 6, 6; B) n=8, 7, 7, 7, 6, 8, 8, 9; C) n=8, 8, 8, 9, 7, 9, 7, 8. [a]p=0.004,

[b]p=0.038, [c]p=0.029, [d]p=0.017. Data graphed as mean ± 95% CI.

Social Conditioning Context Fear Testing

After Day 2's social interaction evaluation of mice's association between their assigned

conspecific and the aversive stimulus they experienced on Day 1, we next assessed their fear

response to the social conditioning context on Day 3. This was accomplished by quantifying

percent of time spent freezing (Table 9, Figure 5).

In the same sex experiment, no significant three-way interaction was detected, nor were

significant two-way interactions of group × sex or time × sex (Table 9). A non-significant trend

for time × group was noted (p=0.073). Time was found to have a significant main effect (Table

9), as is typical for context fear expression testing, with continued exposure to the social

conditioning context in the absence of the conditioned stimulus resulting in gradual fear

extinction. A significant main effect of sex was also found, likely driven by lower freezing levels

across time in Control C and SC mice (Table 9, Figure 5A, 5B). A non-significant trend for

group (p=0.056, Table 9) was also noted. In the different sex experiment, no significant three-

nor two-way interactions were observed. Along with a main effect of time (Table 10), a

significant effect of group was also found (Table 10). Pairwise comparisons revealed significant

differences between groups at specific timepoints (Figure 5). The most consistent of these for

the same sex experiment involved Control C males in the same sex experiment exhibiting

accelerated extinction processes relative to Control C females, and SC males displaying

attenuated freezing behavior compared to SC females (Figure 5A, 5B). In the different sex

experiment, the most robust time course difference was SC males displaying less freezing

behavior relative to Control D males (Figure 5D).

Because of the natural extinction process that can occur during testing, we also specifically

examined behavioral expression of social conditioning context fear during minutes two through

six of testing, to best capture fear expression with minimal confounds from extinction processes

(61,64,65). With this approach, freezing behavior resulted in a significant main effect of group in

the same sex experiment (Table 11, Figure 5C), though pairwise comparisons did not reveal

any significant between-group differences. While no group × sex interaction occurred for the

different sex experiment, a main effect of group was observed (Table 12). After pairwise

comparisons were evaluated, the significant difference detected in the different sex experiment

was the same as that observed in the time course for males (Figure 5D); SC male mice

displayed significantly less freezing relative to Control D male mice (Figure 5F). Integrated with

the results from social interaction testing, these findings indicate that conditioned shifts in social

engagement are distinct from conditioned fear to the social conditioning context.

Table 9. Three-way RM ANOVAs on both sexes' context fear testing with same sex

conspecifics.

Same Sex Experiment – Fear Expression Time Course			
	F Statistic	**p value**	**Partial η^2**
Time	**F(11.34,623.4)=9.498**	**<0.001**	**0.147**
Sex	**F(1,55)=5.052**	**0.029**	**0.084**
Group	*F(3,55)=2.671*	*0.056*	*0.127*
Time × Group	*F(34.00,623.4)=1.388*	*0.073*	*0.070*
Time × Sex	F(11.34,623.4)=1.743	0.059	0.031
Group × Sex	F(3,55)=0.898	0.448	0.047
Time × Group × Sex	F(34.00,623.4)=1.064	0.372	0.055

512   **Bold** indicates highest interaction or main effect with p<0.05; *italics* indicates p<0.10 and partial η^2 >0.060

514   **Table 10.** Three-way RM ANOVAs on both sexes' context fear testing with different sex

515   conspecifics.

Different Sex Experiment – Fear Expression Time Course			
	F Statistic	**p value**	**Partial η^2**
Time	**F(9.434,518.9)=5.590**	**<0.001**	**0.092**
Sex	F(1,55)=0.030	0.862	0.001
Group	**F(3,55)=3.149**	**0.032**	**0.147**
Time × Group	F(28.30,518.9)=0.863	0.672	0.045
Time × Sex	F(9.434,518.9)= 0.575	0.826	0.010
Group × Sex	F(3,55)=0.844	0.476	0.044
Time × Group × Sex	F(28.30,518.9)=0.483	0.990	0.026

516   **Bold** indicates highest interaction or main effect with p<0.05

518   **Table 11.** Two-way ANOVA on both sexes' context fear testing average with same sex

519   conspecifics.

Same Sex Experiment – Fear Expression Average			
	F Statistic	**p value**	**Partial η^2**
Sex	F(1,55)=3.137	0.082	0.054
Group	**F(3,55)=3.603**	**0.019**	**0.164**
Group × Sex	F(3,55)=0.338	0.636	0.030

520   **Bold** indicates highest interaction or main effect with p<0.05

522   **Table 12.** Two-way ANOVA on both sexes' context fear testing average with different sex

523   conspecifics.

Different Sex Experiment – Fear Expression Average			
	F Statistic	**p value**	**Partial η^2**
Sex	F(1,55)=0.000	0.993	0.000
Group	**F(3,55)=4.438**	**0.007**	**0.195**
Group × Sex	F(3,55)=1.416	0.248	0.072

524   **Bold** indicates highest interaction or main effect with p<0.05

Key Socially Conditioned ⊶ Control B ⊶ Control C ⊶ Control D

Figure 5. Day 3 social conditioning context fear behavior as time courses and averages.

Time course (A, B, D, E) and average (C, F) context fear behavior data are shown for mice in all

groups in the same sex (top row) and different sex (bottom row) experiments. Respectively,

numbers of mice in top row: males – 8, 7, 7, 8; females – 8, 8, 8, 9 mice. Numbers of mice in

bottom row: males – 8, 7, 7, 8; females – 8, 9, 8, 8. [a]indicates p=0.025, p=0.009 (left to right,

Panel A) between Control B and SC male mice in the same sex experiment at indicated time

points. [b]indicates p=0.020, p=0.021 (left to right, Panel A) between Control C and D male mice

at indicated time points (Panel A). [c]indicates p=0.031, p=0.026, p=0.005 (left to right, Panel B)

between Control C mice of both sexes within the same sex experiment at indicated time points.

[d]indicates p=0.049, p=0.025, p=0.011, p=0.045, p=0.039 (left to right, Panel B) between SC

mice of both sexes within the same sex experiment at indicated time points. [e]indicates p=0.023,

p=0.046, p=0.040, p=0.048, p=0.019 (left to right, Panel D) between Control D and SC male at

indicated time points. [f]indicates p=0.037 between Control C mice of both sexes within the

different sex experiment at indicated time point. [g]indicates p=0.038 between SC mice of both

sexes within the different sex experiment at indicated time point. [h]indicates p=0.006. Data

graphed as mean ± 95% CI.

Serum Corticosterone

No significant differences were detected in log-transformed corticosterone levels in Control A mice

of either experiment (Table 13, Figure 6A).

Table 13. Two-way ANOVA on corticosterone for Control A mice across experiments and

sexes.

Control A Mice – Log-transformed Corticosterone			
	F Statistic	**p value**	**Partial η^2**
Sex	F(1,26)=1.868	0.183	0.067
Experiment	F(1,26)=2.024	0.167	0.072
Sex × Experiment	F(1,26)=0.044	0.836	0.002

552   In parallel, same sex log-transformed corticosterone levels exhibited no interaction nor main

553   effects, and no significant pairwise comparison results (Table 14, Figure 6B).

555   **Table 14.** Two-way repeated measures ANOVAs on corticosterone for mice of both sexes with

556   same sex targets.

Same Sex Experiment – Log-transformed Corticosterone			
	F Statistic	**p value**	**Partial η^2**
Sex	F(1,54)=0.236	0.629	0.004
Group	F(3,54)=1.465	0.234	0.075
Sex × Group	F(3,54)=1.425	0.245	0.073

Different sex log-transformed corticosterone levels, while not exhibiting a significant sex × group

interaction, did display a significant main effect of sex (Table 15). Pairwise comparisons

indicated that this was primarily driven by SC males exhibiting significantly lower cort than SC

females (Figure 6C), the inverse of what was observed for different sex social interaction

(Figure 5C).

Table 15. Two-way repeated measures ANOVAs on corticosterone for mice of both sexes with

different sex targets.

Different Sex Experiment – Log-transformed Corticosterone			
	F Statistic	**p value**	**Partial η^2**
Sex	**F(1,58)=8.683**	**0.005**	**0.130**
Group	F(3,58)=0.787	0.506	0.039
Sex × Group	F(3,58)=1.101	0.356	0.054

566   **Bold** indicates highest interaction or main effect with p<0.05

Control A

Same Sex

Different Sex

Figure 6. *Day 3 corticosterone levels following social conditioning context fear*

expression testing. Log-transformed corticosterone data are shown for mice in Control A

groups (Panel A), and in the same (Panel B) and different (Panel C) sex experiments. Numbers

of mice graphed within each panel, left to right: A) n=8, 8, 7, 7; B) n=8, 7, 7, 7, 8, 8, 8, 9; C) n=8,

8, 8, 9, 8, 9, 8, 8. [a]p=0.004. Data graphed as mean ± 95% CI.

Discussion

Our goal in executing these experiments was to begin establishing a paradigm for inducing

consistent, reproducible socially paired stress in mice of both sexes. While our results indicate

we have not yet fully optimized this paradigm, we nevertheless have discovered sex-specific

patterns of behavioral and physiological stress responsivity that are useful to fields incorporating

sex as a biological variable and/or seeking to employ cross-sex social stressors in mice.

In the absence of any explicit stressor, our Control A mice displayed distinctive responses to

social stimuli depending both upon the sex of the Control A mouse and the sex of their assigned

conspecific. Freezing – ceasing all movement save breathing – is an innate behavior expressed

by mice in response to a real or perceived threat, enabling threat assessment while minimizing

detection (see review (66)). Despite all Control A mice having already encountered both their

assigned conspecific and the social conditioning context a day prior to freezing measurements

during 'acquisition', freezing remained higher in mice of both sexes in the same sex experiment

versus the different sex experiment. Though these freezing levels were considerably less than

the ~80% freezing exhibited by mice after experiencing five mild foot shocks at the conclusion of

acquisition, the discrepancy in basal freezing levels by Control A mice across experiments is

consistent and visible. This observation indicates that, in the presence of a potential sexual

partner, threat assessment is suppressed. The inverse has been demonstrated in female rats;

that is, fear attenuates sexual behaviors, at least in part through amygdala and hypothalamus

estrogen receptor signaling (67). Indirect evidence in male rats housed with females post-

context fear conditioning suggests that encounters with different sexes may suppress threat

assessment. These male rats subsequently exhibited attenuated fear expression, involving

dopamine receptor signaling in the hippocampus (68). However, a different group studying

mice reported that ejaculation by males was required for retention of extinguished social fear

conditioning (47). Continued evaluations are needed to determine whether our observations –

under conditions where copulation is impossible – that male exposure to a female can diminish

threat assessment behaviors are reproducible. An alternative explanation for our findings is that

locomotor activity was elevated, but additional studies would be necessary to determine this.

That only Control A males, but not Control A females, exhibited elevated social interaction time

with female conspecifics versus male conspecifics suggests this possibility is unlikely.

Corticosterone levels in Control A mice were relatively low as expected, given these mice were

never exposed to a mild foot shock. Similarly, neither sex nor experiment affected

corticosterone levels, suggesting no enduring stress was disproportionately experienced after

repeated same- or different-sex conspecific encounters across the four day study timeline.

Despite some interim deviations in fear acquisition across groups receiving mild foot shocks

(Controls B-D; SC mice) in each experiment, all mice ultimately ended their acquisition at similar

freezing levels. Males exhibited greater variability, as indexed by 95% CIs, across groups in

both experiments than females. This aligns with evidence that male rodent data are just as (69–

72) or more variable (54,73,74) than female data. Such relative consistency in ultimate terminal

freezing levels during social conditioning acquisition is important when interpreting subsequent

differences in social engagement, social conditioning context fear, and corticosterone levels in

each experiment; these are not confounded by social conditioning acquisition discrepancies.

Most telling from our social interaction findings are the opposing patterns of reduced social

interaction across sexes in each experiment. In the same sex experiment, socially conditioned

males exhibited reduced social engagement compared to females; in the different sex

experiment, socially conditioned females displayed attenuated social exploration versus males.

These results indicate that encoding of socially aversive experiences depends upon both the

sex of the mouse experiencing the aversive event and upon the sex of the associated

conspecific. Such sex-specific behavior patterns correspond to evidence that accumbal

dopamine signaling dynamics in mice are sex-dependent, both upon the sex of the studied

mouse and the sex of their assigned conspecific (75). Our findings align with the success of

using social defeat to stress male mice. Additionally, they indicate that future studies seeking to

suppress female mice's social interaction behavior through a socially associated stimulus will

meet with more success if using male, rather than female, conspecifics. Indeed, others have

met with challenges when trying to elicit behavioral shifts in female mice, even after 4 weeks of

stress that involved only female conspecifics (76). One group reported that female mice

consistently prefer a socially paired food reinforcer, even if the pairing was with a same sex

conspecific that had just previously undergone acute stress (mild foot shock) (77). Because

males' food preference was not studied, and neither did females encounter stressed different

sex conspecifics, it remains unclear if these effects are sex-specific, and if they would

generalize to different sex interactions. Additionally, future studies will need to determine how

socially conditioning female mice to male conspecifics subsequently affects interactions with

female conspecifics, and vice versa.

Regarding social conditioning context fear exhibited by mice when tested in the absence of any

conspecifics, these did not align with changes in social interaction. For the purposes of

developing a sex-inclusive social conditioning paradigm, this is preferable. Our goal was for

mice to associate the aversive mild foot shock with their assigned conspecific, and any social

buffering of social conditioning context fear could counteract our goal. This is why our use of

conspecifics of a different strain, that were never housed in the same room as all Control and

SC mice, ensured they had never previously been encountered by the mice tested here. Given

the critical contribution of attachment to social buffering (see review (78)), our data indicate we

can be reasonably confident there were no social buffering effects upon social conditioning

context fear in same sex experiment mice. In the different sex experiment, we did find only in

SC males that their freezing was attenuated, but only relative to Control D mice. This could be

related to our postulated threat assessment reduction relating to Control A mice, in that males in

the vicinity of a possible sexual partner could experience impaired encoding of context cues.

Considering corticosterone measurements were taken thirty minutes after social conditioning

context fear testing, we were interested to discover that these levels were inverted with respect

to social interaction in the different sex experiment. In other words, socially conditioned males

exhibited increased interaction with female conspecifics and lower corticosterone levels, as

compared to socially conditioned females displaying reduced interaction with male conspecifics

and elevated corticosterone levels. Further, corticosterone levels did not appear to correspond

to expressions of social conditioning context fear, despite this test being most temporally

proximal to blood collection. Thus, the different sex social conditioning procedure employed

here appears sufficient to elicit both behavioral and physiological responses. However, though

corticosterone levels appear overall elevated in same sex experiment mice, there were not

group- or sex-specific differences. Most striking, corticosterone levels in same sex SC mice

looked nearly identical, meaning this physiological stress marker did not map onto the

reductions in male versus female social interaction behavior in that experiment. This could be

that the same sex experiment's stress induction was not as enduring as that in the different sex

experiment, or it might mean that different sex social conditioning affects more neurobiological

processes than same sex social conditioning.

We may have been overzealous in our inclusion of controls. In our efforts to account for

temporally proximal social encounters with non-cage mates (Controls C and D), we potentially

muddied the waters when it comes to interpreting how social conditioning affects behavior and

circulating corticosterone. That said, we were surprised by what our Control A mice revealed,

and think that this information helps provide context for interpreting our social conditioning

findings. Nevertheless, the overall absence of social interaction differences between Controls

B-D and SC mice (save for different sex females) indicates that this protocol requires additional

optimization to achieve its overarching goal. Evidence for social transmission of stress in

rodents suggests an alternative interpretation (79). Instead, it could be that because the

conspecific for each SC mouse did not experience the same stress (mild foot shock), this in turn

affects the SC mouse's perception of their aversive experience (51), likely in a sex-specific

manner. Even if so, the present findings still generate useful information about environmental

threat assessment in social contexts, and highlight heretofore unrecognized sex-specific social

learning influences under aversive conditions.

Another limitation is that we could not obtain time course corticosterone measurements without

introducing additional confounding stressors to repeatedly sample blood. Daily blood sampling

would reveal if our corticosterone observations were attributable to the initial conditioning day,

the social interaction test, or social conditioning context fear test, and/or if these differences are

transient or persistent. These experiments were all performed in sexually naïve mice, so we

recognize that sexually experienced mice (for example (80)) of both sexes might respond,

behaviorally and physiologically, in a manner distinct from what we report here. Earlier

practices of dividing mice into 'susceptible' and 'resilient' groups following social stress have

more recently, and justifiably, fallen out of favor, given such responses are artificially elicited,

context-specific, and neither phenotype can be universally deemed 'better' or 'worse' (27,81).

We therefore evaluated all SC mice as a single group. Yet, possible bimodal distribution of

social interaction behavior in SC mice looks likely in all but the different sex SC males.

Increasing numbers to establish definitive thresholds with sufficient power for phenotypic

grouping might prove useful, particularly with the same sex setup. Finally, our studies used a

brief protocol spanning only four days, intended for high throughput and maximal comparison to

other social stress literature. We therefore cannot speak to whether these outcomes persist for

weeks or months, timeframes that are ethologically relevant for social learning and memory.

Here, we have reported a novel foray into sex-inclusive efforts to develop a controllable social

stress paradigm in mice. These experiments have provided utilizable insights regarding how

mice of both sexes associate a same- or different-sex conspecific with an aversive experience.

We report that these associations seem strongest in males after same sex encounters, and in

females after different sex encounters. Such distinctive responses are probably evolutionarily

favorable. Given the behaviorally exhibited associations could inversely correspond to

circulating corticosterone levels under different sex conditions, we propose that future studies

focused on socially stressing females utilize male conspecifics for optimal outcomes.

Acknowledgements

This work was supported by R15 MH118705 to AMJ and TLG, and by Kent State University.

We gratefully acknowledge the mice used in this study, the unrivaled veterinary care by Stan

Dannemiller, DVM, MS, DACLAM, and the dedicated work of our vivarium caretakers. Figure 1

was created with BioRender.com (Toronto, ON).

Availability of data

The dataset supporting the conclusions of this article is included with the article as an additional

file, *Supplemental Dataset*.

References

1. Bloomberg L, Meyers J, Braverman MT. The Importance of Social Interaction: A New

Perspective on Social Epidemiology, Social Risk Factors, and Health. Heal Educ Behav.

1994;21(4):447–63.

2. Lee J, Raycraft L, Johnson AW. The dynamic regulation of appetitive behavior through lateral

hypothalamic Orexin and Melanin Concentrating Hormone expressing cells. Physiol Behav.

2020;113234.

3. Monfils MH, Agee LA. Insights from social transmission of information in rodents. Genes,

Brain Behav. 2019;18(1):e12534.

4. Kamps D, Royer J, Dugan E, Kravits T, Gonzalez-Lopez A, Garcia J, et al. Peer Training to

Facilitate Social Interaction for Elementary Students with Autism and Their Peers. Except Child.

2001;68(2):173–87.

5. Leung L. Loneliness, social support, and preference for online social interaction: the

mediating effects of identity experimentation online among children and adolescents. Chin J

Commun. 2011;4(4):381–99.

6. Jasnow AM, Huhman KL. Activation of GABAA receptors in the amygdala blocks the

acquisition and expression of conditioned defeat in Syrian hamsters. Brain Res. 2001;920(1–

2):142–50.

7. Ago Y, Tanaka T, Ota Y, Kitamoto M, Imoto E, Takuma K, et al. Social crowding in the night-

time reduces an anxiety-like behavior and increases social interaction in adolescent mice.

Behav Brain Res. 2014;270:37–46.

8. Wilson CA, Koenig JI. Social interaction and social withdrawal in rodents as readouts for

investigating the negative symptoms of schizophrenia. Eur Neuropsychopharmacol.

2014;24(5):759–73.

9. Venniro M, Zhang M, Caprioli D, Hoots JK, Golden SA, Heins C, et al. Volitional social

interaction prevents drug addiction in rat models. Nat Neurosci. 2018;21(11):1520–9.

10. Association AP. Diagnostic and Statistical Manual of Mental Disorders: DSM-5-TR [Internet].

American Psychiatric Association Publishing; (G - Reference,Information and Interdisciplinary

Subjects Series). Available from: https://books.google.com/books?id=PIGizgEACAAJ

11. Bergamini G, Mechtersheimer J, Azzinnari D, Sigrist H, Buerge M, Dallmann R, et al.

Chronic social stress induces peripheral and central immune activation, blunted mesolimbic

dopamine function, and reduced reward-directed behaviour in mice. Neurobiol Stress.

2018;8:42–56.

12. Caldwell HK, Aulino EA, Rodriguez KM, Witchey SK, Yaw AM. Social Context, Stress,

Neuropsychiatric Disorders, and the Vasopressin 1b Receptor. Front Neurosci. 2017;11:567.

13. Veney SL, Caldwell HK. Neuroendocrinology of Behavior and Emotions, Environmental and

Social Factors Affecting Behavior. Mastercl Neuroendocr. 2024;109–29.

14. Gilman TL, DaMert JP, Meduri JD, Jasnow AM. Grin1 deletion in CRF neurons sex-

dependently enhances fear, sociability, and social stress responsivity. Psychoneuroendocrino.

2015;58:33–45.

15. Garbarino VR, Santos TA, Nelson AR, Zhang WQ, Smolik CM, Javors MA, et al. Prenatal

metformin exposure or organic cation transporter 3 knock-out curbs social interaction

preference in male mice. Pharmacol Res. 2019;140:21–32.

16. Sahani V, Hurd YL, Bachi K. Neuroscience of Social Stress. Curr Top Behav Neurosci.

2022;54:483–515.

17. Venniro M, Marino RAM, Chow JJ, Caprioli D, Epstein DH, Ramsey LA, et al. The Protective

Effect of Social Reward on Opioid and Psychostimulant Reward and Relapse: Behavior,

Pharmacology, and Brain Regions. J Neurosci. 2022;42(50):9298–314.

18. Coccurello R, D'Amato FR, Moles A. Chronic social stress, hedonism and vulnerability to

obesity: Lessons from Rodents. Neurosci Biobehav Rev. 2009;33(4):537–50.

19. Donovan M, Mackey CS, Platt GN, Rounds J, Brown AN, Trickey DJ, et al. Social isolation

alters behavior, the gut-immune-brain axis, and neurochemical circuits in male and female

prairie voles. Neurobiol Stress. 2020;13:100278.

20. Smith AS, Wang Z. Salubrious effects of oxytocin on social stress-induced deficits. Horm

Behav. 2012;61(3):320–30.

21. Kondrakiewicz K, Kostecki M, Szadzińska W, Knapska E. Ecological validity of social

interaction tests in rats and mice. Genes, Brain Behav. 2019;18(1):e12525.

22. Kuske JX, Trainor BC. Neuroscience of Social Stress. Curr Top Behav Neurosci.

2021;54:95–124.

23. Blanchard RJ, McKittrick CR, Blanchard DC. Animal models of social stress: effects on

behavior and brain neurochemical systems. Physiol Behav. 2001;73(3):261–71.

24. Lopez J, Bagot RC. Defining Valid Chronic Stress Models for Depression With Female

Rodents. Biol Psychiatry. 2021;90(4):226–35.

25. Pantoja-Urbán AH, Richer S, Mittermaier A, Giroux M, Nouel D, Hernandez G, et al. Gains

and Losses: Resilience to Social Defeat Stress in Adolescent Female Mice. Biol Psychiatry.

2024;95(1):37–47.

26. Furman O, Tsoory M, Chen A. Differential chronic social stress models in male and female

mice. Eur J Neurosci. 2022;55(9–10):2777–93.

27. Lyons DM, Ayash S, Schatzberg AF, Müller MB. Ecological validity of social defeat stressors

in mouse models of vulnerability and resilience. Neurosci Biobehav Rev. 2023;145:105032.

28. Peña CJ, Nestler EJ, Bagot RC. Environmental Programming of Susceptibility and

Resilience to Stress in Adulthood in Male Mice. Front Behav Neurosci. 2019;13:40.

29. Latsko MS, Farnbauch LA, Gilman TL, Lynch JF, Jasnow AM. Corticosterone may interact

with peripubertal development to shape adult resistance to social defeat. Horm Behav.

2016;82:38–45.

30. McCormick CM, Merrick A, Secen J, Helmreich DL. Social Instability in Adolescence Alters

the Central and Peripheral Hypothalamic-Pituitary-Adrenal Responses to a Repeated

Homotypic Stressor in Male and Female Rats. J Neuroendocr. 2007;19(2):116–26.

31. McCormick CM, Green MR, Cameron NM, Nixon F, Levy MJ, Clark RA. Deficits in male

sexual behavior in adulthood after social instability stress in adolescence in rats. Horm Behav.

2013;63(1):5–12.

32. Sterley TL, Baimoukhametova D, Füzesi T, Zurek AA, Daviu N, Rasiah NP, et al. Social

transmission and buffering of synaptic changes after stress. Nat Neurosci. 2018;21(3):393–403.

33. Bernberg E, Andersson IJ, Gan L ming, Naylor AS, Johansson ME, Bergström G. Effects of

social isolation and environmental enrichment on atherosclerosis in ApoE−/− mice. Stress.

2008;11(5):381–9.

34. Kercmar J, Büdefeld T, Grgurevic N, Tobet SA, Majdic G. Adolescent social isolation

changes social recognition in adult mice. Behav Brain Res. 2011;216(2):647–51.

35. Pan Y, Mou Q, Huang Z, Chen S, Shi Y, Ye M, et al. Chronic social defeat alters behaviors

and neuronal activation in the brain of female Mongolian gerbils. Behav Brain Res.

2023;448:114456.

36. Iñiguez SD, Flores-Ramirez FJ, Riggs LM, Alipio JB, Garcia-Carachure I, Hernandez MA, et

al. Vicarious Social Defeat Stress Induces Depression-Related Outcomes in Female Mice. Biol

Psychiatry. 2018;83(1):9–17.

37. Miyashita T, Yamaguchi T, Motoyama K, Unno K, Nakano Y, Shimoi K. Social stress

increases biopyrrins, oxidative metabolites of bilirubin, in mouse urine. Biochem Biophys Res

Commun. 2006;349(2):775–80.

38. Pizarro JM, Lumley LA, Medina W, Robison CL, Chang WE, Alagappan A, et al. Acute

social defeat reduces neurotrophin expression in brain cortical and subcortical areas in mice.

Brain Res. 2004;1025(1–2):10–20.

39. Jasnow AM, Shi C, Israel JE, Davis M, Huhman KL. Memory of social defeat is facilitated by

cAMP response element-binding protein overexpression in the amygdala. Behav Neurosci.

2005;119(4):1125.

40. Dulka BN, Lynch JF, Latsko MS, Mulvany JL, Jasnow AM. Phenotypic responses to social

defeat are associated with differences in cued and contextual fear discrimination. Behav

Process. 2015;118:115–22.

41. Bonnefil V, Dietz K, Amatruda M, Wentling M, Aubry AV, Dupree JL, et al. Region-specific

myelin differences define behavioral consequences of chronic social defeat stress in mice.

eLife. 2019;8:e40855.

42. Wang W, Liu W, Duan D, Bai H, Wang Z, Xing Y. Chronic Social Defeat Stress Mouse

Model: Current View on Its Behavioral Deficits and Modifications. Behav Neurosci.

2021;135(3):326–35.

43. Golden SA, Covington HE, Berton O, Russo SJ. A standardized protocol for repeated social

defeat stress in mice. Nat Protoc. 2011;6(8):1183–91.

44. Toth I, Neumann ID, Slattery DA. Social Fear Conditioning: A Novel and Specific Animal

Model to Study Social Anxiety Disorder. Neuropsychopharmacology. 2012;37(6):1433–43.

45. Zoicas I, Slattery DA, Neumann ID. Brain Oxytocin in Social Fear Conditioning and Its

Extinction: Involvement of the Lateral Septum. Neuropsychopharmacology. 2014;39(13):3027–

35.

46. Zerda SH de la, Netser S, Magalnik H, Briller M, Marzan D, Glatt S, et al. Social recognition

in laboratory mice requires integration of behaviorally-induced somatosensory, auditory and

olfactory cues. Psychoneuroendocrinology. 2022;143:105859.

47. Grossmann CP, Sommer C, Fahliogullari IB, Neumann ID, Menon R. Mating-induced

release of oxytocin in the mouse lateral septum: Implications for social fear extinction.

Psychoneuroendocrinology. 2024;166:107083.

48. Wiltgen BJ, Sanders MJ, Behne NS, Fanselow MS. Sex Differences, Context Preexposure,

and the Immediate Shock Deficit in Pavlovian Context Conditioning With Mice. Behav Neurosci.

2001;115(1):26–32.

49. Brown KL, Kennard JA, Sherer DJ, Comalli DM, Woodruff-Pak DS. The context preexposure

facilitation effect in mice: A dose–response analysis of pretraining scopolamine administration.

Behav Brain Res. 2011;225(1):290–6.

50. Siegfried B, Frischknecht HR, Riggio G, Waser PG. Preexposure to a Nonaggressive

Opponent Prevents Low-Intensity, Social-Conflict Analgesia in Mice. Behav Neurosci.

1987;101(3):423–6.

51. Guzmán YF, Tronson NC, Guedea A, Huh KH, Gao C, Radulovic J. Social modeling of

conditioned fear in mice by non-fearful conspecifics. Behav Brain Res. 2009;201(1):173–8.

52. Sterley TL, Bains JS. Social communication of affective states. Curr Opin Neurobiol.

2021;68:44–51.

53. Russell W, Burch R. The Principles of Humane Experimental Technique. Nature [Internet].

1959 Nov 1;184(4700):1675–6. Available from: https://doi.org/10.1038/1841675b0

54. Levy DR, Hunter N, Lin S, Robinson EM, Gillis W, Conlin EB, et al. Mouse spontaneous

behavior reflects individual variation rather than estrous state. Curr Biol. 2023;33(7):1358-

1364.e4.

55. Prendergast BJ, Onishi KG, Zucker I. Female mice liberated for inclusion in neuroscience

and biomedical research. Neurosci Biobehav Rev. 2014;40(Endocrinology 153 2012):1–5.

56. Chari T, Griswold S, Andrews NA, Fagiolini M. The Stage of the Estrus Cycle Is Critical for

Interpretation of Female Mouse Social Interaction Behavior. Front Behav Neurosci.

2020;14:113.

57. Bahadır-Varol A, Esen EC, Eren-Koçak E. Repeated collection of vaginal smear causes

stress in mice. Arch Neuropsychiatry. 2022;59(4):325–9.

58. Becegato M, Meurer YSR, Paiva-Santos MA, Lima AC, Marinho GF, Bioni VS, et al.

Impaired discriminative avoidance and increased plasma corticosterone levels induced by

vaginal lavage procedure in rats. Physiol Behav. 2021;232:113343.

59. Wanisch K, Wotjak CT, Engelmann M. Long-lasting second stage of recognition memory

consolidation in mice. Behav Brain Res. 2008;186(2):191–6.

60. Cai DJ, Aharoni D, Shuman T, Shobe J, Biane J, Song W, et al. A shared neural ensemble

links distinct contextual memories encoded close in time. Nature. 2016;534(7605):115–8.

61. Lynch JF, Winiecki P, Gilman TL, Adkins JM, Jasnow AM. Hippocampal GABAB(1a)

Receptors Constrain Generalized Contextual Fear. Neuropsychopharmacol. 2017;42(4):914–

24.

62. Abel T, Lattal KM. Molecular mechanisms of memory acquisition, consolidation and

retrieval. Curr Opin Neurobiol. 2001;11(2):180–7.

63. Sheppard PAS, Koss WA, Frick KM, Choleris E. Rapid actions of oestrogens and their

receptors on memory acquisition and consolidation in females. J Neuroendocr. 2018;30(2).

64. Weber BL, Nicodemus MM, Hite AK, Spalding IR, Beaver JN, Scrimshaw LR, et al.

Heterotypic Stressors Unmask Behavioral Influences of PMAT Deficiency in Mice. Int J Mol Sci.

2023;24(22):16494.

65. Beaver JN, Weber BL, Ford MT, Anello AE, Ruffin KM, Kassis SK, et al. Generalization of

contextual fear is sex-specifically affected by high salt intake. PLOS ONE.

2023;18(7):e0286221.

66. Blanchard DC, Blanchard RJ. Ethoexperimental Approaches to the Biology of Emotion.

Annu Rev Psychol. 1988;39(1):43–68.

67. Moëne OL, Stavarache M, Ogawa S, Musatov S, Ågmo A. Estrogen receptors α and β in the

central amygdala and the ventromedial nucleus of the hypothalamus: Sociosexual behaviors,

fear and arousal in female rats during emotionally challenging events. Behav Brain Res.

2019;367:128–42.

68. Bai H, Cao J, Liu N, Xu L, Luo J. Sexual behavior modulates contextual fear memory

through dopamine D1/D5 receptors. Hippocampus. 2009;19(3):289–98.

69. Becker JB, Prendergast BJ, Liang JW. Female rats are not more variable than male rats: a

meta-analysis of neuroscience studies. Biol Sex Differ. 2016;7(1):34.

70. Smarr B, Rowland NE, Zucker I. Male and female mice show equal variability in food intake

across 4-day spans that encompass estrous cycles. PLoS ONE. 2019;14(7):e0218935.

71. Kazavchinsky L, Dafna A, Einat H. Individual variability in female and male mice in a test-

retest protocol of the forced swim test. J Pharmacol Toxicol Methods. 2019;95:12–5.

72. Beery AK. Inclusion of females does not increase variability in rodent research studies. Curr

Opin Behav Sci. 2018;23:143–9.

73. Smarr BL, Grant AD, Zucker I, Prendergast BJ, Kriegsfeld LJ. Sex differences in variability

across timescales in BALB/c mice. Biol Sex Differ. 2017;8(1):7.

74. Smarr B, Kriegsfeld LJ. Female mice exhibit less overall variance, with a higher proportion

of structured variance, than males at multiple timescales of continuous body temperature and

locomotive activity records. Biol Sex Differ. 2022;13(1):41.

75. Dai B, Sun F, Tong X, Ding Y, Kuang A, Osakada T, et al. Responses and functions of

dopamine in nucleus accumbens core during social behaviors. Cell Rep. 2022;40(8):111246.

76. Díez-Solinska A, Azkona G, Muñoz-Culla M, Beitia-Oyarzabal G, Goñi-Balentziaga O,

Gómez-Lazaro E, et al. The role of sociability in social instability stress: Behavioral,

neuroendocrine and monoaminergic effects. Physiol Behav. 2023;270:114306.

77. Kietzman HW, Trinoskey-Rice G, Blumenthal SA, Guo JD, Gourley SL. Social

incentivization of instrumental choice in mice requires amygdala-prelimbic cortex-nucleus

accumbens connectivity. Nat Commun. 2022;13(1):4768.

78. Kiyokawa Y, Hennessy MB. Comparative studies of social buffering: A consideration of

approaches, terminology, and pitfalls. Neurosci Biobehav Rev. 2018;86:131–41.

79. Brandl HB, Pruessner JC, Farine DR. The social transmission of stress in animal collectives.

Proc Royal Soc B. 2022;289(1974):20212158.

80. Newman EL, Covington HE, Suh J, Bicakci MB, Ressler KJ, DeBold JF, et al. Fighting

Females: Neural and Behavioral Consequences of Social Defeat Stress in Female Mice. Biol

Psychiatry. 2019;86(9):657–68.

81. Koolhaas JM, Boer SF de, Buwalda B, Meerlo P. Social stress models in rodents: Towards

enhanced validity. Neurobiology Stress. 2017;6:104–12.