Control of pili synthesis and putrescine homeostasis in *Escherichia coli*

Running title: *E. coli* surface motility and putrescine homeostasis

Iti Mehta[1#], Jacob Hogins[1#], Sydney Hall[1], Gabrielle Vragel[1], Sankalya Ambagaspitiye[1Δ],

Philippe Zimmern[2], Larry Reitzer[1]*

[1] Department of Biological Sciences, The University of Texas at Dallas, Richardson, Texas

[2] Department of Urology, University of Texas Southwestern Medical Center, Dallas, Texas, USA

[Δ] Current address: Department of Biosystems Technology, Faculty of Technology, University of

Sri Jayewardenepura, Homagama, 10206, Sri Lanka.

*Author for correspondence: Tel: 972-883-2502; Fax: 972-883-2409; reitzer@utdallas.edu

# Authors contributed equally and should be considered co-first authors

Keywords: polyamines, putrescine, surface motility, pili, *Escherichia coli*, putrescine

homeostasis

Summary

Polyamines are biologically ubiquitous cations that bind to nucleic acids, ribosomes, and phospholipids and, thereby, modulate numerous processes, including surface motility in *Escherichia coli*. We characterized the metabolic pathways that contribute to polyamine-dependent control of surface motility in the commonly used strain W3110 and the transcriptome of a mutant lacking a putrescine synthetic pathway that was required for surface motility. Genetic analysis showed that surface motility required type 1 pili, the simultaneous presence of two independent putrescine anabolic pathways, and modulation by putrescine transport and catabolism. An immunological assay for FimA—the major pili subunit, reverse transcription quantitative PCR of *fimA*, and transmission electron microscopy confirmed that pili synthesis required putrescine. Comparative RNAseq analysis of a wild type and Δ*speB* mutant which exhibits impaired pili synthesis showed that the latter had fewer transcripts for pili structural genes and for *fimB* which codes for the phase variation recombinase that orients the *fim* operon promoter in the ON phase, although loss of *speB* did not affect the promoter orientation. Results from the RNAseq analysis also suggested (a) changes in transcripts for several transcription factor genes that affect *fim* operon expression, (b) compensatory mechanisms for low putrescine which implies a putrescine homeostatic network, and (c) decreased transcripts of genes for oxidative energy metabolism and iron transport which a previous genetic analysis suggests may be sufficient to account for the pili defect in putrescine synthesis mutants. We conclude that pili synthesis requires putrescine and putrescine concentration is controlled by a complex homeostatic network that includes the genes of oxidative energy metabolism.

Introduction

Polyamines are flexible aliphatic cations that are found in virtually all organisms. *E. coli* contains putrescine (1,4-diaminobutane) and lesser amounts of spermidine (N-(3-aminopropyl)-1,4-diaminobutane), and cadaverine (1,5-diaminopentane) (1, 2). Both the pathways and enzymes of polyamine synthesis are redundant (Figure 1). Strains that lack eight of these genes do not contain detectable polyamines, grow slowly aerobically, and do not grow anaerobically (3). Polyamines interact with nucleic acids and phospholipids, can affect chromosome and ribosome structure, ribosome-mRNA interactions, and protein and nucleic acid elongation rates (4). Polyamines modulate, but are not required for, protein synthesis (5) and expression of hundreds of genes (6). A major mechanism of polyamine-dependent regulation is translational control of genes for 18 major transcription regulators: four of the seven σ factors – σ^{18} (FecI), σ^{24} (RpoE), σ^{38} (RpoS), and σ^{54} (RpoN); three histone-like DNA-binding proteins (Fis, H-NS, and StpA); and CpxR, Cya, Cra, and UvrY (5, 6).

Polyamines control flagella-dependent surface motility of *Proteus mirabilis* and pili-dependent surface motility (PDSM) of *E. coli*: the latter has been reported to require the polyamine spermidine (7, 8). While reconstructing putrescine-deficient strains to characterize the basis for the anaerobic growth defect, we noticed that a mutant unable to synthesize spermidine moved normally, while mutants lacking enzymes of putrescine synthesis moved poorly. We characterized the role of polyamines on *E. coli* surface motility and pili synthesis by genetic analysis, ELISA assays, RT-qPCR, electron microscopy, and RNAseq. We showed that PDSM and pili synthesis required two independent pathways of putrescine synthesis, but did not require spermidine, and was modulated by putrescine transport and catabolism. Results from an RNAseq analysis are

consistent with multiple mechanisms of putrescine-dependent control, and, unexpectedly, suggests

a putrescine homeostatic network that rewires metabolism to maintain intracellular putrescine.

Results

W3110 surface motility requires pili

An assessment of surface motility for *E. coli* K-12 and eight derivatives showed five strains

covered a plate in 12-18 hours, while four did not traverse a plate in 36 hours (9). The slow-moving

strains generated genetically stable fast-moving variants which suggests that the former were

ancestral. We chose our lab strain of W3110 for further study because (a) it exhibited the least

variability in spreading diameter and generated fewer fast-moving variants, and (b) we had

previously analyzed this strain for the energy requirements for surface motility (10). We refer to

our slow-moving lab strain hereafter as W3110 but not all W3110 strains move similarly: the

W3110 from the genetic stock center is a fast-moving variant (9).

Several results suggested that W3110 requires pili for surface motility. First, deletion of

fimA, which codes for the major type 1 pili subunit, abolished the oscillatory pattern and reduced,

but did not eliminate, surface motility, but deletion of *fliC* which encodes the major flagellum

subunit did not affect surface motility (Figure 2A). Second, electron microscopy confirmed that

W3110 isolated from surface motility plates possessed pili (Figure 2B). None of >500 cells

expressed flagella. A mixed population of elongated (3-4 μm) and non-elongated (<2 μm) cells

were visible, and pili were mostly associated with elongated cells. Finally, W3110 surface motility

requires glucose (10) which should prevent flagella synthesis via inhibition of cyclic AMP

synthesis (11). We confirmed that glucose suppressed swimming motility which implies

suppression of flagella synthesis in W3110 during surface motility assays (Figure 2C). A W3110

derivative with deletions of both *fliC* and *fimA* still exhibited outward movement on a surface

motility plate (Figure 2A), but electron microscopy showed the absence of appendages (Figure

2C). In summary, genetics, electron microscopy, and a glucose requirement show that W3110

surface motility requires pili.

Results from W3110 sequencing found an *IS1* insertion in *fimE* which could explain its

greater movement that is less variable and greater genetic stability. The FimB and FimE

recombinases are the major components of phase variation that determines the orientation of the

promoter that initiates transcription of the *fimAICDFGH* operon which codes for the pili structural

proteins. The FimB recombinase favors the productive ON orientation, while the more active FimE

recombinase favors the unproductive OFF orientation (12-14). The insertion in *fimE* is the likely

basis for the relative stability of W3110.

PDSM required two independent putrescine synthesis pathways.

We examined PDSM requirements for polyamine anabolic pathways. A major putrescine

biosynthetic pathway is ornithine decarboxylation either by the constitutive SpeC or the low-pH

inducible SpeF. The Δ*speC* and Δ*speF* mutants moved as well as the parental strain, but a Δ*speC*

Δ*speF* double mutant moved less well (Figure 3). SpeA (arginine decarboxylase) and SpeB

(agmatinase) catalyze an alternate two-step putrescine synthetic pathway (Figure 1) (15): deletion

of either gene reduced surface motility (Figure 3). The Δ*speA*, Δ*speB*, and Δ*speC* Δ*speF* strains

grew as well as the parental strain (Supplemental Figure 1A) and had normal swimming motility

(Supplemental Figure 1B). A Δ*speE* mutant, which lacks the enzyme for spermidine synthesis,

moved on a surface as well as the parental strain (Figure 3).

Supplemental putrescine at 1 mM restored surface motility of a *speB* mutant (Figure 4A)

and a *speA* mutant (not shown). Spermidine did not restore wild type movement and, also, caused

an unusual pattern of movement in the parental strain (Figure 4A). These exogenous concentrations

of putrescine and spermidine had no effect on growth rates in liquid motility medium (not shown).

In summary, PDSM required the simultaneous presence of two independent pathways of putrescine

synthesis but did not require spermidine.

Loss of putrescine transport systems affected surface motility.

E. coli has several polyamine transport systems and one has been reported to require for

surface motility (16, 17). Mutants lacking the PlaP and PotF putrescine transporters had 50% and

40% reduced surface motility, respectively, and lost the concentric ring pattern (Figure 5A). Loss

of the PotE and PuuP transporters affected neither the movement diameter (Figure 5B) nor

concentric ring formation (Figure 5A). We note that *potF* and *plaP* are more highly expressed than

potE and *puuP* (85, 64, 17, and 6 counts per million transcripts, respectively) for W3110 grown in

liquid motility medium (18). We conclude that extracellular putrescine and its transport contribute

to PDSM.

Putrescine catabolism affected surface motility.

Because putrescine catabolism modulates intracellular putrescine concentrations (19, 20),

we examined PDSM in putrescine catabolic mutants. The major putrescine catabolic pathway is

initiated with putrescine glutamylation (Figure 6A, right column), and a second pathway initiates

with putrescine deamination (Figure 6A, left column) (20). Deletion of either *puuA* or *patA* which

code for the first enzymes of their respective pathways did not affect surface motility (Figure 6B).

A double mutant moved less well which was unexpected because the proposed higher intracellular

putrescine should have stimulated PDSM. Consistent with the stimulatory effect of putrescine, loss

of *puuR* which codes for the repressor of PuuA-initiated pathway genes (20), impaired PDSM

(Figure 6B). One possible explanation to reconcile these seemingly contradictory results is that an

optimal polyamine concentration stimulates PDSM, and a high concentration is inhibitory. The

next section describes experiments to test this possibility.

High putrescine reduced expression of pili genes.

High putrescine inhibits translation of some mRNAs, and high spermidine, a product of

putrescine metabolism, inhibits growth (21, 22). If the phenotype of the Δ*patA* Δ*puuA* double

mutant results from spermidine toxicity, then loss of spermidine synthase (SpeE) should reverse

the phenotype. This prediction was not met which argues against spermidine toxicity (Figure

6C). For the *speB* mutant, 1 mM putrescine stimulated PDSM, and 4 mM putrescine was slightly

inhibitory (Figure 7A and B). The concentric ring pattern was observed with 1 mM, but not 4

mM putrescine. RT-qPCR to test transcriptional regulation of pili expression by putrescine

showed optimal *fimA* transcription at 1 mM putrescine in the *speB* mutant (Figure 7C). Indirect

enzyme linked immunosorbent assays (ELISAs) against FimA suggested that the level of FimA

in the *speB* mutant at 1 mM putrescine was higher than with 4 mM putrescine, but the difference

was not statistically significant (Figure 7D). Variation of supplemental putrescine did not affect

FimA in the parental strain (Figure 7C and 7D). Transmission electron microscopy (TEM)

showed that pili expression was highest for the *speB* mutant at 1 mM putrescine (Figure 8),

moderate at 0.1- and 4-mM putrescine, and undetectable without putrescine. The cells grown

with 4 mM putrescine were shorter and thinner, which suggests stress, possibly mimicking

osmotic stress. From these results and those in the previous section, we conclude that pili

expression required an optimal putrescine concentration.

RNAseq analysis identifies *fim* operon expression and energy metabolism as targets of

putrescine control.

RNAseq analysis was used to further confirm or determine whether putrescine affected

(a) transcription of the *fim* operon that codes for the pili structural genes, (b) *fim* operon promoter

orientation, i.e. phase variation, or (c) energy metabolism: surface motility of W3110 has been

shown to require glucose metabolism and oxidative phosphorylation (10).

RNAseq analysis. We grew parental W3110 and its Δ*speB* derivative with and without 1

mM putrescine. R^2 values of pairwise comparisons showed that the transcriptome of the *speB*

mutant grown without putrescine differed from the other transcriptomes which were similar

(Table 1 and Supplemental Figure 2). A multidimensional scaling plot and a plot of the 100 most

variable genes visualize these comparisons (Figure 9A and B). When compared to the parental

strain, the *speB* mutant grown without putrescine had 310 downregulated genes and 159

upregulated genes with at least four-fold differential expression and FDR < 0.05. Transcripts for

the putrescine-induced *puuAP* and *puuDRCBE* operons, which specify genes of the major

putrescine catabolic pathway, were reduced from 1.6- to 14-fold fold (FDR ≤ 0.02) in the *speB*

mutant (Supplemental Table 1). Because higher intracellular putrescine results in higher

catabolic gene expression, these results imply lower intracellular putrescine in the Δ*speB* mutant.

Transcription of the *fim* operon is reduced in the Δ*speB* mutant. Deletion of *speB* reduced

transcripts for genes of the *fimA* operon (Figure 10A). Numerous transcription factors activate

the *fim* operon (23, 24), and, of these, loss of *speB* reduced *hns* transcripts, increased *fis*, *lrp*, and

qseB transcripts, but had no effect on *ihfA* and *ihfB* (Figure 10B). Loss of *hns*, *ihfA*, and *lrp*

impaired PDSM (Figure 10C and D), which confirms the requirement for their products for pili

synthesis in our strain background.

Putrescine does not control *fim* operon phase variation in W3110. The phase ON-favoring

FimB and phase OFF-favoring FimE recombinases control the orientation of the *fim* operon

promoter. Our version of W3110 has an insertion in *fimE* which means that FimB is the major

factor that controls phase variation. Loss of *speB* reduced transcripts from *fimB* which could

account for loss of motility. Using specific primers, PCR analysis showed that loss of *speB*, loss

of *hns*, or the presence of putrescine did not alter the ratio of ON to OFF (Supplemental Figure

3). This experiment determines the steady state phase ON to OFF ratio, which less FimB might

not affect if FimB is in excess. In strains with FimE, the rate of phase ON orientation will be

much slower and even small changes in FimB activity could conceivably affect the switch

orientation. We conclude that putrescine does not control phase variation, at least in our strain of

W3110.

Evidence for a putrescine homeostatic network. Altered transcript levels in the *speB*

mutant suggest compensatory mechanisms for low putrescine (Supplemental Table 1 for

transcript differences for all genes, Table 2 for selected genes, and Figure 11 for a diagrammatic

representation of transcript differences). The *speB* mutant had more transcripts from (a) genes of

arginine and ornithine carboxylase, (b) all genes of ornithine and arginine synthesis from

glutamate, and (c) genes for three separate putrescine transport systems; and fewer transcripts

from the *sap* operon which codes for a putrescine exporter (25) and *rpmH* and *rpsT*, whose

products inhibit ornithine and arginine decarboxylase (26). The net effect of these changes

should be to increase intracellular putrescine (solid arrows in Figure 11). In other words, these

changes are compensatory for low putrescine. In addition to these changes, the *speB* mutant had

60-fold and 7-fold more transcripts for *mgtA* and *phoBR* which code for the major magnesium

transporter and the regulators of phosphate assimilation, respectively. These changes may also be

compensatory and are discussed below.

<u>Putrescine affects energy metabolism which could account for loss of motility.</u> Loss of

speB affected transcripts for genes of the major energy metabolism pathways (Table 2): fewer

transcripts for genes coding for all citric acid cycle enzymes, NADH dehydrogenase I (the entire

nuo operon), cytochrome oxidase (the *cyo* operon), and several genes of menaquinone (the

menFDHBCE operon), ubiquinone, and heme synthesis. The *speB* mutant also had fewer

transcripts for all genes of iron acquisition, which is consistent with diminished oxidative energy

generation. A *speB* mutant had fewer transcripts for *ptsH* (Hpr of the phosphotransferase system

of carbohydrate transport), *ptsG* (enzyme IIBC component of glucose transport), *gltA* (citrate

synthase), *sdhA* (a succinate dehydrogenase subunit), and *sucC* (a succinyl-CoA synthetase

subunit), and their loss has been shown to either impair or eliminate W3110 surface motility

(10). The cumulative effect of these differences could account for the motility defect in the *speB*

mutant.

Discussion

Several aspects of polyamine metabolism affect PDSM in *E. coli*. Strains with loss of one

or two polyamine synthesis genes — *speA* and *speB* single mutants and a *speC speF* double mutant

— resulted in PDSM defects. This observation is remarkable because nine enzymes and several

pathways are involved in polyamine synthesis. Our putrescine synthesis mutants had no growth

defect which is not unexpected since deletions of eight biosynthetic genes were required to show

a strong phenotype: impaired aerobic growth and elimination of anaerobic growth (3). Strains with

defects in the two most active putrescine transport systems had altered PDSM. The specific defects

suggest the involvement of extracellular putrescine and arginine in intracellular putrescine

homeostasis. The PDSM defect of putrescine transport mutants not only implicates extracellular

putrescine but also suggests secretion of putrescine which is not added to the medium. The PDSM

defect of *speA* and *speB* mutants implicates extracellular arginine because SpeA is mostly

periplasmic (27). Defective putrescine catabolism also affected PDSM. The role of both putrescine

synthesis and degradation suggests rapid adjustments to fluctuating putrescine concentrations and

environmental conditions: putrescine catabolism is a core response to several stresses (19).

Results from an RNAseq analysis showed that transcription of the *fim* operon was reduced

in the *speB* mutant. Of the known transcription factors that control the *fim* operon, only the gene

for H-NS had fewer transcripts. In contrast, transcripts for other relevant transcription factor genes

were elevated in the *speB* mutant. Our results are consistent with H-NS mediating putrescine-

dependent control, but we cannot exclude a general regulatory dysfunction mediated by altered

nucleoprotein complexes at the *fim* operon promoter. The elevated transcripts for some

transcription factors may be inhibitory.

The transcriptomic results showed that the *speB* mutant had more transcripts for genes

whose products will maintain intracellular putrescine, which suggests compensatory mechanisms

for low intracellular putrescine (Figure 11). The *speB* mutant also had more transcripts for

magnesium and phosphate transport genes. The elevated transcripts for magnesium transport

genes could compensate for low putrescine to the extent that magnesium can directly replace

putrescine, e.g., binding to nucleotides and RNA (28). Two recent studies observe an inverse

correlation between intracellular magnesium and polyamines in *Salmonella* and proposed that

either magnesium or the polyamines maintained an overall divalent cation homeostasis (29, 30).

The rationale for an increase in anionic phosphate assimilation is less clear, although the major

phosphate permease transports metal-phosphates, including magnesium-phosphate (31), which

could help to compensate for low putrescine. Regardless of the rationale, the substantial changes

in gene expression due to low putrescine suggest multiple compensatory mechanisms.

A major observation of the RNAseq analysis is that loss of *speB* lowered transcripts for

many genes of energy metabolism and all genes of iron acquisition. This gene expression pattern

suggests that low putrescine diverts metabolism away from energy generation and toward

putrescine synthesis, i.e., a prioritization of putrescine synthesis over aerobic energy metabolism.

Oxidative energy metabolism and putrescine synthesis compete for α-ketoglutarate and the

metabolic diversion could also be a potential compensatory mechanism for low putrescine.

Another way of looking at this relationship is that intracellular putrescine positively correlates

with oxidative energy metabolism. Deletion of *ptsH*, *ptsG*, *gltA*, *sdhA*, and *sucC* have been

shown to impair W3110 PDSM which suggests that a major, possibly the primary, effect of low

putrescine is reduced energy generation.

In contrast to our results, a thorough and well-designed study showed that *E. coli* PDSM

requires spermidine instead of putrescine (8). The differences between the studies were strain

backgrounds, media (glucose-LB vs glucose-tryptone), and incubation temperature (33º vs 37º).

We found PDSM results for 37º incubations were highly variable and resulted in numerous fast-

moving variants. Since growth at the higher temperature is faster, and faster growth is associated

with higher putrescine (32), we propose that growth at 37º results in an inhibitory putrescine

concentration. In this case, spermidine would be stimulatory because of its known inhibition of the

putrescine biosynthetic enzymes SpeA and SpeC (33, 34) and subsequent reduction of intracellular

putrescine to an optimal concentration. Regardless of the explanation, strains differences are not

unexpected because of decades of strain passage for the strains employed with resulting laboratory

evolution that can affect any of the numerous factors that control the intracellular putrescine

concentration.

The transcript changes observed in our study were different from those in the only other

systematic study of putrescine's effect on *E. coli* gene expression (6). Both studies had almost

identical experimental designs: 1.0- or 1.14-mM putrescine was added to a *speB* or *speB speC*

mutant, respectively. Differences in strains, growth media (minimal media vs tryptone-containing

media), and growth temperature (33° vs 37°) could account for the discrepancy in results. The most

crucial difference could be the growth media: Igarashi and colleagues grew bacteria in a minimal

medium, while we grew bacteria in an amino acid-containing medium that was required for surface

motility. We propose that the slower growth in minimal medium, possibly correlated with higher

guanosine tetraphosphate which is associated with slower growth, counters and obscures many

growth-stimulatory putrescine effects. In contrast, the growth stimulation from the amino acids in

our medium would not counter putrescine effects and would essentially allow detection of effects

caused solely by changes in putrescine concentrations. Regardless of the explanation, common

results can be interpreted as core responses to putrescine limitation. Both studies observed that low

putrescine increased transcripts for genes of arginine synthesis and magnesium transport, and

reduced transcripts for the *sdhCDAB-sucABCD* operon which codes for three TCA cycle enzymes.

The link between putrescine and arginine synthesis is probably mediated by the ArgR repressor

because *argR* transcripts are 2.5-fold lower (FDR = 0.0004) in the *speB* mutant (Supplemental

Table 1).

Several results implicate putrescine in *E. coli* virulence during urinary tract infections.

First, pili-dependent binding to the urothelium is an essential step for uropathogenic *E. coli* (35).

Second, putrescine is present in urine from infected, but not healthy, individuals (36, 37). Third,

urinary putrescine can result from uropathogenic *E. coli* growth in urine (37) and urothelial

pathophysiology (38). Putrescine's possible role in virulence is seemingly contradicted by the

observation that a *speB* mutant outcompetes the parental CFT073 in the bladder of mice, i.e., less

putrescine synthesis enhanced virulence (39). A majority of uropathogenic *E. coli* are associated

with phylogenetic group B2—including CFT073, whereas most *E. coli* lab strains, such as W3110,

are group A. Group B2 strains lack genes for the major putrescine catabolic pathway and have

significantly higher transcripts for genes coding for SpeA and SpeB (18). Both observations

suggest that the cytoplasmic putrescine concentration is higher in group B2 strains. We propose

that the putrescine concentration in group B2 strains is normally in the inhibitory range for pili

synthesis, and that loss of SpeB reduces putrescine to a level that stimulates pili synthesis.

Regardless of the explanation, putrescine control of pili synthesis and other processes in

uropathogenic *E. coli* are worth examination.

Experimental Procedures

Strains. All strains used for growth rate determinations and motility assays were derivatives of *E.*

coli K-12 strain W3110 and are listed in Table 3. Great variations exist in standard lab strains,

including W3110, from different labs. We initially tested nine strains derived from *E. coli* K-12,

mostly from the Coli Genetic Stock Center (9), and chose our lab strain because of relative genetic

stability and more quantitatively reproducible results. Our strain originated from the lab of Jon

Beckwith in the 1960s via the lab of Boris Magasanik where it had been stored in a room

temperature stab until the mid-1980s, at which time it was revived and frozen at -80⁰ C. To

construct the mutant strains, the altered allele was obtained from the Keio collection in which the

gene of interest had been deleted and replaced with an antibiotic resistance gene (40). The marked

deletion allele was transferred into W3110 by P1 transduction (41). The antibiotic gene was

removed as described which generated an in-frame deletion (42).

Media and Growth Conditions. For P1 transductions and plasmid transformation experiments,

cells were grown in standard LB liquid medium (1% tryptone, 1% NaCl and 0.5% yeast extract,

pH 7.0) at 37°C. Antibiotics were used for selection at concentrations of 25 µg/mL (both

chloramphenicol and kanamycin). For growth analysis, bacteria were grown in liquid motility

medium (0.5% glucose, 1% tryptone, and 0.25% NaCl) which we refer to as GT medium. Starter

cultures (typically 6–12 h incubation) were grown in GT medium, harvested by centrifugation,

washed twice with phosphate buffered saline and re-suspended in GT medium before inoculation.

The cells were then grown at 37° C in aerobic conditions (shaking at 240 rpm) and the turbidity

was measured every thirty minutes. Cell growth was measured in Klett units using a Scienceware

Klett colorimeter with a KS-54 filter. 100 Klett units represent an OD_{600} value of about 0.7.

Surface motility assay. For a standard surface motility assay, single colonies from fresh plates

(streaked out from frozen stocks a day before) were inoculated in GT medium for six hours at

37° C in aerobic conditions (shaking at 240 rpm). 30 ml of autoclaved GT medium with 0.45%

agar (Eiken, Tokyo, Japan) was poured into sterile polystyrene petri-dish (100 mm X 15 mm)

and allowed to solidify at room temperature for approximately six hours. Then, 1 µL of the pre-

motility growth medium was inoculated at the center of the agar plate and incubated at 33° C for

36 hrs. Each experiment was performed in triplicate and pictures were taken after 36 hours.

Surface motility was extremely sensitive to humidity. Opening the incubator before 36 hours

resulted in movement cessation. The surface motility assay was performed at 33⁰ C because

results from incubations at 37⁰ C were highly variable and the cultures more frequently generated

genetically stable fast-moving variants.

Swim assay. The media and culturing for swimming motility is identical to that for surface

motility, except that the plates contained 0.25% agar and were solidified at room temperature for

about one hour before inoculation. Then, 1 µL of the culture was stabbed in the center of this

swim agar plate and incubated at 33° C for 20 hours. Each experiment was performed in

triplicate.

Transmission electron microscopy. Cells from surface motility plates were collected and fixed

with 2.5% glutaraldehyde. Bacteria were absorbed onto Foamvar carbon-coated copper grids for

1 min. Grids were washed with distilled water and stained with 1% phosphotungstic acid for 30

s. Washed and stained grids were dried at 37⁰C for 10 minutes. Samples were viewed on a JEOL

1200 EX transmission electron microscope at University of Texas Southwestern Medical Center

(Figure 2) and University of Texas at Dallas (Figure 8).

ELISA analysis. Cells were grown overnight in 5 mL GT media followed by a 2-hour growth in

GT media with 0.0, 0.1, 1.0, or 4.0 mM putrescine. Cells were then lysed using a 24-gauge

needle. The cell lysate was diluted to a concentration of 1 mg/mL protein (based on A280) in pH

9.6 coating buffer (3 g Na_2CO_3, 6 g $NaHCO_3$, 1000 mL distilled water) and then coated onto the

walls of a 96-well plate by incubating overnight at 4°C. The wells were rinsed 3 times with PBS

and further blocked using coating buffer with 1% bovine albumin overnight at 4°C. The

following day, wells were rinsed 3 times with PBS, and primary antibodies to FimA or RpoD

were added following supplier directions. After a 2-hour room temperature incubation, secondary

antibodies conjugated to HRP were added and incubated for a further two hours. HRP was

activated using the TMB substrate kit (Thermo-34021) following the provided protocol.

Expression was read on a BioTek plate reader based on the provided kit protocol. Data was

blanked to wells only treated with bovine albumin then normalized to RpoD. Data was generated

from technical and biological triplicates.

DNA isolation and genome assembly and annotation for W3110. DNA was isolated based on

previous established protocols (43). Short reads were sequenced at the SeqCenter (Pittsburgh,

Pennsylvania) using Illumina tagmentation-based and PCR-based DNA prep and custom IDT

indices targeting inserts of 280 bp without further fragmentation or size selection steps. The

Illumina NovaSeq X Plus sequencer was ran producing 2X151 paired-end reads.

Demultiplexing, quality control, and adapter trimmer was performed with bcl-convert (v4.2.4).

Total Illumina Reads (R1+R2): 4059078 with 553087045 bps >Q30.

Long reads were prepared using the PCR-free Oxford Nanopore Technologies ligation

sequencing kit with the NEBNext companion module. No fragmentation or size selection was

performed. Long read libraries were performed using R10.4.1 flow cell. The 400bps sequencing

mode with a minimum read length of 200bp were selected. Gupply (v6.5.7) was used at the

super-accurate basecalling, demultiplexing, and adapter removal. Total long reads: 53773 with

89.607 % of bps > Q20.

To generate the completed genome, porechop (v0.2.4) was used to trim residual adapter

sequences from long reads. Flye (v2.9.2) was used to generate the *de novo* assembly under the

nano-hq model. Reads longer than the estimated N_{50} based on a genome size of 6Mbp initiated

the assembly. Subsequent polishing using the short read data was performed using Pilon (v1.24)

under default parameters. Long read contigs with an average short read coverage of 15X or less

were removed from the assembly. The assembled contig was confirmed to be circular via

circulator (v1.5.5). Annotation was performed using prokka (v 1.14.6). Finally, statistics were

recorded using QUAST (v5.2.0). The final genome contained 1 contig of 4750347 bp with a

sequencing depth of 123.92x. The N_{50} was 4750347. This genome can be accessed via the

accession number CP165600 or via the BioProject PRJNA1142534 via NCBI. The completed

genome has an estimated average nucleotide identity of 99.9694 to the W3110 genome deposited

to NCBI (genome assembly ASM1024v1).

RNA isolation and quality control. Cells were grown for 2 hours in 1 mL GT media. 60 µL

were then added to 1 mL of fresh GT, with or without 1 mM putrescine, and grown for another

two hours. After growth, the cells were centrifuged, and frozen at -80ºC. The isolation and

analysis protocol has been previously published (18). Cell pellets were thawed, resuspended in

0.7 mL of buffer RLT (Qiagen RNeasy kit), and mechanically lysed using a bead beater

(FastPrep-24 Classic from MP Biomedical) set at the highest setting for three 45-s cycles with a

5-min rest period on ice between cycles. Cell lysates were used to isolate RNA using a Qiagen

RNeasy mini kit. RNA was quantified, DNase treated, and re-quantified on a nanodrop. RNA

that passed the first quality check was analyzed on RNA-IQ (Qubit). RNA with an IQ higher

than 7.5 was used. RNA that passed both checks were run on an agarose gel to check for RNA

integrity. RT-qPCR was used to ensure there was no DNA contamination. RNA libraries that

passed all these quality checks were submitted to the genomics core facility at The University of

Texas at Dallas for RNA sequencing. The core performed rRNA removal (RiboMinus

Transcriptome Isolation Kit or RiboCop bacterial rRNA depletion—Lexogen), library

preparation (Stranded Total RNA Prep—Illumina), and single-end Illumina sequencing.

Reverse transcription quantitative PCR. RNA was isolated as described above. One

microgram of RNA was reverse transcribed using LunaScript RT Super Mix (NEB E3010) and

another microgram was subjected to the same reaction as the reverse transcribed RNA, except

without reverse transcriptase (negative control) following manufacturer instructions. The cDNA

was then diluted to 10 ng/µL for qPCR. PCR reactions were composed of 10 ng cDNA, 8 µL of

nuclease free water, 10 nM primers (rpoD: Forward: TCGTGTTGAAGCAGAAGAAGCG;

Reverse: TCGTCATCGCCATCTTCTTCG) (fimA: Forward: ATGGTGGGACCGTTCACTTT;

Reverse: GGCAACAGCGGCTTTAGATG), and PowerUp SYBR Green 2X master mix for

qPCR (ThermoFisher A25777) in a 20 µL reaction. A Quantstudio 3 qPCR machine was used to

generate critical threshold (Ct) values. The Ct values were then analyzed using the $2^{-\Delta\Delta C_T}$ method

(44) using *rpoD* as a library control.

RNAseq Analysis. Transcripts were aligned to the W3110 genome and CDS (CP165600)

available on NCBI using CLC genomics workbench (Qiagen version 22.00) for RNA-seq

analysis. The gene expression counts generated as this output were used to perform differentially

expressed gene (DEG) analysis using EdgeR (version 3.38.4) to analyze read counts (45, 46).

DEGs had an adjusted P-value less than 0.05.

Data availability: The raw RNA seq reads were deposited in GenBank: BioProject accession

number PRJNA1126736. The genome was deposited in GenBank: BioProject accession number

PRJNA1142534. The genome accession number is CP165600.

Acknowledgements: This work was funded through private donations. PEZ received

endowment support from the Cain Foundation through the Felecia and John Cain Distinguished

Chair in Women's Health in honor of Philippe E. Zimmern, MD. The authors declare no potential

conflicts of interests.

The authors (JH, SH, GV, and LR) would like to thank Prof. Kelli Palmer of The

University of Texas at Dallas for the use of her space, supplies, and tissue homogenizer. JH

would like to further thank Prof. Palmer for the use of her CLC genomics program. The authors

would like to thank the Genome Center at The University of Texas at Dallas for the services to

support our research. The authors would like to thank the Olympus Discovery Center/Imaging

Core facility at UT Dallas for providing equipment and support (Figure 8), and the UT

Southwestern Medical Center for electron microscopy (Figure 2).

References:

1. Tabor CW, Tabor H. 1984. Polyamines. Annual review of biochemistry 53:749-90.
2. Tabor CW, Tabor H. 1984. Polyamines. Annu Rev Biochem 53:749-90.

3. Chattopadhyay MK, Tabor CW, Tabor H. 2009. Polyamines are not required for aerobic growth of
Escherichia coli: preparation of a strain with deletions in all of the genes for polyamine
biosynthesis. J Bacteriol 191:5549-52.
4. Igarashi K, Kashiwagi K. 2000. Polyamines: mysterious modulators of cellular functions. Biochem
Biophys Res Commun 271:559-64.
5. Igarashi K, Kashiwagi K. 2015. Modulation of protein synthesis by polyamines. IUBMB Life
67:160-9.
6. Yoshida M, Kashiwagi K, Shigemasa A, Taniguchi S, Yamamoto K, Makinoshima H, Ishihama A,
Igarashi K. 2004. A unifying model for the role of polyamines in bacterial cell growth, the
polyamine modulon. J Biol Chem 279:46008-13.
7. Armbruster CE, Hodges SA, Mobley HL. 2013. Initiation of swarming motility by *Proteus mirabilis*
occurs in response to specific cues present in urine and requires excess L-glutamine. J Bacteriol
195:1305-19.
8. Kurihara S, Suzuki H, Tsuboi Y, Benno Y. 2009. Dependence of swarming in *Escherichia coli* K-12
on spermidine and the spermidine importer. FEMS Microbiol Lett 294:97-101.
9. Ambagaspitiye S, Sudarshan S, Hogins J, McDill P, De Nisco NJ, Zimmern PE, Reitzer L. 2019.
Fimbriae and flagella mediated surface motility and the effect of glucose on nonpathogenic and
uropathogenic *Escherichia coli* BioRxiv doi.org:10.1101/840991
10. Sudarshan S, Hogins J, Ambagaspitiye S, Zimmern P, Reitzer L. 2021. The nutrient and energy
pathway requirements for surface motility of nonpathogenic and uropathogenic *Escherichia coli*.
J Bacteriol. doi:10.1128/jb.00467-20doi:10.1128/jb.00467-20.
11. Yokota T, Gots JS. 1970. Requirement of adenosine 3', 5'-cyclic phosphate for flagella formation
in *Escherichia* coli and *Salmonella typhimurium*. J Bacteriol 103:513-6.
12. Conway C, Beckett MC, Dorman CJ. 2023. The DNA relaxation-dependent OFF-to-ON biasing of
the type 1 fimbrial genetic switch requires the Fis nucleoid-associated protein. Microbiology
(Reading). 169(1):doi:10.1099/mic.0.001283.
13. Kelly A, Conway C, T OC, Smith SG, Dorman CJ. 2006. DNA supercoiling and the Lrp protein
determine the directionality of fim switch DNA inversion in *Escherichia coli* K-12. J Bacteriol
188:5356-63.
14. O'Gara JP, Dorman CJ. 2000. Effects of local transcription and H-NS on inversion of the fim switch
of *Escherichia coli*. Mol Microbiol 36:457-66.
15. Tabor CW, Tabor H. 1985. Polyamines in microorganisms. Microbiol Rev 49:81-99.
16. Igarashi K, Ito K, Kashiwagi K. 2001. Polyamine uptake systems in *Escherichia coli*. Res Microbiol
152:271-8.
17. Kurihara S, Suzuki H, Oshida M, Benno Y. 2011. A novel putrescine importer required for type 1
pili-driven surface motility induced by extracellular putrescine in *Escherichia coli* K-12. J Biol
Chem 286:10185-92.
18. Hogins J, Xuan Z, Zimmern PE, Reitzer L. 2023. The distinct transcriptome of virulence-associated
phylogenetic group B2 *Escherichia coli*. Microbiol Spectr. doi:10.1128/spectrum.02085-
23e0208523. doi:10.1128/spectrum.02085-23.
19. Schneider BL, Hernandez VJ, Reitzer L. 2013. Putrescine catabolism is a metabolic response to
several stresses in *Escherichia coli*. Mol Microbiol 88:537-50.
20. Schneider BL, Reitzer L. 2012. Pathway and enzyme redundancy in putrescine catabolism in
Escherichia coli. J Bacteriol 194:4080-8.
21. Fukuchi J, Kashiwagi K, Yamagishi M, Ishihama A, Igarashi K. 1995. Decrease in cell viability due
to the accumulation of spermidine in spermidine acetyltransferase-deficient mutant of
Escherichia coli. J Biol Chem 270:18831-5.

22. Sakamoto A, Sahara J, Kawai G, Yamamoto K, Ishihama A, Uemura T, Igarashi K, Kashiwagi K,
Terui Y. 2020. Cytotoxic mechanism of excess polyamines functions through translational
repression of specific proteins encoded by polyamine modulon. Int J Mol Sci.
21(7):doi:10.3390/ijms21072406.
23. Schwan WR. 2011. Regulation of *fim* genes in uropathogenic *Escherichia coli*. World J Clin Infect
Dis 1:17-25.
24. Karp PD, Ong WK, Paley S, Billington R, Caspi R, Fulcher C, Kothari A, Krummenacker M,
Latendresse M, Midford PE, Subhraveti P, Gama-Castro S, Muniz-Rascado L, Bonavides-Martinez
C, Santos-Zavaleta A, Mackie A, Collado-Vides J, Keseler IM, Paulsen I. 2018. The EcoCyc
Database. EcoSal Plus. 8(1):doi:10.1128/ecosalplus.ESP-0006-2018.
25. Sugiyama Y, Nakamura A, Matsumoto M, Kanbe A, Sakanaka M, Higashi K, Igarashi K, Katayama T,
Suzuki H, Kurihara S. 2016. A novel putrescine exporter SapBCDF of *Escherichia coli*. J Biol Chem
291:26343-26351.
26. Keseler IM, Mackie A, Santos-Zavaleta A, Billington R, Bonavides-Martínez C, Caspi R, Fulcher C,
Gama-Castro S, Kothari A, Krummenacker M, Latendresse M, Muñiz-Rascado L, Ong Q, Paley S,
Peralta-Gil M, Subhraveti P, Velázquez-Ramírez DA, Weaver D, Collado-Vides J, Paulsen I, Karp PD.
2016. The EcoCyc database: reflecting new knowledge about *Escherichia coli* K-12. Nucleic Acids
Research 45:D543-D550.
27. Tabor H, Tabor CW. 1969. Partial separation of two pools of arginine in *Escherichia coli*;
preferential use of exogenous rather than endogenous arginine for the biosynthesis of 1,4-
diaminobutane. J Biol Chem 244:6383-7.
28. Miyamoto S, Kashiwagi K, Ito K, Watanabe S, Igarashi K. 1993. Estimation of polyamine
distribution and polyamine stimulation of protein synthesis in *Escherichia coli*. Arch Biochem
Biophys 300:63-8.
29. Iwadate Y, Golubeva YA, Slauch JM. 2023. Cation homeostasis: Coordinate regulation of
polyamine and magnesium levels in *Salmonella*. mBio 14:e0269822.
30. Duprey A, Groisman EA. 2020. DNA supercoiling differences in bacteria result from disparate
DNA gyrase activation by polyamines. PLoS Genet 16:e1009085.
31. van Veen HW, Abee T, Kortstee GJ, Konings WN, Zehnder AJ. 1994. Translocation of metal
phosphate via the phosphate inorganic transport system of *Escherichia coli*. Biochemistry
33:1766-70.
32. Tweeddale H, Notley-McRobb L, Ferenci T. 1998. Effect of slow growth on metabolism of
Escherichia coli, as revealed by global metabolite pool ("metabolome") analysis. J Bacteriol
180:5109-16.
33. Applebaum DM, Dunlap JC, Morris DR. 1977. Comparison of the biosynthetic and biodegradative
ornithine decarboxylases of *Escherichia coli*. Biochemistry 16:1580-4.
34. Wu WH, Morris DR. 1973. Biosynthetic arginine decarboxylase from *Escherichia coli*. Purification
and properties. J Biol Chem 248:1687-95.
35. Flores-Mireles AL, Walker JN, Caparon M, Hultgren SJ. 2015. Urinary tract infections:
epidemiology, mechanisms of infection and treatment options. Nat Rev Microbiol 13:269-84.
36. Puebla-Barragan S, Renaud J, Sumarah M, Reid G. 2020. Malodorous biogenic amines in
Escherichia coli-caused urinary tract infections in women-a metabolomics approach. Sci Rep
10:9703.
37. Satink HP, Hessels J, Kingma AW, van den Berg GA, Muskiet FA, Halie MR. 1989. Microbial
influences on urinary polyamine excretion. Clin Chim Acta 179:305-14.
38. Keay SK, Birder LA, Chai TC. 2014. Evidence for bladder urothelial pathophysiology in functional
bladder disorders. Biomed Res Int 2014:865463.

39. Alteri CJ, Smith SN, Mobley HL. 2009. Fitness of *Escherichia coli* during urinary tract infection
requires gluconeogenesis and the TCA cycle. PLoS Pathog 5:e1000448.
40. Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M, Wanner BL,
Mori H. 2006. Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the
Keio collection. Mol Syst Biol 2:2006 0008.
41. Miller JH. 1972. Experiments in Molecular Genetics. Cold Spring Harbor Laboratory, Cold Spring
Harbor, NY.
42. Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in *Escherichia coli*
K-12 using PCR products. Proc Natl Acad Sci U S A 97:6640-5.
43. Hogins J, Zimmern PE, Reitzer L. 2023. Genome sequences of seven clade B2 *Escherichia coli*
strains associated with recurrent urinary tract infections in postmenopausal women. Microbiol
Resour Announc doi:10.1128/mra.00035-23:e0003523.
44. Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods 25:402-8.
45. McCarthy DJ, Chen Y, Smyth GK. 2012. Differential expression analysis of multifactor RNA-Seq
experiments with respect to biological variation. Nucleic Acids Res 40:4288-97.
46. Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. Bioinformatics 26:139-40.

Table 1. Pairwise statistical comparisons of transcriptomes. Abbreviation: putr is putrescine

Strain 1	Strain 2	R^2
Wild-type (1 mM putr)	wild-type (0 mM putr)	0.950
Wild-type (1 mM putr)	Δ*speB* (1 mM putr)	0.954
Δ*speB* (0 mM putr)	Δ*speB* (1 mM putr)	0.820
Δ*speB* (0 mM putr)	wild-type (0 mM putr)	0.796
Δ*speB* (0 mM putr)	wild-type (1 mM putr)	0.786

566 Table 2. Differentially expressed genes that are proposed to contribute to putrescine homeostasis.

567 A positive number means more transcripts in the Δ*speB* (lower putrescine) strain.

Gene (function)	log$_2$FC	FDR
Polyamine synthesis		
speA (putrescine)	0.99	0.003
speC (putrescine)	1.43	3E-4
speD (spermidine)	-0.83	0.01
speF (putrescine)	1.05	0.007
rpmH (inhibitor of SpeA and SpeC)	-2.77	2E-4
rpsT (inhibitor of SpeA and SpeC)	-1.92	8E-4
Polyamine transport		
potABCD* (spermidine)	-2.17	3E-4
potE (putrescine)	2.07	2E-4
potFGHI* (putrescine)	1.41	4E-4
plaP (putrescine)	1.11	0.005
sapBCDF* (putrescine export)	-1.18	8E-4
Arginine synthesis		
argA_2 (synthesis)	1.77	0.001
argB (synthesis)	1.87	0.002
argC (synthesis)	2.86	0.005
argD (synthesis)	1.41	0.001
argF (synthesis)	1.63	0.01
argG (synthesis)	2.32	0.002
argH (synthesis)	0.97	0.001
argI (synthesis)	1.95	0.018
argR (repressor of arginine regulon)	-1.35	4E-4
Glutamate generation		
glsA (glutamine degradation)	2.16	5E-4

glnG (regulation of ammonia assimilation)	-1.59	0.022
TCA cycle/electron transport		
acnA (TCA cycle)	-1.04	8E-4
acnB (TCA cycle)	-1.96	2E-4
fumA (TCA cycle)	-1.50	6E-4
fumB (TCA cycle)	-1.49	0.003
gltA (TCA cycle)	-1.64	2E-4
icdA (TCA cycle)	-1.32	0.002
sdhCDAB-sucABCD* (TCA cycle)	-3.88	5E-5
nuo operon* (electron transport)	-0.82	0.004
menFDHBCE* (electron transport)	-0.8	0.005
ubiEJB* (electron transport)	-0.93	0.004
cyoABCD* (electron transport)	-1.41	9E-4
hemCD* (heme)	-2.05	2E-4
Iron transport		
exbBD* (transport of all iron chelates)	-2.33	4E-5
tonB (transport of all iron chelates)	-1.9	4E-5
entCEBA* (enterochelin iron)	-5.2	7E-6
fecABCDE * (ferric citrate)	-4.4	3E-5
fecIR (regulators, ferric citrate)	-4.4	6E-6
feoABC* (ferrous iron)	-3.4	7E-4
fur (regulator, iron assimilation)	-0.40	0.11
Magnesium and phosphate transport		
mgtA (magnesium)	5.90	5E-6
phoQ (regulator, magnesium)	0.79	0.009
pstSCAB* (phosphate)	3.61	0.01

*phoBR** (regulator, phosphate assimilation)	2.80	0.02

571 * Values for transcripts of the first gene of operon are given. Results for other genes of the

572 operon are in Supplemental Table 1

575 Table 3. Strains

Strain Name	Genotype	Reference
BLS77	W3110 Δ*puuR*::*cat*	(20)
BLS80	W3110 Δ*puuA*::*cat*	(20)
BLS88	W3110 Δ*patA* Δ*puuA*	(20)
CP2	W3110 Δ*patA*	(20)
IM26	W3110 Δ*speB*::*kan fliC-lacZ*	This study
IM27	W3110 Δ*speE*::*kan fliC-lacZ*	This study
IM28	W3110 Δ*speF*::*cat fliC-lacZ*	This study
IM29	W3110 Δ*speC*::*kan fliC-lacZ*	This study
IM34	W3110 Δ*speC* Δ*speF*::*cat fliC-lacZ*	This study
IM60	W3110 Δ*cadA*::*kan fliC-lacZ*	This study
IM61	W3110 Δ*speA*::*kan*	This study
IM62	W3110 Δ*patA* Δ*puuA* Δ*speE*::*kan*	This study
IM63	W3110 Δ*potE*::*kan*	This study
IM64	W3110 Δ*potF*::*kan*	This study
IM65	W3110 Δ*plaP*::*kan*	This study
IM66	W3110 Δ*puuP*::*kan*	This study
J15	W3110 Δ*speB* Δ*fliC*::*kan*	This study
SA1	W3110 Δ*fliC*::*kan*	This study
SA2	W3110 Δ*fimA*::*kan*	This study
SA3	W3110 Δ*fliC* Δ*fimA*	This study

SA4	W3110 Δ*hns::kan*	This study
SA5	W3110 Δ*ihfA::kan*	This study
SA6	W3110 Δ*lrp::kan*	This study
W3110-LR referred to as W3110	*lacI*q *lacL8*	Lab strain

Figure legends:

Figure 1 Pathways and enzymes of polyamine synthesis.

Figure 2. Surface motility of W3110. (A) Surface motility of parental and derivative strains lacking

the major subunits of the flagella (FliC), pili (FimA), or both. (B) TEM images of cells of W3110

and W3110 Δ*fliC* Δ*fimA* taken from the movement's edge directly from surface motility plates.

(C) Swimming motility with and without 0.5% glucose.

Figure 3. Genetics of surface motility. (A) PDSM of mutants with defects in polyamine anabolic

genes. All assays were performed in triplicate, and representative images are shown. (B)

Diameter of surface movement of polyamine mutants after 36 hours. Error bars represent

standard deviations for three independent replicates. Statistical analysis was performed using the

Dunnett test of significance: *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$. In this figure the Δ*speB*

mutant was IM26.

Figure 4: Nutritional supplementation of Δ*speB* mutant. (A) Putrescine and spermidine

supplementation of wild type and Δ*speB* strains. (B) Diameter of surface movement of

polyamine mutants after 36 hours. Error bars represent standard deviations for three independent

replicates. Statistical analysis was performed using the Sadik test of significance: *, $P<0.05$; **,

$P<0.01$; ***, $P<0.001$. All assays were performed in triplicates and representative images are

shown. In this figure the $\Delta speB$ mutant was IM26.

Figure 5: Putrescine transport and surface motility. (A) Representative images of surface motility

for strains defective in putrescine transport genes. (B) Diameter of surface movement of

polyamine mutants after 36 hours. Error bars represent standard deviations for three independent

replicates. Statistical analysis was performed using the Dunnett test of significance: *, $P<0.05$;

**, $P<0.01$; ***, $P<0.001$. All assays were performed in triplicate and representative images are

shown.

Figure 6: Putrescine catabolism and surface motility. (A) Pathways and enzymes of putrescine

catabolism. (B) Motility diameter of putrescine catabolic mutants after 36 hours. Error bars

represent standard deviations for three independent replicates. Statistical analysis was performed

using the Dunnett test of significance: *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$. All assays were

performed in triplicates and representative images are shown. (C) Effect of loss of $speE$ on the

$patA\ puuA$ catabolic double mutant.

Figure 7: Surface motility and pili expression in W3110 and $\Delta speB$. (A) W3110 $\Delta speB$ surface

motility with 0.0, 0.1, 1.0, and 4.0 mM exogenous putrescine. All assays were performed in

triplicate. Representative images are shown. (B) Average diameter of 3 separate surface motility

plates. The parental strain without putrescine is shown for reference. Significance was

determined by comparing the diameters of the Δ*speB* mutants in the different concentrations of

putrescine compared to the parental W3110. One way ANOVA was used with Dunnett

hypothesis testing to determine p values, ***, *p <0.001*. (C) Reverse transcriptase-quantitative

PCR using primers targeting the *fimA* gene from cells grown in 0.0, 0.1, 1.0, and 4.0 mM of

supplemented putrescine. Double deltas were generated by normalizing the parental and the

Δ*speB* mutant RNA libraries to *rpoD* then comparing *fimA* expression. Two-way ANOVA was

used to determine significance followed by Sadik hypothesis testing. ***, $p<0.001$. (D) Enzyme

linked immunosorbent assays using antibodies targeting pili (FimA) from cells grown with 0.0,

0.1, 1.0, and 4.0 mM of supplemented putrescine. Two-way ANOVA was used to determine

significance followed by FDR adjusting. ***, $p<0.001$. In this figure the Δ*speB* mutant was J15.

Figure 8: Transmission electron micrographs of Δ*speB* mutant cells after surface motility.

Representative images are shown and cells were removed from motility plates with the following

exogenous putrescine concentrations: (A) none, (B) 0.1 mM, (C) 1.0 mM, and (D) 4.0 mM. No

pili were observed with 0 and 0.1 mM putrescine. Optimal pili production was observed with 1.0

mM putrescine. The bar represents one micron. In this figure the Δ*speB* mutant was J15.

Figure 9: Visual representation of the results from transcriptomic sequencing of W3110 and the

Δ*speB* mutant's gene expression in media with and without 1.0 mM putrescine. (A)

Multidimensional scaling plot of the parental and Δ*speB* mutant transcriptomes. When grown

with 1.0 mM putrescine (red), the Δ*speB* mutant transcriptome (triangles) are nearly identical to

the parental transcriptomes when grown without and with putrescine (blue and red circles,

respectively). When grown without putrescine, the Δ*speB* mutant transcriptome (blue triangles)

is greatly skewed from transcriptomes of the parental strain and the Δ*speB* mutant grown with

putrescine. (B) Heatmap of the top 100 most variable genes further demonstrates the

distinctiveness of the Δ*speB* mutant grown without putrescine and the similarities of the Δ*speB*

mutant transcriptome when grown with 1.0 mM putrescine supplementation and the parental

grown with or without putrescine. In this figure the Δ*speB* mutant was J15.

Figure 10: Transcriptomic sequencing of genes for the *fim* region and regulators that control *fim*

gene expression. (A) Expression of the *fim* genes in the parental W3110 and the *speB* mutant

with and without putrescine supplementation. Values were calculated by dividing the individual

replicates' counts per million (CPM) value for each gene by the mean CPM of that gene in

W3110 grown without putrescine and multiplying by 100 to yield the percent expression. Arrows

below the genes signify the known operons: *fimB* and *fimE* belong to single gene operons, while

fimAICDFGH belong to one operon. One way ANOVA was used to determine significance using

Dunnett hypothesis testing. **, $p < 0.01$; ***, $p < 0.001$. (B) Expression of some regulators

known to affect *fim* gene expression. Values were calculated as described in (A). ***, $p < 0.001$.

(C) Surface motility of W3110 and three regulator mutants (Δ*hns*, Δ*ihfA*, and Δ*lrp*). A gene

found to be significantly different by this transcriptomic analysis (*hns*) was confirmed to be

important in surface motility. In this figure the Δ*speB* mutant was J15. (D) Diameter of surface

movement of regulatory mutants after 36 hours. Error bars represent standard deviations for three

independent replicates. Statistical analysis was performed using the Dunnett test of significance:

*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$. All assays were performed in triplicate.

Figure 11: The deduced diversion of metabolism in a *speB* mutant away from energy metabolism

toward putrescine synthesis compared to the parental strain. The effects of low putrescine are

shown. Genes and processes (transport and transcriptional regulators) in red have fewer

transcripts, while those in black or blue have more. The dashed lines represent proposed

reduction in metabolic flux because of fewer transcripts from genes coding for the enzymes

involved. Operons are shown except for the larger operons or regulons. For example, *pur* is

meant to represent the unlinked genes that code for enzymes of purine synthesis. Table 2 or

Supplemental Table 1 should be consulted for specific genes and the quantitative change in

transcripts.

Supplemental Figure 1: Growth and swimming of polyamine anabolic mutants.

Supplemental Table 1. Differential gene expression analysis of W3110 and Δ*speB*. J15 refers to

the Δ*speB* mutant. In the tab labeled "W3110_0_vs_J15_1_ Putrescine" W3110 was grown

without putrescine, J15 was grown with 1.0 mM putrescine; positive fold change values refer to

increases in J15 gene expression.

Supplementary Figure 2. Expression graphs comparing the logCPM of the transcriptomes of the

average of the three replicates of W3110 with and without 1 mM putrescine and the *speB*

mutant with and without 1 mM putrescine. A regression line was calculated and the correlation

between each set of transcriptomes noted on the graph. Higher R^2 values indicate greater

similarity between the transcriptomes.

Supplemental Figure 3. Phase variation in parental W3110, W3110 Δ*speB*, W3110 Δ*hns*, and

MG1655

Figure 1

Figure 1 Pathways and enzymes of polyamine synthesis.

Figure 2

A   Surface Motility

W3110   Δ*fliC*   Δ*fimA*   Δ*fliC* Δ*fimA*

B   Transmission Electron Microscopy

W3110   W3110 Δ*fliC* Δ*fimA*

C   Swimming Motility

- Glucose   + Glucose

Figure 2. Surface motility of W3110. (A) Surface motility of parental and derivative strains lacking the major subunits of the flagella (FliC), pili (FimA), or both. (B) TEM images of cells of W3110 and W3110 Δ*fliC* Δ*fimA* taken from the movement's edge directly from surface motility plates. (C) Swimming motility with and without 0.5% glucose.

Figure 3

A   Surface Motility of Polyamine Biosynthesis Mutants

W3110   ΔspeC   ΔspeF   ΔspeC ΔspeF

ΔspeA   ΔspeB   ΔspeE

B   Polyamine Biosynthesis Mutants
Surface Motility Diameters

Figure 3 Legend

Figure 3. Genetics of surface motility. (A) PDSM of mutants with defects in polyamine anabolic genes. All assays were performed in triplicate, and representative images are shown. (B) Diameter of surface movement of polyamine mutants after 36 hours. Error bars represent standard deviations for three independent replicates. Statistical analysis was performed using the Dunnett test of significance: *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$. In this figure the $\Delta speB$ mutant was IM26.

Figure 4

A

Surface Motility with Putrescine and Spermidine Supplementation

B

**Putrescine and Spermidine Supplementation
Surface Motility Diameters**

Figure 4 Legend

Figure 4: Nutritional supplementation of Δ*speB* mutant. (A) Putrescine and spermidine supplementation of wild type and Δ*speB* strains. (B) Diameter of surface movement of polyamine mutants after 36 hours. Error bars represent standard deviations for three independent replicates. Statistical analysis was performed using the Sadik test of significance: *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$. All assays were performed in triplicates and representative images are shown. In this figure the Δ*speB* mutant was IM26.

Surface Motility of Putrescine Export Mutants

A

$\Delta plaP$ $\Delta potF$ $\Delta potE$ $\Delta puuP$

Putrescine Export Mutant
Surface Motility Diameters

B

Figure 5: Putrescine transport and surface motility. (A) Representative images of surface motility for strains defective in putrescine transport genes. (B) Diameter of surface movement of polyamine mutants after 36 hours. Error bars represent standard deviations for three independent replicates. Statistical analysis was performed using the Dunnett test of significance: *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$. All assays were performed in triplicate and representative images are shown.

Figure 6

A

B

Putrescine Catabolism Double Mutant Surface Motility Diameters

C

Surface Motility of Putrescine Catabolism Double Mutants

W3110   ΔpatA ΔpuuA   ΔpatAΔpuuA ΔspeE

Figure 6: Putrescine catabolism and surface motility. (A) Pathways and enzymes of putrescine catabolism. (B) Motility diameter of putrescine catabolic mutants after 36 hours. Error bars represent standard deviations for three independent replicates. Statistical analysis was performed using the Dunnett test of significance: *, P<0.05; **, P<0.01; ***, P<0.001. All assays were performed in triplicates and representative images are shown. (C) Effect of loss of *speE* on the *patA puuA* catabolic double mutant.

Figure 7

Surface Motility of the W3110 Δ*speB* Δ*fliC* With and Without Putrescine Supplementation

A

0.0mM Putrescine   0.1mM Putrescine

1.0mM Putrescine   4.0mM Putrescine

B

W3110 Δ*speB* Δ*fliC* With and Without Putrescine Supplementation Surface Motility Diameters

Concentration of Putrescine in Millimolar

C

RT-qPCR of *fimA*

D

ELISA of FimA

Figure 7 Legend

Figure 7: Surface motility and pili expression in W3110 and ΔspeB. (A) W3110 ΔspeB surface motility with 0.0, 0.1, 1.0, and 4.0 mM exogenous putrescine. All assays were performed in triplicate. Representative images are shown. (B) Average diameter of 3 separate surface motility plates. The parental strain without putrescine is shown for reference. Significance was determined by comparing the diameters of the ΔspeB mutants in the different concentrations of putrescine compared to the parental W3110. One way ANOVA was used with Dunnett hypothesis testing to determine p values, ***, p <0.001. (C) Reverse transcriptase-quantitative PCR using primers targeting the *fimA* gene from cells grown in 0.0, 0.1, 1.0, and 4.0 mM of supplemented putrescine. Double deltas were generated by normalizing the parental and the ΔspeB mutant RNA libraries to *rpoD* then comparing *fimA* expression. Two-way ANOVA was used to determine significance followed by Sadik hypothesis testing. ***, p<0.001. (D) Enzyme linked immunosorbent assays using antibodies targeting pili (FimA) from cells grown with 0.0, 0.1, 1.0, and 4.0 mM of supplemented putrescine. Two-way ANOVA was used to determine significance followed by FDR adjusting. ***, p<0.001. In this figure the ΔspeB mutant was J15.

Figure 8

**Transmission Electron Microscopy of W3110ΔspeB
with and without Putrescine Supplementation**

Figure 8 Legend

Figure 8: Transmission electron micrographs of Δ*speB* mutant cells after surface motility. Representative images are shown and cells were removed from motility plates with the following exogenous putrescine concentrations: (A) none, (B) 0.1 mM, (C) 1.0 mM, and (D) 4.0 mM. No pili were observed with 0 and 0.1 mM putrescine. Optimal pili production was observed with 1.0 mM putrescine. The bar represents one micron. In this figure the Δ*speB* mutant was J15.

A

B

Figure 9 Legend

Figure 9: Visual representation of the results from transcriptomic sequencing of W3110 and the Δ*speB* mutant's gene expression in media with and without 1.0 mM putrescine. (A) Multidimensional scaling plot of the parental and Δ*speB* mutant transcriptomes. When grown with 1.0 mM putrescine (red), the Δ*speB* mutant transcriptome (triangles) are nearly identical to the parental transcriptomes when grown without and with putrescine (blue and red circles, respectively). When grown without putrescine, the Δ*speB* mutant transcriptome (blue triangles) is greatly skewed from transcriptomes of the parental strain and the Δ*speB* mutant grown with putrescine. (B) Heatmap of the top 100 most variable genes further demonstrates the distinctiveness of the Δ*speB* mutant grown without putrescine and the similarities of the Δ*speB* mutant transcriptome when grown with 1.0 mM putrescine supplementation and the parental grown with or without putrescine. In this figure the Δ*speB* mutant was J15.

Figure 10

Legend:
- ■ W3110 0.0mM Putrescine
- ■ (red) *speB* 0.0mM Putrescine
- ▤ W3110 1.0mM Putrescine
- ▤ (red) *speB* 1.0mM Putrescine

C **Surface Motility of Regulator Mutants**

W3110   Δ*hns*   Δ*ihfA*   Δ*lrp*

D **Regulator Mutant Surface Motility Diameters**

Figure 10 Legend

Figure 10: Transcriptomic sequencing of genes for the *fim* region and regulators that control *fim* gene expression. (A) Expression of the *fim* genes in the parental W3110 and the *speB* mutant with and without putrescine supplementation. Values were calculated by dividing the individual replicates' counts per million (CPM) value for each gene by the mean CPM of that gene in W3110 grown without putrescine and multiplying by 100 to yield the percent expression. Arrows below the genes signify the known operons: *fimB* and *fimE* belong to single gene operons, while *fimAICDFGH* belong to one operon. One way ANOVA was used to determine significance using Dunnett hypothesis testing. **, $p < 0.01$; ***, $p < 0.001$. (B) Expression of some regulators known to affect *fim* gene expression. Values were calculated as described in (A). ***, $p < 0.001$. (C) Surface motility of W3110 and three regulator mutants (Δhns, $\Delta ihfA$, and Δlrp). A gene found to be significantly different by this transcriptomic analysis (*hns*) was confirmed to be important in surface motility. In this figure the $\Delta speB$ mutant was J15. (D) Diameter of surface movement of regulatory mutants after 36 hours. Error bars represent standard deviations for three independent replicates. Statistical analysis was performed using the Dunnett test of significance: *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$. All assays were performed in triplicate.

Figure 11 Legned

Figure 11: The deduced diversion of metabolism in a *speB* mutant away from energy metabolism toward putrescine synthesis compared to the parental strain. The effects of low putrescine are shown. Genes and processes (transport and transcriptional regulators) in red have fewer transcripts, while those in black or blue have more. The dashed lines represent proposed reduction in metabolic flux because of fewer transcripts from genes coding for the enzymes involved. Operons are shown except for the larger operons or regulons. For example, *pur* is meant to represent the unlinked genes that code for enzymes of purine synthesis. Table 2 or Supplemental Table 1 should be consulted for specific genes and the quantitative change in transcripts.