Control of pili synthesis and putrescine homeostasis in *Escherichia coli*

Running title: *E. coli* surface motility and putrescine homeostasis

Iti Mehta[1#], Jacob Hogins[1#], Sydney Hall[1], Gabrielle Vragel[1], Sankalya Ambagaspitiye[1], Philippe

Zimmern[2], Larry Reitzer[1]*

[1] Department of Biological Sciences, UT Dallas, Richardson, Texas

[2] Department of Urology, the University of Texas Southwestern, Dallas, Texas, USA

*Author for correspondence: Tel: 972-883-2502; Fax: 972-883-2409; reitzer@utdallas.edu

# Authors contributed equally and should be considered co-first authors

Keywords: polyamines, putrescine, surface motility, pili, *Escherichia coli*, putrescine

homeostasis

Summary

Polyamines are biologically ubiquitous cations that bind to nucleic acids, ribosomes, and phospholipids and, thereby, modulate numerous processes, including surface motility in *Escherichia coli*. We characterized the metabolic and transcription factors that contribute to polyamine-dependent control of surface motility. Genetic analysis showed that surface motility required type 1 pili and the simultaneous presence of two independent putrescine anabolic pathways and that putrescine transport and catabolism modulated surface motility. The results of an immunological assay for FimA—the major pili subunit, reverse transcription quantitative PCR of *fimA*, and transmission electron microscopy confirmed that pili synthesis required putrescine. RNAseq analysis of a parental and putrescine-deficient mutant and a follow-up genetic analysis suggested that H-NS was the putrescine-responsive regulator of pili synthesis and surface motility. The RNAseq results also showed that low putrescine increased transcripts for genes of arginine synthesis and transport systems for magnesium and phosphate, and decreased transcripts of genes for oxidative energy metabolism and iron transport. We conclude that (a) putrescine controls pili synthesis and surface motility possibly via the transcriptional regulator H-NS, and (b) a complex homeostatic network, which can rewire metabolism, controls putrescine concentrations. During a urinary tract infection, the polyamine putrescine is present in urine and pili are required for the binding of bacteria to the bladder epithelium. Because of its control of pili synthesis and energy metabolism, putrescine is likely to contribute to the establishment and maintenance of urinary tract infections associated with uropathogenic *Escherichia coli*.

Introduction

Polyamines are flexible aliphatic cations that are found in virtually all organisms. *E. coli* contains putrescine (1,4-diaminobutane) and lesser amounts of spermidine (N-(3-aminopropyl)-1,4-diaminobutane), and cadaverine (1,5-diaminopentane) (1, 2). Both the pathways and enzymes of polyamine synthesis are redundant (Fig. 1). Strains that lack eight of these genes do not contain detectable polyamines, grow slowly aerobically, and do not grow anaerobically (3). Polyamines interact with nucleic acids and phospholipids, can affect chromosome and ribosome structure, ribosome-mRNA interactions, and protein and nucleic acid elongation rates (4). Polyamines modulate, but are not required for, protein synthesis (5) and expression of hundreds of genes (6). A major mechanism of polyamine-dependent regulation is translational control of genes for 18 major transcription regulators: four of the seven σ factors – σ^{18} (FecI), σ^{24} (RpoE), σ^{38} (RpoS), and σ^{54} (RpoN); three histone-like DNA-binding proteins (Fis, H-NS, and StpA); and CpxR, Cya, Cra, and UvrY (5, 6).

Polyamines control flagella-dependent surface motility of *Proteus mirabilis* and pili-dependent surface motility (PDSM) of *E. coli*: the latter has been reported to require the polyamine spermidine (7, 8). While reconstructing putrescine-deficient strains to characterize the basis for the anaerobic growth defect, we noticed that a mutant unable to synthesize spermidine moved normally, while mutants lacking enzymes of putrescine synthesis moved poorly. We characterized the role of polyamines on *E. coli* surface motility and pili synthesis by genetic analysis, ELISA assays, RT-qPCR, electron microscopy, and RNAseq. We showed that PDSM and pili synthesis required two independent pathways of putrescine synthesis, but did not require spermidine, and was modulated by putrescine transport and catabolism. In addition to confirming that putrescine controlled pili synthesis, results from an RNAseq analysis identified regulators that mediated

putrescine-dependent control, and, unexpectedly, suggested a putrescine homeostatic network that

rewires metabolism to maintain intracellular putrescine.

Results

Pili-dependent surface motility (PDSM) of W3110.

While characterizing mutants lacking putrescine synthetic enzymes, we found that some

strains were defective in surface motility. Because any single laboratory strain may have properties

that result from unintended laboratory evolution, we assessed nine strains for their surface motility

pattern. Five strains covered a plate in 12-18 hours, while four did not traverse a plate in 36 hours.

The slow-moving strains generated genetically stable fast-moving variants which suggests that the

former were ancestral. Our lab strain of W3110 moved slowly, and among the slow movers showed

the least variability in spreading diameter and generated fewer fast-moving variants. Its outward

movement formed concentric rings, with alternating phases of movement and

consolidation/growth (Fig 2A). Deletion of *fimA*, which codes for the major type 1 pili subunit,

abolished the oscillatory pattern and reduced, but did not eliminate, surface motility, but deletion

of *fliC* which encodes the major flagellum subunit did not affect surface motility (Fig. 2A).

Electron microscopy confirmed that W3110-LR isolated from surface motility plates possessed

pili (Fig. 2B). None of >500 cells expressed flagella. A mixed population of elongated (3-4 μm)

and non-elongated (<2 μm) cells were visible, and pili were mostly associated with elongated cells.

Furthermore, W3110 surface motility requires glucose (9) which should prevent flagella synthesis

via inhibition of cyclic AMP synthesis (10). We confirmed that glucose controlled flagella

synthesis in W3110 by its suppression of swimming motility (Fig 2C). A W3110-LR derivative

with deletions of both *fliC* and *fimA* still exhibited outward movement on a surface motility plate

(Fig 2A), but electron microscopy showed the absence of appendages (Fig. 2C). In summary,

genetics, electron microscopy, and a glucose requirement show that W3110 surface motility

requires pili.

PDSM required two independent putrescine synthesis pathways.

We examined PDSM requirements for polyamine anabolic pathways. A major putrescine

biosynthetic pathway is ornithine decarboxylation either by the constitutive SpeC or the low-pH

inducible SpeF. The Δ*speC* and Δ*speF* mutants moved as well as the parental strain, but a Δ*speC*

Δ*speF* double mutant moved less well (Figure 3). SpeA (arginine decarboxylase) and SpeB

(agmatinase) catalyze an alternate two-step putrescine synthetic pathway (Fig 1) (11): deletion of

either gene reduced surface motility (Figure 3). The Δ*speA*, Δ*speB*, and Δ*speC* Δ*speF* strains grew

as well as the parental strain (Supplemental Figure 1A) and had normal swimming motility

(Supplemental Figure 1B). A Δ*speE* mutant, which lacks the enzyme for spermidine synthesis,

moved on a surface as well as the parental strain (Fig 3).

Supplemental putrescine at 1 mM restored surface motility of a *speB* mutant (Figure 4A)

and a *speA* mutant (not shown). Spermidine did not restore wild type movement and, also, caused

an unusual pattern of movement in the parental strain (Figure 4B). These exogenous concentrations

of putrescine and spermidine had no effect on growth rates in liquid motility medium (not shown).

In summary, PDSM required the simultaneous presence of two independent pathways of putrescine

synthesis but did not require spermidine.

Loss of putrescine transport systems affected surface motility.

Putrescine transport may maintain an appropriate intracellular putrescine concentration,

because *E. coli* secretes putrescine (12). *E. coli* has several putrescine transport systems (13, 14).

Mutants lacking the PlaP and PotF putrescine transporters had 50% and 40% reduced surface

motility, respectively, and lost the concentric ring pattern (Figure 5A). Loss of the PotE and PuuP

transporters affected neither the movement diameter (Figure 5B) nor concentric ring formation

(Figure 5A). We note that *potE* and *plaP* are more highly expressed than *potE* and *puuP* (85, 64,

17, and 6 counts per million transcripts, respectively) for W3110 grown in liquid motility medium

(15). We conclude that extracellular putrescine and its transport contribute to PDSM.

Putrescine catabolism affected surface motility.

Because putrescine catabolism modulates intracellular putrescine concentrations (16, 17),

we examined PDSM in putrescine catabolic mutants. The major putrescine catabolic pathway is

initiated with putrescine glutamylation (reactions 6-9, and 4-5, Fig 6A), and a second pathway

initiates with putrescine deamination (reactions 1-5, Fig 6A) (17). Deletion of either *puuA* or *patA*

which code for the first enzymes of their respective pathways did not affect surface motility (Fig

6B). A double mutant moved less well which was unexpected because the proposed higher

intracellular putrescine should have stimulated PDSM. Consistent with the stimulatory effect of

putrescine, loss of *puuR* which codes for the repressor of PuuA-initiated pathway genes (17),

impaired PDSM (Fig 6B). One possible explanation to reconcile these seemingly contradictory

results is that an optimal polyamine concentration stimulates PDSM, and a high concentration is

inhibitory. Experiments that are described in the next section test this possibility.

High putrescine reduced expression of pili genes.

High putrescine inhibits translation of some mRNAs, and high spermidine, a product of

putrescine metabolism, inhibits growth (18, 19). If the phenotype of the Δ*patA* Δ*puuA* double

mutant results from spermidine toxicity, then loss of spermidine synthase (SpeE) should reverse

the phenotype. This prediction was not met which argues against spermidine toxicity (Fig 6C).

For the *speB* mutant, 1 mM putrescine stimulated PDSM, and 4 mM putrescine was slightly

inhibitory but eliminated the ring pattern (Fig 7AB). The concentric ring pattern was observed

with 1 mM, but not 4 mM putrescine. Indirect enzyme linked immunosorbent assays (ELISAs)

against FimA showed that the level of FimA in the *speB* mutant at 1 mM putrescine was higher

than with 4 mM putrescine (Fig. 7C). Variation of supplemental putrescine did not affect FimA in

the parental strain (Fig. 7C).

To test transcriptional regulation of pili expression by putrescine, we examined transcript

levels by RT-qPCR. Optimal *fimA* transcription occurred at 1 mM putrescine in the *speB* mutant

(Fig 7D). Transmission electron microscopy (TEM) showed that pili expression was highest for

the *speB* mutant at 1 mM putrescine (Fig. 8), moderate at 0.1- and 4-mM putrescine, and

undetectable without putrescine. The cells grown with 4 mM putrescine were shorter and thinner,

which suggests a stress, possibly mimicking osmotic stress. From these results and those in the

previous section, we conclude that pili expression required an optimal putrescine concentration.

RNAseq analysis identified H-NS as a possible mediator of putrescine-dependent control.

To determine whether putrescine affects transcription of genes whose products control

fim operon expression (20), we performed RNAseq analysis of the parental and *speB* strains

grown with and without 1 mM putrescine. The transcriptomes of the parental strain both with

and without putrescine and the *speB* mutant with 1 mM putrescine were highly similar with R^2

values of 0.950 and 0.954, respectively, based on expression graphs comparing the

transcriptomes of the parental strain in both conditions to the *speB* mutant strain (Supplemental

Figure 2). Compared to the transcriptome of the *speB* mutant grown without putrescine, the

transcriptomes of the parental strain without putrescine ($R^2 = 0.796$), the parental with 1 mM

putrescine ($R^2 = 0.786$), and the *speB* mutant with 1 mM putrescine ($R^2 = 0.820$) differed

substantially. A multidimensional scaling plot and a plot of the 100 most variable genes further

visualize these interactions (Fig 9A and 9B).

When compared to the parental strain, the *speB* mutant grown without putrescine had 310

downregulated genes and 159 upregulated genes with at least four-fold differential expression

and FDR < 0.05. Transcripts for the putrescine-responsive *puuAP* and *puuDRCBE* operons,

which specify genes of the major putrescine catabolic pathway, were reduced from 1.6- to 14-

fold (FDR ≤ 0.02) in the *speB* mutant (Supplemental Table 1), which implies lower intracellular

putrescine. Deletion of *speB* reduced transcripts for genes of the *fimA* operon and *fimE*, but not

of *fimB* (Fig 10A). Furthermore, loss of *speB* affected some of the known positive effectors that

control *fim* operon expression which include Fis, H-NS, IHF, Lrp, and QseB (21). Transcripts

from *hns* were reduced, transcripts from *ihfA* and *ihfB* were not affected, and transcripts from *fis*,

lrp, and *qseB* were increased when *speB* was deleted (Fig 10B). Loss of *speB* also increased

transcripts from *lrhA* which codes for a positive effector of *fimE*. Loss of several of these

regulators eliminated PDSM which was expected (Fig 10C). H-NS may mediate putrescine-

dependent control of pili synthesis. However, given the numerous effects of loss of *speB*, a

generalized dysregulation of several major transcription factors may also produce a less-than-

optimal constellation of regulators.

Evidence for a homeostatic network that maintains intracellular putrescine.

The gene expression changes resulting from loss of *speB* unexpectedly suggested compensatory mechanisms for low putrescine. The transcript differences for all genes are provided in Supplemental Table 1, and for selected genes in Table 1. A gene expression pattern is summarized in Fig 11. The *speB* mutant had more transcripts from (a) genes of arginine and ornithine carboxylase, (b) all genes of ornithine and arginine synthesis from glutamate, and (c) genes for three separate putrescine transport systems. The *speB* mutant also had fewer transcripts from the *sap* operon which codes for a putrescine exporter (12), and from *rpmH* and *rpsT*, whose products inhibit ornithine and arginine decarboxylase (22). The net effect of these changes should be to increase intracellular putrescine: follow the solid arrows in the left half of Fig. 11.

Loss of *speB* also affected pathways of energy metabolism. For instance, glutamate and α-ketoglutarate are at a branch point in metabolism which can either provide putrescine precursors or generate energy. The *speB* mutant had fewer transcripts for metabolic genes coding for all citric acid cycle enzymes, NADH dehydrogenase I (the entire *nuo* operon), cytochrome oxidase (the *cyo* operon), and several genes of menaquinone (the *menFDHBCE* operon), ubiquinone, and heme synthesis. In addition to these differences, the *speB* mutant also had fewer transcripts for all genes of iron acquisition, which is consistent with diminished energy generation. In sum, this pattern suggests that low putrescine diverts metabolic flux from energy-generating pathways to putrescine synthesis.

One of the most differentially expressed gene between the parental and *speB* mutant strains was *mgtA* which had 60-fold more transcripts in the *speB* mutant and codes for the major magnesium transporter. The *speB* mutant also had more transcripts for several phosphate

195 assimilation genes, including *phoB* which codes for the response regulator of phosphate

196 limitation.

Discussion

PDSM and pili synthesis required putrescine synthesis and were affected by putrescine

degradation and transport, as well as several transcription factors, which implies that numerous

factors control intracellular putrescine concentrations. The effect of loss of putrescine transport

systems on PDSM implicates extracellular putrescine on putrescine homeostasis. A requirement

for the SpeA-SpeB pathway implicates extracellular arginine since SpeA is mostly periplasmic

(23). The role of both putrescine synthesis and degradation suggests rapid adjustments to

fluctuating putrescine concentrations. Numerous transcription factors control pili synthesis. Of

these, only the gene for H-NS had fewer transcripts in the *speB* mutant, which is consistent with

H-NS mediating putrescine-dependent control. This conclusion is not considered strong because

of the only two-fold transcript reduction, and because of numerous changes in transcripts for other

potential pili regulations in the *speB* mutant, albeit in the wrong direction. Therefore, we cannot

exclude a general regulatory dysfunction mediated by depleted levels of putrescine as the basis for

loss of pili synthesis.

An unexpected observation was that the *speB* mutant had more transcripts for genes whose

products will maintain intracellular putrescine, i.e., compensatory mechanisms for low

intracellular putrescine (Figure 11). Conversely, loss of *speB* lowered transcripts for many genes

of energy metabolism and all genes of iron acquisition. This gene expression pattern suggests a

diversion of metabolism away from energy generation and toward putrescine synthesis, i.e., a

prioritization of putrescine synthesis over aerobic energy metabolism.

We also propose that other changes are also compensatory mechanisms for low intracellular

putrescine, specifically, more transcripts for genes coding for magnesium and phosphate transport

systems in the *speB* mutant. To the extent that magnesium can directly replace putrescine, e.g.,

binding to nucleotides and RNA (24), magnesium could compensate for low putrescine. Our

observations confirm those of J. Slauch and colleagues on an inverse correlation between

intracellular magnesium and polyamines in *Salmonella* (25). They proposed an overall divalent

cation homeostasis maintained by either magnesium or the polyamines (25). A rationale for the

effect on phosphate metabolism is not obvious but considering the effects of low putrescine on

energy metabolism, increased phosphate transport may be an adjustment to changes in energy

metabolism. In any case, the substantial changes in gene expression due to low putrescine suggest

multiple compensatory mechanisms, and a homeostatic gene network which because of effects on

metabolic genes can rewire metabolism.

In contrast to our results, a thorough and well-designed study showed that *E. coli* PDSM

requires spermidine instead of putrescine (8). The differences between the studies were strain

backgrounds, media (glucose-LB vs glucose-tryptone), and incubation temperature (33º vs 37º).

We found PDSM results for 37º incubations were highly variable and resulted in numerous fast-

moving variants. Since growth at the higher temperature is faster, and faster growth is associated

with higher putrescine, we propose that growth at 37º results in an inhibitory putrescine

concentration. In this case, spermidine would be stimulatory because of its known inhibition of the

putrescine biosynthetic enzymes SpeA and SpeC (26, 27) and subsequent reduction of intracellular

putrescine to an optimal concentration.

The transcript changes observed in our study were different from those in the only other

systematic study of putrescine's effect on *E. coli* gene expression (6). Both studies had almost

identical experimental design: 1.0- or 1.14-mM putrescine was added to either a *speB* or *speB speC*

mutant, respectively. Differences in strains, growth media (minimal media vs tryptone-containing

media), and growth temperature (33^0 vs 37^0) could account for the discrepancy in results. The most

crucial difference could be the growth media: Igarashi and colleagues grew bacteria in a minimal

medium, while we grew bacteria in an amino acid-containing medium that was required for surface

motility. We propose that the slower growth in minimal medium correlates with higher guanosine

tetraphosphate and in turn counters and obscures many growth-stimulatory putrescine effects.

In contrast, the growth stimulation from the amino acids in our medium would not counter

putrescine effects and would essentially allow detection of effects caused solely by changes in

putrescine concentrations. Regardless of the explanation, common results can be interpreted as

core responses to putrescine limitation. Both studies observed that low putrescine increased

transcripts for genes of arginine synthesis and magnesium transport, and reduced transcripts for

the *sdhCDAB-sucABCD* operon which codes for TCA cycle enzymes. The link between putrescine

and arginine synthesis is probably mediated by the ArgR repressor because *argR* transcripts are

2.5-fold lower (FDR = 0.0004) in the *speB* mutant (Supplemental Table 1). Cyclic AMP may

mediate putrescine-dependent control of the *sdh-suc* operon because putrescine controls

translation of *cya* mRNA, which codes for adenylate cyclase, and cyclic-AMP controls the

sdhCDAB-sucABCD operon. As noted above, the link between putrescine and magnesium

transport is a recent observation (25).

Several results implicate putrescine in *E. coli* virulence during urinary tract infections.

First, pili-dependent binding to the urothelium is an essential step for uropathogenic *E. coli* (28).

Second, putrescine is present in urine from infected, but not healthy individuals (29, 30). Third,

urinary putrescine can result from UPEC growth in urine (30) and urothelial pathophysiology (31).

Putrescine's possible role in virulence is seemingly contradicted by the observation that a *speB*

mutant outcompetes the parental CFT073 in the bladder of mice, i.e., less putrescine synthesis

enhanced virulence (32). A majority of uropathogenic *E. coli* are associated with phylogenetic

group B2—including CFT073, whereas most *E. coli* lab strains, such as W3110, are group A.

Group B2 strains lack genes for the major putrescine catabolic pathway and have significantly

higher transcripts for genes coding for SpeA and SpeB (15). Both observations suggest that the

cytoplasmic putrescine concentration is higher in group B2 strains. We propose that the putrescine

concentration in group B2 strains is normally in the inhibitory range for pili synthesis, and that

loss of SpeB reduces putrescine to a level that stimulates pili synthesis. Regardless of the

explanation, putrescine is likely to be important during urinary tract infections, but aspects of

putrescine control of pili synthesis and other processes could be different in uropathogenic *E. coli*

and is worth further examination.

Experimental Procedures

Strains. All strains used for growth rate determinations and motility assays were derivatives of *E.*

coli K-12 strain W3110 and are listed in Table 2. Great variations exist in standard lab strains,

including W3110, from different labs. We tested nine strains, most from the Coli Genetic Stock

Center, and chose our lab strain because of relative genetic stability and more quantitatively

reproducible results. Our strain originated from the lab of Jon Beckwith in the 1960s via the lab of

Boris Magasanik where it had been stored in a room temperature stab until the mid-1980s, at which

time it was revived and frozen at -80⁰ C. Although mutations may have accumulated during

storage, its transcriptome is similar to that of other strains of the same phylogenetic group when

grown in motility medium (15). To construct the mutant strains, the altered allele was obtained

from the Keio collection in which the gene of interest had been deleted and replaced with an

antibiotic resistance gene (33). The marked deletion allele was transferred into W3110 by P1

transduction (34). The antibiotic gene was removed as described which generated an in-frame

deletion (35).

Media and Growth Conditions. For P1 transductions and plasmid transformation experiments,

cells were grown in standard LB liquid medium (1% tryptone, 1% NaCl and 0.5% yeast extract,

pH 7.0) at 37°C. Antibiotics were used for selection at concentrations of 25 µg/mL (both

chloramphenicol and kanamycin). For growth analysis, bacteria were grown in liquid motility

medium (0.5% glucose, 1% tryptone, and 0.25% NaCl) which we refer to as GT medium. Starter

cultures (typically 6–12 h incubation) were grown in GT medium, harvested by centrifugation,

washed twice with phosphate buffered saline and re-suspended in GT medium before inoculation.

The cells were then grown at 37° C in aerobic conditions (shaking at 240 rpm) and the turbidity

was measured every thirty minutes. Cell growth was measured in Klett units using a Scienceware

Klett colorimeter with a KS-54 filter. 100 Klett units represent an OD_{600} value of about 0.7.

Surface motility assay. For a standard surface motility assay, single colonies from fresh plates

(streaked out from frozen stocks a day before) were inoculated in GT medium for six hours at

37° C in aerobic conditions (shaking at 240 rpm). 30 ml of autoclaved GT medium with 0.45%

agar (Eiken, Tokyo, Japan) was poured into sterile polystyrene petri-dish (100 mm X 15 mm)

and allowed to solidify at room temperature for approximately six hours. Then, 1 µL of the pre-

motility growth medium was inoculated at the center of the agar plate and incubated at 33° C for

36 hrs. Each experiment was performed in triplicate and pictures were taken after 36 hours.

Surface motility was extremely sensitive to humidity. Opening the incubator before 36 hours

resulted in movement cessation. The surface motility assay was performed at 33º C because

results from incubations at 37º C were highly variable and the cultures more frequently generated

genetically stable fast-moving variants.

Swim assay. The media and culturing for swimming motility is identical to that for surface

motility, except that the plates contained 0.25% agar and were solidified at room temperature for

about one hour before inoculation. Then, 1 µL of the culture was stabbed in the center of this

swim agar plate and incubated at 33° C for 20 hours. Each experiment was performed in

triplicate.

Transmission electron microscopy. Cells from surface motility plates were collected and fixed

with 2.5% glutaraldehyde. Bacteria were absorbed onto Foamvar carbon-coated copper grids for

1 min. Grids were washed with distilled water and stained with 1% phosphotungstic acid for 30

s. Washed and stained grids were dried at 37ºC for 10 minutes. Samples were viewed on a JEOL

1200 EX transmission electron microscope at the University of Texas Southwestern (Figure 2)

and the University of Texas at Dallas (Figure 8).

ELISA analysis. Cells were grown overnight in 5 mL GT media followed by a 2-hour growth in

GT media with 0.0, 0.1, 1.0, or 4.0 mM putrescine. Cells were then lysed using a 24-gauge

needle. The cell lysate was diluted to a concentration of 1 mg/mL protein (based on A280) in pH

9.6 coating buffer (3 g Na_2CO_3, 6 g $NaHCO_3$, 1000 mL distilled water) and then coated onto the

walls of a 96-well plate by incubating overnight at 4°C. The wells were rinsed 3 times with PBS

and further blocked using coating buffer with 1% bovine albumin overnight at 4°C. The

following day, wells were rinsed 3 times with PBS, and primary antibodies to FimA or RpoD

were added following supplier directions. After a 2-hour room temperature incubation, secondary

antibodies conjugated to HRP were added and incubated for a further two hours. HRP was

activated using the TMB substrate kit (Thermo-34021) following the provided protocol.

Expression was read on a BioTek plate reader based on the provided kit protocol. Data was

blanked to wells only treated with bovine albumin then normalized to RpoD. Data was generated

from technical and biological triplicates.

DNA isolation and genome assembly and annotation for W3110. DNA was isolated based on

previous established protocols (36). Short reads were sequenced at the SeqCenter (Pittsburgh,

Pennsylvania) using Illumina tagmentation-based and PCR-based DNA prep and custom IDT

indices targeting inserts of 280 bp without further fragmentation or size selection steps. The

Illumina NovaSeq X Plus sequencer was ran producing 2X151 paired-end reads.

Demultiplexing, quality control, and adapter trimmer was performed with bcl-convert (v4.2.4).

Total Illumina Reads (R1+R2): 4059078 with 553087045 bps >Q30.

Long reads were prepared using the PCR-free Oxford Nanopore Technologies ligation

sequencing kit with the NEBNext companion module. No fragmentation or size selection was

performed. Long read libraries were performed using R10.4.1 flow cell. The 400bps sequencing

mode with a minimum read length of 200bp were selected. Gupply (v6.5.7) was used at the

super-accurate basecalling, demultiplexing, and adapter removal. Total long reads: 53773 with

89.607 % of bps > Q20.

To generate the completed genome, porechop (v0.2.4) was used to trim residual adapter

sequences from long reads. Flye (v2.9.2) was used to generate the *de novo* assembly under the

nano-hq model. Reads longer than the estimated N_{50} based on a genome size of 6Mbp initiated

the assembly. Subsequent polishing using the short read data was performed using Pilon (v1.24)

under default parameters. Long read contigs with an average short read coverage of 15X or less

were removed from the assembly. The assembled contig was confirmed to be circular via

circulator (v1.5.5). Annotation was performed using prokka (v 1.14.6). Finally, statistics were

recorded using QUAST (v5.2.0). The final genome contained 1 contig of 4750347 bp with a

sequencing depth of 123.92x. The N_{50} was 4750347. This genome can be accessed via the

accession number CP165600 or via the BioProject PRJNA1142534 via NCBI.

The completed genome has an estimated average nucleotide identity of 99.9694 to the W3110

genome deposited to NCBI (genome assembly ASM1024v1).

RNA isolation and quality control. Cells were grown for 2 hours in 1 mL GT media. 60 μL

were then added to 1 mL of fresh GT, with or without 1 mM putrescine, and grown for another

two hours. After growth, the cells were centrifuged, and frozen at -80ºC. The isolation and

analysis protocol has been previously published (15). Cell pellets were thawed, resuspended in

0.7 mL of buffer RLT (Qiagen RNeasy kit), and mechanically lysed using a bead beater

(FastPrep-24 Classic from MP Biomedical) set at the highest setting for three 45-s cycles with a

5-min rest period on ice between cycles. Cell lysates were used to isolate RNA using a Qiagen

RNeasy mini kit. RNA was quantified, DNase treated, and re-quantified on a nanodrop. RNA

that passed the first quality check was analyzed on RNA-IQ (Qubit). RNA with an IQ higher

than 7.5 was used. RNA that passed both checks were run on an agarose gel to check for RNA

integrity. RT-qPCR was used to ensure there was no DNA contamination. RNA libraries that

passed all these quality checks were submitted to the genomics core facility at the University of

Texas at Dallas for RNA sequencing. The core performed rRNA removal (RiboMinus

Transcriptome Isolation Kit or RiboCop bacterial rRNA depletion—Lexogen), library

preparation (Stranded Total RNA Prep—Illumina), and single-end Illumina sequencing.

Reverse transcription quantitative PCR. RNA was isolated as described above. One

microgram of RNA was reverse transcribed using LunaScript RT Super Mix (NEB E3010) and

another microgram was subjected to the same reaction as the reverse transcribed RNA, except

without reverse transcriptase (negative control) following manufacturer instructions. The cDNA

was then diluted to 10 ng/µL for qPCR. PCR reactions were composed of 10 ng cDNA, 8 µL of

nuclease free water, 10 nM primers (rpoD: Forward: TCGTGTTGAAGCAGAAGAAGCG;

Reverse: TCGTCATCGCCATCTTCTTCG) (fimA: Forward: ATGGTGGGACCGTTCACTTT;

Reverse: GGCAACAGCGGCTTTAGATG), and PowerUp SYBR Green 2X master mix for

qPCR (ThermoFisher A25777) in a 20 µL reaction. A Quantstudio 3 qPCR machine was used to

generate critical threshold (Ct) values. The Ct values were then analyzed using the $2^{-\Delta\Delta C_T}$ method

(37) using *rpoD* as a library control.

RNAseq Analysis. Transcripts were aligned to the W3110 genome and CDS (CP165600)

available on NCBI using CLC genomics workbench (Qiagen version 22.00) for RNA-seq

analysis. The gene expression counts generated as this output were used to perform differentially

expressed gene (DEG) analysis using EdgeR (version 3.38.4) to analyze read counts (38, 39).

DEGs had an adjusted P-value less than 0.05.

Data availability: The raw RNA seq reads were deposited in GenBank: BioProject accession

number PRJNA1126736. The genome was deposited in GenBank: BioProject accession number

PRJNA1142534. The genome accession number is CP165600.

Acknowledgements: This work was funded through private donations. PEZ received

endowment support from the Cain Foundation through the Felecia and John Cain Distinguished

Chair in Women's Health in honor of Philippe E. Zimmern, MD. The authors declare no potential

conflicts of interests.

The authors (JH, SH, and GV) would like to thank Prof. Kelli Palmer of the University of

Texas at Dallas for the use of her space, supplies, and tissue homogenizer. JH would like to

further thank Prof. Palmer for the use of her CLC genomics program. The authors would like to

thank the Genome Center at The University of Texas at Dallas for the services to support our

research. The authors would like to thank the Olympus Discovery Center/Imaging Core facility

at UT Dallas for providing equipment and support (Fig. 8), and the UT Southwestern Medical

Center for electron microscopy (Fig. 2).

References:

420   1.   Tabor CW, Tabor H. 1984. Polyamines. Annual review of biochemistry 53:749-90.
421   2.   Tabor CW, Tabor H. 1984. Polyamines. Annu Rev Biochem 53:749-90.
422   3.   Chattopadhyay MK, Tabor CW, Tabor H. 2009. Polyamines are not required for aerobic growth of
423   *Escherichia coli*: preparation of a strain with deletions in all of the genes for polyamine
424   biosynthesis. J Bacteriol 191:5549-52.
425   4.   Igarashi K, Kashiwagi K. 2000. Polyamines: mysterious modulators of cellular functions. Biochem
426   Biophys Res Commun 271:559-64.

5.   Igarashi K, Kashiwagi K. 2015. Modulation of protein synthesis by polyamines. IUBMB Life
67:160-9.
6.   Yoshida M, Kashiwagi K, Shigemasa A, Taniguchi S, Yamamoto K, Makinoshima H, Ishihama A,
Igarashi K. 2004. A unifying model for the role of polyamines in bacterial cell growth, the
polyamine modulon. J Biol Chem 279:46008-13.
7.   Armbruster CE, Hodges SA, Mobley HL. 2013. Initiation of swarming motility by *Proteus mirabilis*
occurs in response to specific cues present in urine and requires excess L-glutamine. J Bacteriol
195:1305-19.
8.   Kurihara S, Suzuki H, Tsuboi Y, Benno Y. 2009. Dependence of swarming in *Escherichia coli* K-12
on spermidine and the spermidine importer. FEMS Microbiol Lett 294:97-101.
9.   Sudarshan S, Hogins J, Ambagaspitiye S, Zimmern P, Reitzer L. 2021. The nutrient and energy
pathway requirements for surface motility of nonpathogenic and uropathogenic *Escherichia coli*.
J Bacteriol. doi:10.1128/jb.00467-20doi:10.1128/jb.00467-20.
10.   Yokota T, Gots JS. 1970. Requirement of adenosine 3', 5'-cyclic phosphate for flagella formation
in *Escherichia* coli and *Salmonella typhimurium*. J Bacteriol 103:513-6.
11.   Tabor CW, Tabor H. 1985. Polyamines in microorganisms. Microbiol Rev 49:81-99.
12.   Sugiyama Y, Nakamura A, Matsumoto M, Kanbe A, Sakanaka M, Higashi K, Igarashi K, Katayama T,
Suzuki H, Kurihara S. 2016. A novel putrescine exporter SapBCDF of *Escherichia coli*. J Biol Chem
291:26343-26351.
13.   Igarashi K, Ito K, Kashiwagi K. 2001. Polyamine uptake systems in *Escherichia coli*. Res Microbiol
152:271-8.
14.   Kurihara S, Suzuki H, Oshida M, Benno Y. 2011. A novel putrescine importer required for type 1
pili-driven surface motility induced by extracellular putrescine in *Escherichia coli* K-12. J Biol
Chem 286:10185-92.
15.   Hogins J, Xuan Z, Zimmern PE, Reitzer L. 2023. The distinct transcriptome of virulence-associated
phylogenetic group B2 *Escherichia coli*. Microbiol Spectr. doi:10.1128/spectrum.02085-
23e0208523. doi:10.1128/spectrum.02085-23.
16.   Schneider BL, Hernandez VJ, Reitzer L. 2013. Putrescine catabolism is a metabolic response to
several stresses in *Escherichia coli*. Mol Microbiol 88:537-50.
17.   Schneider BL, Reitzer L. 2012. Pathway and enzyme redundancy in putrescine catabolism in
Escherichia coli. J Bacteriol 194:4080-8.
18.   Fukuchi J, Kashiwagi K, Yamagishi M, Ishihama A, Igarashi K. 1995. Decrease in cell viability due
to the accumulation of spermidine in spermidine acetyltransferase-deficient mutant of
Escherichia coli. J Biol Chem 270:18831-5.
19.   Sakamoto A, Sahara J, Kawai G, Yamamoto K, Ishihama A, Uemura T, Igarashi K, Kashiwagi K,
Terui Y. 2020. Cytotoxic mechanism of excess polyamines functions through translational
repression of specific proteins encoded by polyamine modulon. Int J Mol Sci 21.
20.   Schwan WR. 2011. Regulation of *fim* genes in uropathogenic *Escherichia coli*. World J Clin Infect
Dis 1:17-25.
21.   Karp PD, Ong WK, Paley S, Billington R, Caspi R, Fulcher C, Kothari A, Krummenacker M,
Latendresse M, Midford PE, Subhraveti P, Gama-Castro S, Muniz-Rascado L, Bonavides-Martinez
C, Santos-Zavaleta A, Mackie A, Collado-Vides J, Keseler IM, Paulsen I. 2018. The EcoCyc
Database. EcoSal Plus. 8(1):doi:10.1128/ecosalplus.ESP-0006-2018.
22.   Keseler IM, Mackie A, Santos-Zavaleta A, Billington R, Bonavides-Martínez C, Caspi R, Fulcher C,
Gama-Castro S, Kothari A, Krummenacker M, Latendresse M, Muñiz-Rascado L, Ong Q, Paley S,
Peralta-Gil M, Subhraveti P, Velázquez-Ramírez DA, Weaver D, Collado-Vides J, Paulsen I, Karp PD.

2016. The EcoCyc database: reflecting new knowledge about *Escherichia coli* K-12. Nucleic Acids
Research 45:D543-D550.

23.   Tabor H, Tabor CW. 1969. Partial separation of two pools of arginine in *Escherichia coli*;
preferential use of exogenous rather than endogenous arginine for the biosynthesis of 1,4-
diaminobutane. J Biol Chem 244:6383-7.

24.   Miyamoto S, Kashiwagi K, Ito K, Watanabe S, Igarashi K. 1993. Estimation of polyamine
distribution and polyamine stimulation of protein synthesis in *Escherichia coli*. Arch Biochem
Biophys 300:63-8.

25.   Iwadate Y, Golubeva YA, Slauch JM. 2023. Cation homeostasis: Coordinate regulation of
polyamine and magnesium levels in *Salmonella*. mBio 14:e0269822.

26.   Applebaum DM, Dunlap JC, Morris DR. 1977. Comparison of the biosynthetic and biodegradative
ornithine decarboxylases of *Escherichia coli*. Biochemistry 16:1580-4.

27.   Wu WH, Morris DR. 1973. Biosynthetic arginine decarboxylase from *Escherichia coli*. Purification
and properties. J Biol Chem 248:1687-95.

28.   Flores-Mireles AL, Walker JN, Caparon M, Hultgren SJ. 2015. Urinary tract infections:
epidemiology, mechanisms of infection and treatment options. Nat Rev Microbiol 13:269-84.

29.   Puebla-Barragan S, Renaud J, Sumarah M, Reid G. 2020. Malodorous biogenic amines in
Escherichia coli-caused urinary tract infections in women-a metabolomics approach. Sci Rep
10:9703.

30.   Satink HP, Hessels J, Kingma AW, van den Berg GA, Muskiet FA, Halie MR. 1989. Microbial
influences on urinary polyamine excretion. Clin Chim Acta 179:305-14.

31.   Keay SK, Birder LA, Chai TC. 2014. Evidence for bladder urothelial pathophysiology in functional
bladder disorders. Biomed Res Int 2014:865463.

32.   Alteri CJ, Smith SN, Mobley HL. 2009. Fitness of *Escherichia coli* during urinary tract infection
requires gluconeogenesis and the TCA cycle. PLoS Pathog 5:e1000448.

33.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M, Wanner BL,
Mori H. 2006. Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the
Keio collection. Mol Syst Biol 2:2006 0008.

34.   Miller JH. 1972. Experiments in Molecular Genetics. Cold Spring Harbor Laboratory, Cold Spring
Harbor, NY.

35.   Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in *Escherichia coli*
K-12 using PCR products. Proc Natl Acad Sci U S A 97:6640-5.

36.   Hogins J, Zimmern PE, Reitzer L. 2023. Genome sequences of seven clade B2 *Escherichia coli*
strains associated with recurrent urinary tract infections in postmenopausal women. Microbiol
Resour Announc doi:10.1128/mra.00035-23:e0003523.

37.   Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods 25:402-8.

38.   McCarthy DJ, Chen Y, Smyth GK. 2012. Differential expression analysis of multifactor RNA-Seq
experiments with respect to biological variation. Nucleic Acids Res 40:4288-97.

39.   Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. Bioinformatics 26:139-40.

Table 1. Differentially expressed genes that are proposed to contribute to putrescine homeostasis.

A positive number means more transcripts in the Δ*speB* (lower putrescine) strain.

Gene (function)	log$_2$FC	FDR
Polyamine synthesis		
speA (putrescine)	0.99	0.003
speC (putrescine)	1.43	3E-4
speD (spermidine)	-0.83	0.01
speF (putrescine)	1.05	0.007
rpmH (inhibitor of SpeA and SpeC)	-2.77	2E-4
rpsT (inhibitor of SpeA and SpeC)	-1.92	8E-4
Polyamine transport		
*potABCD** (spermidine)	-2.17	3E-4
potE (putrescine)	2.07	2E-4
*potFGHI** (putrescine)	1.41	4E-4
plaP (putrescine)	1.11	0.005
*sapBCDF** (putrescine export)	-1.18	8E-4
Arginine synthesis		
argA_2 (synthesis)	1.77	0.001
argB (synthesis)	1.87	0.002
argC (synthesis)	2.86	0.005
argD (synthesis)	1.41	0.001
argF (synthesis)	1.63	0.01
argG (synthesis)	2.32	0.002
argH (synthesis)	0.97	0.001
argI (synthesis)	1.95	0.018
argR (repressor of arginine regulon)	-1.33	4E-4

Glutamate generation		
glsA (glutamine degradation)	2.16	5E-4
glnG (regulation of ammonia assimilation)	-1.59	0.022
TCA cycle/electron transport		
acnA (TCA cycle)	-1.04	8E-4
acnB (TCA cycle)	-1.96	2E-4
fumA (TCA cycle)	-1.50	6E-4
fumB (TCA cycle)	-1.49	0.003
gltA (TCA cycle)	-1.64	2E-4
icdA (TCA cycle)	-1.32	0.002
*sdhCDAB-sucABCD** (TCA cycle)	-3.88	5E-5
nuo operon* (electron transport)	-0.82	0.004
*menFDHBCE** (electron transport)	-0.8	0.005
*ubiEJB** (electron transport)	-0.93	0.004
*cyoABCD** (electron transport)	-1.41	9E-4
*hemCD** (heme)	-2.05	2E-4
Iron transport		
*exbBD** (transport of all iron chelates)	-2.33	4E-5
tonB (transport of all iron chelates)	-1.9	4E-5
*entCEBA** (enterochelin iron)	-5.2	7E-6
*fecABCDE** (ferric citrate)	-4.4	3E-5
fecIR (regulators, ferric citrate)	-4.4	6E-6
*feoABC** (ferrous iron)	-3.4	7E-4
fur (regulator, iron assimilation)	-0.40	0.11
Magnesium and phosphate transport		
mgtA (magnesium)	5.90	5E-6

phoQ (regulator, magnesium)	0.79	0.009
pstSCAB* (phosphate)	3.61	0.01
phoBR* (regulator, phosphate assimilation)	2.80	0.02

520 * Values for transcripts of the first gene of operon are given. Results for other genes of the

521 operon are in Supplemental Table 1

524 Table 1. Strains

Strain Name	Genotype	Reference
BLS77	W3110 Δ*puuR*::*cat*	(17)
BLS80	W3110 Δ*puuA*::*cat*	(17)
BLS88	W3110 Δ*patA* Δ*puuA*	(17)
CP2	W3110 Δ*patA*	(17)
IM26	W3110 Δ*speB*::*kan fliC-lacZ*	This study
IM27	W3110 Δ*speE*::*kan fliC-lacZ*	This study
IM28	W3110 Δ*speF*::*cat fliC-lacZ*	This study
IM29	W3110 Δ*speC*::*kan fliC-lacZ*	This study
IM34	W3110 Δ*speC* Δ*speF*::*cat fliC-lacZ*	This study
IM60	W3110 Δ*cadA*::*kan fliC-lacZ*	This study
IM61	W3110 Δ*speA*::*kan*	This study
IM62	W3110 Δ*patA* Δ*puuA* Δ*speE*::*kan*	This study
IM63	W3110 Δ*potE*::*kan*	This study
IM64	W3110 Δ*potF*::*kan*	This study
IM65	W3110 Δ*plaP*::*kan*	This study
IM66	W3110 Δ*puuP*::*kan*	This study
J15	W3110 Δ*speB* Δ*fliC*::*kan*	This study
SA1	W3110 Δ*fliC*::*kan*	This study

SA2	W3110 Δ*fimA::kan*	This study
SA3	W3110 Δ*fliC* Δ*fimA*	This study
SA4	W3110 Δ*hns::kan*	This study
SA5	W3110 Δ*ihfA::kan*	This study
SA6	W3110 Δ*lrp::kan*	This study
W3110-LR	*lacI*^{q} *lacL8*	Lab strain

Figure legends:

Figure 1 Pathways and enzymes of polyamine synthesis.

Figure 2. Surface motility of W3110-LR. (A) Surface motility of parental and derivative strains lacking the major subunits of the flagella (FliC), pili (FimA), or both. (B) Swimming motility with and without 0.5% glucose. (C) TEM images of cells of W3110-LR and W3110-LR Δ*fliC* Δ*fimA* taken from the movement's edge directly from surface motility plates.

Figure 3. Genetics of surface motility. (A) PDSM of mutants with defects in polyamine anabolic genes. (B) Diameter of surface movement of polyamine mutants after 36 hours. Error bars represent standard deviations for three independent replicates. Statistical analysis was performed using the Dunnett test of significance: *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$. All assays were performed in triplicate and representative images are shown. In this figure the Δ*speB* mutant was IM26.

Figure 4: Nutritional supplementation of Δ*speB* mutant. (A) Putrescine and spermidine

supplementation of wild type and Δ*speB* strains. (B) Diameter of surface movement of

polyamine mutants after 36 hours. Error bars represent standard deviations for three independent

replicates. Statistical analysis was performed using the Sadik test of significance: *, $P<0.05$; **,

$P<0.01$; ***, $P<0.001$. All assays were performed in triplicates and representative images are

shown. In this figure the Δ*speB* mutant was IM26.

Figure 5: Putrescine transport and surface motility. (A) Representative images of surface motility

for strains defective in putrescine transport genes. (B) Diameter of surface movement of

polyamine mutants after 36 hours. Error bars represent standard deviations for three independent

replicates. Statistical analysis was performed using the Dunnett test of significance: *, $P<0.05$;

**, $P<0.01$; ***, $P<0.001$. All assays were performed in triplicate and representative images are

shown.

Figure 6: Putrescine catabolism and surface motility. (A) Pathways and enzymes of putrescine

catabolism. (B) Motility diameter of putrescine catabolic mutants after 36 hours. Error bars

represent standard deviations for three independent replicates. Statistical analysis was performed

using the Dunnett test of significance: *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$. All assays were

performed in triplicates and representative images are shown. (C) Effect of loss of *speE* on the

patA puuA catabolic double mutant.

Figure 7: Surface motility and pili expression in W3110 and Δ*speB*. (A) W3110 Δ*speB* surface

motility with 0.0, 0.1, 1.0, and 4.0 mM exogenous putrescine. All assays were performed in

triplicate. Representative images are shown. (B) Average diameter of 3 separate surface motility

plates. The parental strain without putrescine is shown for reference. Significance was

determined by comparing the diameters of the ΔspeB mutants in the different concentrations of

putrescine compared to the parental W3110. One way ANOVA was used with Dunnett

hypothesis testing to determine p values, ***, *p <0.001*. (C) Reverse transcriptase-quantitative

PCR using primers targeting the *fimA* gene from cells grown in 0.0, 0.1, 1.0, and 4.0 mM of

supplemented putrescine. Double deltas were generated by normalizing the parental and the

ΔspeB mutant RNA libraries to *rpoD* then comparing *fimA* expression. Two-way ANOVA was

used to determine significance followed by Sadik hypothesis testing. ***, $p<0.001$. (D) Enzyme

linked immunosorbent assays using antibodies targeting pili (FimA) from cells grown with 0.0,

0.1, 1.0, and 4.0 mM of supplemented putrescine. Two-way ANOVA was used to determine

significance followed by FDR adjusting. ***, $p<0.001$. In this figure the ΔspeB mutant was J15.

Figure 8: Transmission electron micrographs of ΔspeB mutant cells after surface motility.

Representative images are shown and cells were removed from motility plates with the following

exogenous putrescine concentrations: (A) none, (B) 0.1 mM, (C) 1.0 mM, and (D) 4.0 mM. No

pili were observed with 0 and 0.1 mM putrescine. Optimal pili production was observed with 1.0

mM putrescine. Shorter and less pili were observed with 4.0 mM. The arrows in (D) point to

apparent pili fragments. The bar represents one micron. In this figure the ΔspeB mutant was J15.

Figure 9: Visual representation of the results from transcriptomic sequencing of W3110 and the

ΔspeB mutant's gene expression in media with and without 1.0 mM putrescine. (A)

Multidimensional scaling plot of the parental and ΔspeB mutant transcriptomes. When grown

with 1.0 mM putrescine (red), the Δ*speB* mutant transcriptome (triangles) are nearly identical to

the parental transcriptomes when grown without and with putrescine (blue and red circles,

respectively). When grown without putrescine, the Δ*speB* mutant transcriptome (blue triangles)

is greatly skewed from transcriptomes of the parental strain and the Δ*speB* mutant grown with

putrescine. (B) Heatmap of the top 100 most variable genes further demonstrates the

distinctiveness of the Δ*speB* mutant grown without putrescine and the similarities of the Δ*speB*

mutant transcriptome when grown with 1.0 mM putrescine supplementation and the parental

grown with or without putrescine. In this figure the Δ*speB* mutant was J15.

Figure 10: Transcriptomic sequencing of genes for the *fim* region and regulators that control *fim*

gene expression. (A) Expression of the *fim* genes in the parental W3110 and the speB mutant

with and without putrescine supplementation. Values were calculated by dividing the individual

replicates' counts per million (CPM) value for each gene by the mean CPM of that gene in

W3110 grown without putrescine and multiplying by 100 to yield the percent expression. Arrows

below the genes signify the known operons: *fimB* and *fimE* belong to single gene operons, while

fimAICDFGH belong to one operon. One way ANOVA was used to determine significance using

Dunnett hypothesis testing. **, $p < 0.01$; ***, $p < 0.001$. (B) Expression of some regulators

known to affect *fim* gene expression. Values were calculated as described in (A). ***, $p < 0.001$.

(C) Surface motility of W3110 and three regulator mutants (Δ*hns*, Δ*ihfA*, and Δ*lrp*). A gene found

to be significantly different by this transcriptomic analysis (*hns*) was confirmed to be important in

surface motility.

In this figure the Δ*speB* mutant was J15.

Figure 11: The deduced diversion of metabolism in a *speB* mutant away from energy metabolism

toward putrescine synthesis compared to the parental strain. The effects of low putrescine are

shown. Genes and processes (transport and transcriptional regulators) in red have fewer

transcripts, while those in black or blue have more. The dashed lines represent proposed

reduction in metabolic flux because of fewer transcripts from genes coding for the enzymes

involved. Operons are shown except for the larger operons or regulons. For example, pur is

meant to represent the unlinked genes that code for enzymes of purine synthesis. Table 1 or

Supplemental Table 1 should be consulted for specific genes and the quantitative change in

transcripts.

Supplemental Figure 1: Growth and swimming of polyamine anabolic mutants. (A) Growth

curves of wild type and putrescine deficient mutants in a shaking aerobic culture. (B)

Representative images of swimming.

Supplemental Table 1. Supplemental Table 1: Differential gene expression analysis of W3110

and Δ*speB*. J15 refers to the Δ*speB* mutant. In the tab labeled "W3110_0_vs_J15_1_ Putrescine"

W3110 was grown without putrescine, J15 was grown with 1.0 mM putrescine; positive fold

change values refer to increases in J15 gene expression.

Supplementary Figure 2: Expression graphs comparing the logCPM of the transcriptomes of the

average of the three replicates of the W3110 with and without 1 mM putrescine and the speB

mutant with and without 1 mM putrescine. A regression line was calculated and the correlation

between each set of transcriptomes noted on the graph. Higher R2 values indicate greater

similarity between the transcriptomes.

Figure 1

Figure 1 Pathways and enzymes of polyamine synthesis.

Figure 2

A **Surface Motility**

W3110   ΔfliC   ΔfimA   ΔfliC ΔfimA

B   **Swimming Motility**

C   **Transmission Electron Microscopy**

W3110   W3110 ΔfliC ΔfimA

- Glucose   + Glucose

Figure 2. Surface motility of W3110-LR. (A) Surface motility of parental and derivative strains lacking the major subunits of the flagella (FliC), pili (FimA), or both. (B) Swimming motility with and without 0.5% glucose. (C) TEM images of cells of W3110-LR and W3110-LR ΔfliC ΔfimA taken from the movement's edge directly from surface motility plates.

Figure 3

A

W3110   ΔspeC   ΔspeA   ΔspeB

ΔspeE   ΔspeF   ΔspeC ΔspeF

B

Figure 3. Genetics of surface motility. (A) PDSM of mutants with defects in polyamine anabolic genes. (B) Diameter of swarm colonies of polyamine mutants after 36 hours. Error bars represent standard deviations for three independent replicates. Statistical analysis was performed using the Dunnett test of significance: *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$. All assays were performed in triplicates and representative images are shown. In this figure "Δ*speB*" refers to IM26.

Figure 4

A

No Exogenous Putrescine   1.0 mM Putrescine   1.0mM Spermidine

W3110

$\Delta speB$

B

Figure 4 Legend

Figure 4: Nutritional supplementation of Δ*speB* mutant. (A) Putrescine and spermidine supplementation of wild type and Δ*speB* strains. (B) Diameter of swarm colonies of polyamine mutants after 36 hours. Error bars represent standard deviations for three independent replicates. Statistical analysis was performed using the Sadik test of significance: *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$. All assays were performed in triplicates and representative images are shown. In this figure "Δ*speB*" refers to IM26.

Figure 5

A

| $\Delta plaP$ | $\Delta potF$ | $\Delta potE$ | $\Delta puuP$ |

B

Figure 5: Putrescine transport and surface motility. (A) Representative images of swarming for strains defective in putrescine transport genes. (B) Diameter of swarm colonies of polyamine mutants after 36 hours. Error bars represent standard deviations for three independent replicates. Statistical analysis was performed using the Dunnett test of significance: *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$. All assays were performed in triplicates and representative images are shown.

Figure 6

A

B

C

W3110   ΔpatA ΔpuuA   ΔpatAΔpuuA ΔspeE

Figure 6 Legend

Figure 6: Putrescine catabolism and surface motility. (A) Pathways and enzymes of putrescine catabolism. (B) Motility diameter of putrescine catabolic mutants after 36 hours. Error bars represent standard deviations for three independent replicates. Statistical analysis was performed using the Dunnett test of significance: *, $P<0.05$; **, $P<0.01$; ***, $P<0.001$. All assays were performed in triplicates and representative images are shown. (C) Effect of loss of *speE* on the *patA puuA* catabolic double mutant.

Figure 7

A

0.0mM Putrescine   0.1mM Putrescine

1.0mM Putrescine   4.0mM Putrescine

B

C

RT-qPCR of *fimA*

D

ELISA

Figure 7 Legend

Figure 7: Surface motility and pili expression in W3110 and Δ*speB*. (A) W3110 Δ*speB* surface motility with 0.0, 0.1, 1.0-, and 4.0-mM exogenous putrescine. All assays were performed in triplicate. Representative images are shown. (B) Average diameter of 3 separate surface motility plates. The parental strain without putrescine is shown for reference. Significance was determined by comparing the diameters of the Δ*speB* mutants in the different concentrations of putrescine compared to the parental W3110. One way ANOVA was used with Dunnett hypothesis testing to determine p values. ***, *p* <*0.001*. (C) Reverse transcriptase-quantitative PCR using primers targeting the *fimA* gene from cells grown in 0.0, 0.1, 1.0, and 4.0 mM of supplemented putrescine. Double deltas were generated by normalizing the parental and the Δ*speB* mutant RNA libraries to *rpoD* then comparing *fimA* expression. Two-way ANOVA was used to determine significance followed by Sidak hypothesis testing. ***, *p*<0.001. (D) Enzyme linked immunosorbent assays using antibodies targeting pili (FimA) from cells grown with 0.0, 0.1, 1.0, and 4.0 mM of supplemented putrescine. Two-way ANOVA was used to determine significance followed by FDR adjusting. ***, *p*<0.001. In this figure "Δ*speB*" refers to J15.

Figure 8

Figure 8 Legend

Figure 8: Transmission electron micrographs of Δ*speB* mutant cells after surface motility. Representative images are shown and cells were removed from motility plates with the following exogenous putrescine concentrations: (A) none, (B) 0.1 mM, (C) 1.0 mM, and (D) 4.0 mM. No pili were observed with 0, and 0.1 mM putrescine. Optimal pili production was observed with 1.0 mM putrescine. Shorter and less pili were observed with 4.0 mM. The arrows in (D) point to apparent pili fragments. The bar represents one micron. The bar represents one micron. In this figure "Δ*speB*" refers to J15.

Figure 9: Visual representation of the results from transcriptomic sequencing of W3110 and the

ΔspeB mutant's gene expression in media with and without 1.0 mM putrescine. (A)

Multidimensional scaling plot of the parental and ΔspeB mutant transcriptomes. When grown

with 1.0 mM putrescine (red), the ΔspeB mutant transcriptome (triangles) are nearly identical to

the parental transcriptomes when grown without and with putrescine (blue and red circles,

respectively). When grown without putrescine, the ΔspeB mutant transcriptome (blue triagles) is

greatly skewed from transcriptomes of the parental strain and the ΔspeB mutant grown with

putrescine. (B) Heatmap of the top 100 most variable genes further demonstrates the

distinctiveness of the ΔspeB mutant grown without putrescine and the similarities of the ΔspeB

mutant transcriptome when grown with 1.0 mM putrescine supplementation and the parental

grown with or without putrescine. In this figure "ΔspeB" refers to J15.

Figure 10

A

fim Gene Expression

B

Regulator Gene Expression

C

W3110   Δhns   ΔihfA   Δlrp

Figure 10: Transcriptomic sequencing of genes for the *fim* region and regulators that control *fim*

gene expression. (A) Expression of the *fim* genes in the parental W3110 and the speB mutant

with and without putrescine supplementation. Values were calculated by dividing the individual

replicates' counts per million (CPM) value for each gene by the mean CPM of that gene in

W3110 grown without putrescine and multiplying by 100 to yield the percent expression. Arrows

below the genes signify the known operons: *fimB* and *fimE* belong to single gene operons, while

fimAICDFGH belong to one operon. One way ANOVA was used to determine significance using

Dunnet hypothesis testing. **, p <0.01; ***, p <0.001. (B) Expression of some regulators known

to affect *fim* gene expression. Values were calculated as described in (A). Two way ANOVA was

used to determine significance using Sadik hypothesis testing. ***, p <0.001. (C) Surface

motility of W3110 and three regulator mutants (Δ*hns*, Δ*ihfA*, and Δ*lrp*). A gene found to be

significant by the transcriptomic analysis (*hns*) was confirmed to be important in surface

motility.

In this figure "Δ*speB*" refers to J15.

Figure 11 Legned

Figure 11: The deduced diversion of metabolism in a *speB* mutant away from energy metabolism toward putrescine synthesis compared to the parental strain. The effects of low putrescine are shown. Genes and processes (transport and transcriptional regulators) in red have fewer transcripts, while those in black or blue have more. The dashed lines represent proposed reduction in metabolic flux because of fewer transcripts from genes coding for the enzymes involved. Operons are shown except for the larger operons or regulons. For example, pur is meant to represent the unlinked genes that code for enzymes of purine synthesis. Table 1 or Supplemental Table 1 should be consulted for specific genes and the quantitative change in transcripts.