Title

Arp2/3 iso-complexes limit t-tubule overgrowth to ensure myofiber formation and synchronous contraction

Authors and affiliations

Ana Raquel Pereira*[1], Ana da Rosa Soares*[1], Tianyang Liu[2], Josie Liane Ferreira[2,3], Graciano Leal[1], Inês Faleiro[1], Naoko Kogata[4], Michael Way[4,5], Carolyn A. Moores[2], Edgar R. Gomes[1]

* these authors contributed equally to this work

[1] Instituto de Medicina Molecular, Faculdade de Medicina, Universidade de Lisboa, Avenida Professor Egas Moniz, 1649-028 Lisboa, Portugal

[2] Institute of Structural and Molecular Biology, Birkbeck College, London, WC1E 7HX, UK

[3] Institute of Structural and Molecular Biology, Division of Biosciences, University College London, London WC1E 6BT, UK

[4] Cellular signalling and cytoskeletal function laboratory, The Francis Crick Institute, 1 Midland Road, London, NW1 1AT, UK

[5] Department of Infectious Disease, Imperial College, London W2 1PG, UK

Summary

T-tubules are long plasma membrane invaginations that form contact sites with two adjacent endoplasmic reticulum compartments to generate a skeletal muscle triad. Although the importance of t-tubules is well established, and abnormalities are found in multiple muscle disorders, the mechanisms of t-tubule growth are currently unknown. We found that the branched actin nucleator Arp2/3 regulates t-tubule organisation in myofibers. Cortical actin limits t-tubule growth, suggesting that it must be locally disassembled to allow the t-tubule to grow. T-tubule overgrowth is limited by Arpc5-containing Arp2/3 iso-complexes, but not those containing Arpc5L. Loss of Arpc5 results in malformed t-tubule clusters, impaired triads, and asynchronous muscle contraction, as found in multiple muscle disorders. Therefore, limiting t-tubule overgrowth has a role in triad formation and myofiber contraction, providing a possible pathophysiological mechanism for these muscle disorders.

38   **Graphical abstract**

39
40

41   **Keywords**

42   Myofiber; Arp2/3 iso-complexes; Arpc5; cortical actin; t-tubule; muscle contraction

Introduction

The contractile properties of skeletal muscle and its connection to the skeleton are essential for body movement, posture, and breathing. These vital functions are affected by multiple disorders and decline with age. Sarcomeres are the contractile units of muscle cells and are organised in myofibril bundles. Myofibrils contract synchronously upon calcium release from the endoplasmic reticulum (ER), also known as sarcoplasmic reticulum, due to its specialisation in muscle cells. Muscle contraction is initiated at the neuromuscular junction by local depolarisation of the plasma membrane, which propagates throughout the muscle cell (myofiber) (Franzini-Armstrong, 2018). Since myofibers are the longest and thickest cells in the human body, the depolarisation signal must be transmitted throughout their entire volume (Ward et al., 2009). This is achieved by plasma membrane invaginations named transverse tubules (t-tubules) that surround myofibrils, and form membrane contact sites with two adjacent endoplasmic reticulum compartments, in a structure termed triad (Al-Qusairi and Laporte, 2011; Kawaguchi and Fujita, 2024; Porter and Palade, 1957). Depolarisation of the t-tubule membrane triggers calcium release from the ER, a phenomenon termed excitation-contraction coupling (EC-coupling). Disruptions in triad assembly and morphology result in EC-coupling abnormalities, characteristic of multiple muscle disorders (Dowling et al., 2014). While the structure and function of the mature triad are well-defined, the underlying mechanisms governing the biogenesis and remodelling of t-tubules and triads – processes that require extensive yet precisely targeted remodelling of muscle cell membranes - remain elusive.

During mammalian development, t-tubules emerge in the embryo, forming a tubular network and becoming transversely organised only after birth (Takekura et al., 2001). Recent observations suggest that t-tubule biogenesis is initiated from plasma membrane invaginations mediated by Bin1, a BAR domain-containing protein, and caveolin-3-containing caveolae (Lemerle et al., 2023). Caveolin-3 is then released from the t-tubules by Cavin4 to remodel the t-tubule network (Lo et al., 2021). Endosomal membrane traffic and stabilisation at the sarcomeres have been proposed and implicated in t-tubule network formation and maintenance (Al-Qusairi and Laporte, 2011; Hall et al., 2020). However, how t-tubules grow and how this growth impacts triad function is unknown.

Interactions between the plasma membrane and cortical actin are pivotal in membrane remodelling during fundamental cellular processes such as endocytosis and cell migration (Svitkina, 2020). The cortical actin cytoskeleton consists of both linear and branched actin filaments, the latter of which is nucleated by the Arp2/3 complex composed of 7 evolutionary conserved subunits (Campellone et al., 2023). Notably,

some Arp2/3 subunits are encoded by more than one gene in higher Eukaryotes, yielding possibly eight unique Arp2/3 complexes in mammals with different properties (Abella et al., 2016; Cao et al., 2024; Galloni et al., 2021; Pizarro-Cerdá et al., 2017). It has been previously shown that this diversity in Arp2/3 iso-complexes is tailored for specific cellular functions such as cell migration, viral transport, or T-cell activation (Abella et al., 2016; Fäßler et al., 2023; Sadhu et al., 2023). In skeletal muscle, we previously demonstrated that Arp2/3 complexes containing Arpc5L drive nuclei positioning to the periphery of myofiber by promoting desmin-dependent myofibril crosslinking to generate centripetal forces on the nucleus (Roman et al., 2017). We also showed that Arpc5 plays a crucial role in triad organisation by an unknown mechanism.

Here we explored how Arpc5-containing iso-complexes are involved in triad organisation. We found that t-tubules are dynamic, and Arpc5 and cortical actin limit t-tubule overgrowth. Disruption of Arpc5, but not Arpc5L containing Arp2/3 complex, results in defects in triad formation and asynchronous muscle contraction, as observed in multiple muscle disorders.

Results

T-tubules are dynamic in mature and developing myofibers

To understand how the t-tubule network is established in myofibers, we first
evaluated t-tubule dynamics. For this, we used matured myofibers isolated from adult
mice or *in vitro* differentiating myofibers, incubated with a plasma membrane lipid dye
(CellMask), which labels t-tubules, and performed time-lapse microscopy (Takekura et
al., 2001). We observed that t-tubules exhibit growing and shrinking movements in both
isolated myofibers (Figure 1A, Video S1) and developing myofibers (Figure 1B, Video
S2). The mean growing speed of t-tubules in developing myofibers was 1.2 µm/sec
(Figure 1B). T-tubules are known to establish membrane contact sites with the
endoplasmic reticulum (ER), forming triads (Porter and Palade, 1957). Moreover, the ER
of undifferentiated cells is highly dynamic (Lee and Chen, 1988; Nixon-Abell et al., 2016).
We therefore wondered if the growing t-tubules were associated with ER or triad
dynamics. We first looked at ER (GFP-KDEL) and t-tubule dynamics in differentiating
myofibers. Although we observed that the ER of developing myofibers is dynamic, ER
movement did not correlate with growing t-tubules and vice-versa (Figure 1C, Video S3).
To address if t-tubules are dynamic at the triad, we imaged the movement of t-tubules
and Triadin-GFP, an ER transmembrane protein that localises explicitly at triads
(Sébastien et al., 2020). Triadin-GFP exhibits a negligible average speed of 0.015
µm/sec compared to the t-tubule growing average speed of 1.2 µm/sec (Figure 1D, Video
S4). Additionally, in 100% of growing t-tubules, we did not observe Triadin-GFP moving
with the t-tubule (N=39). These results indicate that t-tubules grow in developing and
fully matured myofibers, and this growth is independent of ER and triad dynamics.

Inhibition of the Arp2/3 complex promotes t-tubule growth

Our previous results reported that triad organisation during myofiber formation
depends on the Arp2/3 complex (Roman et al., 2017). Building upon these initial
observations, we hypothesise that the Arp2/3 complex is required for t-tubule growth. To
this end, we established a cellular system to study t-tubule dynamics. For this purpose,
we generated a stable cell line of C2C12 myoblasts expressing GFP-Bin1 (GFP-Bin1
myoblasts) since expression of Bin1 is sufficient to induce t-tubules, as previously
reported (Video S5) (Lee et al., 2002). We treated GFP-Bin1 myoblasts with the Arp2/3
inhibitor CK-666, and contrary to our hypothesis, we observed a significant increase in
the number of growing t-tubules after CK666 treatment (Figure 2A; Video S5 and S6).
This effect was not only observed in GFP-Bin1 myoblasts since treatment of developing

myofibers with CK666 also increased the number of growing t-tubules (Figure 2B; Video

S7 and S8). Together, these results show that Arp2/3 inhibition promotes t-tubule

overgrowth.

We previously showed that the Arp2/3 complex containing Arpc5, but not its

isoform Arpc5L, is involved in triad organisation (Roman et al., 2017). To test if Arpc5

and Arpc5L had differential roles on t-tubule growth, we performed knock-down (KD)

experiments with siRNA targeting either Arpc5 or Arpc5L in GFP-Bin1 myoblasts or

developing myofibers (Figure S1). Quantification of the number of growing t-tubules

revealed that depletion of Arpc5, but not Arpc5L, results in significantly more growing t-

tubules than control in both GFP-Bin1 myoblasts and in developing myofibers (Figure

2C and 2D; Video S9-S14). Overall, these results suggest that Arp2/3 complexes

containing the Arpc5 but not the Arpc5L limit t-tubule growth.

F-Actin does not accumulate near growing t-tubules

It has been previously shown that actin cytoskeleton and Arp2/3 can modulate

membrane tubulation in multiple organelles (Allard et al., 2021). Since our results

suggest that the Arp2/3 complex has a role in limiting t-tubule growth, we next wondered

if there was actin enrichment at the growing t-tubules. To this end, we imaged actin

filaments (F-actin) using either LifeAct-mCherry or mScarlet-Utrophin reporters in GFP-

Bin1 myoblasts (Figure 3A and 3C). As expected, we observed F-actin throughout the

cytoplasm, forming actin bundles (actin cables or stress fibers), filopodia, and

lamellipodia, and enriched around some intracellular vesicles (Video S15 and S16). To

quantify actin dynamics in the vicinity of growing t-tubules, we measured the average

intensity of F-actin fluorescence at a small ROI close to the growing t-tubule over time.

We did not observe F-actin enrichment during t-tubule growth, suggesting that F-actin is

not nucleated near growing t-tubules (Figure 3B and 3D; Video S15 and S16). To further

investigate the organisation of F-actin close to t-tubules, we performed cryo-correlative

Light and Electron Microscopy (cryo-CLEM) and cryo-electron tomography (cryo-ET) of

GFP-Bin1 myoblasts in regions containing t-tubules (Figure S2, 3E-G). We observed

unbranched F-actin forming bundles or meshes throughout the different volumes (Figure

3H, S3A and S3B, Video S17-S19). However, when we analysed slices of the volume at

different z positions, no distinct arrangement of actin filaments was observed near the t-

tubules when compared to other cytoplasmic regions (Figure 3I, S3C-J). Importantly, we

observed very few actin branches around t-tubules (Figure 3I), and we did not observe

more branched actin close to the t-tubules when compared to other regions of the

cytoplasm (Figure S3C-J, compare branched actin in blue versus red ROI). This
contrasts to what is observed for clathrin-mediated endocytosis, where branched actin
is enriched near the endocytic membrane (Collins et al., 2011; Serwas et al., 2022).
Overall, these data suggest that branched F-actin does not accumulate near growing t-
tubules, leading us to explore contributions by the Arp2/3 network to t-tubule dynamics
elsewhere in the cell.

Perturbation of cortical actin promotes t-tubule growth

T-tubules are contiguous with the plasma membrane, and the t-tubule surface area
is about seven times the myofiber surface area (Peachey, 1965). This suggests that
during development, a large amount of membrane must be available to invaginate and
form t-tubules. Plasma membrane morphology is regulated by cortical actin, the major
component of the cell cortex that forms an actomyosin network underneath the plasma
membrane (Chugh and Paluch, 2018; Svitkina, 2020). Since the Arp2/3 complex is also
involved in cortical actin remodelling, we hypothesised that perturbing cortical actin
impacts t-tubule growth (Bovellan et al., 2014). To test this hypothesis, we interfered with
cortical actin by inhibiting non-muscle myosin-2, using para-aminoblebbistatin
(blebbistatin), and quantified the number of growing t-tubules before and after treatment
(Laplaud et al., 2021; Várkuti et al., 2016). We found that both in GFP-Bin1 myoblasts
and in developing myofibers, blebbistatin treatment led to a significant increase in the
number of growing t-tubules (Figure 4A and 4B; Video S20-23). We also interfered with
cortical actin by inducing a glycerol hyperosmotic shock, which leads to a rapid
remodelling of cortical actin due to a reduction of cell volume (Di Ciano et al., 2002). We
quantified the number of growing t-tubules before and after osmotic shock in both
myoblasts (Figure 4C; Video S24 and S25) and developing myofibers (Figure 4D; Video
S26 and S27), and observed a significant increase in the number of growing t-tubules
upon osmotic shock. These results show that perturbations of the cortical actomyosin
network promote t-tubule growth, suggesting that cortical actin limits t-tubule growth.

Arpc5 depletion leads to t-tubule clusters and impaired triad formation

Our data are consistent with a model in which Arp2/3 complexes containing Arpc5
mediate cortical actin, limiting t-tubule growth. In fully differentiated myofibers, t-tubules
establish two-sided connections with the ER, forming triads that become transversely
organised (Al-Qusairi and Laporte, 2011; Kawaguchi and Fujita, 2024; Porter and
Palade, 1957). Since Arpc5, but not Arpc5L is involved in transverse triad organisation
(Roman et al., 2017), we investigated the role of Arpc5 isoforms in the t-tubule network

of fully differentiated myofibers. To this end, we assessed t-tubule transverse alignment
in myofibers depleted for either Arpc5 or Arpc5L in fully differentiated myofibers, which
have transverse triads (Falcone et al., 2014; Roman et al., 2017). In control and Arpc5L
KD conditions, we observed transversely aligned t-tubules forming an evenly distributed
network throughout the myofiber (Figure 5A, Scramble and si Arpc5L). However, in
Arpc5 KD myofibers, we observed that t-tubules were not transversely aligned, and t-
tubules formed clusters (Figure 5A and 5C). T-tubule clusters were not an artefact of
CellMask as they were also observed upon immunostaining against Bin1, a t-tubule-
specific protein (Figure 5B). To better characterise t-tubule clusters, we performed
transmission electron microscopy (TEM) of stained and resin-embedded differentiated
myofibers. We observed a higher proportion of enlarged t-tubules in Arpc5 KD myofibers
when compared to the control (Figure S4A yellow arrows, S4B). Additionally, we noticed
that these enlarged t-tubules still form contact sites with the ER (Figure S4A blue arrows)
but only with one ER compartment. Accordingly, we observed a reduction of triads in
Arpc5 KD compared to the control (Figure S4A green arrows, S4C). Finally, we
quantified the number of Junctophilin-1 (JPH1) puncta, a marker of triads, in fully
differentiated myofibers and observed that Arpc5 KD myofibers, but not Arpc5L or
control, exhibited a reduced number of JPH1 puncta, suggesting impaired triad formation
(Figure 5D and 5E). Together, these results show that lack of Arpc5, which induces t-
tubule overgrowth, contributes to disruption of the t-tubule network, leading to abnormal
t-tubule-ER contact sites and triad impairment.

Arpc5 is required for synchronous excitation-contraction coupling

Triads are fundamental for coupling the excitation of the plasma membrane and t-
tubules to the release of calcium from the ER into the cytosol, resulting in muscle
contraction. This process is known as excitation-contraction coupling (EC-coupling).
Since we observed that Arpc5 depletion induces t-tubule clusters and impaired triads,
we assessed the impact of the loss of Arpc5 on EC-coupling in myofibers. We first
measured cytosolic calcium levels upon electrical pulses by imaging differentiated
myofibers infected with an AAV expressing the calcium reporter GCaMP (Chen et al.,
2013). As expected in the controls, we observed GCaMP fluorescence transients
corresponding to calcium release and uptake from the ER, synchronised with electrical
stimulation (Figure 6A and 6B, scramble). In contrast, most myofibers depleted for Arpc5,
but not Arpc5L exhibited abnormalities in calcium transients, which were not
synchronised with the electrical stimulation pulses (Figure 6A and 6B). We then
investigated whether this asynchrony between excitation and calcium transients impairs

contraction. For this purpose, we performed single-myofiber contraction tracing and
found that myofibers could still contract when Arpc5 and Arpc5L are depleted, as in the
scramble control (Figure 6C and 6D). However, there was a dramatic asynchrony
between electrical stimulation and myofiber contraction in the Arpc5 KD myofibers but
not in the Arpc5L and scramble conditions (Figure 6C and 6E). This observation
suggests that lack of Arpc5 results in defects in EC-coupling that leads to asynchronous
muscle contraction, highlighting the role of Arpc5 containing Arp2/3 complexes and t-
tubule overgrowth in triad function.
243

Discussion

Discussion

T-tubules and their connection with the endoplasmic reticulum to form triads are

fundamental to muscle contraction and are disrupted in multiple muscle disorders. We

found that t-tubules are dynamic, and Arpc5-containing Arp2/3 iso-complexes limit their

growth through the modulation of cortical actomyosin. T-tubule overgrowth upon

disruption of Arpc5, but not Arpc5L, results in t-tubule clusters, impaired triads, and

asynchronous muscle contraction, thus providing a pathophysiological mechanism for

muscle disorders with impaired triad function (Dowling et al., 2014).

Keith Porter and George Palade originally identified triads, which were the first

description of a plasma membrane contact site (Porter and Palade, 1957). We found that

t-tubules are dynamic, even in matured myofibers isolated from adult mice, although to

a lesser extent than in developing myofibers. T-tubule dynamics are independent of ER

dynamics and triads, which further supports that ER organisation precedes t-tubule

network formation, as observed during muscle development (Flucher et al., 1993). T-

tubule dynamics in matured myofibers may be involved in triad homeostasis and repair

during muscle injury.

Since the t-tubule area comprises up to seven times more of the myofiber cell

membrane, considerable membrane remodelling is required to form a t-tubule network

during development (Peachey, 1965). Our results suggest that cortical actin prevents t-

tubule overgrowth since perturbation of cortical actin through non-muscle myosin-2

inhibition, hyperosmotic shock, and Arp2/3 complex inhibition lead to more growing t-

tubules. Therefore, we hypothesise that to grow a t-tubule, cortical actin must be

disassembled to allow plasma membrane release and availability for the t-tubule to grow.

Upon stretch release of adherent cells, tubulation of the plasma membrane has also

been observed, suggesting that membrane release facilitates tubulation, although the

role of cortical actin has not been addressed (Kosmalska et al., 2015). The formation of

tubules from membranes directly depends on membrane release as elegantly

demonstrated using giant lipid vesicles. Excess membrane area created by a decrease

in vesicle volume leads to membrane release and tubulation towards the lumen of the

vesicle, similar to t-tubule growth (Li et al., 2011). In myofibers, disruption of cortical actin

probably results in the excess plasma membrane required for t-tubule growth.

Furthermore, the observed tubulation in lipid vesicles does not involve any external

forces (Li et al., 2011). In myofibers, during t-tubule growth, we did not detect any

enrichment of actin, using multiple F-actin fluorescent probes. Using cryo-CLEM, we also

did not observe evidence for interaction between F-actin and t-tubules, nor accumulation

of branched actin. Although we cannot rule out that external forces are involved in t-

tubule growth, our data does not support a need for actin-related force generation.

Although tubulation can occur in the absence of proteins in lipid vesicles upon a decrease
in membrane volume, Bin1 modulates membrane tubulation in the absence of actin
without a reduction of membrane volume (Picas et al., 2014). Bin1, together with other
proteins such as dynamin-2, are therefore of importance for t-tubule network
remodelling, probably providing the required forces to maintain tubulation, together with
the cortical actin membrane release mechanism we describe here (Lionello et al., 2022;
Picas et al., 2014).

Branched cortical actin is nucleated by the Arp2/3 complex at the cell cortex and is
crosslinked by non-muscle myosin-2 to generate cortical tension essential for plasma
membrane remodelling and cell shape (Chugh and Paluch, 2018; Svitkina, 2020).
Downregulation of the Arp2/3 complex reduces cortical actin and weakens the coupling
of the plasma membrane with the cell cortex (Bovellan et al., 2014). Our results suggest
that Arp2/3 iso-complexes containing Arpc5 or Arpc5L might regulate cortical actin in a
specific manner. However a role for a specific Arp2/3 isocomplex on cortical actin has
yet to be described. During lamellipodia formation, Arp2/3 iso-complexes containing
Arpc5 or Arpc5L have a distinct role in actin network architecture and branched actin
structure modulation (Fäßler et al., 2023). We speculate that similar distinct actin network
arrangements might occur at the cell cortex to limit t-tubule formation through Arpc5 but
not Arpc5L containing Arp2/3 complexes.

Our work implicates a role in limiting t-tubule dynamics for effective triad function
and synchronous contraction. T-tubule overgrowth might result in defects on membrane
contacts with the ER, impairing the formation and maintenance of functional triads.
Interfering with the t-tubule network by depletion of Cavin4 also leads to t-tubule cluster
accumulation and disruption of triads (Lo et al., 2021). Also, t-tubule overgrowth and triad
abnormalities have been described in aged human muscle (Boncompagni et al., 2006).
Furthermore, several muscle disorders with impaired triads also exhibit an overgrowth
of t-tubule clusters, particularly in Mitsugummin25, Dysferlin, Myotubularin, or Dynamin-
2 mutants (Al-Qusairi and Laporte, 2011; Dowling et al., 2014). Therefore, modulation of
t-tubule growth and cluster formation can be a common target to ameliorate or prevent
these diseases.

Methods

T-tubule labelling in matured myofibers

All procedures using animals were approved by the Institutional Ethics Committee and followed the guidelines of the Portuguese National Research Council Guide (DGAV). Ex vivo isolated single myofibers from Extensor Digitorum Longus (EDL) muscles were generated using the protocol previously described (Falcone et al., 2014). EDL muscle was explanted from 8- to 12-week-old male or female C57BL/6J mice and then digested in DMEM (Invitrogen) containing 0.2% type I collagenase (Sigma) for 2 h at 37°C. Mechanical dissociation of fibers was performed using a thin Pasteur pipette in a transilluminating-fluorescent stereomicroscope. Single fibers were collected and plated in ibidi-35mm-dishes pre-coated with 1% matrigel in Iscove's Modified Dulbecco's Media (IMDM, Invitrogen) with Glutamax. After plating, cells were covered with an upper coating of ice-cold 50% matrigel coating in IMDM with Glutamax. Plated myofibers were maintained in 10% horse serum (Fisher Scientific) plus IMDM with Glutamax for the duration of the experiment. To visualize t-tubule dynamics, plated myofibers were labelled for 30 minutes with CellMask Orange (1 μM, Invitrogen).

In vitro myofibers

All animal procedures were approved by the institutional ethics committee and followed the guidelines of the National Research Council Guide for the care and use of laboratory animals. In vitro myofibers were differentiated as previously described (Pimentel et al., 2017) from primary myoblasts isolated from 5 to 7-day-old mixed-gender C57BL/6 mice. Tibialis anterior (TA), EDL, Gastrocnemius and Quadriceps muscles were dissected and kept in ice-cold PBS. Muscles were digested mechanically using surgical scissors and incubated for 90 min at 37ºC with agitation in collagenase type V (Sigma) and dispase II (Roche) in DPBS. The digestion was stopped by mixing 10% FBS 1% Penicillin-Streptomycin IMDM with Glutamax. The cells suspension was centrifuged for 5 min at 75 x g to remove debris followed by a second centrifugation for 5 min at 350 x g. After centrifugation, cells were resuspended in the same medium and pre-plated for 4 hours to allow for fibroblasts adherence. The cells remaining in suspension were collected, centrifuged for 5 min at 350 x g and plated at a density of 220 000 cells/ml in 20% FBS IMDM with Glutamax in ibidi 35 mm dishes pre-coated with 1% matrigel in IMDM with Glutamax. After three days of proliferation, medium was replaced by 10% horse serum IMDM with Glutamax supplemented with 100 ng/ml recombinant agrin to induce

346 differentiation. An ice-cold upper coating of 50% matrigel in 10% horse serum IMDM with

347 Glutamax was added the following day and fresh medium was added. Experiments were

348 performed either at 4-5 or 7-8 days after differentiation.

350 siRNAs, plasmids and transfections

Triadin-GFP (Trisk95-GFP, (Fourest-Lieuvin et al., 2012)) or GFP-KDEL (Campbell-

Valois et al., 2012) plasmids were transfected on in vitro myofibers three days after

plating with 1 µg of DNA per ibidi-35mm-dish of the appropriate plasmid using

Lipofectamine 2000 (Thermo), according to manufacturer's protocol.

GFP-Bin1 myoblasts were transfected with 1 ug plasmid/ibidi-35mm-dish of LifeAct-

mCherry (Janota et al., 2022) or mScarlet-I-UtrCH (gift from Dorus Gadella, Addgene

plasmid #98823; http://n2t.net/addgene:98823; RRID:Addgene_98823 (Chertkova et al.,

2017)) one day after plating, using Lipofectamine 2000 (Thermo), according to

manufacturer's protocol. Cells were imaged by vtiSIM microscope on the next day (see

below).

For Arpc5 isoforms depletion, in vitro myofibers were transfected with Thermo Fisher

Silencer® Select siRNAs for Arpc5 (ID#s85623) and Arpc5L (ID#s211945), 3 days after

plating with 20nM/ibidi-35mm-dish of the appropriate siRNA using Lipofectamine

RNAiMAX. Silencer® Select Scramble(ID #4390843) was used as a negative control.

366 Generating the GFP-Bin1 myoblast stable cell line

GFP-Bin1 stable cell line of C2C12 myoblasts (GFP-Bin1 myoblasts) was generated

using a pLVX-puromycin vector containing Bin1. The lentiviral vector was constructed

with VectorBuilder. The vector ID is VB220728-1365nbr, which can be used to retrieve

detailed information about the vector on vectorbuilder.com. Lentivirus were produced in

HEK 293T cells. To generate stable cell lines expressing exogenous GFP-Bin1, C2C12

myoblasts were plated in a 6-well plate. The cells with a 10% confluency were carefully

covered with 0.5 ml of filtered medium containing the lentiviruses and 0.5 ml of complete

medium was added to myoblasts (1 ml total volume). The medium containing lentiviruses

was left over day, then 1 ml of complete fresh medium was added. Cells were allowed

to grow until 80% of confluency. Subsequently, the sorted myoblasts were allowed to

expand, and the process was repeated for three rounds of sorting to achieve a

homogeneous cell population.

GFP-Bin1 myoblasts were cultured in Dulbecco's modified Eagle's medium (DMEM) with

glutamine (Gibco) supplemented with 10% fetal bovine serum (FBS; Gibco) and

penicillin-streptomycin (100 U/ml:100 µg/ml; Gibco), at 37°C under 5% CO_2.

Fluorescence microscopy

Airyscan images were acquired using a Zeiss LSM980 Airyscan2 microscope, equipped

with a 63x Plan-Apochromat DIC oil objective (NA 1.4). Imaging was performed at 37°C

with 5% of CO_2 using the ZEN 3.8 blue edition. To measure t-tubule growth, images

were recorded every three seconds. Airyscan reconstruction was performed using

automatic filter settings.

For instant structured illumination microscopy (iSIM) imaging, the vtiSIM module

(Visitech) equipped on a Nikon Tie microscope with a 100x AC Oil CFI SR HP Apo TIRF

objective, and a Photometrics Prime BSI Express Scientific sCMOS camera was used.

Timelapse imaging was performed at 37°C with 5% of CO_2.

To visualize t-tubule dynamics in myofibers, the membrane lipid dye CellMask Orange

(1 µM, Invitrogen) was added to the myofibers 5-10 min before imaging. T-tubule and/or

ER dynamics were recorded by timelapse imaging using Airyscan microscopy (see

above).

To visualize actin dynamics in the vicinity of growing t-tubules, GFP-Bin1 myoblasts

transfected with LifeAct-mCherry or mScarlet-I-UtrCH were imaged by vtiSIM timelapse

imaging every second for three minutes.

To visualize t-tubule network and t-tubule clusters, developed in vitro myofibers

(differentiation day 7) were incubated with 1 µM of CellMask Deep red (Invitrogen).

Spinning disk confocal imaging was performed using a Zeiss Axio Observer microscope

equipped with a Yokogawa CSU-X1 scanning unit, an Evolve 512 EMCCD camera and

a 63x Plan-Apochromat DIC oil objective (N.A. 1.4), using the ZEN 3.2 blue edition

software. Cells were kept at 37°C and 5% of CO_2. Myofibers with more than three t-

tubule clusters were visually quantified using Fiji software and the percentage of cells

with t-tubule clusters was plotted using graphpad.

Drug treatments

Drug treatments

Cells were treated with 100 µM of CK666 (Sigma) for Arp2/3 inhibition. A final
concentration of 125 µM of para-aminoblebbistatin (Motorpharma) was used to block
myosin-II. The number of growing t-tubules was performed immediately after drug
treatments for GFP-Bin1 myoblasts and after five minutes for myofibers. T-tubule
dynamics were recorded by timelapse imaging using Airyscan microscopy. GFP-Bin1
myoblasts treated with para-aminoblebbistatin were recorded by timelapse imaging
using vtiSIM microscopy.

Glycerol treatment

Myoblasts and developing myofibers were exposed to 225 mM and 45 mM of glycerol,
respectively, to induce hyperosmotic shock. The number of growing t-tubules was
performed three minutes after glycerol treatment for GFP-Bin1 myoblasts and after five
minutes for myofibers. Myofiber t-tubule dynamics were recorded by timelapse imaging
using Airyscan microscopy. GFP-Bin1 myoblasts t-tubule dynamics were recorded by
timelapse imaging using vtiSIM microscopy.

Immunostaining
Developed myofibers (differentiation day 7) were fixed in 4% paraformaldehyde in PBS
for 10 minutes at room temperature and permeabilized with 0.5% Triton X-100 for 5
minutes. Cells were blocked in a solution containing 10% goat serum, 5% BSA and
VisUBlock Mouse on Mouse Blocking Reagent (R&D Systems, VB001). Primary and
secondary antibodies were diluted in 0.1% saponin. Cells were incubated with primary
antibodies overnight at 4°C and washed three times for 10 minutes with PBS. Cells were
incubated with secondary antibodies for 1 hour at room temperature and washed three
times for 10 minutes with PBS. Cells were mounted with Fluoromount-G (Invitrogen).
Antibodies used for immunostaining were: anti-JPH1 Antibody (1:200 dilution,
Invitrogen, 40-5100); and anti-rabbit Alexa Fluor 488 secondary antibody (1:400, Life
Technologies). Phalloidin conjugated with an Alexa Fluor 647 A22287(1:200, Life
Technologies) was used to stain filamentous actin. Myofibers were imaged using the
vtiSIM microscope.

Image analysis

All images were processed in Fiji (Schindelin et al., 2012). For cell drift correction we used the Fiji plugin Fast4Dreg (Pylvänäinen et al., 2023).

For t-tubule growth speed measurements, manual tracking of the tip of the t-tubule during the timelapse was performed using Fiji manual tracking.

The number of growing t-tubules was manually recorded using the Fiji cell counter tool. This value was then normalized for the cell area. All quantifications were performed within two-minute timelapses.

To measure Triadin-GFP mean speed in developing myofibers, we used Airyscan images of Triadin-GFP timelapses acquired every three seconds for two minutes. Since Triadin-GFP localizes as distinct spots, we used the Fiji plugin Trackmate (Ershov et al., 2022) to track Triadin-GFP spots over time in myofibers. To detect Triadin-GFP spots we used the LoG detector with an estimated object diameter of 0.5 µm and a quality threshold of 5, 10 or 20. We then linked the spots to generate tracks, by using the LAP tracker with a max distance of 0.5 µm and no gap closing. To have enough datapoints to calculate speed trajectories, we only considered tracks that lasted more than 10 frames (30 seconds). The resulting tracked Triadin-GFP spots were then used to calculate mean speed.

To quantify actin dynamics in the vicinity of growing t-tubules, we manually selected a ROI around GFP-Bin1 growing t-tubules (as the ones represented in Figure 2A and 2C), including a few timepoints before the GFP-Bin1 signal was observed using Fiji software. We then extracted at each timepoint the average of fluorescence intensity per µm^2 for GFP-Bin1, LifeAct-mcherry or mScarlet-I-UtrCH using the Fiji software. Values were normalized for the initial corresponding average intensity value and these normalized intensity values per µm^2 over time were plotted using graphpad.

To automatically count the number of Junctophilin 1 (JPH1) puncta, a maximum projection of the Z-stack images that were in focus was first performed using Fiji. Then, a rectangular ROI with an area of 175.552 µm^2 was drawn along the myofiber length. To detect the puncta automatically, we used the LoG detector tool from the FIJI TrackMate plugin (Schindelin et al., 2012) LoG filters for puncta detection were a blob diameter of 0.5 µm and an intensity threshold of 5. To analyse the number of JPH1 puncta per µm^2, the number of detected puncta was normalized for ROI area and plotted.

Western Blot

Cell extracts were collected in 100 mM Tris-HCl pH 8 with 1% SDS. Samples were denatured in Laemmli buffer and heated at 95°C for 5 min. Total protein was resolved on a 10% precast polyacrylamide gel (Bio-Rad) and transferred to a nitrocellulose membrane for 75 min at 100 V. The membranes were blocked with Tris-buffered saline with 0.1% Tween 20 (TBS-T) containing 5% skim milk, followed by incubation with primary antibodies, overnight at 4°C. Membranes were washed and incubated with HRP-conjugated secondary antibodies for 1 hour at room temperature and washed three times before image acquisition. Images were acquired using Amersham ImageQuant™800 (Cytiva). Antibodies used: anti-Arpc5 (1:500 Santa Cruz Biotechnology, sc-166760); anti-Arpc5L (1:500, Proteintech, 22025-1-AP); anti-Vinculin (1:1000, Sigma, V9131); anti-rabbit HRP (1:5000, Thermo Scientific) and anti-mouse HRP (1:5000, Thermo Scientific).

Cryo-ET sample preparation

GFP-Bin1 myoblasts were grown to approximately 60% confluency and then diluted to a confluency of 10%. EM grids (Quantifoil R3.5/1 Au 200 mesh) were glow-discharged using a plasma cleaner (PLASMAFLO PDC-FMG-2, Harrick Plasma) and then placed in a 35mm MatTek dish in a flow hood. 300 µl of 5 µg/ml Fibronectin (Sigma-Aldrich, 341635) was added to the dish, covering the EM grids. The dish with the grids inside was then sterilized under UV light for 1.5 hours. After simultaneous coating and sterilization, the fibronectin was removed from the MatTek dish. The grids were then washed twice with PBS and stored in approximately 300 µl of medium (DMEM high glucose+Pyruvate+L-Glutamine; 10% FBS + 1% PenStrep).

Once the cells were ready, the medium in the grid-containing dish was removed, and diluted cells were added to the dish. The cells were seeded on the EM grids at 37°C 5% CO_2 overnight (~15 hours). The following day, grids were inspected under an inverted biological microscope (Medline Scientific™, 3650.0000) to check the cell density and ensure grid integrity. Grids with minimally broken areas and well-distributed cells were plunge frozen using Leica GP2 with back blotting for 15 seconds and an additional movement of 0.5 mm at 37°C and 90% humidity.

Correlative light and electron microscopy

Frozen grids were first loaded onto a Titan Krios microscope (Thermo Fisher Scientific) operated at an accelerating voltage of 300 kV with a K3 detector and a BioQuantum energy filter (Gatan). Grid atlas images were collected to check the ice quality and the cell density using EPU software (Thermo Fisher Scientific). After locating the spread cells, grids were transferred onto a cryo-light microscope (Leica THUNDER Imager EM Cryo CLEM). Z stack fluorescence and bright field images were taken of the located spread cells using LAS X software at 50X magnification. With the fluorescence images processed using THUNDER technology, individual tubular fluorescence signals were observed. The fluorescence images were saved for fluorescence microscopy (FM) and electron microscopy (EM) image correlation. The grids were then transferred back to a Titan Krios microscope (Thermo Fisher Scientific) operated at an accelerating voltage of 300 kV with Selectris X image filter (Thermo Fisher Scientific) and Falcon 4i camera (Thermo Fisher Scientific). Medium magnification atlases were taken and combined on each fluorescence-labelled cell for manual correlation and locating the regions of interest. Empty holes and dark features were used as markers to facilitate the manual correlation (Supplementary Figure 2B and 2C). After correlation, ROI with fluorescence signals were targeted for cryo-ET tilt series data collection. 114 tilt series were collected using TOMO5 software with a dose symmetric tilt scheme +/- 60° with the step size of 3° at the magnification of ×42,000, pixel size of 3.0 Å, dose rate of 12.6 e⁻/pixel/s, exposure time of 2.69s and total dose per micrograph of 3.66 e⁻/ $Å^2$.

Tomogram reconstruction

Tomograms were reconstructed using RELION 4.1 tomography pipeline (Zivanov et al., 2022). Raw data were imported into RELION followed by motion correction with RELION's own implementation of MotionCor2 (Zheng et al., 2017) and CTF correction with ctffind 4.1 (Rohou and Grigorieff, 2015). Tilt images with significant contamination and grid bars were manually excluded from further data processing in a Napari-based viewer (Exclude Tilt Images job). Then each tilt series was aligned using IMOD [CM4] patch tracking with Patch size 100 nm and Patch overlap 50%. The aligned tilt series were used to generate tomograms with a binned pixel size of 10 Å. Tomograms were denoised using CryoCARE to improve the signal-to-noise level (Buchholz et al., 2019). CryoCARE training was performed with 1200 sub-volumes per tomogram and a sub-volume dimension of 72 pixel. Upon inspecting the tomograms, we found that 58

contained vesicular as opposed to tubular membrane structures (as shown in the
Supplementary Figure 2G) and were excluded. 23 of the tomograms were discarded due
to the presence by non-vitreous ice and could not be used for segmentation. Additionally,
27 tomograms did not correlate well with fluorescence signals because of the ambiguity
of fluorescence and the lack of correlation markers, which posed challenges for manual
correlation. 6 tomograms displayed tubular structures at locations where fluorescence
was observed. From these, the 3 highest quality tomograms were selected for
segmentation and further analysis.

Tomogram Segmentation

Tomograms were processed using IsoNet to further improve the signal-to-noise level
(Liu et al., 2022). The following parameters were used: CTD deconvolve snrfalloff of 0.5,
mask making density_percentage of 100, std_percentage of 100, and z_crop of 0.17.
Then, IsoNet-processed tomograms were cropped to remove regions lacking t-tubule
structures to simplify segmentation. Segmentation was performed using the neural
network-based software Dragonfly (Dragonfly 2024.1 [Computer software]. Comet
Technologies Canada Inc., Montreal, Canada; software available at
https://dragonfly.comet.tech/). For each tomogram, actin and membrane structures were
manually annotated on 5-12 sub-areas across different slices of the tomogram. The
segmented data were then used to train multiple deep-learning models. The model
segmenting the highest number of actin filaments on test slices was selected to segment
the entire tomogram. The UNet++ model (2.5D, 3 slices, depth level of 5, and initial filter
count of 32) was used for tomograms 1 and 2. Additionally, a pretrained UNet model
provided by the Dragonfly team (depth level of 5 and initial filter count of 64) was utilized
for tomogram 3. The segmentations were manually cleaned and corrected in Dragonfly.
The segmented actin and membrane structures were rendered into a 3D contour mesh
and smoothed over five to ten iterations for 3D visualization in Figure 3I and
Supplementary Figure 3. Movies were created by exporting the segmented actin and
membrane structures as binary files, converting these into two MRC files, and then
importing them along with a contrast-inverted 3D tomogram into ChimeraX for
visualization (Pettersen et al., 2021).

Tomogram Analysis

For tomogram analysis, sub-tomograms proximal to the t-tubules and distant from the t-tubules were selected. The sub-volumes near t-tubules were constrained to regions no more than 200 nm from the t-tubule structures. Actin branches – inferred from the relative position of two connected filaments - within these boxes were identified, labelled, and counted. For control purposes, sub-volumes located more than 200 nm away from t-tubules were also identified, and suspected actin branches within these control sub-volumes were similarly labelled and counted. In total, 7 sub-volumes were selected to determine the number of actin branches near the t-tubules, and 3 sub-volumes were selected as controls.

Single-myofiber contraction and calcium transient assays

In vitro myofibers were plated on 35mm fluorodishes (World Precision Instruments). After 7-8 days of differentiation, developed myofibers were imaged using a Nikon Eclipse Ti microscope. Myofibers were kept at 37°C and CO_2 5% using an Okolab stage incubator throughout all experimental procedures. Electrical field stimulation using the MyoPacer system (Ionoptix) was used to induce excitation-contraction of myofibers using 10 Volts at 0.5 Hz frequency pulses with a duration of 20 msec.

To measure cytosolic calcium transients during electrical pulses, myofibers at differentiation day 3-4 were infected using 1 µl/ml of the AAV9-pAAV.CAG.GCaMP6s.WPRE.SV40 (Addgene viral prep # 100844-AAV9; the plasmid was a gift from Douglas Kim & GENIE Project (Chen et al., 2013)). After overnight incubation, the media was replaced with differentiation media (IMDM with Glutamax supplemented with 100 ng/ml recombinant agrin). After 7-8 days of differentiation, GCaMP fluorescence during electrical field stimulation was recorded using a 60x Plan Apo DM Ph3 (NA 1.4) objective, and an Andor Sona CSC-00324 camera at a 20 msec frame rate with the NIS-Elements AR 5.42.05 software. Fiji software was then used to generate subcellular GCaMP fluorescence kymographs by drawing a 475.7 µm² square ROI along the myofiber that was then was maximum projected, generating GCaMP fluorescence kymographs. The number of cells that presented GCaMP transients synchronous with the electrical pulse was then quantified from these kymographs, and plotted.

For single-myofiber contraction tracing, electrical field stimulation was performed as mentioned above. Individual cell contractions were recorded with a MyoCam-S3™ camera (Ionoptix) and a 40x CFI Plan Apochromat Lambda D objective (NA 0.95), using

the IonWizard 7.7.3.180 software. Contraction kinetics were based on CytoMotion (Ionoptix) Pixel Correlation changes relative to the reference frame taken at rest (i.e. before electrical pulses). To generate single-myofiber contraction tracings, pixel correlation (A.U.) curves were plotted using the Cytosolver Cloud (Ionoptix). Myofibers that during the electrical stimuli did not contract, contract, present synchronous or asynchronous contraction during the electrical stimulus were then quantified and plotted.

Transmission Electron Microscopy (TEM) of *in vitro* myofibers

Developed myofibers (7 days of differentiation) were fixed for 1 h at 4 °C in 0.1 M Sorensen's Phosphate Buffer pH 7.4 containing 2.5% (v/v) glutaraldehyde (Science Services EMS), 2% (v/v) formaldehyde (Science Services EMS) and 50 mM of calcium chloride (Sigma). Samples were postfixed for 1 h at 4°C in the dark with 0.1 M Sorensen's Phosphate Buffer pH 7.4 containing 1% (w/v) osmium tetroxide (Science Services EMS) and 0.8% (w/v) potassium ferrocyanide (Sigma-Aldrich). After five washes with distilled water, samples were stained for 1h at room temperature with 0.5% (w/v) uranyl acetate (Analar). Dehydration was performed using an ethanol gradient (30% 10 minutes room temperature, 50% 10 minutes room temperature, 75% overnight 4°C, 90% 10 minutes room temperature, and 100% three times 10 minutes each room temperature). Samples were embedded in Embed-812 epoxy resin (Science Services EMS) and hardened at 60 °C for at least 24 hours. Ultrathin sections (70 nm) were obtained and collected in homemade formvar-carbon coated copper-palladium slot grids and counter-stained with 1% (w/v) uranyl acetate and Reynolds lead citrate, for 5 minutes each. Regions of interest were mapped with SerialEM software and tiles of differentiated cells were acquired using a FEI Tecnai G2 Spirit BioTWIN (120 kV) transmission electron microscope with an Olympus-SIS Veleta CCD Camera.

Stitching was performed using the IMOD blendemont function (Kremer et al., 1996). Stitched tiles were opened in Fiji and we counted the number of triads (one t-tubule connected to two ER) in each tile, and normalized it to the cell area. To measure t-tubule area, we identified the membrane objects that had a low electron-dense lumen, and that were contacting the ER (green or blue arrowheads in Figure S4A). We then measured its area by drawing a region of interest along the t-tubule membrane, using the drawing tool from Fiji software.

Data representation and statistics

All graphs were generated and statistical analyses were conducted using GraphPad
Prism. Normality testing was carried out using the Shapiro-Wilk test, with an alpha
level=0.05. For normal distributions, t-student tests were performed. For datasets that
did not pass the normality test, statistical comparisons were performed using the Mann-
Whitney test. For paired samples that did not pass the normality test, the Wilcoxon test
was used. Statistical significance was defined as * $p < 0.05$.

Acknowledgments

We thank all the members of the Gomes, Way and Moores laboratories for their continued support, scientific discussions and helpful comments on this work. We thank A.L. Sousa from the Electron Microscopy Facility at the Instituto Gulbenkian de Ciência for her technical expertise, sample processing, and imaging of differentiated myofibers by transmission electron microscopy (TEM). We also thank the Bioimaging Unit (funded by PPBI-POCI-01-0145-FEDER-022122) and the Rodent Facility of Instituto de Medicina Molecular João Lobo Antunes for their technical support. We thank N. Lukoyanova and S. Chen for electron microscope support and D. Houldershaw for computational support at Birkbeck.

Funding information

This work was supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 810207). Michael Way is additionally supported by the Francis Crick Institute, which receives its core funding from Cancer Research UK (CC2096), the UK Medical Research Council (CC2096), and the Wellcome Trust (CC2096). Ana Raquel Pereira was supported by the Fundação para a Ciência e Tecnologia (https://doi.org/10.54499/2022.07289.CEECIND/CP1727/CT0003). We acknowledge Diamond Light Source for access and support of the cryo-EM facilities at the UK's national Electron Bio-imaging Centre (eBIC) funder proposal EM20287-56, funded by the Wellcome Trust, MRC and BBSRC. Sample optimization and cryo-CLEM experiments were conducted in the ISMB EM facility (Birkbeck College, University of London) with financial support from the Wellcome Trust (202679/Z/16/Z, 206166/Z/17/Z) and UKRI-Medical Research Council (MR/X013359/1). For the purpose of Open Access, the authors have applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission.

Author contributions

A.R.P. and A.R.S. carried out experiments and analysed data. T.L., J.L.F. and C.A.M designed the cryo-CLEM experiments, T.L and J.L.F. performed collected the cryo-ET data, T.L. performed the 3D reconstruction and segmentation the experiments and, T.L and C.A.M analysed the data. G.L. performed the t-tubule cluster imaging. I.F. generated GFP-Bin1 lentivirus expressing plasmid. N.K. and M.W. provided unpublished data and tools. A.R.P, A.R.S, and E.R.G designed experiments and wrote the manuscript with assistance from other authors. All authors participated in the critical review and revision of the manuscript. E.R.G, M.W and C.A.M supervised the project and secured funding.

Declaration of Interest

The authors declare no competing interests.

References

Abella, J.V.G., Galloni, C., Pernier, J., Barry, D.J., Kjær, S., Carlier, M.-F., Way, M., 2016. Isoform diversity in the Arp2/3 complex determines actin filament dynamics. Nat. Cell Biol. 18, 76–86. https://doi.org/10.1038/ncb3286

Allard, A., Lopes dos Santos, R., Campillo, C., 2021. Remodelling of membrane tubules by the actin cytoskeleton. Biol. Cell 113, 329–343. https://doi.org/10.1111/boc.202000148

Al-Qusairi, L., Laporte, J., 2011. T-tubule biogenesis and triad formation in skeletal muscle and implication in human diseases. Skelet. Muscle 1, 26. https://doi.org/10.1186/2044-5040-1-26

Boncompagni, S., d'Amelio, L., Fulle, S., Fanò, G., Protasi, F., 2006. Progressive Disorganization of the Excitation–Contraction Coupling Apparatus in Aging Human Skeletal Muscle as Revealed by Electron Microscopy: A Possible Role in the Decline of Muscle Performance. J. Gerontol. Ser. A 61, 995–1008. https://doi.org/10.1093/gerona/61.10.995

Bovellan, M., Romeo, Y., Biro, M., Boden, A., Chugh, P., Yonis, A., Vaghela, M., Fritzsche, M., Moulding, D., Thorogate, R., Jégou, A., Thrasher, A.J., Romet-Lemonne, G., Roux, P.P., Paluch, E.K., Charras, G., 2014. Cellular control of cortical actin nucleation. Curr. Biol. CB 24, 1628–1635. https://doi.org/10.1016/j.cub.2014.05.069

Buchholz, T.-O., Krull, A., Shahidi, R., Pigino, G., Jékely, G., Jug, F., 2019. Content-aware image restoration for electron microscopy. Methods Cell Biol. 152, 277–289. https://doi.org/10.1016/bs.mcb.2019.05.001

Campbell-Valois, F.-X., Trost, M., Chemali, M., Dill, B.D., Laplante, A., Duclos, S., Sadeghi, S., Rondeau, C., Morrow, I.C., Bell, C., Gagnon, E., Hatsuzawa, K., Thibault, P., Desjardins, M., 2012. Quantitative proteomics reveals that only a subset of the endoplasmic reticulum contributes to the phagosome. Mol. Cell. Proteomics MCP 11, M111.016378. https://doi.org/10.1074/mcp.M111.016378

Campellone, K.G., Lebek, N.M., King, V.L., 2023. Branching out in different directions: Emerging cellular functions for the Arp2/3 complex and WASP-family actin nucleation factors. Eur. J. Cell Biol. 102, 151301. https://doi.org/10.1016/j.ejcb.2023.151301

Cao, L., Huang, S., Basant, A., Mladenov, M., Way, M., 2024. CK-666 and CK-869 differentially inhibit Arp2/3 iso-complexes. EMBO Rep. 1–19. https://doi.org/10.1038/s44319-024-00201-x

Chen, T.-W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A., Schreiter, E.R., Kerr, R.A., Orger, M.B., Jayaraman, V., Looger, L.L., Svoboda, K., Kim, D.S., 2013. Ultrasensitive fluorescent proteins for imaging neuronal activity. Nature 499, 295–300. https://doi.org/10.1038/nature12354

Chertkova, A.O., Mastop, M., Postma, M., Van Bommel, N., Van Der Niet, S., Batenburg, K.L., Joosen, L., Gadella, T.W.J., Okada, Y., Goedhart, J., 2017. Robust and Bright Genetically Encoded Fluorescent Markers for Highlighting Structures and Compartments in Mammalian Cells. https://doi.org/10.1101/160374

Chugh, P., Paluch, E.K., 2018. The actin cortex at a glance. J. Cell Sci. 131, jcs186254. https://doi.org/10.1242/jcs.186254

Collins, A., Warrington, A., Taylor, K.A., Svitkina, T., 2011. Structural Organization of the Actin Cytoskeleton at Sites of Clathrin-Mediated Endocytosis. Curr. Biol. 21, 1167–1175. https://doi.org/10.1016/j.cub.2011.05.048

Di Ciano, C., Nie, Z., Szászi, K., Lewis, A., Uruno, T., Zhan, X., Rotstein, O.D., Mak, A., Kapus, A., 2002. Osmotic stress-induced remodeling of the cortical cytoskeleton. Am. J. Physiol. Cell Physiol. 283, C850-865. https://doi.org/10.1152/ajpcell.00018.2002

Dowling, J.J., Lawlor, M.W., Dirksen, R.T., 2014. Triadopathies: An Emerging Class of Skeletal Muscle Diseases. Neurotherapeutics 11, 773–785. https://doi.org/10.1007/s13311-014-0300-3

Ershov, D., Phan, M.-S., Pylvänäinen, J.W., Rigaud, S.U., Le Blanc, L., Charles-Orszag, A., Conway, J.R.W., Laine, R.F., Roy, N.H., Bonazzi, D., Duménil, G., Jacquemet, G., Tinevez, J.-Y., 2022. TrackMate 7: integrating state-of-the-art segmentation algorithms into tracking pipelines. Nat. Methods 19, 829–832. https://doi.org/10.1038/s41592-022-01507-1

Falcone, S., Roman, W., Hnia, K., Gache, V., Didier, N., Lainé, J., Auradé, F., Marty, I., Nishino, I., Charlet-Berguerand, N., Romero, N.B., Marazzi, G., Sassoon, D., Laporte, J., Gomes, E.R., 2014. N-WASP is required for Amphiphysin-2/BIN1-dependent nuclear positioning and triad organization in skeletal muscle and is involved in the pathophysiology of centronuclear myopathy. EMBO Mol. Med. 6, 1455–1475. https://doi.org/10.15252/emmm.201404436

Fäßler, F., Javoor, M.G., Datler, J., Döring, H., Hofer, F.W., Dimchev, G., Hodirnau, V.-V., Faix, J., Rottner, K., Schur, F.K.M., 2023. ArpC5 isoforms regulate Arp2/3 complex-dependent protrusion through differential Ena/VASP positioning. Sci. Adv. 9, eadd6495. https://doi.org/10.1126/sciadv.add6495

Flucher, B.E., Takekura, H., Franzini-Armstrong, C., 1993. Development of the excitation-contraction coupling apparatus in skeletal muscle: association of sarcoplasmic reticulum and transverse tubules with myofibrils. Dev. Biol. 160, 135–147. https://doi.org/10.1006/dbio.1993.1292

Fourest-Lieuvin, A., Rendu, J., Osseni, A., Pernet-Gallay, K., Rossi, D., Oddoux, S., Brocard, J., Sorrentino, V., Marty, I., Fauré, J., 2012. Role of triadin in the organization of reticulum membrane at the muscle triad. J. Cell Sci. 125, 3443–3453. https://doi.org/10.1242/jcs.100958

Franzini-Armstrong, C., 2018. The relationship between form and function throughout the history of excitation–contraction coupling. J. Gen. Physiol. 150, 189–210. https://doi.org/10.1085/jgp.201711889

Galloni, C., Carra, D., Abella, J.V.G., Kjær, S., Singaravelu, P., Barry, D.J., Kogata, N., Guérin, C., Blanchoin, L., Way, M., 2021. MICAL2 enhances branched actin network disassembly by oxidizing Arp3B-containing Arp2/3 complexes. J. Cell Biol. 220, e202102043. https://doi.org/10.1083/jcb.202102043

Hall, T.E., Martel, N., Ariotti, N., Xiong, Z., Lo, H.P., Ferguson, C., Rae, J., Lim, Y.-W., Parton, R.G., 2020. In vivo cell biological screening identifies an endocytic capture mechanism for T-tubule formation. Nat. Commun. 11, 3711. https://doi.org/10.1038/s41467-020-17486-w

Janota, C.S., Pinto, A., Pezzarossa, A., Machado, P., Costa, J., Campinho, P., Franco, C.A., Gomes, E.R., 2022. Shielding of actin by the endoplasmic reticulum impacts nuclear positioning. Nat. Commun. 13, 2763. https://doi.org/10.1038/s41467-022-30388-3

Kawaguchi, K., Fujita, N., 2024. Shaping transverse-tubules: central mechanisms that play a role in the cytosol zoning for muscle contraction. J. Biochem. (Tokyo) 175, 125–131. https://doi.org/10.1093/jb/mvad083

Kosmalska, A.J., Casares, L., Elosegui-Artola, A., Thottacherry, J.J., Moreno-Vicente, R., González-Tarragó, V., del Pozo, M.Á., Mayor, S., Arroyo, M., Navajas, D., Trepat, X., Gauthier, N.C., Roca-Cusachs, P., 2015. Physical principles of membrane remodelling during cell mechanoadaptation. Nat. Commun. 6, 7292. https://doi.org/10.1038/ncomms8292

Kremer, J.R., Mastronarde, D.N., McIntosh, J.R., 1996. Computer Visualization of Three-Dimensional Image Data Using IMOD. J. Struct. Biol. 116, 71–76. https://doi.org/10.1006/jsbi.1996.0013

Laplaud, V., Levernier, N., Pineau, J., Roman, M.S., Barbier, L., Sáez, P.J., Lennon-Duménil, A.-M., Vargas, P., Kruse, K., du Roure, O., Piel, M., Heuvingh, J., 2021. Pinching the cortex of live cells reveals thickness instabilities caused by

myosin II motors. Sci. Adv. 7, eabe3640.
https://doi.org/10.1126/sciadv.abe3640

Lee, C., Chen, L.B., 1988. Dynamic behavior of endoplasmic reticulum in living cells.
Cell 54, 37–46. https://doi.org/10.1016/0092-8674(88)90177-8

Lee, E., Marcucci, M., Daniell, L., Pypaert, M., Weisz, O.A., Ochoa, G.-C., Farsad, K.,
Wenk, M.R., De Camilli, P., 2002. Amphiphysin 2 (Bin1) and T-tubule
biogenesis in muscle. Science 297, 1193–1196.
https://doi.org/10.1126/science.1071362

Lemerle, E., Lainé, J., Benoist, M., Moulay, G., Bigot, A., Labasse, C., Madelaine, A.,
Canette, A., Aubin, P., Vallat, J.-M., Romero, N.B., Bitoun, M., Mouly, V., Marty,
I., Cadot, B., Picas, L., Vassilopoulos, S., 2023. Caveolae and Bin1 form ring-
shaped platforms for T-tubule initiation. eLife 12, e84139.
https://doi.org/10.7554/eLife.84139

Li, Y., Lipowsky, R., Dimova, R., 2011. Membrane nanotubes induced by aqueous
phase separation and stabilized by spontaneous curvature. Proc. Natl. Acad.
Sci. 108, 4731–4736. https://doi.org/10.1073/pnas.1015892108

Lionello, V.M., Kretz, C., Edelweiss, E., Crucifix, C., Gómez-Oca, R., Messaddeq, N.,
Buono, S., Koebel, P., Massana Muñoz, X., Diedhiou, N., Cowling, B.S., Bitoun,
M., Laporte, J., 2022. BIN1 modulation in vivo rescues dynamin-related
myopathy. Proc. Natl. Acad. Sci. 119, e2109576119.
https://doi.org/10.1073/pnas.2109576119

Liu, Y.-T., Zhang, H., Wang, H., Tao, C.-L., Bi, G.-Q., Zhou, Z.H., 2022. Isotropic
reconstruction for electron tomography with deep learning. Nat. Commun. 13,
6482. https://doi.org/10.1038/s41467-022-33957-8

Lo, H.P., Lim, Y.-W., Xiong, Z., Martel, N., Ferguson, C., Ariotti, N., Giacomotto, J.,
Rae, J., Floetenmeyer, M., Moradi, S.V., Gao, Y., Tillu, V.A., Xia, D., Wang, H.,
Rahnama, S., Nixon, S.J., Bastiani, M., Day, R.D., Smith, K.A., Palpant, N.J.,
Johnston, W.A., Alexandrov, K., Collins, B.M., Hall, T.E., Parton, R.G., 2021.
Cavin4 interacts with Bin1 to promote T-tubule formation and stability in
developing skeletal muscle. J. Cell Biol. 220, e201905065.
https://doi.org/10.1083/jcb.201905065

Nixon-Abell, J., Obara, C.J., Weigel, A.V., Li, D., Legant, W.R., Xu, C.S., Pasolli, H.A.,
Harvey, K., Hess, H.F., Betzig, E., Blackstone, C., Lippincott-Schwartz, J.,
2016. Increased spatiotemporal resolution reveals highly dynamic dense
tubular matrices in the peripheral ER. Science 354, aaf3928.
https://doi.org/10.1126/science.aaf3928

Peachey, L.D., 1965. THE SARCOPLASMIC RETICULUM AND TRANSVERSE
TUBULES OF THE FROG'S SARTORIUS. J. Cell Biol. 25, 209–231.

Pettersen, E.F., Goddard, T.D., Huang, C.C., Meng, E.C., Couch, G.S., Croll, T.I.,
Morris, J.H., Ferrin, T.E., 2021. UCSF ChimeraX: Structure visualization for
researchers, educators, and developers. Protein Sci. Publ. Protein Soc. 30, 70–
82. https://doi.org/10.1002/pro.3943

Picas, L., Viaud, J., Schauer, K., Vanni, S., Hnia, K., Fraisier, V., Roux, A., Bassereau,
P., Gaits-Iacovoni, F., Payrastre, B., Laporte, J., Manneville, J.-B., Goud, B.,
2014. BIN1/M-Amphiphysin2 induces clustering of phosphoinositides to recruit
its downstream partner dynamin. Nat. Commun. 5, 5647.
https://doi.org/10.1038/ncomms6647

Pimentel, M.R., Falcone, S., Cadot, B., Gomes, E.R., 2017. In Vitro Differentiation of
Mature Myofibers for Live Imaging. J. Vis. Exp. JoVE 55141.
https://doi.org/10.3791/55141

Pizarro-Cerdá, J., Chorev, D.S., Geiger, B., Cossart, P., 2017. The Diverse Family of
Arp2/3 Complexes. Trends Cell Biol. 27, 93–100.
https://doi.org/10.1016/j.tcb.2016.08.001

Porter, K.R., Palade, G.E., 1957. Studies on the endoplasmic reticulum. III. Its form
and distribution in striated muscle cells. J. Biophys. Biochem. Cytol. 3, 269–

300. https://doi.org/10.1083/jcb.3.2.269

Pylvänäinen, J.W., Laine, R.F., Saraiva, B.M.S., Ghimire, S., Follain, G., Henriques, R.,
Jacquemet, G., 2023. Fast4DReg – fast registration of 4D microscopy datasets.
J. Cell Sci. 136, jcs260728. https://doi.org/10.1242/jcs.260728

Rohou, A., Grigorieff, N., 2015. CTFFIND4: Fast and accurate defocus estimation from
electron micrographs. J. Struct. Biol. 192, 216–221.
https://doi.org/10.1016/j.jsb.2015.08.008

Roman, W., Martins, J.P., Carvalho, F.A., Voituriez, R., Abella, J.V.G., Santos, N.C.,
Cadot, B., Way, M., Gomes, E.R., 2017. Myofibril contraction and cross-linking
drive nuclear movement to the periphery of skeletal muscle. Nat. Cell Biol. 19,
1189–1201. https://doi.org/10.1038/ncb3605

Sadhu, L., Tsopoulidis, N., Hasanuzzaman, M., Laketa, V., Way, M., Fackler, O.T.,
2023. ARPC5 isoforms and their regulation by calcium-calmodulin-N-WASP
drive distinct Arp2/3-dependent actin remodeling events in CD4 T cells. eLife
12, e82450. https://doi.org/10.7554/eLife.82450

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J.,
Hartenstein, V., Eliceiri, K., Tomancak, P., Cardona, A., 2012. Fiji: an open-
source platform for biological-image analysis. Nat. Methods 9, 676–682.
https://doi.org/10.1038/nmeth.2019

Sébastien, M., Aubin, P., Brocard, Jacques, Brocard, Julie, Marty, I., Fauré, J., 2020.
Dynamics of triadin, a muscle-specific triad protein, within sarcoplasmic
reticulum subdomains. Mol. Biol. Cell 31, 261–272.
https://doi.org/10.1091/mbc.E19-07-0399

Serwas, D., Akamatsu, M., Moayed, A., Vegesna, K., Vasan, R., Hill, J.M.,
Schöneberg, J., Davies, K.M., Rangamani, P., Drubin, D.G., 2022. Mechanistic
insights into actin force generation during vesicle formation from cryo-electron
tomography. Dev. Cell 57, 1132-1145.e5.
https://doi.org/10.1016/j.devcel.2022.04.012

Svitkina, T.M., 2020. Actin Cell Cortex: Structure and Molecular Organization. Trends
Cell Biol. 30, 556–565. https://doi.org/10.1016/j.tcb.2020.03.005

Takekura, H., Flucher, B.E., Franzini-Armstrong, C., 2001. Sequential docking,
molecular differentiation, and positioning of T-Tubule/SR junctions in
developing mouse skeletal muscle. Dev. Biol. 239, 204–214.
https://doi.org/10.1006/dbio.2001.0437

Várkuti, B.H., Képiró, M., Horváth, I.Á., Végner, L., Ráti, S., Zsigmond, Á., Hegyi, G.,
Lenkei, Z., Varga, M., Málnási-Csizmadia, A., 2016. A highly soluble, non-
phototoxic, non-fluorescent blebbistatin derivative. Sci. Rep. 6, 26141.
https://doi.org/10.1038/srep26141

Ward, S.R., Eng, C.M., Smallwood, L.H., Lieber, R.L., 2009. Are Current
Measurements of Lower Extremity Muscle Architecture Accurate? Clin. Orthop.
467, 1074–1082. https://doi.org/10.1007/s11999-008-0594-8

Zheng, S.Q., Palovcak, E., Armache, J.-P., Verba, K.A., Cheng, Y., Agard, D.A., 2017.
MotionCor2: anisotropic correction of beam-induced motion for improved cryo-
electron microscopy. Nat. Methods 14, 331–332.
https://doi.org/10.1038/nmeth.4193

Zivanov, J., Otón, J., Ke, Z., von Kügelgen, A., Pyle, E., Qu, K., Morado, D., Castaño-
Díez, D., Zanetti, G., Bharat, T.A.M., Briggs, J.A.G., Scheres, S.H.W., 2022. A
Bayesian approach to single-particle electron cryo-tomography in RELION-4.0.
eLife 11, e83724. https://doi.org/10.7554/eLife.83724

Main Figures

Figure 1

Figure 1 – T-tubules and ER are dynamic outside of the triad. (**A**) Top image corresponds to
a timelapse snapshot of a matured isolated myofiber, stained with CellMask to label t-tubules and
imaged by timelapse every 3 seconds for 2 minutes. Bottom image corresponds to kymographs
from regions of interest (ROI) 1 and 2 of representative dynamic t-tubules. The plot represents
the number of growing t-tubules per cell area during 2 minutes (N=9 isolated fibers, from three
independent experiments). (**B**) Top image corresponds to a timelapse snapshot of an in vitro
developing myofiber stained with CellMask to label t-tubules and imaged by timelapse every 3
seconds for 2 minutes. At this stage of development (differentiation day 4), t-tubules are starting
to form and have a longitudinal orientation. Bottom image corresponds to kymographs from ROIs
1 and 2 of representative dynamic t-tubules. The plots represent the number of growing t-tubules
per cell area during 2 min (N=72 cells, from three independent experiments) and t-tubule growth
mean speed (N=58 growing t-tubules, from three independent experiments). (**C**) Left image
corresponds to a timelapse snapshot of an in vitro developing myofiber (differentiation day 4)
expressing GFP-KDEL (ER, green), stained with CellMask to label t-tubules (magenta) and
imaged by timelapse every 3 seconds for 2 minutes. Right image corresponds to kymograph
from red square ROI showing representative T-tubule and ER dynamics. (**D**) Left image
corresponds to a timelapse snapshot of an in vitro developing myofiber (differentiation day 4)
expressing Triadin-GFP (green), stained with CellMask (magenta) to label t-tubules and imaged
by timelapse every 3 seconds for 2 minutes. Right image corresponds to kymograph from red
ROI showing that Triadin-GFP is static when t-tubules are growing (white arrow). The plot
represents Triadin-GFP mean speed in five different cells from two independent repeats (N=783).
A-D scale bars 5 µm; C kymograph scale bar 0.5 µm. Blue lines on plots represent mean values.

Figure 2

Figure 2 – Inhibiting Arp2/3 with CK666 or through Arpc5 depletion promotes t-tubule
growth. (A) Timelapse snapshots of GFP-Bin1 before or after CK666 addition. The plot
corresponds to the number of growing t-tubules per cell during 2 min before or after CK666
addition. Statistics: Wilcoxon test; N=22; results are from three independent experiments. **(B)**
Timelapse snapshots of in vitro developing myofibers (differentiation day 4) with CellMask labeled
t-tubules before or after CK666 addition. Red X indicates the site of t-tubule growth during 2 min.
The plot corresponds to the number of growing t-tubules per cell area during 2 min in the same
myofibers before or after CK666 addition. Statistics: Wilcoxon test; N=38 myofibers, results are
from three independent experiments. (**C**) Timelapse snapshots of GFP-Bin1 myoblasts
transfected with scramble (Control), Arpc5 (si Arpc5) or Arpc5L (si Arpc5L) siRNA. The plot
corresponds to the number of growing t-tubules per cell area during 2 min. Statistics: Mann-
Whitney; Control N=25; siArpC5 N=27; siArpC5L N=25. Results from three independent
experiments. (**D**) Timelapse snapshots of in vitro developing myofibers (differentiation day 4) with
CellMask-labelled t-tubules from scramble (Control); Arpc5 depleted (siArp5) or Arpc5L depleted
(siArpC5L) myofibers. Red X indicates sites of t-tubule growth during 2 min. The plot corresponds
to the number of growing t-tubules per cell area during 2 min. Statistics: Mann-Whitney; Control
N=29; siArpc5 N=24; siArpC5L N=22; results are from three independent experiments. Blue lines
on plots represent mean values. Scale bars 5 µm.

Figure 3

Figure 3 – F-actin is not locally nucleated at growing t-tubules. (A) Left image corresponds
to a timelapse snapshot of GFP-Bin1 myoblasts (green) and LifeAct-mCherry (magenta). Right
image corresponds to kymographs from the red rectangle ROI showing a t-tubule growing over
time in the absence of F-actin changes. (**B**) Plots of LifeAct-mCherry and GFP-Bin1 average
fluorescence intensity over time. Mean and SEM are represented. N=19 from two independent
experiments. (**C**) Left image corresponds to a timelapse snapshot of GFP-Bin1 myoblasts (green)
and mScarlet-I-UtrCH (magenta). Right image corresponds to kymographs from the red rectangle
ROI showing a t-tubule growing over time in the absence of F-actin changes. (**D**) Plots of the
mScarlet-I-UtrCH and GFP-Bin1 average fluorescence intensity over time. Mean and SEM are
represented. N=16 from two independent experiments. (**E**) Representative low magnification
cryo-EM image showing part of one cell on a TEM grid. The cyan rectangle indicates the ROI
imaged using epifluorescence and brightfield microscopy (panel F). (**F**) Merged epifluorescence
(green) and brightfield (grey, FM) image of the ROI. in E, showing multiple GFP-Bin1 t-tubules at
the cell periphery. The orange rectangle indicates the ROI imaged using TEM (panel G). (**G**)
Medium magnification TEM image of the ROI in F, showing the t-tubules observed in F. (**H**) 3D
representation of cropped tomogram segmentation of the white rectangle ROI from G, with actin
filaments (purple) and t-tubule membrane (green). (**I**) Slices of the tomogram in H at different z
positions (from left to right: z=-25nm, 30nm and 135nm relative to the central slice) with actin
filaments (purple), t-tubule membrane (green), and potential actin branches (yellow arrows).
Scale bar: 5 µm in A, B and E, 1 µm in F, 500 nm in G and 100 nm in H and I.

 Figure 4

Figure 4 – Cortical actin disruption promotes t-tubule growth. (**A**) Timelapse snapshots of GFP-Bin1 myoblasts before or after para-aminoblebbistatin addition (Blebb). The plot corresponds to the number of growing t-tubules per cell area for 2 min before or after para-aminoblebbistatin addition. Statistics: Wilcoxon test; N=22 cells, results are from three independent experiments. (**B**) Timelapse snapshots of in vitro developing myofibers (differentiation day 4) with CellMask labeled t-tubules before or after para-aminoblebbistatin addition (Blebb). Red X indicates the site of t-tubule growth during 2 min. The plot corresponds to the number of growing t-tubules per cell area during 2 min in the same myofibers before or after para-aminoblebbistatin addition. Statistics: Wilcoxon test; N=31 myofibers, results are from three independent experiments. (**C**) Timelapse snapshots of GFP-Bin1 myoblasts before or after glycerol addition. The plot corresponds to the number of growing t-tubules per cell area for 2 min before or after glycerol addition. Statistics: Wilcoxon test; N=20 cells, results are from three independent experiments. (**D**) Timelapse snapshots of in vitro developing myofibers (differentiation day 4) with CellMask labeled t-tubules before or after glycerol addition. Red X indicates the site of t-tubule growth during 2 min. The plot corresponds to the number of growing t-tubules per cell area during 2 min in the same myofibers before or after glycerol addition. Statistics: Wilcoxon test; N=20 myofibers, results are from three independent experiments. Scale bars 5 µm. Blue lines on plots represent mean value.

1002

1003 **Figure 5**

1004

Figure 5 – Arpc5 depletion leads to t-tubule clusters and reduction Junctophilin1 puncta
in differentiated myofibers. (A) Representative images of CellMask labelled t-tubules from
control (scramble), Arpc5 (si Arpc5), or Arpc5L (si Arpc5L) depleted myofibers (differentiation day
7). Magnifications corresponding to the red boxes are shown below each image. **(B)**
Representative images of Bin1 immunostained t-tubules from control (scramble), Arpc5 (si
Arpc5), or Arpc5L (siArpc5L) depleted myofibers (differentiation day 7). Magnifications
corresponding to the red boxes are shown below each image. **(C)** Quantification of the
percentage of myofibers with t-tubule clusters.Control N=42 myofibers; si Arpc5 N= 41 myofibers;
si Arpc5L N=50 myofibers, from two independent experiments. (**D**) Representative images of
control (scramble), Arpc5 (si Arpc5), or Arpc5L (siArpc5L) depleted myofibers (differentiation day
7) immunostained for Junctophilin 1 (JPH1, cyan), Bin1 (Yellow); and stained with phalloidin
(Phall, magenta). (**E**) Quantification of the number of Junctophilin 1 (JPH1) puncta per μm^2 in
scramble (Control); Arpc5 (si Arpc5) or in Arpc5L depleted myofibers (si Arpc5L). Statistics:
Mann-Whitney statistical test, siArpc5 P-value = 0,0005; siArpc5L P-value = 0,3161; Control N=44
myofibers; si Arpc5 N=41 myofibers; si Arpc5L N=51 myofibers. Results are from three
independent experiments. Scale bars 5 µm. Blue lines on plots represent mean values.
1021

 Figure 6

Figure 6 – Arpc5 depletion leads to asynchronous muscle contraction (A) Myofibers
(differentiation day 7-8) expressing GCaMP were stimulated with 10 V at a frequency of 0.5 Hz
and subcellular GCaMP fluorescence kymographs from 8 different myofibers of control
(Scramble), Arpc5 (siArpc5) and Arpc5L (si Arpc5L) are shown. Orange marks indicate electrical
pulses. (**B**) Quantification of the percentage of myofibers from A with GCaMP fluorescence
transients synchronised with the electrical pulse. Control N=40, si Arpc5 N=62, si Arpc5L N=52.
Results are from three independent experiments. (**C**) Representative contraction tracing
recordings from control (Scramble), Arpc5 (si Arpc5), and Arpc5L (si Arpc5L) depleted myofiber.
Orange marks indicate electrical pulses. (**D**) Quantification of the percentage of contracting
myofibers from C. Control N=43, siArpc5 N=36, siArpc5L N=28. Results are from three
independent experiments. (**E**) Quantification of the percentage of myofibers from **C** that contract
in synchrony with the electrical impulse. Control N=41, siArpc5 N=35, siArpc5L N=28. Results
are from three independent experiments.

Supplementary Figures

1039

1040 **Figure S1**

1041

A

B

1042
1043
Figure S1 - Arpc5 and Arpc5L depletion . Related to Figure 2. (**A**) Western blots of GFP-Bin1
myoblasts knocked down for scramble (Scr) Arpc5 or Arpc5L isoforms. Membranes were blotted
using antibodies against Arpc5 or Arpc5L with vinculin as a loading control. (**B**) Western blots of
myofibers (differentiation day 4 or day 7), knocked down for scramble (Scr), Arpc5 or Arpc5L.
Membranes were blotted using antibodies against Arpc5 and Arpc5L isoforms and vinculin, used
as a loading control.
1050

1051 **Figure S2**

1052
1053
Figure S2 - Correlative light and electron microscopy workflow. Related to Figure3.
(**A**) Cells were imaged using TEM to check the ice thickness and locate spread cells for FM
imaging. In the example image, one spread cell is indicated with a yellow arrow, with the
approximate cell periphery outlined in black in the zoomed view. (**B**) Cells imaged using cryo-
fluorescence microscopy (FM) with the bright-field image of the area in the blue box from A
showing the GFP-Bin1-labeled t-tubule structures at the cell periphery. (**C**) Cells were re-imaged
using medium magnification TEM images in tiles and combined for FM-TEM correlation. The
empty holes and black features were used for manual correlation, with empty holes indicated by
white circles and the dark feature marked by a white arrow. (**D**) A zoomed-in view of the medium
mag image collected from the orange region in C. Tubular membrane structures are highlighted
with white curvy lines. The shapes of these tubular membrane structures resemble those seen in
the FM image in B. Regions of interest (ROI) were selected for tilt series data collection, indicated
by white boxes. (**E**) and (**F**) The central slice along the z-axis of the 3D tomogram at positions 1
and 2, respectively. (**G**) The bright-field and FM image showing fluorescently-labelled vesicular
structure. A tomographic slice of the reconstruction inside the white box showing nested sets of
membranes is shown on the right. Scale bar: 100 µm and 5 µm in A, 2 µm in B and C,500 nm in
D, 100 nm in E and F. 2 µm and 100 nm in G.

Figure S3

1073
Figure S3 - Quantitative analysis of cryo-electron tomograms. Related to Figure 3. (**A**) 3D
representation of cropped tomogram segmentation in Supplementary Fig1E, sub-volumes
proximal to and distant from T-tubule membrane used for counting branch number indicated with
red boxes and blue boxes, respectively. (**B**) 3D representation of cropped tomogram

segmentation in Supplementary Fig1F, sub-volumes proximal to and distant from T-tubule membrane used for counting branch number indicated with red boxes and blue boxes, respectively. (**C**) Slices and segmentations of the tomogram in A in red box1 at different z positions (from left to right: z=-35nm, 5nm and 34nm relative to the central slice) with actin filaments in purple, T-tubules in green, and potential actin branches indicated with yellow arrows. Scale bar: 100 nm. (**D**) Slices and segmentations of the tomogram in A in red box2 at different z positions (from left to right: z=-26nm, 3nm and 28nm relative to the central slice) with actin filaments in purple, T-tubules in green.Scale bar: 100 nm. (**E**) Slices and segmentations of the tomogram in A in red box3 at different z positions (from left to right: z=1 nm, 14 nm and 34nm relative to the central slice) with actin filaments in purple, T-tubules in green.Scale bar: 100 nm. (**F**) Slices and segmentations of the tomogram in in A blue box1 at different z positions (from left to right: z=-27 nm, 3 nm and 22 nm relative to the central slice) with actin filaments in purple, T-tubules in green, and potential actin branches indicated with yellow arrows. Scale bar: 100 nm. (**G**) Slices and segmentations of the tomogram in A in blue box2 at different z positions (from left to right: z=-13nm, 10nm and 31nm relative to the central slice) with actin filaments in purple, T-tubules in green, and potential actin branches indicated with yellow arrows. Scale bar: 100 nm. (**H**) Slices and segmentations of the tomogram in B in red box1 at different z positions (from left to right: z=-15nm, 11nm and 34nm relative to the central slice) with actin filaments in purple, T-tubules in green, and potential actin branches indicated with yellow arrows. Scale bar: 100 nm. (**I**) Slices and segmentations of the tomogram in B in red box2 at different z positions (from left to right: z=-22 nm, 7 nm and 30 nm relative to the central slice) with actin filaments in purple, T-tubules in green, and potential actin branches indicated with yellow arrows. Scale bar: 100 nm. (**J**) Slices and segmentations of the tomogram in B in blue box at different z positions (from left to right: z=-14 nm, 16 nm and 39 nm relative to the central slice) with actin filaments in purple, T-tubules in green, and potential actin branches indicated with yellow arrows. Scale bar: 100 nm.

1104 **Figure S4**

1105

Figure S4 – **Transmission Electron Microscopy (TEM) of stained and resin-embedded**
myofibers. Related to Figure 5. (**A**) TEM of scramble (Control) or Arpc5 (si Arpc5) depleted
myofibers (differentiation day 7). Green arrowheads indicate triads (one t-tubule in contact with
two adjacent ER compartments). Blue arrows indicate t-tubules connected to the ER only on one
side, not forming a triad. Yellow arrowheads indicate enlarged t-tubules. Scale bars 500 nm.
Magnifications corresponding to the red ROIs are shown as insets, Scale bars 100 nm. (**B**) Plot
representing the area of each t-tubule observed in cells from A. Control N=162 t-tubules, si Arpc5
N=56 t-tubules. A total of 6 cells were quantified per condition, two biological replicates. Mann-
Whitney test (p<0.0001). (**C**) Quantifications of the number of triads normalized for cell area (10^3
μm^2). t-test (p=0.0286) Blue lines in graphs represent mean values. Black lines in graphs
represent SEM.

Video legends

Video S1 – Timelapse movie of t-tubule dynamics in a myofiber isolated from the EDL muscle of an adult mice. Isolated fiber was stained with CellMask to label t-tubules. Scale bar 5 µm. Related to Figure 1.

Video S2 – Timelapse movie of t-tubule dynamics in a developing in vitro myofiber. Cells were stained with CellMask to label t-tubules. Scale bar 5 µm. Related to Figure 1.

Video S3 – Timelapse movie of Endoplasmic Reticulum (GFP-KDEL, green) and t-tubule (CellMask, magenta) dynamics in a developing myofiber. Scale bar 5 µm. Related to Figure 1.

Video S4 – Timelapse movie of Triadin (green) and t-tubule (magenta) dynamics in a developing in vitro myofiber. Cells expressing Triadin-GFP stained with CellMask to label t-tubules. Scale bar 5 µm. Related to Figure 1.

Video S5 – Timelapse movie of t-tubule dynamics in GFP-Bin1 myoblasts before CK666. Scale bar 5 µm. Related to Figure 2.

Video S6 – Timelapse movie of t-tubule dynamics in GFP-Bin1 myoblasts after CK666. Scale bar 5 µm. Related to Figure 2.

Video S7 – Timelapse movie of t-tubule dynamics in a developing myofiber before adding CK666. Cells were stained with CellMask to label t-tubules. Scale bar 5 µm. Related to Figure 2.

Video S8 – Timelapse movie of t-tubule dynamics in a developing myofiber after a 5 minute incubation with CK666. Scale bar 5 µm. Related to Figure 2.

Video S9 – Timelapse movie of t-tubule dynamics in GFP-Bin1 myoblasts depleted for scramble control. Scale bar 5 µm. Related to Figure 2.

Video S10 – Timelapse movie of t-tubule dynamics in GFP-Bin1 depleted for Arpc5. Scale bar 5 µm. Related to Figure 2.

Video S11 – Timelapse movie of t-tubule dynamics in GFP-Bin1 myoblasts depleted for Arpc5L. Scale bar 5 µm. Related to Figure 2.

Video S12 – Timelapse movie of t-tubule dynamics in a developing myofiber in scramble cells (Control). T-tubules were stained with CellMask. Scale bar 5 µm. Related to Figure 2.

Video S13 – Timelapse movie of t-tubule dynamics in a developing myofiber depleted for Arpc5. Scale bar 5 µm. Related to Figure 2.

Video S14 – Timelapse movie of t-tubule dynamics in a developing myofiber depleted for Arpc5L. Scale bar 5 µm. Related to Figure 2.

Video S15 – F-actin (LifeAct-mCherry, magenta) and t-tubules (Bin1-GFP, green) in GFP-Bin1 myoblasts. Scale bar 5 µm. Related to Figure 3.

Video S16 – F-actin (mScarlet-i-UtrCH, magenta) and t-tubules (Bin1-GFP, green) in GFP-Bin1 myoblasts. Scale bar 5 µm. Related to Figure 3.

Video S17 – Cryo-electron tomogram and segmentation model of actin cytoskeleton around t-tubule site related to Figure 3I. Actin cytoskeleton coloured in light purple. T-tubule membrane coloured in green.

Video S18 – Cryo-electron tomogram and segmentation model of actin cytoskeleton around t-tubule site related to Figure S3A. Actin cytoskeleton coloured in light purple. T-tubule membrane coloured in green.

1179
1180 **Video S19** – Cryo-electron tomogram and segmentation model of actin cytoskeleton around t-
1181 tubule site related to Figure S3B. Actin cytoskeleton coloured in light purple. T-tubule membrane
1182 coloured in green.
1183
1184 **Video S20** – Timelapse movie of t-tubule dynamics in GFP-Bin1 myoblasts before para-
1185 aminoblebbistatin (Blebb). Scale Bar 5 µm. Related to Figure 4.
1186
1187 **Video S21** – Timelapse movie of t-tubule dynamics in GFP-Bin1 myoblasts after para-
1188 aminoblebbistatin (Blebb). Scale Bar 5 µm. Related to Figure 4.
1189
1190 **Video S22** – Timelapse movie of t-tubule dynamics in a developing in vitro myofiber before adding
1191 para-aminoblebbistatin. Cells were stained with CellMask to label t-tubules. Scale bar 5 µm.
1192 Related to Figure 4.
1193
1194 **Video S23** – Timelapse movie of t-tubule dynamics in a developing in vitro myofiber after a 5
1195 minute incubation with para-aminoblebbistatin. Scale bar 5 µm. Related to Figure 4.
1196
1197 **Video S24** - Timelapse movie of t-tubule dynamics in GFP-Bin1 myoblasts before glycerol. Scale
1198 Bar 5 µm. Related to Figure 4.
1199
1200 **Video S25** - Timelapse movie of t-tubule dynamics in GFP-Bin1 myoblasts after glycerol. Scale
1201 Bar 5 µm. Related to Figure 4.
1202
1203 **Video S26** - Timelapse movie of t-tubule dynamics in a developing myofiber before glycerol. Cells
1204 were stained with CellMask to label t-tubules. Scale bar 5 µm. Related to Figure 4.
1205
1206 **Video S27** - Timelapse movie of t-tubule dynamics in a developing in vitro myofiber after glycerol.
1207 Cells were stained with CellMask to label t-tubules. Scale bar 5 µm. Related to Figure 4.
1208
1209
1210