Quantitative and Kinetic Proteomics Reveal ApoE Isoform-dependent Proteostasis

Adaptations in Mouse Brain

Nathan R. Zuniga [1]; Noah E. Earls [1]; Ariel E. A. Denos [1]; Jared M. Elison [1]; Benjamin S. Jones

[1]; Ethan G. Smith [1]; Noah G. Moran [1]; Katie L. Brown [1]; Gerome M. Romero [1]; Chad D. Hyer [1];

*Kimberly B. Wagstaff [1]; Haifa M. Almughamsi [1,2]; Mark K. Transtrum [3]; John C. Price [*1]*

[1] Department of Chemistry and Biochemistry, College of Computational, Physical, and

Mathematical Sciences, Brigham Young University, Provo, UT, USA

[2] Department of Chemistry, College of Science, Taif University, Taif, Saudi Arabia

[3] Department of Physics and Astronomy, College of Computational, Physical, and Mathematical

Sciences, Brigham Young University, Provo, UT, USA

*Correspondence to drjohncprice@gmail.com

KEYWORDS:

Amyloidβ; ApoE; Neurodegeneration; Alzheimer's Disease; Proteomics; Endosome; Proteasome;

Isotope Labeling; Mitochondria;

ABSTRACT:

Apolipoprotein E (ApoE) polymorphisms modify the risk of neurodegenerative disease with the

ApoE4 isoform increasing and ApoE2 isoform decreasing risk relative to the 'wild-type control'

ApoE3 isoform. To elucidate how ApoE isoforms alter the proteome, we measured relative protein

abundance and turnover in transgenic mice expressing a human ApoE gene (isoform 2, 3, or 4).

This data provides insight into how ApoE isoforms affect the *in vivo* synthesis and degradation of

a wide variety of proteins. We identified 4849 proteins and tested for ApoE isoform-dependent

changes in the homeostatic regulation of ~2700 ontologies. In the brain, we found that ApoE4 and

ApoE2 both lead to modified regulation of mitochondrial membrane proteins relative to the wild-

type control ApoE3. In ApoE4 mice, this regulation is not cohesive suggesting that aerobic

respiration is impacted by proteasomal and autophagic dysregulation. ApoE2 mice exhibited a

matching change in mitochondrial matrix proteins and the membrane which suggests coordinated

maintenance of the entire organelle. In the liver, we did not observe these changes suggesting that

the ApoE-effect on proteostasis is amplified in the brain relative to other tissues. Our findings

underscore the utility of combining protein abundance and turnover rates to decipher proteome

regulatory mechanisms and their potential role in biology.

INTRODUCTION:

Apolipoprotein E (ApoE) is one of the lipoproteins used for the transport of lipids and cholesterol

throughout the body. ApoE is also the primary transporter of lipids in the brain. The three major

subtypes of human ApoE—ApoE2, ApoE3, and ApoE4— differ by 2 amino acids and exhibit

allelic frequencies of 8.4%, 77.9%, and 13.7%, respectively. [2, 3] The ApoE3 allele is considered

the normal or wild-type, and the behavior of the E2 or E4 isoforms differs from E3 in measurable

ways. The ApoE2 protein isoform, characterized by an R158C substitution relative to the ApoE3,

has been associated with decreased affinity for the LDL receptor[4, 5], while the ApoE4 protein

isoform, which features a C112R substitution relative to ApoE3, favors binding to very-low-

density lipoprotein receptors[4, 5]. Thus, these seemingly minor genotypic changes may lead to

profound biochemical consequences.

Both ApoE2 and E4 modulate disease risk relative to ApoE3. Ferrer et al. observed a 3 – 15-fold

increase in Alzheimer's Disease (AD) prevalence in carriers of the ApoE4 allele relative to ApoE3

carriers and a decreased risk in individuals expressing the ApoE2 allele.[6] Although ApoE2

expression protects against AD, its expression is associated with the increased incidence of familial

type III hyperlipoproteinemia—a disorder characterized by an inability to metabolize lipids

including cholesterol and triglycerides.[7] ApoE isoforms have also been implicated in the

development of Parkinson's disease[8], vascular pathology[9], and most recently, COVID-19

prognosis[10].

Some mechanistic details have been identified for how the ApoE alleles modulate an individual's

risk for disease. ApoE is a transporter of amyloidβ, a widely recognized biomarker in AD

development.[11] ApoE-isoforms modulate brain mRNA expression, presumably in response to

changes in lipid availability[11] as well as direct transcriptional effects.[12] Here we used both

quantitative and kinetic proteomics to explore the impact of human ApoE genotypes in the

proteome of mice. Both approaches leverage liquid chromatography and mass spectrometry (LC-

MS) to identify and quantify thousands of proteins (Figure 1A).[1] We apply a simplified kinetic

model of proteostasis (Figure 1B), which combines turnover rate and concentration measurements

to reveal ApoE isoform-dependent effects on protein synthesis and degradation. Our analysis

identifies key brain-specific proteostasis changes, as evidenced by pathway-level changes in

synthesis and degradation. Building upon a significant body of literature and this proteome scale

study, we propose a unifying mechanism wherein ApoE alleles systemically impact cellular

proteostasis through alterations in endosomal trafficking, mitochondrial function, and proteo-

lysosomal activity.

Experimental Design and Statistical Rationale

Cohort Grouping and Analysis Rationale

A total of 72 homozygous ApoE transgenic mice, with an equal distribution of female and male

individuals were included. This cohort included 24 ApoE2, 24 ApoE3, and 24 ApoE4 (refer to

Table S1 for details). The sample groups for protein turnover rate measurements of each ApoE

genotype and gender, were two independent blocks of six mice. These six mice were selected

based on the metabolic labeling duration, namely Day 0, Hour 6, Day 1, Day 4, Day 16, and Day

32 post-exposure to deuterium.

The kinetic analysis utilized peptide identifications from LC-MS/MS acquisition files to extract

isotope envelope information from LC-MS (MS1 only) data. Notably, this process heavily relies

on peptide retention time. To facilitate this, MS/MS data and MS data were collected within the

same sample worklist. The initial four timepoints (Day 0, Hour 6, Day 1, and Day 4) were used to

generate LC-MS/MS fragmentation spectra and identify peptide sequences with observed charge

and retention time.

To streamline sample processing and turnover rate measurements, mice were organized into four

gender-specific groups of 18 mice (n=6 per genotype, Figure S1). This grouping strategy

accommodated instrument availability and minimized retention time deviations associated with

extensive sample worklists. Additionally, from each group, a subset of four mice per genotype,

comprising the first four timepoints (Day 0, Hour 6, Day 1, Day 4), were selected for LFQ

proteomics. This selection yielded a total of 16 mice per ApoE genotype for an area under the LC

curve (Abundance) fold change (FC) calculations, with an equal distribution of 8 females and 8

males in the four gender-specific groups. Thus, out of the initial 72 mice, 48 were used to generate

the "Abundance" FC values (Figure S1).

To broaden proteome coverage, each brain homogenate sample was fractionated into cytosolic

and membrane components, which were prepared and analyzed separately using the workflow

described below. This fractionation led to the creation of eight datasets for our analysis. Each

dataset underwent individual processing using the Peaks Studio software (Bioinformics Solutions

Inc.) for protein abundance and Deuterater software [1] for turnover rate measurements. Protein-

level abundance fold change relative to control (FC values), turnover rate FC values, and statistical

analysis (P-value) for each comparison (e.g., E2vsE3) were calculated for each dataset

independently to minimize inter-set variance caused by sample prep discrepancies, instrument

noise, buffer compositions, and sample run variables. It is worth noting that due to problems in

sample processing, the Hour 6 sample was omitted from a single ApoE4 dataset (D16 was

substituted for LFQ analysis), and Day 4 was omitted from a single ApoE3 LFQ dataset (See Table

S1, Figure S1).

While FC and P-value calculations were conducted at the protein level, this study mainly focuses

on how proteins with shared functional characteristics are regulated in an ApoE isoform-specific

manner. To achieve this, the StringDB multiprotein tool[13] was employed to identify functional

groups (ontologies) represented in the final data sets (Abundance FC, Turnover FC). Every protein

Abundance FC value was calculated with a minimum of three biological abundance measurements

in experimental (ApoE2, or ApoE4) and control (ApoE3); see the 'Protein ΔAbundance Analysis'

section for more details. The null hypothesis (H0) posited that proteins' collective gene expression

ratio in an ontology would remain unchanged (H0: Abundance FC = 1) across ApoE genotypes.

Consequently, we tested the alternative hypothesis that ApoE genotype alters the regulation of

functionally related protein groups (Ha: Abundance FC $\neq$ 1) using a one-sample t-test. This

analytical approach captured changes occurring across the broader functional proteome rather than

focusing solely on identifying individually significant proteins. Python code created for both

protein- and ontology-level calculations is available in the GitHub repository, as detailed in the

Supplementary Data section of this paper.

Proteostasis Model and Analysis Rational

Figure S2: Protein Homeostasis Model with commonly observed sources and sinks of protein

A protein homeostasis model must account for common sources and sinks of protein mass

(Figure S2). In this model we assume there is a large circulating pool of free amino acids affected

by diet, metabolites, and waste expulsion. Amino acids become the precursors for protein synthesis

in an initial tRNA charging step, which then polymerize in an mRNA-dependent step before

folding into functional proteins. Multiple competing processes subsequently influence the

resulting protein concentration. First, degradation returns the protein to the constituent amino acids

in the free pool. The functional proteins may also transition (reversibly) into an

aggregate/condensate state that undergoes a separate degradation process. Finally, protein

concentration may be affected by importing or exporting proteins.

Our goal is to use chemical kinetics and translate this diagram into a mathematical model with

a few tunable parameters identifiable from experimental data. Unfortunately, a complete

mathematical translation of this system leads to a model with too many parameters to draw

meaningful conclusions. We, therefore, make several simplifying assumptions regarding which

processes are dominant to restrict the parameters to an identifiable subset.

First, the mice in this study are healthy adults, so we assume the protein concentrations are in

steady-state with no protein aggregate. We assume that the pool of free amino acids is large so

that the rate bottleneck in the tRNA-charging/synthesis steps is synthesis and that import/export is

negligible. We also assume the protein pool is well mixed, assuring the random selection of protein

for degradation (unregulated). Although a reasonable starting point for the present study, we

acknowledge that these assumptions are poor for surprisingly large sections of the proteome where

reversible aggregation [14, 15], multistage regulation of synthesis rates [16-18], exchange of protein

subunits in complexes [19, 20] and nonrandom degradation [21-24] are biologically important. There has

yet to be presented a standardized model that successfully accounts for all of these confounding

variables, so we elect to maintain the aforementioned assumptions as a starting point for the

modeling. Previous literature reports have presented mathematical modeling of protein turnover

rates using similar starting points.[25-32] This then results in Equation 1 where the time-dependent

change in a protein concentration is the difference between the synthesis and degradation rates.

$$\textbf{Equation 1}: \frac{dP}{dt} = k_{syn} - k_{deg}[P]$$

This model assumes that the concentration of an individual protein ([P]) in every location is

under the control of a zero-order synthesis rate (k_{syn}) and a concentration-dependent degradation

(k_{deg}) step. The assumption of zero-order synthesis suggests that the precursor is stable and

unresponsive to protein concentration, while a first-order rate for degradation suggests that there

is no regulation of degradation other than protein concentration. In general, the rates k_{syn} and k_{deg}

need not be constant as they are under the control of numerous exogenous factors that may vary in

time. However, we now formalize our final assumption: protein homeostasis. This assumption is

that the multiple processes regulating each protein concentration are in dynamic equilibrium so

that these rates are constant for a given experimental condition. Under these assumptions, both

synthesis and degradation for a given protein are equal, ensuring that the number of proteins

produced is equal to the number of proteins lost. Therefore, during the measurements d[P]/dt=0,

leading to the relationship:

$$\textbf{Equation 2}: [P] = \frac{k_{syn}}{k_{deg}}$$

Our hypothesis is that the expression of the ApoE polymorphysms (ApoEx = ApoE2 or ApoE4)

creates a unique steady state or proteostasis across the proteome that can differ from the

concentration of the human wild-type control (ApoE3). The change in protein abundance

(Equation 3) between the two conditions allows us to infer how the ratio of the rates is changed in

the experimental cohorts, but neither parameter is individually identifiable.

$$\textbf{Equation 3}: \mathit{\Delta Abundance} = \frac{d[P]}{dApoE} = \left(\frac{k_{syn}}{k_{deg}}\right)_{ApoEx} - \left(\frac{k_{syn}}{k_{deg}}\right)_{ApoE3}$$

Using metabolic isotope labeling we can add rate information that will distinguish between

changes in synthesis and changes in degradation. Assume that at t=0 our model simplifications

are true, but that the amino acid pool is replaced with a deuterated version. Proteins synthesized

after t=0 are isotopically labeled, and we can measure the time-dependent replacement of old

unlabeled for new labeled proteins. To make this mathematically explicit, we denote the

concentration of normal proteins by [P] and the concentration of deuterated proteins by [P^D]. These

two concentrations now satisfy the initial value problems.

Equation 4: $\frac{d[P]}{dt} = -k_{deg}[P]$, $[P](0) = \frac{k_{syn}}{k_{deg}}$

and

Equation 5: $\frac{d[P^D]}{dt} = k_{syn} - k_{deg}[P^D]$, $[P^D](0) = 0$

These are true because normal proteins are no longer being synthesized (Equation 4) and [P^D]

have no initial concentration (Equation 5). These ordinary differential equations can be solved in

closed-form using standard techniques. The solutions are:

Equation 6: $[P]_t = \frac{k_{syn}}{k_{deg}}\left(e^{(-k_{deg}t)}\right)$

Equation 7: $[P^D]_t = \frac{k_{syn}}{k_{deg}}\left(1 - e^{(-k_{deg}t)}\right)$

Notice that these equations satisfy $[P]_t + [P^D]_t = k_{syn}/k_{deg}$, which is independent of time as it must

be in homeostasis. However, the measurable fraction of deuterated protein over time is given by.

Equation 8: $\frac{[P^D]}{[P]+[P^D]} = 1 - e^{-k_{deg}t}$,

Equation 8 seems to suggest that the degradation rate is the measurable driving force behind the

turnover of old protein and the replacement by labeled protein. However, because the processes

of synthesis and degradation are exactly balanced in the proteostasis condition, we can just as

easily identify the turnover rate as the per-molar synthesis rate: $k_{syn}/[P]$ or as it is commonly called

fractional synthesis[33]. It is important to emphasize that these rates are only properly defined in

homeostasis. Because there is no assurance that homeostasis is equally applied to all proteins

simultaneously[18, 21, 34], we find it conceptually preferable to define the turnover rate as the mean

of the per-molar synthesis and degradation rates:

Equation 9: $k_{turnover} = \frac{1}{2}\left(k_{syn}/[P] + k_{deg}\right)$

As stated above, each experimental mouse cohort will have a unique homeostasis with a

protein-specific synthesis and degradation rate. Using ApoE3 as our normal control we can

assess how the average of the synthesis and degradation rates have changed with the E2 and E4

polymorphisms (ApoEx).

Equation 10:

$\Delta Turnover = \frac{1}{2}\left(\left(\frac{k_{syn}(ApoE_x)}{[P](ApoE_x)} - \frac{k_{syn}(ApoE_3)}{[P](ApoE_3)}\right) + \left(k_{deg}(ApoE_x) - k_{deg}(ApoE_3)\right)\right)$

This means that if, for example, the ApoE polymorphism increases a proteins concentration

(+$\Delta Abundance$, Equation 3) the $\Delta Turnover$ (Equation 10) will highlight whether the change in

proteostasis was driven by an increase in $k_{syn}/[P](ApoEx)$ or a decrease in $k_{deg}(ApoEx)$ because the

sign of $\Delta Turnover$ will be different for each possibility. Together the unique $\Delta Abundance$ and

$\Delta Turnover$ for each protein identifies whether the differences in proteostasis are primarily due to

changes in synthesis or degradation. Graphing these values produces a plot where each quadrant

(Figure 1C) has meaning. For example, a positive x-axis (+Δabundance) and y-axis (+Δturnover)

suggest that synthesis increases (Syn↑). Conversely, a protein with lower expression levels (-

Δabundance) between ApoE genotypes, could result from less synthesis (Syn↓) if turnover rate

decreases (-Δturnover) or more degradation (Deg↑) if the protein turnover rate increases

(+Δturnover). Since each measurement has independent noise, a nonrandom grouping of multiple

proteins from a functionally related ontology within a quadrant is an important metric of

confidence that the cell is regulating protein expression to change biochemical functions (Figure

1D and 1E).

Mouse Handling

All animal handling experiments were authorized by the Brigham Young University Institutional

Animal Care and Use Committee (IACUC protocol #191102). The mouse model employed for

this study consists of C57BL/6 transgenic mice with homozygous genotypes for each of the three

human ApoE alleles (ApoE2, ApoE3, ApoE4, n=24/allele, see Supplemental Table S1) under the

GFAP promoter (JAX# 004632, 004633, 004631). Notably, this model has provided valuable

insight into genotype-specific effects of ApoE in a large number of other experiments[35-46]. This

study does not encompass differences from wild-type mice. The transgenic mice were selected

with deliberate focus on ApoE isoforms rather than wild-type conditions, or age differences. The

findings reported in this publication use fold change relative to ApoE3 to minimize the GFAP

promoter variable as reported previously[47-49]. Because mouse ApoE has a low sequence identity

(77%) and a different transcription promoter, the human ApoE3 model is the best control for

comparison.   While we recognize the limits of a transgenic model, this study provides valuable

identification of *in vivo* patterns which can be confirmed in future ApoE knock-in mice models

and human studies. These results refer solely to the effects of ApoE isoform differences, rather

than Alzheimer's Disease. Any claims regarding Alzheimer's disease are made solely to highlight

similarities between current ApoE/AD research and our observations to create a holistic

mechanistic hypothesis.

Mice were randomly selected for replicate designation and timepoint based on availability. They

were all 6–8-month-old, retired breeders with no signs of disease or neurological dysfunction.

There were no exclusions among this group. Specific cohort denominations and animal numbers

can be found in Table S1. Blinding was not used during any portion of this experiment as it was

necessary to compare groups at each point. Mice were housed together in the same room of the

facility at the same time. Mice had *ad libitum* access to water and standard nutritional rodent feed

(Teklad 8604) while housed in a temperature-controlled environment of ~24 °C. This environment

included a 12-hr circadian cycle. To initiate turnover rate measurements, mice received an

intraperitoneal (IP) injection of sterile D_2O 0.9% w/v saline (35 μl/g body weight) calculated to

increase internal D_2O concentrations to an initial 5% of overall water weight (w/w). Mice were

then given 8% D_2O as the sole hydration source for the remainder of the experiment. This was

done to maintain overall internal water at 5% D_2O enrichment. Mice were sacrificed according to

the following timepoints post IP injection: day 0 (no D_2O injection), hour 6, day 1, day 4, day 16,

and day 32. Mice were euthanized via CO_2 asphyxiation followed by bilateral thoracotomy. Blood

was collected via cardiac puncture for D_2O enrichment calculations. Brains were divided sagittally

into respective hemispheres. Relevant organs including brain and liver were flash frozen on blocks

of solid CO_2. Tissues were stored frozen at -80 °C until processing.

Tissue Preparation

Singular brain hemispheres and liver sections were homogenized in lysis buffer (25mM

Ammonium Bicarbonate treated with diethylpyrocarbonate and ThermoScientific Halt Protease &

Phosphatase Inhibitor Cocktail) for 60 sec using a MP FastPrep-24 homogenizer. Homogenized

samples were centrifuged for 15 minutes at 14,000xg to separate them into cytosolic and

membrane isolates. The membrane pellet was resuspended in lysis buffer and centrifuged for 15

minutes at 14,000xg a total of three times to remove cytosolic components. Each fraction was

resuspended in 5% SDS. Aliquot concentration was measured via a Pierce™ BCA Protein Assay

Kit purchased from ThermoFisher Scientific, and 50 μg of protein were prepared according to S-

Trap™ documentation (cytosol and membrane fractions were prepared separately). Proteins were

digested with trypsin Lys/C overnight at 36°C. Resultant peptides were dehydrated in a

ThermoScientific Savant SPD131DDA SpeedVac Concentrator and resuspended at a final

concentration of 1 μg/μL in buffer A (3% acetonitrile, 0.1% formic acid).

LC-MS

Samples were separated and measured via liquid chromatography-mass spectrometry (LC-MS)

on an Ultimate 3000 RSLC in connection with a Thermo Easy-spray source and an Orbitrap Fusion

Lumos. Peptides were pre-concentrated with buffer A (3% acetonitrile, 0.1% formic acid) onto a

PepMap Neo Trap Cartridge (particle size 5 μm, inner diameter 300 μm, length 5 mm) and

separated with an EASY-Spray™ HPLC Column (particle size 2 μm, inner diameter 75 μm, length

25 mm) with increasing buffer B (80% acetonitrile, 0.1% formic acid) gradient:

0-5 min, 0 to 5% B; 5-87 min, 5 to 22% B; 87-102 min, 22 to 32% B; 102-112 min, 32 to 95%

B; 112-122 min, 95% B; 122-125 min, 95 to 2% B; 125 to 127 min, 2% B; 127-129 min, 2 to

100% B; 129-132 min, 100% B; 132-133 min, 100 to 2% B; 133-135 min, 2% B; 135-137 min, 2

to 100% B; 137-140 min, 100% B; 140-142 min, 100 to 0% B; 142-144 min, 0% B.

The MS-based data-dependent acquisition method was set to a 3 second cycle time. MS1 scans

were acquired by the Orbitrap at a resolution of 120,000. Precursors with a charge > 1 and < 6

were selected for MS2 fragmentation. MS2 scans of CID precursor fragments were detected with

the linear ion trap at a scan rate of 33.333 Da/sec with a dynamic injection time. CID collisions

were set to 30% for 10ms. A 60 second dynamic exclusion window was enabled; isotopes and

unassigned charge states were excluded. The deuterium labeling information was collected

separately in an MS1-only acquisition with the Orbitrap at a resolution of 60,000 as previously

described by Naylor et al.[1] The mass spectrometry proteomics data have been deposited to the

ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier

PXD044460

Raw Data Processing for Peptide Identification and Label-free Quantitation

Raw files were searched against the 2022 Uniprot/Swissprot *mus musculus* FASTA (containing 17144 entries) using Peaks Studio v.11 (Bioinformics Solutions Inc.) for label-free quantitation (LFQ) analysis. During the data refinement step, the feature "associate feature with chimera [DDA]" was selected to deconvolute scans with co-eluted isobaric peptides. The parent mass error tolerance was set to ±15 ppm and the fragment mass error tolerance was set to 0.5 Da. Cysteine carbamidomethylation was set as a fixed modification, and both methionine oxidation and pyro-glu from glutamine were set as variable modifications in the search. Digest mode was set to semi-specific for the trypsin-lysC enzyme mix allowing for ≤ 3 missed cleavages and the peptide length range was set to 6 – 45 amino acids. The false discovery rate (FDR) for peptide matches was set to 1%, and protein ID significance was set to -10log(P-value) ≥ 15 for each identified protein.

Peaks Studio (Bioinformics Solutions Inc.) was also used to search raw files for use in Deuterater[1] software. The raw files were searched against the 2021 Uniprot/Swissprot *mus musculus* FASTA (containing 17144 entries). Peptide searches were performed using trypsin/lysC semi-specific digest with a tolerance of ±20ppm and missed cleavages ≤ 3. Carbamidomethylation was set as a fixed modification and pyro-glu from glutamine and methionine oxidation were set as variable modifications. Within the Peaks Studio DB module, proteins were identified with two or more unique peptides at an FDR of 2% and significance was set to -10log(P-value) ≥ 15 for each identified protein.

Protein ΔAbundance Analysis

The group of mice used in this paper were divided into two male and two female groups for analysis. Each group produced a dataset for cytosolic proteins and another dataset for membrane

proteins. Please refer to the *Experimental Design and Statistical Rationale* section for more

information resulting in a total of eight datasets.

Data filtering, normalization, and quantitative calculations were performed independently for

each dataset following standardized metrics for data quality and analysis following the process

described by Aguilan et al. [50] Each Peaks Studio DB *protein.csv* output dataset contains the

proteins identified in the analysis and the expression values (relative abundance) for each protein

in each sample are labeled as "Area". This output was filtered to retain only the top proteins in

each protein group and proteins with at most one missing protein "Area" value per genotype (i.e.,

$n - 1$/genotype/dataset). Subsequently, protein "Area" values in the dataset underwent $\log_2$

transformation. The distribution of these protein "Area" values was mean centered by subtracting

by the average protein "Area" from each protein "Area" within the sample. To ensure

comparability across samples, the distribution width was also normalized between samples by

calculating the correlation slope between these total average protein "Area" values across all

samples and the individual sample values. Each protein "Area" in a sample was then divided by

the corresponding sample slope. For samples with a missing protein "Area" value, imputation was

carried out using the scikit-learn KNN imputer function module in python with the two closest

neighbors.[51]

Protein fold change (FC) values, which represent the relative change in protein abundance values

("Area") compared to a reference, were calculated, and used as a metric of change in abundance

(Δabundance). For this study, FCs were calculated for protein expression values in ApoE2 mice

and ApoE4 mice with ApoE3 expression values as reference, respectively. As per Aguilan et al.'s

methodology, an F-test was employed to assess the variance between protein expression values

before performing p-value calculations for statistical significance. To evaluate the statistical

significance of expression levels in each comparison, a two-sample independent t-test

(homoscedastic) was employed for proteins with an insignificant F-test result and a two-sample

independent t-test (heteroscedastic) for proteins with a significant F-test result. Both the F-test and

t-test calculations were conducted with the Scipy python package.[52]

Protein FC values were averaged across all datasets for each respective comparison. This

produced a single set of "Area" (expression value) FCs for each comparison. Please note that both

the ApoE2 vs ApoE3 (E2vsE3) comparison and the ApoE4 vs ApoE3 (E4vsE3) use the same list

of quantified proteins. As outlined by Van den Berg, protein FC values from individual

comparisons were range scaled using the following formula prior to ontology exploration [53]:

$$\textbf{Equation 11}: \ x'_{ij} = \frac{x_{ij} - \bar{x}_\iota}{x_{i_{max}} - x_{i_{min}}}$$

Where x'_{ij}, x_{ij}, $\bar{x}_\iota$, x_{imax}, and x_{imin} are the scaled FC value, non-scaled FC value, mean FC, largest

FC, and smallest FC, respectively. Range scaling was selected because it captures relative change

in protein expression while considering the full range of values specific to the dataset. These scaled

FC values will be utilized in functional analyses as described in the *Ontology-level Calculations*

section below. The python script created for the steps outlined in this quantitative analysis can be

found in the GitHub repository as detailed in the *supplementary data* section of this paper. The

change in abundance was validated in samples using Data Independent Acquisition (DIA)[54] to test

for reproducibility of the ontology level changes (Supplementary table S6).

Protein ΔTurnover Rate calculation

Protein turnover rate values were calculated using Deuterater[1] v5. This software uses an accurate

mass and time database to extract peptide isotope patterns from LC-MS (MS1) centroided data

utilizing feature identifications (e.g. retention time, mass, peptide ID, etc.) obtained from MS/MS

data (refer to the *Raw Data Processing for Peptide Identification and Label-free Quantitation*

section above).

Isotope patterns were extracted from MS1 raw data with an extraction retention time window of

1.5 min and an m/z error limit of $\leq$ 30 ppm. The n-value represents the number of available

positions on a peptide where deuterium can replace hydrogen. In the *theory generation* step,

peptides with data missing in an extracted file are removed, and the n-value is calculated for

remaining extracted peptides based on known quantities for each amino acid[55, 56]. Subsequently,

Fraction New measures the amount of turnover rate for each peptide in a file by calculating

changes in neutromer abundance and spacing[1]. These calculations were performed using the

average between M0 and the highest isotope peak for peptides meeting specified criteria, including

peptide n-values greater than 5, a minimum peptide sequence length of 6, and a minimum allowed

M0 change of 0.04. In the *Rate Calculation* stage, the data from the *Fraction New* step is fitted to

a kinetic rate curve using **Equation 8** from our proteostasis model. Turnover rates were calculated

for peptides that met a specified criterion, including a minimum of 3 non-zero peptide timepoints,

and measurement deviation of less than 0.1, as previously reported[19]. The asymptote value is

assumed to be 1 in the first iteration of analysis for proteins, but not for lipids where multiple pools

of the same lipid are frequently observed[57].

After the Deuterater[1] analysis, all proteins with a valid turnover rate value (Rsq $\geq$ 0.6, combined

unique peptides > 1, combined rate > 0) grouped by allele cohort, and the average turnover rate

value was calculated for each protein in the cohort, respectively. These protein turnover rate values

were $\log_2$ transformed, and the protein turnover rate FC was calculated as the difference of the $\log_2$

rates. The ApoE2 mice and ApoE4 mice were compared to the reference ApoE3 mice, resulting

in a single set of protein turnover rate FCs for each comparison. To standardize the protein

turnover rate FCs, auto scaling [53] was applied, where x'_{ij}, x_{ij}, $\overline{x_i}$, and s_i are the scaled turnover rate

FC value, non-scaled turnover FC rate value, mean turnover FC rate and turnover rate FC standard

deviation, respectively:

$$\textbf{Equation 12}: x'_{ij} = \frac{x_{ij}-\overline{x_i}}{s_i}$$

Auto scaling was implemented because it considers both the mean and standard deviation to

reduce the effects of outliers and variation in the data while preserving the ability to focus on small

changes. It is important to note that because of the signal to noise requirements fewer proteins had

valid turnover rate FC values than quantifiable abundances. Consequently, proteins with turnover

rate FCs represent a smaller subset population in comparison proteins with expression value FCs

calculated from "Area" values. These protein turnover rate values were used for ontology analysis

as outlined in the *Ontology-level Calculations* section below. For further reference, the python

script created to process the *calculated_rates* output from Deuterater[1] can be found in the GitHub

repository, as detailed in the supplementary data section of this paper.

Ontology-level Calculations

The StringDB[13] multiprotein tool was employed to calculate group FC values for functionally-

related protein groups (ontologies) regardless of statistical significance. To streamline the analysis

and reduce the number of redundant term ID (ontologies), ontologies were selected only from the

following established: *GO Process*, *GO Function*, *GO Component*, *KEGG*, *Reactome*, and

WikiPathways. To quantify the representation of each ontology, the "observed gene count" was

divided by the "background gene count" to calculate the "Ontology_coverage (%)" for each

ontology. Only ontologies with ≥25% were included in this analysis. This latter criterion ensures

that the identified ontologies are sufficiently represented in the data (Table S4 and S5).

The "matching proteins in your network (labels)" was used to associate each observed protein

in the ontology with the calculated "Area" FC and turnover rate FC, respectively, for both the

E2vsE3 and the E4vsE3 comparison. Next, the average protein "Area" FC and turnover rate FC

was calculated for each identified ontology by averaging the FC values of proteins within that

category. This step summarized the collective expression and turnover rate changes of proteins

within specific functional groups for each comparison.

To assess the statistical significance of the FC values within each ontology, a one-sample t-test

was performed with null hypothesis (H0) stating the average ΔAbundance (**Equation 3**) for the

ontology is equal to 0, and the alternative hypothesis (Ha) indicating that it is not equal to 0. This

statistical test is used to determine whether the observed changes in protein expression for the

ontology were statistically significant. To account for multiple comparisons and maintain a

controlled false discovery rate (FDR), the Benjamini-Hochberg correction (BH-PV) was

calculated for the resulting p-values (FDR = 0.25). Ontologies with a BH-PV < 0.05 were

considered statistically significant. In the case of highly similar ontology with significant changes,

the ontology with the most proteins was selected to represent the results. The Python code used to

analyze StringDB and calculate the FCs can be found in the GitHub repository, as detailed in the

supplementary data section of this paper.

RESULTS

Proteome Ontology Analysis

Figure 1. Testing for changes in regulation of the brain proteome **A.** Homozygous ApoE transgenic mice (ApoE2, E3, or E4, n = 24 each) were given 8% D2O drinking water and sacrificed at specific timepoints ranging from day 0 to day 32. Tissues were prepared using the S-Trap protocol and analyzed via LC-MS. Data-dependent acquisition was used to collect data for LFQ analysis and MS1 level data was used to calculate turnover rate values for each protein. Resulting spectra from MS/MS acquisitions were analyzed by Peaks Studio (Bioinformatics Solutions Inc.) peptide-protein identification (IDs) and quantitation while Deuterater[1] software was used for turnover rate calculation. **B.** Proteostasis model where protein expression levels and turnover rates are a function of synthesis and degradation. **C.** Regulation of synthesis and degradation can be inferred from Δabundance (x-axis) and Δturnover (y-axis) and visualized using a proteostasis plot. **D & E.** Proteostasis plot showing 276 and 288 protein ontologies with significant Δabundance (red circles) in ApoE2 (panel D) and ApoE4 (panel E)mice relative to ApoE3 mice (E4vsE3) (BH-PV < 0.05).

In our analysis, we identified 4,849 proteins in the brain tissue across the three ApoE-isoform

groups (n = 47). From these, we determined protein abundance and turnover rate FCs for

comparisons: ApoE2 vs. ApoE3 (E2vsE3) and ApoE4 vs. ApoE3 (E4vsE3). Here, ApoE3 serves

as the reference 'normal' control. We quantified 3,532 abundance FCs for both the E2vsE3 and

E4vsE3 comparisons (Supplementary Table S2). A smaller number of turnover rate FCs were

obtained: 1,430 for E2vsE3 and 1,405 FCs for E4vsE3 (Supplementary Table S2) because of the

more rigorous statistical filtering criteria.

For this analysis, we focused on ontologies from six databases: *GO Function*, *GO Component*,

GO Process, WikiPathways, Reactome, and *KEGG*. Using the StringDB results, we calculated the

average abundance FC (Δabundance) and turnover rate FC (Δturnover) for the proteins observed

in each ontology (Please refer to the *Ontology-level Calculations* in the methods section). This

yielded ~2700 ontology-level comparisons between average Δabundance and Δturnover

calculations for both E2vsE3 and E4vsE3 (Supplementary Table S4). The interpretation (Figure

1C) relies on the traditional understanding of protein turnover, contextualizing changes in protein

expression. It offers a lens to assess the variances in the steady states of ApoE genotypes[58]. Using

a one-sample t-test, we discerned which ontologies deviated significantly from a median

Δabundance of 0. In the E2vsE3 comparison, we identified 284 protein ontologies with notable

Δabundance (BH-PV < 0.05) (Figure 1D). For the E4vsE3 comparison, 287 protein ontologies had

significant Δabundance (BH-PV < 0.05) (Figure 1-E).

The box plots in Figures 2 through 5 encapsulate the Δabundance and Δturnover for ontologies

with marked Δabundance shifts. To maximize visibility and to accommodate for space limitations,

these boxplots do not contain outlier points but supplementary Figures 2 through 5 contain the

boxplots with outlier points. Given that some ontologies are repetitive, proteins depicted in the box

plots might appear in multiple ontologies with analogous names/functions. When faced with such

redundancies, we typically chose the ontology with superior coverage (Observed/Total) for

representation. As a convention, each ontology is presented in an "*ontology name (n)*" format,

where (n) indicates the count of quantified proteins within that ontology. Overlap between similar

ontologies is shown in the heatmap.

ApoE Isoforms Modulate Synthesis and Degradation of Endocytic Vesicle Components

We observed that multiple ontologies with significant Δabundance were associated with

endocytosis and vesicular processing (Figure 2). Specifically, the general *Endocytosis (158)*

ontology demonstrated increased Δabundance and decreased Δturnover, suggesting reduced

degradation in both ApoE2 and ApoE4 compared to ApoE3. In the context of ApoE2, *Clathrin-*

mediated endocytosis (82), *Clathrin binding (35)*, and *Clathrin coat (26)* mirrored the same

↓degradation effect observed in endocytosis, while *SNARE complex (32)* showed diminished

Δabundance and Δturnover, suggesting a decline in protein synthesis compared to ApoE3.

Moreover, ApoE2 expression led to significant alterations in several regulatory ontologies tied to

endocytosis and vesicular processes, such as: *Endocytic recycling (34)* (↑synthesis), *Early*

endosome (↑synthesis), and *Regulation of endocytosis (15)* (↑synthesis). In ApoE2, proteins

related to *Lysosome Vesicle Biogenesis (18)* have lower degradation while *Regulation of*

Endocytosis (107) had increased synthesis leading to higher abundance of these protein groups and

presumably more efficient endolysosomal function.

In both ApoE2 (E2vsE3) and ApoE4 (E4vsE3) we noted diminished degradation of general

lysosome (146) proteins. Within this general ontology, the *lysosomal membrane (70)* ontology had

diminished Δabundance and Δturnover only in the ApoE4 group, suggesting less synthesis of the

membrane components compared to ApoE3. This is consistent with large lysosomal vesicles stored

in ApoE4 cells[59, 60]. In ApoE4 mice there was higher Δabundance and Δturnover (↑synthesis) of

Phosphatidylinositol binding (90) relative to ApoE3. Conversely, there was a decline in both

Δabundance and Δturnover (↓synthesis) of *SNARE interactions in vesicular transport (19),*

Synaptobrevin 2-SNAP-25-syntaxin-1a complex (5), and *SNARE complex (32).*

Figure 2. ApoE4 expression disrupts endosomal maturation and ApoE2 increases lysosomal capacity. **A-B.** Bar plot displaying Δabundance (orange) and Δturnover (blue) of proteins detected in all experimental cohorts for significant* ontologies related to endolysosomal trafficking in the E2vsE3 comparison (A) and in the E4vsE3 comparison (B). **C.** Heatmap displaying % of proteins shared across the endolysosomal ontologies with significant* Δabundance. *(*BH-PV < 0.05).*

ApoE Isoforms Modulate Synthesis and Degradation of Mitochondrial Components

Our analysis identified significant Δabundance (BH-PV < 0.05) changes for multiple ontologies related to mitochondrial components (Figure 3). In the E4vsE3 comparison, these ontologies included mitochondrial membranes, protein transport, and morphology (Figure 3A and Figure S3A). Each of these ontologies displayed a negative Δabundance coupled with a positive Δturnover, signifying ↑degradation. We also detected ↓synthesis of *mitochondrial calcium ion transmembrane transport (12)* and *mitophagy (18)*. In contrast, ApoE4 *mitochondrial matrix (159)* also had ↑degradation. (Figure 3B and Figure S3B) The percentage of overlapping proteins in each mitochondrial component ontology is displayed in Figure 3C. The key finding from these ontologies is that within the ApoE2 mice there is a coherent increase in the degradation of all mitochondrial components consistent with an increase in mitochondrial degradation as an entire unit. In contrast, the ApoE4 tissues show discordant changes in matrix versus membrane proteins suggesting that mitochondrial maintenance is more piecemeal and that mitophagy may be less efficient as previously suggested in the literature [61]. Both the ApoE2 and the ApoE4 results are synergistic with the changes in lysosome dynamics discussed above. In ApoE2 more efficient lysosomal processing will facilitate mitophagy based quality control while the inhibited lysosomal processing would inhibit mitophagy and make the ApoE4 more reliant upon individual protein replacement strategies.

Figure 3. ApoE genotype differentially regulates mitochondrial proteostasis.   **A-B.** Bar plot displaying Δabundance (orange) and Δturnover (blue) for ontologies of proteins detected in all experimental cohorts related to mitochondrial components with significant* Δabundance in the E2vsE3 comparison (A) and in the E4vsE3 comparison (B). **C.** Heatmap displaying % of proteins shared across the mitochondrial ontologies with significant* Δabundance. *(**BH-PV < 0.05**).*

ApoE4 Disrupts Metabolic Pathway Control

We observed significant changes in Δabundance (BH-PV < 0.05) across multiple ontologies related to energy production (Figure 4). ApoE2 resulted in lower expression of levels of *Pyruvate metabolism (32)*, *Citrate cycle (TCA cycle) (26)*, and *Glycolysis/Gluconeogenesis (41)*. These reductions were primarily attributed to decreased synthesis (↓synthesis), a trend that was also evident in the *Oxidative stress and redox pathway (48)* proteins which protect the cell from reduced oxygen species. Notably, *Fatty acid beta-oxidation (25)* demonstrated reduced Δabundance coupled with increased Δturnover (↑degradation) in ApoE2 (E2vsE3), suggesting a potential decrease in fatty acid catabolism and an increase in the use of fatty acids for building complex lipids. In contrast, in the ApoE4 mice, major energy production pathways such as *Fructose and mannose metabolism (21), Pyruvate metabolism (32),* and *Glycogen metabolism (22)* all exhibited increased Δabundance and Δturnover, pointing towards enhanced synthesis of enzymes involved in carbohydrate metabolism in the E4vsE3 comparison and an increased reliance on carbohydrates for energy similar to previous observations [62, 63].

Figure 4. ApoE2 and ApoE4 expression drive changes in cellular fuel selection **A-B.** Bar plot displaying Δabundance (orange) and Δturnover (blue) for ontologies of proteins detected in all experimental cohorts belonging to oxidative phosphorylation with significant* Δabundance in the E2vsE3 comparison (A) and in the E4vsE3 comparison (B). **C.** Heatmap displaying % of proteins shared across the oxidative phosphorylation ontologies with significant* Δabundance. *(*BH-PV < 0.05)*.

ApoE Isoforms and Ubiquitin-Proteasome Pathway Activity

The proteasome related ontologies exhibited significant changes in regulation due to ApoE isoforms (Figure 5). For both the E2vsE3 and E4vsE3 comparisons, we identified pronounced increases in Δabundance and reductions in Δturnover (↓degradation) associated with the *proteasome complex (47)* Furthermore, an increased Δabundance and Δturnover (↑synthesis) of proteins involved in the *Regulation of ubiquitin-dependent protein catabolic process (62)* was statistically significant in both comparisons (BH-PV 0.05).

The *proteasome regulatory particle, base subcomplex (11)* displayed ↑synthesis in the E2vsE3 comparison and ↓degradation in the E4vsE3 comparison. Meanwhile, proteins within the *Proteasome regulatory particle, lid subcomplex (6)*, demonstrated significant Δabundance due to ↓degradation in the E4vsE3 comparison with ApoE4 expression. However, these changes were not significant in the E2vsE3 comparison. Additionally, in ApoE4 we noted ↑synthesis in the *Negative regulation of proteasomal ubiquitin-dependent protein catabolic process (21)*, ↓degradation *Deubiquitination (79)*, and ↓synthesis *Hsp70 protein binding (23)*. These observations suggest a nuanced regulation of the ubiquitin-proteasome system (UPS) in association with ApoE isoforms. Hsp70 proteins are often deemed pivotal regulators of proteasome activity[64]. These changes suggest a significant reduction in the protein quality control for ApoE4 tissue (Figure 5C).

Figure 5. Proteasomal activity decreases with ApoE4 expression.   **A-B.** Bar plot displaying Δabundance (orange) and Δturnover (blue) of proteins detected in all experimental cohorts for several ontologies related to proteasomal activity with significant* Δabundance in the E2vsE3 comparison (A) and in the E4vsE3 comparison (B). **C.** Heatmap displaying % of proteins shared across the ontologies with significant* Δabundance. *(**BH-PV < 0.05**).

517
Quantifying ApoE-dependent Shifts in Liver Proteostasis

The liver is the largest producer of ApoE in the body and is also a major receptor of ApoE and

its associated cargo[9, 65, 66]. Therefore, we tested whether the liver tissue from these same

experimental mice would show matching ApoE allele-specific shifts in proteome regulation.

In ApoE2 liver there was not a significant change in any of the endocytic processes relative to

ApoE3 (Table S5). Multiple mitochondrial ontologies in the liver changed in significant ways and

nearly 60% of their proteostasis changes are equivalent to the brain. Most changes in the

mitochondria in the liver with ApoE2 expression involve increased degradation of mitochondrial

components, though there is some reduced synthesis for the mitochondrial envelope and

transmembrane transport. Interestingly, where ApoE2 the brain contains decreased degradation of

proteasomal components, in the liver we observed increased synthesis and greater more

proteasomal capacity similar to published studies[67-69].

In ApoE4/E3 liver comparisons there was not a significant change in any of the endocytic

processes (Table S5) with the exception of endosomal protein localization. In the brain, this

ontology had increased synthesis, while the liver promotes decreased degradation, both of which

result in an increased concentration. As for ApoE4 mitochondrial components, most changes in

the liver involve increased degradation. All of the significant increased degradation ontologies

observed in the brain were observed with the similar increased degradation in the liver, though not

all were significant. Likewise 80% of the significant mitochondrial liver ontologies had the same

proteostasis changes in the brain (Table S5). The 20% differences were due to certain NADH and

ATP synthesis electron transport chain ontologies that were increased synthesis in the brain and

increased degradation in the liver. Similar to the liver comparison of ApoE2 with the proteasome,

in ApoE4 liver data there were no shared proteasome proteostasis changes with the brain [Figure

S7 and S8]. These data suggest that most of the ApoE effects observed in the brain are not global.

DISCUSSION

Exploring ApoE-genotype Effects Through the Lens of Proteostasis

Compared to the neutral ApoE3 allele, expression of ApoE4 heightens the risk for

neurodegeneration, while the expression of ApoE2 is protective[6, 9, 63, 67, 70-73]. We conducted an

experiment to identify how protein homeostasis changes with ApoE genotype in the tissues of

human-ApoE transgenic mice (Fig 1A). Homeostasis is the dynamic control of concentration and

quality in the cell (Figure S2). Traditionally, the dynamic control of protein concentration is

conceptualized as the balance between synthesis and degradation[26] while protein turnover rate is

defined as the time required for a protein to be replaced.[24, 32, 58, 74] We present this as simplified

equations that relate to synthesis and degradation (Fig 1B, see '*Proteostasis Model and Analysis*

Rational' section for more detail). Therefore, changes in protein expression levels (Δabundance)

paired with changes in protein turnover rate (Δturnover) can highlight the regulatory balance of

synthesis versus degradation between conditions (Figure 1B-C).

For instance, to increase [protein] in the experimental condition (resulting in a positive

Δabundance), cells can either elevate synthesis or diminish degradation. Alternatively, to decrease

[protein] (leading to a negative Δabundance), cells might reduce synthesis or increase degradation.

Using the change in protein turnover rates (turnover = ½(synthesis + degradation), *Proteostasis*

Model section Equation 10) we can deduce whether changes in synthesis or degradation led to

changes in abundance (Figure 1B). Therefore a positive Δabundance indicates increased synthesis

when Δturnover is positive or reduced degradation when Δturnover is negative. Conversely, a

negative Δabundance signals increased degradation when Δturnover is positive or decreased

synthesis when Δturnover is negative (Figure 1C). We used this model to identify the ApoE-

dependent changes in proteostasis regulation (Figure 6). Below we discuss how our results unify

a diverse set of literature observations where ApoE-dependent modifications of endosome

trafficking, as well as lysosomal, mitochondrial, and proteosomal function have been reported.

Figure 6. Model comparing the observed changes in proteostasis for ApoE2 and ApoE4. The arrows are color coded to represent the different pathways impacted in both ApoE2 and ApoE4 when compared to ApoE3

ApoE isoforms modify Endocytic/Endosomal trafficking

Previous research has highlighted the dysregulation of endocytic pathways associated with

ApoE4 expression[59, 60, 70, 75-77]. We detected notable ApoE4-dependent changes in several

ontologies related to endocytosis (Figure 2 and Figure S3). This is in line with what is known about

how ApoE isoforms modify affinity for cell surface receptors, such as LDLR and APOER2[4, 5],

initiating the endocytosis of ApoE along with its content. After this endocytic event, ApoE-laden

endosomes undergo various maturation stages, wherein contents are earmarked for recycling,

delivery, or degradation.

ApoE4 has a higher binding affinity to receptors[4, 5] and is known to induce a trafficking anomaly

in the early endosome[78], then lead to accumulation and enlargement of lysoendosomal

compartments[59, 60]. Following an endocytic event, the clathrin coat dismantles, allowing vesicles

to transit to various destinations for cargo release. This fusion mechanism leans heavily on SNARE

and SNAP-receptor proteins, also pivotal for exocytosis. In our study, we observed diminished

synthesis of SNARE and SNAP ontologies in ApoE4 mice which may disrupt vesicle fusion

between organelles[79] and in response to exocytic sequences[80]. Our study revealed a reduced

degradation of proteins associated with *Clathrin-mediated endocytosis (82),* increased synthesis

of *PICALM (90),* and reduced synthesis of the *lysosomal membrane (70)* (Figure 2B, Figure S3B)

in the presence of ApoE4. Priyanka et al. also noted a decline in clathrin-mediated endocytosis in

astrocytes with ApoE4 expression[81] while *in vivo* studies identified alterations in early endosome

populations in 18- and 25-month-old ApoE4 mice[82]. This mechanism further incorporates

phosphatidylinositol binding proteins like PICALM[83]. Before undergoing lysosomal degradation,

endosomes transition to the late endosomal phase. Our findings suggest that ApoE4 expression

reduces the synthesis of both late endosomal and lysosomal membranes. Although the general

lysosome (146) ontology exhibits increased degradation with ApoE4 expression, if we look

specifically at the membrane components of this ontology then the ApoE4 specifically has less

total protein due to lower synthesis. These results are consistent with previous observations of

large-volume lysosomes which would have a low membrane surface/volume ratios accumulating

in the in ApoE4 cells [59, 60]. Collectively, these results underscore multiple points of failure due to

ApoE4-associated inhibition of endosomal maturation and stalled lysosomal functions as

previously observed [78, 82, 84].

The E2vsE3 tissue had similarities in vesicle-centric ontologies. Notably, there was a decline in

the degradation of endocytosis and clathrin protein-related ontologies, and SNARE complexes saw

reduced synthesis. This implies that ApoE2 also modifies vesicle endocytosis. However, the

changes suggested a more streamlined regulation of endolysosomal events with ApoE2 (E2vsE3).

This again agrees with literature reports of modified receptor binding with ApoE2 having lower

affinity while ApoE4 has a higher affinity[4, 5]. This coupled with lower degradation of proteins

within the lysosome vesicle biogenesis ontology for the E2vsE3 comparison, a process intrinsically

tied to endosomal trafficking and central to lysosomal adaptation[85], suggests a tighter control of

endocytic events and better lysosomal quality with ApoE2 expression. These observations agree

with previous research on astrocytes indicating ApoE2 expression increases lysosomal activity

relative to ApoE3 and ApoE4 expression[86].

ApoE-dependent changes in Mitochondrial Proteostasis

We observed ApoE-dependent changes in mitochondrial proteostasis that were consistent with

modified autophagy and lysosomal function. In ApoE4 (E4vsE3) mice, we measured elevated

degradation in *mitochondrial membrane (406)*, *mitochondrial inner membrane (303)* (Figure 3

and Figure S4), *Cristae formation (11)*, *Mitochondrial fusion (13)*, and *mitochondrial transport*

(72) with no accompanying change in general *mitochondrial matrix (159)* and decreased synthesis

of *mitophagy (18).* Mitochondrial membrane complexes play critical functions in cellular

homeostasis—such as energy production, calcium level modulation, apoptosis, and the regulation

of reactive oxygen species (ROS) [87]. Prior research has documented alterations in the

mitochondrial membrane's integrity in the context of neurodegenerative diseases [88, 89].

Most mitochondrial proteins are encoded on the nuclear DNA and are transported into the

mitochondria through translocases (TIM and TOM)[90, 91]. These translocases interact with the many

inner mitochondrial membrane (IMM) folds that make up the cristae via the mitochondrial cristae

organizing system (MICOS).[92] Our observations indicate a change in mitochondrial protein

import, especially evident in the higher degradation of the *TOM complex (10)*, *cristae formation*

(11), and the *MICOS complex* (7) ontologies. (See Figure 5B and Figure S5B). The MICOS also

plays a vital role in cristae organization and the function of respiratory complexes.[93] Disruptions

in MICOS have been documented to modify cristae structure[94], and recent studies associate altered

MICOS protein expressions with ApoE4 manifestation[89]. The MICOS literature also report

evidence of mitochondrial fusion and fission imbalance in Neuro-2a cells expressing ApoE4.[89]

Previous analysis of AD brains indicated diminished protein levels connected to mitochondrial

fusion/fission[95], which our data supports as a degradation driven loss of fusion proteins (See Figure

3B and Figure S3B).

Mitochondria and the endoplasmic reticulum (ER) collectively form the mitochondria-

associated membrane (MAM), which has implications in AD pathology[96-98]. These MAMs

regulate oxidative phosphorylation, calcium levels, protein degradation, and mitochondrial

membrane organization. Our dataset elucidates an ApoE4-induced *MAM (57)*, marked by

increased degradation contrasted against ApoE3 (Figure 3 and Figure S4). Our results support

ApoE4-related MAM instability by diminished synthesis of chaperone complexes, mitophagy, and

calcium transport.

In contrast, ApoE2 mice display increased degradation of mitochondrial membrane ontologies

with a matching increase in the degradation of the matrix proteins. Although both ApoE2 and

ApoE4 mice revealed changes in the mitochondrial membrane and transport, our ApoE2 findings

suggests that there is a cohesive organelle-wide response involving both membrane and matrix

proteins. Drawing from our preliminary insights into endolysosomal systems discussed above, we

postulate this might be evidence of superior mitophagy in ApoE2. Additionally, we theorize, as

described in the literature,[61-63, 89, 99] that the alterations in mitochondrial proteostasis induced by

ApoE4 culminate in mitochondrial dysfunction.

ApoE disrupts ATP production

There is an increasing body of research on ApoE genotype-specific effects on ATP

production[100-102]. Moreover, compromised bioenergetic pathways are identified as a distinct

characteristic of neurodegeneration[103-106]. Several studies highlight an ApoE-related shift towards

glycolysis and diminished oxygen consumption in brain tissues[103, 107]. Our data align with these

observations, revealing a heightened synthesis of ontologies suggesting that ApoE4 expression

leads to a more pronounced reliance on glycolytic pathways compared to ApoE3 (see Figure 4B,

Figure S4B). These ontologies include *Fructose and mannose metabolism (21), Pyruvate*

metabolism (32), and *Glycogen metabolism (22)*. We posit that this increased reliance on

carbohydrate metabolism is a consequence of the lack of cohesive mitochondrial maintenance.

Another study focusing on glycolytic and OXPHOS activities found that ApoE4 expression leads

to compromised mitophagy and elevated ROS levels in brain cells, a trend our data supports[108].

The impaired mitophagy was attributed to cholesterol buildup in lysosomes. While we haven't

analyzed cholesterol or ROS levels, our data does indicate reduced synthesis in *mitophagy (18)*,

lysosomal membrane (70) and *Detoxification of Reactive Oxygen Species (20)* proteins —potential

indicators of disrupted mitophagy and ROS balance (See Figure 3B, Figure S3B)

In cell culture ApoE2 expression has been associated with enhanced ATP production and

heightened glycolytic activity[103, 107]. In contrast, various studies have shown that ApoE4

expression was associated with diminished ATP production[100, 109-111]. In ApoE2 tissue we observed

decreased abundance in ontologies such as the *TCA cycle (25)*, *Pyruvate metabolism (32),* and

Glycolysis / Gluconeogenesis (41) due to diminished synthesis and augmented degradation. Our

study is averaging together all cell types in the brain and therefore may diverge from cell type-

specific experiments [100]. Collectively, our data accentuates the isoform-specific alterations in

diverse metabolic pathways, and suggests that isolating single cell types from the brain may be a

critical method to test for metabolic changes in response to ApoE isoforms.

Linkages between the Proteasome and Mitochondrial Homeostasis

The proteasome is a key component of proteostasis maintenance and is essential in clearing out

misfolded proteins and saving cells from misfolding stress response.[112] Reduced proteasome

activity has consistently been implicated as a major player in the pathophysiology of

neurodegeneration[23, 61, 67, 69, 113]. ApoE has been shown to directly regulate the cleavage of amyloid

precursor protein (APP) to form amyloidogenic Aβ peptides with ApoE4 allowing increased Aβ

peptide cleavage and plaque deposition.[114] This buildup of Aβ is relevant to proteasomal function

and has been shown to directly inhibit proteasome function leading to increased accumulation of

amyloid plaques.[115, 116]

The proteasome also plays a major role in the mitochondrial quality control, especially in

response to misfolded proteins that disrupt mitochondrial activity[117]. Interestingly, a growing body

of literature suggests that proteasome function can also be disrupted by mitochondrial dysfunction.

For example, oxidation of the 26S subunit of the proteasome due to increased mitochondrial

oxidative stress has been shown to increase the accumulation of ubiquitinated substrates and

decrease proteasomal activity[118]. Notably, we discerned a significant reduction in the synthesis of

HSP70 proteins in E4 which latch onto misfolded or compromised proteins before proteasomal

degradation[68]. This interconnection of the proteasome and mitochondria as well as their consistent

implication in neurodegenerative disease has led some researchers to suggest that dysfunction in

either the proteasome or mitochondria are "two sides of the same coin" leading to a futile cycle of

mitochondrial and proteasomal insult.[119]

Our investigation revealed reduced degradation of both the proteasome and its regulatory

complex in ApoE4 mice (refer to Figure 5 and Figure S6). Additionally, while ApoE4 expression

increased the synthesis of ontologies linked to ubiquitin-proteasome regulation, it also elevated the

synthesis specifically for its negative regulation. Such trends align with existing literature

delineating the impact of ApoE isoforms on proteasomal dysregulation in Alzheimer's disease[67,]

[120]. Alongside our observation of compromised mitochondrial activity in ApoE4 mice, these

findings imply a heightened susceptibility to both mitochondrial and proteasomal damage. The

concurrent malfunction of mitochondria and proteasomes has been historically correlated with

neuronal apoptotic pathways and neurodegeneration, thereby underlining a mechanism through

which ApoE4 exacerbates the risk of neurodegenerative ailments such as Alzheimer's[121].

ApoE2 also exhibited a notable decline in the degradation of proteins associated with ubiquitin

and proteasomal processes, paralleling the ApoE4 response (see Figure 5 and Figure S6). While

both ApoE2 and ApoE4 amplify the regulation of ubiquitin-dependent catabolism, the base

complex of the proteasome (responsible for facilitating the unfolding and admission of

ubiquitinated polypeptides into the proteasome's degradation chamber[122]) reduced degradation of

deubiquitinating proteins was more prevalent in ApoE4. This implies larger proteasome pool in

ApoE4, albeit with more regulation. It's worth speculating that the ApoE2-mediated inhibition of

APP cleavage to form Aβ[114] might also be instrumental in curbing proteasomal dysfunction.

These patterns suggest that both phenomena might be interconnected, with proteasomal

dysfunction potentially instigating ApoE4-associated reductions in ATP production. Beyond the

ubiquitin-proteasome system, autophagy is instrumental in clearing defective mitochondria via

lysosomal degradation. As previously mentioned, our findings substantiate the ApoE4-associated

dysregulation of MAM structures, which potentially results in disrupted mitochondrial

morphology and impaired energy production. Our working hypothesis postulates that ApoE4

expression precipitates a decline in proteasomal activity, culminating in the accrual of

dysfunctional mitochondria and a diminished capacity to eliminate these via autophagy. Drawing

from our data on lysosomal components in ApoE4 and existing literature[78, 123], we surmise that

suboptimal endocytic regulation might directly impact autophagy and the proteasomal oversight

of mitochondrial proteostasis. Conversely, ApoE2 expression is purportedly linked to enhanced

proteasomal capability and autophagy through lysosomal degradation, resulting in fortified

mitochondrial proteostasis. Our objective is to delve deeper into these discoveries and authenticate

our hypothesis across various ApoE models.

Liver Proteostasis Changes Compared to Brain

ApoE is an important lipid transporter that is expressed and integral to many parts of the body

beyondthe brain. Previous literature has shown that despite ApoE2's protective effect against

Alzheimer's Disease, it increases risk for cardiovascular health[66]. This leads us to question whether

the mechanism by which ApoE2 protects against Alzheimer's Disease may in turn be detrimental

and disease causing to other tissue. To determine whether ApoE does indeed elicit a global

response across tissue, we tested and analyzed the liver tissue from our experimental mice in the

same manner as the brain.

The changes in ApoE2/E3 liver proteostasis and ApoE2/E3 brain proteostasis were not

equivalent. Although many trends were similar between brain and liver, few changes were

statistically significant in the liver. Mitochondrial protein localization, transportation, and

organization had shared proteostasis changes between tissues, but general trends from other

ontological changes suggest mitochondrion turnover to be reduced in the liver compared to the

brain (See Figure S2).

ApoE4/E3 liver and ApoE4/E3 brain proteostasis changes were also not the same. This is

principally due to significant proteostasis changes among the endosome, metabolism, and

proteasome pathways in the brain, and that most of which were not significant in the liver (see

table S5). However, comparisons between the ApoE4 liver and brain contained several of the same

proteostasis changes for mitochondria, suggesting there may be some degree of shared effects due

to ApoE4.

Thus, while some cellular pathways may be affected similarly, a global effect specific to ApoE

allele is not supported by our data. We propose the lack of a global ApoE effect is because most

tissues have a large number of apolipoproteins participating in lipid transport [9, 66]. The brain has

limited apolipoproteins compared to other tissues, in that it is limited to only the apolipoproteins

it creates (primarily ApoE and ApoJ). This is due to the blood-brain-barrier, which prevents

apolipoprotein transfer between the brain and the rest of the body. Since lipid trafficking in the

liver has access to multiple apoliproproteins we posit that this may dilute the effect of ApoE

isoforms on the pathways within liver tissue.

CONCLUSION:

In this study, we demonstrated how combining protein abundance and turnover rate unveils

novel insights into the cellular mechanisms governing synthesis and degradation. Utilizing

multifaceted proteomics data, we tracked ontology-level variations among distinct ApoE

genotypes in healthy adult mice. Our findings present *in vivo* evidence that harmonizes with

existing literature, identifying how ApoE4 interrupts endosomal trafficking leading to autophagy

and proteasome activity defects and lower mitochondrial quality in the brain. Concurrently, our

data suggests that ApoE2 enhances brain mitochondrial health by amplifying turnover in the brain

(Figure 6).

SUPPORTING INFORMATION:

Figure S1. Detailed workflow chart describing both the mouse model and stages of analysis.

Figure S2: Kinetic model of protein homeostasis identifying the common sources and sinks of protein in a cell

Figure S3. Abundance and turnover FCs for ontologies related to endolysosomal processes in

A) E2vsE3 and B) E4vsE3.

Figure S4. Abundance and turnover FCs for ontologies related to mitochondrial components in

A) E2vsE3 and B) E4vsE3.

Figure S5. Abundance and turnover FCs for ontologies related to cellular metabolism in A)

E2vsE3 and B) E4vsE3.

Figure S6. Abundance and turnover FCs for ontologies rela ted to protein degradation in A)

E2vsE3 and B) E4vsE3.

Figure S7. Model for comparison of the ApoE2 brain and liver homeostasis shifts relative to

ApoE3. The ApoE2 brain model (left, blue), is the same as in Figure 6. The green model (right)

shows the modifications in ApoE2 liver tissue.

Figure S8. Model for comparison of the ApoE4 brain and liver homeostasis shifts relative to

ApoE3. The ApoE4 brain model (right, red), is the same as in Figure 6. The green model (left)

shows the modifications in ApoE4 liver tissue.

The data generated in this investigation can be accessed via the ProteomeXchange Consortium

via the PRIDE partner repository[124] (http://www.proteomexchange.org/) with the accession

number PXD044460. The repository includes the raw LC-MS files used for both quantitative and

kinetic files used in data analysis. In addition, Peaks Studio (Bioinformics Solutions Inc.) outputs

containing peptide and protein level identification data for both quantitative and kinetic

measurements are included in the repository.

Project Name: Quantitative and Kinetic Proteomics Reveal ApoE Isoform-dependent Proteostasis
Adaptations in Mouse Brain
Project accession: PXD044460
790
Reviewer account details:
Username: reviewer_pxd044460@ebi.ac.uk
Password: C9CUGtZN

The output files from Deuterater software including turnover rate values are found within the

repository while the code is found here (https://github.com/JC-Price/DeuteRater/releases).

Lastly, the code used in the ontology analysis can be found by following this link to the GitHub

repository (https://github.com/natepine/ApoE_Proteomics.git).

The following files are available free of charge via the Internet

.xls file containing Tables S1-S5 and data for all quantifiable proteins and measurable ontologies.

.pdf file containing supplementary figures S1-S7, including a diagram that explains the mouse

model in addition to box plots with additional ontologies.

AUTHOR INFORMATION:

Corresponding Author

John C. Price - Department of Chemistry and Biochemistry, Brigham Young University, Provo,

Utah 84602, United States; Email: drjohncprice@gmail.com

Author Contributions

This manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript. A CRediT statement is provided below

Nathan R. Zuniga: Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Resources, Software, Supervision, Validation, Visualization, Writing - Original Draft, Writing - Review & Editing. **Noah E. Earls**: Data Curation, Formal Analysis, Investigation, Validation, Visualization, Writing- Original Draft, Writing - Review Editing. **Ariel E. A. Denos:** Data Curation, Formal Analysis, Supervision, Validation, Visualization, Writing – Original Draft, Writing – Review & Editing. **Jared M. Elison:** Formal Analysis, Investigation, Validation, Visualization, Writing – Original Draft, Writing – Review & Editing. **Benjamin S. Jones:** Formal Analysis, Investigation, Validation, Writing - Review & Editing. **Ethan G. Smith:** Formal Analysis, Investigation, Validation. **Noah G. Moran:** Investigation, Validation, Writing - Review & Editing. **Katie L. Brown** Investigation, Validation, Writing - Review & Editing. **Gerome M. Romero:** Data Curation, Formal Analysis, Investigation. **Chad D. Hyer:** Formal Analysis, Writing - Original Draft. **Kimberly B. Wagstaff:** Investigation, Validation. **Haifa M. Almughamsi:** Methodology, Investigation. **Mark K. Transtrum:** Methodology, Writing – Review & Editing. **John C. Price:** Conceptualization, Funding Acquisition, Methodology, Project Administration, Resources, Supervision, Writing – Original Draft, Writing - Review & Editing.

Funding Sources

This work was made possible by a grant from the Fritz B. Burns Foundation to John C. Price; the National Institutes of Health [R01AG066874] to John C. Price; Brigham Young University

Undergraduate Research Awards to Chad Hyer, Noah Earls, Noah Moran, Ethan Smith, Katie

Brown, Kimberly Wagstaff, Jared Elison, and Benjamin Jones; A grant from Deanship of

Scientific Research, Taif University to Haifa Almughamsi.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT:

The mice used in this research were provided by Johnathan J. Wisco, Ph. D. We gratefully

acknowledge the services provided by BYU's live animal facility and the Fritz B. Burns Biological

Mass Spectrometry Facility.

ABBREVIATIONS:

AD (Alzheimer's Disease), Aβ (Amyloidβ), ApoE (Apolipoprotein E), APP (Amyloid Precursor

Protein), BCA (Bicinchoninic Acid), BH-PV (Benjamini-Hochberg P-value), CID (Collision

induced dissociation), DDA (Data dependent acquisition), E2/E3/E4 (ApoE2/ApoE3/ApoE4),

ETC (Electron Transport Chain), FC (Fold Change), FDR (False discovery rate), GFAP (Glial

fibrillary acidic protein), IACUC (Institutional Animal Care and Use Committee), IP

(Intraperitoneal), LDL (Low density lipoprotein), LFQ (Label free quantification), MS (Mass-

Spectrometry), OXPHOS (Oxidative Phosphorylation), PDH (Pyruvate dehydrogenase), TCA

(Tricarboxylic Acid), TIM (mitochondrial inner membrane translocase), TOM (mitochondrial

outer membrane translocase).

REFERENCES:

1.   Naylor, B. C.; Porter, M. T.; Wilson, E.; Herring, A.; Lofthouse, S.; Hannemann, A.;
Piccolo, S. R.; Rockwood, A. L.; Price, J. C., DeuteRater: a tool for quantifying peptide isotope
precision and kinetic proteomics. *Bioinformatics* **2017,** *33* (10), 1514-1520.
2.   Liu, C. C.; Kanekiyo, T.; Xu, H.; Bu, G. J., Apolipoprotein E and Alzheimer disease:
risk, mechanisms and therapy (vol 9, pg 106, 2013). *Nat Rev Neurol* **2013,** *9* (4), 184-184.
3.   Husain, M. A.; Laurent, B.; Plourde, M., APOE and Alzheimer's Disease: From Lipid
Transport to Physiopathology and Therapeutics. *Front Neurosci* **2021,** *15*, 630502.
4.   Feussner, G.; Albanese, M.; Valencia, A., Three-dimensional structure of the LDL
receptor-binding domain of the human apolipoprotein E2 (Arg136-->Cys) variant.
Atherosclerosis **1996,** *126* (2), 177-84.
5.   Mahley, R. W., Central Nervous System Lipoproteins: ApoE and Regulation of
Cholesterol Metabolism. *Arterioscler Thromb Vasc Biol* **2016,** *36* (7), 1305-15.
6.   Farrer, L. A.; Cupples, L. A.; Haines, J. L.; Hyman, B.; Kukull, W. A.; Mayeux, R.;
Myers, R. H.; Pericak-Vance, M. A.; Risch, N.; van Duijn, C. M., Effects of Age, Sex, and
Ethnicity on the Association Between Apolipoprotein E Genotype and Alzheimer Disease: A
Meta-analysis. *JAMA* **1997,** *278* (16), 1349-1356.
7.   Breslow, J. L.; Zannis, V. I.; Sangiacomo, T. R.; Third, J. L. H. C.; Tracy, T.; Glueck,
C. J., Studies of Familial Type-Iii Hyperlipoproteinemia Using as a Genetic-Marker the Apoe
Phenotype E2/2. *J Lipid Res* **1982,** *23* (8), 1224-1235.
8.   Huang, X. M.; Chen, P. C.; Poole, C., APOE-epsilon 2 allele associated with higher
prevalence of sporadic Parkinson disease. *Neurology* **2004,** *62* (12), 2198-2202.
9.   Phillips, M. C., Apolipoprotein E Isoforms and Lipoprotein Metabolism. *Iubmb Life*
2014, *66* (9), 616-623.
10.   Kuo, C. L.; Pilling, L. C.; Atkins, J. L.; Masoli, J. A. H.; Delgado, J.; Kuchel, G. A.;
Melzer, D., APOE e4 Genotype Predicts Severe COVID-19 in the UK Biobank Community
Cohort. *J Gerontol a-Biol* **2020,** *75* (11), 2231-2232.
11.   Tcw, J.; Qian, L.; Pipalia, N. H.; Chao, M. J.; Liang, S. A.; Shi, Y.; Jain, B. R.;
Bertelsen, S. E.; Kapoor, M.; Marcora, E.; Sikora, E.; Andrews, E. J.; Martini, A. C.; Karch,
C. M.; Head, E.; Holtzman, D. M.; Zhang, B.; Wang, M.; Maxfield, F. R.; Poon, W. W.;
Goate, A. M., Cholesterol and matrisome pathways dysregulated in astrocytes and microglia.
Cell **2022,** *185* (13), 2213-2233.e25.
12.   Theendakara, V.; Peters-Libeu, C. A.; Spilman, P.; Poksay, K. S.; Bredesen, D. E.;
Rao, R. V., Direct Transcriptional Effects of Apolipoprotein E. *J Neurosci* **2016,** *36* (3), 685-
700.
13.   Szklarczyk, D.; Gable, A. L.; Lyon, D.; Junge, A.; Wyder, S.; Huerta-Cepas, J.;
Simonovic, M.; Doncheva, N. T.; Morris, J. H.; Bork, P.; Jensen, L. J.; Mering, C. V.,
STRING v11: protein-protein association networks with increased coverage, supporting
functional discovery in genome-wide experimental datasets. *Nucleic Acids Res* **2019,** *47* (D1),
D607-D613.
14.   Andersen, J. S.; Lam, Y. W.; Leung, A. K.; Ong, S. E.; Lyon, C. E.; Lamond, A. I.;
Mann, M., Nucleolar proteome dynamics. *Nature* **2005,** *433* (7021), 77-83.
15.   Kuechler, E. R.; Budzynska, P. M.; Bernardini, J. P.; Gsponer, J.; Mayor, T., Distinct
Features of Stress Granule Proteins Predict Localization in Membraneless Organelles. *J Mol Biol*
2020, *432* (7), 2349-2368.

16.   Buccitelli, C.; Selbach, M., mRNAs, proteins and the emerging principles of gene
expression control. *Nat Rev Genet* **2020,** *21* (10), 630-644.
17.   Martin-Perez, M.; Villen, J., Determinants and Regulation of Protein Turnover in Yeast.
Cell Syst **2017,** *5* (3), 283-294 e5.
18.   Schwanhausser, B.; Busse, D.; Li, N.; Dittmar, G.; Schuchhardt, J.; Wolf, J.; Chen,
W.; Selbach, M., Global quantification of mammalian gene expression control. *Nature* **2011,** *473*
(7347), 337-42.
19.   Mathis, A. D.; Naylor, B. C.; Carson, R. H.; Evans, E.; Harwell, J.; Knecht, J.;
Hexem, E.; Peelor, F. F., 3rd; Miller, B. F.; Hamilton, K. L.; Transtrum, M. K.; Bikman, B.
T.; Price, J. C., Mechanisms of In Vivo Ribosome Maintenance Change in Response to Nutrient
Signals. *Mol Cell Proteomics* **2017,** *16* (2), 243-254.
20.   Price, J. C.; Guan, S.; Burlingame, A.; Prusiner, S. B.; Ghaemmaghami, S., Analysis of
proteome dynamics in the mouse brain. *Proc Natl Acad Sci U S A* **2010,** *107* (32), 14508-13.
21.   McShane, E.; Sin, C.; Zauber, H.; Wells, J. N.; Donnelly, N.; Wang, X.; Hou, J.;
Chen, W.; Storchova, Z.; Marsh, J. A.; Valleriani, A.; Selbach, M., Kinetic Analysis of Protein
Stability Reveals Age-Dependent Degradation. *Cell* **2016,** *167* (3), 803-815 e21.
22.   Meadow, M. E.; Broas, S.; Hoare, M.; Alimohammadi, F.; Welle, K. A.; Swovick, K.;
Hryhorenko, J. R.; Martinez, J. C.; Biashad, S. A.; Seluanov, A.; Gorbunova, V.; Buchwalter,
A.; Ghaemmaghami, S., Proteome Birthdating Reveals Age-Selectivity of Protein
Ubiquitination. *Mol Cell Proteomics* **2024,** *23* (7), 100791.
23.   Wang, F.; Durfee, L. A.; Huibregtse, J. M., A cotranslational ubiquitination pathway for
quality control of misfolded proteins. *Mol Cell* **2013,** *50* (3), 368-78.
24.   Waterlow, J. C.; Waterlow, J. C., *Protein turnover*. CABI Pub.: Wallingford, UK ;
Cambridge, MA, 2006; p x, 301 p.
25.   Iizuka, R.; Yamagishi-Shirasaki, M.; Funatsu, T., Kinetic study of de novo chromophore
maturation of fluorescent proteins. *Analytical Biochemistry* **2011,** *414* (2), 173-178.
26.   Beynon, R. J., The dynamics of the proteome: strategies for measuring protein turnover
on a proteome-wide scale. *Briefings in Functional Genomics* **2005,** *3* (4), 382-390.
27.   Dörrbaum, A. R.; Kochen, L.; Langer, J. D.; Schuman, E. M., Local and global
influences on protein turnover in neurons and glia. *Elife* **2018,** *7*.
28.   Dhondt, I.; Petyuk, V. A.; Bauer, S.; Brewer, H. M.; Smith, R. D.; Depuydt, G.;
Braeckman, B. P., Changes of Protein Turnover in Aging Caenorhabditis elegans. *Molecular and*
Cellular Proteomics **2017,** *16* (9), 1621-1633.
29.   Visscher, M.; De Henau, S.; Wildschut, Mattheus H. E.; van Es, Robert M.; Dhondt, I.;
Michels, H.; Kemmeren, P.; Nollen, Ellen A.; Braeckman, Bart P.; Burgering, Boudewijn M.
T.; Vos, Harmjan R.; Dansen, Tobias B., Proteome-wide Changes in Protein Turnover Rates in
C. elegans Models of Longevity and Age-Related Disease. *Cell Reports* **2016,** *16* (11), 3041-
3051.
30.   Jonathon, J. O. B.; Vikram, N.; Yao, W.; Phillip, S.; Celeste, M. S.; Nicole, H.;
Megan, S.; Ramin, R.; Aleksandr, G.; Adam, B.; Matthew, K.; Baby, M.-M.; Chunlian, Z.;
Ganesh, K.; Carmela, S.; Vladimir, J.; Fiona, M.; Bryson, B.; Rochelle, B., Precise Estimation
of In Vivo Protein Turnover Rates. *bioRxiv* **2020**.
31.   Claydon, A. J.; Beynon, R., Proteome Dynamics: Revisiting Turnover with a Global
Perspective. *Mol Cell Proteomics* **2012,** *11* (12), 1551-1565.
32.   Ross, A. B.; Langer, J. D.; Jovanovic, M., Proteome Turnover in the Spotlight:
Approaches, Applications, and Perspectives. *Mol Cell Proteomics* **2021,** *20*.

33.   Miller, B. F.; Reid, J. J.; Price, J. C.; Lin, H. L.; Atherton, P. J.; Smith, K., CORP: The
use of deuterated water for the measurement of protein synthesis. *J Appl Physiol (1985)* **2020,**
128 (5), 1163-1176.
34.   Naylor, B. C.; Anderson, C. N. K.; Hadfield, M.; Parkinson, D. H.; Ahlstrom, A.;
Hannemann, A.; Quilling, C. R.; Cutler, K. J.; Denton, R. L.; Adamson, R.; Angel, T. E.;
Burlett, R. S.; Hafen, P. S.; Dallon, J. C.; Transtrum, M. K.; Hyldahl, R. D.; Price, J. C.,
Utilizing Nonequilibrium Isotope Enrichments to Dramatically Increase Turnover Measurement
Ranges in Single Biopsy Samples from Humans. *J Proteome Res* **2022,** *21* (11), 2703-2714.
35.   Black, E.; Rasch, A.; Wimmer, T.; Li, A.; Araujo, A.; Cieslak, S.; Steed, K. S.;
Adhikari, R. D.; Wisco, J. J.; Hutchinson, B. A., The effects of age, genotype and diet on
hippocampal subfield iron dysregulation and Alzheimer's disease biomarkers in an ApoE mouse
model. *European Journal of Neuroscience* **2023,** *57* (6), 1033-1047.
36.   Xu, J.; He, K.; Zhang, K.; Yang, C.; Nie, L.; Dan, D.; Liu, J.; Zhang, C. E.; Yang,
X., Low-Dose Copper Exposure Exacerbates Depression-Like Behavior in ApoE4 Transgenic
Mice. *Oxidative Medicine and Cellular Longevevity* **2021,** *2021*, 1-20.
37.   Cieslak, S. G., The effects of L-Cysteine on Alzheimer's disease pathology in APOE2,
APOE3, and APOE4 homozygous mice. *Journal of Biomedical Nanotechnology* **2020,** *5*.
38.   Chaudhari, K.; Wong, J. M.; Vann, P. H.; Como, T.; O'Bryant, S. E.; Sumien, N.,
ApoE Genotype-Dependent Response to Antioxidant and Exercise Interventions on Brain
Function. *Antioxidants (Basel)* **2020,** *9* (6).
39.   Adhikari, R.; Steed, K. S.; Hutchinson, B.; Wang, H.; Mendoza, M.; Staudte, R.;
Atmojo, M.; Cox, P.; Hancock, T.; Barkdull, K., Hippocampal T2 signal loss and decreased
radial arm maze performance in transgenic murine model for AD. *Brain Nerves* **2020,** *5*, 1-8.
40.   Pandey, R. S.; Graham, L.; Uyar, A.; Preuss, C.; Howell, G. R.; Carter, G. W., Genetic
perturbations of disease risk genes in mice capture transcriptomic signatures of late-onset
Alzheimer's disease. *Molecular Neurodegeneration* **2019,** *14* (1), 50.
41.   He, K.; Nie, L.; Zhou, Q.; Rahman, S. U.; Liu, J.; Yang, X.; Li, S., Proteomic Profiles
of the Early Mitochondrial Changes in APP/PS1 and ApoE4 Transgenic Mice Models of
Alzheimer's Disease. *Journal of Proteome Research* **2019,** *18* (6), 2632-2642.
42.   Meng, F.-T.; Zhao, J.; Fang, H.; Zhang, L.-F.; Wu, H.-M.; Liu, Y.-J., Upregulation of
Mineralocorticoid Receptor in the Hypothalamus Associated with a High Anxiety-like Level in
Apolipoprotein E4 Transgenic Mice. *Behavior Genetics* **2017,** *47* (4), 416-424.
43.   Jankowsky, J. L.; Zheng, H., Practical considerations for choosing a mouse model of
Alzheimer's disease. *Molecular Neurodegeneration* **2017,** *12* (1).
44.   Dorey, E.; Bamji-Mirza, M.; Najem, D.; Li, Y.; Liu, H.; Callaghan, D.; Walker, D.;
Lue, L.-F.; Stanimirovic, D.; Zhang, W., Apolipoprotein E Isoforms Differentially Regulate
Alzheimer's Disease and Amyloid-β-Induced Inflammatory Response in vivo and in vitro.
Journal of Alzheimer's Disease **2017,** *57* (4), 1265-1279.
45.   Chaudhari, K.; Wong, J. M.; Vann, P. H.; Sumien, N., Exercise, but not antioxidants,
reversed ApoE4-associated motor impairments in adult GFAP-ApoE mice. *Behavioural Brain*
Research **2016,** *305*, 37-45.
46.   Meng, F.-T.; Zhao, J.; Fang, H.; Liu, Y.-J., The influence of chronic stress on anxiety-
like behavior and cognitive function in different human GFAP-ApoE transgenic adult male mice.
Stress **2015,** *18* (4), 419-426.

47.   Fagan, A. M.; Watson, M.; Parsadanian, M.; Bales, K. R.; Paul, S. M.; Holtzman, D.
M., Human and murine ApoE markedly alters A beta metabolism before and after plaque
formation in a mouse model of Alzheimer's disease. *Neurobiol Dis* **2002,** *9* (3), 305-18.
48.   Fryer, J. D.; Simmons, K.; Parsadanian, M.; Bales, K. R.; Paul, S. M.; Sullivan, P. M.;
Holtzman, D. M., Human apolipoprotein E4 alters the amyloid-beta 40:42 ratio and promotes the
formation of cerebral amyloid angiopathy in an amyloid precursor protein transgenic model. *J*
Neurosci **2005,** *25* (11), 2803-10.
49.   Liao, F.; Zhang, T. J.; Jiang, H.; Lefton, K. B.; Robinson, G. O.; Vassar, R.; Sullivan,
P. M.; Holtzman, D. M., Murine versus human apolipoprotein E4: differential facilitation of and
co-localization in cerebral amyloid angiopathy and amyloid plaques in APP transgenic mouse
models. *Acta Neuropathol Commun* **2015,** *3*, 70.
50.   Aguilan, J. T.; Kulej, K.; Sidoli, S., Guide for protein fold change and p-value
calculation for non-experts in proteomics. *Molecular Omics* **2020,** *16* (6), 573-582.
51.   Pedregosa, F.; Varoquaux, G.; Gramfort, A.; Michel, V.; Thirion, B.; Grisel, O.;
Blondel, M.; Prettenhofer, P.; Weiss, R.; Dubourg, V.; Vanderplas, J.; Passos, A.;
Cournapeau, D.; Brucher, M.; Perrot, M.; Duchesnay, E., Scikit-learn: Machine Learning in
Python. *J Mach Learn Res* **2011,** *12*, 2825-2830.
52.   Virtanen, P.; Gommers, R.; Oliphant, T. E.; Haberland, M.; Reddy, T.; Cournapeau,
D.; Burovski, E.; Peterson, P.; Weckesser, W.; Bright, J.; van der Walt, S. J.; Brett, M.;
Wilson, J.; Millman, K. J.; Mayorov, N.; Nelson, A. R. J.; Jones, E.; Kern, R.; Larson, E.;
Carey, C. J.; Polat, İ.; Feng, Y.; Moore, E. W.; VanderPlas, J.; Laxalde, D.; Perktold, J.;
Cimrman, R.; Henriksen, I.; Quintero, E. A.; Harris, C. R.; Archibald, A. M.; Ribeiro, A. H.;
Pedregosa, F.; van Mulbregt, P.; Vijaykumar, A.; Bardelli, A. P.; Rothberg, A.; Hilboll, A.;
Kloeckner, A.; Scopatz, A.; Lee, A.; Rokem, A.; Woods, C. N.; Fulton, C.; Masson, C.;
Häggström, C.; Fitzgerald, C.; Nicholson, D. A.; Hagen, D. R.; Pasechnik, D. V.; Olivetti, E.;
Martin, E.; Wieser, E.; Silva, F.; Lenders, F.; Wilhelm, F.; Young, G.; Price, G. A.; Ingold,
G.-L.; Allen, G. E.; Lee, G. R.; Audren, H.; Probst, I.; Dietrich, J. P.; Silterra, J.; Webber, J.
T.; Slavič, J.; Nothman, J.; Buchner, J.; Kulick, J.; Schönberger, J. L.; de Miranda Cardoso,
J. V.; Reimer, J.; Harrington, J.; Rodríguez, J. L. C.; Nunez-Iglesias, J.; Kuczynski, J.; Tritz,
K.; Thoma, M.; Newville, M.; Kümmerer, M.; Bolingbroke, M.; Tartre, M.; Pak, M.; Smith,
N. J.; Nowaczyk, N.; Shebanov, N.; Pavlyk, O.; Brodtkorb, P. A.; Lee, P.; McGibbon, R. T.;
Feldbauer, R.; Lewis, S.; Tygier, S.; Sievert, S.; Vigna, S.; Peterson, S.; More, S.; Pudlik, T.;
Oshima, T.; Pingel, T. J.; Robitaille, T. P.; Spura, T.; Jones, T. R.; Cera, T.; Leslie, T.; Zito,
T.; Krauss, T.; Upadhyay, U.; Halchenko, Y. O.; Vázquez-Baeza, Y.; SciPy, C., SciPy 1.0:
fundamental algorithms for scientific computing in Python. *Nature Methods* **2020,** *17* (3), 261-
272.
53.   van den Berg, R. A.; Hoefsloot, H. C.; Westerhuis, J. A.; Smilde, A. K.; van der Werf,
M. J., Centering, scaling, and transformations: improving the biological information content of
metabolomics data. *BMC Genomics* **2006,** *7*.
54.   Pino, L. K.; Just, S. C.; MacCoss, M. J.; Searle, B. C., Acquiring and Analyzing Data
Independent Acquisition Proteomics Experiments without Spectrum Libraries. *Mol Cell*
Proteomics **2020,** *19* (7), 1088-1103.
55.   Price, J. C.; Khambatta, C. F.; Li, K. W.; Bruss, M. D.; Shankaran, M.; Dalidd, M.;
Floreani, N. A.; Roberts, L. S.; Turner, S. M.; Holmes, W. E.; Hellerstein, M. K., The effect of
long term calorie restriction on in vivo hepatic proteostatis: a novel combination of dynamic and
quantitative proteomics. *Mol Cell Proteomics* **2012,** *11* (12), 1801-14.

56. Price, J. C.; Holmes, W. E.; Li, K. W.; Floreani, N. A.; Neese, R. A.; Turner, S. M.;
Hellerstein, M. K., Measurement of human plasma proteome dynamics with (2)H(2)O and liquid
chromatography tandem mass spectrometry. *Anal Biochem* **2012,** *420* (1), 73-83.
57. Carson, R. H.; Lewis, C. R.; Erickson, M. N.; Zagieboylo, A. P.; Naylor, B. C.; Li, K.
W.; Farnsworth, P. B.; Price, J. C., Imaging regiospecific lipid turnover in mouse brain with
desorption electrospray ionization mass spectrometry. *J Lipid Res* **2017,** *58* (9), 1884-1892.
58. Chou, C. J.; Affolter, M.; Kussmann, M., A Nutrigenomics View of Protein Intake:
Macronutrient, Bioactive Peptides, and Protein Turnover. In *Progress in Molecular Biology and*
Translational Science, Bouchard, C.; Ordovas, J. M., Eds. Academic Press: 2012; Vol. 108, pp
51-74.
59. Ji, Z. S.; Miranda, R. D.; Newhouse, Y. M.; Weisgraber, K. H.; Huang, Y.; Mahley, R.
W., Apolipoprotein E4 potentiates amyloid beta peptide-induced lysosomal leakage and
apoptosis in neuronal cells. *J Biol Chem* **2002,** *277* (24), 21821-8.
60. Hook, V.; Yoon, M.; Mosier, C.; Ito, G.; Podvin, S.; Head, B. P.; Rissman, R.;
O'Donoghue, A. J.; Hook, G., Cathepsin B in neurodegeneration of Alzheimer's disease,
traumatic brain injury, and related brain disorders. *Biochim Biophys Acta Proteins Proteom*
2020, *1868* (8), 140428.
61. Schmukler, E.; Solomon, S.; Simonovitch, S.; Goldshmit, Y.; Wolfson, E.;
Michaelson, D. M.; Pinkas-Kramarski, R., Altered mitochondrial dynamics and function in
APOE4-expressing astrocytes. *Cell Death Dis* **2020,** *11* (7), 578.
62. Qi, G. Y.; Mi, Y. S.; Shi, X. J.; Gu, H. W.; Brinton, R. D.; Yin, F., ApoE4 Impairs
Neuron-Astrocyte Coupling of Fatty Acid Metabolism. *Cell Rep* **2021,** *34* (1).
63. Area-Gomez, E.; Larrea, D.; Pera, M.; Agrawal, R. R.; Guilfoyle, D. N.; Pirhaji, L.;
Shannon, K.; Arain, H. A.; Ashok, A.; Chen, Q. Y.; Dillman, A. A.; Figueroa, H. Y.;
Cookson, M. R.; Gross, S. S.; Fraenkel, E.; Duff, K. E.; Nuriel, T., APOE4 is Associated with
Differential Regional Vulnerability to Bioenergetic Deficits in Aged APOE Mice. *Sci Rep-Uk*
2020, *10* (1).
64. Fernández-Fernández, M. R.; Gragera, M.; Ochoa-Ibarrola, L.; Quintana-Gallardo, L.;
Valpuesta, J. M., Hsp70 - a master regulator in protein degradation. *FEBS Letters* **2017,** *591*
(17), 2648-2660.
65. Lin, Y. T.; Seo, J.; Gao, F.; Feldman, H. M.; Wen, H. L.; Penney, J.; Cam, H. P.;
Gjoneska, E.; Raja, W. K.; Cheng, J.; Rueda, R.; Kritskiy, O.; Abdurrob, F.; Peng, Z.; Milo,
B.; Yu, C. J.; Elmsaouri, S.; Dey, D.; Ko, T.; Yankner, B. A.; Tsai, L. H., APOE4 Causes
Widespread Molecular and Cellular Alterations Associated with Alzheimer's Disease Phenotypes
in Human iPSC-Derived Brain Cell Types. *Neuron* **2018,** *98* (6), 1141-1154.e7.
66. Lumsden, A. L.; Mulugeta, A.; Zhou, A.; Hypponen, E., Apolipoprotein E (APOE)
genotype-associated disease risks: a phenome-wide, registry-based, case-control study utilising
the UK Biobank. *EBioMedicine* **2020,** *59*, 102954.
67. Bonet-Costa, V.; Pomatto, L. C.-D.; Davies, K. J. A., The Proteasome and Oxidative
Stress in Alzheimer's Disease. *Antioxidants & Redox Signaling* **2016,** *25* (16), 886-901.
68. Fernandez-Fernandez, M. R.; Gragera, M.; Ochoa-Ibarrola, L.; Quintana-Gallardo, L.;
Valpuesta, J. M., Hsp70 - a master regulator in protein degradation. *FEBS Lett* **2017,** *591* (17),
2648-2660.
69. Paradise, V.; Sabu, M.; Bafia, J.; Sharif, N. A.; Nguyen, C.; Konrad-Vicario, K. D.;
Mukim, R. D.; Wang, X.; Corjuc, B. T.; Fu, J.; Maldonado, G.; Ndubisi, J.; Strickland, M.;
Figueroa, H.; Almeida, D.; Hyman, B.; Holtzman, D. M.; Nuriel, T.; Ramachandran, K. V.,

Dysregulation of neuroproteasomes by ApoE isoforms drives endogenous Tau aggregation. *bioRxiv* **2023**, 2022.11.29.518293.

70.   Chen, Y.; Durakoglugil, M. S.; Xian, X. D.; Herz, J., ApoE4 reduces glutamate receptor function and synaptic plasticity by selectively impairing ApoE receptor recycling. *P Natl Acad Sci USA* **2010,** *107* (26), 12011-12016.

71.   Dumanis, S. B.; DiBattista, A. M.; Miessau, M.; Moussa, C. E.; Rebeck, G. W., APOE genotype affects the pre-synaptic compartment of glutamatergic nerve terminals. *J Neurochem* **2013,** *124* (1), 4-14.

72.   Hunsberger, H. C.; Pinky, P. D.; Smith, W.; Suppiramaniam, V.; Reed, M. N., The role of APOE4 in Alzheimer's disease: strategies for future therapeutic interventions. *Neuronal Signal* **2019,** *3* (2), NS20180203.

73.   Zhong, N.; Ramaswamy, G.; Weisgraber, K. H., Apolipoprotein E4 Domain Interaction Induces Endoplasmic Reticulum Stress and Impairs Astrocyte Function*. *J Biol Chem* **2009,** *284* (40), 27273-27280.

74.   Marcotti, S.; Sánchez-Sánchez, B. J.; Serna-Morales, E.; Dragu, A.; Díaz-de-la-Loza, M.-d.-C.; Matsubayashi, Y.; Stramer, B. M., Protocol for intervention-free quantification of protein turnover rate by steady-state modeling. *STAR Protocols* **2021,** *2* (1).

75.   Xian, X.; Pohlkamp, T.; Durakoglugil, M. S.; Wong, C. H.; Beck, J. K.; Lane-Donovan, C.; Plattner, F.; Herz, J., Reversal of ApoE4-induced recycling block as a novel prevention approach for Alzheimer's disease. *Elife* **2018,** *7*.

76.   Nuriel, T.; Peng, K. Y.; Ashok, A.; Dillman, A. A.; Figueroa, H. Y.; Apuzzo, J.; Ambat, J.; Levy, E.; Cookson, M. R.; Mathews, P. M.; Duff, K. E., The Endosomal–Lysosomal Pathway Is Dysregulated by APOE4 Expression in Vivo. *Frontiers in Neuroscience* **2017,** *11*.

77.   Heeren, J.; Grewal, T.; Laatsch, A.; Becker, N.; Rinninger, F.; Rye, K. A.; Beisiegel, U., Impaired recycling of apolipoprotein E4 is associated with intracellular cholesterol accumulation. *J Biol Chem* **2004,** *279* (53), 55483-55492.

78.   Xian, X.; Pohlkamp, T.; Durakoglugil, M. S.; Wong, C. H.; Beck, J. K.; Lane-Donovan, C.; Plattner, F.; Herz, J., Reversal of ApoE4-induced recycling block as a novel prevention approach for Alzheimer's disease. *Elife* **2018,** *7*, e40048.

79.   Bonifacino, J. S.; Glick, B. S., The mechanisms of vesicle budding and fusion. *Cell* **2004,** *116* (2), 153-166.

80.   Lin, R. C.; Scheller, R. H., Mechanisms of synaptic vesicle exocytosis. *Annual Review of Cell and Developmental Biology* **2000,** *16*, 19-49.

81.   Narayan, P.; Sienski, G.; Bonner, J. M.; Lin, Y. T.; Seo, J.; Baru, V.; Haque, A.; Milo, B.; Akay, L. A.; Graziosi, A.; Freyzon, Y.; Landgraf, D.; Hesse, W. R.; Valastyan, J.; Barrasa, M. I.; Tsai, L. H.; Lindquist, S., PICALM Rescues Endocytic Defects Caused by the Alzheimer's Disease Risk Factor APOE4. *Cell Reports* **2020,** *33* (1).

82.   Nuriel, T.; Peng, K. Y.; Ashok, A.; Dillman, A. A.; Figueroa, H. Y.; Apuzzo, J.; Ambat, J.; Levy, E.; Cookson, M. R.; Mathews, P. M.; Duff, K. E., The Endosomal-Lysosomal Pathway Is Dysregulated by APOE4 Expression in Vivo. *Frontiers in Neuroscience* **2017,** *11*, 702.

83.   Ando, K.; Nagaraj, S.; Küçükali, F.; de Fisenne, M.-A.; Kosa, A.-C.; Doeraene, E.; Lopez Gutierrez, L.; Brion, J.-P.; Leroy, K., PICALM and Alzheimer’s Disease: An Update and Perspectives. *Cells* **2022,** *11* (24).

84. Chen, Y.; Durakoglugil, M. S.; Xian, X. D.; Herz, J., ApoE4 reduces glutamate receptor
function and synaptic plasticity by selectively impairing ApoE receptor recycling. *Proceedings*
of the National Academy of Sciences **2010,** *107* (26), 12011-12016.
85. Yang, C.; Wang, X., Lysosome biogenesis: Regulation and functions. *Journal of Cell*
Biology **2021,** *220* (6).
86. de Leeuw, S. M.; Kirschner, A. W. T.; Lindner, K.; Rust, R.; Budny, V.; Wolski, W.
E.; Gavin, A.-C.; Nitsch, R. M.; Tackenberg, C., APOE2, E3, and E4 differentially modulate
cellular homeostasis, cholesterol metabolism, and inflammatory response in isogenic iPSC-
derived astrocytes. *Stem Cell Reports* **2022,** *17* (1), 110-126.
87. Giorgi, C.; Marchi, S.; Pinton, P., The machineries, regulation and cellular functions of
mitochondrial calcium. *Nature Reviews Molecular Cell Biology* **2018,** *19* (11), 713-730.
88. Calvo-Rodriguez, M.; Bacskai, B. J., Mitochondria and Calcium in Alzheimer's Disease:
From Cell Signaling to Neuronal Cell Death. *Trends in Neurosciences* **2021,** *44* (2), 136-151.
89. Mahley, R. W., Apolipoprotein E4 targets mitochondria and the mitochondria-associated
membrane complex in neuropathology, including Alzheimer's disease. *Current Opinion in*
Neurobiology **2023,** *79*, 102684.
90. Rath, S.; Sharma, R.; Gupta, R.; Ast, T.; Chan, C.; Durham, T. J.; Goodman, R. P.;
Grabarek, Z.; Haas, M. E.; Hung, W. H. W.; Joshi, P. R.; Jourdain, A. A.; Kim, S. H.;
Kotrys, A. V.; Lam, S. S.; McCoy, J. G.; Meisel, J. D.; Miranda, M.; Panda, A.; Patgiri, A.;
Rogers, R.; Sadre, S.; Shah, H.; Skinner, O. S.; To, T.-L.; Walker, Melissa A.; Wang, H.;
Ward, P. S.; Wengrod, J.; Yuan, C.-C.; Calvo, S. E.; Mootha, V. K., MitoCarta3.0: an updated
mitochondrial proteome now with sub-organelle localization and pathway annotations. *Nucleic*
Acids Research **2021,** *49* (D1), D1541-D1547.
91. Mick, David U.; Dennerlein, S.; Wiese, H.; Reinhold, R.; Pacheu-Grau, D.; Lorenzi,
I.; Sasarman, F.; Weraarpachai, W.; Shoubridge, Eric A.; Warscheid, B.; Rehling, P.,
MITRAC Links Mitochondrial Protein Translocation to Respiratory-Chain Assembly and
Translational Regulation. *Cell* **2012,** *151* (7), 1528-1541.
92. Aiyar, R. S.; Bohnert, M.; Duvezin-Caubet, S.; Voisset, C.; Gagneur, J.; Fritsch, E. S.;
Couplan, E.; von der Malsburg, K.; Funaya, C.; Soubigou, F.; Courtin, F.; Suresh, S.;
Kucharczyk, R.; Evrard, J.; Antony, C.; St.Onge, R. P.; Blondel, M.; di Rago, J.-P.; van der
Laan, M.; Steinmetz, L. M., Mitochondrial protein sorting as a therapeutic target for ATP
synthase disorders. *Nature Communications* **2014,** *5* (1).
93. Anand, R.; Reichert, A. S.; Kondadi, A. K., Emerging Roles of the MICOS Complex in
Cristae Dynamics and Biogenesis. *Biology* **2021,** *10* (7).
94. van der Laan, M.; Horvath, S. E.; Pfanner, N., Mitochondrial contact site and cristae
organizing system. *Current Opinion in Cell Biology* **2016,** *41*, 33-42.
95. Xinglong, W.; Bo, S.; Hyoung-gon, L.; Xinyi, L.; George, P.; Mark, A. S.; Xiongwei,
Z., Impaired Balance of Mitochondrial Fission and Fusion in Alzheimer's Disease. *The*
Journal of Neuroscience **2009,** *29* (28), 9090-9103.
96. Schon, E. A.; Area-Gomez, E., Mitochondria-associated ER membranes in Alzheimer
disease. *Molecular and Cellular Neuroscience* **2013,** *55*, 26-36.
97. Area-Gomez, E.; Del Carmen Lara Castillo, M.; Tambini, M. D.; Guardia-Laguarta, C.;
de Groof, A. J.; Madra, M.; Ikenouchi, J.; Umeda, M.; Bird, T. D.; Sturley, S. L.; Schon, E.
A., Upregulated function of mitochondria-associated ER membranes in Alzheimer disease. *Embo*
Journal **2012,** *31* (21), 4106-4123.

98.   Schon, Eric A.; Przedborski, S., Mitochondria: The Next (Neurode)Generation. *Neuron* **2011,** *70* (6), 1033-1053.

99.   Nakamura, T.; Watanabe, A.; Fujino, T.; Hosono, T.; Michikawa, M., Apolipoprotein E4 (1–272) fragment is associated with mitochondrial proteins and affects mitochondrial function in neuronal cells. *Molecular Neurodegeneration* **2009,** *4* (1), 35.

100.   Williams, H. C.; Farmer, B. C.; Piron, M. A.; Walsh, A. E.; Bruntz, R. C.; Gentry, M. S.; Sun, R. C.; Johnson, L. A., APOE alters glucose flux through central carbon pathways in astrocytes. *Neurobiology of Disease* **2020,** *136*.

101.   Orr, A. L.; Kim, C.; Jimenez-Morales, D.; Newton, B. W.; Johnson, J. R.; Krogan, N. J.; Swaney, D. L.; Mahley, R. W., Neuronal Apolipoprotein E4 Expression Results in Proteome-Wide Alterations and Compromises Bioenergetic Capacity by Disrupting Mitochondrial Function. *Journal of Alzheimer's Disease* **2019,** *68*, 991-1011.

102.   Brandon, C. F.; Holden, C. W.; Nicholas, D.; Margaret, A. P.; Grant, K. N.; David, J. C.; Adeline, E. W.; Rebika, K.; Lyndsay, E. A. Y.; Jude, C. K.; Gabriela, H.; Elizabeth, J. A.; Rachel, M.; Brandon, J. A.; Vedant, A. G.; Philip, A. K.; Matthew, S. G.; Josh, M. M.; Ramon, C. S.; Lance, A. J., APOE4 Lowers Energy Expenditure and Impairs Glucose Oxidation by Increasing Flux through Aerobic Glycolysis. *Molecular Neurodegeneration* **2020,** *16*.

103.   Zhang, X.; Wu, L.; Swerdlow, R. H.; Zhao, L., Opposing Effects of ApoE2 and ApoE4 on Glycolytic Metabolism in Neuronal Aging Supports a Warburg Neuroprotective Cascade against Alzheimer's Disease. *Cells* **2023,** *12* (3).

104.   Valla, J.; Yaari, R.; Wolf, A. B.; Kusne, Y.; Beach, T. G.; Roher, A. E.; Corneveaux, J. J.; Huentelman, M. J.; Caselli, R. J.; Reiman, E. M., Reduced Posterior Cingulate Mitochondrial Activity in Expired Young Adult Carriers of the APOE ε4 Allele, the Major Late-Onset Alzheimer's Susceptibility Gene. *Journal of Alzheimer's Disease* **2010,** *22* (1), 307-313.

105.   Long, W.; Xin, Z.; Liqin, Z., Human ApoE Isoforms Differentially Modulate Brain Glucose and Ketone Body Metabolism: Implications for Alzheimer's Disease Risk Reduction and Early Intervention. *Journal of Neuroscience* **2018,** *38* (30), 6665-6681.

106.   Small, G. W.; Ercoli, L. M.; Silverman, D. H. S.; Huang, S. C.; Komo, S.; Bookheimer, S. Y.; Lavretsky, H.; Miller, K.; Siddarth, P.; Rasgon, N. L.; Mazziotta, J. C.; Saxena, S.; Wu, H. M.; Mega, M. S.; Cummings, J. L.; Saunders, A. M.; Pericak-Vance, M. A.; Roses, A. D.; Barrio, J. R.; Phelps, M. E., Cerebral metabolic and cognitive decline in persons at genetic risk for Alzheimer's disease. *Proceedings of the National Academy of Sciences* **2000,** *97* (11), 6037-6042.

107.   Wu, L.; Zhang, X.; Zhao, L., Human ApoE Isoforms Differentially Modulate Brain Glucose and Ketone Body Metabolism: Implications for Alzheimer's Disease Risk Reduction and Early Intervention. *Journal of Neuroscience* **2018,** *38* (30), 6665-6681.

108.   Lee, H.; Cho, S.; Kim, M.-J.; Park, Y. J.; Cho, E.; Jo, Y. S.; Kim, Y.-S.; Lee, J. Y.; Thoudam, T.; Woo, S.-H.; Lee, S.-I.; Jeon, J.; Lee, Y.-S.; Suh, B.-C.; Yoon, J. H.; Go, Y.; Lee, I.-K.; Seo, J., ApoE4-dependent lysosomal cholesterol accumulation impairs mitochondrial homeostasis and oxidative phosphorylation in human astrocytes. *Cell Reports* **2023,** *42* (10), 113183.

109.   Chin, D.; Hagl, S.; Hoehn, A.; Huebbe, P.; Pallauf, K.; Grune, T.; Frank, J.; Eckert, G. P.; Rimbach, G., Adenosine triphosphate concentrations are higher in the brain of APOE3-compared to APOE4-targeted replacement mice and can be modulated by curcumin. *Genes & Nutrition* **2014,** *9* (3).

110.   Yin, J.; Nielsen, M.; Carcione, T.; Li, S.; Shi, J., Apolipoprotein E regulates mitochondrial function through the PGC-1α-sirtuin 3 pathway. *Aging (Albany NY)* **2019,** *11* (23), 11148-11156.

111.   Fang, W.; Xiao, N.; Zeng, G.; Bi, D.; Dai, X.; Mi, X.; Ye, Q.; Chen, X.; Zhang, J., APOE4 genotype exacerbates the depression-like behavior of mice during aging through ATP decline. *Translational Psychiatry* **2021,** *11* (1).

112.   Mishra, R.; Upadhyay, A.; Prajapati, V. K.; Mishra, A., Proteasome-mediated proteostasis: Novel medicinal and pharmacological strategies for diseases. *Medicinal Research Reviews* **2018,** *38* (6), 1916-1973.

113.   Song, J. Y.; Wang, X. G.; Zhang, Z. Y.; Che, L.; Fan, B.; Li, G. Y., Endoplasmic reticulum stress and the protein degradation system in ophthalmic diseases. *Peerj* **2020,** *8*.

114.   Hou, X.; Zhang, X.; Zou, H.; Guan, M.; Fu, C.; Wang, W.; Zhang, Z.-R.; Geng, Y.; Chen, Y., Differential and substrate-specific inhibition of γ-secretase by the C-terminal region of ApoE2, ApoE3, and ApoE4. *Neuron* **2023,** *111* (12), 1898-1913.e5.

115.   Thibaudeau, T. A.; Anderson, R. T.; Smith, D. M., A common mechanism of proteasome impairment by neurodegenerative disease-associated oligomers. *Nature Communications* **2018,** *9* (1), 1097.

116.   Tseng, B. P.; Green, K. N.; Chan, J. L.; Blurton-Jones, M.; LaFerla, F. M., Aβ inhibits the proteasome and enhances amyloid and tau accumulation. *Neurobiology of Aging* **2008,** *29* (11), 1607-1618.

117.   Kodroń, A.; Mussulini, B. H.; Pilecka, I.; Chacińska, A., The ubiquitin-proteasome system and its crosstalk with mitochondria as therapeutic targets in medicine. *Pharmacological Research* **2021,** *163*.

118.   Wang, X.; Yen, J.; Kaiser, P.; Huang, L., Regulation of the 26S Proteasome Complex During Oxidative Stress. *Science Signaling* **2010,** *3* (151), ra88.

119.   Ross, J. M.; Olson, L.; Coppotelli, G. Mitochondrial and Ubiquitin Proteasome System Dysfunction in Ageing and Disease: Two Sides of the Same Coin? *International Journal of Molecular Sciences* [Online], 2015, p. 19458-19476.

120.   Paradise, V.; Sabu, M.; Bafia, J.; Sharif, N. A.; Nguyen, C.; Konrad-Vicario, K. D.; Mukim, R. D.; Wang, X.; Corjuc, B. T.; Fu, J.; Maldonado, G.; Ndubisi, J.; Strickland, M.; Figueroa, H.; Almeida, D.; Hyman, B.; Holtzman, D. M.; Nuriel, T.; Ramachandran, K. V., Dysregulation of neuroproteasomes by ApoE isoforms drives endogenous Tau aggregation. *bioRxiv* **2023**.

121.   Sun, F.; Kanthasamy, A.; Anantharam, V.; Kanthasamy, A. G., Mitochondrial accumulation of polyubiquitinated proteins and differential regulation of apoptosis by polyubiquitination sites Lys-48 and -63. *Journal of Cellular and Molecular Medicine* **2009,** *13* (8b), 1632-1643.

122.   Dambacher, C. M.; Worden, E. J.; Herzik, M. A., Jr.; Martin, A.; Lander, G. C., Atomic structure of the 26S proteasome lid reveals the mechanism of deubiquitinase inhibition. *Elife* **2016,** *5*.

123.   Fote, G. M.; Geller, N. R.; Efstathiou, N. E.; Hendricks, N.; Vavvas, D. G.; Reidling, J. C.; Thompson, L. M.; Steffan, J. S., Isoform-dependent lysosomal degradation and internalization of apolipoprotein E requires autophagy proteins. *Journal of Cell Science* **2022,** *135* (2).

124.   Perez-Riverol, Y.; Csordas, A.; Bai, J.; Bernal-Llinares, M.; Hewapathirana, S.; Kundu, D. J.; Inuganti, A.; Griss, J.; Mayer, G.; Eisenacher, M.; Pérez, E.; Uszkoreit, J.;

1256   Pfeuffer, J.; Sachsenberg, T.; Yilmaz, S.; Tiwary, S.; Cox, J.; Audain, E.; Walzer, M.;
1257   Jarnuczak, A. F.; Ternent, T.; Brazma, A.; Vizcaíno, J. A., The PRIDE database and related
1258   tools and resources in 2019: improving support for quantification data. *Nucleic Acids Research*
1259   **2019,** *47* (D1), D442-D450.

1260

1261   For Table of Contents Only

1262