Stable isotope composition of Antarctic marine mammals collected at the turn of the 20th century: assessing the baseline for long-term changes in the marine ecosystem.

Evgeny Genelt-Yanovskiy[1], Anna Genelt-Yanovskaya[1], Maria Fontanals-Coll[2], Kweku Afrifa Yamoah[2], Oliver Craig[2], Richard Sabin[3], James Scourse[1]

[1]Department of Earth and Environmental Sciences, University of Exeter, Penryn Campus, Cornwall, TR10 9FE UK.
[2]University of York, BioArCh, Environment Building 2nd Floor, Wentworth Way, Heslington, York, YO10 5DD, UK.
[3]The Natural History Museum, Cromwell Road, London SW7 5BD, UK.

ORCID IDs
EGY 0000-0001-7115-2181; AGY 0000-0002-6050-2569; MFC 0000-0002-4308-0907; KAY 0000-0001-7488-0459; OC 0000-0002-4296-8402; RS 0000-0003-0699-7596; JS 0000-0003-2658-8730

Corresponding authors: e.genelt-yanovskiy@exeter.ac.uk, j.scourse@exeter.ac.uk

Abstract

Great Antarctic expeditions, and subsequently developed seal hunting and whaling industries left an enormous legacy in natural history museum collections. Stable isotopes of carbon δ^{13}C and nitrogen δ^{15}N from specimens preserved in the collections can provide valuable information on long-term changes in foraging ecology of species and specify the baselines for the future comparisons with modern samples. To provide the basis for analysing the impact of whaling on marine ecosystem structuring, we conducted the bulk isotope analysis from the specimens of baleen whales (*Balaenoptera musculus* and *B. physalus*), beaked whale *(Berardius arnuxii)*, and seals *(Arctocephalus australis* and *Hydrurga leptonyx*) collected between 1843 and 1951, preserved in the collection of Natural History Museum, London. Having controlled for the Suess effect, δ^{13}C values in *B. musculus*, *B. physalus* and *H. leptonyx* were different before and after the onset of industrial whaling (1904). Bone collagen δ^{15}N values and corresponding trophic position consistently changed in *B. musculus* and *A. australis,* suggesting potential temporal shifts in their diets. This study highlights the use of museum specimens for tracing the historical trends that could be associated with changes in the population structure and distribution of species and indicate long-term variability in their foraging ecology.

Keywords

Antarctic, cetacean, pinniped, bone collagen, museum collections, carbon, nitrogen, stable isotopes, Suess effect, trophic position, foraging ecology

Introduction

The Southern Ocean is highly productive (1), and the waters surrounding the Antarctic Peninsula characterized by high krill abundance are usually considered one of the most

productive areas of the Antarctic (2,3). High productivity is also typical of the Subantarctic islands including South Georgia located beyond the northern extent of sea ice (4), as well as the Patagonian shelf and southern tip of South America due to northward branches of the high nutrient Antarctic Circumpolar Current (5). High densities of krill, *Euphasia superba* in the offshore areas and *E. crystallorophias* in the coastal waters around Antarctica attract huge numbers of seabirds (penguins and Procellariiformes) and marine mammals, including pinnipeds and cetaceans (6,7).

Antarctic marine ecosystems are currently experiencing rapid change, especially around the Antarctic Peninsula, identified as one of the most rapidly warming regions of the Southern Hemisphere (8,9). The Antarctic seas were effectively pristine prior to discovery of the continent in the eighteenth century followed by the development of seal hunting and subsequently whaling which resulted in a severe decline in commercially important species. Intensive seal hunting brought Antarctic fur seals *Arctocephalus gazella* almost to extinction by the late 1700-early 1800s in South Georgia, and the whaling industry peaked in the early 1900s when the permanent whaling station was opened at Grytviken (4,10). Following international moratoria on industrial whaling activities, whale populations have been recovering over recent decades after a long period of decline, but recent census studies indicate that their abundance has still not attained pre-whaling numbers (11,12).

The significant decline in the abundance of marine mammals in the late 19[th] – early 20[th] centuries caused by industrial-scale exploitation may well have had a significant impact on the marine ecosystem of Antarctica, as well as on the structure and distribution of populations of the species affected. Intensive whaling and seal hunting in Antarctica coincided with the period known as "Heroic Age of Antarctic exploration", a series of naval and land expeditions between the late 19th century and 1920s. This period left an enormous legacy in natural history museum collections around the world, including the specimens of species targeted by industrial whaling and sealing. Museum specimens are increasingly important for gaining ecological and biological data, and as carriers of geochemical and molecular information, that are important for studying rare, elusive and even extinct species. Historical museum "specimens of opportunity" are often challenging to use due to uncertainties in time and location of collections, and in the quality of post-collection curation, but similarly to the subfossil specimens from excavations of archaeological or paleontological sites can provide snapshots of past genetic diversity, spatial origin and diet (13–15). Stable isotope analysis, most commonly the δ^{13}C carbon and δ^{15}N nitrogen isotopes of marine species, can indicate latitudinal range shifts, migrations between nearshore and offshore environments as well as changes in trophic structure (16).

Historical specimens of Antarctic whales from various collections have become increasingly important in isotopic and genetic studies, being used to describe niche partitioning between fin (*Balaenoptera physalus)* and sei (*B. borealis*) whales (17), and changes in mitochondrial DNA diversity in blue (*B. musculus)*, humpback (*Megaptera novaeangliae*) and fin whales (18). Isotopic analyses of collections representing Antarctic fur seal (*A. gazella*) in Antarctica (19) and South American fur seal (*A. australis*) in Patagonia (20) indicate long-term latitudinal and longitudinal shifts in their prey distributions. In this study, we measured δ^{13}C and δ^{15}N compositions of the oldest Antarctic and Subantarctic marine mammal bone specimens of opportunity preserved in the Natural History Museum in London (NHMUK). These specimens represent the species most targeted by sealing and whaling activity and thus collected during the expeditions prior to and during the development of industrial scale harvesting. Following

the timeline and southward trend of exploration of Subantarctic and Antarctic waters, the oldest specimens in the dataset correspond to the period of survey of the Strait of Magellan by the HMS *Alert* Expedition (1876–1884) and the James Clark Ross Expedition onboard HMS *Erebus* and HMS *Terror* (1838-1843). The most recent specimens deposited in the NHM were collected by the *Southern Harvester* in the 1960s, the last UK whaling vessel operating in Antarctic and Subantarctic waters; these provide a snapshot from the later phases of 20[th] century whaling activity. By focusing primarily on well-studied whale species, particularly blue whale *B. musculus* and fin whale *B. physalus*, as well as two species of seals – leopard seal *Hydrurga leptonyx* and South American fur seal *A. australis* - we explore the following research questions: (i) How have δ^{13}C and δ^{15}N stable isotopes varied over the period of intensive whaling; (ii) Do δ^{13}C values show any historical trends that could be associated with changes in the distribution of species; and (iii) Can the observed trends in δ^{15}N indicate long-term changes in trophic position of the species. We also analysed δ^{13}C and δ^{15}N bulk isotopic signatures of several specimens of Arnoux's beaked whale *Berardius arnuxii* from the same period. While this species has rarely been exploited, this study provides a trophic snapshot of early 20[th] century populations of this elusive Antarctic species.

Materials and methods

Sampling

Samples were collected at the mammal collection of Natural History Museum (London) using handheld cutting tools and a Dremel tool with either dental bur or carbide cutting disk. Before sampling, the surface of the target fragment of the bone was cleaned using Dremel with the bristle polishing brush bit. In total, 42 samples of bone fragments were analysed (Table 1).

Table 1. Bone material and number of specimens used in the study

Species	n	Years of collections	Bone fragments sampled
Balaenoptera musculus	11	1879 - 1951	Periotic; ear bone; lumbar vertebra; shoulder blade; pelvic
Balaenoptera physalis	4	1912 - 1951	Periotic; ear bone; pelvic
Berardius arnuxii	5	1922 - 1931	Rib; sternum; skull
Arctocephalus australis	10	1876 - 1951	Skull
Hydrurga leptonyx	12	1839 - 1963	Skull

The initial weight of the bone samples was about 500 mg. Collagen was extracted and prepared for stable isotope analysis at the BioArCh laboratories in the Department of Archaeology, University of York (UK). The bone samples (approx. 200 to 300 mg) were demineralized using 0.6 M HCl at 4°C for several weeks, then rinsed with ultrapure H_2O (milli-Q) and gelatinized with 0.001 M HCl at 80°C for 48 h. The supernatant containing the collagen was filtered using polyethylene Ezee filters (Elkay Laboratories, 9 ml, pore size 60–90 μm) and then 30 kDa Amicon Ultra-4 centrifugal filter units (Millipore, MA, USA). Samples were then frozen for 24–48 hours at −20°C, lyophilized, and weighed into tin capsules (0.5 mg per duplicate) for stable isotope analysis.

Figure 1. Locations of samples of marine mammals analysed in the study. Map redrawn from Google Earth. Detailed description of locations and NHM accession numbers are available in Table 1 Supplementary material 2.

Bulk EA-IRMS analysis

The BioArCh laboratories at the University of York's Department of Archaeology conducted duplicate measurements of stable carbon and nitrogen isotope ratios. Using a Sercon EA-GSL elemental analyzer linked to a Sercon 20-22 continuous flow isotope ratio mass spectrometer (Sercon, Crewe, UK), the ratios of $^{13}C:^{12}C$ and $^{15}N:^{14}N$ were assessed relative to standards (V-PDB for carbon and AIR for nitrogen) and reported in delta notation (δ) in parts per mil (‰). Stable carbon and nitrogen isotope ratios were measured using a Sercon HS2022 continuous flow isotope ratio mass spectrometer at BioArCh, University of York (UK).

Stable carbon and nitrogen isotope ratios were calibrated relative to VPDB and AIR scales using the standard reference materials IsoAnalytical Cane ($\delta^{13}C$ = -11.64‰ ± 0.03), Sigma Methionine ($\delta^{13}C$ = -35.83‰ ± 0.03, $\delta^{15}N$ = -0.76‰ ± 0.05) and IAEA N2 ($\delta^{15}N$ = 20.41‰ ± 0.12). Uncertainty was monitored using the standard reference materials Sigma fish gel ($\delta13C$ = -15.27‰ ± 0.04, $\delta15N$ = 15.21‰ ± 0.13), IsoAnalytical Alanine ($\delta^{13}C$ = -23.33‰ ± 0.10, $\delta^{15}N$ = -5.56‰ ± 0.14) and IsoAnalytical Soy ($\delta^{13}C$ = -25.22‰ ± 0.03, $\delta^{15}N$ = 0.99‰ ± 0.07). Precision (u(Rw)) was determined to be ± 0.165‰ for $\delta^{13}C$ and ± 0.124‰ for $\delta^{15}N$ based on repeated measurement of calibration standards, check standards and sample replicates. According to calibration and check standards (Table 1 Supplementary material 1), measurement precision (ssrm, pooled measured standard deviation of calibration and check standards) was ± 0.151‰ for $\delta^{13}C$ (df = 28) and ± 0.124‰ for $\delta^{15}N$ (df = 24). All the samples (42/42) were analysed in duplicate (Table 2 Supplementary material 1). The measurement precision specific to samples (srep, pooled standard deviation of sample replicates) was ± 0.094‰ for $\delta^{13}C$ (df = 47) and ± 0.050‰ for $\delta^{15}N$ (df = 47). Accuracy (u(bias)) was determined to be ± 0.091‰ for $\delta^{13}C$ and ± 0.159‰ for $\delta^{15}N$ based on the difference between observed and reported δ values (RMSbias) and long-term standard deviations (u(Cref)) of check standards. Total analytical standard uncertainty (uc) was estimated to be ± 0.188‰ for $\delta^{13}C$ and ± 0.204‰ for $\delta^{15}N$ (21).

 Data analysis
Basic descriptive statistics to explore variation for both $\delta^{13}C$ and $\delta^{15}N$ within the bone samples
were performed using R (R Core Team, 2024) with RStudio (RStudio Team, 2024). Raw $\delta^{13}C$
values were corrected to consider the oceanic Suess effect – the depletion in the $\delta^{13}C$ isotopic
composition of dissolved inorganic carbon (DIC) in oceans due to the increase in
anthropogenic CO_2 released into the atmosphere since the Industrial Revolution (Keeling,
1979; Misarti et al., 2009; Keeling et al., 2017). Calculations were performed using the
following equation introduced by Hilton et al. (2006), modified by Misarti et al. (2009) and
later presented within the SuessR package (22):
175
176 Suess effect correction factor = *a* * exp(*b**0.027)
177
Where *a* is a constant reflecting the maximum observed rate of $\delta^{13}C$ decline in surface waters
DIC in a specific region, *b* is the year of sample collection minus 1850 (taken as the
environmental onset of the Industrial Revolution) and 0.027 is the parameter value obtained
by Hilton et al. (2006) after fitting an exponential curve to the global ocean $\delta^{13}C$ data from
1945 to 1997 published by Gruber et al. (1999). The values of total accumulation of Suess
effect for the constant *a* were obtained from Eide et al. (2017) for the surface waters (0 meters
depth layer in a database available under the doi:10.1594/PANGAEA.872004). These values
were obtained and averaged for five regions (Table 2). Per-year ratios of *a* were obtained by
dividing these values by 140, i.e., the number of years between the year of 1850 (Industrial
Revolution) and 1990 (start of the main decade of collecting data in Eide et al.,2017).
188
189 **Table 2.** Oceanic Suess effect correction values extracted from doi:10.1594/PANGAEA.872004
190 (see Eide et al., 2017 for details). N – number of point estimates of total accumulated Suess
191 effect in surface (0 meter) water layer. Total CSE – mean value of total Oceanic Suess effect for
192 each region. The a-value – expected per-year $\delta^{13}C$ decline (‰).
193

	Latitude range		Longitude range		n	Total CSE	a-value
Patagonia / Strait of Magellan	-49	-55	-77	-67	19	-0.91	-0.0065
Falkland Islands	-50.8	-52.5	-62	-57.4	2	-0.93	-0.0066
South Georgia	-53	-56	-38.7	-33.9	13	-0.73	-0.0052
South Shetlands	-61	-63	-63	-57	9	-0.75	-0.0054
Ross Sea	-71	-77	171	-160	173	-0.71	-0.0051

194
195 Bone $\delta^{15}N$ signatures were converted to trophic position (TP) using the following equation
196 (23,24):
197
198 TP = 2 + ($\delta^{15}N_{sample}$ − $\delta^{15}N_{POM}$)/TEF,
199
200 where Dm is $\delta^{15}N$ value in marine mammal bone, POM is $\delta^{15}N$ value of marine particulate
201 organic matter, and TEF is trophic enrichment factor in $\delta^{15}N$ for marine mammals (23,25,26).
202 Data on POM were obtained from previously published modern datasets (8,27–30).
203
204 Consumer and source isotope data were analysed using the MixSIAR package in R (31–33),
205 where source data were composed of mean and standard deviation (SD) from potential prey
206 sources. The consumer data per blue whale and South American fur seal were divided into

207 three consecutive datasets, each representing a period of collection, and each dataset was
208 treated as having no covariates, i.e. including all individuals without separation by their IDs.
209
210 Table 3. Stable isotope signatures of potential prey items for the blue whale and South
211 American fur seal, mined from the literature (see text for references) and used for the MixSIAR
212 modeling.

Prey group / Species	Foraging area	n	δ13C		δ15N	
			mean	SE	mean	SE
Euphasia superba	South Shetland Islands	24	-26.87	0.94	3.63	0.38
Euphasia superba	South Georgia	26	-23.87	1.22	4.27	0.54
Thysanoessa spp./ Euphasia frigida	Polar front	7	-23.25	1.05	5.10	0.20
Euphasia superba	Antarctic shelf waters	43	-31.20	0.10	2.85	0.75
Euphasia lucens	Subantarctic (SE Patagonian shelf waters)	7	-19.80	0.70	7.30	0.80
Squid (*Doryteuthis gahi*)		10	-19.00	0.60	13.60	0.70
Demersal fish (*Patagonotothen spp.*)		5	-19.30	0.60	14.80	1.00
Small planktonic fish (*Sprattus fuegensis*)		25	-20.20	1.00	13.10	0.90
Piscivorous fish (Gadidae)		6	-14.90	0.30	17.20	0.40

213
For both species, source isotopic data were obtained based on recent dietary studies of the
species concerned and spatial variability of isotopic signatures of potential prey items (34–37)
(Table 3). The models ran three Markov Chain Monte Carlo (MCMC) chains with 50000
iterations, a burn in of 25000, and were thinned by 25 ('short' MCMC run parameters).
MixSIAR reports results as mean, SD and credible intervals for each posterior density
distribution per prey source.
220

Results

Stable isotope data

The raw $\delta^{13}C$ and $\delta^{15}N$ values varied significantly in two of three whale species studied (Figure
2, Table 4). The $\delta^{13}C$ values varied significantly in blue whale specimens between -23.67 to -
13.23, and similarly $\delta^{15}N$ values varied between 5.26 and 17.05. In fin whale, $\delta^{13}C$ values
ranged between -21.66 to -15.43, and $\delta^{15}N$ values ranged between 6.13 to 11.64. In Arnoux's
beaked whale $\delta^{13}C$ values were less variable, with a narrow range between -19.62 to -18.88,
and $\delta^{15}N$ values between 11.55 and 13.94.
229

230 The raw $\delta^{13}C$ values in South American fur seal demonstrated a narrow range between -
231 12.97‰ to -11.42‰, while raw $\delta^{15}N$ values varied between 16.73‰ to 23.56‰. In leopard
232 seal, $\delta^{13}C$ values showed more pronounced variation (-22.02 to -15.8), compared to variation
233 in $\delta^{15}N$ between 7.45 and 11.27. The Suess-corrected $\delta^{13}C$ values demonstrated the same
234 trends and did not deviate more than 0.12 ‰ from raw values (Figure 2, Table 4).
235

236 The complete set of individual $\delta^{13}C$ and $\delta^{15}N$ values, and detailed information about the origin
237 of each specimen analysed in this study is presented in Table 1 Supplementary material 2.
238

239 **Table 4.** Mean and range stable isotope values from the bone collagen values of marine
240 mammals collected in Subantarctic waters between the 1880s and 1960s.

species	n	Raw δ¹³C ± SE	Suess-corrected δ¹³C	δ¹⁵N ± SE	C:N mass ratio
Balaenoptera musculus blue whale	11	-19.86 ± 1.00	-19.88 ± 1.00	7.73 ± 1.07	3.58 ± 0.11
Balaenoptera physalus fin whale	4	-19.01 ± 0.70	-19.13 ± 1.32	8.36 ± 0.58	3.55 ± 0.10
Berardius arnuxii Arnoux's beaked whale	5	-19.29 ± 0.15	-19.34 ± 0.15	12.46 ± 0.53	3.33 ± 0.03
Arctocephalus australis South American fur seal	10	- 12.31 ± 0.22	-12.33 ± 0.15	19.37 ± 0.49	3.33 ± 0.02
Hydrurga leptonyx leopard seal	12	-19.74 ± 0.50	- 19.76 ± 0.50	10.12 ± 0.30	3.44 ± 0.04

Carbon to nitrogen (C:N) atomic ratio ranged from 3.24 to 4.3 in various species. So, while
most of the samples fell within the expected range (from 2.9 to 3.6) for unaltered bone
samples (DeNiro, 1985), several specimens demonstrated C:N values outside this range,
particularly three blue whale specimens collected in 1914 near South Georgia (3.72, 4.17, 4.3),
one fin whale also collected in 1914 near South Georgia (3.85), and one leopard seal collected
near the Falklands before 1885 (3.84) (Table 1 Supplementary Material 2).

248
249

250
Figure 2. δ¹³C (Suess corrected) and δ¹⁵N biplot in the bone of marine mammals sampled in
Subantarctic waters between the 1880s and 1960s. Solid line ellipses correspond to a
multivariate t-distribution, dashed line ellipses correspond to a multivariate normal
distribution.

pre- and post-1904 stable isotopic values
In total, 20 specimens pre-dating the opening of the whaling station at South Georgia were
analyzed, including three specimens of blue whale, six specimens of South American fur seal,
and 10 specimens of leopard seal (Table 5). The mean raw δ¹³C value in blue whale over this

period was -15.86 ± 2.2, and mean δ15N value 12.3 ± 2.42 (range 9.07 – 17.05), South
American fur seal δ13C -12.26 ± 0.21 and δ15N 20.32 ± 0.7, and leopard seal -19.71 ± 0.59 and
δ13C value 10.32 ± 0.21. Suess corrected δ13C values deviated by 0.003‰ and 0.11‰ with a
mean value of 0.029‰ across all samples and species studied (Table 5).

Table 5. Stable isotope ratios in bone collagen of NHM specimens pre-dating the opening of
whaling station at South Georgia.

NHMUK ID	Species	Location	Year*	Raw δ13C	Suess corrected δ13C	δ15N	C:N mass ratio
ZD.1880.7.28.19	*B.musculus*	Strait of Magellan*	March 1879	-20.29	-20.31	9.07	3.58
ZD.1879.8.21.9	*B.musculus*	Strait of Magellan	March 1879	-14.04	-14.06	10.79	3.49
ZD.1879.8.21.8	*B.musculus*	Strait of Magellan	February 1879	-13.23	-13.23	17.05	3.45
ZD.1852.3.11.5	*A.australis*	Falkland Islands	before 1850	-12.97	-12.98	21.13	3.34
ZD.1879.8.21.5	*A.australis*	Strait of Magellan	1878 to 1880	-12.08	-12.09	17.50	3.31
ZD.1880.7.28.12	*A.australis*	Strait of Magellan	1878 to 1880	-11.96	-11.97	19.29	3.35
ZD.1880.7.28.13	*A.australis*	Strait of Magellan	1878 to 1880	-11.42	-11.43	20.40	3.29
ZD.1880.7.28.14	*A.australis*	Patagonian west coast	1878 to 1880	-12.49	-12.51	23.56	3.34
ZE.1950.11.14.3	*A.australis*	Patagonian west coast	04/01/1876	-12.01	-12.02	19.84	3.30
ZE.1950.11.14.4	*A.australis*	Patagonian west coast	04/01/1876	-12.91	-12.92	20.48	3.48
ZD.1843.1.8.4	*H.leptonyx*	Antarctic Seas	1843	-22.02	-22.02	9.90	3.36
ZD.1846.4.15.23	*H.leptonyx*	Antarctic Seas	1843	-18.38	-18.39	10.85	3.34
ZD.1846.4.15.24	*H.leptonyx*	Antarctic Seas	1843	-21.08	-21.09	10.33	3.39
ZD.1876.1.28.81	*H.leptonyx*	Kerguelen	13/10/1874	-15.80	-15.80	9.11	3.32
ZD.1880.7.28.5	*H.leptonyx*	Falkland Islands	1878 to 1880	-19.00	-19.02	11.27	3.51
ZD.1885.10.20.1	*H.leptonyx*	Falkland Islands	before 1885	-18.55	-18.57	10.71	3.84
ZD.1893.9.14.1	*H.leptonyx*	South Georgia	before 1893	-19.39	-19.40	10.79	3.35
ZD.1908.2.20.54	*H.leptonyx*	Ross Sea	1901 to 1904	-21.20	-21.22	9.64	3.51
ZD.1908.2.20.56	*H.leptonyx*	Ross Sea	1901 to 1904	-20.56	-20.58	10.57	3.38
GERM.325.e	*H.leptonyx*	Antarctic Seas	1839 to 1843	-21.15	-21.15	9.97	3.42

*'Strait of Magellan' dataset consists of material from the Strait itself and surrounding
archipelagoes along the Chilean Pacific coastline / West coast of Patagonia (see Table 1
Supplementary material 2 for details).

Among the more recent specimens, the δ13C mean value for blue whale was -20.33 ± 0.51,
and mean δ15N value was 6.01 ± 0.31 (N = 8). In South American fur seal, the mean δ13C value

was -12.42 ± 0.16 and δ15N mean value 17.16 ± 0.23 (N=3). In leopard seal, the mean δ13C
value -was 19.85 ± 0.8 and δ15N 9.12 ± 1.67 (N = 2).

Differences were most pronounced in δ15N values for South American fur seal (PERMANOVA
F=14.61, p = 0.041) (Figure 3), yet no difference in δ13C values for this species was found
between the end of the 19th and early 20th centuries. Despite the large variability in δ13C values
in general, no significant changes in either δ13C or δ15N in leopard seals were found. Changes
in stable isotope signatures after 1904 in blue whale were significant for both carbon and
nitrogen with more pronounced changes in δ15N (PERMANOVA F=38.94, p=0.0011);
differences in δ13C were also significant (PERMANOVA F=12.73, p=0.0046).

284
285   **Figure 3.** Suess corrected δ13C and δ15N biplot summarizing within-species variability of
286   samples collected before (triangles) and after (squares) the opening of whaling station at
287   South Georgia in 1904.

288
289   *Estimation of trophic positions using δ15N values*

Among the baleen whale individuals, the calculated trophic position (TP) varied between 3.1 and 5.3 in fin whale and between 2.4 and 5.7 in individual blue whales, but averaged TP was 3.4±0.3 across all specimens (Figure 4A). In both fin and blue whale specimens pronounced spatial heterogeneity was found showing that TP was higher at the South Shetland Isles compared to South Georgia. In Arnoux's beaked whale TP varied between 2.9 to 7.3 with higher values at the South Shetlands and lower values at the Falklands.

South American fur seal TP varied from 5.1 to 8.4 with higher values in the Straits of Magellan and lower values in the Falkland Islands. Leopard seal TP varied between 2.6 and 6.3. Highest TP values in leopard seal were found in the Ross Sea, lowest values in specimens from the Falklands.

Figure 4. Averaged trophic position (**A,** solid-filled bars) and corresponding averaged bone collage δ^{15}N values (**A,** semi-transparent bars) of five marine mammal species in the various regions studied. Changes of trophic position estimates (**B**) for South American fur seal (*A. australis*), leopard seal (*H. leptonyx*), and blue whale (*B. musculus*) between the 1850s and 1960s.

Comparison of TP in different time periods (Figure 4B) of sampling using PERMANOVA analysis indicates significant TP differences in blue whale between the material from the 1913-1914 and 1925-1951 (PERMANOVA F=1.91, p=0.0367). Due to high within-sample variability, no differences between the 1879 samples and the subsequently collected material were revealed using PERMANOVA. The South American fur seal material from the 1870-1880s is significantly different from material collected during 1925-1951 (PERMANOVA F=8.7, p=0.0549). Contrasting with these two species, no significant temporal changes in TP of leopard seal between 1839 and 1963 is revealed (PERMANOVA F=1.37, p=0.0663).

Source prediction by MixSIAR

The Bayesian mixing modelling approach was applied to the datasets on blue whale and South American fur seal, because these species demonstrated significant changes in trophic position over the period between 1850s and 1950s. Based on stable isotope data of four species of krill from contrasting locations around study area, MixSIAR estimated that the major source of food of blue whales collected in the Magellan Strait in the late 19th-century originated in Subantarctic waters, matching the geographic origin of the bone specimens (Figure 5A). MixSIAR model predictions for specimens representing the two following periods (Figure 5B,C) were non-identical. While all these samples were collected in waters surrounding South Georgia, MixSIAR indicated primary foraging grounds matching the geographic origin of the specimens dated 1913-1914, and a mix of foraging areas in 1925-1951, including the Polar Front region.

Figure 5. Comparison of MixSIAR-estimated dietary source contributions (as Bayesian credibility intervals and posterior densities) for the blue whale *B. musculus* (A to C) and South American fur seal *A. australis* (D to F). Abbreviations indicate potential prey items: *E.s. – Euphasia superba, Th. spp – Thysanoessa spp., E.f. – Euphasia frigida, E.l. – Euphasia lucens, P.spp - Patagonotothen spp., Gad – Gadidae, Sprat - Sprattus fuegensis, Dor – Doryteuthis (Loligo) gahi.*

MixSIAR model predictions for South American fur seal considered only one foraging area, Subantarctic waters. All specimens from the 19th century indicated similar dominant prey sources – piscivorous fish belonging to Gadidae family (Figure 5D,E). Specimens dated

342   between 1924 and 1951 demonstrated a changed pattern of mixed dietary sources in which
343   each species included in the analysis contributed to diet nearly equally.
344

Discussion

The number of studies examining the impacts of industrial whaling during the 20th-century is
increasing (11,17,18,27,38). So, it is a priority to accumulate as much as possible data
representing the state of their populations prior to the onset of industrial-scale whaling.
Despite the several uncertainties in the dataset, particularly the origin and often the unclear
state of preservation of several specimens, this study provides a snapshot of stable isotope
signatures of common Antarctic and Subantarctic marine mammals during the period of
extensive exploration of Antarctic waters in the nineteenth century and at the turn of the
20th-century.
354

Methodological considerations of stable isotope analysis of specimens of opportunity
Museum specimens represent a unique source of information about past populations and rare
species (16), but in some cases can be difficult to handle if their provenance is unclear (39).
Our dataset covers the period between the 1840s and 1960s, and while the geographical
origin and period of collecting is known, and accurate to at least within a year for several
specimens, there are uncertainties associated with the either location or date of collection of
others. The highest level of such uncertainty in the dataset is linked to the *Discovery*
Investigations programme that ran between 1924 and 1951, including several specimens of
blue whale *B. musculus* and South American fur seal *A. australis*, the latter being collected
prior to 1949 (Table 1 Supplementary Material 2). For all the specimens with uncertain year
of collection, the known time interval of the expedition or research program was used in
comparative analyses.
367

The stable isotope analysis approach is based on the relationship that carbon and nitrogen
isotopes in the predator tissues reflect those of its prey in a predictable way (40). Factors such
as nutrient resources, composition of primary producers and regional geographic
characteristics result in distinct isotopic patterns between different regions (41), allowing the
use of stable isotope analysis for investigating large-scale movements of predators (41,42).
Stable isotope analysis has been useful in detecting niche overlap and trophic interactions
between different trophic levels (43). Implementation of δ^{13}C and δ^{15}N isotope analyses, as
well as ^{14}C radiocarbon dating of bone fragments, rely on the preservation of collagen in bone
samples that can be evaluated using carbon to nitrogen (C:N) atomic ratio (44).
377

378   Five specimens in our dataset demonstrated slight deviations from the commonly accepted
379   range of C:N atomic ratio between 2.9 and 3.6. Since these specimens passed the other quality
380   control indicators (Table 1, Supplementary Material 2), namely percentage of carbon higher
381   than 13% and percentage of nitrogen higher than 4% (45) and did not deviate strongly from
382   the other specimens from same locations and periods, they were retained in our dataset.
383

We used the correction of raw δ^{13}C values for the Suess effect, which is crucial if modern and
historic specimens are included in the analysis, especially if various species in the same region
demonstrate different trajectories of long-term changes of δ^{13}C values. A study of four species
of penguins from subantarctic French Southern Ocean and Antarctic territories (46)
demonstrated that Suess-corrected δ^{13}C values did not change over time in Antarctic species
from Adelie Land, but decreased in Subantarctic and subtropical species. However, Suess-

corrected δ^{13}C values of the Weddell seals from the Ross Sea decreased during the 20th
century (47), indicating changes in the primary productivity of this sector of the Antarctic. In
our dataset, the correction values did not alter the results based on raw δ^{13}C values but were
necessary for comparison of the carbon signatures among specimens from the same species
collected at various locations during different periods. The oceanic Suess effect was evaluated
to be weak near the Antarctic continent and higher in Subantarctic waters (48). Based on the
global database of the full oceanic Suess effect (49), the average correction value for our
dataset was 0.027‰, varying between 0.004‰ (for samples dated 1839) and 0.11‰ (for
samples dated 1963). When the date of specimen collection was unknown, we used the
earliest date within the period indicated.

Evaluation of dietary preferences is conventionally based on δ^{15}N isotope analysis. By
analysing the nitrogen signatures of potential prey items or most common taxa in the
ecosystem, it is possible to evaluate the trophic relationships between consumers of different
levels, even if stomach content data for the particular population or species are scarce (50–
54). Conventional bulk nitrogen isotope analysis is based on assumption that isotopic
composition of a consumer reflect the isotopic composition of primary producers at the
baseline of the food web (40). Since δ^{15}N values from bulk stable isotope analysis cannot be
compared directly among tissues derived from food webs in different areas (26,55), but the
comparisons are possible through calculating trophic position based on δ^{15}N values of lower
trophic levels, or particulate organic matter (POM) in aquatic ecosystems (8,30,56). In recent
decade, stable isotope analysis of specific compounds within tissues, such as individual amino
acids, became increasingly popular in archaeological, trophic and animal migration studies
(57,58), and when possible, comparisons of bulk and CSIA are being implemented (59).

Since specimens representing different periods were often collected at different locations (all
species except the South American fur seal), and also because the complexity of food webs in
Antarctica varies with latitude (60–62), we calculated the trophic position for all four species
in focus based on regional variation of the modern δ^{15}N values of POM from the
corresponding regions (See Materials and Methods for details). Despite the obvious
constraints mentioned above, this approach (Figure 4) allowed to provide more accurate
geographic comparisons of nitrogen signatures.

In addition, we applied Bayesian mixing models (implemented in MixSIAR package) to
evaluate the dietary shifts in two species - blue whale and South American fur seal. Mixing
models are extensively used to evaluate the percentage contribution of potential prey items
to the diet of consumers (32,63–68). This method is increasingly being used to reconstruct
dietary preferences of past populations from museum specimens using modern data on diet
and isotopic signatures of major prey items (51,54,69).

The geographic differences in isotopic signatures raise questions about population
connectivity and foraging strategies in highly mobile marine species. We discuss below how
the observed variation in carbon and nitrogen stable isotope signatures of common marine
mammal bone collagen samples can be linked with rapid changes in Antarctic and Subantarctic
ecosystems during the period of their acquisition by the NHM.

Spatial and temporal variation in stable isotope values
We found pronounced variation in bone collagen δ^{13}C and δ^{15}N values across the two species
of baleen whales (*B. musculus* and *B. physalus*) and in leopard seal (*H. leptonyx*). In South

American fur seal (*A. australis*) we observed temporal changes in $\delta^{15}N$ and no variation in $\delta^{13}C$
values. Only a minor variation in isotopic signatures of Arnoux's beaked whale (*B. arnuxii*) was
found. The observed intraspecific variation in isotopic signatures mostly corresponded to the
geographic origin of specimen collection, but also may potentially indicate temporal variation
in foraging strategies and distribution of highly mobile species.
444
The blue whale specimens demonstrated the largest within-species variation in both carbon
and nitrogen isotopes in our dataset. Among these specimens, highest variation was observed
among the 1870s data from the Magellan Strait. Two of three 1870s specimens demonstrated
$\delta^{13}C$ signatures significantly different from the more recent specimens, and all three deviated
from the post-1904 specimens in their $\delta^{15}N$ signatures. The averaged trophic position of blue
whale was similar in different regions based on the background of the pronounced geographic
variation in $\delta^{15}N$ values. The post-1904 isotope data correspond well with 1940s baleen plate
data of whales collected near the Ross Sea (56).
453
The potential explanation for the unexpectedly high individual TP value (5.7) obtained for the
specimen dated 1879 from Strait of Magellan could indicate that this whale does not belong
to the region where it was collected. We used POM values to calculate TP of the three blue
whales from the Strait of Magellan, but this approach could be an oversimplification for the
highly migratory animals like blue whales. Studies undertaken between the 1920s and 1950s
interpreted the Eastern South Pacific blue whales as Antarctic blue whales regularly migrating
between higher and lower latitudes. Following the discovery of nursery and feeding grounds
in Southern Chile (70), Eastern South Pacific and Antarctic populations of blue whales were
considered as different subspecies, but satellite tracking data indicate that blue whales from
Chile migrate as far north as the Galapagos Islands (70). Blue whales in the Southern
Hemisphere were hunted nearly to extinction during the 20th century, and the modern
population estimates are considered to be less than 1% of the pre-whaling period (71). Thus,
the modern within-species structure of blue whales in the Southern Hemisphere is likely to
be substantially different to the pre-whaling period. Recent genetic studies reveal that many
Antarctic blue whales demonstrate signatures of admixture with populations from higher
latitudes (38), providing a plausible basis for the interpretation of past connectivity between
these populations.
471
Krill (Euphasiacea) is of the primary prey sources of baleen whales, of which the Antarctic krill
Euphasia superba is considered the main prey item of Antarctic blue whales (72,73). Blue
whales seasonally forage in productive coastal areas, and separate populations rely on
different areas with contrasting oceanographic characteristics and krill species which can be
traced using stable isotope analysis (74). As the geographic origin of several of the studied
blue whale specimens are unknown, the Bayesian mixing models for the blue whale were
applied as source data using several of the most common krill species from various latitudes
from the Antarctic shelf to Subantarctic waters around South Patagonia and the Falkland
Islands (Figure 5). This approach indicates isotopic signatures matching the origin of
specimens for both 1870s specimens from Strait of Magellan and 1913-1914 specimens from
South Georgia. In contrast, trophic signatures of blue whales caught in the open sea during
the 1925-1950 period were mixed. The latter can either indicate mixed geographic origin of
the studied specimens that were later landed at South Georgia, but also can be explained by
a shift in the foraging areas of blue whales escaping the whaling regions.
486

Our dataset included four fin whale specimens collected between 1912 and 1951 at South
Georgia, the South Shetlands and in the South Atlantic. These specimens also demonstrated
a pronounced variation in carbon and nitrogen isotopes, and while these results can be
considered preliminary, they corroborate a recent baleen plate isotopic study concluding that
the blue and fin whale in the Southern Ocean occupy a similar trophic level but demonstrate
niche differentiation (56).
493
The δ^{13}C values of Arnoux's beaked whale did not vary between the specimens from the
Falkland Islands and South Shetland Islands. Though there is a minor variation in δ^{15}N values,
this dataset is not sufficient for evaluating diet variation. The isotopic signatures obtained in
our study are also similar to the beaked whale from Falkland Islands (37), and indicate a
narrow isotopic niche of beaked, at least during 1920s-1930s. While most sightings of this rare
species are from the fast-ice edges of the Antarctic seas and the nearshore of South Africa,
New Zealand, and Australia (75), our data confirm the previous hypotheses of the association
of this species with deep offshore marine areas.
502
Our data indicate changes in the diet of the South American fur seal from the Falkland Islands
against a background of consistent regional δ^{13}C values. Estimation of trophic position from
the δ^{15}N values reveal a significant change in the trophic preferences of fur seal at the Falkland
Islands from the 19th to the 20th century. The diet of South American fur seal includes both
coastal and open-sea shelf prey items (76). Currently the dominant prey items of fur seals at
the Falklands are Falkland herring *Sprattus fugensis*, and several other taxa including
Patagonian squid *Doryteuthis gahi*, and several rock cod species *Patagonotothen spp.* (77).
While piscivorous offshore fish species are rarely found in stomachs of fur seals (76,77), these
were included among other potential sources in our mixing model (Figure 5), primarily as an
outgroup, and because recent studies based on satellite tracking data indicated long-distance
foraging migrations in several colonies of South American fur seal (34). Calculation of trophic
position and illustration of dietary preferences using the mixing models both indicate that
South American fur seals could have had more selective feeding in the 19[th]-century, and more
opportunistic feeding in the 20[th]-century, possibly due to a decline in preferred prey species.
Seals from 1870s were presumably foraging for larger prey of higher trophic levels. As several
fish taxa in Patagonian waters have recently experienced overfishing (78–80), our data
indicate that observed changes in the diet of South American fur seal represent an adaptation
to changes in the pelagic ecosystem.
521
Leopard seal is a species capable of foraging on taxa at various trophic levels, including other
seal species, penguins, fish, and krill (81–83). Huge variation in dietary preferences should
affect the TP in different locations, seasons and even particular individuals. The highest TP in
leopard seal was typical of specimens from the Ross Sea (5.97 $\pm$ 0.19), with samples with the
lowest TP from the Falkland Islands (2.66 $\pm$ 0.16). TP of leopard seal can be explained by
contrasting foraging strategies of this species in these two regions. As with the other species,
we divided the TP data into three periods, but no significant differences between seals from
these periods were found because differences within each temporal subsets was higher that
between the three subsets (Figure 4). According to these data, TP of leopard seal was nearly
twice lower compared to the South American fur seal. This finding is remarkable considering
that fur seal pups are one of the common prey items for the leopard seal.
533

CONCLUSIONS

Bulk stable isotope analysis of museum specimens in this study provides a record of Subantarctic and Antarctic marine environments during the period of active exploration and exploitation of marine resources in the region at the turn of the 20th century. Here we show how this dataset can be used to address the questions about the influence of industrial whaling, seal hunting and even development of fishing on the foraging of whales and seals. This study discusses the possible constraints for using this dataset in the standard ecological analyses aimed at understanding the long-term changes in Antarctic marine ecosystem using several keystone species.

Our results suggest that the major foraging grounds of baleen whales could have been different before the extensive whaling started in 1900s from South Georgia and South Shetland Islands with more contacts between the populations from Southern Ocean and South Pacific. South American Fur seal population at the Falkland Islands could have had foraged for a prey of higher trophic levels before the onset of industrial fishing in the region, and the leopard seal demonstrated high degree of adaptation to the local availability of prey objects. Compared to the other species, Arnoux's beaked whale, a rare species that was not targeted by whaling industry, did not demonstrate pronounced variability in stable isotope signatures. These hypotheses, raised on the basis of conventional bulk stable isotope analysis, should be further verified on the basis of larger sample sizes for each species, comparison with modern data from Antarctic and other regions within the species ranges, and using more advanced methods such as compound-specific isotope analysis that can more accurately distinguish environmental and dietary signatures.

Funding

This study was implemented within the SEACHANGE project, funded by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation Programme (grant agreement No 856488). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

We thank C. Ullmann (University of Exeter) for the technical support of sampling at NHM, and Phaedra Kokkini (NHMUK) for the support during the work at NHM collection and transfer of samples for the subsequent analysis, and all the ERC SEACHANGE project collaborators for their useful comments during the preparation of the manuscript.

References

1. Pinkerton MH, Boyd PW, Deppeler S, Hayward A, Höfer J, Moreau S. Evidence for the Impact of Climate Change on Primary Producers in the Southern Ocean. Front Ecol Evol. 2021 May 17;9:592027.

2.   Mascioni M, Almandoz GO, Ekern L, Pan BJ, Vernet M. Microplanktonic diatom
assemblages dominated the primary production but not the biomass in an Antarctic fjord.
Journal of Marine Systems. 2021 Dec;224:103624.

3.   Vernet M, Martinson D, Iannuzzi R, Stammerjohn S, Kozlowski W, Sines K, et al. Primary
production within the sea-ice zone west of the Antarctic Peninsula: I—Sea ice, summer
mixed layer, and irradiance. Deep Sea Research Part II: Topical Studies in Oceanography.
2008 Sep;55(18–19):2068–85.

4.   Atkinson A, Whitehouse M, Priddle J, Cripps G, Ward P, Brandon M. South Georgia,
Antarctica: a productive, cold water, pelagic ecosystem. Mar Ecol Prog Ser. 2001;216:279–
308.

5.   Franco BC, Ruiz-Etcheverry LA, Marrari M, Piola AR, Matano RP. Climate Change Impacts
on the Patagonian Shelf Break Front. Geophysical Research Letters. 2022 Feb
28;49(4):e2021GL096513.

6.   Bestley S, Ropert-Coudert Y, Bengtson Nash S, Brooks CM, Cotté C, Dewar M, et al. Marine
Ecosystem Assessment for the Southern Ocean: Birds and Marine Mammals in a Changing
Climate. Front Ecol Evol. 2020 Nov 4;8:566936.

7.   Hammerschlag N, Williams L, Fallows M, Fallows C. Disappearance of white sharks leads
to the novel emergence of an allopatric apex predator, the sevengill shark. Sci Rep. 2019
Feb 13;9(1):1908.

8.   Seyboth E, Botta S, Mendes CRB, Negrete J, Dalla Rosa L, Secchi ER. Isotopic evidence of
the effect of warming on the northern Antarctic Peninsula ecosystem. Deep Sea Research
Part II: Topical Studies in Oceanography. 2018 Mar;149:218–28.

9.   Vaughan DG, Marshall GJ, Connolley WM, Parkinson C, Mulvaney R, Hodgson DA, et al.
Recent Rapid Regional Climate Warming on the Antarctic Peninsula. Climatic Change.
2003;60(3):243–74.

10. Headland R. The Island of South Georgia. CUP Archive; 1984.

11. Calderan S, Black A, Branch T, Collins M, Kelly N, Leaper R, et al. South Georgia blue whales
five decades after the end of whaling. Endang Species Res. 2020 Nov 19;43:359–73.

12. Hamabe K, Matsuoka K, Kitakado T. Estimation of abundance and population dynamics of
the Antarctic blue whale in the Antarctic Ocean south of 60°S, from 70°E to 170°W. Marine
Mammal Science. 2023 Apr;39(2):671–87.

13. Aaris-Sørensen K, Rasmussen KL, Kinze C, Petersen KS. Late Pleistocene and Holocene
whale remains (Cetacea) from Denmark and adjacent countries: Species, distribution,
chronology, and trace element concentrations. Marine Mammal Science. 2010
Apr;26(2):253–81.

14. De Bruyn M, Hall BL, Chauke LF, Baroni C, Koch PL, Hoelzel AR. Rapid Response of a Marine
Mammal Species to Holocene Climate and Habitat Change. Przeworski M, editor. PLoS
Genet. 2009 Jul 10;5(7):e1000554.

15. English PA, Green DJ, Nocera JJ. Stable Isotopes from Museum Specimens May Provide
Evidence of Long-Term Change in the Trophic Ecology of a Migratory Aerial Insectivore.
Front Ecol Evol. 2018 Feb 12;6:14.

16. Smith KJ, Trueman CN, France CAM, Sparks JP, Brownlow AC, Dähne M, et al. Stable
Isotope Analysis of Specimens of Opportunity Reveals Ocean-Scale Site Fidelity in an
Elusive Whale Species. Front Conserv Sci. 2021 May 25;2:653766.

17. Buss DL, Hearne E, Loy RHY, Manica A, O'Connell TC, Jackson JA. Evidence of resource
partitioning between fin and sei whales during the twentieth-century whaling period. Mar
Biol. 2022 Nov;169(11):150.

18. Sremba AL, Martin AR, Wilson P, Cypriano-Souza AL, Buss DL, Hart T, et al. Diversity of
mitochondrial DNA in 3 species of great whales before and after modern whaling. Alter E,
editor. Journal of Heredity. 2023 Nov 15;114(6):587–97.

19. De Lima R, Cebuhar J, Negrete J, Ferreira A, Secchi E, Botta S. Ecosystem shifts inferred
from long-term stable isotope analysis of male Antarctic fur seal *Arctocephalus gazella*
teeth. Mar Ecol Prog Ser. 2022 Aug 25;695:203–16.

20. Szteren D, Aurioles-Gamboa D, Labrada-Martagón V, Hernández-Camacho CJ, De María
M. Historical age-class diet changes in South American fur seals and sea lions in Uruguay.
Mar Biol. 2018 Apr;165(4):59.

21. Szpak P, Metcalfe JZ, Macdonald RA. Best practices for calibrating and reporting stable
isotope measurements in archaeology. Journal of Archaeological Science: Reports. 2017
Jun;13:609–16.

22. Clark CT, Cape MR, Shapley MD, Mueter FJ, Finney BP, Misarti N. SuessR: Regional
corrections for the effects of anthropogenic CO_2 on $\delta^{13}C$ data from marine organisms.
Methods Ecol Evol. 2021 Aug;12(8):1508–20.

23. Hobson KA, Welch HE. Determination of trophic relationships within a high Arctic marine
food web using $\delta^{13}C$ and $\delta^{15}N$ analysis. Marine ecology progress series. 1992;9–18.

24. Lesage V, Hammill MO, Kovacs KM. Marine mammals and the community structure of the
Estuary and Gulf of St Lawrence, Canada: evidence from stable isotope analysis. Marine
Ecology Progress Series. 2001;210:203–21.

25. Hobson KA, Piatt JF, Pitocchelli J. Using stable isotopes to determine seabird trophic
relationships. Journal of animal ecology. 1994;786–98.

26. Post DM. Using stable isotopes to estimate trophic position: models, methods, and
assumptions. Ecology. 2002;83(3):703–18.

27. Jackson J, Kennedy A, Moore M, Andriolo A, Bamford C, Calderan S, et al. Have whales
returned to a historical hotspot of industrial whaling? The pattern of southern right whale
Eubalaena australis recovery at South Georgia. Endang Species Res. 2020 Nov 5;43:323–
39.

28. Bruno DO, Riccialdelli L, Acha EM, Fernández DA. Seasonal variation of autochthonous and
allochthonous carbon sources for the first levels of the Beagle Channel food web. Journal
of Marine Systems. 2023 Mar;239:103859.

29. Büring T, Van Der Grient J, Pierce G, Bustamante P, Scotti M, Jones JB, et al. Unveiling the
wasp-waist structure of the Falkland shelf ecosystem: the role of *Doryteuthis gahi* as a
keystone species and its trophic influences. J Mar Biol Ass. 2024;104:e2.

30. Espinasse B, Pakhomov E, Hunt B, Bury S. Latitudinal gradient consistency in carbon and
nitrogen stable isotopes of particulate organic matter in the Southern Ocean. Mar Ecol
Prog Ser. 2019 Nov 21;631:19–30.

31. R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna,
Austria: R Foundation for Statistical Computing; 2024. Available from: https://www.R-
project.org/

32. Stock BC, Jackson AL, Ward EJ, Parnell AC, Phillips DL, Semmens BX. Analyzing mixing
systems using a new generation of Bayesian tracer mixing models. PeerJ. 2018;6:e5096.

33. Stock B, Semmens B. MixSIAR GUI user manual v3. 1. Scripps Institution of Oceanography,
UC San Diego, San Diego, California, USA. 2016;

34. Baylis A, Tierney M, Orben R, Staniland I, Brickle P. Geographic variation in the foraging
behaviour of South American fur seals. Mar Ecol Prog Ser. 2018 May 28;596:233–45.

35. Ciancio JE, Pascual MA, Botto F, Frere E, Iribarne O. Trophic relationships of exotic
anadromous salmonids in the southern Patagonian Shelf as inferred from stable isotopes.
Limnology & Oceanography. 2008 Mar;53(2):788–98.

36. Schmidt K, McClelland JW, Mente E, Montoya JP, Atkinson A, Voss M. Trophic-level
interpretation based on d15N values: implications of tissue-specific fractionation and
amino acid composition. Mar Ecol Prog Ser. 2004;266:43–58.

37. Riccialdelli L, Paso Viola M, Panarello H, Goodall R. Evaluating the isotopic niche of beaked
whales from the southwestern South Atlantic and Southern Oceans. Mar Ecol Prog Ser.
2017 Oct 13;581:183–98.

38. Attard CRM, Sandoval-Castillo J, Lang AR, Vernazzani BG, Torres LG, Baldwin R, et al. Global
conservation genomics of blue whales calls into question subspecies taxonomy and refines
knowledge of population structure. Animal Conservation. 2024 Mar 15;acv.12935.

39. Keighley X, Bro-Jørgensen MH, Ahlgren H, Szpak P, Ciucani MM, Sánchez Barreiro F, et al.
Predicting sample success for large-scale ancient DNA studies on marine mammals.
Molecular Ecology Resources. 2021 May;21(4):1149–66.

40. Deniro MJ, Epstein S. Influence of diet on the distribution of nitrogen isotopes in animals.
Geochimica et Cosmochimica Acta. 1981 Mar;45(3):341–51.

41. Graham S, Mendelssohn I. Multiple levels of nitrogen applied to an oligohaline marsh
identify a plant community response sequence to eutrophication. Mar Ecol Prog Ser. 2010
Nov 4;417:73–82.

42. McMahon KW, Hamady LL, Thorrold SR. A review of ecogeochemistry approaches to
estimating movements of marine animals. Limnology & Oceanography. 2013
Mar;58(2):697–714.

43. Franco-Trecu V, Aurioles-Gamboa D, Inchausti P. Individual trophic specialisation and niche
segregation explain the contrasting population trends of two sympatric otariids. Mar Biol.
2014 Mar;161(3):609–18.

44. Brock F, Wood R, Higham TFG, Ditchfield P, Bayliss A, Ramsey CB. Reliability of Nitrogen
Content (%N) and Carbon:Nitrogen Atomic Ratios (C:N) as Indicators of Collagen
Preservation Suitable for Radiocarbon Dating. Radiocarbon. 2012;54(3–4):879–86.

45. Guiry EJ, Szpak P. Quality control for modern bone collagen stable carbon and nitrogen
isotope measurements. Soto D, editor. Methods Ecol Evol. 2020 Sep;11(9):1049–60.

46. Jaeger A, Cherel Y. Isotopic Investigation of Contemporary and Historic Changes in Penguin
Trophic Niches and Carrying Capacity of the Southern Indian Ocean. Thrush S, editor. PLoS
ONE. 2011 Feb 2;6(2):e16484.

47. Hückstädt LA, McCarthy MD, Koch PL, Costa DP. What difference does a century make?
Shifts in the ecosystem structure of the Ross Sea, Antarctica, as evidenced from a sentinel
species, the Weddell seal. Proc R Soc B. 2017 Aug 30;284(1861):20170927.

48. McNeil BI, Matear RJ, Tilbrook B. Does carbon 13 track anthropogenic CO_2 in the Southern
Ocean? Global Biogeochemical Cycles. 2001 Sep;15(3):597–613.

49. Eide M, Olsen A, Ninnemann US, Eldevik T. A global estimate of the full oceanic ^{13}C Suess
effect since the preindustrial. Global Biogeochemical Cycles. 2017 Mar;31(3):492–514.

50. García-Seoane R, Viana IG, Bode A. Using MixSIAR to quantify mixed contributions of
primary producers from amino acid δ15N of marine consumers. Marine Environmental
Research. 2023 Jan;183:105792.

51. Hiler W, Trauth SE, Wheeler B, Jimenez A, Radanovic M, Milanovich JR, et al. Stable Isotope
Analysis of Ozark Hellbender (*Cryptobranchus alleganiensis bishopi*) Living and Preserved
Museum Tissue Reveals a Shift in Their Generalist Diet Composition. Ecologies. 2021 Apr
6;2(2):187–202.

52. Ogilvy C, Constantine R, Bury SJ, Carroll EL. Diet variation in a critically endangered marine
predator revealed with stable isotope analysis. R Soc open sci. 2022 Aug;9(8):220470.

53. Varela JL, Rojo-Nieto E, Sorell JM, Medina A. Using stable isotope analysis to assess trophic
relationships between Atlantic bluefin tuna (*Thunnus thynnus*) and striped dolphin
(*Stenella coeruleoalba*) in the Strait of Gibraltar. Marine Environmental Research. 2018
Aug;139:57–63.

54. Velasquez-Vacca A, Seminoff JA, Jones TT, Balazs GH, Cardona L. Isotopic ecology of
Hawaiian green sea turtles (*Chelonia mydas*) and reliability of δ13C, δ15N, and δ34S
analyses of unprocessed bone samples for dietary studies. Mar Biol. 2023 Jul;170(7):81.

55. Quillfeldt P, Masello JF. Compound-specific stable isotope analyses in Falkland Islands
seabirds reveal seasonal changes in trophic positions. BMC Ecol. 2020 Dec;20(1):21.

56. Smith MEK, Ososky JJ, Hunt KE, Cioffi WR, Read AJ, Friedlaender AS, et al. Historical baleen
plates indicate that once abundant Antarctic blue and fin whales demonstrated distinct
migratory and foraging strategies. Ecology and Evolution. 2024 May;14(5):e11376.

57. Magozzi S, Thorrold SR, Houghton L, Bendall VA, Hetherington S, Mucientes G, et al.
Compound-Specific Stable Isotope Analysis of Amino Acids in Pelagic Shark Vertebrae
Reveals Baseline, Trophic, and Physiological Effects on Bulk Protein Isotope Records. Front
Mar Sci. 2021 Sep 1;8:673016.

58. Soncin S, Talbot HM, Fernandes R, Harris A, Von Tersch M, Robson HK, et al. High-
resolution dietary reconstruction of victims of the 79 CE Vesuvius eruption at
Herculaneum by compound-specific isotope analysis. Sci Adv. 2021 Aug
27;7(35):eabg5791.

59. Canseco JA, Niklitschek EJ, Quezada-Romegialli C, Yarnes C, Harrod C. Comparing trophic
position estimates using bulk and compound specific stable isotope analyses: applying
new approaches to mackerel icefish *Champsocephalus gunnari*. PeerJ. 2024 May
17;12:e17372.

60. Norkko A, Thrush SF, Cummings VJ, Gibbs MM, Andrew NL, Norkko J, et al. TROPHIC
STRUCTURE OF COASTAL ANTARCTIC FOOD WEBS ASSOCIATED WITH CHANGES IN SEA ICE
AND FOOD SUPPLY. Ecology. 2007 Nov;88(11):2810–20.

61. Rodriguez ID, Marina TI, Schloss IR, Saravia LA. Marine food webs are more complex but
less stable in sub-Antarctic (Beagle Channel, Argentina) than in Antarctic (Potter Cove,
Antarctic Peninsula) regions. Marine Environmental Research. 2022 Feb;174:105561.

62. Saporiti F, Bearhop S, Vales D, Silva L, Zenteno L, Tavares M, et al. Latitudinal changes in
the structure of marine food webs in the Southwestern Atlantic Ocean. Mar Ecol Prog Ser.
2015 Oct 28;538:23–34.

63. Genelt-Yanovskaya AS, Polyakova NV, Ivanov MV, Nadtochii EV, Ivanova TS, Genelt-
Yanovskiy EA, et al. Tracing the Food Web of Changing Arctic Ocean: Trophic Status of
Highly Abundant Fish, Gasterosteus aculeatus (L.), in the White Sea Recovered Using
Stomach Content and Stable Isotope Analyses. Diversity. 2022 Nov 6;14(11):955.

64. Giménez J, Marçalo A, Ramírez F, Verborgh P, Gauffier P, Esteban R, et al. Diet of bottlenose
dolphins (*Tursiops truncatus*) from the Gulf of Cadiz: Insights from stomach content and
stable isotope analyses. Rosenfeld CS, editor. PLoS ONE. 2017 Sep 12;12(9):e0184673.

65. Guerrero AI, Pinnock A, Negrete J, Rogers TL. Complementary use of stable isotopes and
fatty acids for quantitative diet estimation of sympatric predators, the Antarctic pack-ice
seals. Oecologia. 2021 Nov;197(3):729–42.

66. Gül G, Demirel N. Ontogenetic shift in diet and trophic role of *Raja clavata* inferred by
stable isotopes and stomach content analysis in the Sea of Marmara. Journal of Fish
Biology. 2022 Sep;101(3):560–72.

67. Haug T, Biuw M, Kovacs KM, Lindblom L, Lindstrøm U, Lydersen C, et al. Trophic
interactions between common minke whales (*Balaenoptera acutorostrata*) and their prey

during summer in the northern Barents Sea. Progress in Oceanography. 2024 Jun;224:103267.

68. Mazoudier SQ, Kingsford MJ, Strickland JK, Pitt KA. Stable isotopes reveal sargassum rafts provide a trophic subsidy to juvenile pelagic fishes. Estuarine, Coastal and Shelf Science. 2023 Dec;295:108548.

69. Torres-Poché Z, Mora MA, Boutton TW, Morrow ME. Diet sources of the endangered Attwater's prairie-chicken in Texas: evidence from δ^{13} C, δ^{15} N, and Bayesian mixing models. Ecosphere. 2020 Oct;11(10):e03269.

70. Hucke-Gaete R, Bedriñana-Romano L, Viddi FA, Ruiz JE, Torres-Florez JP, Zerbini AN. From Chilean Patagonia to Galapagos, Ecuador: novel insights on blue whale migratory pathways along the Eastern South Pacific. Johnston D, editor. PeerJ. 2018 Apr 30;6:e4695.

71. Shabangu FW, Stafford KM, Findlay KP, Rankin S, Ljungblad D, Tsuda Y, et al. Overview of the SOWER cruise circumpolar acoustic survey data and analyses of Antarctic blue whale calls. J Cetacean Res Manage. 2024;21–41.

72. Kawaguchi S, Atkinson A, Bahlburg D, Bernard KS, Cavan EL, Cox MJ, et al. Climate change impacts on Antarctic krill behaviour and population dynamics. Nat Rev Earth Environ. 2023 Dec 19;5(1):43–58.

73. Miller EJ, Potts JM, Cox MJ, Miller BS, Calderan S, Leaper R, et al. The characteristics of krill swarms in relation to aggregating Antarctic blue whales. Sci Rep. 2019 Nov 11;9(1):16487.

74. Busquets-Vass G, Newsome SD, Calambokidis J, Serra-Valente G, Jacobsen JK, Aguíñiga-García S, et al. Estimating blue whale skin isotopic incorporation rates and baleen growth rates: Implications for assessing diet and movement patterns in mysticetes. Boyce MS, editor. PLoS ONE. 2017 May 31;12(5):e0177880.

75. Feij B, Kühn S, Meijboom A, Van Franeker JA, Bornemann H, Kelly N, et al. Distribution of Arnoux's beaked whales (*Berardius arnuxii*). Marine Mammal Science. 2024 Jul 13;e13158.

76. Dassis M, Farenga M, Bastida R, Rodríguez D. At-sea behavior of South American fur seals: Influence of coastal hydrographic conditions and physiological implication. Mammalian Biology. 2012 Jan;77(1):47–52.

77. Baylis AMM, Arnould JPY, Staniland IJ. Diet of South American fur seals at the Falkland Islands. Marine Mammal Science. 2014 Jul;30(3):1210–9.

78. Bingham M. The decline of Falkland Islands penguins in the presence of a commercial fishing industry. Revista chilena de historia natural. 2002;75(4):805–18.

79. Heredia-Azuaje H, Niklitschek E, Sepúlveda M, Harrod C, Guerrero A, Peña G, et al. Salmon farming, overfishing and southern sea lion: Not so opportunistic responses of a top predator to human perturbations in the Patagonian Fjords. Estuarine, Coastal and Shelf Science. 2024 Apr;299:108669.

80. Tschopp A, Herrera VP, García NA, Crespo EA, Coscarella MA. Temporal changes in the diet
composition of Shorttail Yellownose skate, when exposed to overfishing conditions in
northern and central Patagonia, Argentina. Hydrobiologia. 2024 Jun;851(11):2695–710.

81. Boveng PL, Hiruki LM, Schwartz MK, Bengtson JL. Population growth of antarctic fur seals:
limitation by a top predator, the leopard seal? Ecology. 1998 Dec;79(8):2863–77.

82. Krause DJ, Goebel ME, Kurle CM. Leopard seal diets in a rapidly warming polar region vary
by year, season, sex, and body size. BMC Ecol. 2020 Dec;20(1):32.

83. Sperou ES, Crocker DE, Borras-Chavez R, Costa DP, Goebel ME, Kanatous SB, et al. Large
and in charge: cortisol levels vary with sex, diet, and body mass in an Antarctic predator,
the leopard seal. Front Mar Sci. 2023 Jun 23;10:1179236.

813