**TITLE**

Revealing function-altering MECP2 mutations in individuals with autism spectrum disorder

using yeast and *Drosophila*.

**AUTHOR LIST**

Eric Chen[1]*, Jessica Schmitt[1]*, Graeme McIntosh[1]*, Barry P. Young[1], Tianshun Lian[1], Jie Liu[1],

Kexin K. Chen[2], J. Beatrice Liston[1], Lily MacDonald[1], Bill Wang[1], Sonia Medina Giro[1],

Benjamin Boehme[1], Mriga Das[1], Seevasant Indran[1], Jesse T. Chao[3], Sanja Rogic[2], Paul

Pavlidis[2], Douglas W. Allan[1]#, and Christopher J.R. Loewen[1]#

[1] Department of Cellular and Physiological Sciences, The Life Sciences Institute, The University

of British Columbia, 2350 Health Sciences Mall, Vancouver, BC, Canada, V6T 1Z3

[2] Department of Psychiatry and Michael Smith Labs, The Michael Smith Institute, The

University of British Columbia, 301-2185 East Mall, Vancouver, BC, Canada, V6T 1Z4

[3] Department of Medical Biophysics, Sunnybrook Research Institute, The University of Toronto,

2075 Bayview Avenue, E240, Toronto, Ontario, Canada, M4N 3M5

(co-first *, co-corresponding **#)**

(email: christopher.loewen@ubc.ca, doug.allan@ubc.ca)

Running title: Altered MECP2 function in autism

Keywords: MECP2, autism spectrum disorder, ASD, Rett syndrome, variant prediction, budding

yeast, Drosophila, genetic interactions, Sentinel Interaction Mapping, VariCarta, transgenesis

**ABSTRACT**

Pathogenic variants in MECP2 commonly lead to Rett syndrome, where MECP2's function as a

DNA cytosine methylation reader is believed critical. MECP2 variants are also catalogued in

individuals with autism spectrum disorder (ASD), including nine missense variants which had no

known clinical significance at the start of this study. To assess these nine variants as risk alleles

for ASD, we developed MECP2 variant functional assays using budding yeast and *Drosophila*.

We calibrated these assays with known pathogenic and benign variants. Our data predict that

four ASD variants are loss of function and five are functional. Protein destabilization offers

insight into the altered function of some of these variants. Notably, yeast and *Drosophila* lack

DNA methylation, yet all Rett pathogenic and ASD variants located in the methyl DNA binding

domain that we analyzed proved to be loss of function, suggesting a clinically-relevant role for

non-methyl DNA-binding by MECP2.

**INTRODUCTION**

The X-linked protein MECP2 was first identified in screens for proteins recruited to methylated

DNA (Lewis et al. 1992), and loss of function (LoF) alleles of MECP2 were subsequently found

as causative in the neurodevelopmental disorder, Rett syndrome (Amir et al. 1999). Since then

variants in MECP2 have been found in most cases of Rett, but also in other neurodevelopmental

disorders, including autism spectrum disorder (ASD) (Wan et al. 1999; Amir et al. 2000;

Nomura 2005; Chahrour and Zoghbi 2007). The origin and phenotypic consequence of MECP2-

damaging variants differ notably between sexes. In females, the vast majority of these variants

are *de novo*, while in males, they can also be inherited from unaffected mothers (Ananth et al.

2024). Variants that typically result in Rett syndrome in females tend in males to cause severe

neonatal encephalopathy, often leading to death in the first year of life. However, in males with

two populations of X chromosomes, such as those with Klinefelter syndrome (47XXY) or

somatic mosaicism, these variants result in a milder Rett-like condition known as male RTT

encephalopathy. Males can also be affected by variants that are seemingly benign in females,

which can lead to moderate to severe intellectual disability (Pascual-Alonso et al. 2021).

MECP2 has largely been examined as a DNA cytosine methylation reader via its methyl cytosine

binding domain (MBD) that recruits co-repressor complexes to DNA to repress transcription via

its transcriptional repressor domain (TRD) (Jones et al. 1998; Ng and Bird 1999; Lunyak et al.

2002; Harikrishnan et al. 2005). The MBD, and to a lesser extent the TRD, harbors most Rett

LoF variants (Ehrhart et al. 2021), resulting in variants which are believed to reduce binding to

methylated DNA or reduce recruitment of co-regulator complexes, respectively (Ballestar et al.

2000; Yusufzai and Wolffe 2000). While numerous MECP2 variants have been found in patients

that exhibit Rett with ASD, recent reports have annotated five MECP2 missense variants in ASD

without an associated Rett diagnosis (R91W, G195S, A202G, E282G, S411R) (Schaaf et al.

2011; Wang et al. 2016; Wen et al. 2017; Guo et al. 2018; Leblond et al. 2021). However, it is

unclear whether any of these variants are function altering and clinically significant. Two of

them, R91W and A202G, are classified by ClinVar (Landrum et al. 2018) as variants of

uncertain significance (VUS) in cases of severe neonatal onset encephalopathy with

microcephaly. E282G is classified by ClinVar as likely pathogenic. The remaining two variants

(G195S and S411R) have no current ClinVar classification.

While MECP2 is understood to primarily be a methyl-cytosine DNA (DNAm) binding protein, it

has become clear it also binds unmethylated DNA with a preference for GC-rich sequences

(Georgel et al. 2003; Hansen et al. 2010; Skene et al. 2010; Cheng et al. 2014). This has been

proposed as an alternate hypothesis for why MECP2 appears to preferentially bind methylated

GC-rich regions of the genome, as opposed to binding DNAm methylation *per se*. It can be

challenging to segregate DNAm-dependent from DNAm-independent functions of MeCP2 *in*

*vivo*, as DNAm is both abundant and essential in mammals. Consequently, it is still unclear what

fraction of MeCP2's functions are dependent on its interaction with methylated and/or

unmethylated DNA. Such ambiguity leaves unresolved the question of whether disease relevant

variants in MECP2 selectively disrupt binding to methylated versus unmethylated DNA.

There are extremely low levels of methylated cytosine in the *Drosophila* genome, and none in

*Saccharomyces cerevisiae* (Lyko et al. 2000; Buitrago et al. 2021). Thus, fly and yeast offer

avenues to explore the interactions of MECP2 with unmethylated DNA, and to test if Rett and

ASD variants alter MECP2 binding to unmethylated DNA. Moreover, while neither organism

has an ortholog of *MECP2*, both organisms are robust genetic models that have proven well

suited to comparing the function of large numbers of gene variants (Post et al. 2020; Young et al.

2020; Ganguly et al. 2021; Her et al. 2024). MECP2 has been expressed in *Drosophila*, where it

associates with chromatin and mediates transcriptional repression via interactions with orthologs

of known MECP2 physical interactors in humans, including Sin3A, N-CoR, REST as well as

other chromatin modifiers (Cukier et al. 2008). In fly neurons, MECP2 impacts dendritic

branching and behavioural output (Vonhoff et al. 2012; Williams, White, et al. 2016) and C-

terminal truncation mutants (R294X specifically) promote neuronal apoptosis (Williams, Mehler,

et al. 2016). In yeast, MECP2 has recently been shown to bind chromatin and mediate

widespread transcriptional repression (Brown et al. 2025 Mar 18).

Our goal in this study was to use the *Drosophila* and budding yeast models to test the functional

impact of these ASD MECP2 variants to determine whether they may confer autism risk, by

virtue of their altered function. We calibrated our assays with MECP2 variants classified by

ClinVar as pathogenic or benign, in accordance with American College of Medical Genetics and

Genomics (ACMG) guidelines for experimental assays (Brnich et al. 2019); first to validate our

assays for variant assessment, and second to provide a rubric for determining if test ASD variants

are likely pathogenic or likely benign. All calibration variants functioned as expected in the

*Drosophila* assay, indicating that we could functionally assess any test variants. The yeast assay

could be calibrated appropriately for the MBD domain, but the MBD proved to be the only

domain showing the expected separation between benign and pathogenic calibration variants.

Our analyses indicate four of the nine test ASD variants were LoF, making these variants

candidate risk alleles for ASD. Interestingly, these variants mostly proved to be less severe LoF

than Rett-associated pathogenics, providing a possible explanation for how MECP2 mutations

can lead to ASD, but not Rett.

**METHODS**

**MECP2 isoform selection**

*MECP2* has two major isoforms, E1 and E2. We selected to work with *MECP2* isoform E2

(ENST00000303391, NM_004992) based on the recommendation by the ClinGen

Rett/Angelman-like Expert Panel (McKnight et al. 2022). This transcript is annotated as the

APPRIS (Annotation of Principal and Alternative Splice Isoforms (Rodriguez et al. 2013))

principal isoform in Ensembl genome browser, based on the annotations for protein structure,

function and cross-species conservation, and is the highest expressed transcript in the brain

(GTEx portal; **gtexportal.org**).

**Variant aggregation, annotation and prioritization**

We utilized our in-house computational pipeline to aggregate *MECP2-E2* missense variants from

ClinVar and VariCarta and perform comprehensive annotation. We collected ClinVar variants

(downloaded in January 2022) that have at least one gold review star, i.e., the review status

(CLNREVSTAT) has to have one of the following designations: "criteria provided, single

submitter", "criteria provided, conflicting interpretations", "criteria provided, multiple

submitters, no conflicts" or "reviewed by expert panel". For each of these variants, we also

obtain its interpretation (clinical significance; CLNSIG) and reported condition (disease name;

CLNDN). The ASD missense variants were obtained from VariCarta, a database of variants

found in individuals diagnosed with ASD and reported in the scientific literature, that we created

(Belmadani et al. 2019). We annotated the aggregated variants using the Ensembl VEP tool

(McLaren et al. 2016). We also added protein domain annotation relevant in the context of Rett

Syndrome (Good et al. 2021).

*MECP2-E2* **cloning**

A *MECP2-E2* cDNA clone was obtained (Horizon, Clone Id: 3956518, # MHS6278-

202830559), and *MECP2* coding sequences were amplified with iProof DNA polymerase (Bio-

Rad, California USA, #1725301) using the following primers:

GA-attL1-MECP2-522-F (tgtacaaaaaagcaggctccggcaccatggtagctgggatgttag)

GA-attL1-MECP2-522-R (ttgtacaagaaagctgggtctatcagctaactctctcggtc)

Amplicons were cloned in PCR8-GW-TOPO (Invitrogen, ThermoFisher Scientific, Waltham,

USA, catalog #K250020) using NEBuilder HiFi DNA Assembly Master Mix (New England

Biolabs, USA,  # M5520). The resulting construct was named MECP2-pENTRY. A sequence of

CCGGCACC was introduced in primer GA-attL1-MECP2-522-F (underlined) preceding the

start codon for *MECP2-E2*. After LR cloning of MECP2-pEntry into pDEST vectors, this

CCGGCACC sequence can function as a Kozak sequence to initiate translation, or as an in-

frame linker for N-terminal peptide sequences. Missense variants were generated in MECP2-

pENTRY either using the Q5 mutagenesis kit (New England Biolabs, USA, # E0554) in house,

or were made by GenScript (New Jersey, USA).  For MECP2 expression clone construction,

MECP2-pENTRY plasmids were subcloned in pAG426GAL-ccdB (a kind gift from a gift from

Susan Lindquist; Addgene plasmid # 14155) or pGW-HA.attB (Bischof et al. 2007) using the

Gateway LR Clonase reaction (Thermo Fisher Scientific, Waltham, USA, catalog # 11791020).

All transgenic *UAS-MECP2* variants (on pGW-HA.attB) *Drosophila* strains used were verified

by Sanger sequencing (UBC NAPS, Vancouver, Canada) or whole amplicon sequencing

(Plasmidsaurus, CA, USA) (Supplemental Data 1).

**Sentinel interaction mapping (SIM) in yeast**

SIM screening for *MECP2* was performed as described previously (Young et al. 2020). A singer

RoToR HDA colony arraying robot was used to copy arrays of yeast cells. All steps were

performed at a density of 1,536 colonies per plate and plates were incubated for 24 hours at 30

°C prior to the next pinning step, except where indicated otherwise. The SGA query yeast strain

Y7092 (Tong et al. 2001) was transformed with plasmid pSI7338-MECP2 and arrayed on

synthetic defined medium lacking uracil. The haploid yeast deletion mutant array (DMA) was

arrayed across a total of four plates on YPD medium containing 200 mg/L G418 sulfate

(YPD+G418). The query array was mated to the DMA by subsequently pinning both arrays on to

fresh YPD plates. Diploids were selected by pinning colonies to synthetic defined medium

lacking uracil containing 200 mg/L of G418 sulfate and subsequently pinned to YPD+G418

plates. Each plate was then pinned in triplicate to enriched sporulation media (Tong et al. 2001)

and incubated at 22 °C for seven days. Following this, *MAT*a haploid cells were germinated by

pinning to synthetic complete medium lacking histidine, arginine and lysine with the addition of

50 mg/L canavanine sulfate, 50 mg/L thialysine and incubating for two days. To select for

deletion strains carrying the MECP2-Ref plasmid, colonies were pinned to two pairs of plates

("control" and "experimental") containing synthetic complete medium lacking histidine,

arginine, lysine and uracil, containing 50 mg/L canavnine sulfate, 50 mg/L thialysine, 200 mg/L

G418 sulfate ("HURK+G418" medium). The experimental set was supplemented with 10 nM -

estradiol to induce expression of *MECP2*; this was omitted from the control set. The final plates

used for imaging were generated by copying each of these plates once more to the same medium.

Plates were scanned using a CanoScan 8600F flatbed scanner and subsequently analyzed using

*Balony* software (Young and Loewen 2013) to compare the normalized colony size for each

strain in the presence or absence of estradiol. Genetic interactions were defined as those strains

where the ratio of experimental:control colony size was below a threshold determined by *Balony*

in all three replicates and with a p-value $< 0.05$. The strains *pho85Δ, rad50Δ, slx5Δ, swi4Δ,*

*swi6Δ,* and *xrs2Δ* were confirmed to be correct by PCR amplification of down-tag barcodes with

the standard primers B-U1 and B-D1 and sequencing.

**Yeast quantitative liquid growth assays**

Yeast strain JHY716 was transformed with various plasmids containing *MECP2* variants placed

downstream of the *GAL1* promoter. Four independent clones were picked from transformation

plates and transferred to SD-URA media containing 2% raffinose, 2% galactose to induce

expression of *MECP2*. For each clone, a 2 mL culture in SD-URA with 2% raffinose, 2%

galactose was grown overnight at 30 °C in a shaking incubator. The following morning, cultures

were diluted to an OD600 of ~0.25 in the same media. After 6 hours of incubation at 30 °C, log

phase cultures were diluted to an OD600 of 0.125. Three aliquots of 200 µl were taken from each

culture and transferred to a well of a flat-bottomed 96-well dish. Growth of cultures was

measured overnight at 30 °C with shaking at 800 rpm in an Agilent Logphase 600 plate reader.

All plates contained strains expressing wild type *MECP2* and the corresponding empty vector as

controls. The exponential growth constant k was determined for each well by fitting to an

exponential growth equation during the exponential phase (typically between 5 and 10 hours post

dilution).

**Protein abundance assays**

To determine the abundance of MECP2 variants in yeast, strains were grown in SD-URA media

containing 2% raffinose to early log phase. MECP2 expression was induced by the addition of

2% galactose to the growth medium. Cells were harvested three hours later by centrifugation and

stored at -80 °C for subsequent processing. Cell pellets were resuspended in Laemmli sample

buffer, and heated to 95 C for 5 minutes. Cell lysis was achieved by the addition of 50 µl acid-

washed glass beads followed by two rounds of disruption in a MP Biomedical FastPrep-24

homogenizer (maximum speed for 60 s with 5 minutes cooling between cycles). Samples (0.2

OD600 equivalents) were separated by SDS-PAGE on Bio-Rad 4-15% pre-cast gels and

transferred to nitrocellulose membranes. MECP2 was immunoblotted using Sigma antibody

(M9317), with PGK1 (Abcam AB113687) used as a loading control. For visualization,

secondary antibodies conjugated to Cy3 (MECP2) and AlexaFluor-488 (PGK1) were used and

imaged using a Bio-Rad ChemiDoc MP Imaging System. MECP2 band volume was normalized

to PGK1 band volume for each variant.

***Drosophila* strains**

All flies were maintained on standard cornmeal food at 25 °C and 70% humidity unless

otherwise specified. *P{GAL4}A9* (BL8761) (referred to as *A9-GAL4*) was obtained from

Bloomington *Drosophila* Stock Centre (Indiana, USA). *UAS-MECP2* reference (*UAS-MECP2-*

*ref*) and variant sequences (collectively termed *UAS-MECP2.#*) were integrated into embryos of

genotype  *y[1] w[] P{y[+7.7]=nos-phiC31\int.NLS}X; P{y[+t7.7]=CaryP}attP40* (referred to as

attP40) by Genome ProLab (Quebec, Canada), and maintained as *w[1118]; P{y[+t7.7]*

*w[+mC]=UAS-MECP2.#.GW.*}attP40*, where # represents the variant and * represents

integrant line number (Supplemental Data 1).

**Crossing scheme**

Experimenters were double blinded to genotype throughout. Homozygous *UAS-MECP2* males

and a control *attP40* line were crossed to *A9-GAL4* virgins (ratio 4 males : 12 females), and

maintained at room temperature (~21 °C ) for 72 hours. Crosses were then transferred into fresh

'assay' vials, and transferred each 24 hrs each for three days, generating three 'assay' vials per

cross. After removal of adults from each vial, vials were placed into a 29 °C incubator (70%

humidity). Three days after eclosion, adult flies were sorted by sex and placed in pure

isopropanol at -20 °C. All progeny flies for comparison were tested in a single batch.

**Wing imaging**

Experimenters were double blinded to genotype throughout. Flies were removed from

isopropanol, dried on paper towels, and 10-25 left wings were removed per genotype per assay

vial (one wing per fly). Wings were mounted on microscope slides in Canadian Balsam Fir

(1.01691.0025, CAS-No: 8007-47-4, Sigma Aldrich, MA, USA), coverslipped and sealed with

nail polish. Females have larger wings and the X-linked *A9-GAL4* driver is expressed at higher

levels in males, due to X-chromosome male-hyperactivation. Thus, it is important to analyze

males and females separately. While the impact of MECP2 variants in the wing was similar

between males and females, the phenotype was substantially stronger in males due to the higher

GAL4 levels; hence, higher MECP2 levels. To simplify analysis, we therefore only fully

analyzed male wings. Slides were compressed while drying for 24 hrs. Mounted wings were

scanned using a Panoramic Midi II digital slide scanner (3DHISTECH, Budapest, Hungary).

Scanned fly wing images were cropped at 15x zoom using CaseViewer (3DHISTECH,

Budapest,Hungary). Images were imported into Imaris v10.0 software (Bitplane, MA, USA) for

measurement of wing vein area per wing area, using Labkit machine learning and Fiji software

(University of Wisconsin-Madison, USA). In Labkit, we developed

"BIGSMOOTHCLASSYBOI", a machine learning algorithm to automate measurement of wing

vein area for calculation of wing vein area/wing area (details provided in Supplemental Methods

1).

**Statistical analysis of variant effects**

Yeast growth constants (k) and fly relative wing vein area data were analyzed using methods

similar to Post et al. (Post et al. 2020). Briefly, linear mixed-effects models were fit to each data

set using the *lmerTest* R package (Kuznetsova et al. 2017) to estimate the phenotypic effects of

each variant compared to the *MECP2-ref* constructs, treating experimental batches as random

effects (for R scripts see Supplemental Methods 2 and 3). For visualization, values were

standardized to the range [0-1] where 0 represents the mean phenotype measured for an empty

vector, and 1 represents the mean phenotype with the *MECP2-ref* construct. Four biological

replicates, each with three technical replicates were performed for yeast variant growth assays,

while for *Drosophila* wing assays 10-25 wings, each from a different animal were imaged per

genotype.

**RESULTS**

*Selection of variants*

We selected benign and pathogenic variants with clear clinical classifications to calibrate our

functional assays, and nine test (ASD) variants (Figure 1, Supplemental Data 1). We focused on

ClinVar variants with "pathogenic", "likely_pathogenic", "benign" and "likely_benign" clinical

interpretation as calibration variants. Priority was given to those with either "reviewed by expert

panel" or "criteria provided, multiple submitters" review status. A few exceptions were made.

D121E, a single-submitter benign variant, and T148A, originally classified as benign, but now

listed as conflicting were selected because there were no other benign variants in the MBD that

matched the aforementioned criteria. When selecting pathogenic variants, in addition to using

review status as a criterion, we also prioritized the ones with "Rett syndrome" condition

annotation as well as the ones that have been frequently observed in Rett patients (Good et al.

2021). In total, we selected 37 calibration variants, 10 benign and 27 pathogenic. We selected

ASD MECP2 variants from VariCarta as test variants as long as they did not have a clear clinical

interpretation in ClinVar, for the total of 9 test (ASD) variants. Since the time of our initial

selection, six test (ASD) variants have been given updated classifications in ClinVar: P152L,

P217L and E282G becoming pathogenic/ likely pathogenic, and G195S, P376S and P388S

becoming benign (Figure 1 displays the initial classifications).

***MECP2 produces a slow growth phenotype in yeast***

*MECP2-ref* was cloned into a high-copy 2-micron plasmid designed to give controlled

expression in response to addition of estradiol. Using an anti-MECP2 antibody we monitored

MECP2 protein levels by Western blot in response to increasing estradiol (Figure 2A). In the no-

estradiol condition we detected low levels of MECP2 protein, which increased with increasing

estradiol and then saturated at ~10 nM. Next, we measured the growth rate of WT yeast

containing either the *MECP2-ref* plasmid or an empty vector control in response to estradiol

addition (Figure 2B). Compared to empty vector which had little effect in response to increasing

estradiol, yeast containing MECP2 showed a dose-dependent decrease in exponential growth

rate. We compared this to another vector designed for strong overexpression utilizing the

*Gal1/10* promoter in a 2-micron vector and observed an even greater decrease in growth rate in

the presence of MECP2 (Figure 2C). Thus, expression of MECP2 reduced the fitness of WT

yeast in a dose-dependent manner.

*MECP2 expression affects DNA-related processes in yeast*

To understand how MECP2 might be functioning in yeast and if this is related to the normal

function of MECP2 we employed a genetic interaction mapping approach we previously

developed to study the function of human genes in yeast, called Sentinel Interaction Mapping

(SIM) (Young et al. 2020). For SIM we express the human gene of interest in a collection of

nearly 4,800 yeast deletion mutants and systematically identify genetic interactions, which

indicate functional associations between the human gene and yeast cellular pathways. Hence,

these genetic interactions will inform us on the function of the human gene in yeast and if this is

relevant to its function in humans. We performed a SIM screen with *MECP2-ref* in which

MECP2, under control of the estradiol responsive promoter in the 2-micron vector, was

expressed in ~ 4,800 different yeast deletion mutant strains in high-density arrays (see Methods

for details). Colony sizes were scored using Balony software and genetic interactions were

identified by comparing control arrays containing no estradiol (uninduced) to experimental

arrays containing 10 nM estradiol (induced). We chose 10 nM estradiol because MECP2 protein

expression saturated at this concentration and the effect on growth was not too severe, which

would otherwise hinder identification of genetic interactions.

The SIM screen identified 123 aggravating genetic interactions in 3/3 biological replicates

(Supplemental Data 2). We examined functional enrichment for these interactions using Gene

Ontology (GO) analysis (Figure 3A; also see Supplemental Data 3 for gene lists). This revealed a

network of interconnected functions related to DNA and chromosome stability, with the most

significant enrichment being for the terms "Cellular response to stress" (30 genes) and "DNA

damage response" (24 genes). Functions related to DNA binding (27 genes), cell cycle (21

genes), Golgi (8 genes) and nitrogen compound utilization (17 genes) were also enriched for.

The large enrichment for DNA-related functions was consistent with MECP2 being a DNA

binding protein with roles in transcriptional repression and chromatin organization. We focused

on a subset of interactions from GO terms most relevant to MECP2 function. Shown in Figure

3B is the effect on growth of the individual mutants from the screen upon MECP2 induction for

the GO terms "Chromatin organization", "DNA binding" and "DNA damage response". Because

these are aggravating interactions, MECP2 induction (10 nM estradiol) resulted in decreased

growth rate of the mutants relative to no induction (0 nM estradiol). Growth was normalized to

account for the effect of MECP2 induction on WT yeast growth.

Looking more closely at these genes, we identified five out of the ten members of the *RAD52*

epistasis group, *RAD50/51/52/54* and *XRS2* (*RAD55/57/59, RDH54* and *MRE11* were not

identified) (Figure 3B). Mutants in this group are defective in the repair of DNA damage,

maintenance of telomere length and meiotic and mitotic recombination (Symington 2002). We

also identified three out of four of the subunits of the highly conserved MRX endo/exonuclease

complex, which functions in repair of DNA double strand breaks, detection of damaged DNA,

DNA damage checkpoint activation, telomerase recruitment and suppression of chromosomal

defects (Syed and Tainer 2018). We identified Rad50, Xrs2 and Sae2, but not *MRE11* in our

screen (Figure 3B). *RAD50* and *XRS2* were among the strongest interactions such that MECP2

expression reduced growth of the *rad50Δ* and *xrs2Δ* mutants to almost undetectable levels.

We also identified both subunits of the SBF transcription factor complex, Swi4 and Swi6. Swi6

is also a component of the MBF complex (MBF comprises Swi6 and Mbp1), which along with

SBF functions to regulate transcription of the G1/S gene cluster as well as transcription of genes

involved in DNA synthesis and repair (Bähler 2005). *swi6Δ* cells grow slowly, have a delayed

G1/S transition and are sensitive to DNA damaging agents. Expression of MECP2 in the *swi6Δ*

mutant severely reduced growth indicating a strong interaction. Additionally, we identified a

strong interaction with the cyclin-dependent kinase *PHO85*, which regulates many cellular

processes in response to stress and also plays an important role in G1/S transition via interaction

with various G1 cyclins (Jiménez et al. 2013). *Δpho85* cells grow slowly, have increased mitotic

chromosome loss and shortened telomeres. Taken together, these strong aggravating interactions

with genes with central functions in DNA synthesis and repair, chromatin organization and G1/S

cell cycle progression suggested MECP2 expression severely antagonized these processes which

resulted in reduced fitness.

***Assessment of MECP2 variants in yeast***

We decided to simply exploit the robust slow growth phenotype observed upon MECP2

expression in WT yeast to determine the effects of MECP2 variants, anticipating variants that

interfered with MECP2 function would no longer reduce growth. We measured yeast growth in

liquid culture for all 46 variants using the galactose-inducible 2-micron vector, because this

vector produced the largest growth defect. We calculated exponential growth rates for WT yeast

expressing MECP2 variants and using a linear mixed-effects model estimated the effects of the

variants on MECP2 function (raw data is provided in Supplemental Data 4). Shown in Figure 4A

is the model's estimates of MECP2 function, arranged by variant amino acid position. Values

were standardized to the range [0-1] where 0 represents the mean phenotype measured for an

empty vector, and 1 represents the mean phenotype with the *MECP2-ref* construct. Hence,

variants with values near to 1 functioned similarly to MECP2-ref, whereas values greater or less

than 1 indicated increased or decreased function, respectively (error is plotted as 95% confidence

intervals).

Benign calibration variants had little effect on function as expected across all MECP2 domains.

The one exception was the D121E variant which was nearly complete LoF, for which we have

additional supporting evidence in fly leading to reclassification of this variant as pathogenic (see

below). For the MBD domain, all seven pathogenic calibration variants were complete or near-

complete LoF (we define a variant is complete LoF when its 95% confidence interval overlaps

with 0). This supported that our assay was disease-relevant for variants in the MBD. L100V,

R106W, R133C, and T158M have been previously reported to have reduced DNA binding *in*

*vitro* and *in vivo* (Tillotson and Bird 2020), suggesting our assay reported on this function of the

MBD. Residue R133 is part of the DNA interface and interacts directly with guanine of the

mCpG island, and T158 is part of the Asx-ST motif that interacts with the DNA backbone

(Tillotson and Bird 2020). The two test (ASD) variants in this domain, R91W and P152L, were

found to have decreased function, although not as severe as the calibrating pathogenics. In

contrast, calibrating pathogenics outside the MBD generally did not show decreased function.

This indicated the yeast assay could not reliably detect altered function for variants outside the

MBD.

***Protein level of variants in yeast***

We measured protein levels for all variants in the MBD to determine if decreased stability/

increased turnover might be a factor in our assessment of function. We also included additional

non-MBD calibrating benigns from the TRD (T228S) and the CTD (S356G) for comparison; and

P388S because this variant showed the most reduced function of all non-MBD variants. MECP-2

protein levels from yeast extracts were measured on Western blots and normalized to MECP2-ref

(Figure 4B and Supplemental Figure 1). For the MBD, 8 out of 13 MBD variants had reduced

levels relative to MECP2-ref, while 5 were not significantly different from MECP2-ref. Variants

with decreased protein levels ranged from ~65% (T148A) to ~37% (R106W) of MECP2-ref. The

two calibrating benigns outside the MBD, T228S and S356G, did not show decreased protein

levels, whereas the test (ASD) variant P388S had substantially reduced protein levels (~ 25% of

MECP2-ref).

Comparing protein levels to function of variants there appeared to be little correlation. Some

variants had normal protein levels yet were non-functional (L100V, P101S, D121E, R133C).

Other variants showed both decreased protein levels and decreased function (R106W, T158M,

G163W, P388S). This suggested in some cases the variant affected MECP2 DNA binding

directly leading to decreased function (e.g., R133C), whereas in other cases decreased function

might also be a result of destabilization/ increased turnover (e.g., R106W). To investigate this in

more detail we looked to variants outside the MBD, because these should not directly affect

DNA binding. P388S was the only variant outside the MBD that showed moderate LoF (~ 50%;

Figure 3A) and we found it to have greatly reduced protein levels, whereas a variant in the same

domain, S356G, was functional and had normal protein levels. The modest decrease in function

of P388S relative to its low protein levels, suggested our functional assay was relatively

insensitive to destabilization. The T148A variant, which had reduced protein levels (~65% of

MECP2-ref), but was completely functional in the assay, further supported this conclusion.

Overall, these data suggested the yeast functional assay primarily reported on the intrinsic DNA

binding ability of the MECP2 MBD.

***Using a fly wing vein assay for assessing MECP2 variant function***

Expression of *MECP2* in *Drosophila* tissues has a wide range of phenotypes (Cukier et al. 2008;

Vonhoff et al. 2012; Williams, White, et al. 2016; Williams, Mehler, et al. 2016). From these

reports, we chose to assess MECP2 variants in developing wing tissues. This was because (1) the

reported ectopic wing vein phenotype could be exploited here to provide a quantitative readout

of MECP2 variant protein function, and (2) *MECP2* was shown to genetically interact with

known MECP2 co-repressor complexes in the context of this specific phenotype (Cukier et al.

2008), suggesting the wing vein phenotype was relevant to MECP2 function in humans. These

provided strong precedence for using this wing vein assay for assessing the function of ASD

variants in MECP2.

We integrated *UAS-MECP2-ref* into the *attP40* locus on the *pGW.HA.attB* transgenic vector

backbone to control for genome position effect and copy number (Bischof et al. 2007), to reduce

variability in expression levels between integrants. In pilot experiments we tested a range of

conditions including wing imaginal disc *GAL4* drivers, incubation temperatures and sex of

progeny (as females have larger wings). We found that the ideal set of variables to produce a

robust ectopic wing vein phenotype were the *A9-GAL4* driver in males (X-chromosome

hyperactivation in male flies results in higher GAL4/UAS activity) raised at 29 °C (Figure 5,

arrows). We also noted *MECP2-ref* expression resulted in smaller wings (Figure 5). Thus, we

could proceed with assessing the effects of variants using this assay.

***Assessment of 25 MECP2 variants using the wing vein assay***

In total 25 transgenic lines were generated, including eight calibrating benign, eight calibrating

pathogenic and nine test (ASD) variants. For all transgenic lines we passaged mating adults

through three consecutive 'assay' vials. Adult progeny were then collected 3 days post-eclosion

from each assay vial, and a single wing from a minimum of 10 adult male progeny per assay vial

was removed and slide-mounted, for a total of 30 wings per variant. Mounted wings were

scanned and total wing vein area / wing area was analyzed using a Labkit machine learning

algorithm that semi-automated the quantitation of wing vein area (see Methods).

Next, we examined wing vein phenotypes induced by expression of all 25 MECP2 variants, with

*attP40* and *MECP2-ref* controls in parallel. Representative images for controls, selected

calibrating benign and pathogenic variants and all test (ASD) variants are shown in Figure 5

(also see Supplemental Figures 2 and 3 for representative images of all MECP2 variants and the

masked area that was considered to contain vein tissue). Benign calibration variants all showed

increased ectopic wing veins (T203M and T228S are shown) whereas calibrating pathogenics

showed very little (L100V and T158M are shown), and test (ASD) variants had a range of

outcomes (Figure 5). For example, P376S showed increased ectopic wing veins similar to

*MECP2-ref* whereas P152L had few extra wing veins.

To quantify this phenotype, we trained two machine learning algorithms to quantify (1) the total

area of the wing vein region and (2) the total area of the wing (see Supplemental Data 5 for

details). Then using a linear mixed effects model these two values were used to calculate the

estimated effect of variants relative to *MECP2-ref* and were normalized to *MECP2-ref* (at 1) and

*attp40* (at 0) (Figure 6, error bars are 95 % confidence intervals). All benign calibration variants

functioned similar to *MECP2-ref*, except for D121E. This variant was classified in ClinVar as

benign by a single contributor, but we find both yeast and fly assays characterized it as a strong

LoF, hence we predict D121E is likely pathogenic. All pathogenic calibration variants showed

varying degrees of LoF, indicating the wing vein assay reported on disease relevant functions for

MECP2. Most were strong LoF variants, except for P225R, which had reduced function but not a

complete LoF. This successful calibration of the assay indicated clinically pathogenic and benign

variants can be correctly classified in the fly model. We were particularly confident for variants

in the MBD and TRD domains, because we tested multiple calibrating pathogenics in these

domains, and both DNA and co-repressor interactions through these domains are validated in this

assay. In the same experimental batch as the calibrating benign and pathogenic variants

(controlling for food, temperature and humidity), we determined the wing vein phenotype for all

nine test (ASD) variants. We found that both MBD variants R91W and P152L were LoF. The

TRD variants P217L and E282G exhibited a partial LoF with data distributed similarly to the

TRD pathogenic variant P225R. Finally, the Interdomain (G195S, A202G) and CTD variants

(P376S, P388S and S411R) all exhibited normal function, similar to the calibrating benigns.

Thus, we conclude that only test (ASD) variants in the MBD and TRD domains were LoF.

Finally, we compared yeast and fly assay results for MBD variants only, because the yeast assay

could only reliably measure MBD variants. We observed remarkable similarity in results

between the two assays. Of particular interest was that both test (ASD) variants R91W and

P152L appeared more functional in both yeast and fly assays than the pathogenic calibration

variants. This suggested ASD variants may retain more DNA binding capacity than Rett variants,

given these assays likely reported on the intrinsic ability of MECP2 to bind DNA.

**DISCUSSION**

***Clinical classification of variants***

Using our assays calibrated with variants classified in ClinVar as pathogenic and benign, we are

able to provide functional assessments for nine test (ASD) variants, five of which have been

found in cases of ASD without Rett (underlined): R91W, P152L, G195S, A202G, P217L,

E282G, P376S, P388S, S411R (Figure 1). We found that R91W and P152L were LoF variants

(yeast and fly) and we classify these as likely pathogenic. Our classification for P152L agrees

with the updated pathogenic ClinVar classification and we re-classify R91W from a VUS in

ClinVar. G195S had a modest gain of function (fly), hence we classify it as likely benign. Our

likely pathogenic classification for G195S agrees with the updated benign ClinVar classification.

A202G was found to be functional (fly). The ClinVar classification for this variant is conflicting

(one benign and one VUS report) and we propose this variant be classified as likely benign.

P217L and E282G showed moderately reduced function (fly) consistent with their updated likely

pathogenic ClinVar classifications. P376S, P388S and S411R were found to be functionally

normal (fly). For P376S and P388S this is consistent with their updated benign classifications in

ClinVar. However, since P388S appeared to be a destabilized protein in yeast we suggest P388S

is worthy of further examination. We classify S411R as likely benign providing the first

classification for this variant. Importantly, we provide functional evidence to classify two

variants found in ASD but not Rett as likely pathogenic (R91W, E282G), suggesting certain

mutations in MECP2 may cause ASD while others will cause Rett. Further, our data suggest

variants found in ASD but not Rett may be less severe than Rett causing mutations, since the

ASD-only variants we tested were more functional than Rett causing variants.

Finally, we provide evidence for reclassification of two variants, D121E and T148A, currently

classified in ClinVar as benign and conflicting, respectively. D121E was LoF in both yeast and

*Drosophila* assays, hence we propose D121E be reclassified as likely pathogenic. Interestingly,

our data are consistent with *in vitro* studies showing a ~7-fold reduction in the DNA binding

capacity of the D121E variant (Free et al. 2001). This variant is neither in VariCarta nor

RettBASE and it has one report in ClinVar, but this report lacks a description for how a benign

classification was made. Interestingly, the computational predictors SIFT, PolyPhen and CADD

all predict this variant is likely to be LoF (Supplemental Data 1), supporting our likely

pathogenic classification. For T148A, the current ClinVar classification is conflicting. We find

this variant to be functional (yeast), but to have modestly decreased protein levels (~65% of

MECP2-ref), suggesting it may have reduced stability. SIFT, PolyPhen and CADD all predict

this variant to be damaging. We propose T148A be reclassified as likely benign, but suggest

further characterization is warranted.

***Calibrating functional assays by protein domain***

Often proteins contain multiple domains which contribute distinct functions to the protein. Our

results indicate that when developing functional assays for human proteins in evolutionarily

distant model organisms it is important to calibrate assays for each domain. This can be achieved

through inclusion of calibration variants with known clinical/biochemical designations covering

each domain. MEPC2 is a multi-domain protein and is largely unstructured outside the MBD

domain (Ghosh et al. 2010). We initially focused on identifying calibrating variants in the MBD

and TRD domains, as these domains harbour many of the clinically relevant variants.

The yeast assay only calibrated for the MBD and failed to calibrate for all other domains,

indicating we could only assess functional effects of variants found within the MBD. A likely

explanation is that the MECP2 MBD binds DNA in yeast without the need of accessory factors.

Indeed, Brown *et al.*, show this is likely to be the case (Brown et al. 2025 Mar 18). The TRD

mediates interactions with co-repressor complexes, including Sin3A, N-Cor and REST, and Rett-

causing mutations disrupt these interactions (Lyst et al. 2013). It is likely the assay failed to

calibrate for TRD variants because these complexes are either not present in yeast (N-Cor and

REST), or the TRD did not interact with the yeast ortholog (Sin3A/HDAC). The results of the

MECP2 SIM screen also provided valuable insight into MECP2's function in yeast. The strong

aggravating genetic interactions we observed with genes involved in DNA damage response and

DNA repair pathways (e.g., *RAD50*) suggest MECP2 expression either directly or indirectly

results in DNA damage. These effects of MECP2 likely disrupted the cell cycle (supported by

the interactions with cell cycle genes, e.g., *PHO85*) leading to overall decreased cell growth.

Since variants in the MBD known to disrupt DNA binding failed to inhibit growth, this further

suggested DNA binding by MECP2 contributed to defects in DNA and/or chromatin.

*Drosophila* was successful in stratifying pathogenic and benign calibration variants in both the

MBD and TRD domains, indicating we can make high confidence predictions about test ASD

variants located in these domains. For variants in the MBD, it is likely these disrupt DNA

binding, although DNA binding by MECP2 has yet to be demonstrated in *Drosophila*. For the

TRD, our data suggest MECP2 is able to interact with *Drosophila* orthologs of co-repressor

complexes. First, R306C prevents interaction with co-repressors and was complete LoF in our

assay (Lyst et al. 2013). Second, co-repressor complex mutants suppress phenotypes resulting

from MECP2 over-expression in the *Drosophila* wing (Cukier et al. 2008). For example,

mutation of the Polycomb group co-repressor ortholog of Asxl1, *Asx*, prevents ectopic wing vein

growth resulting from MECP2 expression (Cukier et al. 2008). For the Interdomain and CTD

domain, testing additional pathogenic variants in these domains would increase confidence in our

predictions for test (ASD) variants within these domains. Thus, when developing functional

assays in surrogate model systems with considerable evolutionary distance from humans, it is

important to consider calibrating assays for each domain before making conclusions about the

effects of test variants.

***MECP2 protein stability***

The stability of a handful of MECP2 variants has been studied both using *in vitro* assays and *in*

*vivo* mouse models (Kucukkal et al. 2015; Yang et al. 2016; Sperlazza et al. 2017; Tillotson and

Bird 2020). We were able to compare seven variants located in the MBD between our yeast

results and these studies (Supplemental Table 1). R106W, P152R and T158M are all destabilized

in mouse models and *in vitro* systems, and based on protein levels in yeast were predicted to be

also destabilized in yeast also. R133C is found to be stable in *in vitro* systems and is predicted to

be stable in yeast, but is destabilized in mouse. L100V and P101S are destabilized *in vitro*, but

are predicted to be stable in yeast. R133H is stable *in vitro*, but is predicted to be destabilized in

yeast. Hence, the discrepancies between these data emphasize the value of comparing variant

function across multiple species and assay systems. MECP2 turnover in cells is regulated by

ubiquitylation by the E3 ligase RNF4 (Wang 2014), indicating it may be an *in vivo*-specific

modifier of variant stability. Interestingly, we identified the yeast orthologs of RNF4, *SLX5* and

*SLX8*, as aggravating genetic interactions in our SIM screen (Figure 3B), suggesting MECP2

levels may also be regulated by this E3 ligase in yeast. Thus, regulated turnover *in vivo* is likely

an important factor contributing to MECP2 variant function and may vary across assay systems.

Interaction of MECP2 with DNA and co-repressors also influences MECP2 intrinsic stability

(Ghosh et al. 2010), hence differences in these interactions could be another factor contributing

to differences in MECP2 variant function/stability between assay systems.

***MECP2 and DNA methylation***

It is noteworthy that all pathogenic calibration variants in the MBD were LoF in our functional

assays despite there being no, or extremely low levels of DNA cytosine methylation in yeast and

*Drosophila*, respectively. Our data correlate well with Brown *et al.* (Brown et al. 2025 Mar 18),

showing that MECP2 binds the unmethylated yeast genome and mediates widespread

transcriptional suppression. This raises the possibility that ASD and Rett MECP2 variants may

not be specifically deficient in their ability to bind methylated DNA, but may be generally

deficient in binding to unmethylated DNA and/or chromatin. It is challenging to specify

differential impacts of specific Rett variants on methyl versus non-methyl DNA binding in

models where DNA methylation is present. We propose this may have obscured the impact that

pathogenic variants have on non-methylated DNA binding and its importance in healthy and

disease states, and we propose that the yeast and fly models offer valuable platforms to further

examine these mechanisms.

## DATA AVAILABILITY STATEMENT

Strains and plasmids are available upon request. The authors affirm that all data necessary for

confirming the conclusions of the article are present within the article, figures, tables and

supplemental data files.

## ACKNOWLEDGEMENTS

We thank Dr. Janel Kopp, UBC for access to the Panoramic Midi II digital slide scanner. Susan

Lindquist for the generous gift of the *pAG426GAL-ccdB* vector. Stocks obtained from the

Bloomington *Drosophila* Stock Center (NIH P40OD018537) were used in this study. We thank

*Drosophila* Genomics Resource Center, supported by NIH grant 2P40OD010949 for the *esc*

cDNA. We used FlyBase (releases 2020 through 2024) for general information on genes,

phenotypes and stocks. We would like to thank Gino Laberge at Genome ProLab for the kind

gift of balancer stocks and also their transgenic services (Genome ProLab Inc, Quebec, Canada).

We thank Libby Natola and Guillaume Poirier-Morency (Pavlidis Lab) for contributions to the

variant annotation pipeline. This work was funded by an operating grant from The Simons

Foundation Autism Research Initiative (2021 Genomics of ASD: Pathway to Genetic Therapies:

AWD-021951. SIMOFOUN 2021).

## REFERENCES

Amir RE, Van den Veyver IB, Schultz R, Malicki DM, Tran CQ, Dahle EJ, Philippi A, Timar L, Percy AK, Motil KJ, et al. 2000. Influence of mutation type and X chromosome inactivation on Rett syndrome phenotypes. Ann Neurol. 47(5):670–679.

Amir RE, Van den Veyver IB, Wan M, Tran CQ, Francke U, Zoghbi HY. 1999. Rett syndrome is caused by mutations in X-linked MECP2, encoding methyl-CpG-binding protein 2. Nat Genet. 23(2):185–188. doi:10.1038/13810.

Ananth A, Fu C, Neul JL, Benke T, Marsh E, Suter B, Ferdinandsen K, Skinner SA, Annese F,
Percy AK. 2024. MECP2 Variants in Males: More Common than Previously Appreciated.
Pediatr Neurol. 161:263–267. doi:10.1016/j.pediatrneurol.2024.09.022.

Bähler J. 2005. Cell-Cycle Control of Gene Expression in Budding and Fission Yeast. Annu Rev
Genet. 39(Volume 39, 2005):69–94. doi:10.1146/annurev.genet.39.110304.095808.

Ballestar E, Yusufzai TM, Wolffe AP. 2000. Effects of Rett Syndrome Mutations of the Methyl-
CpG Binding Domain of the Transcriptional Repressor MeCP2 on Selectivity for Association
with Methylated DNA. Biochemistry. 39(24):7100–7106. doi:10.1021/bi0001271.

Belmadani M, Jacobson M, Holmes N, Phan M, Nguyen T, Pavlidis P, Rogic S. 2019. VariCarta:
A Comprehensive Database of Harmonized Genomic Variants Found in Autism Spectrum
Disorder Sequencing Studies. Autism Res Off J Int Soc Autism Res. 12(12):1728–1736.
doi:10.1002/aur.2236.

Bischof J, Maeda RK, Hediger M, Karch F, Basler K. 2007. An optimized transgenesis system
for Drosophila using germ-line-specific phiC31 integrases. Proc Natl Acad Sci U S A.
104(9):3312–3317. doi:10.1073/pnas.0611511104.

Brnich SE, Abou Tayoun AN, Couch FJ, Cutting GR, Greenblatt MS, Heinen CD, Kanavy DM,
Luo X, McNulty SM, Starita LM, et al. 2019. Recommendations for application of the functional
evidence PS3/BS3 criterion using the ACMG/AMP sequence variant interpretation framework.
Genome Med. 12:3. doi:10.1186/s13073-019-0690-2.

Brown JAR, Ling MYM, Ausió J, Howe LJ. 2025 Mar 18. Human MeCP2 binds to promoters
and inhibits transcription in an unmethylated yeast genome. Genetics.:iyaf043.
doi:10.1093/genetics/iyaf043.

Buitrago D, Labrador M, Arcon JP, Lema R, Flores O, Esteve-Codina A, Blanc J, Villegas N,
Bellido D, Gut M, et al. 2021. Impact of DNA methylation on 3D genome structure. Nat
Commun. 12(1):3243. doi:10.1038/s41467-021-23142-8.

Chahrour M, Zoghbi HY. 2007. The Story of Rett Syndrome: From Clinic to Neurobiology.
Neuron. 56(3):422–437. doi:10.1016/j.neuron.2007.10.001.

Cheng T-L, Wang Z, Liao Q, Zhu Y, Zhou W-H, Xu W, Qiu Z. 2014. MeCP2 suppresses nuclear
microRNA processing and dendritic growth by regulating the DGCR8/Drosha complex. Dev
Cell. 28(5):547–560. doi:10.1016/j.devcel.2014.01.032.

Cukier HN, Perez AM, Collins AL, Zhou Z, Zoghbi HY, Botas J. 2008. Genetic Modifiers of
MeCP2 Function in Drosophila. PLoS Genet. 4(9):e1000179. doi:10.1371/journal.pgen.1000179.

Ehrhart F, Jacobsen A, Rigau M, Bosio M, Kaliyaperumal R, Laros JFJ, Willighagen EL,
Valencia A, Roos M, Capella-Gutierrez S, et al. 2021. A catalogue of 863 Rett-syndrome-
causing MECP2 mutations and lessons learned from data integration. Sci Data. 8(1):10.
doi:10.1038/s41597-020-00794-7.

Free A, Wakefield RID, Smith BO, Dryden DTF, Barlow PN, Bird AP. 2001. DNA Recognition
by the Methyl-CpG Binding Domain of MeCP2 *. J Biol Chem. 276(5):3353–3360.
doi:10.1074/jbc.M007224200.

Ganguly P, Madonsela L, Chao JT, Loewen CJR, O'Connor TP, Verheyen EM, Allan DW.
2021. A scalable Drosophila assay for clinical interpretation of human PTEN variants in
suppression of PI3K/AKT induced cellular proliferation. PLoS Genet. 17(9):e1009774.
doi:10.1371/journal.pgen.1009774.

Georgel PT, Horowitz-Scherer RA, Adkins N, Woodcock CL, Wade PA, Hansen JC. 2003.
Chromatin compaction by human MeCP2. Assembly of novel secondary chromatin structures in
the absence of DNA methylation. J Biol Chem. 278(34):32181–32188.
doi:10.1074/jbc.M305308200.

Ghosh RP, Nikitina T, Horowitz-Scherer RA, Gierasch LM, Uversky VN, Hite K, Hansen JC,
Woodcock CL. 2010. Unique physical properties and interactions of the domains of methylated
DNA binding protein 2. Biochemistry. 49(20):4395–4410. doi:10.1021/bi9019753.

Good KV, Vincent JB, Ausió J. 2021. MeCP2: The Genetic Driver of Rett Syndrome
Epigenetics. Front Genet. 12:620859. doi:10.3389/fgene.2021.620859.

Guo H, Wang T, Wu H, Long M, Coe BP, Li H, Xun G, Ou J, Chen B, Duan G, et al. 2018.
Inherited and multiple de novo mutations in autism/developmental delay risk genes suggest a
multifactorial model. Mol Autism. 9(1):64. doi:10.1186/s13229-018-0247-z.

Hansen JC, Ghosh RP, Woodcock CL. 2010. Binding of the Rett syndrome protein, MeCP2, to
methylated and unmethylated DNA and chromatin. IUBMB Life. 62(10):732–738.
doi:10.1002/iub.386.

Harikrishnan KN, Chow MZ, Baker EK, Pal S, Bassal S, Brasacchio D, Wang L, Craig JM,
Jones PL, Sif S, et al. 2005. Brahma links the SWI/SNF chromatin-remodeling complex with
MeCP2-dependent transcriptional silencing. Nat Genet. 37(3):254–264. doi:10.1038/ng1516.

Her Y, Pascual DM, Goldstone-Joubert Z, Marcogliese PC. 2024. Variant functional assessment
in Drosophila by overexpression: what can we learn? Genome. 67(6):158–167. doi:10.1139/gen-
2023-0135.

Jiménez J, Ricco N, Grijota-Martínez C, Fadó R, Clotet J. 2013. Redundancy or specificity? The
role of the CDK Pho85 in cell cycle control. Int J Biochem Mol Biol. 4(3):140–149.

Jones PL, Veenstra GJ, Wade PA, Vermaak D, Kass SU, Landsberger N, Strouboulis J, Wolffe
AP. 1998. Methylated DNA and MeCP2 recruit histone deacetylase to repress transcription. Nat
Genet. 19(2):187–191. doi:10.1038/561.

Kucukkal TG, Yang Y, Uvarov O, Cao W, Alexov E. 2015. Impact of Rett Syndrome Mutations
on MeCP2 MBD Stability. Biochemistry. 54(41):6357–6368. doi:10.1021/acs.biochem.5b00790.

Kuznetsova A, Brockhoff PB, Christensen RHB. 2017. lmerTest Package: Tests in Linear Mixed
Effects Models. J Stat Softw. 82:1–26. doi:10.18637/jss.v082.i13.

Landrum MJ, Lee JM, Benson M, Brown GR, Chao C, Chitipiralla S, Gu B, Hart J, Hoffman D,
Jang W, et al. 2018. ClinVar: improving access to variant interpretations and supporting
evidence. Nucleic Acids Res. 46(D1):D1062–D1067. doi:10.1093/nar/gkx1153.

Leblond CS, Le T-L, Malesys S, Cliquet F, Tabet A-C, Delorme R, Rolland T, Bourgeron T.
2021. Operative list of genes associated with autism and neurodevelopmental disorders based on
database review. Mol Cell Neurosci. 113:103623. doi:10.1016/j.mcn.2021.103623.

Lewis JD, Meehan RR, Henzel WJ, Maurer-Fogy I, Jeppesen P, Klein F, Bird A. 1992.
Purification, sequence, and cellular localization of a novel chromosomal protein that binds to
Methylated DNA. Cell. 69(6):905–914. doi:10.1016/0092-8674(92)90610-O.

Lunyak VV, Burgess R, Prefontaine GG, Nelson C, Sze S-H, Chenoweth J, Schwartz P, Pevzner
PA, Glass C, Mandel G, et al. 2002. Corepressor-Dependent Silencing of Chromosomal Regions
Encoding Neuronal Genes. Science. 298(5599):1747–1752. doi:10.1126/science.1076469.

Lyko F, Ramsahoye BH, Jaenisch R. 2000. DNA methylation in Drosophila melanogaster.
Nature. 408(6812):538–540. doi:10.1038/35046205.

Lyst MJ, Ekiert R, Ebert DH, Merusi C, Nowak J, Selfridge J, Guy J, Kastan NR, Robinson ND,
de Lima Alves F, et al. 2013. Rett syndrome mutations abolish the interaction of MeCP2 with the
NCoR/SMRT co-repressor. Nat Neurosci. 16(7):898–902. doi:10.1038/nn.3434.

McKnight D, Bean L, Karbassi I, Beattie K, Bienvenu T, Bonin H, Fang P, Chrisodoulou J, Friez
M, Helgeson M, et al. 2022. Recommendations by the ClinGen Rett/Angelman-like expert panel
for gene-specific variant interpretation methods. Hum Mutat. 43(8):1097–1113.
doi:10.1002/humu.24302.

McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GRS, Thormann A, Flicek P, Cunningham F.
2016. The Ensembl Variant Effect Predictor. Genome Biol. 17(1):122. doi:10.1186/s13059-016-
0974-4.

Ng HH, Bird A. 1999. DNA methylation and chromatin modification. Curr Opin Genet Dev.
9(2):158–163. doi:10.1016/s0959-437x(99)80024-0.

Nomura Y. 2005. Early behavior characteristics and sleep disturbance in Rett syndrome. Brain
Dev. 27:S35–S42. doi:10.1016/j.braindev.2005.03.017.

Pascual-Alonso A, Martínez-Monseny AF, Xiol C, Armstrong J. 2021. MECP2-Related
Disorders in Males. Int J Mol Sci. 22(17):9610. doi:10.3390/ijms22179610.

Post KL, Belmadani M, Ganguly P, Meili F, Dingwall R, McDiarmid TA, Meyers WM,
Herrington C, Young BP, Callaghan DB, et al. 2020. Multi-model functionalization of disease-
associated PTEN missense mutations identifies multiple molecular mechanisms underlying
protein dysfunction. Nat Commun. 11(1):2073. doi:10.1038/s41467-020-15943-0.

Rodriguez JM, Maietta P, Ezkurdia I, Pietrelli A, Wesselink J-J, Lopez G, Valencia A, Tress
ML. 2013. APPRIS: annotation of principal and alternative splice isoforms. Nucleic Acids Res.
41(D1):D110–D117. doi:10.1093/nar/gks1058.

Schaaf CP, Sabo A, Sakai Y, Crosby J, Muzny D, Hawes A, Lewis L, Akbar H, Varghese R,
Boerwinkle E, et al. 2011. Oligogenic heterozygosity in individuals with high-functioning autism
spectrum disorders. Hum Mol Genet. 20(17):3366–3375. doi:10.1093/hmg/ddr243.

Skene PJ, Illingworth RS, Webb S, Kerr ARW, James KD, Turner DJ, Andrews R, Bird AP.
2010. Neuronal MeCP2 is expressed at near histone-octamer levels and globally alters the
chromatin state. Mol Cell. 37(4):457–468. doi:10.1016/j.molcel.2010.01.030.

Sperlazza MJ, Bilinovich SM, Sinanan LM, Javier FR, Williams DC. 2017. Structural Basis of
MeCP2 Distribution on Non-CpG Methylated and Hydroxymethylated DNA. J Mol Biol.
429(10):1581–1594. doi:10.1016/j.jmb.2017.04.009.

Syed A, Tainer JA. 2018. The MRE11–RAD50–NBS1 Complex Conducts the Orchestration of
Damage Signaling and Outcomes to Stress in DNA Replication and Repair. Annu Rev Biochem.
87(Volume 87, 2018):263–294. doi:10.1146/annurev-biochem-062917-012415.

Symington LS. 2002. Role of *RAD52* Epistasis Group Genes in Homologous Recombination and
Double-Strand Break Repair. Microbiol Mol Biol Rev. 66(4):630–670.
doi:10.1128/MMBR.66.4.630-670.2002.

Tillotson R, Bird A. 2020. The Molecular Basis of MeCP2 Function in the Brain. J Mol Biol.
432(6):1602–1623. doi:10.1016/j.jmb.2019.10.004.

Tong AH, Evangelista M, Parsons AB, Xu H, Bader GD, Pagé N, Robinson M, Raghibizadeh S,
Hogue CW, Bussey H, et al. 2001. Systematic genetic analysis with ordered arrays of yeast
deletion mutants. Science. 294(5550):2364–2368. doi:10.1126/science.1065810.

Vonhoff F, Williams A, Ryglewski S, Duch C. 2012. Drosophila as a model for MECP2 gain of
function in neurons. PloS One. 7(2):e31835. doi:10.1371/journal.pone.0031835.

Wan M, Lee SSJ, Zhang X, Houwink-Manville I, Song H-R, Amir RE, Budden S, Naidu S,
Pereira JLP, Lo IFM, et al. 1999. Rett Syndrome and Beyond: Recurrent Spontaneous and
Familial MECP2 Mutations at CpG Hotspots. Am J Hum Genet. 65(6):1520–1529.
doi:10.1086/302690.

Wang T, Guo H, Xiong B, Stessman HAF, Wu H, Coe BP, Turner TN, Liu Y, Zhao W,
Hoekzema K, et al. 2016. De novo genic mutations among a Chinese autism spectrum disorder
cohort. Nat Commun. 7(1):13316. doi:10.1038/ncomms13316.

Wang Y. 2014. RING finger protein 4 (RNF4) derepresses gene expression from DNA
methylation. J Biol Chem. 289(49):33808–33813. doi:10.1074/jbc.C114.611558.

Wen Z, Cheng T-L, Li G-Z, Sun S-B, Yu S-Y, Zhang Y, Du Y-S, Qiu Z. 2017. Identification of
autism-related MECP2 mutations by whole-exome sequencing and functional validation. Mol
Autism. 8:43. doi:10.1186/s13229-017-0157-5.

Williams AA, Mehler VJ, Mueller C, Vonhoff F, White R, Duch C. 2016. Apoptotic Activity of
MeCP2 Is Enhanced by C-Terminal Truncating Mutations. PLoS ONE. 11(7):e0159632.
doi:10.1371/journal.pone.0159632.

Williams AA, White R, Siniard A, Corneveaux J, Huentelman M, Duch C. 2016. MECP2
impairs neuronal structure by regulating KIBRA. Neurobiol Dis. 91:284–291.
doi:10.1016/j.nbd.2016.03.019.

Yang Y, Kucukkal TG, Li J, Alexov E, Cao W. 2016. Binding Analysis of Methyl-CpG Binding
Domain of MeCP2 and Rett Syndrome Mutations. ACS Chem Biol. 11(10):2706–2715.
doi:10.1021/acschembio.6b00450.

Young BP, Loewen CJR. 2013. Balony: a software package for analysis of data generated by
synthetic genetic array experiments. BMC Bioinformatics. 14:354. doi:10.1186/1471-2105-14-
354.

Young BP, Post KL, Chao JT, Meili F, Haas K, Loewen C. 2020. Sentinel interaction mapping -
a generic approach for the functional analysis of human disease gene variants using yeast. Dis
Model Mech. 13(7):dmm044560. doi:10.1242/dmm.044560.

Yusufzai TM, Wolffe AP. 2000. Functional consequences of Rett syndrome mutations on human
MeCP2. Nucleic Acids Res. 28(21):4172–4179.

**AUTHOR CONTRIBUTIONS**

DWA, CJRL, BPY, SMG, JTC, MD, SR, MD, PP contributed to conception of the work.

DWA, CJRL, GM, EC, BPY, SI, TL, JL, JBL, SMG, BB, MD, JS, SR, PP contributed to design

of the work.

DWA, CJRL, GM, EC, BPY, JBL, LM, JS, JBL, SR, PP contributed to acquisition, analysis, or

interpretation of data.

DWA, SR, PP, GM, EC, JBL, MD, KKC, BB, BW contributed to creation of new software used.

DWA, CJRL, JS, BPY, JTC, JBL, GM, EC, LM, JL, JBL, JTC, SR, PP contributed to drafting

the paper or substantively revised it.

**FIGURE CAPTIONS**

**Figure 1:** The forty-six MECP2 variants examined in this study. Domain locations of variants as well as variant class, variant inheritance, and whether they are found in Rett Syndrome and/or ASD. NTD – N-terminal domain, MBD – Methyl-CpG-binding domain, ID – Interdomain, TRD – Transcription repressor domain, CTD – C-terminal domain.

**Figure 2:** MECP2 expression and effect on growth in yeast. (A) Untagged MECP2-ref expressed from a 2-micron plasmid utilizing the estradiol-responsive promoter at various estradiol concentrations in WT yeast. Equivalent amounts of yeast were loaded into each lane and western blotted with anti-MECP2 antibody. M – molecular weight markers, arrowhead – full length MECP2. (B) Effect of MECP2-ref expression on growth of WT yeast from a plasmid utilizing the estradiol-responsive promoter at various estradiol concentrations. +EV – empty vector in wild type yeast, +MECP2-ref – MECP2-ref plasmid in wild type yeast. (C) Similar to B except using a 2-micron plasmid containing a galactose-responsive promoter and grown in galactose. **** – indicates p value < 0.0001. ns – not significant.

**Figure 3:** Results of SIM genetic interaction screen with MECP2-ref. (A) Functional enrichment for genetic interactions is shown where nodes indicate enriched GO terms and edges indicate statistically significant associations between terms. Node size and edge thickness indicate statistical significance (minimum $p < 0.05$, Bonferonni corrected). Highlighted labels indicate most significant terms within networks. Network generated using Cytoscape and the ClueGO plug-in. (B) Effect of MECP2-ref on yeast growth in mutants from the GO categories "DNA

damage response", "Chromatin organization" and "DNA binding" that were identified in the

SIM screen. Normalized colony size indicates the size of the colonies on the array plates after

accounting for the effect of MECP2-ref expression in WT yeast, in non-inducing (0 nM

estradiol) and inducing (10 nM estradiol) conditions.

**Figure 4:** Analysis of 46 MECP2 variants using the yeast assay. (A) Estimated functional effects

of MECP2 variants expressed in WT yeast. Data is normalized to MECP2-ref (=1) and empty

vector (=0). Error bars indicate 95% confidence intervals. (B) Protein stability of MECP2

variants in WT yeast. Protein levels were normalized to MECP2-ref and plotted on a log scale.

Error bars indicate S.D. (n = 3). * indicates $p < 0.05$, ** indicates $p < 0.01$. Colour coding same

as in (A). All variants are located in MBD except T228S (TRD), S356G (CTD) and P388S

(CTD).

**Figure 5:** *Drosophila* wing phenotypes induced by MECP2 variants. Representative images of

wings are shown expressing the indicated controls and variants. *attp40* – no MECP2, *MECP2-ref*

– WT MECP. All images are the same scale. Arrows indicate ectopic vein tissue for *UAS-*

*MECP2-ref* only.

**Figure 6:** Analysis of MECP2 variant wing vein phenotypes. We show our quantitation of total

wing vein per wing area for each genotype, normalized to *MECP2-ref* (WT MECP2) = 1 and

*attp40* (no MECP2) = 0. Error bars indicate 95% confidence intervals.

## SUPPLEMENTAL FIGURE CAPTIONS

**Supplemental Figure 1**: Representative MECP2 Western blots. Arrow indicates the full-length MECP2 protein. Antibody labelling is as follows: red - anti-MECP2, green - anti-PGK1.

**Supplemental Figure 2**: Representative wing phenotypes for control, benign and pathogenic MECP2 calibration variants tested in *Drosophila*. For each variant, raw images are on the left and the masked image used to quantitate total wing vein per wing is on the right.

**Supplemental Figure 3**: Representative wing phenotypes for test (ASD) MECP2 variants tested in *Drosophila*. For each variant, raw images are on the left and the masked image used to quantitate total wing vein per wing is on the right.

## SUPPLEMENTAL METHODS CAPTIONS

**Supplemental Methods 1:** Detailed outline of wing vein quantitation methods.

**Supplemental Methods 2:** Yeast statistical model R script.

**Supplemental Methods 3:** Fly statistical model R script.

## SUPPLEMENTAL TABLE CAPTIONS

**Supplemental Table 1:** MECP2 protein stability summary table.

## SUPPLEMENTAL DATA CAPTIONS

**Supplemental Data 1:** Detailed information for each variant, primers used in cloning, the molecular genetic reagents made for each organism, and the *Drosophila* stocks generated.

**Supplemental Data 2:** Yeast SIM screen raw data.

**Supplemental Data 3:** GO analysis results for yeast SIM screen listing genes in each GO term.

**Supplemental Data 4:** Raw data for yeast variant assays used in statistical model.

**Supplemental Data 5:** Raw data for *Drosophila* variant assays used in statistical model.

## Figure 1

# Figure 2

A

B

C

# Figure 3

A

organelle fission

meiotic cell cycle

meiosis I cell
cycle process

mitotic
recombination

homologous
recombination

DNA recombination

telomere maintenance

recombinational
repair

double-strand break repair via
homologous recombination

DNA metabolic
process

**cellular response to
stress**

nuclear
chromosome

DNA repair

double-strand
break repair

**negative regulation of nitrogen
compound metabolic process**

**negative regulation
of cellular metabolic
process**

**Golgi apparatus
subcompartment**

**DNA damage
response**

telomere organization

chromosome organization

chromosome

chromatin organization

**DNA binding**

chromatin remodeling

protein-DNA complex
organization

protein-containing
complex organization

establishment of
cell polarity

mitotic cell cycle

mitotic cell cycle process

**cell cycle process**

regulation of cell cycle

B

Normalized colony size

**β**-estradiol   ● 0 nM   ■ 10 nM

WT, ade12Δ, bck1Δ, cla4Δ, ctf4Δ, dia2Δ, doc1Δ, esc2Δ, gas1Δ, gcn2Δ, gcn5Δ, hht2Δ, hir1Δ, hur1Δ, ies1Δ, mck1Δ, mit1Δ, mms4Δ, msn4Δ, ngg1Δ, nhp10Δ, nup170Δ, nup84Δ, pho85Δ, rad50Δ, rad51Δ, rad52Δ, rad54Δ, rlf2Δ, rmi1Δ, rpb9Δ, rtg1Δ, rtg3Δ, rtt106Δ, sae2Δ, scs2Δ, set2Δ, sfl1Δ, sfp1Δ, sif2Δ, slx5Δ, slx8Δ, smi1Δ, snf1Δ, srs2Δ, sub1Δ, swi4Δ, swi6Δ, ume6Δ, wss1Δ, xrs2Δ, yer134cΔ, yox1Δ

● DNA damage response   ● Chromatin organization   ● DNA binding

# Figure 4

# Figure 5

100 μm

# Figure 6

# Supplemental Figure 1

# Supplemental Figure 2

**CONTROLS**

*attP40*          *MECP2-ref*

**BENIGN**

*H52R*          *D121E*

*T196S*          *T203M*

*T228S*          *S356G*

*E394K*          *R458H*

**PATHOGENIC**

*L100V*          *P101S*

*R133C*          *K135E*

*P152R*          *T158M*

*P225R*          *R306C*

# Supplemental Figure 3

**TEST (ASD)**

Supplemental Table 1: Summary of MECP2 protein stability results

| Variant | Yeast* (this study) | Yang [1] (in vitro, WT = 1) | Kucukkal [2] (in vitro) ($\Delta\Delta G$) | Sperlazza [3] (NMR) (+ slight, ++ more destabilized) | Tillotson [4] (mouse models, WT = 1) |
|---|---|---|---|---|---|
| R91W | 1 | | | | |
| L100V | 1 | 0.5 | +1.25 | | |
| P101S | 1 | | | + | |
| R106W | 0.37 | 1 | +0.1 | ++ | 0.1 (TAVI) |
| D121E | 1 | | | | |
| R133C | 1 | 1 | +0.4 | | 0.6 (eGFP) |
| R133H | 0.56 | 1 | +0.25 | | |
| K135E | 0.50 | | | | |
| T148A | 0.65 | | | | |
| P152L | 0.65 | | | | |
| P152R | 0.41 | 0.25 | +1.5 | | |
| T158M | 0.48 | 1 | +0.25 | + | 0.3 (eGFP), 0.25 (TAVI) |
| G163W | 0.66 | | | | |
| P225R | | | | | 0.22 |
| T228S | 1 | | | | |
| R306C | | | | | 1.0 |
| P322L | | | | | 0.03 |
| S356G | 1.54 | | | | |
| P388S | 0.25 | | | | |

* Values not significantly different than MECP2-Ref are set to 1, all others are significant (p<0.05)

References:
1. Yang, Y., Kucukkal, T. G., Li, J., Alexov, E. & Cao, W. Binding Analysis of Methyl-CpG Binding Domain of MeCP2 and Rett Syndrome Mutations. *ACS Chem. Biol.* **11**, 2706–2715 (2016).
2. Kucukkal, T. G., Yang, Y., Uvarov, O., Cao, W. & Alexov, E. Impact of Rett Syndrome Mutations on MeCP2 MBD Stability. *Biochemistry* **54**, 6357–6368 (2015).
3. Sperlazza, M. J., Bilinovich, S. M., Sinanan, L. M., Javier, F. R. & Williams, D. C. Structural Basis of MeCP2 Distribution on Non-CpG Methylated and Hydroxymethylated DNA. *J. Mol. Biol.* **429**, 1581–1594 (2017).
4. Tillotson, R. & Bird, A. The Molecular Basis of MeCP2 Function in the Brain. *J. Mol. Biol.* **432**, 1602–1623 (2020).