TITLE

A multi-model approach defines function altering MECP2 missense variants identified in

individuals with autism spectrum disorder.

AUTHOR LIST

Eric Chen[1]*, Jessica Schmitt[1]*, Graeme McIntosh[1]*, Ajay Singh[2]*, Barry Young[1], Tianshun

Lian[1], Jie Liu[1], Vedanta Khan[2], Ke Chen[3,] J Beatrice Liston[1], Lily MacDonald[1], Bill Wang[1],

Sonia Medina Giro[1], Benjamin Boehme[1], Mriga Das[1], Seevasant Indran[1], Sanja Rogic[3], Paul

Pavlidis[3], Jesse T Chao[2], Douglas W Allan[1]#, and Christopher JR Loewen[1]#

[1]Department of Cellular and Physiological Sciences, The Life Sciences Institute, The University

of British Columbia, Vancouver, BC, Canada

[2]Department of Medical Biophysics, Sunnybrook Research Institute, The University of Toronto,

Toronto, ON, Canada

[3]Department of Psychiatry, The Michael Smith Institute, The University of British Columbia,

Vancouver, BC, Canada

(co-first *, co-corresponding **#)**

ABSTRACT

MECP2 is commonly mutated in Rett syndrome, where MECP2's function as a DNA cytosine

methylation reader is believed critical. MECP2 variants are also catalogued in individuals with

autism spectrum disorder (ASD), including nine missense variants with no known clinical

significance. To assess these nine as risk alleles for ASD, we developed MECP2 variant function

assays using yeast, *Drosophila* and human cell lines. We calibrated these assays with known

reference pathogenic and benign variants. Our data predict that four ASD variants are loss of

function (LoF) and five are functional. Protein destabilization or nuclear delocalization offers

insight into the altered function of a number of these variants. Notably, yeast

and *Drosophila* lack DNA methylation, yet all Rett reference pathogenic and ASD variants in the

methyl DNA binding domain that we analyzed proved to be LoF, suggesting a clinically-relevant

role for non-methyl DNA-binding by MECP2.

INTRODUCTION

The X-linked protein MECP2 was first identified in screens for proteins recruited to methylated DNA [1], and LoF alleles of MECP2 were subsequently found as causative in the neurodevelopmental disorder, Rett syndrome [2]. Since then, variants in MECP2 have been found in most cases of Rett, but also in other neurodevelopmental disorders including autism spectrum disorder (ASD) [3–6]. MECP2 has largely been examined as a DNA cytosine methylation reader via its methyl cytosine binding domain (MBD) that recruits co-repressor complexes to DNA to repress transcription via its transcriptional repressor domain (TRD) [7–10]. The MBD, and to a lesser extent the TRD, harbors most Rett LoF variants [11], resulting in variants which are believed to reduce binding to methylated DNA or reduce recruitment of co-regulator complexes, respectively [12,13]. While numerous MECP2 variants have been found in patients that exhibit Rett with ASD, recent reports have annotated six MECP2 missense variants in ASD without an associated Rett diagnosis (R91W, P152L, G195S, A202G, E282G, S411R) [14–18]. It is unclear whether any of these variants are function altering and clinically significant. Two of them, R91W and A202G, are classified by ClinVar as variants of uncertain significance (VUS) in cases of severe neonatal onset encephalopathy with microcephaly, and E282G is classified as likely pathogenic. The remaining four variants have no current ClinVar classification.

While MECP2 is understood to primarily be a methyl-cytosine DNA (DNAm) binding protein, it has become clear that it also binds unmethylated DNA with a preference for GC-rich sequences [19–22]. This has been proposed as an alternate hypothesis for why MECP2 appears to preferentially bind methylated GC-rich regions of the genome, as opposed to binding DNAm methylation *per se*. It can be challenging to segregate DNAm-dependent from DNAm-independent functions of

MeCP2 *in vivo*, as DNAm is both abundant and essential in mammals. Consequently, it is still

unclear what fraction of MeCP2's functions are dependent on its interaction with methylated

and/or unmethylated DNA. Such ambiguity leaves unresolved the question of whether disease-

relevant variants in MECP2 may selectively disrupt binding to methylated versus unmethylated

DNA.

There are extremely low levels of methylated cytosine in the *Drosophila* genome, and none in

Saccharomyces cerevisiae [23,24]. Thus, *Drosophila* and yeast offer avenues to explore the

interactions of MECP2 with unmethylated DNA, and to test if Rett and ASD variants alter

MECP2 binding to unmethylated DNA. Moreover, while neither organism has an ortholog of

MECP2, both organisms are robust genetic models that have proven well-suited to comparing the

function of large numbers of gene variants [25–27]. In *Drosophila*, human MECP2 associates with

chromatin and mediates transcriptional repression via interactions with orthologs of known

MECP2 physical interactors in humans, including Sin3A, N-Cor, REST as well as other

chromatin modifiers [28]. In fly neurons, MECP2 impacts dendritic branching and behavioural

output [29,30] and C-terminal truncation mutants (R294X specifically) promotes neuronal apoptosis

[31]. In yeast, an accompanying article shows that MECP2 binds chromatin and mediates

widespread transcriptional repression (Brown et al.).

Our goal was to use the *Drosophila* and budding yeast models to test the functional impact of

these ASD MECP2 variants to determine whether they may confer autism risk, by virtue of their

altered function. Further, we complemented these functional studies with molecular

characterization of MECP2 variant stability and nuclear localization in human cell lines. We

calibrated our assays with MECP2 variants classified by ClinVar as pathogenic or benign, in

accordance with ACMG guidelines for experimental assays [32]; first to validate our assays for

variant assessment, and second to provide a rubric for determining if test variants are likely

pathogenic or likely benign. All reference variants functioned as predicted in the *Drosophila*

assay, indicating that we could functionally assess any test variants. In yeast, reference benign

variants functioned as predicted, but only reference pathogenic variants in the MBD functioned

as predicted; thus we consider the yeast assay to be exclusively predictive for test variants in the

MBD. Our analyses indicate that four of the nine test ASD variants are LoF, making these

variants candidate risk alleles for ASD. Interestingly, these variants mostly proved to be less

severe LoF than Rett pathogenics, providing a possible explanation for how MECP2 mutations

can lead to ASD but not Rett.

METHODS

MECP2 isoform selection

MECP2 is known to have two major isoforms, E1 and E2. We selected to work with MECP2

isoform E2 (ENST00000303391, NM_004992) based on the recommendation by the ClinGen

Rett/Angelman-like Expert Panel [33]. This transcript is annotated as the APPRIS principal isoform

in Ensembl genome browser, based on the annotations for protein structure, function and cross-

species conservation, and is the highest expressed transcript in the brain (GTEx portal).

Variant aggregation, annotation and prioritization

We utilized our in-house computational pipeline to aggregate MECP2-E2 missense variants from

ClinVar and VariCarta and perform comprehensive annotation. We collected ClinVar variants

(downloaded in January 2022) that have at least one gold review star, i.e. the review status

(CLNREVSTAT) has to have one of the following designations: "criteria provided, single

submitter", "criteria provided, conflicting interpretations", "criteria provided, multiple

submitters, no conflicts" or "reviewed by expert panel". For each of these variants, we also

obtain its interpretation (clinical significance; CLNSIG) and reported condition (disease name;

CLNDN). The ASD missense variants were obtained from VariCarta, a database of variants

found in individuals diagnosed with ASD and reported in the scientific literature, that we created

[34]. We annotated the aggregated variants using the Ensembl VEP tool [35]. We also added protein

domain annotation relevant in the context of Rett Syndrome [36].

MECP2-E2 cloning

A *MECP2-E2* cDNA clone was obtained (Horizon, Clone Id: 3956518, # MHS6278-

202830559), and MECP2 coding sequences were amplified with iProof DNA polymerase (Bio-

Rad, California USA, #1725301) using the following primers:

GA-attL1-MECP2-522-F (tgtacaaaaaagcaggctccggcaccatggtagctgggatgttag)

GA-attL1-MECP2-522-R (ttgtacaagaaagctgggtctatcagctaactctctcggtc)

Amplicons were cloned in PCR8-GW-TOPO (Invitrogen, ThermoFisher Scientific, Waltham,

USA, catalog #K250020) using NEBuilder HiFi DNA Assembly Master Mix (New England

Biolabs, USA, # M5520). The resulting construct was named *MECP2-pENTRY*. A sequence of

CCGGCACC was introduced in primer GA-attL1-MECP2-522-F (underlined) preceding the

start codon for *MECP2-E2*. After LR cloning of MECP2-pEntry into pDEST vectors, this

CCGGCACC sequence can function as a Kozak sequence to initiate translation, or as an in-

frame linker for N-terminal peptide sequences. Missense variants were generated in *MECP2-*

pENTRY either using the Q5 mutagenesis kit (New England Biolabs, USA, # E0554) in house, or

were made by GenScript (New Jersey, USA). For MECP2 expression clone construction,

MECP2-pENTRY plasmids were subcloned in *pAG426GAL-ccdB* (a kind gift from a gift from

Susan Lindquist; Addgene plasmid # 14155), *pGW-HA.attB* [37] or *pDEST-CMV-N-EGFP*

(Addgene Plasmid #122842) respectively using the Gateway LR Clonase reaction (Thermo

Fisher Scientific, Waltham, USA, catalog # 11791020). All transgenic *UAS-MECP2* variants (on

pGW-HA.attB) *Drosophila* strains used were verified by Sanger sequencing (UBC NAPS,

Vancouver, Canada) or whole amplicon sequencing (Plasmidsaurus, CA, USA) (Supplemental

Data 1). All HEK variants (on *pDEST-CMV-N-EGFP*) were verified by Sanger (UBC NAPS,

Vancouver, Canada) or whole plasmid sequencing (Plasmidsaurus, CA, USA) (Supplemental

Data 1).

Synthetic dosage lethality screening in yeast

Synthetic dosage lethality screening of MECP2 was performed as described previously [27], with

the following modifications. Yeast strain Y7092 was transformed with plasmid pSI7338-

MECP2. Following mating of this strain with the yeast deletion collection, diploid selection,

sporulation and *MATa* haploid selection, MECP2 expression was induced by plating on media

containing 10 nM beta-estradiol. After incubation for one day at 30 °C, these plates were

replicated on identical media. Plates were imaged and analyzed the following day. Control plates

were generated by replicating haploid cells on the same media in the absence of beta-estradiol.

Plate images were analyzed using *Balony* software as previously described [38].

Yeast quantitative liquid growth assays

Yeast strain JHY716 was transformed with various plasmids containing MECP2 variants placed

downstream of the *GAL1* promoter. Four independent clones were picked from transformation

plates and transferred to SD-URA media containing 2% raffinose, 2% galactose to induce

expression of MECP2. For each clone, a 2 mL culture in SD-URA with 2% raffinose, 2%

galactose was grown overnight at 30 °C in a shaking incubator. The following morning, cultures

were diluted to an OD600 of ~0.25 in the same media. After 6 hours of incubation at 30 °C, log

phase cultures were diluted to an OD600 of 0.125. Three aliquots of 200 µl were taken from each

culture and transferred to a well of a flat-bottomed 96-well dish. Growth of cultures was

measured overnight at 30 °C with shaking at 800 rpm in an Agilent Logphase 600 plate reader.

All plates contained strains expressing wild type MECP2 and the corresponding empty vector as

controls. The exponential growth constant k was determined for each well by fitting to an

exponential growth equation during the exponential phase (typically between 5 and 10 hours post

dilution).

Protein abundance assays

To determine the abundance of MECP2 variants in yeast, strains were grown in SD-URA media

containing 2% raffinose to early log phase. MECP2 expression was induced by the addition of

2% galactose to the growth medium. Cells were harvested three hours later by centrifugation and

stored at -80 °C for subsequent processing. Cell pellets were resuspended in Laemmli sample

buffer, and heated to 95 C for 5 minutes. Cell lysis was achieved by the addition of 50 µl acid-

washed glass beads followed by two rounds of disruption in a MP Biomedical FastPrep-24

homogenizer (maximum speed for 60 s with 5 minutes cooling between cycles). Samples (0.2

OD600 equivalents) were separated by SDS-PAGE on Bio-Rad 4-15% pre-cast gels and

transferred to nitrocellulose membranes. MECP2 was immunoblotted using Sigma antibody

(M9317), with PGK1 (Abcam AB113687) used as a loading control. For visualization,

secondary antibodies conjugated to Cy3 (MECP2) and AlexaFluor-488 (PGK1) were used and

imaged using a Bio-Rad ChemiDoc MP Imaging System. MECP2 band volume was normalized

to PGK1 band volume for each variant.

Drosophila **strains**

All flies were maintained on standard cornmeal food at 25C and 70% humidity unless otherwise

specified. *P{GAL4}A9* (BL8761) (referred to as *A9-GAL4*) and *Asx[XF23]*(BL6041) were

obtained from *Bloomington* Drosophila Stock Centre (Indiana, USA). *UAS-MECP2* reference

(*UAS-MECP2-ref*) and variant sequences (collectively termed *UAS-MECP2.#*) were integrated

into embryos of genotype *y[1] w[] P{y[+7.7]=nos-phiC31\int.NLS}X;*

P{y[+t7.7]=CaryP}attP40 (referred to as attP40) by Genome ProLab (Quebec, Canada), and

maintained as w[1118]; P{y[+t7.7] w[+mC]=UAS-MECP2.#.GW.}*attP40**, where # represents

the variant and * represents integrant line number (Supplemental Data 1).

Crossing scheme

Experimenters were double blinded to genotype throughout. Homozygous *UAS-MECP2* males

and a control *attP40* line were crossed to *A9-GAL4* virgins (ratio 4 males : 12 females), and

maintained at room temperature (~21C) for 72 hours. Crosses were then transferred into fresh

'assay' vials, and transferred each 24 hrs each for three days, generating three 'assay' vials per

cross. After removal of adults from each vial, vials were placed into a 29C incubator (70%

humidity). Three days after eclosion, adult flies were sorted by sex and placed in pure

isopropanol at -20C. All transgenic flies for comparison were tested in a single batch.

Wing imaging

Experimenters were double blinded to genotype throughout. Flies were removed from

isopropanol, dried on paper towels, and 10-25 left wings were removed per genotype per assay

vial (one wing per fly). Wings were mounted on microscope slides in Canadian Balsam Fir

(1.01691.0025, CAS-No: 8007-47-4, Sigma Aldrich, MA, USA), coverslipped and sealed with

nail polish. Slides were compressed while drying for 24 hrs. Mounted wings were scanned using

a Panoramic Midi II digital slide scanner (3DHISTECH, Budapest, Hungary). Scanned fly wing

images were cropped at 15x zoom using CaseViewer (3DHISTECH, Budapest,Hungary). Images

were imported into Imaris v10.0 software (Bitplane, MA, USA) for measurement of wing vein

area per wing area, using Labkit machine learning and Fiji software (University of Wisconsin-

Madison, USA). In Labkit, we developed "BIGSMOOTHCLASSYBOI", a machine learning

algorithm to automate measurement of wing vein area for calculation of wing vein area/wing

area (details provided in Supplemental Methods 1).

Statistical analysis of variant effects

Yeast growth constants (k) and fly relative wing vein area data were analyzed using methods

similar to Post et al. [26]. Briefly, linear mixed-effects models were fit to each data set using the

lmerTest R package [39] to estimate the phenotypic effects of each variant compared to the

MECP2-ref constructs, treating experimental batches as random effects. For visualization, values

were standardized to the range [0-1] where 0 represents the mean phenotype measured for an

empty vector, and 1 represents the mean phenotype with the *MECP2-ref* construct. Four

biological replicates, each with three technical replicates were performed for yeast variant

growth assays, while for *Drosophila* wing assays 10-25 wings each from a different animal were

imaged per genotype.

HEK cell culture and variant transfection

HEK 293T cells were maintained in Dulbecco's Modified Eagle Medium (DMEM)

supplemented with 10% fetal bovine serum (FBS) and 1% penicillin-streptomycin in a

humidified incubator at 37°C with 5% CO_2. To transfect variants, cells were seeded in 24-well,

glass bottom plates (CellVis) and grown to ~80% confluency. Each MECP2E2 variant plasmid

was transfected using Lipofectamine 2000 (Thermo Fisher Scientific) according to the

manufacturer's protocol. 24 hours post-transfection, cells were fixed with 4% paraformaldehyde

and stained with 4',6-diamidino-2-phenylindole (DAPI) nuclear stain for 5 minutes. Prolong

Diamond Antifade mountant (Thermo Fisher Scientific) was added to each well and coverslips

were placed on top. The plates were then given 24 hours to set. Each variant was tested in three

separate transfections.

HEK cell microscopy

The plates were imaged using the Olympus scanR high-content screening microscope. For image

acquisition, the objective (20x) was focused on the centre plane of the DAPI channel, and 5 Z-

stack images were captured at a step width of 0.6 μm. Deconvolution was performed for each

image field and the maximum projection was used for quantification. Image analysis, including

nuclear and cytoplasmic segmentations, was carried out using the scanR software. We developed

a "cyto-ring assay" which uses a U-Net convolutional neural network to segment the nuclear

compartment based on DAPI staining, and calculate the cytoplasmic compartment based on the

nuclear segmentation. Two features, including protein abundance and protein translocation, were

extracted after segmentation from GFP-expressing cells. Protein abundance was calculated by

adding the total GFP intensity in the nuclear and cytoplasmic compartment. Protein translocation

was calculated as the ratio of GFP intensities within the nucleus divided by that in the cytoplasm.

Measurement for each variant was normalized to that of WT on the corresponding plate.

HEK cell data analysis

Statistical analysis was performed using GraphPad Prism 10. Multiple unpaired T-tests followed

by Holm-Sidak Correction were used to compare the mean values of protein abundance of each

variant to that of WT, as well as the protein translocation ratio of each variant to WT before

normalization.

RESULTS

Selection of variants

We selected 'reference' benign and pathogenic variants with clear clinical classifications to

calibrate our functional assays, and nine test (ASD) variants (Figure 1, Supplemental Data 1).

We focused on ClinVar variants with "pathogenic", "likely_pathogenic", "benign" and

"likely_benign" clinical interpretation as reference variants. Priority was given to those with

either "reviewed by expert panel" or "criteria provided, multiple submitters" review status. A

few exceptions were made. D121E, a single-submitter benign variant, and T148A, originally

classified as benign, but now listed as conflicting were selected because they were the only

benign variants found in the MBD. When selecting reference pathogenic variants, in addition to

using review status as a criterion, we also prioritized the ones with "Rett syndrome" condition

annotation as well as the ones that have been frequently observed in Rett patients [36]. In total, we

selected 37 reference variants, 10 benign and 27 pathogenic. We selected all VariCarta MECP2

missense variants as ASD variants as long as they didn't have a clear clinical interpretation in

ClinVar, for the total of 9 test ASD variants, with the goal to determine the functional

consequences of these variants. Since the time of our initial selection, four test ASD variants

have been given updated classifications in ClinVar, P217L and E282G becoming likely

pathogenic, and P376S and P388S becoming benign.

MECP2 produces a slow growth phenotype in yeast

MECP2-ref was cloned into a plasmid designed to give controlled expression in response to

addition of estradiol. Using an anti-MECP2 antibody we monitored MECP2 protein levels by

Western blot in response to increasing estradiol (Figure 2A). In the no-estradiol condition we

detected low levels of MECP2 protein, which increased with increasing estradiol and then

saturated at ~10nm. At the highest expression level, MECP2 levels were comparable to a more

conventional vector employing the TDH3 promoter in a multicopy 2 micron vector (Figure 2A).

Next we measured the growth rate of WT yeast containing either the MECP2-ref plasmid or an

empty vector control in response to estradiol addition (Figure 2B). Compared to empty vector

which had little effect in response to increasing estradiol, yeast containing MECP2 showed a

dose-dependent decrease in exponential growth rate. We observed a similar result with a more

conventional vector employing a galactose-inducible promoter (Supplemental Figure 1). Thus,

expression of MECP2 reduced the fitness of WT yeast.

MECP2 impacts DNA-related functions in yeast

To understand how MECP2 might be functioning in yeast and if this is related to the normal

function of MECP2 we employed a genetic interaction mapping approach we previously

developed to study the function of human genes in yeast, called Sentinel Interaction Mapping

(SIM) [27]. Using SIM we express the human gene of interest in a collection of nearly 4,800 yeast

deletion mutants and systematically identify aggravating and alleviating genetic interactions

between the human and yeast genes. The known functions of the interacting yeast genes inform

us on the function of the human gene in yeast and if it is relevant to its known function in

humans. SIM also has the capacity to uncover novel functions for human genes. We performed a

SIM screen with MECP2-ref in which MECP2 under control of an estradiol responsive promoter

was expressed in ~ 4,800 different yeast deletion mutant strains in high-density arrays. We

identified 132 interactions in 3/3 biological replicates (Supplemental Data 2). These interactions

showed enrichment for functions falling under the gene ontology categories DNA Damage

Response, DNA Binding and Cell Cycle Process, consistent with the known function of MECP2

as a DNA binding protein with roles in transcriptional repression (Figure 2C). Hence, these

interactions suggested that the decrease in fitness observed upon MECP2 expression was a result

of disruption of these or related processes in yeast.

Functionalization of MECP2 variants in yeast

We decided to simply exploit the robust slow growth phenotype observed in WT yeast to

measure the effects of MECP2 variants, anticipating that variants that interfered with MECP2

function would no longer slow down growth. We measured yeast growth in liquid culture for all

46 variants using a galactose-inducible vector, calculated exponential growth rates, and using a

linear mixed-effects model estimated the effects of the variants on MECP2 function. Raw data is

provided in Supplemental Data 3. Shown in Figure 3A are the model's estimates of MECP2

function arranged by variant amino acid position. Reference benign variants had little effect on

function as expected across all domains. The one exception was the D121E variant which was

nearly complete LoF, for which we have additional supporting evidence in fly leading to

reclassification of this variant as pathogenic (see below). For the MBD domain, all seven

reference pathogenic variants located in the MBD were complete or near-complete LoF. This

supported that our assay was disease-relevant for variants in the MBD. L100V, R106W, R133C,

and T158M have been previously reported to have reduced DNA binding *in vitro* and *in vivo* [40],

suggesting that our assay reported on this function of the MBD. R133 is part of the DNA

interface and interacts directly with guanine of the mCpG island, and T158 is part of the Asx-ST

motif that interacts with the DNA backbone [40]. The two test ASD variants in this domain, R91W

and P152L, were found to have decreased function, although not as severe as the reference

pathogenics. In contrast, reference pathogenics outside the MBD generally did not show

decreased function. Given the general consistency of the reference annotation with the assay

results for the MDB variants, our interpretation is that our assay was not sensitive to variants

outside the MBD.

Protein stability of variants in yeast

We measured protein stability for a subset of variants with a focus on the MBD to determine if

stability was a factor in our assessment of function. We also included additional non-MBD

reference benigns from the TRD (T228S) and the CTD (S356G) for comparison, and P388S

because this variant showed decreased function suggesting it was destabilized. MECP-2 protein

levels from yeast extracts were measured on Western blots and normalized to MECP2-ref

(Figure 3B and Supplemental Figure 1). In the MBD, 8 out of 13 MBD variants were

destabilized relative to MECP2-ref, while 5 were not significantly different from Ref. Stability

ranged from 75% (T148A) to 37% (R106W). The two reference benigns outside the MBD,

T228S and S356G, were stable. Now, comparing stability to function of MBD variants there

appeared to be little correlation. Some variants were stable yet non-functional (L100V, P101S,

D121E, R133C), while the T148A variant was destabilized yet functional. Some variants showed

both destabilization and decreased function (R106W, T158M, G163W). This suggested that in

some cases the variant affected MECP2 DNA binding directly (e.g. R133C), whereas in other

cases decreased function was also a result of destabilization. It also suggested the functional

assay was fairly insensitive to destabilization.

To determine the sensitivity of our assay to destabilization we looked to variants outside the

MBD, because these should not directly affect DNA binding. We identified one variant out of a

total of 29, P388S, that showed moderate LoF (approx. 50%), suggesting it might be

destabilized. We measured the stability of P388S and found it was highly destabilized (approx.

25% stable). A functional variant in the same domain, S356G, was not destabilized. Given the

modest decrease in function we observed for P388S relative to its high degree of destabilization,

we conclude our functional assay was relatively insensitive to destabilization. This was also

supported by T148A, which was 67% stable, but was completely functional. Overall, these data

suggested that the yeast functional assay primarily reported on the intrinsic DNA binding ability

of the MBD.

Known MECP2 pathogenic and benign variants function predictably in Drosophila

Expression of *MECP2* in *Drosophila* tissues has a wide range of phenotypes [28–31]. From these

reports, we chose to assess MECP2 variants in developing wing tissues; the reported ectopic

wing vein phenotype gives a quantitative readout of MECP2 protein function, and MECP2

genetically interacts with known co-repressor complexes relevant to MECP2 function in this

assay [28].

We integrated *UAS-MECP2* variants (on the *pGW.HA.attB* transgenic vector backbone) into the

same locus, *attP40*, to control for genome position effect and copy number[37]. After pilot

experiments testing a range of wing imaginal disc *GAL4* drivers and incubation temperatures, we

established assay conditions in which *A9-GAL4* virgin females were crossed to homozygous

UAS-MECP2 males, as well as a control *attP40* line, and raised animals at 29C. We transferred

mating adults through three consecutive 'assay' vials. Adult progeny were then collected 3 days

post-eclosion, and a single wing from a minimum of 10 adult male progeny per assay vial was

removed and slide-mounted, for a total of 30 wings per variant. Mounted wings were scanned

and total wing vein area / wing area per wing was analyzed using a Labkit machine learning

algorithm that semi-automated the quantitation of wing vein area (see Methods).

First, we wished to validate our assay by repeating previous observations, by examining the

effect of MECP2-ref expression in developing wing tissue [28]. In agreement, we observed ectopic

wing vein tissue in wings upon expression of *MECP2-ref* (Figure 4). Further, this phenotype was

strongly suppressed in a heterozygotic background for *Asx[XF23],* consistent with known roles

for MECP2 in chromatin regulation and gene expression [20].

Next, we examined wing vein phenotypes induced by expression of all MECP2 variants and

controls in parallel. We tested eight reference benign and eight reference pathogenic variants.

Representative images for selected variants as well as *attP40* negative controls and MECP2-ref

are shown in Figure 5A (see Supplemental Figure 2 for representative images of MECP2 variants

and also the masked area that was considered to contain vein tissue). Benign variants all showed

increased ectopic wing vein whereas reference pathogenics showed very little, and test (ASD)

variants had a range of outcomes. To quantify this phenotype, we trained two machine learning

algorithms to quantify (1) the total area of the wing vein region and (2) the total area of the wing

(see Supplemental Data 3 for details). Using a linear mixed effects model these two values were

used to calculate the estimated effect of variants relative to MECP2-ref and were normalized to

the vein area/wing area for MECP2-ref (at 1) and attp40 (at 0) (Figure 5B).

All reference benign variants functioned similar to MECP2-ref, except for D121E. This variant

was classified in ClinVar as benign by a single contributor, but we find both yeast and fly assays

characterize it as a strong LoF, hence we predict that D121E is pathogenic. All reference

pathogenic variants showed varying degrees of a LoF or reduced function phenotype, indicating

the wing vein assay reported on disease relevant functions for MECP2, especially for the MBD

and TRD domains for which we have multiple calibrating pathogenics. Most were strong LoF

variants, except for P225R, which had reduced function but not a complete LoF. This successful

calibration of the assay indicated that clinically pathogenic and benign variants can be correctly

classified in the fly model.

In the same experimental batch as the reference pathogenic and benign variants described above,

we determined the wing vein phenotype for MECP2 test (ASD) variants identified in ASD that

have no clear classification. We found that both MBD variants R91W and P152L were LoF. The

TRD variants P217L and E282G exhibited a partial LoF with data distributed similarly to the

TRD pathogenic variant P225R. Finally, the Interdomain (G195S, A202G) and CTD variants

(P376S, P388S and S411R) all exhibited normal function that compared to the reference benign

variants. Thus, we conclude that only ASD variants in the MBD and TRD domains were LoF.

Finally, we compared yeast and fly assay results for MBD variants (because the yeast assay

calibrated for MBD variants only). We observed remarkable similarity in results between the two

assays (compare Figures 3A and 5B). Of particular interest was that both test (ASD) variants

(R91W and P152L) appeared more functional in both yeast and fly assays than reference

pathogenic variants. Given that T158M is DNA-binding defective and is complete LoF in our

assays, this suggested that ASD variants retain more DNA binding capacity than Rett

pathogenics.

MECP2 localization and stability in human cells

To investigate whether MECP2 variants had altered nuclear localization we expressed GFP-

MECP2 in HEK cells (Raw data available in Supplemental Data 4). GFP-MECP2 localized

predominantly to the nucleus (Figure 7A) as expected. Using high-content imaging we assessed

the nuclear localization of all 46 variants using our imaging pipeline (Figure 7B). We identified

17 variants that showed altered nuclear localization. We were particularly interested in variants

that had decreased nuclear localization. 14 of the 17 showed decreased nuclear localization and

of these only 2 were reference benigns, whereas the remaining 12 were either reference

pathogenics or test (ASD). 9 of these were located in the TRD. The variant showing the greatest

delocalization was the pathogenic variant R255G, and consistent with this R255 is known to be

part of the nuclear localization signal [41]. Three test (ASD) variants showed substantial nuclear

delocalization, A202G, P217L and P388S, suggesting that nuclear delocalization of MECP2 is a

mechanism in ASD (see Figure 7A for representative images). Using this same image data we

also estimated protein stability of the variants (Figure 7C). 11 variants showed decreased

stability, of which 8 were reference pathogenics and 3 were test (ASD) (R91W, P217L, P376S).

Interestingly, 8 of the 11 were variants in the TRD domain, suggesting this region was important

for MECP2 stability. Twelve out of thirteen MBD variants were found to be stable, which

disagreed with the yeast stability measurements that found many MBD variants were

destabilized. This also disagreed with published *in vitro* and *in vivo* stability studies (see

Discussion), suggesting the HEK assay was less sensitive to destabilization in the MBD. Possible

explanations are that the N terminal GFP tag stabilized MBD variants, perhaps by altering

proteasome-mediated degradation [42], or that high expression driven by the CMV promoter

masked MBD destabilization. Because variants in the TRD are much more distal to the GFP tag

it may have had a lesser effect on stabilizing TRD variants.

DISCUSSION

Clinical classification of variants

Using our assays calibrated with known reference pathogenic and benign variants, we are able to

provide functional assessments for nine test ASD variants, six of which have been found in cases

of ASD without Rett (underlined): R91W, P152L, G195S, A202G, P217L, E282G, P376S,

P388S, S411R (Figure 7). We found that R91W and P152L were LoF variants (yeast and fly)

and we classify these as likely pathogenic. S411R functioned normally (fly) and G195S had a

modest gain of function (fly), hence we classify these as likely benign. This provides the first

classifications for P152L, G195S and S411R, and re-classifies R91W from a variant of uncertain

significance (VoUS) in ClinVar to a likely pathogenic variant. E282G showed moderately

reduced function (fly) and had reduced nuclear localization (HEK), consistent with its updated

likely pathogenic ClinVar classification (originally had no classification). A202G was found to

be functional (fly), but showed decreased nuclear localization (HEK). The ClinVar classification

for this variant is now conflicting (one benign and one VoUS report; originally it was uncertain)

and we recommend this variant be further tested before a classification can be made. For the

additional three test (ASD) variants that are ASD and Rett, our results largely support their

current ClinVar classifications. We find that R133C and P152R were LoF (yeast and fly),

matching their likely pathogenic ClinVar classifications. P217L showed moderately reduced

function (fly) and had reduced nuclear localization and decreased stability consistent with its

updated likely pathogenic ClinVar classification. P376S was found to be functionally normal

(fly), but had reduced stability (HEK) suggesting its updated benign classification in ClinVar

should be re-examined. Finally, P388S was found to be functionally normal (fly), which

corresponds with its updated benign classification in ClinVar; however our HEK results

indicated P388S had reduced nuclear localization. Because we did not test pathogenic variants in

the CTD in the *Drosophila* assay we would suggest that P388S is worthy of further examination.

D121E, which is classified as benign in ClinVar, was LoF in both yeast and *Drosophila* assays.

Interestingly, our data was consistent with *in vitro* studies showing a ~7-fold reduction in the

DNA binding capacity of the D121E variant [43]. This variant is not in VariCarta or RettBASE. It

has one report in ClinVar, but this report lacks a description for how a benign classification was

made. Interestingly, the computational predictors SIFT, PolyPhen and CADD all predict this

variant is likely to cause LoF. We propose this variant should be re-classified in ClinVar as

likely pathogenic.

Calibrating functional assays by domain

MEPC2 is a multi-domain protein, yet it is largely unstructured outside the MBD [44]. We initially

focused on identifying numerous reference variants in the MBD and TRD, as these domains

harbour many of the clinically relevant variants. The yeast assay only calibrated for the MBD

and failed to calibrate for all other domains indicating we can only assess functional effects of

variants in the MBD. The likely explanation for this is that the MECP2 MBD binds DNA in

yeast without the need of accessory factors. Indeed, in an accompanying paper, Brown et al.

show this to be likely the case. The TRD domain mediates interactions with co-repressor

complexes, including Sin3A, N-Cor and REST, and Rett-causing mutations disrupt these

interactions [45]. It is likely that the assay failed to calibrate for TRD variants because these

complexes either are not present in yeast (N-Cor and REST) or the TRD did not interact with the

yeast orthologs (Sin3A/HDAC). This may also be the case for the other domains for which

pathogenic reference variants failed to exhibit altered function.

Drosophila was successful in stratifying reference pathogenic and benign variants in both the

MBD and TRD domains, indicating we can reliably make predictions about test ASD variants

located in both these domains. For variants in the MBD, it is likely these disrupt DNA binding,

although this has yet to be shown. For the TRD, our data suggest MECP2 is able to interact with

the *Drosophila* orthologs of the co-repressor complexes. First, R306C prevents interaction with

co-repressors and was complete LoF in our assay [45]. Second, co-repressor complex mutants

suppress phenotypes resulting from MECP2 over-expression in the *Drosophila* wing [28]. For

example, the Polycomb group co-repressor ortholog of Asxl1, Asx, shows genetic interactions

with MECP2 expressed in *Drosophila* [28]. We further confirmed mutation of Asx suppressed

wing vein overgrowth resulting from MECP2 overexpression, supporting MECP2 interacts with

this co-repressor. Thus, when developing functional assays in surrogate model systems with

some evolutionary distance from humans, it is important to consider calibrating assays for each

domain before making conclusions about the effects of test variants.

MECP2 protein stability

The stability of a handful of MECP2 variants has been studied both *in vitro* and in *in vivo* mouse

models [40,46–48]. We could compare seven variants located in the MBD between our yeast results

and these studies (Supplemental Table 1). R106W, P152R and T158M were all destabilized in

yeast, mouse and *in vitro* systems. R133C was found stable in yeast and *in vitro* systems, but

destabilized in mouse. In contrast, L100V and P101S were stable in yeast but destabilized *in*

vitro, and R133H was destabilized in yeast but stable *in vitro*. These results highlight the value of

comparing variant function across multiple species and model systems. For example, MECP2

turnover in cells is regulated by ubiquitylation by the E3 ligase RNF4 [42], indicating an *in vivo*-

specific modifier of variant stability. Moreover, the use of GFP-tagged MECP2 in a subset of

studies could conceivably affect protein turnover. Interestingly, we identified the yeast orthologs

of RNF4, Slx5 and Slx8, as aggravating genetic interactions in our SIM screen, suggesting

MECP2 levels may also be regulated similarly in yeast. Thus, intrinsic stability and regulated

turnover *in vivo* are important factors contributing to MECP2 variant protein levels and may vary

across models.

For variants outside the MBD, 14 variants showed altered stability using the HEK cell assay.

Comparing our results to the literature, 3/3 agreed with animal models (P225R and P322L were

destabilized and R306C was stable). This suggested our HEK cell assay was able to detect

destabilization for variants outside the MBD. Interestingly, reference pathogenics in the TRD

predominately showed decreased stability (7/12 variants). Thus, destabilization could be a

mechanism for the pathogenicity of TRD variants. MECP2 is known to be largely unstructured

outside the MBD, however when recruited to DNA, secondary structure significantly increases

(38-59% for the ID and 22-30% for the TRD) [44]. This suggests disruption of interactions with co-

repressors by these variants could be an additional factor leading to destabilization and Rett

and/or ASD phenotypes.

MECP2 nuclear localization

Nuclear localization of MECP2 is mediated by its NLS (amino acids 255-271) which binds to the

nuclear import factors karyopherins 3 and 4, and by the MBD domain in a karyopherin-

independent yet unknown mechanism [41,49]. An engineered mutation in the NLS, R255D, disrupts

MECP2 nuclear localization and interaction with karyopherins [41], but there have been no

missense mutations identified in the NLS in Rett to date, raising the question of the importance

of the NLS to normal brain function. This is of potential interest because of the attractiveness of

karyopherin binding as a therapeutic target. Our finding that R255G is strongly nuclear

delocalized, together with the fact that this variant is found in an individual with intellectual

disability and is predicted to be pathogenic, suggests mutations in the NLS may disrupt MECP2

function. Importantly, we found three test (ASD) variants, A202G, P217L and P388S, that

showed substantially decreased nuclear localization, suggesting disrupted nuclear targeting is a

mechanism in ASD. Because these variants lie outside the NLS suggests nuclear targeting by the

MBD may be affected, although the mechanism remains undefined.

MECP2 and DNA methylation

It is noteworthy that all reference pathogenic variants in the MBD were LoF in our functional

assays despite there being no, or extremely low levels of DNA cytosine methylation in yeast and

Drosophila respectively. Our data correlate well with Brown et al (accompanying article)

showing that MECP2 binds the unmethylated yeast genome and mediates widespread

transcriptional suppression, which is reduced in the pathogenic Rett variant T158M. This raises

the possibility that ASD and Rett MECP2 variants may not be specifically deficient in their

ability to bind methylated DNA, but may be generally deficient in binding to unmethylated DNA

and/or chromatin. We plan on expanding the panel of MECP2 Rett variants tested in our assays,

towards testing whether most if not all Rett pathogenic variants are deficient in non-methylated

DNA binding. It is challenging to specify differential impacts of specific Rett variants on methyl

versus non-methyl DNA binding in models where DNA methylation is present. We propose this

may have obscured the impact that pathogenic variants have on non-methylated DNA binding

and its importance in healthy and disease states, and propose that the yeast and fly models offer

valuable platforms to further examine these mechanisms.

REFERENCES

1. Lewis, J. D. *et al.* Purification, sequence, and cellular localization of a novel chromosomal

protein that binds to Methylated DNA. *Cell* **69**, 905–914 (1992).

2. Amir, R. E. *et al.* Rett syndrome is caused by mutations in X-linked MECP2, encoding

methyl-CpG-binding protein 2. *Nat. Genet.* **23**, 185–188 (1999).

3. Wan, M. *et al.* Rett Syndrome and Beyond: Recurrent Spontaneous and Familial MECP2

Mutations at CpG Hotspots. *Am. J. Hum. Genet.* **65**, 1520–1529 (1999).

4. Chahrour, M. & Zoghbi, H. Y. The Story of Rett Syndrome: From Clinic to Neurobiology.

Neuron **56**, 422–437 (2007).

5. Nomura, Y. Early behavior characteristics and sleep disturbance in Rett syndrome. *Brain*

Dev. **27**, S35–S42 (2005).

6. Amir, R. E. *et al.* Influence of mutation type and X chromosome inactivation on Rett

syndrome phenotypes. *Ann. Neurol.* **47**, 670–679 (2000).

7. Ng, H. H. & Bird, A. DNA methylation and chromatin modification. *Curr. Opin. Genet.*

Dev. **9**, 158–163 (1999).

8. Jones, P. L. *et al.* Methylated DNA and MeCP2 recruit histone deacetylase to repress

transcription. *Nat. Genet.* **19**, 187–191 (1998).

9. Harikrishnan, K. N. *et al.* Brahma links the SWI/SNF chromatin-remodeling complex with

MeCP2-dependent transcriptional silencing. *Nat. Genet.* **37**, 254–264 (2005).

10. Lunyak, V. V. *et al.* Corepressor-Dependent Silencing of Chromosomal Regions Encoding

Neuronal Genes. *Science* **298**, 1747–1752 (2002).

11. Ehrhart, F. *et al.* A catalogue of 863 Rett-syndrome-causing MECP2 mutations and lessons

learned from data integration. *Sci. Data* **8**, 10 (2021).

12. Yusufzai, T. M. & Wolffe, A. P. Functional consequences of Rett syndrome mutations on

human MeCP2. *Nucleic Acids Res.* **28**, 4172–4179 (2000).

13. Ballestar, E., Yusufzai, T. M. & Wolffe, A. P. Effects of Rett Syndrome Mutations of the

Methyl-CpG Binding Domain of the Transcriptional Repressor MeCP2 on Selectivity for

Association with Methylated DNA. *Biochemistry* **39**, 7100–7106 (2000).

14. Leblond, C. S. *et al.* Operative list of genes associated with autism and neurodevelopmental

disorders based on database review. *Mol. Cell. Neurosci.* **113**, 103623 (2021).

15. Wen, Z. *et al.* Identification of autism-related MECP2 mutations by whole-exome

sequencing and functional validation. *Mol. Autism* **8**, 43 (2017).

16. Wang, T. *et al.* De novo genic mutations among a Chinese autism spectrum disorder cohort.

Nat. Commun. **7**, 13316 (2016).

17. Guo, H. *et al.* Inherited and multiple de novo mutations in autism/developmental delay risk

genes suggest a multifactorial model. *Mol. Autism* **9**, 64 (2018).

18. Schaaf, C. P. *et al.* Oligogenic heterozygosity in individuals with high-functioning autism

spectrum disorders. *Hum. Mol. Genet.* **20**, 3366–3375 (2011).

19. Cheng, T.-L. *et al.* MeCP2 suppresses nuclear microRNA processing and dendritic growth

by regulating the DGCR8/Drosha complex. *Dev. Cell* **28**, 547–560 (2014).

20. Georgel, P. T. *et al.* Chromatin compaction by human MeCP2. Assembly of novel secondary

chromatin structures in the absence of DNA methylation. *J. Biol. Chem.* **278**, 32181–32188

(2003).

21. Hansen, J. C., Ghosh, R. P. & Woodcock, C. L. Binding of the Rett syndrome protein,

MeCP2, to methylated and unmethylated DNA and chromatin. *IUBMB Life* **62**, 732–738

(2010).

22. Skene, P. J. *et al.* Neuronal MeCP2 is expressed at near histone-octamer levels and globally

alters the chromatin state. *Mol. Cell* **37**, 457–468 (2010).

23. Lyko, F., Ramsahoye, B. H. & Jaenisch, R. DNA methylation in Drosophila melanogaster.

Nature **408**, 538–540 (2000).

24. Buitrago, D. *et al.* Impact of DNA methylation on 3D genome structure. *Nat. Commun.* **12**,

3243 (2021).

25. Ganguly, P. *et al.* A scalable Drosophila assay for clinical interpretation of human PTEN

variants in suppression of PI3K/AKT induced cellular proliferation. *PLoS Genet.* **17**,

e1009774 (2021).

26. Post, K. L. *et al.* Multi-model functionalization of disease-associated PTEN missense

mutations identifies multiple molecular mechanisms underlying protein dysfunction. *Nat.*

Commun. **11**, 2073 (2020).

27. Young, B. P. *et al.* Sentinel interaction mapping - a generic approach for the functional

analysis of human disease gene variants using yeast. *Dis. Model. Mech.* **13**, dmm044560

(2020).

28. Cukier, H. N. *et al.* Genetic Modifiers of MeCP2 Function in Drosophila. *PLoS Genet.* **4**,

e1000179 (2008).

29. Vonhoff, F., Williams, A., Ryglewski, S. & Duch, C. Drosophila as a model for MECP2 gain

of function in neurons. *PloS One* **7**, e31835 (2012).

30. Williams, A. A. *et al.* MECP2 impairs neuronal structure by regulating KIBRA. *Neurobiol.*

Dis. **91**, 284–291 (2016).

31. Williams, A. A. *et al.* Apoptotic Activity of MeCP2 Is Enhanced by C-Terminal Truncating

Mutations. *PLoS ONE* **11**, e0159632 (2016).

32. Brnich, S. E. *et al.* Recommendations for application of the functional evidence PS3/BS3

criterion using the ACMG/AMP sequence variant interpretation framework. *Genome Med.*

12, 3 (2019).

33. McKnight, D. *et al.* Recommendations by the ClinGen Rett/Angelman-like expert panel for

gene-specific variant interpretation methods. *Hum. Mutat.* **43**, 1097–1113 (2022).

34. Belmadani, M. *et al.* VariCarta: A Comprehensive Database of Harmonized Genomic

Variants Found in Autism Spectrum Disorder Sequencing Studies. *Autism Res. Off. J. Int.*

Soc. Autism Res. **12**, 1728–1736 (2019).

35. McLaren, W. *et al.* The Ensembl Variant Effect Predictor. *Genome Biol.* **17**, 122 (2016).

36. Good, K. V., Vincent, J. B. & Ausió, J. MeCP2: The Genetic Driver of Rett Syndrome

Epigenetics. *Front. Genet.* **12**, 620859 (2021).

37. Bischof, J., Maeda, R. K., Hediger, M., Karch, F. & Basler, K. An optimized transgenesis

system for Drosophila using germ-line-specific phiC31 integrases. *Proc. Natl. Acad. Sci. U.*

S. A. **104**, 3312–3317 (2007).

38. Young, B. P. & Loewen, C. J. R. Balony: a software package for analysis of data generated

by synthetic genetic array experiments. *BMC Bioinformatics* **14**, 354 (2013).

39. Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. lmerTest Package: Tests in Linear

Mixed Effects Models. *J. Stat. Softw.* **82**, 1–26 (2017).

40. Tillotson, R. & Bird, A. The Molecular Basis of MeCP2 Function in the Brain. *J. Mol. Biol.*

432, 1602–1623 (2020).

41. Baker, S., Lombardi, L. & Zoghbi, H. Karyopherin α 3 and Karyopherin α 4 Proteins

Mediate the Nuclear Import of Methyl-CpG Binding Protein 2. *J. Biol. Chem.* **290**, (2015).

42. Wang, Y. RING finger protein 4 (RNF4) derepresses gene expression from DNA

methylation. *J. Biol. Chem.* **289**, 33808–33813 (2014).

43. Free, A. *et al.* DNA Recognition by the Methyl-CpG Binding Domain of MeCP2 *. *J. Biol.*

Chem. **276**, 3353–3360 (2001).

44. Ghosh, R. P. *et al.* Unique physical properties and interactions of the domains of methylated

DNA binding protein 2. *Biochemistry* **49**, 4395–4410 (2010).

45. Lyst, M. J. *et al.* Rett syndrome mutations abolish the interaction of MeCP2 with the

NCoR/SMRT co-repressor. *Nat. Neurosci.* **16**, 898–902 (2013).

46. Yang, Y., Kucukkal, T. G., Li, J., Alexov, E. & Cao, W. Binding Analysis of Methyl-CpG

Binding Domain of MeCP2 and Rett Syndrome Mutations. *ACS Chem. Biol.* **11**, 2706–2715

(2016).

47. Kucukkal, T. G., Yang, Y., Uvarov, O., Cao, W. & Alexov, E. Impact of Rett Syndrome

Mutations on MeCP2 MBD Stability. *Biochemistry* **54**, 6357–6368 (2015).

48. Sperlazza, M. J., Bilinovich, S. M., Sinanan, L. M., Javier, F. R. & Williams, D. C.

Structural Basis of MeCP2 Distribution on Non-CpG Methylated and Hydroxymethylated

DNA. *J. Mol. Biol.* **429**, 1581–1594 (2017).

49. Lyst, M. J. *et al.* Affinity for DNA Contributes to NLS Independent Nuclear Localization of

MeCP2. *Cell Rep.* **24**, 2213–2220 (2018).

DATA SHARING STATEMENT

673 Strains and plasmids are available upon request. The authors affirm that all data necessary for

674 confirming the conclusions of the article are present within the article, figures, and tables.

ACKNOWLEDGEMENTS

We thank Dr. Janel Kopp, UBC for access to the Panoramic Midi II digital slide scanner. Susan

Lindquist for the generous gift of the *pAG426GAL-ccdB* vector. Stocks obtained from the

Bloomington *Drosophila* Stock Center (NIH P40OD018537) were used in this study. We thank

Drosophila Genomics Resource Center, supported by NIH grant 2P40OD010949 for the esc

cDNA. We used FlyBase (releases 2020 through 2024) to find information on

phenotypes/function/stocks/gene expression (etc). We would like to thank Gino Laberge at

Genome ProLab for the kind gift of balancer stocks and also their transgenic services (Genome

ProLab Inc, Quebec, Canada). We thanks Libby Natola and Guillaume Poirier-Morency

(Pavlidis Lab) for contributions to the variant annotation pipeline. This work was funded by an

operating grant from The Simons Foundation Autism Research Initiative (2021 Genomics of

ASD: Pathway to Genetic Therapies: AWD-021951. SIMOFOUN 2021).

AUTHOR CONTRIBUTIONS

DWA, CJRL, BPY, SMG, JTC, MD, SR, MD, PP contributed to conception of the work

DWA, CJRL, GM, EC, BPY, SI, TL, JL, JBL, SMG, BB, MD, JS, AS, JTC SR, PP contributed

to design of the work

DWA, CJRL, GM, EC, BPY, JBL, LM, JS, JBL, AS, VK, JTC, SR, PP contributed to

acquisition, analysis, or interpretation of data

DWA, SR, PP, GM, EC, JBL, MD, BB, BW, JTC contributed to creation of new software used

DWA, CJRL, JS, BPY, JTC, JBL, GM, EC, LM, JL, JBL, AS, JTC, SR, PP contributed to

drafting the paper or substantively revised it

FIGURE CAPTIONS

Figure 1: Forty-six MECP2 variants by amino acid position and class. Reference pathogenic and

benign variants are clinically annotated variants used to calibrate assays and test (ASD) variants

are variants with no clinical classification found in individuals with ASD. NTD – N-terminal

domain, MBD – Methyl-CpG-binding domain, ID – Interdomain, TRD – Transcription repressor

domain, CTD – C-terminal domain.

Figure 2: MECP2 expression and genetic interactions in yeast. (A) Untagged MECP2-ref

expressed either from a plasmid utilizing the estradiol-responsive promoter or TDH3 constitutive

promoter (CEN low-copy or 2 micron high-copy) in WT yeast. Top band corresponds to full-

length MECP2. Western blot labelled with anti-MECP2 antibody. (B) Effect of MECP2-ref

expression on growth of WT yeast from a plasmid utilizing the estradiol-responsive promoter.

EV – empty vector, MECP2 – MECP2-ref. (C) Results of SIM genetic interaction screen with

MECP2-ref. Functional enrichment for genetic interactions is shown where nodes indicate

enriched GO terms and edges indicate statistically significant associations between terms. Node

size and edge thickness indicate statistical significance (minimum $p < 0.05$). Bolded labels

indicate most significant terms within networks.

Figure 3: Analysis of 46 MECP2 variants using the yeast assay. (A) Estimated functional effects

of MECP2 variants expressed in WT yeast. Data is normalized to MECP2-ref (=1) and empty

vector (=0). Error bars indicate 95% confidence intervals. (B) Protein stability of MECP2

variants. Protein levels normalized to MECP2-ref and plotted on a log scale. Error bars indicate

S.D. (n = 3). * indicates $p < 0.05$, ** indicates $p < 0.01$. Colour coding same as in (A). All

variants are located in MBD except T228S (TRD), S356G (CTD) and P388S (CTD).

Figure 4: MECP2 induces ectopic wing-vein formation which is suppressed by Asx loss-of-

function. (A) Crossing scheme to introduce MECP2-ref expression into WT and Asx[XF23]

background. (B) Representative images of wings of the indicated genotypes at the same scale.

Arrows indicate ectopic wing vein.

Figure 5: Analysis of MECP2 variants in the *Drosophila* wing assay. (A) Representative images

of wings expressing the indicated variants. *attp40* – no MECP2, *MECP2-ref* – WT MECP. All

images same scale. (B) Graph showing normalized activity of MECP2 variants where *MECP2-*

ref (WT MECP2) = 1 and *attp40* (no MECP2) = 0. Error bars indicate 95% confidence interval.

Figure 6: MECP2 localization and stability in HEK cells. (A) Representative images of MECP2-

ref (WT) and test (ASD) variants showing altered nuclear localization. (B) Translocation

analysis for MECP2 nuclear:cytoplasmic localization using high-throughput imaging. (C) Protein

abundance analysis for MECP2 using high-throughput imaging.

Figure 7: Summary of results for variants in MECP2 tested in yeast, *Drosophila* and HEK-cell

assays.

SUPPLEMENTAL FIGURE CAPTIONS

Supplemental Figure 1: (A) Growth rate of PGAL1-MECP2-ref compared to empty vector

(EV). Error bars indicate standard deviation (n = 3). (B) Representative MECP2 Western blots.

Arrow indicates the full-length MECP2 protein. Antibody labelling is as follows: red - anti-

MECP2, green - anti-PGK1.

Supplemental Figure 2: Representative wing phenotypes for all MECP2 variants. For each

variant, raw images are on the left and the masked image used to quantitate total wing vein per

wing is on the right.

SUPPLEMENTAL METHODS CAPTIONS

Supplemental Methods 1: Detailed outline of wing vein quantitation methods.

SUPPLEMENTAL DATA CAPTIONS

Supplemental Data 1: Detailed information for each variant, primers used in cloning, the

molecular genetic reagents made for each organism, and the *Drosophila* stocks generated.

Supplemental Data 2: Yeast SIM screen raw data

Supplemental Data 3: Raw data for yeast and *Drosophila* assays

Supplemental Data 4: Raw data for HEK assays

Figure 1

Figure 2

A

B

Figure 3

A

A9-GAL4
or
A9-GAL4: Asx[XF23]/CyO

♀

X →

♂

attP40
or
UAS-MECP2-ref

B *A9-GAL4/Y; attP40/+*

A9-GAL4/Y; UAS-MECP2-ref / +

A9-GAL4/Y; UAS-MECP2-ref / Asx[XF23]

100 µm

Figure 4

Figure 5

Figure 6

Figure 7

Figure S1

Figure S2-1

PATHOGENIC

P225R   *R306C*

TEST (ASD)

R91W   *P152L*

G195S   *A202G*

P217L   *E282G*

P376S   *P388S*

S411R

Figure S2-2

Supplementary Table 1: Summary of MECP2 protein stability results

Variant	Yeast*	HEK*	Yang [1] (in vitro)	Kucukkal [2] (in vitro) ($\Delta\Delta G$)	Sperlazza [3] (NMR) (+ slight, ++ more destabilized)	Tillotson [4] (mouse models)
G7E		0.6				
R91W	1	0.65				
L100V	1	1	0.5	1.25		
P101S	1	1			+	
R106W	0.4	1	1	0.1	++	0.1 (TAVI)
D121E	1	1				
R133C	1	1	1	0.4		0.6 (eGFP)
R133H	0.6	1	1	0.25		
K135E	0.5	1				
T148A	0.75	1				
P152L	0.75	1				
P152R	0.4	1	0.25	1.5		
T158M	0.5	1	1	0.25	+	0.3 (eGFP), 0.25 (TAVI)
G163W	0.75	1.25				
A202G		1.4				
P217L		0.35				
P225R		0.7				0.22
P225T		0.25				
T228S	1	1.25				
E282G		1.5				
I303V		1.2				
K304E		0.75				
K304N		0.8				
R306C		1				1.0
R306L		0.2				
R309W		0.5				
P322L		0.3				0.03
S356G	1.5	1				
P376S		0.5				
P388S	0.25	1.6				

* values not significantly different than MECP2-Ref are set to 1, all others are significant

References:

1. Yang, Y., Kucukkal, T. G., Li, J., Alexov, E. & Cao, W. Binding Analysis of Methyl-CpG Binding Domain of MeCP2 and Rett Syndrome Mutations. *ACS Chem. Biol.* **11**, 2706–2715 (2016).