Mesenchymal Meis2 controls whisker development independently from trigeminal sensory innervation

Mehmet Mahsum Kaplan [1], Erika Hudacova [1,2], Miroslav Matejcek [1], Haneen Tuaima [4], Jan Krivanek [4] and Ondrej Machon [1,3*]

* Correspondence: ondrej.machon@iem.cas.cz

[1] Department of Developmental Biology, Institute of Experimental Medicine, Czech Academy of Sciences, Prague, Czech Republic

[2] Department of Cell Biology, Faculty of Science, Charles University, Prague, Czech Republic

[3] Laboratory of Transcriptional Regulation, Institute of Molecular Genetics, Czech Academy of Sciences, Prague, Czech Republic

[4] Department of Histology and Embryology, Faculty of Medicine, Masaryk University, Brno, Czech Republic

Abstract

Hair follicle development is initiated by reciprocal molecular interactions between the placode-forming epithelium and the underlying mesenchyme. Cell fate transformation in dermal fibroblasts generates cell niche for placode induction by activation of signaling pathways WNT, EDA and FGF in epithelium. These successive paracrine epithelial signals initiate dermal condensation in the underlying mesenchyme. Although epithelial signaling from the placode to mesenchyme is better described, little is known about primary mesenchymal signals resulting in placode induction. Here we show that *Meis2* expression in cells derived from the neural crest is critical for whisker formation, and also for branching of trigeminal nerves. While whisker formation is independent of the trigeminal sensory innervation, MEIS2 in mesenchymal dermal cells orchestrates initial steps of epithelial placode formation and subsequent dermal condensation. MEIS2 regulates the expression of transcription factor *Foxd1* which is typical of pre-dermal condensation. However, deletion of *Foxd1* does not affect whisker development. Overall, our data report an early role of mesenchymal MEIS2 during whisker formation and provide evidence that whiskers can normally develop in the absence of sensory innervation or FOXD1 expression.

Keywords

whisker follicle; hair follicle placode; cranial nerves; Meis2; Foxd1, Sox2

Introduction

The local cell environment provides essential cues that determine cell fate decisions, proliferation, and migration. The molecular communication between the developing epithelium and the underlying cell environment is crucial for the formation of multiple structures, which are derived from the embryonic epithelium, such as eye lens, hair follicles including whisker (vibrissae) follicles, teeth, taste papillae, the intestine lumen, or salivary glands. In the developing vertebrate head, these tissue cross-talks employ neural crest-derived mesenchyme with the epithelium, myogenic progenitors, or cartilage. Critical steps in whisker follicle (WF) development are very similar to normal hair follicle (HF) development [1] which has been studied much more extensively. HF morphogenesis is initiated by thickening of the epithelium forming placode (Pc) which subsequently induces condensation of underlying fibroblasts termed dermal condensation (DC) [2]. Beta-catenin dependent WNT signaling is essential for Pc induction [3] which initiates EDA pathway [4], FGF20 [5] and SHH signaling [6] in the epithelium. Abrogation of any of these signaling pathways leads to the arrest of HF development. However, the succession of initial inductive steps in HF formation is still unclear. Pc formation requires hitherto unknown primary signals from dermal fibroblasts (Fb) at pre-DC stage that is characterized by expression of *Foxd1* [7] and *Prdm1* [8]. Recent studies employing single-cell transcriptomics helped elucidate molecular determination of Pc and DC induction. Mok et al. [7] described a wave of transcription factors that define cell fate trajectories, starting in dermal fibroblasts underlying the epithelium before Pc induction (pre-DC stage), towards DC and late DC stage when Pc invagination commences. For instance, expression of *Foxd1* transcription factor was detected in pre-DC fibroblasts that are also characterized by expression of *Twist2, Lef1*, and *Smad3* [7,9]. DC cells predominantly express *Sox2, Foxd1, Ptch1, Bmp4* [10,11]. On the other hand, DC initiation requires canonical WNT and FGF20 signaling in Pc, suggesting a reciprocal molecular crosstalk between the dermal mesenchyme and overlying epithelium [7,10]. Transplantation experiments suggested that mesenchymal cells modulate WNT signaling in Pc by activation of *Rspo1*, however, RSPO1 is was not sufficient for HF induction even in combination of BMP inhibition [12]. Although these data comprehensively map molecular events during HF initiation, identification of the primary inductive signal for Pc induction coming from dermal fibroblasts is still missing.

Epithelial structures such as whiskers or teeth are formed at precisely defined locations. Whiskers are arranged in an accurate rectangular pattern in five rows each containing four to nine follicles. Each whisker is innervated by a separate whisker-to-barrel circuit that is reflected in a clear topographic organization of the barrel cortex [13]. Axons of the trigeminal nerve ending in the infraorbital nerve innervate each whisker germ in the maxillary prominence. It has been speculated that early innervation plays a role in initiating WF formation [1]. A similar hypothesis was proposed for tooth development based on several observations. For instance, a surgical removal of trigeminal (TG) axons innervating teeth led to tooth degeneration [14,15]. Hence, cranial nerves might deliver morphogenetic signals that orchestrate tooth maintenance. The role of innervation during tooth development was also supported by Kaukua et al. [16], who showed that dental mesenchymal stem cells are recruited from the nerve-associated glial cell. Thus, peripheral nerves and accompanying glia contribute to the mesenchymal tooth germ niche. Nonetheless, it is still unclear what cellular mechanisms regulate precise positions of whiskers in the snout. Turing reaction-diffusion mathematical model can be applied for oscillating tissue patterning that is orchestrated by diffusion of short-range activating morphogen that triggers expression of long-range inhibitor. Reaction-diffusion-based instruction of the epithelium by combinatorial WNT, FGF and BMP signaling is capable of HF placode induction in highly restricted regular pattern [17]. However, highly specific rectangular arrangement and relatively long distance among WFs within the developing snout may indicate that other molecular determinants play a role in spatiotemporal control of WF induction. This revives the previous speculations that axonal network may be involved in WF positioning. In this report, we assessed the scenario that the neural network is implicated in the control of accurate geometrical arrangement of WFs.

Here, we generated *Meis2* conditional mutants using the Wnt1-Cre2 driver [18] targeting exclusively the neural crest-derived tissues, including the dermal mesenchyme and cranial nerves, without detectable recombination in the overlying epithelium [19]. Although *Meis2* expression in the epithelium of Wnt1-Cre2; Meis2 [fl/fl] (Meis2 cKO) mutants was preserved, whisker development was arrested at Pc

induction stage. This is documented by failure in induction of expression of placodal genes such as *Shh*,
Edar and *Lef1*. We further show that severely affected branching of trigeminal nerves innervating WFs
cannot cause their developmental arrest because WFs form normally even in a complete absence of
trigeminal nerves. Expression of *Meis2* in dermal fibroblasts is therefore necessary for induction of
whisker placodes followed by dermal condensation. Protein expression analysis suggests that MEIS2
transcription factor is essential for triggering primary inductive signal from the mesenchyme to the
epithelium to initiate WF placode formation. While single-cell transcriptomics and immunohistochemical
validations showed that Meis2 operates upstream of Foxd1, primary pre-DC marker, we report a
dispensable role of Foxd1 during WF development. Overall, our data not only describe an essential role
of mesenchymal Meis2 for the WF development, but also document normal WF formation even in the
complete absence of innervating axons.
101
Results

Meis2 expression in cranial neural crest-derived cells is required for correct trigeminal nerve
projections and whisker development

To study whisker morphogenesis, we used Wnt1-Cre2; *Meis2*^{fl/fl} conditional mutants, in which *Meis2* is
deleted in neural crest-derived cells (hereafter, *Meis2* cKO). Micro-computed tomography (micro-CT) of
E15.5 embryos revealed severely compromised development of whiskers (Figure 1A). Missing whiskers
were also observed whole-mount embryonic heads stained with placodal marker Sox9 antibody after
light-sheet microscopy scanning at as early as E13.5 (Figure 1B). It is noteworthy that we observed in
some cases a small number of Sox9+ WFs in *Meis2* cKO. They most likely represented normal follicles
that probably 'escaped' from Wnt1-Cre2-mediated deletion of *Meis2*. The number of WF 'escapers' in
Meis2 cKO snouts varied among samples. Some mutant embryos completely missed WFs, but some
showed a reduction in their numbers. Overall, quantification of the WF number documented reduction
to 5.7 ± 2.0% at E12.5 and to 17.1 ± 5.9% at E13.5 in the mutants (n=at least 3 embryos for E12.5, n=
at least 8 embryos for E13.5, mean ± sem). Analysis of placode thickness/follicle length and DC area
revealed that these escaper whiskers have normal morphology (Supp Figure 1A, B). To analyze
expression of *Meis2* in the snout, we performed fluorescent immunohistochemistry. MEIS2 protein was
detected in the epithelium, in the most superficial dermis including dermal condensates (DC). In *Meis2*
cKO, mesenchymal expression of Meis2 virtually disappeared (Figure 1C, asterisk) while its epithelial
expression was maintained (Figure 1C, arrowheads), which corresponds to tissue-specific Cre
recombination of Wnt1-Cre2 [18,19]. It is interesting that MEIS2 signal was also remarkably reduced around
sole WF 'escapers' (Figure 1C, arrow) but we cannot exclude that, due to incomplete Cre-mediated
deletion, a minor expression of *Meis2* below the immunohistochemical detectability threshold is
sufficient for WF induction. To verify the tissue-specificity in the snout region, we crossed Wnt1-Cre2
mice with mTmG mouse strain where only cells derived from the neural crest express membrane-
localized GFP. Whole mount immunofluorescence imaging confirmed the Wnt1-Cre2 activity in neural
crest derivates in the craniofacial area, midbrain, and dorsal spinal cord (Figure 1D). In the snout, Cre
activity was restricted to dermal mesenchyme without a detectable recombination in the overlying
epithelium (Figure 1E). Moreover, branches of trigeminal sensory nerves were labelled with GFP (Figure
1E, arrows), consistent with the previously known origin of trigeminal neurons mainly from neural crest
cells [20]. Whiskers are innervated by the axons of the infraorbital nerve, a branch of the trigeminal nerve.
To examine the axonal outgrowths associated with whisker germs, we co-labelled the hair follicle marker
Sox9 with the sensory nerve marker TrkA in whole-mount heads at E13.5. Figure 1F and Supp Video 1
and Supp Video 2 show that missing whiskers in mice lacking *Meis2* in neural crest-derived cells were
accompanied by severely affected trigeminal nerve growth and branching. Because defects in nerve
branching were not limited to infraorbital nerve and were also observed in other trigeminal branches,
this phenotype is most likely not caused by missing WFs themselves. Moreover, the size of the
trigeminal ganglion and the nerve exit from the ganglion appeared grossly normal (Supp Figure 1C).

These observations show that defects in the trigeminal nerve morphology occur during branching and
fasciculation of axonal projections through the craniofacial region. Overall, these results indicate an
essential role of MEIS2 not only in development of the whiskers follicles but also for the branching of
trigeminal nerves that innervate them.

Trigeminal innervation is not required for WF development

Aberrant whisker development in *Meis2* cKO could be explained by an indispensable function of Meis2
in mesenchymal cells or by insufficient trigeminal sensory innervation. The latter scenario takes into
account the possibility that the nerve or nerve-associated Schwann cells produce factors that are
necessary for WF formation. Thus, compromised branching of sensory axons in the *Meis2* cKO snout
might result in insufficient secretion of crucial WF inducing factors. In order to directly assess the function
of sensory innervation in WF development, we analyzed Neurog1[-/-] mice which do not develop some
cranial nerves including trigeminal nerves and therefore lack sensory innervation of the developing
whiskers [21]. Immunostaining of E13.5 100-μm thick snout cryosections for placodal marker Sox9, axonal
marker Tuj1 and sensory nerve marker TrkA antibodies revealed that whiskers developed normally in
Neurog1[-/-] mutants even in the complete absence of innervating sensory axons (Figure 2A). Lack of any
Tuj1 labelled axon fascicules in the proximity of the normally developed WFs in the Neurog1[-/-] snout
further excludes the possibility of non-sensory neurons triggering WF development in the absence of
sensory innervation. In order to assess the role of sensory innervation for overall whisker patterning, we
stained E12.5 and E13.5 snout whole mounts with SOX9 and EDAR antibodies marking initial stages of
WF development [4], i.e. placode formation. These results revealed essentially identical whisker
patterning between control and Neurog1[-/-] mice (Figure 2B). EDAR staining performed at E12.5 10-μm
FFPE sections further confirmed normal placodal expression of EDAR, indicating normal initiation of the
WF development (Figure 2C). WF formation was also validated by micro-CT at E17.5 showing normal
progression of WF development at later stages (Figure 2D, top) even when trigeminal ganglia were
absent (Figure 2D, bottom, arrows). Overall, these data unambiguously show that whisker formation
does not require trigeminal sensory innervation and exclude the possibility that impaired nerves or
nerve-associated cells in *Meis2* cKO resulted in impaired WF induction. Therefore, it can be confidently
concluded that the dermal mesenchyme expressing *Meis2* is evidently critical for WF development.

In addition to whisker phenotype, we analyzed tooth morphology in Neurog1[-/-] mice since the role of
innervation for tooth development represents a long-standing controversy and has not been directly
tested previously[22–27]. Interestingly, we did detect any defects in tooth development (Supp Figure 2A)
which was corroborated also by micro-CT images from E17.5 mice (Supp Figure 2B), reporting an
innervation independent formation of teeth in Neurog1[-/-] mice.

Mesenchymal Meis2 operates upstream of epithelial EDAR signaling during initial steps of WF
formation

Our data strongly suggest that *Meis2* expression in the mesenchyme is required for induction of placodal
expression of Sox9 in WF. Since a number of regulators of HF development were identified to function
upstream of placodal Sox9 expression, we tested these successive molecular events leading to WF
induction. Shh release in basal placodal cells was shown to trigger Sox9 expression in suprabasal
placodal cells. This is achieved by asymmetric cell division resulting in suprabasal daughter cells with
low WNT activity so that Shh can drive Sox9 expression [28]. We performed RNA in situ hybridization to
evaluate *Shh* expression in *Meis2* cKO mice and control littermates at E13.5. These data revealed a
remarkably decreased number of *Shh*-positive foci in *Meis2* cKO mice (Figure 3A, arrow), showing that
placodal expression of *Shh* is regulated by mesenchymal MEIS2 protein. Again, we observed a small
number of Shh-positive WF escapers. EDA/NFκB signaling in placodes has been shown to act upstream
of Shh signaling before Pc invagination [4]. We therefore stained whole mount E12.5 snouts from *Meis2*

cKO mice and controls with Sox9 and EDAR antibodies. As seen in Figure 3B, EDAR protein was
localized to epithelial WF placodes and overlapped with SOX9 staining in controls. In contrast, in Meis2
cKO mutants, both EDAR and SOX9 were hardly detectable, documenting that MEIS2 mesenchymal
function is upstream of EDA signaling in Pc. Whole-mount EDAR staining was confirmed on tissue
sections which detected significantly reduced number of EDAR-positive foci in the *Meis2* cKO
epithelium. Moreover, EDAR-negative regions within the mutant epithelium did now show any signs of
morphological placode formation as inspected by DAPI staining (Figure 3C, Supp Figure 3A). Overall,
these data demonstrate that mesenchymal MEIS2 operates at very early stages of WF formation in
which it participates in forming a primary inductive mesenchymal signal for EDAR+ placode formation.
This early arrest in mutants affects all subsequent steps of WF development including *Shh*, *Sox9*
expression, and Pc and DC formation.

Meis2 deletion in NCC-derived cells affects epithelial but not dermal WNT signaling

It has been shown that Wnt/β-catenin signaling acts upstream of EDA/NFκB signaling as the earliest
molecular pathway identified during HF formation [4]. Epithelial WNT activity is required for activation of
dermal canonical WNT signaling, while dermal WNT signaling reciprocally affects patterned WNT
activity in the epithelium [29]. We therefore assessed canonical WNT signaling in *Meis2* cKO snouts by
several types of WNT signaling readouts. Firstly, we used LEF1 immunofluorescence because its
expression has been widely used as proxy for WNT signaling in HF [30,31] and *Lef1* is also required for
whisker formation [32]. Matos et al. [31] showed that dermal cells expressed Lef1 broadly whereas in the
epithelium, Lef1 expression was concentrated in Pc cells compared to inter follicular regions. During Pc
downgrowth in HF, basal Pc cells displayed LEF1 signal similarly to Shh and EDAR, in contrast to
suprabasal placodal cells lacking LEF1. This is consistent with the view that high WNT activity is seen
in the basal cells whereas suprabasal cells are WNT low [31]. In WF, we similarly detected robust LEF1
signal in Pc that was remarkably lower in interfollicular regions (Figure 4A, Supp Figure 3B, C). However,
in distal snout regions with no WFs, the epithelium showed confluent LEF1 signal (Supp Figure 3C)
suggesting that LEF1 signal is regionally dependent Pc indicator. In contrast to the epithelium, DC
underneath invaginating placodes consistently displayed decreased LEF1 staining compared to peri-
DC neighboring regions (Figure 4A, Supp Figure 3B). In *Meis2* cKO, we rarely detected a patterned
placodal increase in Lef1 staining in the expected sites of WF positions (Figure 4A, Supp Figure 3C).
This shows that failure of WF initiation in *Meis2* cKO is accompanied by missing expression of LEF1 in
Pc indicating that canonical WNT signaling is not induced in the Pc epithelium. On the other hand, LEF1
staining in *Meis2* cKO dermal fibroblasts was confluent and without down-regulation in presumptive DC
sites which again confirms early developmental arrest in mutant WFs.

Secondly, in order to verify WNT activity dynamics in developing WF, we employed Wnt reporter mouse
strain BAT-gal [33] in which multiple TCF-binding sites promote expression of *beta-galactosidase* gene.
We crossed this reporter mouse strain to *Meis2* cKO and stained snout frontal sections with anti-
galactosidase antibody. In contrast to LEF1 staining, BAT-gal reporter staining did not show consistent
upregulation and concentration of WNT activity in Pc and no corresponding downregulation in DC in
controls. It rather exhibited confluent expression in the dermal mesenchyme (Figure 4B) that was not
changed in mutants. BAT-gal reporter therefore did not serve as a reliable WNT readout in developing
WF in our hands probably due to high stability of galactosidase protein compared to RNA [34], which
limited the analysis of the dynamic nature of Wnt activity during WF development.

Thirdly, we tested the expression of selected Wnt ligands in *Meis2* cKO snouts. *Wnt10b* expression is
strongly activated in HF placodes [35] and it also is important for HF progression [36]. Thus, WNT10b seems
the most likely Wnt gene activating canonical pathway in HF which is routinely monitored by *Axin2*
expression in many tissue contexts including HF [10,28]. We performed HCR FISH to assess expression
of *Wnt10b* and WNT target gene *Axin2* in snouts at E12.5. Similarly to LEF1 staining, we observed *Axin2*

increase in wild-type Pc, while DC and dermal mesenchyme displayed uniform *Axin2* expression, which
did not accord with decreased LEF1 in DC (Figure 4C, Supp Figure 3B, D). This might reflect differential
responses of *Lef1* and *Axin2* expression to WNT activity in DC. In *Meis2* cKO, upregulation of *Axin2* in
the epithelium was not detected which most likely reflected missing placodes. On the other hand, mutant
dermal mesenchyme displayed broad *Axin2* expression which was similar to wild-type controls. *Wnt10b*
transcripts were detected in normal Pc in controls while this expression disappeared in missing Pc in
mutants (Figure 4C). Overall, these data suggest that mesenchymal MEIS2 does not affect WNT
signaling in the dermis, whereas it is critical for the patterned WNT upregulation in the epithelium. Since
WNT activation in the epithelium is one of the earliest steps of HF and WF development, this leads to
loss of EDAR and SHH markers in Meis2 cKO WF, and it confirms a very early role of mesenchymal
Meis2 during whisker development.

Meis2 regulates dermal cell proliferation and transition of fibroblast cell fate to DC

We have recently performed single cell RNA sequencing (scRNA seq) datasets for mesenchymal cells
derived from cranial neural crest of E12.5 and E13.5 Meis2 cKO embryos and their control littermates
(Hudacova et al. manuscript in revision, GSE262468). Cell cluster analysis identified one cluster
representing dermal fibroblasts. We subset and further re-clustered using standard Seurat workflow
(methods). We identified 6 clusters where Cluster 2 was defined as DC by cell markers Sox2, Sox18,
Tbx18, Bmp4, Lef1 and Cdkn1a (Figure 5A-B, Supp Figure 4A, Supplementary Data 1). The other
clusters were determined to represent fibroblasts annotated by their markers in the heatmap and module
scores generated for fibroblast (Fb), pre-DC, DC1 and DC2 markers
(https://github.com/kaplanmm/whisker_scRNA/tree/main.) [7,37]. (Supp Figure 4, Supplementary Data 1).
Cluster 2 displayed low expression of Fb module score genes and high expression of DC2 module score
genes confirming again that cluster 2 represents developed DCs. Transcriptomic analysis showed that
the cell number proportion in cluster 2 was substantially lower in *Meis2* cKO mice compared to control
littermates (Figure 5C), reflecting WF phenotype in the mutant. Interestingly, clusters 0 also contained
lower cell proportion in the *Meis2* cKO dataset. On the other hand, the cell number proportion in cluster
1 increased in *Meis2* cKO (Figure 5C). However, cell identity of cluster 1 was unclear because it was
marked by only 27 genes (Supplementary Data 1). Cluster 0 revealed GO terms related to cell cycle
and chromosome segregation/organization according to GO analysis by clusterProfiler R package [38]
using top 100 markers (Supplementary Data 2). Therefore, we conclude that cluster 0 very likely
represents dividing cells. Lower cell number proportion in cluster 0 in *Meis2* cKO suggested a Meis2-
dependent regulation of dermal cell proliferation. This hypothesis was tested by performing EdU chase
experiments where EdU was injected at E12.5 and embryos were harvested after 2 or 18 hours. Indeed,
the 2D area coverage of DAPI+ region by EdU+ region in the upper dermis of *Meis2* cKO snout
decreased from $52.96 \pm 2.05\%$ to $42.38 \pm 2.77\%$ (mean $\pm$ sem, t-test p= 0.0057) in 2-hour pulse and
from $52.17 \pm 0.81\%$ to $45.76 \pm 1.85\%$ (mean $\pm$ sem, t-test p= 0.0052) in 18-hour pulse (Figure 5D, E).
These EdU experiments confirmed scRNA-seq data and indicated a Meis2 function in proliferation of
upper dermal cells which might contribute to the aberrant whisker phenotype of the mutant.

Because cell migration is a critical process for the migration of Fb cell into DC during cell fate acquisition
[39], we plotted cell adhesion and ECM module scores, both of which were found to be increased in
mutants (Figure 5F, G). Thus, such an increase in cell interactions might limit Fb migration efficiency by
decreasing conductivity of the environment for cell migration and ultimately to decreased numbers of
WFs in mutants. Similar results were found by Hudacova et al. (manuscript in revision) in craniofacial
mesenchymal cells in *Meis2* cKO mice. These increases in cell adhesion and ECM gene expression
might also contribute to decreased TG axonal growth and branching described above (Figure 1, Supp
Video 1 and Supp Video 2).

Meis2 controls expression of *Foxd1* in the dermis

Next, we performed differential gene expression analysis between DC cluster of *Meis2* cKO and controls at E12.5 and E13.5. In this scRNA-seq dataset, *Foxd1* was found within the top genes downregulated in *Meis2* cKO mice (avg_log2FC = 1.0585, p_val = 4.55E-06) (Figure 6A, Supplementary Data 3). Interestingly, *Foxd1* expression was described as a hallmark of the earliest molecular changes in dermal fibroblasts prior to DC formation and defined pre-DC stage in HF [7]. Further, *Sox2*, a well-known DC marker, was also strongly downregulated in *Meis2* cKO (Figure 6A). To validate transcriptomic data, we analyzed FOXD1 protein expression in tissue sections of wild-type snouts at E12.5 and E13.5. Similarly to HF, *Foxd1* expression preceded *Sox2* also in WF which occurred before the Pc downgrowth into the dermal niche at E12.5. During initial Pc invagination and early DC stage at E13.5, *Foxd1* and *Sox2* expression overlapped (Figure 6B, arrows). In more progressed WF germs with deeply invaginated Pc, *Foxd1* expression remained in peri-DC regions rather than in DC that was marked by strong SOX2 signal (Figure 6B, right panels, arrowhead). Next, whole-mount staining of FOXD1 in the whole snout highlighted dynamic WF development along proximo-distal axis. It started in proximal columns at E12.5, in which WF formation was more progressed, while distal snout regions initiate WF slightly later (Figure 6C). By E13.5, 3-4 proximal columns displayed discernable WFs (Figure 6D). Strikingly, *Meis2* cKO snouts displayed no FOXD1+ WFs at E12.5 (Figure 6C). At E13.5, we detected around 4 proximal WFs in mutants while controls contained around 17 WFs on each snout side (Figure 6D). FOXD1+ WFs in *Meis2* cKO most likely represented 'escaper' WFs that were described above. Whole-mount FOXD1 immunostaining was further confirmed on tissue sections (Figure 6C, D, lower panels). Normal WF at E13.5 expressed Foxd1 in DC or peri-DC and Sox9 in Pc. In contrast, this progression of WF formation was halted in *Meis2* cKO (Figure 6C, D). These results suggest that Meis2 is the key component of transcription controlling machinery in the dermal mesenchymal that initiate WF.

Foxd1 is dispensable for WF development

Because *Foxd1* expression in DC is strictly regionally specific and its expression is lost in *Meis2* cKO, we hypothesized that Foxd1 transcription factor function is essential for WF formation. We therefore analyzed whiskers in embryos where Foxd1 expression is ablated. We crossed heterozygous *Foxd1*-GFP-CreERT2 mice, in which GFP-Cre-ERT2 cassette was inserted into initiation codon of *Foxd1* gene. Homozygous embryos are therefore null mutants for *Foxd1* gene. WF development was assessed in whole-mount stained embryonic heads at E13.5 using SOX9, FOXD1 and TUJ1 antibodies. Figure 7A shows that SOX9 was detected in invaginating placodes, FOXD1 in surrounding DC mesenchyme and TUJ1 in axonal projections innervating each WF. In parallel, we stained EDAR in WF placodes (Figure 7B). Interestingly, the number and morphology of WF in controls and *Foxd1* mutants were essentially identical. Thus, FOXD1 protein is a very important marker of the early DC cell fate in WF but does not have functional relevance for whisker formation despite its dynamic and regionally specific expression pattern around the developing WFs.

Discussion

We have shown that that mesenchymal expression of transcription factor Meis2 is essential for the initial steps of mouse whisker development. Mouse embryos, in which *Meis2* was deleted in tissues derived from neural crest cells by Wnt1-Cre2 driver, lost majority of whiskers. The aberrant whisker phenotype in the mutants is accompanied by markedly reduced trigeminal nerve branching in the snout. This suggests a novel function of Meis2 in the trigeminal axonal growth, branching and guidance. It may also implicate that whisker formation, or WF positional determination, may depend on early sensory innervation. We report here, however, that whisker formation is entirely independent from sensory innervation. *Neurog1*[-/-] mice completely lack trigeminal nerves including their ganglions but whisker

development is not affected. Moreover, analysis of Neurog1[-/-] mice at E17.5 reveals that sensory
innervation is not required for the further growth and stabilization of WFs. The effect of trigeminal
innervation deficiency on postnatal WF development could not be tested because Neurog1[-/-] mice are
lethal within a day after birth. All these experiments exclude the possibility that whisker phenotype in
Meis2 cKO embryos is an indirect consequence of decreased trigeminal nerve branching. It rather
indicates mesenchymal Meis2 is a pivotal regulator of WF formation.

Our observations suggest that nerve branching phenotype in the mutant is independent from trigeminal
ganglia development or missing whiskers as *Meis2* cKO mice showed normal trigeminal ganglia size
and nerve exit from the ganglia as well as defective TG nerve branches that do not innervate whiskers.
Thus, MEIS2 rather influences trigeminal branching and guidance through the snout. Description of the
exact nature of factors controlling trigeminal nerve branching downstream of MEIS2 warrants further
investigation. A number of molecular pathways, such as Slit/Robo[40,41], Sema/Nrp/Plxn[42–46], Ephrin/Eph[47]
signaling, have been implicated during TG nerve development and Meis2 might act upstream of
mesenchymal expression of ligands and/or trigeminal expression of receptors of these pathways to
control TG nerve development. One possible mechanism might involve Sema3a, an inhibitory factor for
trigeminal nerve growth during tooth innervation and *Sema3a* null mice display hyper-branched
trigeminal nerves[45,48,49]. Our scRNA seq datasets indicated an increase of *Sema3a* expression in *Meis2*
cKO mesenchymal cell populations (data not shown) which could explain decreased nerve branching in
Meis2 cKO mice and could be addressed in future studies.

Essential role of MEIS2 in the dermal mesenchyme during very early steps of WF development is
supported by several key observations. Firstly, although epithelial expression of *Meis2* is preserved in
Meis2 cKO mice, thickening of the epithelium is rarely observed suggesting that the release of WF
placode-inducing dermal factors is under control of mesenchymal Meis2. It is widely accepted that HF
or WF formation is initiated from the dermis by a 'first dermal signal' that has not been identified yet.
Our data show that this process is dependent on MEIS2. Secondly, the absence of MEIS2 results in the
loss of early placodal markers Lef1 and EDAR and pre-DC marker Foxd1 at expected WF locations.
Absence of patterned upregulation of Lef1 and EDAR was documented in HFs after blocking dermal
WNT signaling [29]. Interestingly, our analysis of WNT signaling readouts did now show remarkable
changes in canonical WNT signaling in Meis2 cKO dermis, indicating that normal dermal WNT signaling
is not sufficient to trigger patterned Lef1 and EDAR upregulation in the epithelium when dermal MEIS2
is absent. These epithelial changes probably require additional MEIS2-dependent intermediate steps,
which might act either in parallel with, or downstream of dermal WNT signaling to induce placode
formation. In the dermis, pre-DC marker Foxd1 expression was shown to be dependent on Fgf20
expression in HF Pc and occurs after Pc formation [7]. Our analysis of *Foxd1* expression in wild-type
snout dermis revealed that Foxd1 expression appears prior to morphologically distinguishable WF Pc.
In *Meis2* cKO, FOXD1 signal in the dermis prior to WF Pc formation was not detected which accords
with single-cell RNA seq data. This all suggests that MEIS2 operates upstream of *Foxd1*. However,
direct analysis of Foxd1-null mice did not reveal any phenotype in WF development. This could be
explained by either no function of FOXD1 in this process or a possibly functional compensation by its
paralog FOXD3. Lastly, it has been shown in HF development that dermal fibroblasts proliferate
extensively prior to converting to DC cell fate. DC cells, on the other hand, exit cell cycle and become
quiescent [7,10]. We generally observed reduced proliferation in upper dermal cells in *Meis2* cKO mice.
Lower cell proliferation could contribute to inefficient cell fate change of Fb to DC and might also lead to
insufficient expression and/or release of Pc-inducing factor in the dermis of *Meis2* mutants. Overall,
these observations strongly suggest that absence of *Meis2* in mesenchymal cells affects earliest
processes of WF development, even prior to Pc formation.

It is important to note that we observed small numbers (around 18% at E13.5) of WFs in *Meis2* cKO
expressing normal placodal and DC markers. These rarely formed WF 'escapers' in *Meis2* cKO mice

had normal WF morphology, because their placode thickness, follicle length and DC size were
comparable between controls and mutants at E12.5-E13.5. WF 'escapers' in mutants were probably
not sensitive to Wnt1-Cre2-mediated deletion of *Meis2*. Although immunostaining of MEIS2 in snout
sections indicated that Meis2 expression around WF escapers was hardly detectable in mutants, a
slight but undetectable expression of Meis2 resulting from incomplete gene deletion might account for
normal WF formation. On the other hand, it is also possible that *Meis1* paralogue can partially
substitute missing *Meis2* in its function in the dermal mesenchyme. Alternatively, MEIS2 function in
WF development might be restricted to a short time window as each WF develops at different time
points along proximal-distal axis.

Materials and Methods

Mouse strains

Generation of the floxed allele of *Meis2* gene (*Meis2* fl/fl) was described by Machon et al. (2015).[50]
*Meis2*fl/fl strain was crossed with reporter mouse lines R26R-*EYFP* fl/fl (RRID:IMSR JAX006148) before
crossing with the Wnt1-Cre2 [18]. This reporter enabled monitoring Cre recombination specificity. Wnt1–
Cre2 (RRID:IMSR JAX:022137) was used for specific deletion of the *Meis2* fl/fl gene in neural crest cells.
For lineage tracing experiments, we used Wnt1-Cre2 crossed with mTmG (RRID:IMSR JAX:007676).
During embryo harvesting, Cre recombination was checked by monitoring EYFP fluorescence. In very
rare cases (1-2 embryos in 10 litters), Cre activity in the whole body was observed indicating germline
recombination. Such animals were removed from the analysis and their parents were removed from
crossing   schemes.   Neurogenin1   knockout   mice   were   purchased   from   Jax   Lab
(RRID:IMSR_JAX:017306).   *Foxd1*-GFP-Cre-ERT2   mice   were   purchased   from   Jax   Lab
(RRID:IMSR_JAX:012464). The presence of the vaginal plug was regarded as the embryonic day 0.5
(E0.5). Pregnant mice were anesthetized using isoflurane gas inhalation and euthanized by cervical
dislocation. Embryos were dissected and placed in ice-cold PBS. All mice were genotyped as previously
described.[50] Neurog1 and Foxd1 mutant mice were genotyped according to Jax protocol. All procedures
involving experimental animals were approved by the Institutional Committee for Animal Care and Use
(permission #PP-10/2019) with every effort to minimize the suffering and the number of animals used.
This work did not include human subjects.

Immunohistochemistry and image processing

All embryos used for staining were harvested at either E12.5 or E13.5 embryonic stages and heads
were fixed in 4% paraformaldehyde (PFA) overnight at 4^0C. After PFA fixation samples were prepared
in three ways. (1) For FFPE staining, after dehydration in ethanol and xylene, samples were embedded
in paraffin and frontal 10-µm sections were prepared. Followed by stepwise rehydration, antigen retrieval
in 0.1 M citrate buffer pH 6.0 under pressure boiling for 12 min was carried out. (2) For cryosection
staining, after dehydration in 30% saccharose for 24 hours, samples were embedded in OCT and 100
µm cryo-sections were obtained. (3) For whole mount staining, whole heads were directly processed to
permeabilization and blocking after PFA fixation.

Tissues were blocked in 5% bovine serum albumin BSA in PBS with 0.1% Triton x-100 for FFPE
sections or with 0.5% Triton x-100 for thick cryosections or whole mounts for 1 hour and incubated in
primary antibody solution (1% BSA in PBS and 0.1% Triton x-100 for 10-µm section or 0.5% Triton x-
100 for 100-µm sections and whole mounts). Primary antibody incubation was overnight for FFPE
section, two nights for cryosections and three nights for whole mount staining at 4^0C. Primary antibodies
used: Sox9 (Merck Sigma, AB5535), TrkA (R&D Systems, AF1056), Meis2 (GeneScript, custom), Tuj1
(R&D Systems, MAB1195), EDAR (R&D Systems, AF745), Lef1 (Cell Signaling, C12A5), β-
galactosidase (Abcam, ab9361), Sox2 (R&D Systems, MAB2018), Foxd1 (Abcam, AB129324). Primary
antibody solutions were washed out with (PBS, Triton X-100) and incubated in fluorescent secondary
antibodies solutions for one hour for FFPE sections, three hours for cryosections at room temperature
or overnight for whole mounts at 4^0C. Secondary antibodies: donkey anti-mouse, -rabbit, -goat with

Alexa488, 568 or 647 fluorophores (ThermoFisher Scientific). Primary and secondary antibodies were
used at 1:500 for thick section and whole mount staining and at 1:1000 for thin section staining.
Immunofluorescent images were scanned by spinning disc confocal microscopy using an Olympus
SpinSR10. Imaging was performed for whole mount stainings after tissue clearing for one hour at room
temperature. Acquired z-stacks were maximum intensity projected using ImageJ. Figures were
assembled using Photoshop.

For cell proliferation assays, EdU was injected intraperitoneally to pregnant females at E12.5 for 2 or 18
hours before harvesting (0.5 ml per mouse, 2.5 mg/ml). EdU Base Click kit was used for EdU detection
according to manufactures protocol on FFPE sections.

Micro-CT

Embryos were fixed in 4% PFA for 2 days and soaked in Lugol's iodine for several days. Scanning was
performed on the instrument Bruker Skyscan 1272 with the resolution 3 µm.
434
In situ hybridization

Antisense RNA probe for Shh was cloned into pGEM-T-easy vector (Promega) using primers: Shh-F
TCACAAGTCCTCAGGTTCCG, Shh-R GGGCTTCAGCTGGACTTGAC. Antisense mRNA was
transcribed with T7 polymerase. Whole-mount *in situ* hybridization was performed using standard
protocols. Fluorescent in situ hybridization HCR RNA-FISH (Molecular Instruments) was performed
according to manufacturer's protocols (company cat#??).

Single Cell RNA sequencing analysis

scRNA sequencing datasets are retrieved from integrated Seurat object (GSE262468) and processed
with standard Seurat workflow[51]. Briefly, dermal fibroblast cluster was subset and split by condition which
was followed by SCTransform normalization where mitochondrial, ribosomal and cell cycle genes were
regressed. Seurat objects were then integrated by using IntegrateData function with 3000 integration
features. For UMAP as non-linear dimensional reduction embedding, 30 principal components were
used. We used Clustree package to determine optimal resolution which was set at 0.6 for FindClusters
function according to developers' instructions[52]. SCT assay was used to generate feature plots and
violin plots. Normalized and Scaled data of the RNA assay were used to extract cluster markers using
FindAllMarkers function and to generate heatmaps. FindMarkers function of Seurat package was used
for differential gene expression analysis between conditions with default parameters. clusterProfiler
(4.6.2) was used for GO analysis [38], where GO terms were extracted by enrichGO function. This paper
did not produce original codes. Codes used for analysis and list of genes used to generate module
scores can be accessed at https://github.com/kaplanmm/whisker_scRNA/tree/main.

Quantification and Statistical Analysis

For placode thickness/WF length and DC size analysis, images of DAPI stained 10 µm FFPE snout
sections were captured with Olympus SpinSR10 microscope and 20X objective. With ImageJ software,
maximum intensity projection was applied to z-stack images. DAPI staining was used to visually identify
placodes and DCs. The distance from the outer layer of epithelium in the placodal region to epithelial-
mesenchymal boundary was measured. For 2D DC size, DCs were identified by densely accumulated
cells underneath the placodes. The area of the region covered by these cells was measured using
ImageJ.

The number of WFs were manually counted in embryos that were used throughout the study for whole
mount immunostainings (labeled by SOX9, EDAR and/or FOXD1 antibodies) or *in situ* hybridization
(*Shh* mRNA). Numbers of WFs at both sides of the snout were pooled. The decrease in percentage of
WFs in *Meis2* cKO mice was calculated in comparison to their control littermates. Values from each
experiment were averaged to determine the overall decrease in the percentage of WFs.

For EdU/DAPI area measurements, images of EdU chased and DAPI stained 10 µm FFPE snout
sections were captured with Olympus SpinSR10 microscope equipped with 20X objective. With ImageJ

software, maximum intensity projection was applied to z-stack images. Dermal region between the
epithelium and around 100-µm distance into dermis was used for analysis, while DC domains were
excluded. The selected dermal regions were processed separately for thresholding steps to cover EdU+
and DAPI+ regions. Threshold values were determined visually to cover specific EdU and DAPI staining.
The same threshold values were used for control and *Meis2* cKO mutants. Total area values for EdU
and DAPI were used to calculate 2D EdU coverage of the DAPI regions (EdU/DAPI).

Statistical analyses were performed using GraphPad Prism 10 software. N numbers specified in the
respective figure legends for each analysis. All the data are presented with mean ± sem (standard error
of the mean). Two-tailed student t-test was used to explore statistical differences between two
experimental groups (Control vs *Meis2* cKO).

Acknowledgements

This work was supported by Czech Science Foundation (grants 22-10660S, 23-06160S) and Masaryk
University, Faculty of Medicine (MUNI/A/1598/2023). We thank the Microscopy Service Centre of IEM
CAS and Light Microscopy Core Facility, IMG CAS, Prague, Czech Republic, supported by MEYS –
LM2023050 and RVO – 68378050-KAV-NPUI. This work was also funded by MEYS Czech Republic
(NanoEnviCZ, LM2018124) and EU Structural Funds Pro-NanoEnviCz
(CZ.02.1.01/0.0/0.0/16_013/0001821).

Author contributions

488 MMK designed and performed experiments, bioinformatics and wrote the manuscript. EH performed
489 experiments and bioinformatics. MM performed experiments. XX performed experiments, JK designed
490 experiments, OM designed and performed experiments, wrote manuscript, and provided funding. The
491 authors declare no competing interests.

Figure legends:

Figure 1. **Severe whisker follicle development in Meis2 cKO embryos. (A)** micro-CT images of
control and Meis2 cKO embryos at E15.5 showing aberrant whisker phenotype in *Meis2* cKO mice. **(B)**
Light-sheet microscopy of Sox9 whole-mount immunostaining of WF. Arrow indicates example of an
"escaper" whisker. **(C)** MEIS2 immunofluorescence on frontal frozen section of E13.5 snouts showing
Meis2 expression in dermis, DC and epithelium including Pc. Arrowheads indicate epithelial expression
of *Meis2* in both control and *Meis2* cKO mice. Arrow and asterisk show disappearing *Meis2* expression
in the DC and dermis, respectively, in the mutant. Scale bar: 150 µm. **(D)** Wnt1-Cre; *mTmG* embryos
showing Cre recombination specificity in the craniofacial area, midbrain and dorsal spinal cord.
Recombined cells are labeled by membrane-localized GFP in green and non-recombined cells labeled
by membrane-localized tdTomato in red. **(E)** Frontal sections of Wnt1-Cre; *mTmG* snouts documenting
Cre recombination in the neural crest-derive mesenchyme, cranial nerve projections without
recombination in the overlying epithelium. **(F)** Whole-mount staining of *Meis2* cKO heads at E13.5 with
SOX9 and TRKA (Ntrk1) antibodies showing almost absence of WF in mutants and compromised
branching of the trigeminal nerve. Scale bar: 500 µm.

Figure 2. *Neurog1[-/-]* **(KO) embryos lack the trigeminal nerve but WF development is normal. (A)**
Triple immunostaining of 100-µm sections shows absence of trigeminal nerve projections (Tuj1+, TrkA+)
and normal WF (Sox9+) in Neurog1[-/-] mice. Arrows in black and white images show examples of
invagination of whisker placodes labeled by Sox9 antibody. Scale bar: 500 µm. **(B)** Whole-mount
immunostaining of WFs with SOX9 and EDAR antibodies showing normal WF morphology and
patterning in mutants at E12.5 (top) and 13.5 (bottom). Scale bars: 300 µm. **(C)** EDAR staining of 10-
µm sections at E12.5 showing normal initiation of placode formation in Neurog1 mutants. Scale bar: 150

μm. **(D)** micro-CT images reveal normally developed whiskers (top) at E17.5 in mutants while TGs are
lacking (bottom, arrows).

Figure 3. **Induction and progression of WF development is compromised in *Meis2* cKO. (A)** Whole-
mount *in situ* hybridization of *Shh* mRNA documenting loss of WFs in mutants. Arrow shows an escaper
whisker. **(B)** Whole-mount immunostaining of WFs with SOX9 and EDAR antibodies showing absence
of WF in Meis2 cKO at E12.5. Two examples for each genotype are shown. Scale bar: 300 μm. **(C)**
EDAR staining of 10-μm sections showing placode formation arrest in mutants. Scale bar: 150 μm.

Figure 4. **WNT signaling in the epithelium is affected by deletion of *Meis2* in the dermal**
mesenchyme. (A) LEF1 staining of 10-μm paraffin frontal sections of snouts showing abundance of
LEF1 in the dermal mesenchyme while in the epithelium it is concentrated in placodes in regions of WF
appearance. Similar to missing placodes, LEF1 is also lost in *Meis2* cKO epithelium. Scale bars: 150
μm. **(B)** beta-galactosidase immunostaining of sections from WNT reporter BAT-gal controls and Meis2
cKO;BAT-gal mutants showing widespread galactosidase signal in the dermis and epithelium. Scale
bars: 150 μm. **(C)** *in situ* hybridization HCR-FISH using probes for *Axin2* and *Wnt10b*. In the epithelium,
Wnt10b is detected in placodes and the signal is lost in Meis2 cKO. Similarly, upregulation of *Axin2* is
not observed in the expected WF loci. In the dermal mesenchyme, *Axin2* mRNA is detected throughout
dermis regardless of WFs. This pattern was not changed in *Meis2* cKO. Scale bar: 50 μm.

Figure 5. **Dermal condensation and proliferation are reduced in *Meis2* cKO. (A) Uniform Manifold**
Approximation and Projection (UMAP) diagram of single-cell RNA seq analysis of the dermal
mesenchyme subset with 6-cluster resolution in which cluster 2 (green) represents dermal condensate
(DC) of WFs. **(B)** UMAP representations of typical DC markers *Sox2*, *Sox18*, *Tbx18*, *Bmp4, Lef1* or
Cdkn1 shown by FeaturePlots. **(C)** Relative cell numbers in 6 clusters showing a higher cell count in
cluster 1 (nonspecialized cells) at the expense of clusters 0 (dividing cells) and 2 (DC). **(D)** Analysis of
cell proliferation by EdU incorporation after 2-hour pulse at E12.5 and 18-hour pulse at E13.5. **(E)** Overall
proliferation rate was reduced in *Meis2* cKO from 52.96 ± 2.05% to 42.38 ± 2.77% (mean ± sem, t-test
p= 0.0057) in 2-hour pulse (top) and from 52.17 ± 0.81% to 45.76 ± 1.85% (mean ± sem, t-test p=
0.0052) in 18-hour pulse (bottom). Each data point indicates the average value for one section. n = 2
mice and at least 9 sections. **(F)** Increase of cell adhesion module score in mutants at E13.5 by scRNA
seq analysis. Module scores are generated by indicated genes. **(G)** Increase of extracellular matrix
(ECM) module score in mutants at E13.5 by scRNA seq analysis. Module scores are generated by
indicated genes.

Figure 6. **Expression of the earliest pre-DC and DC marker Foxd1 is lost in *Meis2* cKO. (A)** UMAP
representations of decline in *Foxd1* (top) and *Sox2* (bottom) expression in DC cluster (#2) shown by
FeaturePlots. Expression of both genes is significantly reduced in mutants. **(B)** Immunofluorescence of
FOXD1 and SOX2 at initial stages of WF showing FOXD1 in pre-DC stage while the expression shifts
from DC to peri-DC regions during WF progression. SOX2 appears as a typical DC marker with no
earlier expression. Scale bar: 50 μm. **(C-D)** Whole-mount immunostaining of FOXD1 and SOX9 of heads
from controls and Meis2 cKO at E12.5 **(C)** and E13.5 **(D).** It shows absence of WF including pre-DC
marker FOXD1 at E12.5 stage when proximal columns of WFs have already formed. A few WF escapers
develop in mutants at E13.5 in proximal regions while at least 6 WF columns have formed in controls
Scale bars: 400 μm (C) and 500 μm (D). Lower panels: FOXD1 staining in FFPE sections. Scale bars:
150 μm.

Figure 7. **WFs normally develop in loss-of-function *Foxd1* mutants. (A)** Whole-mount
immunostaining of FOXD1, TUJ1 and SOX9 of heads from controls and Foxd1 null mutants at E13.5
showing normal formation of WFs in mutants in which FOXD1 signal disappears. Normal WF
development is also reflected in normal WF innervation represented by TUJ1 staining. Scale bars: 500

μm. **(B)** Whole-mount immunostaining of EDAR confirmed normal Pc appearance in Foxd1 null mutants.
Scale bars: 500 μm.

Supplementary Figure 1. **Normal sizes of escaper WFs and TGs in Meis2 cKO mice. (A-B)**
Quantification of 2D DC area (A) and placode (Pc) thickness of non-invaginated or length of the
invaginated follicles (B) showing comparable values between control and Meis2 cKO at E12.5 (left) and
E13.5 (right). N = 4 embryos for E12.5 and 3 embryos for E13.5, and at least 30 WFs for controls and
10 for Meis2 cKO mice. P-values for all graphs are larger than 0.05, student t-test. Data are presented
as mean ± sem. **(C)** 100-μm frozen sections through TG ganglia stained with TUJ1 and TRKA
antibodies showing non-defective TG ganglion size or exit of the TG nerve from the ganglia. Scale bar:
250 μm.

Supplementary Figure 2. **Normal tooth development in Neurog1$^{-/-}$ mice. (A)** 100-μm frozen sections
of molar teeth of control and Neurog1$^{-/-}$ mice at E13.5 stained with SOX9, TUJ1 and TRKA antibodies
showing normal phenotype of the mutant. Scale bar: 150 μm. **(B)** microCT images around molar tooth
level at E17.5 in Neurog1$^{-/-}$ mice.

Supplementary Figure 3. **EDAR and LEF1 expression in Meis2 cKO mice. (A)** Examples of 10-μm
FPPE sections stained with EDAR antibody showing affected expression of EDAR in Meis2 cKO mice.
Arrows indicate EDAR positive sites which are observed less frequently in the mutant snout. Scale bar:
500 μm. **(B)** Representative images of normally developed escaper WFs in the mutants by Lef1 staining.
Lef1 expression is similar in just forming placodes (top) and invaginated (bottom). Additionally, Lef1
expression shows a decline in the DC regions compared to peri-DC. Scale bar: 30 μm. **(C)**
Representative micrographs of Lef1 stained 10-μm FFPE snout sections showing normal expression of
Lef1 in the dermis whereas decreased number of placodes and therefore Lef1 expression in the
epithelium of the mutant snout. Scale bar: 500 μm. **(D)** *In situ* hybridization HCR-FISH using probes for
Axin2 and *Wnt10b* in control snout at E12.5 showing placodal expression of *Wnt10b* (green) and *Axin2*
(arrows, red) as well as widespread expression of *Axin2* in the dermis. Scale bar: 60 μm.

Supplementary Figure 4. **Identification of DC cluster in scRNA sequencing datasets. (A)** Heatmap
of top 10 markers for each identified cluster in scRNA seq data. Each column represents individual cells
whereas each row represents indicated marker genes on left. Genes in the black box are top markers
of cluster 2 (DC). **(B)** FeaturePlots and VlnPlots of fibroblasts (Fb), pre-DC, DC1 and DC2 module
scores in each cluster. See Methods for selected genes generating module scores.

References

1.   Wrenn, J.T., and Wessells, N.K. (1984). The early development of mystacial vibrissae in the
mouse. J.Embryol.Exp.Morphol. *83*, 137–156.

2.   Sennett, R., and Rendl, M. (2012). Mesenchymal-epithelial interactions during hair follicle
morphogenesis and cycling. Semin Cell Dev Biol *23*, 917–927.
https://doi.org/10.1016/j.semcdb.2012.08.011.

3.   Huelsken, J., Vogel, R., Erdmann, B., Cotsarelis, G., and Birchmeier, W. (2001). beta-Catenin
controls hair follicle morphogenesis and stem cell differentiation in the skin. Cell *105*, 533–545.
https://doi.org/10.1016/s0092-8674(01)00336-1.

4.   Zhang, Y., Tomann, P., Andl, T., Gallant, N.M., Huelsken, J., Jerchow, B., Birchmeier, W., Paus,
R., Piccolo, S., Mikkola, M.L., et al. (2009). Reciprocal Requirements for EDA/EDAR/NF-κB and
Wnt/β-Catenin Signaling Pathways in Hair Follicle Induction. Developmental Cell *17*, 49–61.
https://doi.org/10.1016/j.devcel.2009.05.011.

5. Huh, S.-H., Närhi, K., Lindfors, P.H., Häärä, O., Yang, L., Ornitz, D.M., and Mikkola, M.L. (2013). Fgf20 governs formation of primary and secondary dermal condensations in developing hair follicles. Genes Dev *27*, 450–458. https://doi.org/10.1101/gad.198945.112.

6. St-Jacques, B., Dassule, H.R., Karavanova, I., Botchkarev, V.A., Li, J., Danielian, P.S., McMahon, J.A., Lewis, P.M., Paus, R., and McMahon, A.P. (1998). Sonic hedgehog signaling is essential for hair development. Current Biology *8*, 1058–1069. https://doi.org/10.1016/S0960-9822(98)70443-9.

7. Mok, K.-W., Saxena, N., Heitman, N., Grisanti, L., Srivastava, D., Muraro, M.J., Jacob, T., Sennett, R., Wang, Z., Su, Y., et al. (2019). Dermal Condensate Niche Fate Specification Occurs Prior to Formation and Is Placode Progenitor Dependent. Dev Cell *48*, 32-48.e5. https://doi.org/10.1016/j.devcel.2018.11.034.

8. Manti, P.G., Darbellay, F., Leleu, M., Coughlan, A.Y., Moret, B., Cuennet, J., Droux, F., Stoudmann, M., Mancini, G.-F., Hautier, A., et al. (2022). The Transcriptional Regulator Prdm1 Is Essential for the Early Development of the Sensory Whisker Follicle and Is Linked to the Beta-Catenin First Dermal Signal. Biomedicines *10*, 2647. https://doi.org/10.3390/biomedicines10102647.

9. Sennett, R., Wang, Z., Rezza, A., Grisanti, L., Roitershtein, N., Sicchio, C., Mok, K.W., Heitman, N.J., Clavel, C., Ma'ayan, A., et al. (2015). An Integrated Transcriptome Atlas of Embryonic Hair Follicle Progenitors, Their Niche, and the Developing Skin. Dev Cell *34*, 577–591. https://doi.org/10.1016/j.devcel.2015.06.023.

10. Gupta, K., Levinsohn, J., Linderman, G., Chen, D., Sun, T.Y., Dong, D., Taketo, M.M., Bosenberg, M., Kluger, Y., Choate, K., et al. (2019). Single-Cell Analysis Reveals a Hair Follicle Dermal Niche Molecular Differentiation Trajectory that Begins Prior to Morphogenesis. Developmental Cell *48*, 17-31.e6. https://doi.org/10.1016/j.devcel.2018.11.032.

11. Ge, W., Tan, S.-J., Wang, S.-H., Li, L., Sun, X.-F., Shen, W., and Wang, X. (2020). Single-cell Transcriptome Profiling reveals Dermal and Epithelial cell fate decisions during Embryonic Hair Follicle Development. Theranostics *10*, 7581–7598. https://doi.org/10.7150/thno.44306.

12. Mäkelä, O.J.M., and Mikkola, M.L. (2023). Mesenchyme governs hair follicle induction. Development *150*, dev202140. https://doi.org/10.1242/dev.202140.

13. Yang, J.W., Kilb, W., Kirischuk, S., Unichenko, P., Stuttgen, M.C., and Luhmann, H.J. (2018). Development of the whisker-to-barrel cortex system. Curr.Opin.Neurobiol. *53*, 29–34. https://doi.org/10.1016/j.conb.2018.04.023.

14. Zhao, H., Feng, J., Seidel, K., Shi, S., Klein, O., Sharpe, P., and Chai, Y. (2014). Secretion of shh by a neurovascular bundle niche supports mesenchymal stem cell homeostasis in the adult mouse incisor. Cell Stem Cell *14*, 160–173. https://doi.org/10.1016/j.stem.2013.12.013.

15. Adameyko, I., and Fried, K. (2016). The Nervous System Orchestrates and Integrates Craniofacial Development: A Review. Front Physiol *7*, 49. https://doi.org/10.3389/fphys.2016.00049.

16. Kaukua, N., Shahidi, M.K., Konstantinidou, C., Dyachuk, V., Kaucka, M., Furlan, A., An, Z., Wang, L., Hultman, I., Ahrlund-Richter, L., et al. (2014). Glial origin of mesenchymal stem cells in a tooth model system. Nature *513*, 551–554. https://doi.org/10.1038/nature13536.

17. Glover, J.D., Wells, K.L., Matthäus, F., Painter, K.J., Ho, W., Riddell, J., Johansson, J.A., Ford,
M.J., Jahoda, C.A.B., Klika, V., et al. (2017). Hierarchical patterning modes orchestrate hair
follicle morphogenesis. PLoS Biol *15*, e2002117. https://doi.org/10.1371/journal.pbio.2002117.

18. Lewis, A.E., Vasudevan, H.N., O'Neill, A.K., Soriano, P., and Bush, J.O. (2013). The widely used
Wnt1-Cre transgene causes developmental phenotypes by ectopic activation of Wnt signaling.
Dev.Biol. *379*, 229–234. https://doi.org/10.1016/j.ydbio.2013.04.026.

19. Fabik, J., Kovacova, K., Kozmik, Z., and Machon, O. (2020). Neural crest cells require Meis2 for
patterning the mandibular arch via the Sonic hedgehog pathway. Biol.Open. *9*.
https://doi.org/10.1242/bio.052043.

20. Leonard, C.E., Quiros, J., Lefcort, F., and Taneyhill, L.A. (2022). Loss of Elp1 disrupts trigeminal
ganglion neurodevelopment in a model of familial dysautonomia. Elife. *11*.
https://doi.org/10.7554/eLife.71455.

21. Ma, Q., Chen, Z., del, B.B., I., de la Pompa, J.L., and Anderson, D.J. (1998). neurogenin1 is
essential for the determination of neuronal precursors for proximal cranial sensory ganglia. Neuron
20, 469–482. https://doi.org/10.1016/s0896-6273(00)80988-5.

22. Løes, S., Kettunen, P., Kvinnsland, I., and Luukko, K. (2002). Mouse rudimentary diastema tooth
primordia are devoid of peripheral nerve fibers. Anat Embryol *205*, 187–191.
https://doi.org/10.1007/s00429-002-0247-8.

23. Lumsden, A.G., and Buchanan, J.A. (1986). An experimental study of timing and topography of
early tooth development in the mouse embryo with an analysis of the role of innervation. Arch Oral
Biol *31*, 301–311. https://doi.org/10.1016/0003-9969(86)90044-0.

24. Stainier, D.Y., and Gilbert, W. (1990). Pioneer neurons in the mouse trigeminal sensory system.
Proc Natl Acad Sci U S A *87*, 923–927. https://doi.org/10.1073/pnas.87.3.923.

25. Maeda, Y., Miwa, Y., and Sato, I. (2019). Distribution of the neuropeptide calcitonin gene-related
peptide-α of tooth germ during formation of the mouse mandible. Ann Anat *221*, 38–47.
https://doi.org/10.1016/j.aanat.2018.09.001.

26. Weil, M., Itin, A., and Keshet, E. (1995). A role for mesenchyme-derived tachykinins in tooth and
mammary gland morphogenesis. Development *121*, 2419–2428.
https://doi.org/10.1242/dev.121.8.2419.

27. Duan, Y., Liang, Y., Yang, F., and Ma, Y. (2022). Neural Regulations in Tooth Development and
Tooth-Periodontium Complex Homeostasis: A Literature Review. Int J Mol Sci *23*, 14150.
https://doi.org/10.3390/ijms232214150.

28. Ouspenskaia, T., Matos, I., Mertz, A.F., Fiore, V.F., and Fuchs, E. (2016). WNT-SHH Antagonism
Specifies and Expands Stem Cells prior to Niche Formation. Cell *164*, 156–169.
https://doi.org/10.1016/j.cell.2015.11.058.

29. Chen, D., Jarrell, A., Guo, C., Lang, R., and Atit, R. (2012). Dermal β-catenin activity in response
to epidermal Wnt ligands is required for fibroblast proliferation and hair follicle initiation.
Development *139*, 1522–1533. https://doi.org/10.1242/dev.076463.

30. DasGupta, R., and Fuchs, E. (1999). Multiple roles for activated LEF/TCF transcription complexes
during hair follicle development and differentiation. Development *126*, 4557–4568.
https://doi.org/10.1242/dev.126.20.4557.

31. Matos, I., Asare, A., Levorse, J., Ouspenskaia, T., de la Cruz-Racelis, J., Schuhmacher, L.-N., and
Fuchs, E. Progenitors oppositely polarize WNT activators and inhibitors to orchestrate tissue
development. eLife *9*, e54304. https://doi.org/10.7554/eLife.54304.

32. Kratochwil, K., Dull, M., Farinas, I., Galceran, J., and Grosschedl, R. (1996). Lef1 expression is
activated by BMP-4 and regulates inductive tissue interactions in tooth and hair development.
Genes Dev *10*, 1382–1394. https://doi.org/10.1101/gad.10.11.1382.

33. Maretto, S., Cordenonsi, M., Dupont, S., Braghetta, P., Broccoli, V., Hassan, A.B., Volpin, D.,
Bressan, G.M., and Piccolo, S. (2003). Mapping Wnt/beta-catenin signaling during mouse
development and in colorectal tumors. Proc.Natl.Acad.Sci.U.S.A *100*, 3299–3304.
https://doi.org/10.1073/pnas.0434590100.

34. Machon, O., Backman, M., Machonova, O., Kozmik, Z., Vacik, T., Andersen, L., and Krauss, S.
(2007). A dynamic gradient of Wnt signaling controls initiation of neurogenesis in the mammalian
cortex and cellular specification in the hippocampus. Dev.Biol. *311*, 223–237.
https://doi.org/10.1016/j.ydbio.2007.08.038.

35. Reddy, S., Andl, T., Bagasra, A., Lu, M.M., Epstein, D.J., Morrisey, E.E., and Millar, S.E. (2001).
Characterization of Wnt gene expression in developing and postnatal hair follicles and
identification of Wnt5a as a target of Sonic hedgehog in hair follicle morphogenesis. Mech.Dev.
107, 69–82. https://doi.org/10.1016/s0925-4773(01)00452-x.

36. Wu, Z., Zhu, Y., Liu, H., Liu, G., and Li, F. (2020). Wnt10b promotes hair follicles growth and
dermal papilla cells proliferation via Wnt/β-Catenin signaling pathway in Rex rabbits. Biosci Rep
40, BSR20191248. https://doi.org/10.1042/BSR20191248.

37. Sennett, R., Wang, Z., Rezza, A., Grisanti, L., Roitershtein, N., Sicchio, C., Mok, K.W., Heitman,
N.J., Clavel, C., Ma'ayan, A., et al. (2015). An Integrated Transcriptome Atlas of Embryonic Hair
Follicle Progenitors, Their Niche, and the Developing Skin. Dev Cell *34*, 577–591.
https://doi.org/10.1016/j.devcel.2015.06.023.

38. Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L., et al.
(2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. The Innovation
2, 100141. https://doi.org/10.1016/j.xinn.2021.100141.

39. Biggs, L.C., Mäkelä, O.J., Myllymäki, S.-M., Das Roy, R., Närhi, K., Pispa, J., Mustonen, T., and
Mikkola, M.L. (2018). Hair follicle dermal condensation forms via Fgf20 primed cell cycle exit,
cell motility, and aggregation. eLife *7*, e36468. https://doi.org/10.7554/eLife.36468.

40. Kubilus, J.K., and Linsenmayer, T.F. (2010). Developmental guidance of embryonic corneal
innervation: roles of Semaphorin3A and Slit2. Dev Biol *344*, 172–184.
https://doi.org/10.1016/j.ydbio.2010.04.032.

41. Ma, L., and Tessier-Lavigne, M. (2007). Dual Branch-Promoting and Branch-Repelling Actions of
Slit/Robo Signaling on Peripheral and Central Branches of Developing Sensory Axons. J.
Neurosci. *27*, 6843–6851. https://doi.org/10.1523/JNEUROSCI.1479-07.2007.

42. Taniguchi, M., Yuasa, S., Fujisawa, H., Naruse, I., Saga, S., Mishina, M., and Yagi, T. (1997).
Disruption of Semaphorin III/D Gene Causes Severe Abnormality in Peripheral Nerve Projection.
Neuron *19*, 519–530. https://doi.org/10.1016/S0896-6273(00)80368-2.

43. Schwarz, Q., Vieira, J.M., Howard, B., Eickholt, B.J., and Ruhrberg, C. (2008). Neuropilin 1 and 2
control cranial gangliogenesis and axon guidance through neural crest cells. Development *135*,
1605–1613. https://doi.org/10.1242/dev.015412.

44. Schwarz, Q., Waimey, K.E., Golding, M., Takamatsu, H., Kumanogoh, A., Fujisawa, H., Cheng,
H.-J., and Ruhrberg, C. (2008). Plexin A3 and plexin A4 convey semaphorin signals during facial
nerve development. Dev Biol *324*, 1–9. https://doi.org/10.1016/j.ydbio.2008.08.020.

45. Ulupinar, E., Datwani, A., Behar, O., Fujisawa, H., and Erzurumlu, R. (1999). Role of semaphorin
III in the developing rodent trigeminal system. Mol Cell Neurosci *13*, 281–292.
https://doi.org/10.1006/mcne.1999.0747.

46. Cheng, H.J., Bagri, A., Yaron, A., Stein, E., Pleasure, S.J., and Tessier-Lavigne, M. (2001). Plexin-
A3 mediates semaphorin signaling and regulates the development of hippocampal axonal
projections. Neuron *32*, 249–263. https://doi.org/10.1016/s0896-6273(01)00478-0.

47. North, H.A., Karim, A., Jacquin, M.F., and Donoghue, M.J. (2010). EphA4 is necessary for
spatially selective peripheral somatosensory topography. Dev Dyn *239*, 630–638.
https://doi.org/10.1002/dvdy.22185.

48. Dent, E.W., Barnes, A.M., Tang, F., and Kalil, K. (2004). Netrin-1 and semaphorin 3A promote or
inhibit cortical axon branching, respectively, by reorganization of the cytoskeleton. J Neurosci *24*,
3002–3012. https://doi.org/10.1523/JNEUROSCI.4963-03.2004.

49. Moe, K., Sijaona, A., Shrestha, A., Kettunen, P., Taniguchi, M., and Luukko, K. (2012).
Semaphorin 3A controls timing and patterning of the dental pulp innervation. Differentiation *84*,
371–379. https://doi.org/10.1016/j.diff.2012.09.003.

50. Machon, O., Masek, J., Machonova, O., Krauss, S., and Kozmik, Z. (2015). Meis2 is essential for
cranial and cardiac neural crest development. BMC.Dev.Biol. *15*, 40.
https://doi.org/10.1186/s12861-015-0093-6.

51. Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., Zheng, S., Butler, A., Lee, M.J., Wilk, A.J.,
Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal single-cell data. Cell *184*,
3573-3587.e29. https://doi.org/10.1016/j.cell.2021.04.048.

52. Zappia, L., and Oshlack, A. (2018). Clustering trees: a visualization for evaluating clusterings at
multiple resolutions. GigaScience *7*. https://doi.org/10.1093/gigascience/giy083.

Figure 1

A Control   Meis2 cKO

E15.5

B E13.5

Sox9

C E13.5

DAPI Meis2

D Wnt1-Cre2;
mTmG

E12.5

E E13.5

F Control   Meis2 cKO

E13.5   Sox9   TrkA

TrkA Sox9

Figure 2

Figure 3

Figure 4

Figure 5

A

B Sox2   Sox18   Tbx18
   Bmp4   Lef1   Cdkn1a

C

D Control   Meis2 cKO
   E12.5 2h
   DAPI / EdU
   E13.5 18h

E EdU/ DAPI (%)   **
   Control   Meis2 cKO

F Cell Adhesion
   Chd2, Cdh11, Vcan, Ncam1, Actg1, Fn1, Tnc, Alcam
   Control E12.5   Meis2 cKO E12.5   Control E13.5   Meis2 cKO E13.5

G ECM
   Bgn, Cd34, Col1a1, Col1a2, Col3a1, Dcn, Lama4, P4hb, Timp2, Vcam1
   Control E12.5   Meis2 cKO E12.5   Control E13.5   Meis2 cKO E13.5

Figure 6

Figure 7