Mesenchymal Meis2 controls whisker development independently from trigeminal sensory innervation

Mehmet Mahsum Kaplan [1], Erika Hudacova [1,2], Miroslav Matejcek [1], Haneen Tuaima [4], Jan Krivanek [4] and Ondrej Machon [1,3*]

* Correspondence: ondrej.machon@iem.cas.cz

[1] Department of Developmental Biology, Institute of Experimental Medicine, Czech Academy of Sciences, Prague, Czech Republic

[2] Department of Cell Biology, Faculty of Science, Charles University, Prague, Czech Republic

[3] Laboratory of Transcriptional Regulation, Institute of Molecular Genetics, Czech Academy of Sciences, Prague, Czech Republic

[4] Department of Histology and Embryology, Faculty of Medicine, Masaryk University, Brno, Czech Republic

Abstract

Hair follicle development is initiated by reciprocal molecular interactions between the placode-forming epithelium and the underlying mesenchyme. Cell fate transformation in dermal fibroblasts generates a cell niche for placode induction by activation of signaling pathways WNT, EDA, and FGF in the epithelium. These successive paracrine epithelial signals initiate dermal condensation in the underlying mesenchyme. Although epithelial signaling from the placode to mesenchyme is better described, little is known about primary mesenchymal signals resulting in placode induction. Here we show that *Meis2* expression in cells derived from the neural crest is critical for whisker formation, and also for branching of trigeminal nerves. While whisker formation is independent of the trigeminal sensory innervation, MEIS2 in mesenchymal dermal cells orchestrates the initial steps of epithelial placode formation and subsequent dermal condensation. MEIS2 regulates the expression of transcription factor *Foxd1*, which is typical of pre-dermal condensation. However, deletion of *Foxd1* does not affect whisker development. Overall, our data suggest an early role of mesenchymal MEIS2 during whisker formation and provide evidence that whiskers can normally develop in the absence of sensory innervation or *Foxd1* expression.

Keywords

whisker follicle; hair follicle placode; cranial nerves; Meis2; Foxd1, Sox2

Introduction

The local cell environment provides essential cues that determine cell fate decisions, proliferation, and migration. The molecular communication between the developing epithelium and the underlying cell environment is crucial for the formation of multiple structures, which are derived from the embryonic epithelium, such as the eye lens, hair follicles including whisker (vibrissae) follicles, teeth, taste papillae, the intestine lumen, or salivary glands. In the developing vertebrate head, these tissue cross-talks employ neural crest-derived mesenchyme with the epithelium, myogenic progenitors, or cartilage. Critical steps in whisker follicle (WF) development are very similar to normal hair follicle (HF) development [1], which has been studied much more extensively. HF morphogenesis is initiated by thickening of the epithelium forming a placode (Pc), which subsequently induces condensation of underlying fibroblasts termed dermal condensation (DC) [2]. Beta-catenin-dependent WNT signaling is essential for Pc induction [3], which initiates the EDA pathway [4], FGF20 [5], and SHH signaling [6] in the epithelium. Abrogation of any of these signaling pathways leads to the arrest of HF development. However, the succession of initial inductive steps in HF formation is still unclear. Pc formation requires hitherto unknown primary signals from dermal fibroblasts (Fb) at the pre-DC stage that is characterized by expression of *Foxd1* [7] and *Prdm1* [8]. Recent studies employing single-cell transcriptomics helped elucidate the molecular determination of Pc and DC induction. Mok et al. [7] described a wave of transcription factors that define cell fate trajectories, starting in dermal fibroblasts underlying the epithelium before Pc induction (pre-DC stage), towards DC and the late DC stage when Pc invagination commences. For instance, expression of *Foxd1* transcription factor was detected in pre-DC fibroblasts that are also characterized by expression of *Twist2, Lef1*, and *Smad3* [7,9]. DC cells predominantly express *Sox2*, *Foxd1, Ptch1,* and *Bmp4* [10,11]. On the other hand, DC initiation requires canonical WNT and FGF20 signaling in Pc, suggesting a reciprocal molecular crosstalk between the dermal mesenchyme and overlying epithelium [7,10]. Transplantation experiments suggested that mesenchymal cells modulate WNT signaling in Pc by activation of RSPO1; however, RSPO1 is not sufficient for HF induction even in combination with BMP inhibition [12]. Although these data comprehensively map molecular events during HF initiation, identification of the primary inductive signal for Pc induction coming from dermal fibroblasts is still missing.

Epithelial structures such as whiskers or teeth are formed at precisely defined locations. Whiskers are arranged in an accurate rectangular pattern in five rows, each containing four to nine follicles. Each whisker is innervated by a separate whisker-to-barrel circuit that is reflected in a clear topographic organization of the barrel cortex [13]. Axons of the trigeminal nerve ending in the infraorbital nerve innervate each whisker germ in the maxillary prominence. It has been suggested that early innervation does not play a role in determining WF pattern as distribution of trigeminal axons is widespread and random in the maxillary process, and no one-to-one spatial correlation between nerves and WFs was found [1,14] . This view has been challenged by the observation of a regular pattern of the nerve plexus in the snout half a day prior to the pattern of WFs [15,16]. However, these studies do not imply whether WFs develop normally *in vivo* in the complete absence of innervation. Although normal vessel ring organization in the whisker follicles of mice lacking trigeminal sensory innervation has been reported [17], a comprehensive description of the WF development was not studied in this context. Therefore, while existing literature points to a scenario with an unlikely role of innervation for the WF formation and patterning [18–20], a more direct and conclusive *in vivo* study is needed to resolve this issue.

Similarly, the influence of the trigeminal nerves for tooth development has not been conclusively demonstrated. Organ culture methods showed no involvement of trigeminal (TG) sensory innervation [21], and tooth germs can trigger their own innervation [22]. Along these lines, diastema tooth primordia develop and subsequently disappear with no TG innervation [23]. However, the existence of earlier pioneering axons in the TG system has been reported; therefore, tooth development might be initiated by interaction with the early arriving pioneering axons [24]. Moreover, roles of neurogenic factors have been shown during tooth development [25–27]. A surgical removal of (TG) axons innervating teeth led to tooth degeneration [28,29]. Hence, cranial nerves might deliver morphogenetic signals that orchestrate

tooth maintenance. The role of innervation during tooth development was also supported by Kaukua et
al. [30], who showed that dental mesenchymal stem cells are recruited from the nerve-associated glial
cell. Taken together, it has been suggested that innervation play a role in tooth development [31].
However, in a model system where innervation is completely missing, tooth development has not been
analyzed.
Thus, peripheral nerves and accompanying glia contribute to the mesenchymal tooth germ
niche. Nonetheless, it is still unclear what cellular mechanisms regulate precise positions of whiskers
in the snout. Turing reaction-diffusion mathematical model can be applied for oscillating tissue
patterning that is orchestrated by diffusion of short-range activating morphogen that triggers
expression of long-range inhibitor. Reaction-diffusion-based instruction of the epithelium by
combinatorial WNT, FGF, and BMP signaling is capable of HF placode induction in a highly restricted
regular pattern [32]. However, a highly specific rectangular arrangement and relatively long distance
among WFs within the developing snout may indicate that other molecular determinants play a role in
spatiotemporal control of WF induction. This revives the previous speculations that an axonal network
may be involved in WF positioning. In this report, we assessed the scenario that the neural network is
implicated in the control of accurate geometrical arrangement of WFs.
Here, we generated *Meis2* conditional mutants using the Wnt1-Cre2 driver [33] targeting
exclusively the neural crest-derived tissues, including the dermal mesenchyme and cranial nerves,
without detectable recombination in the overlying epithelium [34]. Although *Meis2* expression in the
epithelium of Wnt1-Cre2; *Meis2* [fl/fl] (*Meis2* cKO) mutants was preserved, whisker development was
arrested at the Pc induction stage. This is documented by failure in induction of expression of placodal
genes such as *Shh*, *Edar* and *Lef1*. We further show that severely affected branching of trigeminal
nerves innervating WFs cannot cause their developmental arrest because WFs form normally even in
a complete absence of trigeminal nerves. Expression of *Meis2* in dermal fibroblasts is therefore
necessary for induction of whisker placodes followed by dermal condensation. Protein expression
analysis suggests that MEIS2 transcription factor is essential for triggering a primary inductive signal
from the mesenchyme to the epithelium to initiate WF placode formation. While single-cell
transcriptomics and immunohistochemical validations showed that *Meis2* operates upstream of *Foxd1*,
primary pre-DC marker, we report a dispensable role of *Foxd1* during WF development. Overall, our
data not only describe an essential role of mesenchymal *Meis2* for WF development but also
document normal WF formation even in the complete absence of innervating axons.
121
Results

***Meis2* expression in cranial neural crest-derived cells is required for correct trigeminal nerve**
projections and whisker development

To study whisker morphogenesis, we used Wnt1-Cre2; *Meis2*[fl/fl] conditional mutants, in which *Meis2* is
deleted in neural crest-derived cells (hereafter, *Meis2* cKO). Micro-computed tomography (micro-CT)
of E15.5 embryos revealed severely compromised development of whiskers (Figure 1A). Missing
whiskers were also observed whole-mount embryonic heads stained with placodal marker SOX9
antibody after light-sheet microscopy scanning at as early as E13.5 (Figure 1B). It is noteworthy that
we observed in some cases a small number of SOX9+ WFs in *Meis2* cKO. They most likely
represented normal follicles that 'escaped' from Wnt1-Cre2-mediated deletion of *Meis2*. The number
of WF 'escapers' in *Meis2* cKO snouts varied among samples. Some mutant embryos completely
missed WFs, but some showed a reduction in their numbers. Overall, quantification of the WF number
documented a reduction to 5.7 ± 2.0% at E12.5 and to 17.1 ± 5.9% at E13.5 in the mutants (n=at least
3 embryos for E12.5, n= at least 8 embryos for E13.5, mean ± sem). Analysis of placode
thickness/follicle length and DC area revealed that these escaper whiskers have normal morphology
(Supp Figure 1A, B). To analyze expression of *Meis2* in the snout, we performed fluorescent
immunohistochemistry. MEIS2 protein was detected in the epithelium, in the most superficial dermis
including dermal condensates (DC). In *Meis2* cKO, mesenchymal expression of *Meis2* virtually

disappeared (Figure 1C, asterisk, Supp Figure 1C) while its epithelial expression was maintained
(Figure 1C, arrowheads, Supp Figure 1C), which corresponds to tissue-specific Cre recombination of
Wnt1-Cre2 [33,34]. It is interesting that MEIS2 signal was also remarkably reduced around sole WF
'escapers' (Figure 1C, arrow) but we cannot exclude that, due to incomplete Cre-mediated deletion, a
minor expression of *Meis2* below the immunohistochemical detectability threshold is sufficient for WF
induction. To verify the tissue-specificity in the snout region, we crossed Wnt1-Cre2 mice with mTmG
mouse strain, where only cells derived from the neural crest express membrane-localized GFP.
Whole-mount immunofluorescence imaging confirmed the Wnt1-Cre2 activity in neural crest
derivatives in the craniofacial area, midbrain, and dorsal spinal cord (Figure 1D). In the snout, Cre
activity was restricted to dermal mesenchyme without a detectable recombination in the overlying
epithelium (Figure 1E). Moreover, branches of trigeminal sensory nerves were labelled with GFP
(Figure 1E, arrows), consistent with the previously known origin of trigeminal neurons mainly from
neural crest cells [35]. Whiskers are innervated by the axons of the infraorbital nerve, a branch of the
trigeminal nerve. To examine the axonal outgrowths associated with whisker germs, we co-labelled the
hair follicle marker SOX9 with the sensory nerve marker TRKA in whole-mount heads at E13.5. Figure
1F and Supp Video 1 and Supp Video 2 show that missing whiskers in mice lacking *Meis2* in neural
crest-derived cells were accompanied by severely affected trigeminal nerve growth and branching.
Because defects in nerve branching were not limited to the infraorbital nerve and were also observed
in other trigeminal branches, this phenotype is most likely not caused by missing WFs themselves.
Moreover, the size of the trigeminal ganglion and the nerve exit from the ganglion appeared grossly
normal (Supp Figure 1D). These observations suggest that defects in the trigeminal nerve morphology
occur during branching and fasciculation of axonal projections through the craniofacial region. Overall,
these results indicate an essential role of *Meis2* not only in the development of the whisker follicles but
also for the branching of trigeminal nerves that innervate them.

Normal sizes of the escaped WFs in *Meis2* cKO mice at E12.5 and E13.5 (Supp Figure 1A, B) suggest
that WF phenotype in the mutants is not due to a delay in their development since in this case WFs
would appear smaller. In order to further exclude this possibility that WF formation is delayed in *Meis2*
cKO mice, we analyzed whisker phenotype at later embryonic stages. Sections of E15.5 micro-CT
images showed aberrant WF phenotype (Supp Figure 2A). Also at E18.5, the pattern of WFs were
missing in the mutant (Supp Figure 2B). Staining of E18.5 snout section with placodal marker EDAR
antibody revealed that, similarly to E13.5, a small number of escaped WFs in *Meis2* cKO with
comparable depths to control WFs (Supp Figure 2C, arrows). However, this aberrant whisker
phenotype persisted until E18.5. These data indicate that the whisker phenotype in the mutants is not
due to a developmental delay but rather suggest a direct involvement of mesenchymal *Meis2* in
mechanisms initiating WF formation. Interestingly, at E18.5 stage, hair follicles became detectable in
the snout with a similar appearance in the mutants, suggesting a specific role of *Meis2* in WFs. This
could be due to an early role of *Meis2* in the mesenchyme because HFs develop later.

Trigeminal innervation is not required for WF development

Aberrant whisker development in *Meis2* cKO could be explained by an indispensable function of *Meis2*
in mesenchymal cells or by insufficient trigeminal sensory innervation. The latter scenario takes into
account the possibility that the nerve or nerve-associated Schwann cells produce factors that are
necessary for WF formation. Thus, compromised branching of sensory axons in the *Meis2* cKO snout
might result in insufficient secretion of crucial WF-inducing factors. In order to directly assess the
function of sensory innervation in WF development, we analyzed *Neurog1*[-/-] mice, which do not
develop some cranial nerves, including trigeminal nerves, and therefore lack sensory innervation of
the developing whiskers [36]. Immunostaining of E13.5 100-μm thick snout cryosections for placodal
marker SOX9, axonal marker TUJ1, and sensory nerve marker TRKA antibodies revealed that
whiskers developed normally in *Neurog1*[-/-] mutants even in the complete absence of innervating

sensory axons (Figure 2A). Lack of any TUJ1-labelled axon fascicules in the proximity of the normally
developed WFs in the *Neurog1*[-/-] snout further excludes the possibility of non-sensory neurons
triggering WF development in the absence of sensory innervation. In order to assess the role of
sensory innervation for overall whisker patterning, we stained E12.5 and E13.5 snout whole mounts
with SOX9 and EDAR antibodies, marking initial stages of WF development [4], i.e. placode formation.
These results revealed essentially identical whisker patterning between control and *Neurog1*[-/-] mice
(Figure 2B). EDAR staining performed at E12.5 10-µm FFPE sections further confirmed normal
placodal expression of EDAR, indicating normal initiation of the WF development (Figure 2C). WF
formation was also validated by micro-CT at E17.5, showing normal progression of WF development
at later stages (Figure 2D, top) even when trigeminal ganglia were absent (Figure 2D, bottom, arrows).
Overall, these data unambiguously show that whisker formation does not require trigeminal sensory
innervation and exclude the possibility that impaired nerves or nerve-associated cells in *Meis2* cKO
resulted in impaired WF induction. Therefore, it can be confidently concluded that the dermal
mesenchyme expressing *Meis2* is evidently critical for WF development.

In addition to whisker phenotype, we analyzed tooth morphology in *Neurog1*[-/-] mice since the role of
innervation for tooth development represents a long-standing controversy and has not been directly
tested previously[21,23,24,31,37,38]. Interestingly, we did not detect any defects in tooth development (Supp
Figure 3A), which was corroborated also by micro-CT images from E17.5 mice (Supp Figure 3B),
reporting an innervation-independent formation of teeth in *Neurog1*[-/-] mice.

207 Mesenchymal *Meis2* operates upstream of epithelial EDAR signaling during initial steps of WF
208 formation

Our data strongly suggest that *Meis2* expression in the mesenchyme is required for induction of
placodal expression of *Sox9* in WF. Since a number of regulators of HF development were identified to
function upstream of placodal *Sox9* expression, we tested these successive molecular events leading
to WF induction. SHH release in basal placodal cells was shown to trigger *Sox9* expression in
suprabasal placodal cells. This is achieved by asymmetric cell division resulting in suprabasal
daughter cells with low WNT activity so that SHH can drive *Sox9* expression [39]. We performed RNA in
situ hybridization to evaluate *Shh* expression in *Meis2* cKO mice and control littermates at E13.5.
These data revealed a remarkably decreased number of *Shh*-positive foci in *Meis2* cKO mice (Figure
3A, arrow), showing that placodal expression of *Shh* is regulated by mesenchymal MEIS2 protein.
Again, we observed a small number of *Shh*-positive WF escapers (arrow). EDA/NFκB signaling in
placodes has been shown to act upstream of Shh signaling before Pc invagination [4]. We therefore
stained whole mount E12.5 snouts from *Meis2* cKO mice and controls with SOX9 and EDAR
antibodies. As seen in Figure 3B, EDAR protein was localized to epithelial WF placodes and
overlapped with SOX9 staining in controls. In contrast, in *Meis2* cKO mutants, both EDAR and SOX9
were hardly detectable, documenting that *Meis2* mesenchymal function is upstream of EDA signaling
in Pc. Whole-mount EDAR staining was confirmed on tissue sections which detected significantly
reduced number of EDAR-positive foci in the *Meis2* cKO epithelium. Moreover, EDAR-negative
regions within the mutant epithelium did not show any signs of morphological placode formation as
inspected by DAPI staining (Figure 3C, Supp Figure 4A). Overall, these data demonstrate that
mesenchymal *Meis2* operates at very early stages of WF formation in which it participates in forming a
primary inductive mesenchymal signal for EDAR+ placode formation. This early arrest in mutants
affects all subsequent steps of WF development including *Shh* and *Sox9* expression, and Pc and DC
formation.

232 *Meis2* deletion in NCC-derived cells affects epithelial but not dermal WNT signaling

It has been shown that WNT/β-catenin signaling acts upstream of EDA/NFκB signaling as the earliest
molecular pathway identified during HF formation [4]. Epithelial WNT activity is required for activation of
dermal canonical WNT signaling, while dermal WNT signaling reciprocally affects patterned WNT
activity in the epithelium [40]. We therefore assessed canonical WNT signaling in *Meis2* cKO snouts by
several types of WNT signaling readouts. Firstly, we used LEF1 immunofluorescence because its
expression has been widely used as proxy for WNT signaling in HF [41,42] and *Lef1* is also required for
whisker formation [43]. Matos et al. [42] showed that dermal cells expressed *Lef1* broadly whereas in the
epithelium, *Lef1* expression was concentrated in Pc cells compared to inter follicular regions. During
Pc downgrowth in HF, basal Pc cells displayed LEF1 signal similarly to Shh and EDAR, in contrast to
suprabasal placodal cells lacking LEF1. This is consistent with the view that high WNT activity is seen
in the basal cells whereas suprabasal cells are WNT low [42]. In WF, we similarly detected robust LEF1
signal in Pc that was remarkably lower in interfollicular regions (Figure 4A, Supp Figure 3B, C).
However, in distal snout regions with no WFs, the epithelium showed confluent LEF1 signal (Supp
Figure 3C) suggesting that LEF1 signal is regionally dependent Pc indicator. In contrast to the
epithelium, DC underneath invaginating placodes consistently displayed decreased LEF1 staining
compared to peri-DC neighboring regions (Figure 4A, Supp Figure 4B). In *Meis2* cKO, we rarely
detected a patterned placodal increase in LEF1 staining in the expected sites of WF positions (Figure
4A, Supp Figure 4C). Quantification of LEF1-stained E12.5 10μm FFPE sections revealed a significant
decrease in LEF1+ placodal sites per section from 2.89 ± 0.39 in control to 1 ± 0.82 *Meis2* cKO (mean
± sem, p = 0.0002, two-tailed t-test, n = at least 3 mice, 18 sections). This shows that failure of WF
initiation in *Meis2* cKO is accompanied by missing expression of *Lef1* in Pc, indicating that canonical
WNT signaling is not induced in the Pc epithelium. However, average LEF1 intensity in the escaped
placodes of the mutants did not significantly differ from that of control (95.08 ± 6.47 %, p = 0.51, two-
tailed t-test). On the other hand, LEF1 staining in *Meis2* cKO dermal fibroblasts was confluent and
without down-regulation in presumptive DC sites, which again confirms early developmental arrest in
mutant WFs. Analysis of LEF1 staining intensity in both DC (93.60 ± 6.77 %, p = 0.305, two-tailed t-
test, n = 3 mice and 43 DCs for control and 17 DCs for mutants) and non-DC upper dermis (89.91 ±
13.19 %, p = 0.117, two-tailed t-test, n = 3 mice and 37 regions for control and 33 regions for mutants)
again did not reveal any significant difference between control and mutant snouts. These data indicate
that *Meis2* in the mesenchyme does not affect dermal WNT signaling but is required for induction of
epithelial WNT activity during placode formation.

Secondly, in order to verify WNT activity dynamics in developing WF, we employed WNT reporter
mouse strain BAT-gal [44] in which multiple TCF-binding sites promote expression of *beta-galactosidase*
gene. We crossed this reporter mouse strain to *Meis2* cKO and stained snout frontal sections with an
anti-galactosidase antibody. In contrast to LEF1 staining, BAT-gal reporter staining did not show
consistent upregulation and concentration of WNT activity in Pc and no corresponding downregulation
in DC in controls. It rather exhibited confluent expression in the dermal mesenchyme (Figure 4B) that
was not changed in mutants. Thus, BAT-gal reporter therefore did not serve as a reliable WNT readout
in developing WF in our hands, probably due to the high stability of galactosidase protein compared to
RNA [45], which limited the analysis of the dynamic nature of WNT activity during WF development.

Thirdly, we tested the expression of selected WNT ligands in *Meis2* cKO snouts. *Wnt10b* expression
is strongly activated in HF placodes [46] and it also is important for HF progression [47]. Thus, *WNT10b*
seems the most likely *Wnt* gene activating canonical pathway in HF which is routinely monitored by
Axin2 expression in many tissue contexts including HF [10,39]. We performed HCR FISH to assess
expression of *Wnt10b* and WNT target gene *Axin2* in snouts at E12.5. Similarly to LEF1 staining, we
observed *Axin2* increase in wild-type Pc, while DC and dermal mesenchyme displayed uniform *Axin2*
expression, which did not accord with decreased LEF1 in DC (Figure 4C, Supp Figure 4B, D). This
might reflect differential responses of *Lef1* and *Axin2* expression to WNT activity in DC. In *Meis2* cKO,

upregulation of *Axin2* in the epithelium was not detected which most likely reflected missing placodes.
On the other hand, mutant dermal mesenchyme displayed broad *Axin2* expression which was similar
to wild-type controls. *Wnt10b* transcripts were detected in normal Pc in controls while this expression
disappeared in missing Pc in mutants (Figure 4C). Overall, these data suggest that mesenchymal
Meis2 does not affect WNT signaling in the dermis, whereas it is critical for the patterned WNT
upregulation in the epithelium. Since WNT activation in the epithelium is one of the earliest steps of
HF and WF development, this leads to loss of EDAR and *Shh* markers in *Meis2* cKO WF, and it
confirms a very early role of mesenchymal *Meis2* during whisker development.

289 *Meis2* regulates dermal cell proliferation and the transition of fibroblast cell fate to DC

We have recently reported single-cell RNA sequencing (scRNA-seq) datasets for mesenchymal cells
derived from the cranial neural crest of E12.5 and E13.5 *Meis2* cKO embryos and their control
littermates [48]. Cell cluster analysis identified one cluster representing dermal fibroblasts. We subset
and further re-clustered using the standard Seurat workflow (in Methods). We identified 6 clusters,
where Cluster 2 was defined as DC by cell markers *Sox2*, *Sox18*, *Tbx18*, *Bmp4*, *Lef1*, and *Cdkn1a*
(Figure 5A-B, Supp Figure 5A, Supplementary Data 1). The other clusters were determined to
represent fibroblasts annotated by their markers in the heatmap and module scores generated for
fibroblast (Fb), pre-DC, DC1, and DC2 markers
(https://github.com/kaplanmm/whisker_scRNA/tree/main.) [7,49]. (Supp Figure 5, Supplementary Data 1).
Cluster 2 displayed low expression of Fb module score genes and high expression of DC2 module
score genes, confirming again that cluster 2 represents developed DCs. Transcriptomic analysis
showed that the cell number proportion in cluster 2 was substantially lower in *Meis2* cKO mice
compared to control littermates (Figure 5C), reflecting the WF phenotype in the mutant. Interestingly,
clusters 0 also contained a lower cell proportion in the *Meis2* cKO dataset. On the other hand, the cell
number proportion in cluster 1 increased in *Meis2* cKO (Figure 5C). However, the cell identity of
cluster 1 was unclear because it was marked by only 27 genes (Supplementary Data 1). Cluster 0
revealed GO terms related to cell cycle and chromosome segregation/organization according to GO
analysis by clusterProfiler R package [50] using top 100 markers (Supplementary Data 2). Therefore, we
conclude that cluster 0 very likely represents dividing cells. Lower cell number proportion in cluster 0
in *Meis2* cKO suggested a *Meis2*-dependent regulation of dermal cell proliferation. This hypothesis
was tested by performing EdU chase experiments where EdU was injected at E12.5 and embryos
were harvested after 2 or 18 hours. Indeed, the 2D area coverage of DAPI+ region by EdU+ region in
the upper dermis of *Meis2* cKO snout decreased from 52.96 ± 2.05% to 42.38 ± 2.77% (mean ± sem,
t-test p= 0.0057) in 2-hour pulse and from 52.17 ± 0.81% to 45.76 ± 1.85% (mean ± sem, t-test p=
0.0052) in 18-hour pulse (Figure 5D-E). These EdU experiments confirmed scRNA-seq data and
indicated a *Meis2* function in proliferation of upper dermal cells which might contribute to the aberrant
whisker phenotype of the mutant.

Because cell migration is a critical process for the migration of Fb cells into DC during cell fate
acquisition [51], we plotted cell adhesion and ECM module scores, both of which were found to be
increased in mutants (Figure 5F, G). Thus, such an increase in cell interactions might limit Fb
migration efficiency by decreasing conductivity of the environment for cell migration and ultimately to
decreased numbers of WFs in mutants. Similar results were found by Hudacova et al.[48] in craniofacial
mesenchymal cells in *Meis2* cKO mice. These increases in cell adhesion and ECM gene expression
might also contribute to decreased TG axonal growth and branching described above (Figure 1, Supp
Video 1 and Supp Video 2).

325 *Meis2* controls expression of *Foxd1* in the dermis

Next, we performed differential gene expression analysis between the DC cluster of *Meis2* cKO and
controls at E12.5 and E13.5. In this scRNA-seq dataset, *Foxd1* was found within the top genes
downregulated in *Meis2* cKO mice (avg_log2FC = 1.0585, p_val = 4.55E-06) (Figure 6A,
Supplementary Data 3). Interestingly, *Foxd1* expression was described as a hallmark of the earliest
molecular changes in dermal fibroblasts prior to DC formation and defined the pre-DC stage in HF [7].
Further, *Sox2*, a well-known DC marker, was also strongly downregulated in *Meis2* cKO (Figure 6A).
To validate transcriptomic data, we analyzed FOXD1 protein expression in tissue sections of wild-type
snouts at E12.5 and E13.5. Similarly to HF, *Foxd1* expression preceded *Sox2* also in WF which
occurred before the Pc downgrowth into the dermal niche at E12.5. During initial Pc invagination and
early DC stage at E13.5, *Foxd1* and *Sox2* expression overlapped (Figure 6B, arrows). In more
progressed WF germs with deeply invaginated Pc, *Foxd1* expression remained in peri-DC regions
rather than in DC that was marked by a strong SOX2 signal (Figure 6B, right panels, arrowhead).
Next, whole-mount staining of FOXD1 in the whole snout highlighted dynamic WF development along
the proximo-distal axis. It started in proximal columns at E12.5, in which WF formation was more
progressed, while distal snout regions initiate WF slightly later (Figure 6C). By E13.5, 3-4 proximal
columns displayed discernible WFs (Figure 6D). Strikingly, *Meis2* cKO snouts displayed no FOXD1+
WFs at E12.5 (Figure 6C). At E13.5, we detected around 4 proximal WFs in mutants while controls
contained around 17 WFs on each snout side (Figure 6D). FOXD1+ WFs in *Meis2* cKO most likely
represented 'escaper' WFs that were described above. Whole-mount FOXD1 immunostaining was
further confirmed on tissue sections (Figure 6C-D, lower panels). Normal WF at E13.5 expressed
Foxd1 in DC or peri-DC and *Sox9* in Pc. In contrast, this progression of WF formation was halted in
Meis2 cKO (Figure 6C, D). These results suggest that MEIS2 is the key component of transcription
controlling machinery in the dermal mesenchymal that initiates WF.

Foxd1 is dispensable for WF development

Because *Foxd1* expression in DC is strictly regionally specific and its expression is lost in *Meis2* cKO,
we hypothesized that *Foxd1* transcription factor function is essential for WF formation. We therefore
analyzed whiskers in embryos where *Foxd1* expression is ablated. We crossed heterozygous *Foxd1*-
GFP-CreERT2 mice, in which GFP-Cre-ERT2 cassette was inserted into the initiation codon of *Foxd1*
gene. Homozygous embryos are therefore null mutants for *Foxd1* gene. WF development was
assessed in whole-mount-stained embryonic heads at E13.5 using SOX9, FOXD1, and TUJ1
antibodies. Figure 7A shows that SOX9 was detected in invaginating placodes, FOXD1 in surrounding
DC mesenchyme, and TUJ1 in axonal projections innervating each WF. In parallel, we stained EDAR
in WF placodes (Figure 7B). Interestingly, the number and morphology of WF in controls and *Foxd1*
mutants were essentially identical. Thus, FOXD1 protein is a very important marker of the early DC
cell fate in WF but does not have functional relevance for whisker formation despite its dynamic and
regionally specific expression pattern around the developing WFs.

Discussion

We have shown that mesenchymal expression of transcription factor *Meis2* is essential for the initial
steps of mouse whisker development. Mouse embryos, in which *Meis2* was deleted in tissues derived
from neural crest cells by Wnt1-Cre2 driver, lost the majority of whiskers. The aberrant whisker
phenotype in the mutants is accompanied by markedly reduced trigeminal nerve branching in the
snout. This suggests a novel function of *Meis2* in trigeminal axonal growth, branching, and guidance. It
may also implicate that whisker formation, or WF positional determination, may depend on early
sensory innervation. We report here, however, that whisker formation is entirely independent of
sensory innervation. *Neurog1*[-/-] mice completely lack trigeminal nerves, including their ganglions, but

whisker development is not affected. Moreover, analysis of *Neurog1*[-/-] mice at E17.5 reveals that
sensory innervation is not required for the further growth and stabilization of WFs. The effect of
trigeminal innervation deficiency on postnatal WF development could not be tested because *Neurog1*[-/-]
mice are lethal within a day after birth. All these experiments exclude the possibility that the whisker
phenotype in *Meis2* cKO embryos is an indirect consequence of decreased trigeminal nerve
branching. It rather indicates that mesenchymal *Meis2* is a pivotal regulator of WF formation.

Our observations suggest that nerve branching phenotype in the mutant is independent of trigeminal
ganglia development or missing whiskers, as *Meis2* cKO mice showed normal trigeminal ganglia size
and nerve exit from the ganglia as well as defective TG nerve branches that do not innervate
whiskers. Thus, *Meis2* rather influences trigeminal branching and guidance through the snout.
Description of the exact nature of factors controlling trigeminal nerve branching downstream of MEIS2
warrants further investigation. A number of molecular pathways, such as SLIT/ROBO,
SEMA/NRP/PLXN [52–56], EPHRIN/EPH [57] signaling, have been implicated during TG nerve
development, and *Meis2* might act upstream of mesenchymal expression of ligands and/or trigeminal
expression of receptors of these pathways to control TG nerve development. One possible
mechanism might involve SEMA3A, an inhibitory factor for trigeminal nerve growth during tooth
innervation, and *Sema3a* null mice display hyper-branched trigeminal nerves [55,58,59]. Our scRNA seq
datasets indicated an increase of *Sema3a* expression in *Meis2* cKO mesenchymal cell populations
(data not shown), which could explain decreased nerve branching in *Meis2* cKO mice and could be
addressed in future studies.

Essential role of *Meis2* in the dermal mesenchyme during very early steps of WF development is
supported by several key observations. Firstly, although epithelial expression of *Meis2* is preserved in
Meis2 cKO mice, thickening of the epithelium is rarely observed, suggesting that the release of WF
placode-inducing dermal factors is under control of mesenchymal *Meis2*. It is widely accepted that HF
or WF formation is initiated from the dermis by a 'first dermal signal' that has not been identified yet.
Our data show that this process is dependent on *Meis2*. Secondly, the absence of *Meis2* results in the
loss of early placodal markers LEF1 and EDAR and pre-DC marker FOXD1 at expected WF locations.
Absence of patterned upregulation of LEF1 and EDAR was documented in HFs after blocking dermal
WNT signaling [40]. Interestingly, our analysis of WNT signaling readouts did now show remarkable
changes in canonical WNT signaling in *Meis2* cKO dermis, indicating that normal dermal WNT
signaling is not sufficient to trigger patterned LEF1 and EDAR upregulation in the epithelium when
dermal MEIS2 is absent. These epithelial changes probably require additional *Meis2*-dependent
intermediate steps, which might act either in parallel with, or downstream of dermal WNT signaling to
induce placode formation. In the dermis, pre-DC marker FOXD1 expression was shown to be
dependent on *Fgf20* expression in HF Pc and occurs after Pc formation [7]. Our analysis of *Foxd1*
expression in wild-type snout dermis revealed that *Foxd1* expression appears prior to morphologically
distinguishable WF Pc. In *Meis2* cKO, FOXD1 signal in the dermis prior to WF Pc formation was not
detected, which accords with single-cell RNA seq data. This all suggests that *Meis2* operates
upstream of *Foxd1*. However, direct analysis of *Foxd1*-null mice did not reveal any phenotype in WF
development. This could be explained by either no function of *Foxd1* in this process or by a possible
functional compensation by its paralog *Foxd3*. Since *Foxd1* does not affect WF formation and
therefore the absence of *Foxd1* cannot account for lacking WFs in *Meis2* cKO mice, loss of FOXD1 in
these mice evidently is a result of lacking pre-DC and DC structures. Lastly, it has been shown in HF
development that dermal fibroblasts proliferate extensively prior to converting to DC cell fate. DC cells,
on the other hand, exit the cell cycle and become quiescent [7,10]. We generally observed reduced
proliferation in upper dermal cells in *Meis2* cKO mice. Lower cell proliferation could contribute to
inefficient cell fate change of Fb to DC and might also lead to insufficient expression and/or release of
Pc-inducing factor in the dermis of *Meis2* mutants. Overall, these observations strongly suggest that

the absence of *Meis2* in mesenchymal cells affects earliest processes of WF development, even prior
to Pc formation.

It is important to note that we observed small numbers (around 18% at E13.5) of WFs in *Meis2* cKO
expressing normal placodal and DC markers. These rarely formed WF 'escapers' in *Meis2* cKO mice
had normal WF morphology, because their placode thickness, follicle length, and DC size were
comparable between controls and mutants at E12.5-E13.5. WF 'escapers' in mutants were probably
not sensitive to Wnt1-Cre2-mediated deletion of *Meis2*. Although immunostaining of MEIS2 in snout
sections indicated that *Meis2* expression around WF escapers was hardly detectable in mutants, a
slight but undetectable expression of *Meis2* resulting from incomplete gene deletion might account for
normal WF formation. On the other hand, it is also possible that *Meis1* paralogue can partially
substitute for missing *Meis2* in its function in the dermal mesenchyme. Alternatively, *Meis2* function in
WF development might be restricted to a short time window as each WF develops at different time
points along the proximal-distal axis.

Materials and Methods

Mouse strains

Generation of the floxed allele of *Meis2* gene (*Meis2* [fl/fl]) was described by Machon et al. (2015) [60]
Meis2[fl/fl] strain was crossed with reporter mouse lines R26R-*EYFP* [fl/fl] (RRID:IMSR JAX006148)
before crossing with the Wnt1-Cre2 [33]. This reporter enabled monitoring Cre recombination specificity.
Wnt1–Cre2 (RRID:IMSR JAX:022137) was used for specific deletion of the *Meis2* [fl/fl] gene in neural
crest cells. For lineage tracing experiments, we used Wnt1-Cre2 crossed with mTmG (RRID:IMSR
JAX:007676). During embryo harvesting, Cre recombination was checked by monitoring EYFP
fluorescence. In very rare cases (1-2 embryos in 10 litters), Cre activity in the whole body was
observed, indicating germline recombination. Such animals were removed from the analysis, and their
parents were removed from crossing schemes. Neurogenin1 knockout mice were purchased from Jax
Lab (RRID:IMSR_JAX:017306). *Foxd1*-GFP-Cre-ERT2 mice were purchased from Jax Lab
(RRID:IMSR_JAX:012464). The presence of the vaginal plug was regarded as the embryonic day 0.5
(E0.5). Pregnant mice were anesthetized using isoflurane gas inhalation and euthanized by cervical
dislocation. Embryos were dissected and placed in ice-cold PBS. All mice were genotyped as
previously described [60]. *Neurog1* and *Foxd1* mutant mice were genotyped according to the Jax
protocol. All procedures involving experimental animals were approved by the Institutional Committee
for Animal Care and Use (permission #PP-10/2019) with every effort to minimize suffering and the
number of animals used. This work did not include human subjects.

Immunohistochemistry and image processing

All embryos used for staining were harvested at either E12.5 or E13.5 embryonic stages, and heads
were fixed in 4% paraformaldehyde (PFA) overnight at 4^0C. After PFA fixation, samples were prepared
in three ways. (1) For FFPE staining, after dehydration in ethanol and xylene, samples were
embedded in paraffin, and frontal 10-μm sections were prepared. Followed by stepwise rehydration,
antigen retrieval in 0.1 M citrate buffer, pH 6.0, under pressure boiling for 12 min was carried out. (2)
For cryosection staining, after dehydration in 30% saccharose for 24 hours, samples were embedded
in OCT, and 100 μm cryo-sections were obtained. (3) For whole mount staining, whole heads were
directly processed to permeabilization and blocking after PFA fixation.

Tissues were blocked in 5% bovine serum albumin (BSA) in PBS with 0.1% Triton X-100 for FFPE
sections or with 0.5% Triton X-100 for thick cryosections or whole mounts for 1 hour and incubated in
primary antibody solution (1% BSA in PBS and 0.1% Triton X-100 for 10-μm sections or 0.5% Triton
X-100 for 100-μm sections and whole mounts). Primary antibody incubation was overnight for FFPE
sections, two nights for cryosections, and three nights for whole mount staining at 4ºC. Primary
antibodies used: SOX9 (Merck Sigma, AB5535), TRKA (R&D Systems, AF1056), MEIS2 (GeneScript,

custom), TUJ1 (R&D Systems, MAB1195), EDAR (R&D Systems, AF745), Lef1 (Cell Signaling,
C12A5), β-galactosidase (Abcam, ab9361), SOX2 (R&D Systems, MAB2018), FOXD1 (Abcam,
AB129324). Primary antibody solutions were washed out with (PBS, Triton X-100) and incubated in
fluorescent secondary antibody solutions for one hour for FFPE sections, three hours for cryosections
at room temperature or overnight for whole mounts at 4^0C. Secondary antibodies: donkey anti-mouse,
-rabbit, -goat with Alexa488, 568, or 647 fluorophores (ThermoFisher Scientific). Primary and
secondary antibodies were used at 1:500 for thick section and whole mount staining and at 1:1000 for
thin section staining. Immunofluorescent images were scanned by spinning disc confocal microscopy
using an Olympus SpinSR10. Imaging was performed for whole mount stainings after tissue clearing
for one hour at room temperature. Acquired z-stacks were maximum intensity projected using ImageJ.
Figures were assembled using Photoshop.

For cell proliferation assays, EdU was injected intraperitoneally to pregnant females at E12.5 for 2 or
18 hours before harvesting (0.5 ml per mouse, 2.5 mg/ml). EdU Base Click kit was used for EdU
detection according to manufacturer's protocol on FFPE sections.

Micro-CT

483 Embryos were fixed in 4% PFA for 2 days and soaked in Lugol's iodine for several days. Scanning
484 was performed on the instrument Bruker Skyscan 1272 with the resolution 3 μm.
485

In situ hybridization

487 Antisense RNA probe for *Shh* was cloned into pGEM-T-easy vector (Promega) using primers: *Shh*-F
488 TCACAAGTCCTCAGGTTCCG, *Shh*-R GGGCTTCAGCTGGACTTGAC. Antisense mRNA was
489 transcribed with T7 polymerase. Whole-mount *in situ* hybridization was performed using standard
490 protocols [60]. Fluorescent in situ hybridization HCR RNA-FISH (Molecular Instruments) was performed
491 according to manufacturer's protocols.

Single Cell RNA sequencing analysis

scRNA sequencing datasets are retrieved from integrated Seurat objects (GSE262468) [48] and
processed with the standard Seurat workflow [61]. Briefly, dermal fibroblast clusters were subset and
split by condition, which was followed by SCTransform normalization, where mitochondrial, ribosomal,
and cell cycle genes were regressed. Seurat objects were then integrated by using the IntegrateData
function with 3000 integration features. For UMAP as a non-linear dimensional reduction embedding,
30 principal components were used. We used the Clustree package to determine the optimal
resolution, which was set at 0.6 for the FindClusters function according to the developers' instructions
[62]. SCT assay was used to generate feature plots and violin plots. Normalized and Scaled data of the
RNA assay were used to extract cluster markers using the FindAllMarkers function and to generate
heatmaps. The FindMarkers function of the Seurat package was used for differential gene expression
analysis between conditions with default parameters. clusterProfiler (4.6.2) was used for GO analysis
[50], where GO terms were extracted by the enrichGO function. This paper did not produce original
codes. Codes used for analysis and lists of genes used to generate module scores can be accessed
at https://github.com/kaplanmm/whisker_scRNA/tree/main.

Quantification and Statistical Analysis

For placode thickness/WF length and DC size analysis, images of DAPI-stained 10 μm FFPE snout
sections were captured with an Olympus SpinSR10 microscope and a 20X objective. With ImageJ
software, maximum intensity projection was applied to z-stack images. DAPI staining was used to
visually identify placodes and DCs. The distance from the outer layer of epithelium in the placodal
region to the epithelial-mesenchymal boundary was measured. For 2D DC size, DCs were identified
by densely accumulated cells underneath the placodes. The area of the region covered by these cells
was measured using ImageJ.

The number of WFs was manually counted in embryos that were used throughout the study for whole-
mount immunostainings (labeled by SOX9, EDAR, and/or FOXD1 antibodies) or *in situ* hybridization

(*Shh* mRNA). The numbers of WFs at both sides of the snout were pooled. The decrease in the percentage of WFs in *Meis2* cKO mice was calculated in comparison to their control littermates. Values from each experiment were averaged to determine the overall decrease in the percentage of WFs.

For EdU/DAPI area measurements, images of EdU-chased and DAPI-stained 10 μm FFPE snout sections were captured with an Olympus SpinSR10 microscope equipped with a 20X objective. With ImageJ software, maximum intensity projection was applied to z-stack images. The dermal region between the epithelium and around a 100-μm distance into the dermis was used for analysis, while DC domains were excluded. The selected dermal regions were processed separately for thresholding steps to cover EdU+ and DAPI+ regions. Threshold values were determined visually to cover specific EdU and DAPI staining. The same threshold values were used for control and *Meis2* cKO mutants. Total area values for EdU and DAPI were used to calculate 2D EdU coverage of the DAPI regions (EdU/DAPI).

Statistical analyses were performed using GraphPad Prism 10 software. N numbers specified in the respective figure legends for each analysis. All the data are presented with mean ± sem (standard error of the mean). Two-tailed student's t-test was used to explore statistical differences between two experimental groups (Control vs. *Meis2* cKO).

Acknowledgements

This work was supported by the Czech Science Foundation (grants 22-10660S, 23-06160S) and Masaryk University, Faculty of Medicine (MUNI/A/1598/2023). We thank the Microscopy Service Centre of IEM CAS and Light Microscopy Core Facility, IMG CAS, Prague, Czech Republic, supported by MEYS – LM2023050 and RVO – 68378050-KAV-NPUI. This work was also funded by MEYS Czech Republic (NanoEnviCZ, LM2018124) and EU Structural Funds Pro-NanoEnviCz (CZ.02.1.01/0.0/0.0/16_013/0001821).

Author contributions

MMK designed and performed experiments, bioinformatics, and wrote the manuscript. EH performed experiments and bioinformatics. MM performed experiments. HT performed experiments. JK designed experiments. OM designed and performed experiments, wrote the manuscript, and provided funding. The authors declare no competing interests.

Figure legends:

Figure 1. **Severe whisker follicle development in *Meis2* cKO embryos. (A)** Micro-CT images of control and *Meis2* cKO embryos at E15.5 showing aberrant whisker phenotype in *Meis2* cKO mice. **(B)** Light-sheet microscopy of Sox9 whole-mount immunostaining of WF. Arrow indicates example of an "escaper" whisker. **(C)** MEIS2 immunofluorescence on frontal frozen sections of E13.5 snouts showing MEIS2 expression in the dermis, DC, and epithelium including Pc. Arrowheads indicate epithelial expression of MEIS2 in both control and *Meis2* cKO mice. Arrow and asterisk show disappearing MEIS2 expression in the DC and dermis, respectively, in the mutant. Scale bar: 150 μm. **(D)** Wnt1-Cre; *mTmG* embryos showing Cre recombination specificity in the craniofacial area, midbrain, and dorsal spinal cord. Recombined cells are labeled by membrane-localized GFP in green, and non-recombined cells labeled by membrane-localized tdTomato in red. **(E)** Frontal sections of Wnt1-Cre; *mTmG* snouts documenting Cre recombination in the neural crest-derived mesenchyme, cranial nerve projections without recombination in the overlying epithelium. **(F)** Whole-mount staining of *Meis2* cKO heads at E13.5 with SOX9 and TRKA antibodies showing almost absence of WFs in and compromised branching of the trigeminal nerve in mutants. Scale bar: 500 μm.

Figure 2. **Neurog1^{-/-} (KO) embryos lack the trigeminal nerve, but WFs development is normal.**
(A) Triple immunostaining of 100-μm sections shows the absence of trigeminal nerve projections (TUJ1+, TRKA+) and normal WF (SOX9+) in *Neurog1^{-/-}* mice. Arrows in black and white images show examples of invagination of whisker placodes labeled by SOX9 antibody. Scale bar: 500 μm. **(B)** Whole-mount immunostaining of WFs with SOX9 and EDAR antibodies showing normal WF morphology and patterning in mutants at E12.5 (top) and 13.5 (bottom). Scale bars: 300 μm. **(C)** EDAR staining of 10-μm sections at E12.5 showing normal initiation of placode formation in *Neurog1* mutants. Scale bar: 150 μm. **(D)** micro-CT images reveal normally developed whiskers (top) at E17.5 in mutants while TGs are lacking (bottom, arrows).

Figure 3. **Induction and progression of WF development is compromised in *Meis2* cKO. (A)** Whole-mount *in situ* hybridization of *Shh* mRNA documenting loss of WFs in mutants. Arrow shows an escaper whisker. **(B)** Whole-mount immunostaining of WFs with SOX9 and EDAR antibodies showing absence of WFs in *Meis2* cKO at E12.5. Two examples for each genotype are shown. Scale bar: 300 μm. **(C)** EDAR staining of 10-μm sections showing placode formation arrest in mutants. Scale bar: 150 μm.

Figure 4. **WNT signaling in the epithelium is affected by deletion of *Meis2* in the dermal mesenchyme. (A)** LEF1 staining of 10-μm paraffin frontal sections of snouts showing abundance of LEF1 in the dermal mesenchyme, while in the epithelium, it is concentrated in placodes in regions of WF appearance. Similar to missing placodes, LEF1 is also lost in *Meis2* cKO epithelium. Scale bars: 150 μm. **(B)** beta-galactosidase immunostaining of sections from WNT reporter BAT-gal controls and *Meis2* cKO; BAT-gal mutants showing widespread galactosidase signal in the dermis and epithelium. Scale bars: 150 μm. **(C)** *In situ* hybridization HCR-FISH using probes for *Axin2* and *Wnt10b*. In the epithelium, *Wnt10b* is detected in placodes, and the signal is lost in *Meis2* cKO. Similarly, upregulation of *Axin2* is not observed in the expected WF loci. In the dermal mesenchyme, *Axin2* mRNA is detected throughout the dermis regardless of WFs. This pattern was not changed in *Meis2* cKO. Scale bar: 50 μm.

Figure 5. **Dermal condensation and proliferation are reduced in *Meis2* cKO. (A)** Uniform Manifold Approximation and Projection (UMAP) diagram of single-cell RNA-seq analysis of the dermal mesenchyme subset with 6-cluster resolution in which cluster 2 (green) represents dermal condensate (DC) of WFs. **(B)** UMAP representations of typical DC markers *Sox2*, *Sox18*, *Tbx18*, *Bmp4, Lef1*, and *Cdkn1* shown by FeaturePlots. **(C)** Relative cell numbers in 6 clusters showing a higher cell count in cluster 1 (nonspecialized cells) at the expense of clusters 0 (dividing cells) and 2 (DC). **(D)** Analysis of cell proliferation by EdU incorporation after a 2-hour pulse at E12.5 and an 18-hour pulse at E13.5. **(E)** Overall proliferation rate was reduced in *Meis2* cKO from 52.96 ± 2.05% to 42.38 ± 2.77% (mean ± sem, t-test p= 0.0057) in 2-hour pulse (top) and from 52.17 ± 0.81% to 45.76 ± 1.85% (mean ± sem, t-test p= 0.0052) in 18-hour pulse (bottom). Each data point indicates the average value for one section. n = 2 mice and at least 9 sections. **(F)** Increase of cell adhesion module score in mutants at E13.5 by scRNA-seq analysis. Module scores are generated by indicated genes. **(G)** Increase of extracellular matrix (ECM) module score in mutants at E13.5 by scRNA-seq analysis. Module scores are generated by indicated genes.

Figure 6. **Expression of the earliest pre-DC and DC marker *Foxd1* is lost in *Meis2* cKO. (A)** UMAP representations of decline in *Foxd1* (top) and *Sox2* (bottom) expression in DC cluster (#2) shown by FeaturePlots. Expression of both genes is significantly reduced in mutants. **(B)** Immunofluorescence of FOXD1 and SOX2 at initial stages of WF showing FOXD1 in the pre-DC stage while the expression shifts from DC to peri-DC regions during WF progression. SOX2 appears as a typical DC marker with no earlier expression. Scale bar: 50 μm. **(C-D)** Whole-mount immunostaining of FOXD1 and Sox9 of heads from controls and *Meis2* cKO at E12.5 **(C)** and E13.5 **(D).** It shows the absence of WFs including the pre-DC marker FOXD1 at the E12.5 stage when proximal columns of

WFs have already formed. A few WF escapers develop in mutants at E13.5 in proximal regions while
at least 6 WF columns have formed in controls. Scale bars: 400 μm (C) and 500 μm (D). Lower
panels: FOXD1 staining in FFPE sections. Scale bars: 150 μm.

Figure 7. **WFs normally develop in loss-of-function *Foxd1* mutants. (A)** Whole-mount
immunostaining of FOXD1, TUJ1, and SOX9 of heads from controls and *Foxd1*-null mutants at E13.5
showing normal formation of WFs in mutants in which FOXD1 signal disappears. Normal WF
development is also reflected in normal WF innervation represented by TUJ1 staining. Scale bars: 500
μm. **(B)** Whole-mount immunostaining of EDAR confirmed normal Pc appearance in *Foxd1*-null
mutants. Scale bars: 500 μm.

Supplementary Figure 1. **Normal sizes of escaper WFs and TGs in *Meis2* cKO mice. (A-B)**
Quantification of 2D DC area (**A**) and placode (Pc) thickness of non-invaginated or length of the
invaginated follicles (**B**) showing comparable values between control and *Meis2* cKO at E12.5 (left)
and E13.5 (right). N = 4 embryos for E12.5 and 3 embryos for E13.5, and at least 30 WFs for controls
and 10 for *Meis2* cKO mice. P-values for all graphs are larger than 0.05, student t-test. Data are
presented as mean ± sem. **(C)** 3 examples of 100-μm frozen frontal sections through snout stained
with MEIS2 antibody showing disappearing MEIS2 staining in the *Meis2* cKO dermis. Scale bar: 300
μm. **(D)** 100-μm frozen sections through TG ganglia stained with TUJ1 and TRKA antibodies showing
non-defective TG ganglion size or exit of the TG nerve from the ganglia. Scale bar: 250 μm.

Supplementary Figure 2. **Whisker phenotype persists at E18.5 *Meis2* cKO mice. (A)** Micro-CT
images through the snout of E15.5 control and *Meis2* cKO mice showing abnormal whisker phenotype
in the mutant. Arrows show some escaper WFs. **(B)** Brightfield images of control and *Meis2* cKO at
E18.5. **(C)** 2 examples of 100-μm frozen sections of snouts from E18.5 control and *Meis2* cKO stained
with DAPI (blue) and EDAR (red) showing only a few whiskers in the mutant. In contrast, EDAR-
labeled hair follicles appeared to be normal in the mutant. Arrows show some escaper WFs. Scale
bar: 1 mm.

Supplementary Figure 3. **Normal tooth development in *Neurog1*^-/- mice. (A)** 100-μm frozen sections
of molar teeth of control and *Neurog1*^-/- mice at E13.5 stained with SOX9, TUJ1, and TRKA antibodies
showing the normal phenotype of the mutant. Scale bar: 150 μm. **(B)** Micro-CT images around the
molar tooth level at E17.5 in *Neurog1*^-/- mice.

Supplementary Figure 4. **EDAR and LEF1 expression in *Meis2* cKO mice. (A)** 3 examples of 10-μm
FPPE sections stained with EDAR antibody showing affected expression of EDAR in *Meis2* cKO mice.
Arrows indicate EDAR-positive sites which are observed less frequently in the mutant snout. Scale
bar: 500 μm. **(B)** 2 representative images of normally developed escaper WFs in the mutants by LEF1
staining. LEF1 expression is similar in just forming placodes (top) and invaginated (bottom).
Additionally, LEF1 expression shows a decline in the DC regions compared to peri-DC. Scale bar: 30
μm. **(C)** 3 representative micrographs of LEF1-stained 10-μm FFPE snout sections showing normal
expression of LEF1 in the dermis but the decreased number of LEF1+ placodes in the epithelium of
the mutant snout. Scale bar: 500 μm. **(D)** *In situ* hybridization HCR-FISH using probes for *Axin2* and
Wnt10b in control snout at E12.5 showing placodal expression of *Wnt10b* (green) and *Axin2* (arrows,
red) as well as widespread expression of *Axin2* in the dermis. Scale bar: 60 μm.

Supplementary Figure 5. **Identification of DC cluster in scRNA sequencing datasets. (A)**
Heatmap of top 10 markers for each identified cluster in scRNA-seq data. Each column represents
individual cells, whereas each row represents indicated marker genes on the left. Genes in the black
box are top markers of cluster 2 (DC). **(B)** FeaturePlots and VlnPlots of fibroblasts (Fb), pre-DC, DC1,
and DC2 module scores in each cluster. See Methods for selected genes used for generating module
scores.

References

1. Wrenn, J.T., and Wessells, N.K. (1984). The early development of mystacial vibrissae in the mouse. J.Embryol.Exp.Morphol. *83*, 137–156.

2. Sennett, R., and Rendl, M. (2012). Mesenchymal-epithelial interactions during hair follicle morphogenesis and cycling. Semin Cell Dev Biol *23*, 917–927. https://doi.org/10.1016/j.semcdb.2012.08.011.

3. Huelsken, J., Vogel, R., Erdmann, B., Cotsarelis, G., and Birchmeier, W. (2001). beta-Catenin controls hair follicle morphogenesis and stem cell differentiation in the skin. Cell *105*, 533–545. https://doi.org/10.1016/s0092-8674(01)00336-1.

4. Zhang, Y., Tomann, P., Andl, T., Gallant, N.M., Huelsken, J., Jerchow, B., Birchmeier, W., Paus, R., Piccolo, S., Mikkola, M.L., et al. (2009). Reciprocal Requirements for EDA/EDAR/NF-κB and Wnt/β-Catenin Signaling Pathways in Hair Follicle Induction. Developmental Cell *17*, 49–61. https://doi.org/10.1016/j.devcel.2009.05.011.

5. Huh, S.-H., Närhi, K., Lindfors, P.H., Häärä, O., Yang, L., Ornitz, D.M., and Mikkola, M.L. (2013). Fgf20 governs formation of primary and secondary dermal condensations in developing hair follicles. Genes Dev *27*, 450–458. https://doi.org/10.1101/gad.198945.112.

6. St-Jacques, B., Dassule, H.R., Karavanova, I., Botchkarev, V.A., Li, J., Danielian, P.S., McMahon, J.A., Lewis, P.M., Paus, R., and McMahon, A.P. (1998). Sonic hedgehog signaling is essential for hair development. Current Biology *8*, 1058–1069. https://doi.org/10.1016/S0960-9822(98)70443-9.

7. Mok, K.-W., Saxena, N., Heitman, N., Grisanti, L., Srivastava, D., Muraro, M.J., Jacob, T., Sennett, R., Wang, Z., Su, Y., et al. (2019). Dermal Condensate Niche Fate Specification Occurs Prior to Formation and Is Placode Progenitor Dependent. Dev Cell *48*, 32-48.e5. https://doi.org/10.1016/j.devcel.2018.11.034.

8. Manti, P.G., Darbellay, F., Leleu, M., Coughlan, A.Y., Moret, B., Cuennet, J., Droux, F., Stoudmann, M., Mancini, G.-F., Hautier, A., et al. (2022). The Transcriptional Regulator Prdm1 Is Essential for the Early Development of the Sensory Whisker Follicle and Is Linked to the Beta-Catenin First Dermal Signal. Biomedicines *10*, 2647. https://doi.org/10.3390/biomedicines10102647.

9. Sennett, R., Wang, Z., Rezza, A., Grisanti, L., Roitershtein, N., Sicchio, C., Mok, K.W., Heitman, N.J., Clavel, C., Ma'ayan, A., et al. (2015). An Integrated Transcriptome Atlas of Embryonic Hair Follicle Progenitors, Their Niche, and the Developing Skin. Dev Cell *34*, 577–591. https://doi.org/10.1016/j.devcel.2015.06.023.

10. Gupta, K., Levinsohn, J., Linderman, G., Chen, D., Sun, T.Y., Dong, D., Taketo, M.M., Bosenberg, M., Kluger, Y., Choate, K., et al. (2019). Single-Cell Analysis Reveals a Hair Follicle Dermal Niche Molecular Differentiation Trajectory that Begins Prior to Morphogenesis. Developmental Cell *48*, 17-31.e6. https://doi.org/10.1016/j.devcel.2018.11.032.

11. Ge, W., Tan, S.-J., Wang, S.-H., Li, L., Sun, X.-F., Shen, W., and Wang, X. (2020). Single-cell
Transcriptome Profiling reveals Dermal and Epithelial cell fate decisions during Embryonic Hair
Follicle Development. Theranostics *10*, 7581–7598. https://doi.org/10.7150/thno.44306.

12. Mäkelä, O.J.M., and Mikkola, M.L. (2023). Mesenchyme governs hair follicle induction.
Development *150*, dev202140. https://doi.org/10.1242/dev.202140.

13. Yang, J.W., Kilb, W., Kirischuk, S., Unichenko, P., Stuttgen, M.C., and Luhmann, H.J. (2018).
Development of the whisker-to-barrel cortex system. Curr.Opin.Neurobiol. *53*, 29–34.
https://doi.org/10.1016/j.conb.2018.04.023.

14. Davies, A.M., and Lumsden, A.G. (1986). Fasciculation in the early mouse trigeminal nerve is not
ordered in relation to the emerging pattern of whisker follicles. J Comp Neurol *253*, 13–24.
https://doi.org/10.1002/cne.902530103.

15. Van Exan, R.J., and Hardy, M.H. (1980). A spatial relationship between innervation and the early
differentiation of vibrissa follicles in the embryonic mouse. J Anat *131*, 643–656.

16. Maklad, A., Conway, M., Hodges, C., and Hansen, L.A. (2010). Development of Innervation to
Maxillary Whiskers in Mice. The Anatomical Record *293*, 1553–1567.
https://doi.org/10.1002/ar.21194.

17. Oh, W.-J., and Gu, C. (2013). Establishment of Neurovascular Congruency in the Mouse Whisker
System by an Independent Patterning Mechanism. Neuron *80*, 458–469.
https://doi.org/10.1016/j.neuron.2013.09.005.

18. Andrés, F.L., and Van der Loos, H. (1982). Whisker patterns form in cultured non-innervated
muzzle skin from mouse embryos. Neurosci Lett *30*, 37–41. https://doi.org/10.1016/0304-
3940(82)90008-8.

19. Andrés, F.L., and Van der Loos, H. (1983). Cultured embryonic non-innervated mouse muzzle is
capable of generating a whisker pattern. International Journal of Developmental Neuroscience *1*,
319–338. https://doi.org/10.1016/0736-5748(83)90034-5.

20. Lillesaar, C., and Hildebrand, C. (1999). Denervation does not affect the growth of rat vibrissae.
Neuroscience Letters *261*, 69–72. https://doi.org/10.1016/S0304-3940(98)01015-5.

21. Lumsden, A.G., and Buchanan, J.A. (1986). An experimental study of timing and topography of
early tooth development in the mouse embryo with an analysis of the role of innervation. Arch Oral
Biol *31*, 301–311. https://doi.org/10.1016/0003-9969(86)90044-0.

22. Erdélyi, G., Fried, K., and Hildebrand, C. (1987). Nerve growth to tooth buds after homotopic or
heterotopic autotransplantation. Developmental Brain Research *33*, 39–47.
https://doi.org/10.1016/0165-3806(87)90174-X.

23. Løes, S., Kettunen, P., Kvinnsland, I., and Luukko, K. (2002). Mouse rudimentary diastema tooth
primordia are devoid of peripheral nerve fibers. Anat Embryol *205*, 187–191.
https://doi.org/10.1007/s00429-002-0247-8.

24. Stainier, D.Y., and Gilbert, W. (1990). Pioneer neurons in the mouse trigeminal sensory system.
Proc Natl Acad Sci U S A *87*, 923–927. https://doi.org/10.1073/pnas.87.3.923.

25. Sandor, B., Fintor, K., Felszeghy, Sz., Juhasz, T., Reglodi, D., Mark, L., Kiss, P., Jungling, A.,
Fulop, B.D., Nagy, A.D., et al. (2014). Structural and Morphometric Comparison of the Molar
Teeth in Pre-eruptive Developmental Stage of PACAP-Deficient and Wild-Type Mice. J Mol
Neurosci *54*, 331–341. https://doi.org/10.1007/s12031-014-0392-6.

26. Tachibana, R., Tatehara, S., Kumasaka, S., Tokuyama, R., and Satomura, K. (2014). Effect of
Melatonin on Human Dental Papilla Cells. International Journal of Molecular Sciences *15*, 17304–
17317. https://doi.org/10.3390/ijms151017304.

27. Moiseiwitsch, J.R.D., and Lauder, J.M. (1996). Stimulation of murine tooth development in
organotypic culture by the neurotransmitter serotonin. Archives of Oral Biology *41*, 161–165.
https://doi.org/10.1016/0003-9969(95)00117-4.

28. Zhao, H., Feng, J., Seidel, K., Shi, S., Klein, O., Sharpe, P., and Chai, Y. (2014). Secretion of shh
by a neurovascular bundle niche supports mesenchymal stem cell homeostasis in the adult mouse
incisor. Cell Stem Cell *14*, 160–173. https://doi.org/10.1016/j.stem.2013.12.013.

29. Adameyko, I., and Fried, K. (2016). The Nervous System Orchestrates and Integrates Craniofacial
Development: A Review. Front Physiol *7*, 49. https://doi.org/10.3389/fphys.2016.00049.

30. Kaukua, N., Shahidi, M.K., Konstantinidou, C., Dyachuk, V., Kaucka, M., Furlan, A., An, Z.,
Wang, L., Hultman, I., Ahrlund-Richter, L., et al. (2014). Glial origin of mesenchymal stem cells in
a tooth model system. Nature *513*, 551–554. https://doi.org/10.1038/nature13536.

31. Duan, Y., Liang, Y., Yang, F., and Ma, Y. (2022). Neural Regulations in Tooth Development and
Tooth-Periodontium Complex Homeostasis: A Literature Review. Int J Mol Sci *23*, 14150.
https://doi.org/10.3390/ijms232214150.

32. Glover, J.D., Wells, K.L., Matthäus, F., Painter, K.J., Ho, W., Riddell, J., Johansson, J.A., Ford,
M.J., Jahoda, C.A.B., Klika, V., et al. (2017). Hierarchical patterning modes orchestrate hair
follicle morphogenesis. PLoS Biol *15*, e2002117. https://doi.org/10.1371/journal.pbio.2002117.

33. Lewis, A.E., Vasudevan, H.N., O'Neill, A.K., Soriano, P., and Bush, J.O. (2013). The widely used
Wnt1-Cre transgene causes developmental phenotypes by ectopic activation of Wnt signaling.
Dev.Biol. *379*, 229–234. https://doi.org/10.1016/j.ydbio.2013.04.026.

34. Fabik, J., Kovacova, K., Kozmik, Z., and Machon, O. (2020). Neural crest cells require Meis2 for
patterning the mandibular arch via the Sonic hedgehog pathway. Biol.Open. *9*.
https://doi.org/10.1242/bio.052043.

35. Leonard, C.E., Quiros, J., Lefcort, F., and Taneyhill, L.A. (2022). Loss of Elp1 disrupts trigeminal
ganglion neurodevelopment in a model of familial dysautonomia. Elife. *11*.
https://doi.org/10.7554/eLife.71455.

36. Ma, Q., Chen, Z., del, B.B., I, de la Pompa, J.L., and Anderson, D.J. (1998). neurogenin1 is
essential for the determination of neuronal precursors for proximal cranial sensory ganglia. Neuron
20, 469–482. https://doi.org/10.1016/s0896-6273(00)80988-5.

37. Maeda, Y., Miwa, Y., and Sato, I. (2019). Distribution of the neuropeptide calcitonin gene-related
peptide-α of tooth germ during formation of the mouse mandible. Ann Anat *221*, 38–47.
https://doi.org/10.1016/j.aanat.2018.09.001.

38. Weil, M., Itin, A., and Keshet, E. (1995). A role for mesenchyme-derived tachykinins in tooth and
mammary gland morphogenesis. Development *121*, 2419–2428.
https://doi.org/10.1242/dev.121.8.2419.

39. Ouspenskaia, T., Matos, I., Mertz, A.F., Fiore, V.F., and Fuchs, E. (2016). WNT-SHH Antagonism
Specifies and Expands Stem Cells prior to Niche Formation. Cell *164*, 156–169.
https://doi.org/10.1016/j.cell.2015.11.058.

40. Chen, D., Jarrell, A., Guo, C., Lang, R., and Atit, R. (2012). Dermal β-catenin activity in response
to epidermal Wnt ligands is required for fibroblast proliferation and hair follicle initiation.
Development *139*, 1522–1533. https://doi.org/10.1242/dev.076463.

41. DasGupta, R., and Fuchs, E. (1999). Multiple roles for activated LEF/TCF transcription complexes
during hair follicle development and differentiation. Development *126*, 4557–4568.
https://doi.org/10.1242/dev.126.20.4557.

42. Matos, I., Asare, A., Levorse, J., Ouspenskaia, T., de la Cruz-Racelis, J., Schuhmacher, L.-N., and
Fuchs, E. Progenitors oppositely polarize WNT activators and inhibitors to orchestrate tissue
development. eLife *9*, e54304. https://doi.org/10.7554/eLife.54304.

43. Kratochwil, K., Dull, M., Farinas, I., Galceran, J., and Grosschedl, R. (1996). Lef1 expression is
activated by BMP-4 and regulates inductive tissue interactions in tooth and hair development.
Genes Dev *10*, 1382–1394. https://doi.org/10.1101/gad.10.11.1382.

44. Maretto, S., Cordenonsi, M., Dupont, S., Braghetta, P., Broccoli, V., Hassan, A.B., Volpin, D.,
Bressan, G.M., and Piccolo, S. (2003). Mapping Wnt/beta-catenin signaling during mouse
development and in colorectal tumors. Proc.Natl.Acad.Sci.U.S.A *100*, 3299–3304.
https://doi.org/10.1073/pnas.0434590100.

45. Machon, O., Backman, M., Machonova, O., Kozmik, Z., Vacik, T., Andersen, L., and Krauss, S.
(2007). A dynamic gradient of Wnt signaling controls initiation of neurogenesis in the mammalian
cortex and cellular specification in the hippocampus. Dev.Biol. *311*, 223–237.
https://doi.org/10.1016/j.ydbio.2007.08.038.

46. Reddy, S., Andl, T., Bagasra, A., Lu, M.M., Epstein, D.J., Morrisey, E.E., and Millar, S.E. (2001).
Characterization of Wnt gene expression in developing and postnatal hair follicles and
identification of Wnt5a as a target of Sonic hedgehog in hair follicle morphogenesis. Mech.Dev.
107, 69–82. https://doi.org/10.1016/s0925-4773(01)00452-x.

47. Wu, Z., Zhu, Y., Liu, H., Liu, G., and Li, F. (2020). Wnt10b promotes hair follicles growth and
dermal papilla cells proliferation via Wnt/β-Catenin signaling pathway in Rex rabbits. Biosci Rep
40, BSR20191248. https://doi.org/10.1042/BSR20191248.

48. Hudacova, E., Abaffy, P., Kaplan, M.M., Krausova, M., Kubista, M., and Machon, O. (2025).
Single-cell transcriptomic resolution of osteogenesis during craniofacial morphogenesis. Bone *190*,
117297. https://doi.org/10.1016/j.bone.2024.117297.

49. Sennett, R., Wang, Z., Rezza, A., Grisanti, L., Roitershtein, N., Sicchio, C., Mok, K.W., Heitman,
N.J., Clavel, C., Ma'ayan, A., et al. (2015). An Integrated Transcriptome Atlas of Embryonic Hair
Follicle Progenitors, Their Niche, and the Developing Skin. Dev Cell *34*, 577–591.
https://doi.org/10.1016/j.devcel.2015.06.023.

50. Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L., et al.
(2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. The Innovation
2, 100141. https://doi.org/10.1016/j.xinn.2021.100141.

51. Biggs, L.C., Mäkelä, O.J., Myllymäki, S.-M., Das Roy, R., Närhi, K., Pispa, J., Mustonen, T., and
Mikkola, M.L. (2018). Hair follicle dermal condensation forms via Fgf20 primed cell cycle exit,
cell motility, and aggregation. eLife *7*, e36468. https://doi.org/10.7554/eLife.36468.

52. Taniguchi, M., Yuasa, S., Fujisawa, H., Naruse, I., Saga, S., Mishina, M., and Yagi, T. (1997).
Disruption of Semaphorin III/D Gene Causes Severe Abnormality in Peripheral Nerve Projection.
Neuron *19*, 519–530. https://doi.org/10.1016/S0896-6273(00)80368-2.

53. Schwarz, Q., Vieira, J.M., Howard, B., Eickholt, B.J., and Ruhrberg, C. (2008). Neuropilin 1 and 2
control cranial gangliogenesis and axon guidance through neural crest cells. Development *135*,
1605–1613. https://doi.org/10.1242/dev.015412.

54. Schwarz, Q., Waimey, K.E., Golding, M., Takamatsu, H., Kumanogoh, A., Fujisawa, H., Cheng,
H.-J., and Ruhrberg, C. (2008). Plexin A3 and plexin A4 convey semaphorin signals during facial
nerve development. Dev Biol *324*, 1–9. https://doi.org/10.1016/j.ydbio.2008.08.020.

55. Ulupinar, E., Datwani, A., Behar, O., Fujisawa, H., and Erzurumlu, R. (1999). Role of semaphorin
III in the developing rodent trigeminal system. Mol Cell Neurosci *13*, 281–292.
https://doi.org/10.1006/mcne.1999.0747.

56. Cheng, H.J., Bagri, A., Yaron, A., Stein, E., Pleasure, S.J., and Tessier-Lavigne, M. (2001). Plexin-
A3 mediates semaphorin signaling and regulates the development of hippocampal axonal
projections. Neuron *32*, 249–263. https://doi.org/10.1016/s0896-6273(01)00478-0.

57. North, H.A., Karim, A., Jacquin, M.F., and Donoghue, M.J. (2010). EphA4 is necessary for
spatially selective peripheral somatosensory topography. Dev Dyn *239*, 630–638.
https://doi.org/10.1002/dvdy.22185.

58. Dent, E.W., Barnes, A.M., Tang, F., and Kalil, K. (2004). Netrin-1 and semaphorin 3A promote or
inhibit cortical axon branching, respectively, by reorganization of the cytoskeleton. J Neurosci *24*,
3002–3012. https://doi.org/10.1523/JNEUROSCI.4963-03.2004.

59. Moe, K., Sijaona, A., Shrestha, A., Kettunen, P., Taniguchi, M., and Luukko, K. (2012).
Semaphorin 3A controls timing and patterning of the dental pulp innervation. Differentiation *84*,
371–379. https://doi.org/10.1016/j.diff.2012.09.003.

60. Machon, O., Masek, J., Machonova, O., Krauss, S., and Kozmik, Z. (2015). Meis2 is essential for
cranial and cardiac neural crest development. BMC.Dev.Biol. *15*, 40.
https://doi.org/10.1186/s12861-015-0093-6.

61. Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., Zheng, S., Butler, A., Lee, M.J., Wilk, A.J.,
Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal single-cell data. Cell *184*,
3573-3587.e29. https://doi.org/10.1016/j.cell.2021.04.048.

62. Zappia, L., and Oshlack, A. (2018). Clustering trees: a visualization for evaluating clusterings at
multiple resolutions. GigaScience *7*. https://doi.org/10.1093/gigascience/giy083.

A E15.5 Control *Meis2* cKO

B E13.5 SOX9

C E13.5 DAPI MEIS2

D Wnt1-Cre2; mTmG E12.5

E E13.5

F Control *Meis2* cKO

E13.5 SOX9 TRKA

TRKA SOX9