Low concentrations of the food contaminant Deoxynivalenol trigger apoptosis and impair GnRH-induced LH secretion in pituitary gonadotrope-like cells

Guodong Cai[1,3†], Lingchen Yang[2,3†], Francis Marien-Bourgeois[3], Derek Boerboom[3], Gustavo Zamberlam[3] and Imourana Alassane-Kpembi[3*]

[1]College of Veterinary Medicine, Yangzhou University, Yangzhou 225009, Jiangsu, China.

[2]College of Veterinary Medicine, Hunan Agricultural University, Changsha 410128, Hunan, China.

[3] Centre de Recherche en Reproduction et Fertilité (CRRF), Département de Biomédecine vétérinaire, Faculté de Médecine vétérinaire, Université de Montréal, Saint-Hyacinthe H3T 1J4, Québec, Canada

* Corresponding author. E-mail: Imourana.alassane-kpembi@umontreal.ca

Contributing authors: 008400@yzu.edu.cn; lcyang@hunau.edu.cn

[†] These authors contributed equally to this work.

Abstract

The *Fusarium* mycotoxin deoxynivalenol (DON) is one of the most frequently occurring food contaminants. Nearly all individuals are exposed to DON, due to it widespread presence in grains and grain-based products. Chronic DON poisoning is associated with growth retardation, immunotoxicity as well as impaired reproduction and fetal development. At the molecular level, DON alters intracellular signaling by activating mitogen-activated protein kinases (MAPKs) that modulate cell growth, differentiation, and apoptosis. Of note, these MAPKs are also critical mediators of gonadotrophin-releasing hormone (GnRH)-induced synthesis and secretion of follicle-stimulating hormone (FSH) and luteinizing hormone (LH) by pituitary gonadotrope cells. So far, no research has explored the potential endocrine-disrupting effects of DON on pituitary gonadotropins production. Herein, we show the first evidence that DON affects LH production by the immortalized gonadotrope-like cell line LβT2 in a concentration-dependent manner. Taken together, our experiments demonstrated that low concentrations of DON affect GnRH-induced signaling through a mechanism that, at least in

part, involves apoptosis and inhibition of GnRH-induced phosphorylation of ERK-MAPK. Consequently, DON also affects the GnRH-induced expression of *Cga* and *Lhb*, two critical genes for LH synthesis and secretion by gonadotrope cells in mammals. This research broadens our knowledge of the toxicity of DON and brings a new depth to the potential neuroendocrine implications for reproduction.

Keywords: Mycotoxins; deoxynivalenol; pituitary gland; LβT2 cells; MAPK signaling pathway; gonadotropins, LH

1. Introduction

Mycotoxins, as pervasive contaminants of foodstuffs, pose a substantial threat to both food security and human health. Among them, Deoxynivalenol (DON), produced by *Fusarium* fungi species, is one of the most prevalent, and represents a global health concern (Streit et al. 2012). Nearly all individuals worldwide are exposed to DON, and the current exposure levels necessitate taking vigilant steps to guarantee food safety (Mishra et al. 2020). Acute exposure to DON in experimental animal models cause symptoms such as diarrhea, vomiting, leukocytosis, hemorrhage, and even death, while chronic low-dose exposure is associated with anorexia, growth retardation, immunotoxicity and altered neuroendocrine signaling (Pestka 2010).

A negative impact of DON on reproductive functions primarily related to altered gonadal function has also been reported in several species (Pizzo et al. 2015; Sprando et al. 1999; Tassis et al. 2022). Interestingly, male rats exposed to DON presented altered serum levels of the gonadotropins follicle-stimulating hormone (FSH) and luteinizing hormone (LH), suggesting that DON could act directly at the level of the pituitary gland (Sprando et al. 2005).

At the cellular level, the primary mechanism of DON toxicity involves inhibiting protein and nucleic acid synthesis by binding to the ribosome and activating cellular kinases, including the mitogen-activated protein kinases (MAPKs) p38, c-Jun N-terminal kinase, and extracellular-signal-regulated kinase (ERK) (Payros et al. 2016). This mechanism influences various signal transduction pathways, including those related to cell proliferation, stress, inflammation, differentiation and apoptosis (Murphy and Blenis 2006). Most importantly, MAPKs are also critical mediators of GnRH-induced gonadotropin synthesis and secretion by pituitary gonadotrope cells (Kahnamouyi et al. 2018). Gonadotropins play a pivotal role in both male and female reproduction. In males, LH regulates the synthesis of androgens by the Leydig cells, whereas FSH promotes Sertoli cell function and thereby influences spermatogenesis (Ulloa-Aguirre and Lira-Albarrán 2016). In females, FSH is the key hormone for follicle growth, selection of the dominant follicle and oocyte maturation, while a midcycle surge of LH is required for oocyte meiotic maturation, successful ovulation, and luteinization of the follicle (Singh et al. 2023). LH and FSH are dimeric glycoproteins composed of a common alpha subunit (CGα) and distinct β subunits (LHβ and FSHβ), which confer biological specificity (Bernard and Tran 2013). The main determinant of gonadotropin synthesis is the level of expression of *LHB* and

FSHB. The transcription of *LHB* is regulated mainly by hypothalamic gonadotropin-releasing hormone

(GnRH) (Tremblay and Drouin 1999), which binds to its receptor (GnRHr) to activate the extracellular

regulated kinases 1 and 2 (ERK1/ 2) via phosphorylation (Bliss et al. 2009; Liu et al. 2002). Yet, to the

best of our knowledge, no research has explored the potential endocrine-disrupting effects of DON on

pituitary gonadotropin production.

To start addressing this gap, our study investigated the effects of DON on pituitary gonadotrope

cell viability, and its impact on GnRH-stimulated ERK signaling and LH secretion. As gonadotropes

are a rare cell population in the pituitary gland (making their enrichment and primary culture

considerable technical challenges), we employed the immortalized gonadotrope-like cell line LβT2.

This cell line is derived from a mouse pituitary tumor, expresses GnRHr, CGα, LHβ and FSHβ and

secretes LH and FSH upon stimulation (Graham et al. 1999; Turgeon et al. 1996). Therefore, this cell

line serves as a crucial tool for the study of gonadotrope physiology, as well as for the identification of

potential endocrine disruptors of gonadotropin production in mammals.

2. Materials and methods

2.1 Chemicals and Reagents

DON and dimethyl sulfoxide (DMSO) were obtained from Millipore Sigma Canada Ltd. (Ontario,

Canada). GnRH was from Peptide Pros (Miami, USA). Dulbecco's Modified Eagle's Medium

(DMEM), Fetal Bovine Serum (FBS), gentamycin and Trypsin were from Wisent (St-Bruno, Quebec,

Canada).

2.2 Cell culture

After thawing, LβT2 cells were seeded in flasks and cultured in DMEM medium (Wisent, St-

Bruno, Quebec, Canada) supplemented with 10% FBS and 4 µg/mL gentamycin and incubated at 37□

with 5% CO_2. The culture medium was refreshed every two days after cell adhesion to the flask was

achieved. When the cells reached 80% confluence, they were trypsinized and seeded in a specific plate

at a specific density, according to the needs of each experiment, as described below.

2.3 Cytotoxicity assessment

LβT2 cells were distributed in 96-well plates at 5×10^3 cells in 100 μL culture medium per well and

allowed to attach. The following day, the culture medium was aspirated, and the cells were exposed to

100 μL of new medium with graded concentrations of DON: 0 nM (control), 1 nM, 10 nM, 10^2 nM, 10^3

nM and 10^4 nM for 24 or 48h. Cell viability was then determined by CellTiter Blue (Promega, Madison,

WI) as previously described by Cai et *al*. (2023).

2.4 Real-time apoptosis and necrosis assay

The Real Time-Glo™ Annexin V Apoptosis and Necrosis Assay (Promega, Madison, WI, Cat#:

JA1011) was used to simultaneously analyze apoptosis and necrosis. This assay probes the presence of

phosphatidylserine on the outer leaflet of the cell membrane during apoptosis, while producing a

fluorescent signal upon the loss of membrane integrity because of the binding of the cell-impermeant

pro-fluorescent DNA dye it contains.

For this experiment, LβT2 cells were seeded at 3×10^4 cells per well in 100 μL culture medium in

white clear-bottom 96-well plates (Costar, Cambridge, MA, USA) and incubated in an atmosphere of 5%

CO2 at 37 °C. After 48 h, 50 μL of the existing cell medium was carefully withdrawn and replaced by

an equivalent volume of fresh medium without (control group) or with 40 nM DON. In each well, 100

μL of a 2X reagent solution was dispensed, according to the manufacturer's instructions, resulting in a

final liquid volume of 200 μL per well. This ensured a final mycotoxin concentration of 10 nM in the

treated well. Subsequently, the plates were placed in a 37°C incubator with 5% CO2. Measurements

were taken at 2 h, 6 h and 24 h post-treatment, to assess membrane integrity through fluorescence

(excitation/emission wavelength: 485 ± 10 nm / 525 ± 10 nm) and phosphatidylserine translocation

through the luminescent signal generated by the Annexin V-dependent assembly of two luciferase

fragments.

2.5 LβT2 cell treatments for MAPK activation

LβT2 cells were distributed in 24-well plates at 1.5×10^5 cells in 300 μL of medium per well and

cultured for 24 hours. Then the cells were either directly stimulated with graded concentrations of

GnRH from 0.001nM to 0.1nM for 15 min; or pre-treated with DON at concentrations of 1 μM or 10

μM for 15 min before stimulation with the highest concentration of GnRH (0.1nM) for 15 min.

Following GnRH challenge, cells were collected, and cellular proteins were extracted using a

radioimmunoprecipitation assay (RIPA) buffer. The protein concentrations of the samples were

quantified using a bicinchoninic acid (BCA) protein quantification kit from Thermo Fisher Scientific

(Waltham, Massachusetts, USA). Twenty µg of protein from each sample were separated using a

Precast Midi Protein Gel (BIO-RAD, Hercules, California, USA) and then transferred to

polyvinylidene fluoride (PVDF) membranes (BIO-RAD, Hercules, California, USA). These were

probed using primary antibodies specific to total ERK and p38, phosphorylated ERK and p38, and β-

tubulin. Detection was done using a peroxidase-conjugated secondary antibody (Jackson

ImmunoResearch Inc, USA), and the antigen-antibody complexes were visualized using an enhanced

chemiluminescence assay (ECL, Invitrogen, Carlsbad, California, USA).

2.5 LβT2 cell treatments for gene expression and hormonal assay

To determine effects of DON on the expression pattern of gonadotropin synthesis-related genes

and LH secretion, LβT2 cells were seeded and allowed to attach in 6-well plates at 1×10^6 cells per well.

On the next day, the cell supernatant was carefully aspirated and replaced with new culture medium

with graded concentrations of DON ranging from 1 nM to 10 µM for 24 h. Following this pre-

treatment with DON, the cells were stimulated with 1nM GnRH for 6 h. While in one experiment cells

were harvested for total RNA extraction, in another experiment medium was collected for LH

measurement.

2.5.1 Total RNA extraction and real-time PCR

The expression of the gene encoding the GnRH receptor (*Gnrhr*) and the gonadotropin-related

genes *Cga*, *Lhb*, and *Fshb* was analyzed. For this, total RNA was first extracted using Trizol as per the

guidelines provided by the manufacturer (BIO-RAD). Following purity and concentration assessment

on a Nanodrop spectrophotometer, total RNA (1µg) was reverse transcribed into cDNA, using the

Advanced cDNA synthesis Kit (Wisent Bioproducts, St-Bruno, QC, Canada) according to the

manufacturer's instructions. RT-qPCR was run on a CFX96 Touch Real-Time PCR Detection System

(BIO-RAD) using qPCR Mastermix with Supergreen (Wisent Bioproducts, St-Bruno, QC, Canada)

following the manufacturer's protocol. To ensure specificity of PCR products, melting curves were

analyzed and the amplicons were sequenced to verify authenticity. Experimental samples were run in

duplicate and data were expressed relative to *Rpl19* as housekeeping gene, and normalized to a

calibrator sample using the Pfaffl method with correction for amplification efficiency (Pfaffl 2001).

Primer details for both housekeeping and target genes are provided in Table 1.

2.5.2 LH measurement

LH levels in cell culture medium samples were evaluated using the Ultra-Sensitive Mouse and Rat

LH ELISA performed by the Center for Research in Reproduction at the Ligand Assay and Analysis

Core Laboratory of the University of Virginia. The capture monoclonal antibody (anti-bovine LH beta

subunit, 518B7; RRID:AB_2784498) was provided by Janet Roser, University of California. The

detection polyclonal antibody (rabbit LH antiserum, AFP240580Rb; RRID:AB_2784499) was

provided by the National Hormone and Peptide Program (NHPP). HRP-conjugated polyclonal antibody

(goat anti-rabbit) was purchased from DakoCytomation (Glostrup, Denmark; D048701-2;

RRID:AB_2784500). Mouse LH reference prep (AFP5306A; NHPP) was used as the assay standard.

The limit of quantitation was 0.016 ng/ml.

2.6 Statistical analysis

All statistical analyses were performed with Prism software 8.0.2 for Windows (GraphPad). The

reported values are the means ± standard error of the mean (SEM) of at least three independent

experiments, each with triplicate wells per experimental condition. Data were transformed to

logarithms if they were not normally distributed (Shapiro–Wilk test). Two-tailed *t* tests or one-way

ANOVAs followed by Tukey's multiple-comparison tests were run for comparisons between the

experimental conditions. Differences with a *p* value < 0.05 were considered statistically significant.

3   Results

3.1 Low concentrations of DON alter gonadotrope-like cell viability

The cytotoxicity of DON for LβT2 cells was assessed using the CellTiter-Blue® Cell Viability

Assay. The results indicated decreased viability of LβT2 cells following exposure to increased graded

concentrations of DON (Fig. 1). Interestingly, it was observed that viability was significantly different

from control after 24 h (*p*<0.05) or 48 h (*p*<0.01) of exposure even the lowest tested concentration of 1

nM. The decline in cell viability showed a clear concentration-dependent pattern at both 24 and 48 h,

reaching its nadir at 10^4 nM DON for both exposure durations (*p*<0.01). Furthermore, the results also

indicated that with longer exposure time (48h *vs.* 24h), the impact on LβT2 cell viability was more

profound. This shows a concentration and time-dependent reduction in gonadotrope-like cell viability

in response to DON starting from the nanomolar range.

3.2 DON induces apoptosis in gonadotrope-like cells

To further elucidate the mechanism of DON-induced cytotoxicity on LβT2 gonadotrope-like cells,

we then conducted a real-time assay with the objective of determining the temporal dynamics of

translocation of phosphatidylserine and loss of membrane integrity following exposure to this

mycotoxin. Translocation of phosphatidylserine to the outer leaflet of the plasma membrane was

revealed through annexin V binding coupled with a luminescent signal emission (Fig. 2a), while loss of

membrane integrity was associated with a fluorescent signal resulting from DNA staining by a cell-

impermeant reagent (Fig. 2b). Apoptotic cell death is therefore characterized by a temporal lag

between phosphatidylserine translocation and the compromise of membrane integrity, whereas necrotic

cell death is characterized by simultaneous annexin V binding and the disruption of membrane integrity.

For this experiment, we chose 10 nM of DON in the treated group. According to the previous

experiment, at this dose, DON already alters cells viability at both 24 and 48h post-exposure. At the 6-h

exposure time point, it was already possible to observe a significant increased phosphatidylserine

translocation compared to the control ($p<0.01$), while no significant difference in membrane integrity

loss was observed ($p>0.05$). However, both phosphatidylserine translocation and membrane integrity

compromise showed statistically significant differences between the DON-treated cells and the control

at the 24-h exposure. These findings indicate that exposure to 10 nM DON for 6 h induced apoptosis

without triggering necrosis.

3.3 DON attenuates GnRH-induced phosphorylation of Erk but not p38 in gonadotrope-like cells

To start exploring the potential negative impact of DON on pituitary gonadotropin production-

related signaling, we investigated the effects of DON exposure on the activation of MAPKs by GnRH

in gonadotrope-like cells. For this, LβT2 cells were challenged with distinct doses of GnRH ranging

from 0.001nM to 0.1nM for 15 min. However, only GnRH at the dose of 0.1 nM significantly increased

the phosphorylation levels for Erk (Fig. 3; $p<0.01$) and p38 (Fig. 4; $p<0.01$) as expected. Most

interestingly, when LβT2 cells were pre-exposed to DON at concentrations of 1 μM and 10 μM before

challenging with 0.1 nM GnRH, only the phospho-Erk (Fig. 3; $p<0.05$) but not the phospho-p38 (Fig. 3; $p<0.01$) increase was attenuated. Together, these results indicate that DON interferes with GnRH-stimulated ERK signaling.

3.4 DON attenuates the GnRH-induced expression of Cga and Lhb but not Gnrhr in gonadotrope-like cells

To better elucidate the impact of DON on GnRH-induced downstream signaling in gonadotrope-like cells, we also investigated the effects of DON on GnRH-induced expression of some of the genes considered essential for the synthesis of gonadotropins. For this, we first performed a preliminary experiment using the same doses of GnRH employed in the previous experiment. However, we did not observe a GnRH-induced increase of *Gnrhr, Cga* or *Lhb* mRNA levels at any dose (0.001 nM to 0.1 nM, data not shown). When we increased the GnRH dose to 1nM, the expression of these three genes was significantly augmented (Fig.5; $p<0.05$). Interestingly, while the GnRH-induced expression of *Cga* and *Lhb* was significantly attenuated by DON pre-treatment ($p<0.05$), *Gnrhr* was not ($p>0.05$). GnRH did not increase *Fshb* expression, which is a typical outcome when only a single pulse of GnRH is used (Lawson et al. 2007). We nonetheless were able to determine that DON does not affect the basal expression of either *Fshb* ($p>0.05$) nor of that of the other gonadotropin-synthesis related genes ($p>0.05$).

3.5 LH secretion by gonadotrope-like cells is dose-dependently inhibited by DON

Together, all the previous experiments demonstrated that DON affects GnRH-induced signaling in gonadotrope-like cells trough a mechanism that, at least in part, involve apoptosis and attenuation of GnRH-induced Erk phosphorylation. Consequently, DON also affects the GnRH-induced expression of *Cga* and *Lhb*, two critical genes for LH synthesis and secretion by gonadotrope cells. To ultimately determine if DON would indeed affect LH secretion by gonadotropes, we then performed an experiment in which LβT2 cells were pre-exposed to graded concentrations of DON (ranging from 1 to 10^4 nM) and then challenged with 1 nM GnRH (Fig. 6). As expected, GnRH at this concentration significantly induced LH release into the culture medium in comparison to control ($p<0.05$). Interestingly, at 1nM DON was not able to attenuate GnRH-induced LH production ($p>0.05$), but at 10^3 nM and 10^4 nM, DON inhibited it successfully ($p<0.05$). Together, these results clearly indicate the

DON exerts a concentration-dependent negative effect on GnRH-induced LH secretion.

Discussion

The present study provides strong evidence for the potential endocrine-disrupting effects of DON

exposure on pituitary gonadotropin production in mammals. We demonstrate herein for the first time

that exposure to realistic concentrations (i.e. low concentrations observed in the diet) of the type B

trichothecene mycotoxin DON can not only be cytotoxic to gonadotrope-like cells, but can impair

GnRH-induced LH secretion in this cell type in a concentration-dependent manner.

A growing body of literature is pointing to the impact of *Fusarium* mycotoxins on the

hypothalamic–pituitary–gonadal (HPG) axis. The type A trichothecene mycotoxin T-2 was reported to

induce alterations upstream in the rat HPG axis, through increase of hypothalamic GnRH synthesis,

and downstream through precocious onset of the pubertal development of reproductive organs (Yang et

al. 2016). Downstream of the HPG axis, the type B trichothecene DON has been shown to have a

negative impact on bovine granulosa cell steroidogenesis and survival (Guerrero-Netro et al. 2015). In

rats, DON also increased progenitor Leydig cell proliferation, but inhibited their maturation and

function, including steroidogenesis (Yang et al. 2024). Only a few studies have investigated the toxicity

of mycotoxins midstream of the HPG axis. DON inhibited the growth of rat pituitary GH3 cells (Liu et

al. 2020). In porcine gonadotrope cells, the *Fusarium* toxins ZEA and α-ZOL specifically inhibited

FSH synthesis and secretion, but not that of LH (He et al. 2018). These FSH inhibiting effects relied on

the activation of the membrane estrogen receptor GPR30, and subsequent phosphorylation of protein

kinase C (PKC) and MAPK proteins ERK and p38. One of the main mechanisms underlying DON

toxicity involves the activation of MAPKs (Payros et al. 2016). Most studies have reported a rapid

increase in phosphorylation of JNK, ERK and p38 proteins in various cell types following exposure to

DON (Lee et al. 2019; Mishra et al. 2014; Zhang et al. 2020). However, the level of phosphorylated

ERK decreased with increasing DON concentrations in piglet hippocampal nerve cells, indicating that

the pattern of regulation of MAPKs may vary in a cell-type dependent manner (Wang et al. 2018).

GnRH is a central regulator of reproductive functions that acts by binding a G-protein coupled

receptor on the surface of the gonadotrope cells. The GnRH binding transmits signals, mostly via the

MAPK cascade, to increase the synthesis of LH and FSH (Melamed et al. 2012). Given the pivotal role

of MAPKs for both the toxicity of DON and GnRH activation of the synthesis and release of gonadotropin by gonadotrope cells, we wanted to know, midstream in the HPG axis, what type of interference in GnRH-induced gonadotropin production could result from gonadotrope cells exposure to DON.

We observed herein a significant decrease of the viability of the gonadotrope-like cells even at the lowest tested concentration (1nM) of DON. Similar to the cell death mechanisms we previously reported in the intestinal mucosa, the cytotoxicity of DON toward the gonadotrope cells showed here appears to be driven by apoptosis (Payros et al. 2021). This cytotoxicity and the concentration ranges it occurred are relevant since *in vivo* experiments have established that the transport of DON across the blood-brain barrier is effective, with the toxin being detected in the brain within minutes of oral exposure (Behrens et al. 2015; Prelusky et al. 1990). Moreover, as the pituitary gland sits outside the blood-brain barrier, the tiny population of gonadotrope cells that constitute only 7% to 15% of the total number of anterior pituitary cells, are even exposed to higher concentrations of dietary DON compared to other parts of the brain. Considering the provisional maximum tolerable daily intake (PMTDI) of DON established by the Joint FAO/WHO Expert Committee on Food Additives, it is anticipated that the concentrations of DON in blood and cerebrospinal fluid will be 1.5 nM and 0.45 nM respectively (Maresca 2013). DON has been highlighted as of significant risk to human health, owing to the less than 30 safety factor between the doses of this mycotoxin susceptible to be present in the organism, in relation to its provisional maximum tolerable daily intake (PMTDI), and the doses affecting cell functions (Maresca and Fantini 2010; Muri et al. 2009). The 1 to 10 nM DON concentration range in this study showed clear cytotoxicity to LβT2 cells. Assuming that human gonadotrope cells have similar sensitivity to DON, the safety factor would be 0.7 to 20, positioning the pituitary as a prominent target for DON action.

We observed that exposure of gonadotrope-like cells to DON prior to stimulation by GnRH decreases the induced phosphorylation levels of Erk, but not p38. DON regulate MAPKs via the ribotoxic stress response, which involves the double-stranded RNA-activated protein kinase (PKR) and hemoitopoeitic cell kinase (Hck) as upstream transducers (Pestka 2008). Instead of PKR and Hck, GnRH activation of the phosphorylation of Erk and p38-MAPKs is mediated by several isoforms of PKC (Harris et al. 1997; Maccario et al. 2004). Moreover, the profile of the PKC isoforms involved in

GnRH to p38MAPK phosphorylation differs from ERK1/2 activation. On top of the PKCβII, PKCδ

and PKCε isoforms reported for GnRH activation of the phosphorylation of ERK in LβT2 cells, the

GnRH induction of p38MAPK phosphorylation also involves PKCα (Mugami et al. 2018). Our results

suggest that, contrary to PKCβII, PKCδ and PKCε, the activation of MAPKs by GnRH via PKCα may

escape interference by DON. This hypothesis deserves to be investigated in the future.

The DON alteration of GnRH-induced phosphorylation of MAPKs in this study was followed by

impaired induction of key regulatory genes for LH production (*Cga* and *Lhb*), and dose-dependent

inhibition of the secretion of LH itself by gonadotrope cells. Although DON did not alter basal levels

of *Fshb*, it would be inaccurate to conclude that DON has no effect on FSH secretion, since the

experimental design of our study was specifically tailored to stimulate LH secretion rather than FSH

(GnRH one pulse approach). Although produced by the same gonadotrope cell, the release of LH or

FSH depends on specific GnRH pulsatility patterns, with LH release favored by fast GnRH pulse

frequencies and FSH favored by slow pulse frequencies (Tsutsumi and Webster 2009). To allow

simultaneous release of FSH and LH, mouse pituitary cells were pulsed for 2 minutes at a 10 nM

GnRH peak pulse concentration with a 58-minute interpulse interval for 4 hours (Kim and Lawson

2015), which is drastically different from the 0.1 to 1 nM GnRH stimulation for 15 minutes or six

hours used in the present study. Further investigations featuring the suitable pulsatility are needed to

confirm if DON is also involved in the synthesis and release of FSH.

In conclusion, our study shows the first evidence that DON affects GnRH-induced signaling in

gonadotrope-like cells through a mechanism that, at least in part, involves apoptosis and inhibition of

GnRH-induced Erk phosphorylation. Most importantly, DON affected LH production by gonadotrope-

like cells in a concentration-dependent manner. LH is a critical regulator of gonadal function in

mammals and its altered synthesis or secretion can affect fertility in both sexes. This research broadens

our knowledge of DON's neurotoxic effects and brings a new depth to the potential neuroendocrine

implications. This highlights the critical role of DON in compromising overall health, particularly its

detrimental effects on the reproductive system, not only through direct effects on reproductive organs

but also through interference with the control of critical hormones for reproduction. This provides

novel insights and considerations for elucidating the toxicological mechanisms of DON.

Acknowledgments:

The authors thank Dr. Pamela Mellon (UC-San Diego) and Dr. Daniel J. Bernard (McGill University-

Canada) for providing the LβT2 cells. The authors also thank the University of Virginia Ligand Core

that provided assistance in LH measurement in cell culture supernatants. The University of Virginia

Center for Research in Reproduction Ligand Assay and Analysis Core that is supported by the Eunice

Kennedy Shriver NICHD/NIH Grant R24HD102061

Funding: *This work was supported by Foundation J.-Louis Lévesque. Lingchen Yang and Guodong*

Cai received a scholarship from China Scholarship Council.

Financial interests: *Authors declare they have no financial interests.*

Author Contributions: *All authors contributed to the study conception and design. Conceptualization,*

Methodology and Funding acquisition were performed by *Imourana Alassane-Kpembi; Formal*

analysis, Investigation, Data curation and Writing—original draft preparation were performed by

Guodong Cai and Lingchen Yang; Writing—review and editing was performed by Francis Marien-

Bourgeois, Derek Boerboom, Gustavo Zamberlam, and Imourana Alassane-Kpembi. Supervision was

performed by Derek Boerboom, Gustavo Zamberlam and Imourana Alassane-Kpembi. All authors read

and approved the final manuscript.

Conflicts of Interest: *The authors declare no conflict of interest.*

References

Behrens M, Hüwel S, Galla HJ, Humpf HU (2015) Blood-Brain Barrier Effects of the Fusarium
Mycotoxins Deoxynivalenol, 3 Acetyldeoxynivalenol, and Moniliformin and Their Transfer to
the Brain. PloS one 10(11):e0143640 doi:10.1371/journal.pone.0143640

Bernard DJ, Tran S (2013) Mechanisms of activin-stimulated FSH synthesis: the story of a pig and a
FOX. Biology of reproduction 88(3):78 doi:10.1095/biolreprod.113.107797

Bliss SP, Miller A, Navratil AM, et al. (2009) ERK signaling in the pituitary is required for female but
not   male   fertility.   Molecular   endocrinology   (Baltimore,   Md)   23(7):1092-101
doi:10.1210/me.2009-0030

Cai G, Guerrero-Netro HM, Bian J, Oswald IP, Price C, Alassane-Kpembi I (2023) Real-life exposure
to Fusarium toxins deoxynivalenol and zearalenone triggers apoptosis and activates NLRP3
inflammasome   in   bovine   primary   theca   cells.   Mycotoxin   research   39(4):367-377
doi:10.1007/s12550-023-00499-x

Graham KE, Nusser KD, Low MJ (1999) LbetaT2 gonadotroph cells secrete follicle stimulating
hormone   (FSH)   in   response   to   active   A.   The   Journal   of   endocrinology   162(3):R1-5
doi:10.1677/joe.0.162r001

Guerrero-Netro HM, Chorfi Y, Price CA (2015) Effects of the mycotoxin deoxynivalenol on
steroidogenesis   and   apoptosis   in   granulosa   cells.   Reproduction   (Cambridge,   England)
149(6):555-61 doi:10.1530/rep-15-0018

Harris D, Reiss N, Naor Z (1997) Differential activation of protein kinase C delta and epsilon gene
expression by gonadotropin-releasing hormone in alphaT3-1 cells. Autoregulation by protein
kinase   C.   The   Journal   of   biological   chemistry   272(21):13534-40
doi:10.1074/jbc.272.21.13534

He J, Wei C, Li Y, et al. (2018) Zearalenone and alpha-zearalenol inhibit the synthesis and secretion of
pig follicle stimulating hormone via the non-classical estrogen membrane receptor GPR30.
Molecular and cellular endocrinology 461:43-54 doi:10.1016/j.mce.2017.08.010

Kahnamouyi S, Nouri M, Farzadi L, Darabi M, Hosseini V, Mehdizadeh A (2018) The role of mitogen-
activated protein kinase-extracellular receptor kinase pathway in female fertility outcomes: a
focus on pituitary gonadotropins regulation. Therapeutic advances in endocrinology and
metabolism 9(7):209-215 doi:10.1177/2042018818772775

Kim T, Lawson MA (2015) GnRH Regulates Gonadotropin Gene Expression Through NADPH/Dual
Oxidase-Derived   Reactive   Oxygen   Species.   Endocrinology   156(6):2185-99
doi:10.1210/en.2014-1709

Lawson MA, Tsutsumi R, Zhang H, et al. (2007) Pulse sensitivity of the luteinizing hormone beta
promoter is determined by a negative feedback loop Involving early growth response-1 and
Ngfi-A binding protein 1 and 2. Molecular endocrinology (Baltimore, Md) 21(5):1175-91
doi:10.1210/me.2006-0392

Lee JY, Lim W, Park S, Kim J, You S, Song G (2019) Deoxynivalenol induces apoptosis and disrupts
375   cellular homeostasis through MAPK signaling pathways in bovine mammary epithelial cells.
376   Environmental pollution (Barking, Essex : 1987) 252(Pt A):879-887
377   doi:10.1016/j.envpol.2019.06.001

378   Liu A, Hu S, Wu Q, et al. (2020) Epigenetic upregulation of galanin-like peptide mediates
379   deoxynivalenol induced-growth inhibition in pituitary cells. Toxicology and applied
380   pharmacology 403:115166 doi:10.1016/j.taap.2020.115166

381   Liu F, Austin DA, Mellon PL, Olefsky JM, Webster NJ (2002) GnRH activates ERK1/2 leading to the
382   induction of c-fos and LHbeta protein expression in LbetaT2 cells. Molecular endocrinology
383   (Baltimore, Md) 16(3):419-34 doi:10.1210/mend.16.3.0791

384   Maccario H, Junoy B, Poulin B, Boyer B, Enjalbert A, Drouva SV (2004) Protein kinase Cdelta as
385   gonadotropin-releasing hormone target isoenzyme in the alphaT3-1 gonadotrope cell line.
386   Neuroendocrinology 79(4):204-20 doi:10.1159/000078102

387   Maresca M (2013) From the gut to the brain: journey and pathophysiological effects of the food-
388   associated trichothecene mycotoxin deoxynivalenol. Toxins 5(4):784-820
389   doi:10.3390/toxins5040784

390   Maresca M, Fantini J (2010) Some food-associated mycotoxins as potential risk factors in humans
391   predisposed to chronic intestinal inflammatory diseases. Toxicon : official journal of the
392   International Society on Toxinology 56(3):282-94 doi:10.1016/j.toxicon.2010.04.016

393   Melamed P, Savulescu D, Lim S, et al. (2012) Gonadotrophin-releasing hormone signalling
394   downstream of calmodulin. Journal of neuroendocrinology 24(12):1463-75
395   doi:10.1111/j.1365-2826.2012.02359.x

396   Mishra S, Srivastava S, Dewangan J, Divakar A, Kumar Rath S (2020) Global occurrence of
397   deoxynivalenol in food commodities and exposure risk assessment in humans in the last
398   decade: a survey. Critical reviews in food science and nutrition 60(8):1346-1374
399   doi:10.1080/10408398.2019.1571479

400   Mishra S, Tripathi A, Chaudhari BP, Dwivedi PD, Pandey HP, Das M (2014) Deoxynivalenol induced
401   mouse skin cell proliferation and inflammation via MAPK pathway. Toxicology and applied
402   pharmacology 279(2):186-97 doi:10.1016/j.taap.2014.06.003

403   Mugami S, Dobkin-Bekman M, Rahamim-Ben Navi L, Naor Z (2018) Differential roles of PKC
404   isoforms (PKCs) in GnRH stimulation of MAPK phosphorylation in gonadotrope derived
405   cells. Molecular and cellular endocrinology 463:97-105 doi:10.1016/j.mce.2017.04.004

406   Muri SD, van der Voet H, Boon PE, van Klaveren JD, Brüschweiler BJ (2009) Comparison of human
407   health risks resulting from exposure to fungicides and mycotoxins via food. Food and
408   chemical toxicology : an international journal published for the British Industrial Biological
409   Research Association 47(12):2963-74 doi:10.1016/j.fct.2009.03.035

410   Murphy LO, Blenis J (2006) MAPK signal specificity: the right place at the right time. Trends in
411   biochemical sciences 31(5):268-75 doi:10.1016/j.tibs.2006.03.009

Payros D, Alassane-Kpembi I, Laffitte J, et al. (2021) Dietary Exposure to the Food Contaminant Deoxynivalenol Triggers Colonic Breakdown by Activating the Mitochondrial and the Death Receptor Pathways. Molecular nutrition & food research 65(23):e2100191 doi:10.1002/mnfr.202100191

Payros D, Alassane-Kpembi I, Pierron A, Loiseau N, Pinton P, Oswald IP (2016) Toxicology of deoxynivalenol and its acetylated and modified forms. Archives of toxicology 90(12):2931-2957 doi:10.1007/s00204-016-1826-4

Pestka JJ (2008) Mechanisms of deoxynivalenol-induced gene expression and apoptosis. Food Additives & Contaminants: Part A 25(9):1128-1140 doi:10.1080/02652030802056626

Pestka JJ (2010) Deoxynivalenol: mechanisms of action, human exposure, and toxicological relevance. Archives of toxicology 84(9):663-79 doi:10.1007/s00204-010-0579-8

Pfaffl MW (2001) A new mathematical model for relative quantification in real-time RT-PCR. Nucleic acids research 29(9):e45 doi:10.1093/nar/29.9.e45

Pizzo F, Caloni F, Schutz LF, Totty ML, Spicer LJ (2015) Individual and combined effects of deoxynivalenol and α-zearalenol on cell proliferation and steroidogenesis of granulosa cells in cattle. Environmental toxicology and pharmacology 40(3):722-8 doi:10.1016/j.etap.2015.08.025

Prelusky DB, Hartin KE, Trenholm HL (1990) Distribution of deoxynivalenol in cerebral spinal fluid following administration to swine and sheep. Journal of environmental science and health Part B, Pesticides, food contaminants, and agricultural wastes 25(3):395-413 doi:10.1080/03601239009372697

Singh R, Kaur S, Yadav S, Bhatia S (2023) Gonadotropins as pharmacological agents in assisted reproductive technology and polycystic ovary syndrome. Trends in endocrinology and metabolism: TEM 34(4):194-215 doi:10.1016/j.tem.2023.02.002

Sprando RL, Collins TF, Black TN, et al. (2005) Characterization of the effect of deoxynivalenol on selected male reproductive endpoints. Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association 43(4):623-35 doi:10.1016/j.fct.2004.12.017

Sprando RL, Pestka J, Collins TF, et al. (1999) The effect of vomitoxin (Deoxnivalenol) on testicular morphology, testicular spermatid counts and epididymal sperm counts in IL-6KO [B6129-IL6 [TmlKopf] (IL-6 gene deficient)] and WT [B6129F2 (wild type to B6129-IL6 with an intact IL-6 gene)] mice. Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association 37(11):1073-9 doi:10.1016/s0278-6915(99)00103-9

Streit E, Schatzmayr G, Tassis P, et al. (2012) Current Situation of Mycotoxin Contamination and Co-occurrence in Animal Feed—Focus on Europe. 4(10):788-809

Tassis PD, Reisinger N, Nagl V, et al. (2022) Comparative Effects of Deoxynivalenol, Zearalenone and Its Modified Forms De-Epoxy-Deoxynivalenol and Hydrolyzed Zearalenone on Boar Semen

In Vitro. Toxins 14(7) doi:10.3390/toxins14070497

Tremblay JJ, Drouin J (1999) Egr-1 is a downstream effector of GnRH and synergizes by direct
interaction with Ptx1 and SF-1 to enhance luteinizing hormone beta gene transcription.
Molecular and cellular biology 19(4):2567-76 doi:10.1128/mcb.19.4.2567

Tsutsumi R, Webster NJ (2009) GnRH pulsatility, the pituitary response and reproductive dysfunction.
Endocrine journal 56(6):729-37 doi:10.1507/endocrj.k09e-185

Turgeon JL, Kimura Y, Waring DW, Mellon PL (1996) Steroid and pulsatile gonadotropin-releasing
hormone (GnRH) regulation of luteinizing hormone and GnRH receptor in a novel
gonadotrope cell line. Molecular endocrinology (Baltimore, Md) 10(4):439-50
doi:10.1210/mend.10.4.8721988

Ulloa-Aguirre A, Lira-Albarrán S (2016) Clinical Applications of Gonadotropins in the Male. Progress
in   molecular   biology   and   translational   science   143:121-174
doi:10.1016/bs.pmbts.2016.08.003

Wang X, Fan M, Chu X, et al. (2018) Deoxynivalenol induces toxicity and apoptosis in piglet
hippocampal nerve cells via the MAPK signaling pathway. Toxicon : official journal of the
International Society on Toxinology 155:1-8 doi:10.1016/j.toxicon.2018.09.006

Yang J, Ye L, Cui R, et al. (2024) Deoxynivalenol Inhibits Progenitor Leydig Cell Development by
Stimulating Mitochondrial Fission in Rats. Journal of agricultural and food chemistry
72(18):10616-10626 doi:10.1021/acs.jafc.4c01151

Yang R, Wang YM, Zhang L, Zhao ZM, Zhao J, Peng SQ (2016) Prepubertal exposure to T-2 toxin
advances pubertal onset and development in female rats via promoting the onset of
hypothalamic-pituitary-gonadal axis function. Human & experimental toxicology
35(12):1276-1285 doi:10.1177/0960327116629529

Zhang H, Deng X, Zhou C, Wu W, Zhang H (2020) Deoxynivalenol Induces Inflammation in IPEC-J2
Cells by Activating P38 Mapk And Erk1/2. Toxins 12(3) doi:10.3390/toxins12030180

Figure 1

Figure 2

483
484

485   Figure 3

486
487

488   Figure 4

489
490

491   Figure 5

492
493

494 Figure 6

495
496

497 Graphical abstract

498

499

500 Table 1. Primer sequences of RT-qPCR

Gene	Upstream primers（5'→3'）	Downstream primers（5'→3'）
Rpl19	CTGAAGGTCAAAGGGAATGTG	GGACAGAGTCTTGATGATCTC
Gnrhr	TTCGCTACCTCCTTTGTCGT	CACGGGTTTAGGAAAGCAAA
Cga	TCCCTCAAAAAGTCCAGAGC	GAAGAGAATGAAGAATATGCAG
Fshb	GCTACTGCTACACTAGGGATCTGG	TGGTCCTTCAGGACAAGGGT
Lhb	ACTGTGCCGGCCTGTCAACG	AGCAGCCGGCAGTACTCGGA

Figure captions

Figure 1. Cytotoxic effects of DON on LβT2 cells.

Cell viability (%) of was determined by CellTiter Blue in LβT2 cell exposed to DON for 24 and 48 h in

triplicate, and the experiment was repeated 3 times (n=3). Significant differences post 24 h treatments

compared with control, *$p < 0.05$; ** $p < 0.01$. Significant difference post 48 h treatments compared

with control, #$p < 0.05$; ##$p < 0.01$.

Figure 2. DON induces apoptosis in LβT2 cells.

RealTime-Glo™ Annexin V Apoptosis and Necrosis Assay detection of apoptosis (A) and secondary

necrosis (B) in LβT2 cells treated with DON (10 nM) in triplicate, and the experiment was repeated 3

times (n=3). Significant differences post 24 h treatment compared with control, *$p < 0.05$; ** $p < 0.01$.

Figure 3. Effects of DON on the expression and phosphorylation of ERK 1/2 proteins in LβT2 cells

stimulated by GnRH.

LβT2 cells were pre-exposed for 24 h to DON at concentrations of 1 μM and 10 μM before challenging

them with distinct doses of GnRH for 15 min. Protein levels were measured by Western blot and total

and phosphorylated levels of ERK 1/2 were normalized to β-actin. The experiment was repeated 3

times (n=3). Significant effect of GnRH treatment compared with control, *** $p < 0.001$; Significant

effect of DON treatment compared with GnRH alone, ##$p < 0.01$; ###$p < 0.001$.

Figure 4. Effects of DON on the expression and phosphorylation of p38-MAPK protein in LβT2 cells

stimulated by GnRH.

LβT2 cells were pre-exposed for 24 h to DON at concentrations of 1 μM and 10 μM before challenging

them with distinct doses of GnRH for 15 min. Protein levels were measured by Western blot and total

and phosphorylated levels of p38-MAPK were normalized to β-actin. The experiment was repeated 3

times (n=3). Significant effect of GnRH and/or DON treatments compared with controls, *$p < 0.05$;

*** *p < 0.001*.

Figure 5. Effects of DON on *Gnrhr*, *Cga*, *Fshb* and *Lhb* gene expression in LβT2 cells stimulated by

GnRH.

LβT2 cells were pre-exposed for 24 h to DON at concentrations of 1 μM or 10 μM before challenging

them with distinct doses of GnRH for 6 h. Cells were collected for RNA extraction and mRNA levels

of *Gnrhr*, *Cga*, *Fshb* and *Lhb* were assessed by RT-qPCR and normalized to the housekeeping gene

Rpl19. The experiment was repeated 3 times (n=3). Significant effects of GnRH treatment compared

with untreated control, $*p < 0.05$; $**p < 0.01$; $*** \ p < 0.0001$. Significant effect of DON+GnRH

treatment relative to GnRH-treated control, $^{####}p < 0.0001$.

Figure 6. Effects of DON on LH concentration in culture medium of LβT2 cells stimulated by GnRH.

LβT2 cells were treated or not with graded concentrations of DON for 24 hours, prior to stimulation

with 1nM GnRH for 6 hours. Cell culture media samples were collected and then analyzed by Ultra-

Sensitive Mouse and Rat LH ELISA. The experiment was repeated 3 times (n=3). Significant

differences of GnRH and/or DON treatments compared with untreated control, $*p < 0.05$; $** \ p < 0.01$.

Significant effect of DON+GnRH treatment relative to GnRH-treated control, $^{#}p < 0.05$); $^{##} \ p < 0.01$.