The conserved ATPase PCH-2 controls the number and distribution of crossovers by

antagonizing their formation in *C. elegans*

Bhumil Patel, Maryke Grobler, Alberto Herrera, Elias Logari, Valery Ortiz, and Needhi Bhalla[1]

Department of Molecular, Cell and Developmental Biology, University of California, Santa Cruz, Santa

Cruz, CA 95064

[1] corresponding author: nbhalla@ucsc.edu

Department of Molecular, Cell and Developmental Biology

225 Sinsheimer Labs

University of California, Santa Cruz

Santa Cruz, CA 95064

phone: (831) 459-1319

fax: (831) 459-3139

Abstract

Meiotic crossover recombination is essential for both accurate chromosome segregation and the

generation of new haplotypes for natural selection to act upon. This requirement is known as

crossover assurance and is one example of crossover control. While the conserved role of the

ATPase, PCH-2, during meiotic prophase has been enigmatic, a universal phenotype when *pch-2* or

its orthologs are mutated is a change in the number and distribution of meiotic crossovers. Here, we

show that PCH-2 controls the number and distribution of crossovers by antagonizing their formation.

This antagonism produces different effects at different stages of meiotic prophase: early in meiotic

prophase, PCH-2 prevents double strand breaks from becoming crossover-eligible intermediates,

limiting crossover formation at sites of initial double strand break formation and homolog interactions.

Later in meiotic prophase, PCH-2 winnows the number of crossover-eligible intermediates,

contributing to the designation of crossovers and ultimately, crossover assurance. We also

demonstrate that PCH-2 accomplishes this regulation through the meiotic HORMAD, HIM-3. Our data

strongly support a model in which PCH-2's conserved role is to remodel meiotic HORMADs

throughout meiotic prophase to destabilize crossover-eligible precursors, coordinate meiotic

recombination with synapsis, and contribute to the progressive implementation of meiotic

recombination, guaranteeing crossover control.

Introduction

Meiosis is a specialized type of cell division that reduces chromosome number by half, resulting in the production of genetically diverse haploid gametes, so that fertilization during sexual reproduction restores diploidy. This process occurs in two stages: meiosis I, in which homologous chromosomes are partitioned, and meiosis II, in which sister chromatid are segregated. The regulation of meiosis is crucial for ensuring that both genetic recombination and chromosome segregation occur accurately. Errors in human meiosis are associated with birth defects, such as Down and Turner syndromes, infertility and miscarriages, underscoring the importance of understanding meiosis to human health (Hassold and Hunt, 2001).

During prophase I, homologous chromosomes pair and this pairing is stabilized through a process called synapsis, in which a protein structure called the synaptonemal complex (SC) holds homologs together. This close association between homologs during synapsis facilitates crossover formation, the process in which double strand breaks (DSBs) are deliberately introduced into the genome, repaired by meiosis-specific mechanisms that exchange DNA between homologous chromosomes and generate the chiasmata, or linkage, that promotes accurate meiotic chromosome segregation. Therefore, disruptions in pairing, synapsis, or recombination prevent the formation of chiasmata and can lead to meiotic errors such as nondisjunction, resulting in aneuploid gametes.

In addition to the fundamental role that recombination plays in ensuring accurate chromosome segregation, crossover recombination accomplishes another important function: it generates new haplotypes for natural selection to act upon to drive evolution. Thus, to assure a random assortment of alleles on a population level, the distribution of crossovers may be as tightly regulated as their number (Veller et al., 2019). The significance of controlling both crossover number and distribution is clearly illustrated by the existence of mechanisms such as crossover assurance, in which every pair of homologous chromosomes gets at least one crossover; crossover homeostasis, in which the number

of crossovers remains relatively invariant even if the number of recombination precursors change; and

crossover interference, in which the presence of a crossover inhibits the formation of a crossover

nearby (Gray and Cohen, 2016). DSBs typically vastly outnumber crossovers in most organisms and

are introduced gradually throughout early prophase (Joshi et al., 2015; Woglar and Villeneuve, 2018).

Therefore, to accomplish this precise level of control, meiotic crossover recombination and the

decision about which DSBs become crossover-eligible intermediates, and eventually, which crossover-

eligible intermediates get designated as crossovers, is implemented progressively throughout meiotic

prophase (Cole et al., 2012; Joshi et al., 2015; Morgan et al., 2021; Yokoo et al., 2012). In many

systems, transitions from DSBs to crossover-eligible intermediates, and crossover-eligible

intermediates to crossovers, can be molecularly and/or cytologically monitored.

PCH-2, also known as TRIP13 in mammals, is an evolutionarily ancient AAA-ATPase that plays a

significant role in regulating meiosis across different organisms, including *M. musculus* (mice), *S.*

cerevisiae (budding yeast), *D. melanogaster* (fruit flies), and *C. elegans* (worms) (Bhalla, 2023). PCH-

2 and its orthologs structurally remodel a family of proteins with HORMA domains (HORMADs) to

control their function (Gu et al., 2022). HORMADs participate in a variety of signaling events and can

exist in at least three structurally distinct conformations: a "closed" conformation, which they adopt

when they bind a short peptide sequence in their own protein sequence or another protein (also called

a closure motif) and their C-terminus wraps around this motif to stabilize the interaction; an "open"

conformation when unbound, in which their C-terminus is discretely tucked against the HORMA

domain; and an "extended" conformation, which is an intermediate between the two (Gu et al., 2022).

PCH-2 and its orthologs convert the closed version of HORMADs to the open or extended versions,

playing an important role in recycling HORMADs during signaling. During meiosis, closed versions of

meiotic HORMADs assemble on chromosomes to form meiotic chromosome axes, which are essential

for pairing, synapsis and recombination between homologous chromosomes (Kim et al., 2014). The

remodeling of meiotic HORMADs by PCH-2 to an "open" or "extended" conformation is thought to

reduce the levels of HORMADs on chromosomes (Borner et al., 2008; Cuacos et al., 2021; Lambing

et al., 2015; Wojtasz et al., 2009) and/or increase their dynamic association and dissociation (Russo et

al., 2023), modulating and coordinating homolog pairing, synapsis and recombination during

prophase.

The conserved role of the PCH-2/HORMAD module in meiosis has been difficult to characterize, in

part, we have argued, because of the evolutionary innovation that is an inherent aspect of sexual

reproduction (Bhalla, 2023). Moreover, in some systems, such as budding yeast and plants (Herruzo

et al., 2021; Yang et al., 2020), PCH-2 orthologs not only remodel meiotic HORMADs on meiotic

chromosomes but also perform this function in the cytoplasm to make meiotic HORMADs available for

their role(s) in meiotic nuclei. This dual role can complicate functional analyses, particularly in *pch-2*

null mutants. However, all meiotic systems exhibit defects in the number and distribution of crossovers

when PCH-2 function is abrogated by mutation (Deshong et al., 2014; Joshi et al., 2009; Joyce and

McKim, 2009; Lambing et al., 2015; Roig et al., 2010; Zanders and Alani, 2009). Unfortunately, the

lack of a clear pattern when analyzing these defects in recombination in *pch-2* mutants has

contributed to an inability to develop a unified, integrated model of PCH-2 function in the field.

In *C. elegans*, meiotic HORMADs localize to meiotic chromosomes independently of PCH-2 (Deshong

et al., 2014). Moreover, the progressive implementation of meiotic recombination can be cytologically

monitored in meiotic nuclei that are organized both spatially and temporally in the *C. elegans* germline

(Woglar and Villeneuve, 2018; Yokoo et al., 2012). Here, we exploit this system to show that PCH-2 is

required to control the number and distribution of crossovers by antagonizing their formation. This

antagonism produces different consequences depending on the stage of meiotic prophase. In early

meiotic prophase, PCH-2 inhibits DSBs from becoming crossover-eligible intermediates, ensuring that

crossovers are more widely distributed than sites of initial DSB formation and/or homolog interactions.

Later in meiotic prophase, PCH-2 is responsible for winnowing the numbers of crossover-eligible

intermediates on synapsed chromosomes, contributing to the designation of crossovers and ultimately,

crossover assurance. Genetic analysis demonstrates that PCH-2's regulation of crossover-eligible

intermediates is through one of three essential meiotic HORMADs, HIM-3. Finally, we link PCH-2's

effect on early DSBs in early meiotic prophase to cell cycle stage, demonstrating that both limit early

DSBs from becoming crossovers. We propose that PCH-2's remodeling of HIM-3 on meiotic

chromosomes destabilizes crossover-eligible intermediates throughout meiotic prophase, contributing

to the progressive implementation of meiotic recombination to control the number and distribution of

crossovers, also known as crossover control.

Results

PCH-2 controls the number and distribution of crossovers in similar patterns on multiple

chromosomes

We had previously shown that loss of PCH-2 reduced the frequency of double crossovers and genetic

length of both an autosome (chromosome III) and the X chromosome, albeit not uniformly among

genetic intervals (Deshong et al., 2014). To determine whether a more obvious pattern could be

observed, we expanded our analysis by monitoring recombination genetically in both wildtype and

pch-2 mutant animals using five single nucleotide polymorphisms (SNPs) that spanned 95% of

Chromosomes, I, III, IV and the X chromosome (Figure 1). We excluded Chromosome II from our

analysis because of the potential difficulty combining Hawaiian SNPs with the *pch-2* mutation, which is

linked to Chromosome II, and Chromosome V because of our use of the *bcIs39* transgene to identify

cross progeny, which may disrupt recombination on that chromosome.

In wildtype animals, we observed multiple double crossovers, ranging from 1-13, depending on the

chromosome. In a majority of these double crossovers (69%), one crossover was at the end of the

chromosome where pairing and synapsis initiate, also called the Pairing Center (PC) (MacQueen et

al., 2005), suggesting some relationship between where chromosomes might make initial contacts and

the likelihood of double crossovers. On all chromosomes analyzed, we observed no double

crossovers in *pch-2* mutants and this difference was statistically significant for Chromosomes I, III and

X. Moreover, there was a striking and consistent shift of crossovers to the PC end of all four

chromosomes tested. This shift in the distribution of crossovers to the PC ends of chromosomes was

generally accompanied by a reduction in crossovers in the center of chromosomes. In *C. elegans*, the

center of chromosomes are where double strand breaks are less numerous (Nadarajan et al., 2021;

Yu et al., 2016) and where genes are more abundant. In the case of the X chromosome and

chromosome III, recombination at the non-PC end was also reduced and appeared to more closely

resemble the physical map at this end of these chromosomes. Thus, PCH-2 ensures a wider

distribution of crossovers across chromosomes, away from regions that are more likely to undergo

early homolog interactions (PC ends) and with more double strand breaks (both the PC and non-PC

ends of chromosomes), and towards the center of chromosomes, where double strand breaks are less

abundant.

PCH-2 prevents exogenous DSBs early in meiotic prophase from becoming crossovers

We previously showed that PCH-2 promotes crossover formation and crossover assurance through its

regulation of the meiotic HORMAD, HIM-3, in *C. elegans* (Russo et al., 2023). Moreover, this

promoting role seems linked to its localization to the synaptonemal complex (Patel et al., 2023).

However, the observed shift in the distribution of crossovers in *pch-2* mutants suggests that PCH-2

may also play a role in inhibiting crossovers and that this role may be occurring in early meiotic

prophase, when chromosomes are undergoing initial homolog interactions. We had previously

observed that meiotic nuclei in early prophase were more likely to produce crossovers when DSBs

were induced by excision of the *Mos* transposon in *pch-2* mutants than in control animals but

experimental caveats limited our ability to properly interpret this experiment (Deshong et al., 2014).

To explicitly test this possibility, we took advantage of both the spatiotemporal organization of meiotic

nuclei in the *C. elegans* germline, and the observation that nuclei travel in an assembly line process at

a stereotypical pace to late pachytene (Jaramillo-Lambert et al., 2007), where we can cytologically

assess crossover formation by staining for the essential crossover factor, COSA-1 (Yokoo et al.,

2012). We performed irradiation experiments to introduce exogenous DSBs at different timepoints

during meiotic prophase and analyzed PCH-2's role in crossover formation (Figure 2A).

In this experiment, we tested the effects of ionizing radiation at two distinct time points: 8 hours after

irradiation, when we can monitor crossover formation in late pachytene nuclei that received

exogenous DSBs in mid-pachytene, and 24 hours after irradiation, when we monitor crossover

formation in late pachytene nuclei that received exogenous DSBs in leptotene/zygotene, also known

as the transition zone in *C. elegans* (Figure 2A). In order to detect crossovers in late pachytene,

control and *pch-2* mutants germlines were stained for GFP::COSA-1, which allowed us to visualize

crossovers, and DAPI, which allowed us to visualize DNA. In unirradiated control worms, 94% of late

pachytene nuclei have 6 GFP::COSA-1 foci, one per chromosome pair (Figure 2B). In unirradiated

pch-2 mutants, the percentage of nuclei have 6 GFP::COSA-1 foci drops to 82%, primarily because of

the significant increase in meiotic nuclei with less than 6 GFP::COSA-1 foci (17%) (Figure 2B),

consistent with our previous report that loss of PCH-2 leads to a decrease in crossover formation

(Deshong et al., 2014). At 8 hours after irradiation, *pch-2* mutants' defect in crossover assurance

becomes even more pronounced. We observed a significant increase in nuclei containing less than 6

COSA-1 nuclei in *pch-2* mutants (26%), when compared to control (9%) germlines (Figures 2C and

2D). However, we did not observe a significant number of nuclei with greater than 6 COSA-1 foci in

either backgrounds. This observation reinforces previous findings that exogenous DSBs introduced in

mid-pachytene do not affect the number of crossovers in wildtype animals (Yokoo et al., 2012).

Next, we examined crossover formation 24 hours post-irradiation (Figures 2E and 2F). For this time

point, we also detected a significant loss of crossover assurance in *pch-2* mutants, compared to

control worms. However, in contrast to the 8 hour time point, we also saw a significant increase in

nuclei containing less than 6 COSA-1 foci in both control (13%) and *pch-2* mutants (26%), compared

to unirradiated worms (p values = 0.0008 and < 0.0001, respectively, Fischer's Exact Test), indicating

that even in control animals, extra DSBs in early meiotic prophase can disrupt crossover assurance. In

addition, we also observed a significant increase in nuclei with more than 6 COSA-1 foci in *pch-2*

mutants (wildtype: 2%, *pch-2*: 13%), demonstrating that loss of PCH-2 leads to increased crossover

formation when nuclei in the transition zone get more DSBs. In other words, PCH-2 inhibits

exogenous DSBs introduced in early meiotic prophase from becoming crossovers, supporting our

recombination analysis (Figure 1).

PCH-2 prevents SPO-11-induced DSBs from becoming crossovers in early meiotic prophase

Our previous experiment indicated that PCH-2 is preventing exogenous DSBs in early meiotic

prophase from becoming crossovers (Figure 2). To test whether this was also the case for

programmed meiotic DSB formation, we used the auxin-inducible degradation (AID) system to remove

the enzyme that is responsible for programmed meiotic DSBs, SPO-11 (*spo-11::AID::3XFLAG*)

(Dernburg et al., 1998; Keeney et al., 1997). When we treated both control and *pch-2* mutant worms

for 36 hours with auxin (Figure 3A), we observed a complete loss of GFP::COSA-1 foci (Figure 3C).

When these animals were treated with auxin for 48 hours, to allow nuclei without SPO-11 an additional

12 hours to reach diakinesis (Figure 3D), we observed 12 DAPI stained bodies, or univalents, which

are the 6 homolog pairs that have not formed chiasmata (Figure 3F). Thus, we can reliably remove

SPO-11 from meiotic nuclei throughout the germline and prevent crossover formation.

Since meiotic nuclei travel through the germline at about one cell per row per hour (Jaramillo-Lambert

et al., 2007), we took advantage of this precise spatiotemporal resolution to test four different

timepoints in early meiotic prophase to determine if acute depletion of SPO-11 leads to any changes in

crossover formation between control animals and *pch-2* mutants (Figures 3A and D). In this

experiment, we assayed crossover formation with two experimental approaches. Similar to our

irradiation experiment (Figures 2E and 2F), we assayed crossover formation by analyzing the number

of GFP::COSA-1 foci in late pachytene 24, 22, 20 and 18 hours after acutely depleting SPO-11 (Figure

3A). In addition to this approach, we analyzed the presence of bivalents in diakinesis (Figure 3D),

which indicates the successful formation of chiasmata between homologous chromosomes. Since

meiotic nuclei travel from late pachytene to diakinesis in approximately 12 hours (Deshong et al.,

2014), the 36, 34, 32 and 30 hour timepoints to assay bivalent formation correspond to the 24, 22, 20

and 18 hour timepoints, respectively, in which GFP::COSA-1 were analyzed. We also performed a

control experiment, treating worms with ethanol for 24 and 36 hour timepoints to ensure that this

tagged version of SPO-11 was fully functional for crossover formation (Figures 3C and 3F).

At the 24 hour time point (Figures 3A and 3C), when meiotic nuclei in the transition zone do not

receive SPO-11-induced DSBs, we observe a range of GFP::COSA-1 foci in late pachytene in both

control animals and *pch-2* mutants with a majority of nuclei having 0 or 1 focus (control and *pch-2*

averages: 1.8 GFP::COSA-1 foci), indicating a loss of crossover formation, albeit with some

heterogeneity. This heterogeneity likely reflects that it takes 1-4 hours to significantly affect DSB

formation through acute degradation of SPO-11 at this stage of meiosis in *C. elegans* (Hicks et al.,

2022). In addition, there was no statistically significant difference in the number of GFP::COSA-1 foci

between control animals and *pch-2* mutants. However, when we assayed bivalent formation 12 hours

later (see 36 hours auxin treatment in Figures 3D and 3F), we observed the same heterogeneity and a

slight, statistically significant difference in the number of DAPI stained bodies between control animals

(average number of DAPI stained bodies: 10.6) and *pch-2* mutants (average number of DAPI stained

bodies: 9.6), suggesting that even at this very early time point, we can detect a role for PCH-2 in

preventing early SPO-11-induced DSBs from becoming crossovers.

We continued to monitor GFP::COSA-1 and bivalent formation at successive timepoints. At the 22

hour and 34 hour timepoints, where SPO-11 depletion begins later in the transition zone, we saw an

increase in the number of GFP::COSA-1 foci (Figure 3C) and a reduction in the number of DAPI

stained bodies (Figure 3F), suggesting that more DSBs are becoming crossovers. In our analysis of

GFP::COSA-1 foci at this timepoint, there appeared to a bimodal distribution: the majority of nuclei in

both genotypes fell into two clear classes, those with no COSA-1 foci (24% in control animals, 26% in

pch-2 mutants) and those with 6 COSA-1 foci (42% in both control animals and *pch-2* mutants)

(Figure 3C). The number of crossovers continued to increase at the 20 hour and 32 hour timepoints,

eventually reaching wildtype numbers for GFP::COSA-1 and DAPI stained bodies by the 18 hour and

30 hour timepoints, respectively (Figures 3C and 3F).

At the 22 hour and 34 hour timepoints, we did not detect any statistically significant difference in the

number of GFP::COSA-1 foci or DAPI stained bodies between control animals and *pch-2* mutants

(Figures 3C and 3F), even when we analyzed the number of nuclei that had greater than 6

GFP::COSA-1 foci. In contrast, while the number of GFP::COSA-1 foci was not statistically

significantly different between control animals and *pch-2* mutants at the 20 hour time point (Figure

3C), the number of nuclei with greater than 6 GFP::COSA-1 foci was significantly higher in *pch-2*

mutants (control: 3%, *pch-2*: 10%, p value = 0.009, Fischer's Exact Test), similar to our analysis of

GFP::COSA-1 24 hours after irradiation. Even more strikingly, the number of DAPI stained bodies was

significantly lower in *pch-2* animals than control animals at the corresponding 32 hour timepoint

(Figure 3F), indicating that more homolog pairs had successfully formed chiasmata in *pch-2* mutants.

No difference in GFP::COSA-1 foci or bivalent formation was observed at the 18 hour and 30 hours

timepoints, respectively (Figures 3C and 3F). These data indicate that during a narrow time frame of

early meiotic prophase, likely during leptotene/zygotene given the delay in limiting DSB formation with

auxin-induced degradation of SPO-11, PCH-2 prevents SPO-11-induced DSBs from becoming

crossovers and chiasmata.

PCH-2 is required for timely loading and removal of MSH-5 on meiotic chromosomes through

its regulation of HIM-3

After DSB formation, a subset of DSBs are licensed to be repaired through a pro-crossover pathway

and eventually winnowed to a stereotypical number of crossovers (Yokoo et al., 2012). In *C. elegans*,

these crossover-eligible intermediates can be visualized by the loading of the pro-crossover factor

MSH-5, a component of the meiosis-specific MutS γ complex that stabilizes crossover-specific DNA

repair intermediates called joint molecules (Janisiw et al., 2018; Kelly et al., 2000; Snowden et al.,

2004). In the spatiotemporally organized meiotic nuclei of the germline, a functional GFP tagged

version of MSH-5, GFP::MSH-5, begins to form a few foci in leptotene/zygotene (the transition zone),

becoming more numerous in early pachytene before decreasing in number in mid pachytene to

ultimately colocalize with COSA-1 marked sites in late pachytene in a process called designation

(Janisiw et al., 2018; Woglar and Villeneuve, 2018; Yokoo et al., 2012; see control in Figures 4A and

4B). The mechanism through which MSH-5 functions with the SC and other meiosis factors to

ultimately form crossovers has not been established but may involve phosphorylation by cyclin-

dependent kinases (Haversat et al., 2022; Zhang et al., 2021).

Given that we have shown that PCH-2 prevents early DSBs from becoming crossovers (Figures 2 and

3), we tested whether we could cytologically detect this inhibition at the level of MSH-5 behavior in

control animals and *pch-2* mutants. We generated wildtype and *pch-2* mutants with GFP::MSH-5 and

quantified the average number of GFP::MSH-5 foci per row of nuclei from the transition zone to late

pachytene (Figures 4A and 4B). We observed a statistically significant increase in the average number

of GFP::MSH-5 foci per row in the transition zone in *pch-2* mutants (Figures 4A and 4B, p value <

0.0001, Student's t-test), indicating that PCH-2 typically limits MSH-5 loading at this early stage of

meiotic prophase. The average number of GFP::MSH-5 foci per row were similar in number in early

pachytene in both backgrounds. However, we also observed a substantial increase in the average

number of GFP::MSH-5 foci per row in *pch-2* mutants in the mid and late pachytene regions (Figures

4A and 4B), indicating that PCH-2 also promotes the turnover or maturation of GFP::MSH-5 foci at

these later stages of meiotic prophase. Thus, PCH-2 prevents the loading of the crossover-promoting

factor, MSH-5, limiting the formation of crossover-eligible intermediates during early meiotic prophase,

consistent with both our genetic analysis of recombination (Figure 1) and PCH-2's role in preventing

early DSBs from becoming crossovers (Figures 2 and 3). However, PCH-2 also appears to limit the

number of crossover-eligible intermediates in mid and late pachytene, an unexpected observation

given that the most severe recombination defect in *pch-2* mutants is the loss of crossover assurance

(Deshong et al., 2014).

We previously showed that PCH-2 genetically interacts with meiotic HORMADs to control different

aspects of meiosis. A mutant version of the essential meiotic HORMAD, HIM-3^{R93Y}, binds its closure

motif with reduced affinity *in vitro* and we have proposed that an analogous mutation in another

essential meiotic HORMAD, HTP-3^{H96Y}, behaves similarly (Russo et al., 2023). These mutations

suppress different meiotic defects in *pch-2* mutants *in vivo*, indicating that PCH-2 regulates pairing and

synapsis through its regulation of HTP-3 and crossover recombination through its regulation of HIM-3

(Russo et al., 2023). To determine which meiotic HORMAD PCH-2 might be regulating to affect

GFP::MSH-5's loading and removal on meiotic chromosomes, and the potential role for these early

crossover-eligible intermediates, we constructed *gfp::msh-5;htp-3^{H96Y}, gfp::msh-5;pch-2;htp-3^{H96Y},*

gfp::msh-5;him-3^{R93Y} and *gfp::msh-5;pch-2;him-3^{R93Y}* mutants and quantified GFP::MSH-5 foci

throughout the germline. Both *htp-3^{H96Y}* and *him-3^{R93Y}* mutants showed a drastic increase in the

average number of GFP::MSH-5 per row of nuclei throughout the germline (Figures 4C and 4D,

Supplemental Figure 1). For example, compared to control germlines that peaked at approximately an

average of 20 MSH-5 foci per row in early pachytene and then decreased in mid-pachytene, *him-3^{R93Y}*

germlines had closer to an average of 30 GFP::MSH-5 foci per row in early pachytene. Moreover, the

average number of GFP::MSH-5 foci per row increased further in mid and late pachytene, achieving

peaks closer to an average of 35 foci (Figure 4D). These very high numbers of GFP::MSH-5 were

striking but do not produce an increased number of crossovers in *htp-3^{H96Y}* or *him-3^{R93Y}* single mutants

(Russo et al., 2023), raising the possibility that not all of them reflect functional, crossover-specific

intermediates, similar to what has been observed in the absence of synapsis (Woglar and Villeneuve,

2018). These data also suggest that these mutations affect the behavior of GFP::MSH-5 on meiotic

chromosomes.

Quantification of GFP::MSH-5 foci in *pch-2;htp-3*^{H93Y} double mutants was similar to *htp-3*^{H93Y} single

mutants, suggesting that PCH-2's effect on the behavior of GFP::MSH-5 foci was not through its

regulation of HTP-3 (Supplemental Figure 1). In stark contrast to *pch-2;htp-3*^{H93Y} double mutants, the

pch-2;*him-3*^{R93Y} double mutant showed a dramatic decrease in the overall average of MSH-5 foci per

row throughout the germline (Figures 4C and 4D). These averages fell between the averages in

control and *pch-2* mutant animals, particularly in the transition zone and late pachytene (Figure 4D),

consistent with our previous report that this double mutant has fewer defects in crossover formation

than either single mutant (Russo et al., 2023). These data indicate that PCH-2's effect on promoting

the removal of GFP::MSH-5 in both the transition zone and mid to late pachytene is through its

regulation of HIM-3.

pch-2 single mutants exhibit a loss of crossover assurance, as visualized by GFP::COSA-1 foci

(Figure 2B and Deshong et al., 2014). However, our quantification of GFP::MSH-5 in *pch-2* single

mutants in mid to late pachytene showed an increased number of crossover-eligible intermediates

(Figures 4A and 4B), suggesting a complex relationship between having too many crossover-eligible

intermediates and crossover assurance in *C. elegans*. To address this inconsistency, we generated

wildtype animals and *pch-2* mutants with both GFP::MSH-5 and a version of COSA-1 that has been

endogenously tagged at the N-terminus with the epitope tag, OLLAS (Janisiw et al., 2018), a fusion of

the *E. coli* OmpF protein and the mouse Langerin extracellular domain. Consistent with our and

others' previous findings, we observe a gradual reduction in the average number of GFP::MSH-5 foci

per row of nuclei in control animals as these nuclei approach the end of pachytene, ultimately

converging with the average number of OLLAS::COSA-1 foci per row (Figures 4E and 4F; Janisiw et

al., 2018; Woglar and Villeneuve, 2018). In addition, we consistently observed co-localization of

GFP::MSH-5 and OLLAS::COSA-1 foci in control animals, with a few MSH-5 foci persisting without

COSA-1, consistent with crossover designation (Figures 4E and 4F).

By contrast, we detected higher average numbers of GFP::MSH-5 foci per row across all late

pachytene nuclei in *pch-2* mutants and this average never converged upon the average number of

OLLAS::COSA-1 foci per row (Figures 4E and 4F). To test whether there was a correlation between

the number of GFP::MSH-5 foci and the loss of crossover assurance in *pch-2* mutants, we determined

the average number of GFP::MSH-5 foci in meiotic nuclei with 6 OLLAS::COSA-1 foci and those with

less than 6 OLLAS::COSA-1 foci in both control and *pch-2* mutant animals (Figures 4E and 4G,

Supplemental Figure 2). Indeed, we observed that *pch-2* mutant nuclei with less than 6

OLLAS::COSA-1 foci had significantly higher numbers of GFP::MSH-5 foci, compared to both control

and *pch-2* mutant nuclei with 6 OLLAS::COSA-1 foci (Figures 4E and 4G), suggesting that the inability

to reduce the number of crossover-eligible intermediates in *pch-2* mutants, counterintuitively, prevents

designation on some chromosomes and contributes to the observed loss of crossover assurance.

PCH-2 is removed from the synaptonemal complex when crossovers are designated

We have previously shown that PCH-2 localization to the SC is extended when there are partial

defects in synapsis or changes in karyotype, producing an increase in crossovers and a loss of

crossover interference (Deshong et al., 2014; Patel et al., 2023). These data led us to hypothesize that

PCH-2's presence on chromosomes promotes crossover formation. However, our quantification of

GFP::MSH-5 indicates that PCH-2 is required to limit the number of crossover-eligible intermediates,

in direct contrast to our proposed hypothesis (Figure 4). Therefore, we decided to revisit what role

PCH-2 localization to the SC might play in regulating crossover formation. To this end, we localized

PCH-2 in *dsb-2* mutants, in which DSB formation is substantially reduced and fewer crossovers are

formed in *C. elegans* (Rosu et al., 2013). We performed this experiment because, in budding yeast,

similar mutants that reduce DSB formation rely on Pch2 to successfully complete meiosis (Joshi et al.,

2009; Zanders and Alani, 2009).

Because the recombination defect in *dsb-2* mutants worsens with age (Rosu et al., 2013), we looked

at PCH-2 localization in both young (24 hours post L4 larval stage) and older animals (48 hours post

L4). We observed a unique and striking localization pattern in *dsb-2* mutants that we had not observed

before. When stained for PCH-2 and GFP::COSA-1, nuclei retain PCH-2 onto chromosomes far into

late pachytene, past the normal region when PCH-2 typically is removed from chromosomes (Figures

5A and 5B). However, unlike what we have observed in other mutants that have defects in synapsis or

changes in karyotype (Deshong et al., 2014; Patel et al., 2023), this retention of PCH-2 is not uniform

among all late pachytene nuclei in *dsb-2* mutants. Instead, most nuclei lose PCH-2 localization in late

pachytene while some retain it. We tested whether there was a relationship between the retention of

PCH-2 and the number of GFP::COSA-1 foci and found that 96% of nuclei that lose PCH-2 have at

least 1 GFP::COSA-1 focus (Figures 5A and 5B). In older animals (see 48 hours post-L4), this pattern

was even more clear: 77% of nuclei without PCH-2 had 1 or more GFP::COSA-1 focus and 93% of

meiotic nuclei without a GFP::COSA-1 focus retained PCH-2.

These data indicate that PCH-2's removal from the SC is in response to, coincident with or facilitates

crossover designation. To distinguish between these possibilities, we analyzed GFP::COSA-1 foci in

dsb-2::AID (Zhang et al., 2018) and *dsb-2::AID;pch-2* worms, reasoning that if PCH-2's removal is in

response to or coincident with designation, we should not detect any differences in the number of

GFP::COSA-1 foci in *dsb-2::AID* and *dsb-2::AID;pch-2* worms. We performed these experiments with

the auxin-inducible degradation system because of variability in the *dsb-2* mutant background that

affected reproducibility of our experiments.

Upon auxin treatment of *dsb-2::AID* worms, we observed a statistically significant (p value < 0.0001,

Mann-Whitney U test) decrease in the number of GFP::COSA-1 foci, in comparison with ethanol-

treated worms, verifying that we can reliably knock down DSB-2 with the AID system (Figure 5D).

When we performed the same experiment in *dsb-2::AID;pch-2* worms, the average number of

GFP::COSA-1 foci further decreased, indicating that fewer crossovers are designated when DSBs are

reduced and PCH-2 is absent (Figures 5C and 5D). These data argue against the possibility that PCH-

2's removal from the SC is simply in response to or coincident with crossover designation and instead,

suggest that PCH-2's removal from the SC somehow facilitates crossover designation and assurance.

Given the correlation between elevated GFP::MSH-5 foci and the loss of crossover assurance we

observe in *pch-2* mutants, we propose that PCH-2 is retained on meiotic chromosomes to ensure that

extra crossover-eligible intermediates are removed and crossover designation is delayed until

crossover assurance can be guaranteed in *C. elegans*.

PCH-2 and high CHK-2 activity control the fate of early double strand breaks

In *C. elegans*, meiotic cell cycle entry and progression depends on the activity of CHK-2, the meiosis-

specific ortholog of the DNA-damage kinase Chk2/CHEK2 (Baudrimont et al., 2022; Castellano-Pozo

et al., 2020; Kim et al., 2015; MacQueen and Villeneuve, 2001; Zhang et al., 2023) . Meiotic nuclei in

leptotene/zygotene are characterized by high CHK-2 activity, which drops to intermediate activity in

mid-pachytene (Kim et al., 2015; Zhang et al., 2023). In late pachytene, CHK-2 activity is inactivated

by the recruitment of polo-like kinase, PLK-2, to the SC, which enables crossover designation (Zhang

et al., 2023). Thus, CHK-2 activity also has consequences for the progression of meiotic

recombination and DNA repair.

Given that early DSBs are prevented from becoming crossovers by PCH-2 in early meiotic prophase,

we wanted to test if high CHK-2 activity in leptotene/zygotene also contributed to the fate of these

early DSBs. To evaluate this possibility, we used *syp-1^{T452A}* mutants (Sato-Carlton et al., 2018). The

SC component, SYP-1, is phosphorylated by cell cycle kinase, CDK-1, on T452, producing a Polo box

binding motif that recruits the polo-like kinase, PLK-2, to the SC, contributing to the inactivation of

CHK-2 when chromosomes are synapsed (Brandt et al., 2020; Zhang et al., 2023). Therefore, in *syp-*

1^{T452A} mutants, CHK-2 activity remains high throughout most of meiotic prophase in the *C. elegans*

germline, delaying meiotic progression as visualized by the extension of the transition zone (Figure

6A, Supplemental Figure 3).

We quantified the total number of GFP::COSA-1 foci in late pachytene nuclei in *syp-1^{T452A}* mutants

and observed a significant decrease in the average number of GFP::COSA-1 foci when compared to

control animals (Figures 6B and 6C), consistent with previous findings that *syp-1^{T452A}* mutants delay

designation and have fewer crossovers (Zhang et al., 2023). These data are also consistent with the

possibility that some DSBs fail to become crossovers when CHK-2 activity remains high. If this

hypothesis is correct, we predict that the combination of high CHK-2 activity and loss of *pch-2* should

produce more crossovers. We generated *pch-2;syp-1^{T452A}* double mutants to test this hypothesis.

These double mutants exhibited the same delay in meiotic progression as *syp-1^{T452A}* single mutants

(Supplemental Figure 3). When we quantified GFP::COSA-1 in *pch-2;syp-1^{T452A}* double mutants, we

detected a significant increase in the average number of GFP::COSA-1 foci compared to *syp-1^{T452A}*

single mutants (Figure 6C), in strong support of our hypothesis.

To further verify that the *pch-2* mutation suppresses the crossover defect in *syp-1^{T452A}* mutants, we

also monitored bivalent formation in diakinesis nuclei. *syp-1^{T452A}* single mutants exhibit an average of

7.12 DAPI staining bodies in diakinesis, higher than both control animals and *pch-2* single mutants

(Figure 6D). By contrast, *pch-2;syp-1^{T452A}* mutants had an average of 6.05 DAPI staining bodies

(Figure 6D and 6E), directly supporting our COSA-1 analysis and indicating that both PCH-2 function

and high CHK-2 activity collaborate to control the fate of DSBs and prevent some of them from

becoming crossovers in early meiotic prophase.

Discussion

We have shown that PCH-2 antagonizes crossover formation throughout meiotic prophase (Figures 2,

3, and 4) and that this regulation occurs through one of the three essential meiotic HORMADs in *C.*

elegans, HIM-3 (Figure 4). We propose that PCH-2 remodels HIM-3 on meiotic chromosomes to

destabilize crossover-eligible intermediates, visualized in our experiments as GFP-MSH-5 foci, thus

limiting which DSBs will become crossovers (Figure 7). However, this antagonism has different

consequences depending on when during meiotic prophase it occurs, underscoring the importance of

temporal regulation of these events. During leptotene/zygotene, when CHK-2 activity is high (Kim et

al., 2015; Zhang et al., 2023) and PCH-2 is present as foci on chromosomes (Deshong et al., 2014),

PCH-2 prevents crossover formation at some sites of initial DSB formation and early homolog

interactions (Figure 7B). In this way, PCH-2 promotes a wider distribution of crossovers across the

genome (Figure 1). In pachytene, when CHK-2 activity has decreased (Kim et al., 2015; Zhang et al.,

2023) and PCH-2 is localized to the SC (Deshong et al., 2014), PCH-2 winnows crossover-eligible

intermediates marked by GFP-MSH-5, ensuring their designation, colocalization with COSA-1 and

crossover assurance (Figure 7B). When there are defects in recombination, such as partial synapsis

(Deshong et al., 2014), changes in karyotype (Patel et al., 2023) or too few DSBs (Figure 5), PCH-2

persists on the SC to prevent designation, guarantee crossover assurance and some degree of

homeostasis, independent of an additional feedback mechanism that increases DSB formation (Patel

et al., 2023). That persistence of PCH-2 also disrupts crossover interference in two of these situations

(Deshong et al., 2014; Patel et al., 2023) strongly suggests that crossover interference is

mechanistically linked to assurance and homeostasis. For example, PCH-2's persistence on the

synaptonemal complex may maintain an extended period of competency for interhomolog repair, as

has been observed in mutants defective in crossover recombination (Rosu et al., 2011).

him-3^{R93Y}*;pch-2* double mutants have stronger crossover assurance than either single mutant but less

than wildtype animals (Russo et al., 2023), a phenotype which can now be explained by the behavior

of GFP::MSH-5 foci in these double mutants (Figures 4C and 4D). We have shown that HIM-3^{R93Y}

mutant protein can adopt the closed conformation and loads on meiotic chromosomes similar to

wildtype HIM-3 (Russo et al., 2023). In vitro analysis shows that HIM-3^{R93Y} binds its closure motif with

reduced affinity, likely affecting its ability to adopt the closed conformation in vivo (Figure 7A and

Russo et al., 2023). The proposed role of PCH-2 in meiosis is to remodel meiotic HORMADs from the

closed conformation to the extended one (Figure 7A), regulating their association with chromosomes

(Bhalla, 2023). However, since meiotic HORMADs are not visibly depleted from chromosomes during

meiotic progression in *C. elegans* (Couteau et al., 2004; Couteau and Zetka, 2005; Goodyer et al.,

2008; Martinez-Perez and Villeneuve, 2005), this genetic interaction supports a role for PCH-2 in

temporarily reducing the occupancy of meiotic HORMADs on meiotic chromosomes, destabilizing

interactions with partner proteins that modulate the progression and fidelity of meiotic recombination

(Russo et al., 2023). Thus, PCH-2's remodeling of HIM-3 would disrupt protein-protein interactions

that underlie crossover-eligible intermediates, destabilizing them and reducing their number on

chromosomes, contributing to crossover control (Figure 7B). Since meiotic HORMADs are essential

for meiotic chromosome axis structure and function, our data therefore support a role for the meiotic

axis in crossover control, as suggested by previous reports (Chu et al., 2024; Girard et al., 2023;

Lambing et al., 2020; Nabeshima et al., 2004). However, the disassembly of crossover-eligible

intermediates would also release pro-crossover factors present at these sites, such as MSH-5, to

concentrate at other, more stable sites on the SC, facilitating designation during synapsis (Girard et

al., 2023; Yokoo et al., 2012). In this way, PCH-2's mechanism of action may reconcile current

competing models of crossover control (Girard et al., 2023). Moreover, these results raise the

possibility that this behavior of PCH-2 and/or meiotic HORMADs might be regulated by post-

translational modifications associated with crossover control, such as ubiquitination, SUMOylation and

phosphorylation (Gray and Cohen, 2016).

Given that mutation of PCH-2 produces changes in the number and distribution of crossovers across

multiple model systems, we argue that controlling crossover distribution and number is the conserved

role of PCH-2 in meiosis (Bhalla, 2023). We have previously proposed that the conserved role of PCH-

2 is to coordinate meiotic recombination with synapsis (Bhalla, 2023). With these results, we further

refine this model. Early in meiotic prophase, PCH-2 remodels meiotic HORMADs to prevent some

DSBs from becoming crossover-eligible intermediates, widening the recombination landscape beyond

early homolog interactions and/or sites that tend to be more favorable for DSB formation, also known

as "hot spots." For example, this may explain the localization of TRIP13, the PCH-2 ortholog in mice,

to telomeres (Chotiner et al., 2024), sites that experience early homolog interactions due to the

organization of meiotic chromosomes in the bouquet formation (Scherthan et al., 1996). This

antagonism may also expand the regions of the genome that initiate synapsis in organisms that use

DSBs to accomplish this event. This possibility is supported by the observation that loss of TRIP13 in

mammals produce meiotic chromosomes that exhibit partial asynapsis (Roig et al., 2010), particularly

near regions that may act as barriers to SC polymerization (Brown et al., 2005; Roig et al., 2010).

Limiting which DSBs becomes crossover-eligible intermediates in early meiotic prophase also ensures

that meiotic recombination overlaps with synapsis, either completely, as in *C. elegans* (Yokoo et al.,

2012) or partially, as in budding yeast, plants and mice (Capilla-Perez et al., 2021; Cole et al., 2012;

Joshi et al., 2015; Morgan et al., 2021). Once synapsis is complete, PCH-2 continues to remodel

meiotic HORMADs on chromosomes to control the gradual implementation of crossover number and

distribution, reinforcing the important role that synapsis plays in mediating crossover control (Durand

et al., 2022; Libuda et al., 2013).

Unexpectedly, the inability to reduce the number of crossover-eligible intermediates in *pch-2* mutants,

as visualized by GFP::MSH-5 foci, does not produce extra crossovers but a loss of crossover-

assurance in *C. elegans* (Figure 4), a somewhat counterintuitive result. One interpretation of these

data is that crossover-eligible intermediates may be more numerous but absent from some

chromosomes in *pch-2* mutants, explaining the loss of crossover assurance. Since the absence of

crossover intermediates in *C. elegans* is accompanied by premature desynapsis of individual

chromosomes (Machovina et al., 2016; Pattabiraman et al., 2017) and chromosomes in *pch-2* mutants

delay desynapsis (Deshong et al., 2014), we do not favor this interpretation. Instead, we propose that

having too many crossover-eligible intermediates can be as deleterious to crossover assurance as

having too few (Figure 7B). This possibility is further supported by the loss of crossover assurance we

detect in irradiated wildtype worms, which is exacerbated in *pch-2* mutants (Figure 2).

This phenomenon, where crossover-eligible intermediates need to be winnowed to some threshold

number to ensure crossover assurance, may explain the loss of crossover assurance also observed in

Trip13 deficient mice (Roig et al., 2010) and on small chromosomes in budding yeast (Chakraborty et

al., 2017). Alternatively, the counterintuitive relationship between the number of crossover-eligible

precursors and crossover assurance in *pch-2* mutants we observe might reflect an additional layer of

regulation during crossover formation specific to *C. elegans*. Since *C. elegans* chromosomes are

holocentric, crossovers play an additional role organizing chromosomes for the ordered release of

sister chromatid cohesion during meiosis I (Martinez-Perez et al., 2008; Nabeshima et al., 2005) and

extra crossovers can be deleterious to accurate chromosome segregation (Hollis et al., 2020). By

contrast, in *Arabidopsis*, a system that appears to be able to tolerate an extraordinarily high number of

crossovers with little to no effect on chromosome segregation (Durand et al., 2022), PCH2's inability to

localize to the SC produces an increase in crossover formation (Figure 7), as visualized by both MLH1

foci and the formation of chiasmata (Yang et al., 2022). Once again, an overarching theme that

becomes apparent in our model is that PCH-2 may play a common role in different systems, with

dramatic variations in phenotypic consequences given species-specific requirements and constraints.

We were not surprised to see high numbers of double crossovers on almost every chromosome in our

genetic analysis of recombination in wildtype worms, given our previous analysis (Deshong et al.,

2014). However, we were surprised to see that the majority of them were found near Pairing Centers

and sites of synapsis initiation, suggesting a relationship between early homolog interactions and the

formation of double crossovers. When we revisited our previous data, we observed similar patterns. In

addition, we do not detect these double crossovers cytologically in *C. elegans*, even when using the

OLLAS::COSA-1 reporter, which has been reported to identify double crossovers in spermatogenesis

not visualized by GFP::COSA-1 (Cahoon et al., 2023). Crossovers that are cytologically marked by

COSA-1 are known as Class I crossovers, which depend on pro-crossover factors such as MSH-5 and

ZHP-3, rely on synapsis and exhibit crossover control (Gray and Cohen, 2016). These data raise the

intriguing possibility that these double crossovers are the product of the alternate, Class II, pathway of

crossover formation, which relies on a different suite of proteins, does not respond to crossover

control, do not depend on synapsis and contributes to varying degrees in different model systems

(Gray and Cohen, 2016; Youds et al., 2010). Thus, based on the close, functional relationship that

exists between Class I crossovers and synapsis and the apparent antagonistic relationship that exists

between Class II crossovers and synapsis, important corollaries of our model may be that PCH-2

specifically coordinates recombination with synapsis to promote Class I crossovers, limit Class II

crossovers and that Class II crossovers are more likely to form early in meiosis, prior to synapsis.

Therefore, variations in the contribution of the Class II crossover pathways to crossover recombination

and the degree of cross-talk between Class I and Class II pathways among model systems might

reflect the degree to which crossover formation overlaps with synapsis (Bhalla, 2023; Gray and

Cohen, 2016; Yokoo et al., 2012). Furthermore, these corollaries are entirely consistent with PCH-2's

absence from the genome of fission yeast and Tetrahymena (Kops et al., 2020; Wu and Burgess,

2006), both systems in which chromosomes do not undergo meiotic synapsis, crossovers do not

exhibit interference and all crossovers are dependent on the Class II pathway (Hollingsworth and Brill,

2004; Lukaszewicz et al., 2013; Wolfe et al., 1976). However, this aspect of the model needs to be

formally tested.

Our results also have some additional important implications about the regulation of DSB formation

across the genome in *C. elegans*. It is formally possible that PCH-2 controls DSB distribution.

However, the complexity of recombination defects in *pch-2* mutants argues against this possibility.

Instead, we explain the shift in the recombination landscape away from the central regions of

chromosomes and toward PC ends in *pch-2* mutants (Figure 1) as a result of early DSBs becoming

crossovers at the expense of later DSBs. This explanation is better supported by the panel of defects

observed in *pch-2* mutants: DSBs introduced early in meiosis become crossover eligible intermediates

and crossovers (Figures 2, 3 and 4) and while the number of DSBs are constant, DNA repair is

accelerated (Deshong et al., 2014)**.** Moreover, this explanation suggests that when DSBs happen in

meiotic prophase affects where they happen in the genome. Specifically, we propose that

chromosome arms, which are gene poor, receive DSBs early during (or even throughout) DSB

formation and the center of chromosomes, which are gene rich, receive DSBs later. The shift in

recombination to the center of chromosomes when defects in meiosis prolong DSB formation provide

further support to this possibility (Carlton et al., 2006; Deshong et al., 2014). A similar regulation of the

timing of DSB formation has been demonstrated in budding yeast, where the DSB landscape across

the whole genome expands when time in prophase is increased (Lopez Ruiz et al., 2024) and small,

highly recombinogenic, chromosomes, get more DSBs later in meiotic prophase (Murakami et al.,

2020; Subramanian et al., 2019). However, this temporal regulation has not been reported previously

in *C. elegans* and suggests that this phenomenon is more widely conserved. This expansion of the

DSB landscape in *C. elegans* to include the center regions of chromosomes later in meiosis may be a

deliberate attempt for recombination to create new haplotypes for evolution to act on, despite the

relative paucity of DSBs and the observation that they can result in chromosome missegregation

(Altendorfer et al., 2020).

Finally, our work raises some important questions about the functional role(s) of DSBs in meiosis,

aside from their contributions to crossover formation. Hicks and colleagues were the first to report that

early DSBs do not contribute to crossover formation in *C. elegans* (Hicks et al., 2022). Here we show

that these early DSBs are prevented from becoming crossovers by both PCH-2 activity and cell cycle

stage, specifically in leptotene/zygotene, when homologs are initiating pairing and synapsis. In

budding yeast, similar, early-occurring DSBs have been characterized as "scout DSBs" because of

their preference for repair from sister chromatids, versus homologous chromosomes, and their

proposed role in contributing to homolog pairing (Borde and de Massy, 2015; Joshi et al., 2015). The

homolog bias that these "scout DSBs" do display seems dependent on budding yeast *PCH2* (Joshi et

al., 2015) but interpreting this experiment is complicated by the fact that Pch2 is also required to make

the budding yeast meiotic HORMAD, Hop1, available for its loading onto meiotic chromosomes

(Herruzo et al., 2021).

In contrast to budding yeast, *C. elegans* does not rely on DSBs to promote homolog pairing and

initiate synapsis (Dernburg et al., 1998); in worms, *cis*-acting sites called Pairing Centers are essential

for homolog pairing and synapsis (MacQueen et al., 2005). There is certainly some support in the

literature for DSBs playing a role in supporting pairing and synapsis in *C. elegans* (Guo et al., 2022;

Mlynarczyk-Evans et al., 2013; Roelens et al., 2015). However, if these early DSBs were contributing

to pairing and synapsis, we would expect to see a genetic interaction between $htp\text{-}3^{H96Y}$ and *pch-2*

mutations in the installation of GFP::MSH-5 in the transition zone; we previously reported that *htp-*

3^{H96Y} suppresses the acceleration of pairing and synapsis of *pch-2* mutants, particularly when Pairing

Center function is compromised (Russo et al., 2023). Instead, we favor the possibility that these early

DSBs are generated to amplify the signaling of DNA damage kinases to prime them for their role in

recombination, a role that has also been proposed for "scout" DSBs (Joshi et al., 2015). That ATM-1, a

conserved DNA damage kinase that is downstream of CHK-2 in *C. elegans*, relies on DSBs for full

activity, supports this proposed, conserved role (Yu et al., 2023).

Our work provides an important framework to finally understand the role of PCH-2 in controlling the

number and distribution of crossovers, a role that we argue is its conserved role. While specific details

may vary across systems, we propose that PCH-2 remodels meiotic HORMADs throughout meiotic

prophase to destabilize crossover-eligible precursors, coordinating meiotic recombination with

synapsis, contributing to the progressive implementation of meiotic recombination and guaranteeing

crossover assurance, interference and homeostasis.

Methods and Materials

C.elegans **Genetics and Genome Engineering**

The *C. elegans* Bristol N2 was used as the wild-type strain. All strains were maintained at 20ºC under

standard conditions unless stated. Mutant combinations were generated by crossing. The following

mutants and rearrangements were used:

LGII: pch-2(tm1458), meIs8 ([pie-1p::GFP::cosa-1 + unc-119(+)], dsb-2(me96), dsb-

2::AID::3xFLAG])

LGIII: cosa-1(ddr12[OLLAS::cosa-1]); htp-3(vc75)

LGIV: him-3(blt9), spo-11(ie59[spo-11::AID::3xFLAG]), msh-5[ddr22(GFP::msh-5)], ieSi38 [sun-

1p::TIR1::mRuby::sun-1 3'UTR + Cbr-unc-119(+)]

LGV: syp-1(icm85[T452A]), nT1[qIs51], bcIs39 (Plin-15::ced-1::GFP)

Genetic analysis of Recombination

The wildtype Hawaiian CB4856 strain (HI) and the Bristol N2 strain were used to assay recombination

between single nucleotide polymorphisms (SNPs) on Chromosomes I, III, IV and X (Bazan and Hillers,

2011; Wicks et al., 2001). The SNPs, primers, enzymes used for restriction digests and expected

fragment sizes are included in Supplemental Table 1. To measure wild-type recombination, N2 males

containing *bcIs39* were crossed to Hawaiian CB4856 worms. Cross-progeny hermaphrodites were

identified by the presence of *bcIs39* and contained one N2 and one CB4856 chromosome. These

were assayed for recombination by crossing with CB4856 males containing *myo-2::mCherry*. Cross-

progeny hermaphrodites from the resulting cross were isolated as L4s, and then cultured individually

in 96- well plates in liquid S-media complete supplemented with HB101. Four days after initial

culturing, starved populations were lysed and used for PCR and restriction digest to detect N2 and

CB4856 SNP alleles.

For recombination in *pch-2* mutants, strains homozygous for the CB4856 background of the relevant

SNPs were created by backcrossing *pch-2* mutants to worms of the CB4856 background at least eight

times and verifying the presence of Hawaiian SNPs on all chromosomes tested in the recombination

assay. These Hawaiianized *pch-2* mutants were then mated with *pch-2; bcIs39*. Subsequent steps

were performed as in the wild-type worms.

Immunostaining

DAPI staining and immunostaining was performed as in (Russo et al., 2023), 20 to 24 hours post L4

unless otherwise noted. For analyzing bivalents, the same protocol was implemented with the

exception that hermaphrodites were dissected and DAPI stained 48 hours post late L4 stage, unless

otherwise noted.

The following primary antibodies were used at the indicated dilutions: alpaca anti-GFP Booster

(ChromoTek, gb2AF488) was used at 1:1000; rat anti-OLLAS (Invitrogen, PIMA516125) was used at

1:1000; and rabbit anti PCH-2 (Deshong et al., 2014) was used at 1:500. The following secondary

antibodies were used at the indicated dilutions: anti-rabbit Cy3 (Jackson Labs) was used at 1:500 and

anti-rat Cy5 (Jackson Labs) was used at 1:500

Irradiation Experiments

Control and *pch-2* mutant L4's were aged 12-14 hours before being exposed to 1,000 rad (10 Gy) of

X-ray radiation using a Precision MultiRad 160 X-irradiator (Precision X-Ray Inc.). Germlines were

then fixed and stained, 8 hours and 24 hours post irradiation.

Auxin-induced Degradation Experiments

Auxin treatment was performed by transferring young adult worms (aged 12-14 hours post-L4) to

bacteria-seeded plates containing auxin or 99% ethanol at specific time points, except for experiments

in which L4s were transferred directly to bacteria-seeded plates containing auxin or ethanol (the 36

hour and 48 hour time points in Figure 3C and E and the experiments in Figures 5C and D). The

natural auxin indole3-acetic acid (IAA) was purchased from Alfa Aesar (#A10556). A 400 mM stock

solution in ethanol was prepared and was stored at 4ºC for up to one month. Auxin was diluted to

100mM, and 100ul was spread onto NGM plates. Plates were allowed to dry before seeding with fresh

OP50 culture. Plates were left at 20ºC for 2-3 days in the dark to allow for bacterial lawn growth.

Imaging and Quantification

All images were acquired using a DeltaVision Personal DV system (Applied Precision) equipped with a

100X N.A. 1.40 oil-immersion objective (Olympus), resulting in an effective XY pixel spacing of 0.064

or 0.040 µm. Three-dimensional image stacks were collected at 0.2-µm Z spacing and processed by

constrained, iterative deconvolution. Image scaling, analysis and maximum-intensity projections were

performed using functions in the softWoRx software package.

For analysis of GFP::MSH-5 foci and meiotic progression, sum projections were generated using

ImageJ for each image of the germline. ImageJ plugins Cell Counter, ROI Manager, and Find Maxima

were used to identify and quantify GFP::MSH-5 foci by row from transition zone to the end of

pachytene. The threshold value was set depending on background conditions to ensure minimal

signals were identified. Foci were only quantified if they co-localized with DAPI staining. For all

genotypes, three germlines per genotype were analyzed and representative germlines are shown.

Graphing and Statistical Analysis

724 Data was analyzed using Python 3.8 and Prism for statistical significance. All datasets were tested for

725 normality using the Shapiro-Wilk test. For Figures 1, 2 and 5B, Fisher's exact test was used to

726 determine significance. For Figures 3, 4, 5D and 6, Mann-Whitney U test was used to determine

727 significance.

Author contributions

730 B.P., M.G., and N.B. designed the experiments. B.P., M.G., A.H., E.L., and V.O. performed the

731 experiments. B.P. and N.B. analyzed the data. B.P. and N.B. wrote the initial draft of the manuscript

732 and B.P., M.G., A.H. and N.B. revised the manuscript. N.B. acquired funding.

Acknowledgements

We would like to thank Josh Arribere, Pete Carlton, Abby Dernburg, Nicola Silva and Anne Villeneuve

for valuable strains and reagents. We would also like to thank the members of the Bhalla lab for

careful review of the manuscript. This work was supported by the NIH (grant numbers R35GM141835

[N.B.], R25GM051765 [V.O.] and T34GM140956 [A.H. and V.O.]). Some strains were provided by the

CGC, which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440).

References

Altendorfer, E., L.I. Lascarez-Lagunas, S. Nadarajan, I. Mathieson, and M.P. Colaiacovo. 2020.

Crossover Position Drives Chromosome Remodeling for Accurate Meiotic Chromosome

Segregation. *Curr Biol*. 30:1329-1338 e1327.

Baudrimont, A., D. Paouneskou, A. Mohammad, R. Lichtenberger, J. Blundon, Y. Kim, M. Hartl, S.

Falk, T. Schedl, and V. Jantsch. 2022. Release of CHK-2 from PPM-1.D anchorage schedules

meiotic entry. *Sci Adv*. 8:eabl8861.

Bazan, G.C., and K.J. Hillers. 2011. SNP-based mapping of crossover recombination in

Caenorhabditis elegans. *Methods Mol Biol*. 745:207-222.

Bhalla, N. 2023. PCH-2 and meiotic HORMADs: A module for evolutionary innovation in meiosis? *Curr*

Top Dev Biol. 151:317-344.

Borde, V., and B. de Massy. 2015. Meiosis: early DNA double-strand breaks pave the way for inter-

homolog repair. *Dev Cell*. 32:663-664.

Borner, G.V., A. Barot, and N. Kleckner. 2008. Yeast Pch2 promotes domainal axis organization, timely

recombination progression, and arrest of defective recombinosomes during meiosis. *Proc Natl*

Acad Sci U S A. 105:3327-3332.

Brandt, J.N., K.A. Hussey, and Y. Kim. 2020. Spatial and temporal control of targeting Polo-like kinase

during meiotic prophase. *J Cell Biol*. 219.

Brown, P.W., L. Judis, E.R. Chan, S. Schwartz, A. Seftel, A. Thomas, and T.J. Hassold. 2005. Meiotic

synapsis proceeds from a limited number of subtelomeric sites in the human male. *Am J Hum*

Genet. 77:556-566.

Cahoon, C.K., C.J. Uebel, A.M. Villeneuve, and D.E. Libuda. 2023. Epitope tag-specific differences in

the detection of COSA-1 marked crossover sites in C. elegans spermatocytes. *MicroPubl Biol*.

2023.

Capilla-Perez, L., S. Durand, A. Hurel, Q. Lian, A. Chambon, C. Taochy, V. Solier, M. Grelon, and R.

Mercier. 2021. The synaptonemal complex imposes crossover interference and heterochiasmy

in Arabidopsis. *Proc Natl Acad Sci U S A*. 118.

Carlton, P.M., A.P. Farruggio, and A.F. Dernburg. 2006. A link between meiotic prophase progression

and crossover control. *PLoS Genet*. 2:e12.

Castellano-Pozo, M., S. Pacheco, G. Sioutas, A.L. Jaso-Tamame, M.H. Dore, M.M. Karimi, and E.

Martinez-Perez. 2020. Surveillance of cohesin-supported chromosome structure controls

meiotic progression. *Nat Commun*. 11:4345.

Chakraborty, P., A.V. Pankajam, G. Lin, A. Dutta, G.N. Krishnaprasad, M.M. Tekkedil, A. Shinohara,

L.M. Steinmetz, and K.T. Nishant. 2017. Modulating Crossover Frequency and Interference for

Obligate Crossovers in Saccharomyces cerevisiae Meiosis. *G3 (Bethesda)*. 7:1511-1524.

Chotiner, J.Y., N.A. Leu, F. Yang, I.G. Cossu, Y. Guan, H. Lin, and P.J. Wang. 2024. TRIP13 localizes

to synapsed chromosomes and functions as a dosage-sensitive regulator of meiosis. *Elife*. 12.

Chu, L., J. Zhuang, M. Geng, Y. Zhang, J. Zhu, C. Zhang, A. Schnittger, B. Yi, and C. Yang. 2024.

ASYNAPSIS3 has diverse dosage-dependent effects on meiotic crossover formation in

Brassica napus. *Plant Cell*. 36:3838-3856.

Cole, F., L. Kauppi, J. Lange, I. Roig, R. Wang, S. Keeney, and M. Jasin. 2012. Homeostatic control of

recombination is implemented progressively in mouse meiosis. *Nat Cell Biol*. 14:424-430.

Couteau, F., K. Nabeshima, A. Villeneuve, and M. Zetka. 2004. A component of C. elegans meiotic

chromosome axes at the interface of homolog alignment, synapsis, nuclear reorganization,

and recombination. *Curr Biol*. 14:585-592.

Couteau, F., and M. Zetka. 2005. HTP-1 coordinates synaptonemal complex assembly with homolog

alignment during meiosis in C. elegans. *Genes Dev*. 19:2744-2756.

Cuacos, M., C. Lambing, M. Pachon-Penalba, K. Osman, S.J. Armstrong, I.R. Henderson, E.

Sanchez-Moran, F.C.H. Franklin, and S. Heckmann. 2021. Meiotic chromosome axis

remodelling is critical for meiotic recombination in Brassica rapa. *J Exp Bot*. 72:3012-3027.

Dernburg, A.F., K. McDonald, G. Moulder, R. Barstead, M. Dresser, and A.M. Villeneuve. 1998. Meiotic

recombination in C. elegans initiates by a conserved mechanism and is dispensable for

homologous chromosome synapsis. *Cell*. 94:387-398.

Deshong, A.J., A.L. Ye, P. Lamelza, and N. Bhalla. 2014. A quality control mechanism coordinates

795   meiotic prophase events to promote crossover assurance. *PLoS Genet*. 10:e1004291.

796   Durand, S., Q. Lian, J. Jing, M. Ernst, M. Grelon, D. Zwicker, and R. Mercier. 2022. Joint control of

797   meiotic crossover patterning by the synaptonemal complex and HEI10 dosage. *Nat Commun*.

798   13:5999.

799   Girard, C., D. Zwicker, and R. Mercier. 2023. The regulation of meiotic crossover distribution: a coarse

800   solution to a century-old mystery? *Biochem Soc Trans*. 51:1179-1190.

801   Goodyer, W., S. Kaitna, F. Couteau, J.D. Ward, S.J. Boulton, and M. Zetka. 2008. HTP-3 links DSB

802   formation with homolog pairing and crossing over during C. elegans meiosis. *Dev Cell*. 14:263-

803   274.

804   Gray, S., and P.E. Cohen. 2016. Control of Meiotic Crossovers: From Double-Strand Break Formation

805   to Designation. *Annu Rev Genet*. 50:175-210.

806   Gu, Y., A. Desai, and K.D. Corbett. 2022. Evolutionary Dynamics and Molecular Mechanisms of

807   HORMA Domain Protein Signaling. *Annu Rev Biochem*. 91:541-569.

808   Guo, H., E.L. Stamper, A. Sato-Carlton, M.A. Shimazoe, X. Li, L. Zhang, L. Stevens, K.C.J. Tam, A.F.

809   Dernburg, and P.M. Carlton. 2022. Phosphoregulation of DSB-1 mediates control of meiotic

810   double-strand break activity. *Elife*. 11.

811   Hassold, T., and P. Hunt. 2001. To err (meiotically) is human: the genesis of human aneuploidy. *Nat*

812   *Rev Genet*. 2:280-291.

813   Haversat, J., A. Woglar, K. Klatt, C.C. Akerib, V. Roberts, S.Y. Chen, S. Arur, A.M. Villeneuve, and Y.

814   Kim. 2022. Robust designation of meiotic crossover sites by CDK-2 through phosphorylation of

815   the MutSgamma complex. *Proc Natl Acad Sci U S A*. 119:e2117865119.

816   Herruzo, E., A. Lago-Maciel, S. Baztan, B. Santos, J.A. Carballo, and P.A. San-Segundo. 2021. Pch2

817   orchestrates the meiotic recombination checkpoint from the cytoplasm. *PLoS Genet*.

818   17:e1009560.

Hicks, T., S. Trivedi, M. Eppert, R. Bowman, H. Tian, A. Dafalla, C. Crahan, S. Smolikove, and N.

Silva. 2022. Continuous double-strand break induction and their differential processing sustain

chiasma formation during Caenorhabditis elegans meiosis. *Cell Rep*. 40:111403.

Hollingsworth, N.M., and S.J. Brill. 2004. The Mus81 solution to resolution: generating meiotic

crossovers without Holliday junctions. *Genes Dev*. 18:117-125.

Hollis, J.A., M.L. Glover, A.J. Schlientz, C.K. Cahoon, B. Bowerman, S.M. Wignall, and D.E. Libuda.

2020. Excess crossovers impede faithful meiotic chromosome segregation in C. elegans.

PLoS Genet. 16:e1009001.

Janisiw, E., M.R. Dello Stritto, V. Jantsch, and N. Silva. 2018. BRCA1-BARD1 associate with the

synaptonemal complex and pro-crossover factors and influence RAD-51 dynamics during

Caenorhabditis elegans meiosis. *PLoS Genet*. 14:e1007653.

Jaramillo-Lambert, A., M. Ellefson, A.M. Villeneuve, and J. Engebrecht. 2007. Differential timing of S

phases, X chromosome replication, and meiotic prophase in the C. elegans germ line. *Dev*

Biol. 308:206-221.

Joshi, N., A. Barot, C. Jamison, and G.V. Borner. 2009. Pch2 links chromosome axis remodeling at

future crossover sites and crossover distribution during yeast meiosis. *PLoS Genet*.

5:e1000557.

Joshi, N., M.S. Brown, D.K. Bishop, and G.V. Borner. 2015. Gradual implementation of the meiotic

recombination program via checkpoint pathways controlled by global DSB levels. *Mol Cell*.

57:797-811.

Joyce, E.F., and K.S. McKim. 2009. Drosophila PCH2 is required for a pachytene checkpoint that

monitors double-strand-break-independent events leading to meiotic crossover formation.

Genetics. 181:39-51.

Keeney, S., C.N. Giroux, and N. Kleckner. 1997. Meiosis-specific DNA double-strand breaks are

catalyzed by Spo11, a member of a widely conserved protein family. *Cell*. 88:375-384.

Kelly, K.O., A.F. Dernburg, G.M. Stanfield, and A.M. Villeneuve. 2000. Caenorhabditis elegans msh-5

is required for both normal and radiation-induced meiotic crossing over but not for completion

of meiosis. *Genetics*. 156:617-630.

Kim, Y., N. Kostow, and A.F. Dernburg. 2015. The Chromosome Axis Mediates Feedback Control of

CHK-2 to Ensure Crossover Formation in C. elegans. *Dev Cell*. 35:247-261.

Kim, Y., S.C. Rosenberg, C.L. Kugel, N. Kostow, O. Rog, V. Davydov, T.Y. Su, A.F. Dernburg, and K.D.

Corbett. 2014. The chromosome axis controls meiotic events through a hierarchical assembly

of HORMA domain proteins. *Dev Cell*. 31:487-502.

Kops, G., B. Snel, and E.C. Tromer. 2020. Evolutionary Dynamics of the Spindle Assembly Checkpoint

in Eukaryotes. *Curr Biol*. 30:R589-R602.

Lambing, C., P.C. Kuo, A.J. Tock, S.D. Topp, and I.R. Henderson. 2020. ASY1 acts as a dosage-

dependent antagonist of telomere-led recombination and mediates crossover interference in

Arabidopsis. *Proc Natl Acad Sci U S A*. 117:13647-13658.

Lambing, C., K. Osman, K. Nuntasoontorn, A. West, J.D. Higgins, G.P. Copenhaver, J. Yang, S.J.

Armstrong, K. Mechtler, E. Roitinger, and F.C. Franklin. 2015. Arabidopsis PCH2 Mediates

Meiotic Chromosome Remodeling and Maturation of Crossovers. *PLoS Genet*. 11:e1005372.

Libuda, D.E., S. Uzawa, B.J. Meyer, and A.M. Villeneuve. 2013. Meiotic chromosome structures

constrain and respond to designation of crossover sites. *Nature*. 502:703-706.

Lopez Ruiz, L.M., D. Johnson, W.H. Gittens, G.G.B. Brown, R.M. Allison, and M.J. Neale. 2024.

Meiotic prophase length modulates Tel1-dependent DNA double-strand break interference.

PLoS Genet. 20:e1011140.

Lukaszewicz, A., R.A. Howard-Till, and J. Loidl. 2013. Mus81 nuclease and Sgs1 helicase are

essential for meiotic recombination in a protist lacking a synaptonemal complex. *Nucleic Acids*

Res. 41:9296-9309.

Machovina, T.S., R. Mainpal, A. Daryabeigi, O. McGovern, D. Paouneskou, S. Labella, M. Zetka, V.

Jantsch, and J.L. Yanowitz. 2016. A Surveillance System Ensures Crossover Formation in C.

elegans. *Curr Biol*. 26:2873-2884.

MacQueen, A.J., C.M. Phillips, N. Bhalla, P. Weiser, A.M. Villeneuve, and A.F. Dernburg. 2005.

Chromosome sites play dual roles to establish homologous synapsis during meiosis in C.

elegans. *Cell*. 123:1037-1050.

MacQueen, A.J., and A.M. Villeneuve. 2001. Nuclear reorganization and homologous chromosome

pairing during meiotic prophase require C. elegans chk-2. *Genes Dev*. 15:1674-1687.

Martinez-Perez, E., M. Schvarzstein, C. Barroso, J. Lightfoot, A.F. Dernburg, and A.M. Villeneuve.

2008. Crossovers trigger a remodeling of meiotic chromosome axis composition that is linked

to two-step loss of sister chromatid cohesion. *Genes Dev*. 22:2886-2901.

Martinez-Perez, E., and A.M. Villeneuve. 2005. HTP-1-dependent constraints coordinate homolog

pairing and synapsis and promote chiasma formation during C. elegans meiosis. *Genes Dev*.

19:2727-2743.

Mlynarczyk-Evans, S., B. Roelens, and A.M. Villeneuve. 2013. Evidence that masking of synapsis

imperfections counterbalances quality control to promote efficient meiosis. *PLoS Genet*.

9:e1003963.

Morgan, C., J.A. Fozard, M. Hartley, I.R. Henderson, K. Bomblies, and M. Howard. 2021. Diffusion-

mediated HEI10 coarsening can explain meiotic crossover positioning in Arabidopsis. *Nat

Commun*. 12:4674.

Murakami, H., I. Lam, P.C. Huang, J. Song, M. van Overbeek, and S. Keeney. 2020. Multilayered

mechanisms ensure that short chromosomes recombine in meiosis. *Nature*. 582:124-128.

Nabeshima, K., A.M. Villeneuve, and M.P. Colaiacovo. 2005. Crossing over is coupled to late meiotic

prophase bivalent differentiation through asymmetric disassembly of the SC. *J Cell Biol*.

168:683-689.

Nabeshima, K., A.M. Villeneuve, and K.J. Hillers. 2004. Chromosome-wide regulation of meiotic

crossover formation in Caenorhabditis elegans requires properly assembled chromosome

axes. *Genetics*. 168:1275-1292.

Nadarajan, S., E. Altendorfer, T.T. Saito, M. Martinez-Garcia, and M.P. Colaiacovo. 2021. HIM-17

regulates the position of recombination events and GSP-1/2 localization to establish short arm

identity on bivalents in meiosis. *Proc Natl Acad Sci U S A*. 118.

Patel, B., M. Grobler, and N. Bhalla. 2023. Chromosomal fusions, but not chromosomal inversions,

activate a PCH-2 dependent checkpoint that promotes crossover formation in C. elegans.

MicroPubl Biol. 2023.

Pattabiraman, D., B. Roelens, A. Woglar, and A.M. Villeneuve. 2017. Meiotic recombination modulates

the structure and dynamics of the synaptonemal complex during C. elegans meiosis. *PLoS*

Genet. 13:e1006670.

Roelens, B., M. Schvarzstein, and A.M. Villeneuve. 2015. Manipulation of Karyotype in Caenorhabditis

elegans Reveals Multiple Inputs Driving Pairwise Chromosome Synapsis During Meiosis.

Genetics. 201:1363-1379.

Roig, I., J.A. Dowdle, A. Toth, D.G. de Rooij, M. Jasin, and S. Keeney. 2010. Mouse TRIP13/PCH2 is

required for recombination and normal higher-order chromosome structure during meiosis.

PLoS Genet. 6.

Rosu, S., D.E. Libuda, and A.M. Villeneuve. 2011. Robust crossover assurance and regulated

interhomolog access maintain meiotic crossover number. *Science*. 334:1286-1289.

Rosu, S., K.A. Zawadzki, E.L. Stamper, D.E. Libuda, A.L. Reese, A.F. Dernburg, and A.M. Villeneuve.

2013. The C. elegans DSB-2 protein reveals a regulatory network that controls competence for

meiotic DSB formation and promotes crossover assurance. *PLoS Genet*. 9:e1003674.

Russo, A.E., S. Giacopazzi, A. Deshong, M. Menon, V. Ortiz, K.M. Ego, K.D. Corbett, and N. Bhalla.

2023. The conserved AAA ATPase PCH-2 distributes its regulation of meiotic prophase events

through multiple meiotic HORMADs in C. elegans. *PLoS Genet*. 19:e1010708.

Sato-Carlton, A., C. Nakamura-Tabuchi, S.K. Chartrand, T. Uchino, and P.M. Carlton. 2018.

Phosphorylation of the synaptonemal complex protein SYP-1 promotes meiotic chromosome

segregation. *J Cell Biol*. 217:555-570.

Scherthan, H., S. Weich, H. Schwegler, C. Heyting, M. Harle, and T. Cremer. 1996. Centromere and

telomere movements during early meiotic prophase of mouse and man are associated with the

onset of chromosome pairing. *J Cell Biol*. 134:1109-1125.

Snowden, T., S. Acharya, C. Butz, M. Berardini, and R. Fishel. 2004. hMSH4-hMSH5 recognizes

Holliday Junctions and forms a meiosis-specific sliding clamp that embraces homologous

chromosomes. *Mol Cell*. 15:437-451.

Subramanian, V.V., X. Zhu, T.E. Markowitz, L.A. Vale-Silva, P.A. San-Segundo, N.M. Hollingsworth, S.

Keeney, and A. Hochwagen. 2019. Persistent DNA-break potential near telomeres increases

initiation of meiotic recombination on short chromosomes. *Nat Commun*. 10:970.

Veller, C., N. Kleckner, and M.A. Nowak. 2019. A rigorous measure of genome-wide genetic shuffling

that takes into account crossover positions and Mendel's second law. *Proc Natl Acad Sci U S*

A. 116:1659-1668.

Wicks, S.R., R.T. Yeh, W.R. Gish, R.H. Waterston, and R.H. Plasterk. 2001. Rapid gene mapping in

Caenorhabditis elegans using a high density polymorphism map. *Nat Genet*. 28:160-164.

Woglar, A., and A.M. Villeneuve. 2018. Dynamic Architecture of DNA Repair Complexes and the

Synaptonemal Complex at Sites of Meiotic Recombination. *Cell*. 173:1678-1691 e1616.

Wojtasz, L., K. Daniel, I. Roig, E. Bolcun-Filas, H. Xu, V. Boonsanay, C.R. Eckmann, H.J. Cooke, M.

Jasin, S. Keeney, M.J. McKay, and A. Toth. 2009. Mouse HORMAD1 and HORMAD2, two

conserved meiotic chromosomal proteins, are depleted from synapsed chromosome axes with

the help of TRIP13 AAA-ATPase. *PLoS Genet*. 5:e1000702.

Wolfe, J., B. Hunter, and W.S. Adair. 1976. A cytological study of micronuclear elongation during

conjugation in Tetrahymena. *Chromosoma*. 55:289-308.

Wu, H.Y., and S.M. Burgess. 2006. Two distinct surveillance mechanisms monitor meiotic

chromosome metabolism in budding yeast. *Curr Biol*. 16:2473-2479.

Yang, C., B. Hu, S.M. Portheine, P. Chuenban, and A. Schnittger. 2020. State changes of the HORMA

protein ASY1 are mediated by an interplay between its closure motif and PCH2. *Nucleic Acids*

Res. 48:11521-11535.

Yang, C., K. Sofroni, Y. Hamamura, B. Hu, H.T. Elbasi, M. Balboni, L. Chu, D. Stang, M. Heese, and A.

Schnittger. 2022. ZYP1-mediated recruitment of PCH2 to the synaptonemal complex remodels

the chromosome axis leading to crossover restriction. *Nucleic Acids Res*. 50:12924-12937.

Yokoo, R., K.A. Zawadzki, K. Nabeshima, M. Drake, S. Arur, and A.M. Villeneuve. 2012. COSA-1

reveals robust homeostasis and separable licensing and reinforcement steps governing

meiotic crossovers. *Cell*. 149:75-87.

Youds, J.L., D.G. Mets, M.J. McIlwraith, J.S. Martin, J.D. Ward, O.N. NJ, A.M. Rose, S.C. West, B.J.

Meyer, and S.J. Boulton. 2010. RTEL-1 enforces meiotic crossover interference and

homeostasis. *Science*. 327:1254-1258.

Yu, Z., H.J. Kim, and A.F. Dernburg. 2023. ATM signaling modulates cohesin behavior in meiotic

prophase and proliferating cells. *Nat Struct Mol Biol*. 30:436-450.

Yu, Z., Y. Kim, and A.F. Dernburg. 2016. Meiotic recombination and the crossover assurance

checkpoint in Caenorhabditis elegans. *Semin Cell Dev Biol*. 54:106-116.

Zanders, S., and E. Alani. 2009. The pch2Delta mutation in baker's yeast alters meiotic crossover

levels and confers a defect in crossover interference. *PLoS Genet*. 5:e1000571.

Zhang, L., W. Stauffer, D. Zwicker, and A.F. Dernburg. 2021. Crossover patterning through kinase-

regulated condensation and coarsening of recombination nodules. *bioRxiv*.

Zhang, L., W.T. Stauffer, J.S. Wang, F. Wu, Z. Yu, C. Liu, H.J. Kim, and A.F. Dernburg. 2023.

Recruitment of Polo-like kinase couples synapsis to meiotic progression via inactivation of

CHK-2. *Elife*. 12.

 Figure Legends

 Figure 1. PCH-2 controls the number and distribution of crossovers in similar patterns on

 multiple chromosomes. Genetic analysis of meiotic recombination in wildtype and *pch-2* mutants.

 DCO indicates double crossovers. Physical and genetic maps of Chromosome I, III, IV and the X

 chromosome are depicted to scale. Genetic distance is shown in centimorgans. A * indicates a p-value

 < 0.05, a ** indicates a p value < 0.01 and a *** indicates p-value < 0.001.

 Figure 2. PCH-2 prevents exogenous double strand breaks from becoming crossovers in early

 meiotic prophase. A. Illustration of the irradiation experiments in control and *pch-2* mutants. Box

 indicates late pachytene, the area where GFP::COSA-1 foci are analyzed. **B.** Fraction of meiotic

 nuclei with less than 6, 6, or greater than 6 GFP::COSA-1 foci in control animals (yellow, n = 446) and

 pch-2 mutants (blue, n = 552). **C.** Meiotic nuclei in control animals and *pch-2* mutants 8 hours post

 irradiation stained for DAPI (magenta) and GFP::COSA-1 (green). Scale bar is 4 um. **D.** Fraction of

 meiotic nuclei with less than 6, 6, or greater than 6 GFP::COSA-1 foci in control animals (yellow, n =

 143) and *pch-2* mutants (blue, n = 125) 8 hours post irradiation. **E.** Meiotic nuclei in control animals

 and *pch-2* mutants 24 hours post irradiation with DAPI (magenta) and GFP::COSA-1 (green). **F.**

 Fraction of meiotic nuclei with less than 6, 6, or greater than 6 GFP::COSA-1 foci in control animals (n

 = 179) and *pch-2* mutants (n = 378) 24 hours post irradiation. A *** indicates p-value < 0.001, and a

 **** indicates p-value < 0.0001.

 Figure 3. PCH-2 prevents SPO-11-induced double strand breaks from becoming crossovers in

 early meiotic prophase. A. Illustration of the SPO-11 depletion experiment to assay GFP::COSA-1 in

 control animals and *pch-2* mutants at different timepoints of auxin treatment. Each timepoint indicates

 when SPO-11 is depleted in the germline with auxin induced degradation. Box indicates late

 pachytene, the area where GFP::COSA-1 foci are analyzed. **B.** Representative images of meiotic

nuclei in control animals and *pch-2* mutants treated with auxin for 20 hours, stained for DAPI

(magenta) and GFP::COSA-1 (green). Scale bar is 5 um. **C.** Number of GFP::COSA-1 foci in meiotic

nuclei at different timepoints of auxin treatment in control (blue) and *pch-2* mutants (yellow). Error bars

represent standard error of the mean (SEM). N values are as follows: 36 hours on auxin, control (78

nuclei) and *pch-2* (83 nuclei); 24 hours on auxin, control (132 nuclei) and *pch-2* (155 nuclei); 22 hours

on auxin, control (152 nuclei) and *pch-2* (168 nuclei); 20 hours on auxin, control (139 nuclei) and *pch-*

2 (157 nuclei); 18 hours on auxin, control (154 nuclei) and *pch-2* (154 nuclei); and 24 hours on

ethanol, control (86 nuclei) and *pch-2* (143 nuclei). **D.** Illustration of the SPO-11 depletion experiment

to assay bivalents in control animals and *pch-2* mutants at different timepoints of auxin treatment.

Each timepoint indicates when SPO-11 is depleted in the germline with auxin induced degradation.

Box indicates diakinesis, where DAPI stained bodies are analyzed. **E.** Oocytes from control animals

and *pch-2* mutants stained for DAPI (magenta). Scale bar is 4um. **F.** Number of DAPI stained bodies

in meiotic nuclei at different timepoints of auxin treatment in control animals and *pch-2* mutants. N

values are as follows: 48 hours on auxin, control (47 nuclei) and *pch-2* (43 nuclei); 36 hours on auxin,

control (46 nuclei) and *pch-2* (50 nuclei); 34 hours on auxin, control (41 nuclei) and *pch-2* (41 nuclei);

32 hours on auxin, control (51 nuclei) and *pch-2* (47 nuclei); 30 hours on auxin, control (46 nuclei) and

pch-2 (21 nuclei); and 36 hours on ethanol, control (52 nuclei) and *pch-2* (48 nuclei). Error bars

represent SEM and a * indicates p-value < 0.05 and a *** indicates p-value < 0.001.

1015

Figure 4. PCH-2 is required for timely loading and removal of MSH-5 on meiotic chromosomes

through its regulation of HIM-3. A. Representative images of nuclei in different stages of meiotic

prophase in control animals and *pch-2* mutants stained for DAPI (magenta) and GFP::MSH-5 (green)

Scale bar is 5 um. **B.** Scatter plot showing average GFP::MSH-5 foci per row of germline nuclei in

control animals (yellow, 163 nuclei) and *pch-2* mutants (blue, 195 nuclei) from the transition zone (TZ)

to late pachytene (LP), normalized to 100. The line represents a rolling average of four rows. **C.**

Representative images of nuclei in different stages of meiotic prophase in *him-3*[R93Y] mutants (left) and

pch-2;*him-3*[R93Y] double mutants (right), stained for DAPI (magenta) and GFP::MSH-5 (green). **D.**

Scatter plot showing average GFP::MSH-5 foci per row in *him-3^{R93Y}* (brown, 183 nuclei) and *pch-*

2;him-3^{R93Y} mutants (pink, 163 nuclei) from the transition zone (TZ) to late pachytene (LP), normalized

to 100. The line represents a rolling average of four rows. Similar data is provided for a control

germline (opaque yellow, 236 nuclei) for comparison. **E.** Representative images of meiotic nuclei in

control animals and *pch-2* mutants stained for DAPI (blue), GFP::MSH-5 (green), and OLLAS::COSA-

1 (red). Yellow circles indicate GFP::MSH-5 without OLLAS::COSA-1. Scale bar is 4um. **F.** Scatter plot

showing average GFP::MSH-5 (green) and OLLAS::COSA-1 (red) foci per row in the last five rows of

the germline in control animals (36 nuclei) and *pch-2* mutants (45 nuclei). The line represents a rolling

average of 2 rows. **G.** Swarm plot showing number of GFP::MSH-5 foci in control (yellow, 14 nuclei)

and *pch-2* mutant (blue, 29 nuclei) nuclei with less than 6 OLLAS::COSA-1 foci (left) and 6

OLLAS::COSA-1 foci (right). Error bars represent SEM. A * indicates a p-value less than 0.05 and a **

indicates a p-value < 0.01.

1036

Figure 5. PCH-2 is removed when crossovers are designated. A. Representative images of

meiotic nuclei in *dsb-2* animals 24 hours post L4 and 48 hours post L4 stained for DAPI (magenta),

PCH-2 (red), and GFP::COSA-1 (green). Scale bar is 4 um. **B.** Stacked histograms showing

percentage of PCH-2 positive (n = 43 at 24 hours, n = 42 at 48 hours) and negative (n = 194 at 24

hours, n = 130 at 48 hours) nuclei with (lime) and without (dark green) GFP::COSA-1 foci in *dsb-2*

mutants at 24 hours post L4 and 48 hours post L4. **C.** Representative images of meiotic nuclei in *dsb-*

2::AID and *dsb-2::AID;pch-2* mutants treated with auxin and stained for DAPI (magenta) and

GFP::COSA-1 (green). Scale bar is 5 um. **D.** Swarm plot showing the number of GFP::COSA-1 foci in

dsb-2::AID (gray) and *dsb-2::AID;pch-2* (lemon) mutants when treated with ethanol or auxin. N values

are as follows: *dsb-2::AID* (101 nuclei) and *dsb-2::AID;pch-2* (154 nuclei) on ethanol, *dsb-2::AID* (136

nuclei) and *dsb-2::AID;pch-2* (131 nuclei) on auxin. Error bars represent the SEM. A ** indicates a p-

value less than 0.01 and a **** indicates p-value < 0.0001.

1049

Figure 6. PCH-2 and high CHK-2 activity control the fate of early double strand breaks. A. Illustration of CHK-2 activity in wildtype and $syp-1^{T452A}$ germlines. **B.** Representative images of meiotic nuclei late pachytene in $syp-1^{T452A}$ and $pch-2;syp-1^{T452A}$ mutants stained for DAPI (magenta) and GFP::COSA-1 (green). Scale bar is 5 um. **C.** Swarm plot showing number of GFP::COSA-1 foci in control animals (blue), $pch-2$ (yellow), $syp-1^{T452A}$ (maroon), and $pch-2;syp-1^{T452A}$ (light blue) mutants. Error bars represent SEM. **D.** Oocytes from $syp-1^{T452A}$ and $pch-2;syp-1^{T452A}$ mutant worms stained for DAPI (magenta). Scale bar is 4um. **D.** Swarm plot showing number of DAPI stained bodies in control animals (blue, n = 154), $pch-2$ (yellow, n = 89), $syp-1^{T452A}$ (maroon, n = 247)), and $pch-2;syp-1^{T452A}$ (light blue, n = 242) mutants. Error bars represent SEM. A **** indicates p-value < 0.0001.

Figure 7. PCH-2 remodels HIM-3 to disassemble crossover-eligible intermediates, controlling crossover distribution and number. A. Model for how $pch-2$ and $him-3^{R93Y}$ mutations genetically interact to affect the progression of meiotic recombination. HIM-3 adopts the closed conformation upon binding an interacting protein with a closure motif and its conversion to the extended conformation is facilitated by PCH-2's remodeling of its HORMA domain. **B.** Model for how PCH-2 and HIM-3 progressively implement meiotic recombination during different stages of meiotic prophase.

Supplemental Figure 1: PCH-2 does not regulate GFP::MSH-5 loading and removal through HTP-3. A. Representative images of nuclei in different stages of meiotic prophase in $htp-3^{H96Y}$ and $pch-2; htp-3^{H96Y}$ mutants stained for DAPI (magenta) and GFP::MSH-5 (green) Scale bar in all images is 5 um. **B.** Scatter plot showing average GFP::MSH-5 foci per row of germline nuclei in $htp-3^{H96Y}$ (blue, 132 nuclei) and $pch-2; htp-3^{H96Y}$ (green, 161 nuclei) mutants from the transition zone to late pachytene, normalized to 100. The line represents a rolling average of four rows. Similar data is provided for a control germline (opaque yellow, 163 nuclei) for comparison

1075 **Supplemental Figure 2: *pch-2* meiotic nuclei with elevated numbers of GFP::MSH-5 foci show**

1076 **defects in crossover assurance.** Gray scale images of control and *pch-2* mutant nuclei stained for

1077 DAPI, GFP::MSH-5 and OLLAS::COSA-1. Scale bar in image is 4 um.

1078

1079 **Supplemental Figure 3: *syp-1^{T452A}* and *pch-2;syp-1^{T452A}* mutants display a similar defect in**

1080 **meiotic progression. A.** Representative images of *syp-1^{T452A}* and *pch-2;syp-1^{T452A}* mutant germlines.

1081 Encircled nuclei indicate transition zone nuclei. Scale bar indicates 10 um. **B.** Quantification of fraction

1082 of rows of transition zone nuclei in 3 germlines in *syp-1^{T452A}* (maroon) and *pch-2;syp-1^{T452A}* (light blue)

1083 mutants.

1084

1085

1086

Figure 1

A

24 Hours

8 Hours

Leptotene/Zygotene   Mid-Pachytene   Late-Pachytene

B

Unirradiated

control
pch-2

****

Fraction of Meiotic Nuclei

Number of COSA-1 Foci

<6   6   >6

C   8 Hours Post Irradiation

control

pch-2

DAPI
COSA-1

D   8 Hours Post Irradiation

***

Fraction of Meiotic Nuclei

Number of COSA-1 Foci

<6   6   >6

E   24 Hours Post Irradiation

control

pch-2

F   24 Hours Post Irradiation

****   ****

Fraction of Meiotic Nuclei

Number of COSA-1 Foci

<6   6   >6

A Areas of SPO-11 depletion to assay COSA-1 foci

Time Points
- 36 hours
- 24 hours
- 22 hours
- 20 hours
- 18 hours

B
control 20 hr. Auxin
6
6
6
pch-2
7
7
DAPI
COSA-1

C
Number of COSA-1 Foci

control / *pch-2*

36 Hours Auxin | 24 Hours Auxin | 22 Hours Auxin | 20 Hours Auxin | 18 Hours Auxin | 24 Hours Ethanol

D Areas of SPO-11 depletion to assay bivalents

Time Points
- 48 hours
- 36 hours
- 34 hours
- 32 hours
- 30 hours

E
control 32 hr. Auxin
11
pch-2
6
DAPI

F
Number of DAPI Stained Bodies

*

***

control / *pch-2*

48 Hours Auxin | 36 Hours Auxin | 34 Hours Auxin | 32 Hours Auxin | 30 Hours Auxin | 36 Hours Ethanol

A

dsb-2 24 hr.

3
0
2

dsb-2 48 hr.

0
1
1

DAPI
PCH-2
COSA-1

B

24 Hours Post L4   **48 Hours Post L4**

dsb-2   *dsb-2*

████ 0 COSA-1 Foci
████ ≥1 COSA-1 Focus

****   ****

Fraction of Meiotic Nuclei

PCH-2 Positive Nuclei   PCH-2 Negative Nuclei   PCH-2 Positive Nuclei   PCH-2 Negative Nuclei

C

dsb-2::AID + Auxin.

6
2
5

dsb-2::AID;pch-2 + Auxin

3
2
2

DAPI
COSA-1

D

Ethanol Treatment   **Auxin Treatment**

Number of COSA-1 Foci

**

dsb-2::AID   *dsb-2::AID; pch-2*   *dsb-2::AID*   *dsb-2::AID; pch-2*

A

wildtype:

closure motif

PCH-2

extended HIM-3 ⟷ closed HIM-3

pch-2 mutants:

closure motif

extended HIM-3 ⟶ closed HIM-3

him-3^{R93Y} mutants:

closure motif

PCH-2

extended HIM-3 ⟷ closed HIM-3

pch-2;him-3^{R93Y} mutants:

closure motif

extended HIM-3 ⟷ closed HIM-3

B

Figure 7

Leptotene/Zygotene:
high CHK-2 activity
PCH-2 present as foci
on meiotic chromosomes

HIM-3

PCH-2

Mid-pachytene:
intermediate CHK-2 activity
PCH-2 localizes to
the synaptonemal complex

HIM-3

PCH-2

PCH-2

no PCH-2

Late-pachytene:
no CHK-2 activity
PCH-2 absent
crossover designation

no
crossover

⭐ Double strand break

⭐ Crossover-eligible intermediate

● Crossover

Supplementary Table 1: Single nucleotide polymorphisms used for recombination assay

Chromosome and SNP	Primer Name	Primer Genetic Location	Primer Sequence FOR	Primer Sequence REV	Restriction enzyme	N2	HI
IA	F56C11	-19	ATGCCAGTGATAAGGAACGG	TCACATCCCTTGTCGATGAA	DraI	354, 146	500
IB	Y71G12	−12.3	GACAATGACCAATAAGACG	GATCCGTGAAATTGTTCCG	BsrI	440, 125	364, 125, 76
IC	K04F10	0.9	ATCATTCTCCAGGCCACGTTAC	CTGAACTAGTCGAACAAACCCC	NdeI	594	300, 294
ID	T07D10	13.6	CTTGGTGTGGGGAGAGTATAGG	TTTGTCCGGATTGACTCTGC	Sau3AI	303, 63	207, 96, 63
IE	ZK909	28.8	CACAAGTGGTTTGGAAGTACCG	CAACAAAGGGATAGATCACGGG	HindIII	450	236, 214
IIIA	pkP3081	-26.98	AGCAAGAATGAGCCGATTG	GTCGGCCGTTTTCAAATAACTG	TaqI	222, 149	195, 145, 27
IIIB	pkP3095	-5.12	TCTCGTCAATTGTCGCCTG	TTATTTGCAATCCAACGGC	ApoI	308	168, 140
IIIC	pkP3101	-0.9	CCAAGTGCAAACTATGGTGC	ATAAACAATTTCAGTGCCGC	HinfI	495	282, 213
IIID	pkP3035	0.9	CGTAAACTACCAAACTCGGTG	GGTCTACTACAACTATACAGGC	Eco0109I	732	419, 313
IIIE	pkP3080	21.29	CGGTGGTGGTAAAAGTGTAAC	CAACATTCAGGCTGTGCTTTCC	Hpy188III	365, 76, 68, 35	241, 124, 76, 68, 35
IVA	F56B3	-24	TGATGGTGTGTCTGCGTACC	AGAGCTGGAGAGCACGGATA	DraI	301, 128, 71	429, 71
IVB	F52C12	−14.9	ACATTTAGTCACGCGTAGGG	GCCCGAATCTAGCACATAAG	HpaII	191, 137, 22	328, 22
IVC	B0273	1.8	AATACAGCAGTCGTTCCGTTC	TGAACTTCATGAACCAGCTTG	DraI	288, 144	432

IVD	K10D11	6.7	GATTATTT CAGAGGA GCAGAGC	CATAGCAC GTGGAATA ACCAC	HindIII	420	245, 175
IVE	T02D1	16.8	TGCTTAA AGTCATC GTGTCCA C	TGTAAACC GTATCGAA TCCGAC	EarI	174, 235	408
XA	pkP6139	-19.97	AAGAGTG AACCTTT TCCGTGA G	TGATGCAA TTTATACA CACGCC	MseI	401, 31	279, 122, 31
XB	pkP6120	-10.46	TCGTGGC ACCATAA AAGTG	GATTCAGA TCAAACAG AGGTGG	DraI	243	128, 115
XC	pkP6157	-0.14	GGGGTAT AATGAAC CAACCTG	TGTAGGAA CCGTTTGT TTCTTC	ApoI	261, 48	150, 111, 48
XD	pkP6161	9.38	ATCGACC CCAACAA TGCAC	TCCGTCAT CCAAATCT CCG	AseI	542	287, 255
XE	pkP6170	24.07	CGCTGTC ACAATCT CTAAAAT G	AAACCCTC CCCACTTT GTTGTC	ApoI	249, 118, 56	197, 118, 56, 52

A

htp-3^{H96Y}   *pch-2;htp-3*^{H96Y}

DAPI
MSH-5   Transition Zone

Mid Pachytene

Late Pachytene

B

control

htp-3^{H96Y}

pch-2;htp-3^{H96Y}

Average MSH-5 Foci / Row

Location Along the Germline

A

A

syp-1^{T452A}

pch-2;syp-1^{T452A}

DAPI

B