# Cryo-electron tomography of the actin cytoskeleton and membrane organization during lamellipodia formation using optogenetics.

Hironori Inaba[1,2,3], Tsuyoshi Imasaki[4], Kazuhiro Aoyama[5,6], Shogo Yoshihara[1,2], Hiroko Takazaki[7], Takayuki Kato[7], Hidemasa Goto[3], Kaoru Mitsuoka[6], Ryo Nitta[4] & Takao Nakata[1,2,*]

[1]Department of Cell Biology, Graduate School of Medical and Dental Sciences, Tokyo Medical and Dental University (TMDU), Tokyo, Japan.

[2]The Center for Brain Integration Research (CBIR), Tokyo Medical and Dental University (TMDU), Tokyo, Japan.

[3]Department of Histology and Cell Biology, Graduate School of Medicine, Mie University, Mie, Japan.

[4]Division of Structural Medicine and Anatomy, Department of Physiology and Cell Biology, Graduate School of Medicine, Kobe University, Hyogo, Japan.

[5]NanoPort Japan, Application Laboratory, Thermo Fisher Scientific, Tokyo, Japan

[6]Research Center for Ultra-High Voltage Electron Microscopy, Osaka University, Osaka,

Japan.

[7]Institute for Protein Research, Osaka University, Osaka, Japan

[*]Correspondence should be addressed to T.N. (E-mail: info.cbio@tmd.ac.jp)

Takao Nakata, M.D., Ph.D.

Department of Cell Biology, and The Center for Brain Integration Research (CBIR),

Tokyo, Medical and Dental University (TMDU), 1-5-45 Yushima, Bunkyo-ku, Tokyo

113-8510, Japan.

TEL: +81-3-5803-5143

Running title: Cryo-ET of PA-Rac1-induced lamellipodia

# 1 **Abstract**

Lamellipodia are sheet-like cellular protrusions crucial for cell migration and

endocytosis; their ultrastructure has been extensively studied using electron microscopy.

However, the ultrastructure of the actin cytoskeleton during lamellipodia formation

remains underexplored. Here, we employed the optogenetic tool PA-Rac1 combined with

cryo-electron tomography (cryo-ET) to precisely control Rac1 activation and subsequent

freezing via blue light irradiation. This approach enabled detailed structural analysis of

newly formed lamellipodia in cells with intact plasma membranes. We successfully

visualized lamellipodia with varying degrees of extension, representing different stages

of lamellipodia formation. In minor extensions, several unbundled actin filaments formed

"Minor protrusions" at several points along the leading edge. For moderately extended

lamellipodia, cross-linked actin filaments formed small filopodia-like structures, termed

"mini filopodia." In the most extended lamellipodia, filopodia matured at multiple points

along the leading edge, and the number of cross-linked actin filaments running nearly

parallel to the leading edge increased throughout the lamellipodia. These observations

suggest that actin polymerization begins in specific plasma membrane regions, forming

mini filopodia that either mature into full filopodia or detach from the leading edge to

form parallel filaments. This turnover of actin structures likely drives lamellipodial

1    protrusion and stabilizes the base, providing new insights into the structural dynamics of

2    the actin cytoskeleton and the mechanisms driving cell migration.

# 1  Introduction

Lamellipodia, the sheet-like and thin cellular protrusions, are crucial components that

form at the leading edge of migrating cells[1,2]. The actin cytoskeleton plays a fundamental

role in cell shaping and motility[3], and which is equally critical in lamellipodia formation

and retraction. The lamellipodia-like actin network is a fundamental structure observed

in cellular motility, including amoeboid movement and endocytosis[4,5]. These processes

are essential for the immune system[6,7], early development[8], and cancer metastasis and

invasion[9].

The small GTPase Rac1 localizes the periphery of the plasma membrane and is known as

a major upstream factor in lamellipodia formation[10,11]. Rac1 activates the SCAR/WAVE

complex[12], which in turn triggers the activation of the Arp2/3 complex[13]. The Arp2/3

complex, an actin nucleation factor, binds to the sides of existing actin filaments, leading

to the growth of new actin filaments at an angle of approximately $70°$[14]. This actin

branching is believed to be a principal feature of lamellipodia.

Most lamellipodia also possess rib-like structures: microspikes, which do not protrude

from the leading edge, and filopodia, which extend beyond it (Fig. 1a). These are straight

and bundled actin filaments, reported to be polymerized by the formin/mDia family and

crosslinked by fascin[15].

The dynamics and the structure of lamellipodia have been extensively studied over the

past half-century using both light and electron microscopy (LM and EM). Super-

resolution LM, total internal reflection fluorescence microscopy (TIRF), and speckle

microscopy have provided insights into protein localization and actin dynamics within

lamellipodia[16–21]. However, highly crowded actin structures in lamellipodia make it

challenging to resolve single filament structures using LM. In contrast, EM has been

instrumental in revealing the ultra-structure of lamellipodia[22–24]. Nonetheless, the

chemical fixation and negative staining in conventional EM potentially introduce artifacts.

The recent development of cryo-electron tomography (cryo-ET) has overcome these

limitations. With advanced image processing, cryo-ET can visualize native structures of

proteins and protein complexes in cells[25–27]. Given that lamellipodia are thin cellular

structures with a 100–200 nm height, they are ideal targets for cryo-ET. However,

observing lamellipodia with intact cell membranes using cryo-ET remains challenging

due to sample thickness and ice constraints. Consequently, most cryo-ET studies on

lamellipodia have been conducted with fixed and permeabilized cells[28–30]. Moreover,

most studies focus on motile cells like fish epidermal keratocytes, observing the structure

of already established lamellipodia, which overlooks the early stage of lamellipodia

formation. Recently, Chung and colleagues observed lamellipodia with intact cell

membranes in spreading cells, utilizing gal-8 for the extracellular matrix (ECM) to

facilitate lamellipodia extension thinly enough for cryo-ET analysis[31]. Their methods

enable the determination of the polarity of actin filaments and the high-resolution

structure of the Arp2/3 complex within an intact cellular membrane. However,

protrusions during cell spreading may not fully represent lamellipodia formation.

Here, we employed the optogenetic tool PA-Rac1 (photoactivatable-Rac1)[32] to observe

lamellipodia formation using cryo-ET. We utilized blue light irradiation within the

chamber of an automatic plunge freezer, allowing precise control over the timing of Rac1

activation and its subsequent fixation by quick freezing. The frozen specimens were then

subjected to cryogenic correlative light and electron microscopy (cryo-CLEM), and cryo-

ET data of lamellipodia with intact cell membranes was obtained. We analyzed the

ultrastructure of newly formed lamellipodia with varying degrees of extension, with a

focus on the interactions between the plasma membrane and the actin cytoskeleton at the

leading edge. These detailed investigations shed light on the ultrastructure of lamellipodia,

enhancing our understanding of the mechanisms driving cell motility.

# Results

**Light-induced lamellipodia formation on EM grids.**

To study the ultrastructure of the actin cytoskeleton and plasma membrane during lamellipodia formation through cryo-ET, it is critical to vitrify cells immediately after stimulation. This ensures that the resultant changes are distinctly observable in subsequent analyses. Accordingly, we utilized the optogenetic tool PA-Rac1 in this study, which consists of the photosensitive LOV2 domain and a constitutively active Rac1 mutant. Blue light exposure changes the LOV2 structure, enabling Rac1 activation, which induces lamellipodia formation[11,32]. For this study, we choose COS-7 cells for their adhesiveness, ease of transfection, and clear observation of lamellipodia formation on EM grids.

Initially, PA-Rac1-induced lamellipodia formation in COS-7 cells was characterized on glass-bottom dishes using confocal laser scanning microscopy (CLSM, Fig. S1a and Movie 1). Lifeact-mCherry was used as an F-actin marker[33]. Even before activation by light, cells spontaneously formed lamellipodia (Fig. S1a; 0:00). After activation, more prominent protrusions emerged from all around the cells, clearly differentiating the lamellipodia formation induced by PA-Rac1 (Fig. S1a; 2:30). Significant lamellipodia formation induced by PA-Rac1 was observed along the cell periphery, mainly where the

cortex forms inwardly curving arcs, extended over 20 µm in length, and reminiscent of a

crescent shape (Fig. 1b). These 'crescent-shaped edges' were characterized by abundant

cortical actin bundles, marked by the intense fluorescence of Lifeact-mCherry. In line

with the hypotheses that lamellipodia formation is initiated by the Arp2/3 complex-

mediated side branching from existing actin filaments oriented parallel to the

plasmalemma[28], these regions are reasonable sites of vigorous lamellipodia formation.

The initial extension of lamellipodia was typically completed within about 2 minutes of

starting blue light irradiation, followed by repeated formation and retraction. The extent

of lamellipodia expansion varied among cells and within different areas of the same cell,

with the most significant extension reaching lengths of over 5 µm. After the formation of

lamellipodia, the actin retrograde flow was observed (Movie 1). Microspikes and

filopodia, were formed within lamellipodia (Fig. S1a). Interestingly, these structures

moved laterally (Fig. S1b), and their tips occasionally detached from the leading edge

and gradually collapsed. Eventually, some were incorporated into the pre-existing cortical

actin bundles (Fig. S1d). These data suggest that bilateral flow[22] also occurs in PA-Rac1-

induced lamellipodia. Interestingly, the fluorescence intensity of Lifeact-mCherry was

notably high at the anterior region of the lamellipodia, comparable to the intensity in

cortical actin bundles. In contrast, it significantly decreased towards the central regions

within lamellipodia.

Next, lamellipodia formation with PA-Rac1 was induced on EM grids. The formation and

mobility of lamellipodia are largely influenced by the ECM. To effectively observe

lamellipodia through cryo-ET, they must be thinly spread and close to the substrate.

Therefore, EM grids were coated with poly-L-lysine and laminin. Similar to the

observations on glass-bottom dishes, lamellipodia formed around the cells upon blue light

irradiation, mainly from the crescent-shaped cell edges (Figs. 1c, d and Movie 2). The

actin retrograde flow and the formation of microspikes and filopodia were also observed,

along with their lateral movement (Fig. S1c) and collapsing (Fig. S1e). These

observations demonstrate the similarity of lamellipodia formation on EM grids to that

observed on glass-bottom dishes, at least through CLSM. Notably, lamellipodia formation

from the entire cell perimeter is an essential indicator for identifying PA-Rac1-induced

lamellipodia. Another principal indicator is the formation of dotted actin structures within

the interior of cells, approximately 1 μm in diameter (Fig. 1d, yellow arrowhead).

Although their specific nature remains unidentified. These features allowed us to

confidently select the cells with PA-Rac1-induced lamellipodia after plunge freezing.

To determine the optimal timing for freezing, we measured cell area changes post-PA-

Rac1 activation (Fig. 1e). The cell area increased significantly within the first four

minutes after the start of blue-light irradiation. Despite lamellipodia maintained beyond

this time, some areas showed retraction and re-expansion (Movie 2). Based on these

findings, plunge freezing was performed at the two-minute mark after the start of light

irradiation to observe nascent lamellipodia.

**Acquisition of cryo-ET for PA-Rac1-induced lamellipodia.**

To analyze the ultrastructure of PA-Rac1-induced lamellipodia with an intact membrane,

COS-7 cells expressing mVenus-PA-Rac1 and Lifeact-mCherry were cultured on the EM

grids and subjected to plunge freezing immediately after two minutes of blue light

irradiation (Fig. 2). Cells with PA-Rac1-induced lamellipodia were selected based on the

previously mentioned criteria through Leica Cryo-CLEM light microscopy. Subsequently,

these grids were subjected to cryo-EM, and tilt series were collected while correlated with

the cryo-LM images (Fig. S2). Since the plasma membrane tended to deform at the holes

of the Quantifoil (Figs. S3a, b)[34,35], these areas were avoided for analysis. Although the

signal-to-noise (S/N) ratio was reduced on the carbon sheet, there was still sufficient

contrast to segment cellular structures (Figs. S3c, d). After several attempts of grid

preparation and data collection, we successfully prepare well vitrified two grids, and

measured by cryo-EM. We succeeded in analyzing 16 tomograms from three cells,

capturing images from the leading edge to the rear regions of the lamellipodia with

varying degrees of extension (Figs. S2, S4). We measured the variety of potential

lamellipodia regions indicated by white squares in Fig. S4 and analyzed them.

The average height of lamellipodia from three cells ranged from 50 to 250 nm (Fig. 3a),

and filament density was between 2,500 and 15,000 per um$^{-3}$ (Fig. 3b), aligning with

previous studies[31,36]. The lower actin filament density in Cell 2 might be related to

pronounced lamellipodia spreading. Across all lamellipodia, filaments perpendicular to

the leading edge were relatively rare, peaking at angles of around 20–30 degrees and

around 90 degrees to the leading edge (Fig. 3c). When analyzed by individual cells, the

orientation of actin filaments varied from cell to cell (Fig. S5a). The orientation of actin

filaments, including regions outside the lamellipodia, was depicted in Fig. S5b. We also

explored the correlation between the length of actin filaments and their orientation to the

leading edge (Fig. 3d). For filaments shorter than 200 nm, a peak orientation was 15 to

25 degrees. In contrast, longer filaments tend to align more parallel to the leading edge.

The Arp2/3 complex mediates actin branch formation, crucial for organizing the actin

cytoskeleton in lamellipodia[20,23,30]. We identified actin branches by meticulously

reviewing tomograms within the 3dmod software, using the segmented actin filaments as

references. The criteria for identifying branches were based on their position at the ends

of filaments, branching angle around 70 degrees, and detecting densities on the base of

branches, indicative of the Arp2/3 complex. Identified actin branches are represented as

black dots in the segmentation images, with representative examples displayed in Fig. 3e.

The average frequency of the actin branch was one branch per 6.2 µm of filament length

(Fig. 3f). On the other hand, the average frequency of the actin branch outside of

lamellipodia was one branch per 31.7 µm filament length, indicating approximately five

times more branches within the lamellipodia. These data underscore the essential role of

actin branches in lamellipodia formation.

**The architecture of the actin cytoskeleton and protrusive membrane structure of the**

**growing lamellipodia.**

Next, we analyzed the details of the actin cytoskeleton at various stages of lamellipodia

formation, with the assumption that different degrees of extension represent different

stages of lamellipodia formation. First, we examined a lamellipodium with a relatively

low degree of extension, measuring approximately 0.5 µm (Fig. 4). The actin bundle on

the left side in the tomography (Figs. 4d, e and Movie 3) could be the original cortical

actin. Another example of such a degree of extended lamellipodia is shown in Fig. S4c-

1. Focusing on the leading edge, the plasma membrane appeared slightly undulated. Here,

we classified the surface features into three groups based on the ratio of height to width

(H/W) of the protrusions (Fig. S6). To determine H and W, we manually selected all

visibly distinguishable protrusions with a height of 20 nm and above. The width (W) was

measured as the distance between the two endpoints at the base of each protrusion, while

the height (H) was defined as the perpendicular distance from this base line to the apex

of the protrusion. Based on these measurements, we categorized the protrusions as "Large

protrusion" (H/W > 0.5), "Minor protrusions" ($0 <$ H/W $\leq 0.5$). The remaining areas,

where H/W < 0, were classified as "Concave regions" (Fig. S6). At the Minor protrusions,

several actin filaments ran towards the leading edge (Figs. 4f, g). Most of these actin

filaments, located just beneath the protrusive plasma membrane, had their growing ends

oriented toward the leading edge. In these Minor protrusions, the actin cytoskeleton was

not well-organized, lacking cross-linked actin filaments, and slightly curved filaments

were observed. Inside the lamellipodium, actin filaments were oriented in various

directions, showing no distinctive structures.

Next, we studied a lamellipodium extended to approximately 1.5 μm (Fig. 5 and Movie

4). In this lamellipodium, two Large protrusions were observed at the leading edge (Figs.

5f, g). In the Large protrusion highlighted in Fig. 5f, the growing end of actin filaments

localized in a region of the plasma membrane. These filaments formed two clusters, with

some filaments running parallel and extending in different directions, guiding the

membrane protrusion along their paths. In contrast, in the Large protrusion observed in

Fig. 5g, the actin cytoskeleton was arranged in a manner that two or more straight actin

filaments ran parallel and oriented toward the leading edge, similar to that in microspikes

and filopodia but on a smaller scale. The spacing between parallel actin filaments in these

protrusions (Fig. 5h), approximately 8 nm, suggested fascin-mediated cross-linking,

characteristic of microspike and filopodia formation[37,38]. Due to the limited number of

filaments and their shorter length (Fig 5e), hereafter, we refer to this structure as "mini

filopodia." Such structures were also detected in another tomogram (Fig. S4c-2).

Notably, within these Large protrusions, the localized high density areas on the plasma

membrane were observed (Fig. 5f, orange asterisk). These areas suggest the presence of

protein complexes or specialized membranes, offering a promising direction for further

exploration.

Meanwhile, in the Minor protrusion of the leading edge (below box f in Fig. 5e), several

actin filaments ran towards the leading edge without bundling, like the minor protrusions

in Fig. 4. In contrast, some single filaments ran towards the leading edge in the Concave

region, with more prominent filaments running parallel to the leading edge in these

regions.

When focusing on the interior of the lamellipodium (Fig. 5e), long actin filaments parallel

to the leading edge were prominent. Additionally, a few actin filaments color-coded in

blue at the lower left of the tomogram appeared detached from the leading edge but

exhibited an orderly structure like mini filopodia. For other actin filaments, the similar

orientation of neighboring filaments suggests forming an organized actin network in this

degree of extended lamellipodia.

Next, we quantified the distance between the growing ends of actin filaments and the

plasma membrane for each of the three groups, explicitly focusing on actin filaments

oriented toward the leading edge (Figs. S6b–e). Actin filaments adhering to the plasma

membrane (red arrows in Figs.5f, g, and Figs. S6c, d) were relatively common in the

Concave regions, accounting for about half of the observed cases. On the other hand, in

both Large and Minor protrusions, these adherent filaments constituted only about 15%

of the total, with a notable number of filaments showing gaps between their growing ends

and the plasma membrane. These gaps predominantly ranged from 10 to 20 nm, although

no uniform distance was consistently observed.

**Analysis of actin architecture in extended lamellipodia.**

In our tomograms, the lamellipodia in Cell 2 was most extended, measuring over 3 μm

(Fig. 6). In the LM image, a few filopodia or microspikes can be observed within the

lamellipodia (Figs. 6b, c). Part of filopodium was detected in a tomogram (Figs. 6d, e and

Movie 5), where bundles of over ten filaments ran linearly toward the leading edge. When

magnified, these bundles were spaced approximately 8 nm apart (Fig. 6h). Other actin

bundles ran parallel to the leading edge inner lamellipodia (for example, white/black

dotted boxes in Figs. 6f, g and Movie 6). These three to five actin filaments were also

cross-linked at approximately 8 nm (Figs. 6i, j). These data suggest that these actin

bundles, including filopodia and mini filopodia, are formed by the exact mechanism

involving fascin-mediated cross-linking.

Our observations primarily highlight the role of fascin in the prominent actin cross-

linking within lamellipodia. Detailed analysis revealed variations in actin filament density,

with regions of densely packed filaments and areas where they were sparse (Fig. S4).

Many closely situated actin filaments ran in similar directions, hinting at the involvement

of other actin cross-linking proteins, such as alpha-actinin[39], in their organization.

The protrusive membrane at the leading edge was not captured in this cell. In the Concave

regions of the leading edge, a single filament ran toward it (Figs. 6f, g), similar to previous

observations. Additionally, prominent actin filaments ran parallel to the leading edge

beneath the plasma membrane in these regions.

**Comparative analysis of actin dynamics and structure in anterior and posterior**

**regions of lamellipodia.**

Here, we obtained tomograms in various regions from the anterior to the posterior regions

of lamellipodia. The anterior region of the lamellipodia, located about 1 to 1.5 μm from

the leading edge, is a specialized area where the activity of actin polymerization and

depolymerization are exceptionally high[1]. Moreover, proteins that regulate actin

dynamics, such as Ena/VASP, the SCAR/WAVE complex, and profilin, concentrate in the

anterior region. Therefore, the anterior region is frequently analyzed separately from the

more posterior parts of the lamellipodia (Fig. 1a). In Cell 3, the high fluorescence

intensity of Lifeact-mCherry at the anterior regions and significantly lower intensity in

the posterior regions of lamellipodia (Figs. 6b, c), suggesting that there could be structure

differences between these regions. To compare these differences, we classified

tomograms, including other cells, based on distance from the leading edge: 'anterior'

(near) and 'posterior' (far) (Fig. 7a). In the anterior region, the orientation of actin

filaments was similar to the overall one (Fig. 7b). On the other hand, in the posterior

region, the peak at 30 to 40 degrees was more pronounced, with another peak observed

around 90 degrees. Although the density of actin filaments appeared to decrease towards

the rear of lamellipodia (Fig. 7c), this reduction was surprisingly minor compared to the

variance observed in Lifeact-mCherry fluorescence. Both regions contained fascin-

bundled actin filaments. Despite differences in orientations, no obvious distinction in the

actin cytoskeleton between these regions was identified, except for structures associated

with the plasma membrane at the leading edge.

**The ultrastructure of the convergence zone of lamellipodia and cortical region of**

**cells.**

The convergence zone of the lamellipodia, located near the original cortical region of the

cells[40,41], exhibited a predominantly unorganized actin cytoskeleton with minimal

branching and a high degree of filament bendiness (Fig. 8 and S4 b6–b8). The cortical

actin bundles showed continuous filaments extending across entire tomograms, indicating

their composition of actin filaments exceeding 1 μm in length (Fig. 8b). Within these

bundles, dozens of actin filaments twisted together with few branching. Additionally,

microtubules running parallel to the actin bundles suggested a coordinated structural

arrangement.

Beyond these bundled actin filaments, tubular membrane structures were observed

alongside microtubules, indicative of the endoplasmic reticulum (ER; Figs. 8c, d). This

1  arrangement aligns with previous studies in the LM[42], highlighting the role of

2  microtubules and ER in cellular transport and the maintenance of structural integrity.

## Discussion

In the present study, we conducted ultrastructural analyses on lamellipodia formation in cells with intact plasma membranes, using a combination of cryo-ET and optogenetics. The density and the orientation of actin filaments (Fig. 3a, b) were consistent with previous studies[22,31,36], affirming our methodology. Furthermore, the preservation of protrusive membrane structures and internal ER membrane (Fig. 8) suggests the effectiveness of the freezing process.

Our observations raise new fundamental questions on the organizations and roles of the actin cytoskeleton in lamellipodia formation, which could be different from those observed in motile keratocytes[28,29,43] or during spreading cells post-plating[31]. The frequency of actin branches (Fig. 3f) was significantly lower, about one-tenth compared to previous studies[28,31,43]. While it is conceivable that some branches were not detected due to the inherently noisy images produced by cryo-ET, it is unlikely that the number of missed branches would account for a tenfold difference. This discrepancy could be attributed to distinct experimental conditions, including the type of cells and protrusive activity of lamellipodia. The density of actin branching mediated by Arp2/3 is believed to influence the stiffness of the actin cytoskeleton and protrusion speed of lamellipodia[44,45]. Even considering these discrepancies, the higher frequency of actin

branches found within the lamellipodia affirms their critical roles in the organization of

the actin cytoskeleton during lamellipodia formation.

Taking advantage of preserving intact cell membranes, we analyzed the architecture of

the plasma membrane and the actin cytoskeleton at the leading edge in detail. Based on

our observations, we propose a model for the reorganization of the actin cytoskeleton

during lamellipodia formation, as illustrated in Fig. 9, encompassing different stages of

extension.

In the early stage, Minor protrusions form at the leading edge (Fig. 4 and Movie 3).

Several actin filaments grew towards the leading edge but were not bundled (Figs. 4f, g).

Conversely, in Convaved regions, actin filaments grow towards the leading edge

individually. The localized actin polymerization in the specific regions suggests the

involvement of liquid-liquid phase separation (LLPS) of actin-polymerizing factors[46–48].

Meanwhile, no organized actin structures were observed within the inner regions of

lamellipodia at this stage.

In the middle stage, the actin cytoskeleton exhibited a more structured and organized

arrangement compared to the early stage (Fig. 5 and Movie 4). Large protrusions appeared,

with 5 to 10 actin filaments growing to the leading edge, some of which were cross-linked

at approximately 8 nm intervals. These structures, which we refer to as mini filopodia,

resemble microspikes and filopodia, but have fewer and shorter filaments, making them

detectable only with EM. The formation of microspikes and filopodia within lamellipodia

is known to be facilitated by Ena/VASP proteins[49]. The spacing between actin filaments

in mini filopodia suggests bundling by fascin, which enhances Ena/VASP activity[50],

indicating a cooperative mechanism in the protrusion formation. Studies on LLPS have

shown that actin filaments become bundled after nucleation[39,48], suggesting that mini

filopodia form through the growth and bundling of Minor protrusions. At this stage,

Minor protrusions continue to be observed, suggesting that these structures turn over

during lamellipodia formation. Additionally, there was a mini filopodia-like actin

structure near the leading edge that appeared to be separated from it, suggesting it could

have detached after formation. Furthermore, some long actin filaments ran nearly parallel

to the leading edge within the inner regions of lamellipodia.

In the late stage, tomograms did not capture protrusive structures at the leading edge in

this study. Instead, prominent microspikes and filopodia were observed in cryo-LM, and

parts of these structures were captured in a tomogram (Fig. 6). These microspikes and

filopodia are considered to form as a result of further maturation of mini filopodia.

Conversely, many bundled actin filaments running nearly parallel to the leading edge

were observed within the inner regions of lamellipodia (Fig. 6i, j). It is possible that these

bundled actin filaments have originally been part of mini filopodia and later detached

from the leading edge. In living cells, microspikes and filopodia moved laterally due to

bilateral flow, occasionally collapsing (Figs. S1b-e). Such phenomena could also occur

in mini filopodia, suggesting that mini filopodia might either mature into filopodia or

detach from the cell membrane to form filaments running nearly parallel to the leading

edge. Actin filaments running parallel to the leading edge were often longer (Fig. 3d).

This makes sense as the retrograde flow of actin tends to push longer filaments into

obstacles where they stay. This is akin to how longer logs in a river are more likely to

encounter obstacles and align parallel to the flow. Within cells, these obstacles might be

other actin filaments and premature cellular adhesions[51].

The mechanism of lamellipodia formation is as follows: the initiation of actin

polymerization at the specific region of the plasma membrane at the leading edge leads

to the formation of Minor protrusions. These Minor protrusions then develop into mini

filopodia through the growth and bundling of actin filaments, which drive the plasma

membrane forward. These structures can further evolve into filopodia, but sometimes

detach from the leading edge and remain within the lamellipodia as actin bundles parallel

to the leading edge. The turnover of these structures within the lamellipodia is a major

driving force for lamellipodia extension. Our model complements the traditional role of

Arp2/3-mediated branching in lamellipodia extension.

Both in Large and Minor protrusions, gaps existed between the growing ends of actin

filaments and the plasma membrane (Fig. S6e). These gaps are expected, as actin

polymerization requires free space for the attachment of actin monomers to the filament

tips. The emergence of these spaces can be attributed to the random movement of the

plasma membrane, known as Brownian motion[52]. At the growing plus ends of actin

filaments, proteins like Ena/VASP and mDia/formin are believed to function, but the

resolution of our cryo-ET was insufficient to identify any complex densities. Conversely,

in the Concave regions, more actin filaments directly contacted the plasma membrane

than protrusion regions (Figs. S6d, e), likely due to higher tension in these areas.

Moreover, some actin filaments ran parallel to the leading edge beneath the plasma

membrane, likely helping to resist membrane tension.

Cryo-LM played a crucial role in identifying cells with PA-Rac1 induced-lamellipodia

and documenting their states. In particular, Cell 2, which exhibited the most extended

lamellipodia, provided significant information. Tomography No.1 and No.2 captured the

anterior region (Figs. S4b-1,2), with high Lifeact-mCherry fluorescence. In contrast,

tomography No.3 to 5 captured the posterior region (Figs. S4b-3–5), showing diminished

fluorescence. Despite differences in fluorescence intensity, the density of actin filaments

was not as varied (Fig. 3b). This discrepancy could be due to two factors. First, at the tips

of the lamellipodia, actin polymerization and depolymerization are active, with abundant

G-actin[16,53]. Lifeact-mCherry is known to bind not only to F-actin but also to G-actin[33].

Second, actin retrograde flow could carry Lifeact-mCherry away from the posterior

region[54]. Although actin filament orientation differed between regions (Figs. 7a, b), the

limited number of tomograms prevented clear elucidation of specific structural

differences.

In this study, we used the optogenetic tool PA-Rac1 to induce lamellipodia formation.

Optogenetics allows for precisely manipulating intracellular signaling molecules with

light, providing high specificity and time control[55,56]. This method addresses a

fundamental limitation of electron microscopy: the inability to observe temporal

dynamics due to specimen fixation. Although we froze samples at a single time point, the

combination of light irradiation and rapid freezing allows for precise control over reaction

times (Fig. 2), creating pseudo-time-lapse cryo-samples. By examining these frozen

specimens, we will be able to investigate the dynamics of lamellipodia formation at

nanometer resolution. While similar techniques have been applied to neural cells[57,58] and

purified proteins[59], using them for in-cell cryo-ET without staining, as demonstrated here,

offers a novel approach. This novel approach allows us to observe the ultrastructure of

intact, living cells, enhancing our understanding of cellular dynamics.

In summary, the present study integrates optogenetics with cryo-ET to reveal new insights

into the structure of lamellipodia formation, particularly the formation and growth of

protrusive structures. Despite challenges, it demonstrates the promise of combining these

techniques for a deeper understanding of cell motility. Future efforts should aim to refine

these methods to enhance our exploration of cellular mechanisms.

# Methods

## Cell culture

COS-7 cells (RCB0539: RIKEN BRC through the National BioResource Project of the MEXT/AMED, Japan) were cultured in Dulbecco's modified Eagle's medium (DMEM, Nacalai Tesque #08458-65), supplemented with 10 % fetal bovine serum (FBS, Biowest, #S1810) and incubated at 37°C with 5% $CO_2$.

Before seeding the cells, 200 mesh gold holey carbon grids (R1.2/1.3; Quantifoil Micro Tools) were coated with 0.01% (w/v) poly-L-lysine (Nacalai Tesque, #28360-14), applied with 20 nm gold colloidal markers (Sigma-Aldrich, #741965), and coated with thin carbon as described previously[35]. The grids were again coated with 0.01% (w/v) poly-L-lysine for over 1 hour and coated with 5 µg/mL laminin (Thermo Fisher Scientific, #23017015). A total of $1\times10^5$ COS-7 cells were seeded onto the grids, which were placed in 3D-printed grid holders[60] in a four-well plate (SPL Life Sciences, #30004).

  The transfection mixture, composed of 1 µg of pTriEx-mVenus-PA-Rac1[32] (kindly gifted from Klaus Hahn; Addgene plasmid, #22007) and 0.02 µg of Lifeact-mCherry[61], 50 µl of Opti-MEM™ (Thermo Fisher Scientific #31985070), and 3.6 µl of ViaFect™ transfection reagent (Promega, #E4982), was incubated at RT for 15 minutes. This transfection mix was added to the cells just before seeding.

After transfection, the cells were incubated at 37°C with 5% $CO_2$ for 16–20 hours before

vitrification. Before vitrification, the expression of the fluorescent proteins was

confirmed using either a BZ-X700 fluorescence microscope (Keyence) or an EVOS

M5000 imaging system (Thermo Fisher Scientific).

For observing live cell imaging on the glass-bottom dishes, cells were cultured and

transfected according to previously described methods [62]. Briefly, cells were seeded on

glass-bottomed dishes (Greiner Bio-one, #627870) coated with collagen (Cellmatrix Type

IC; Nitta Gelatin, #631-00771) two days before the observations. The following day, cells

were transfected with the plasmid described above, and analyses were conducted a day

later.

**Live cell imaging**

The grids with cultured cells were inverted so that the cell side faced downward and were

then submerged in a glass-bottom dish (Violamo, #GBCD15) filled with Leibovitz's L-

15 (Thermo Fisher Scientific, #11415064). Cells were analyzed under the serum-starved

condition in a sealed, heated chamber during live cell imaging at 37 °C without $CO_2$

supplementation. Images were obtained using a confocal laser scanning microscope

(FV1200, Olympus) on an IX83 microscope (Olympus) equipped with 40×/0.95 NA dry

objective lenses and FV10-ASW software (Olympus). Photoactivation and imaging

utilized standard setting for ECFP(C-Y-R), EYFP(C-Y-R), DsRed2(C-Y-R), and DIC

(Argon laser power: 5% at 458-nm and 1 % at 515-nm; diode laser power: 5% at 559-nm;

DM 458/515/560 dichroic excitation, SDM510 and SDM560 emission filters; 475–500

nm emission window for ECFP, 530–540 nm emission window for EYFP, and 570–670

nm emission window for DsRed2; DIC images were obtained with DsRed2 settings)

using the sequential line capturing mode (scan rate: 10 µs per pixel, pixel size 4.83 µm).

The interval for both imaging and optoactivation was set to 10 seconds. PA-Rac1

activation was induced using the ECFP channels. mCherry images were acquired through

the DsRed2 channel. Separately, mVenus images were captured with the EYFP channels

only after live imaging to confirm the expression of mVenus-PA-Rac1.

Image analysis was conducted using ImageJ/Fiji software (NIH)[63]. The changes in cell

area were quantified using mCherry images. These images were binarized and cell area

was measured using the "Analyze Particles" function.

Adobe Photoshop 2024 and Illustrator 2024 (Adobe Systems) were used for the final

figure preparation.

**Blue light irradiation and vitrification**

Samples were vitrified using either a Leica GP2 plunger (Leica Microsystems) or the

Vitrobot Mark IV System (Thermo Fisher Scientific) set to 37°C and 95% humidity.

Before transfer into the blotting chamber, the medium was manually blotted off, and 5 µl

of PBS was added to the grids. Inside the blotting chamber, blue light irradiation was

performed using a fiber-coupled LED (470 nm, 1000 mA; Thorlabs, #470F1), which was

connected to a ferrule patch cable (φ400 µm core, 0.39 NA; Thorlabs, #M79L01). The

output of LED was regulated by a T-Cube LED driver (1200 mA; Thorlabs, #LEDD1B).

The cable was fed into the blotting chamber through an aperture reserved for sample

application, with the blue light projecting directly from the front of the grid at

approximately 1 cm distance, at maximum power. The irradiation timing was manually

managed to ensure that blotting was completed, and the sample was plunged into liquid

ethane precisely 2 minutes after the onset of irradiation. Vitrification of the grids in liquid

ethane (-185°C) was following backside blotting for 10 seconds or bilateral blotting for

5 sec, using No.2 filter paper and blotting sensor of the Leica GP2 or Vitrobot Mark IV

System, respectively. Samples were stored under liquid nitrogen conditions until the time

of imaging.

**Cryo-Fluorescence Widefield Microscopy**

Fluorescence images of vitrified samples were captured using a THUNDER Imager EM Cryo CLEM widefield microscope (Leica Microsystems), which was equipped with a 50×/0.9 NA dry objective lens, a metal halide light source (EL6000), an air-cooled detector (DFC9000 GT), and a cryo-stage maintained at -190°C. Before observation, grids were placed into AutoGrids™ (Thermo Fisher Scientific). A 2 mm by 2 mm square area was imaged at the grid's center using the LAS X Navigator software (Leica Microsystems). For each field of view, a symmetrical 20 μm Z-stack with 5 μm intervals was captured centered around the autofocus point to create maps. Subsequently, lamellipodia-forming cells were visually selected for further imaging. A 20 μm Z-stack with 0.75 μm intervals was acquired at these locations. The imaging utilized multiple channels: transmitted light brightfield, reflection, GFP (Ex: BP470/40, Em: 525/50), and Texas Red (Ex: BP560/40, Em: 630/75). Image stitching was executed using the LAS X software. Small Volume Computational Clearing (SVCC) was applied to the acquired image stacks to diminish blurring and enhance weaker or subdued signals. Images and mosaic tiles were exported in TIFF format. Additional image processing, including maximum intensity projection, flipping, cropping, and contrast adjustment, was conducted using ImageJ/Fiji and Adobe Photoshop.

**Cryo-electron microscopy**

Cryo-electron tomograms were acquired on a Titan Krios (Thermo Fisher Scientific)

operating at an acceleration voltage of 300 kV and equipped with a Cs corrector (CEOS,

GmbH), a Volta phase plate and BioQuantam K3 direct detector with energy filter (slit

width of 20 eV) (Gatan). The microscope was controlled using the Tomography 5

software (Thermo Fisher Scientific). Initially, an atlas was acquired to identify the square

where the target cells were located. Subsequently, the tomography acquisition position

was determined based on the correlation with fluorescence images. Tilt series were

acquired with a range of up to -70 to 70° in 2° increments. The magnification was ×19,500,

resulting in a pixel size of 3.71Å. The electron dose per image was calculated as 1.44 e$^-$

/Å$^2$, with the total dose for series approximately 100 e$^-$/Å$^2$.

**Image processing**

Before 3D reconstruction, poor quality tilt images caused by obstacles such as grid bars

blocking the beam at high tilt angles, were removed. The tilt series images were then

reconstructed into 3D tomograms by using the IMOD software package[64]. Micrographs

were aligned by cross-correlation, followed by alignment through tracking 20 nm gold

fiducial beads coated on the grid. These aligned micrographs underwent reconstruction

for visual analysis using IMOD SIRT (simultaneous iterative reconstruction technique,

number of iterations: 8) at bin2. CTF correction was not performed on these

reconstructions.

Segmentation of cellular components was initially performed using AMIRA software

(Thermo Fisher Scientific). Before segmentation, the reconstructed images were binned

2 to 3 times (resulting in final bin4 and bin6 from the original images) and processed with

a Gaussian filter. The membrane structure was manually segmented, while actin filaments

and microtubules were segmented using the XFiber module of Amira[65]. Filament tracing

was undertaken after high-contrast structures such as the membrane, grid edge, and outer

cellular space were masked. Microtubule tracing utilized the following parameters, as

described previously[66]: cylinder length: 600 Å, angular sampling: 5, mask cylinder radius:

140 Å, outer cylinder radius: 125 Å, inner cylinder radius: 75 Å, and missing wedge

according to the individual tilt series. For actin filaments, these parameters were used:

cylinder length: 500 Å, angular sampling: 6 Å, mask cylinder radius: 45 Å, outer cylinder

radius: 35 Å, inner cylinder radius: 0 Å. The Trace Correlation Lines module was

configured with these parameter values: minimum seed correlation: 75–125 (tomogram

dependent), minimum continuation quality: 60–90 (tomogram dependent), direction

coefficient: 0.3, minimum distance: 70 Å, minimum length: 350 Å, search corn minimum

step size (%): 10. Segments and point coordinates were extracted into separate Excel

sheets from Amira and reformatted to IMOD/Etomo-style files using a MATLAB-script

("amira_reformat_to_coordinates.m" in "computational toolbox for ultrastructural

quantitative analysis of filament networks in cryo-ET data"[36]). Tracing errors were

manually corrected in 3dmod in IMOD, and the branch points of actin filaments were

also determined manually in 3dmod.

Quantitative analysis of actin filaments was conducted using the ultrastructural analysis

toolbox[36].

Visualization of the 3D model was performed in 3dmod. Color-coded maps illustrating

the angular distribution of actin filaments relative to the leading edge direction were

generated using the ultrastructural analysis toolbox. Three-dimensional images of

membranes and microtubules images created in 3dmod were meticulously merged with

the color maps in Adobe Photoshop.

The distance between the membrane and the tips of actin filaments was measured using

ImageJ/Fiji.

Graphs were created using Graphpad Prism 10 (GraphPad Software) and MATLAB

(MathWorks).

**Data availability**

The reconstructed tomograms shown in Figs. 4–6 have been deposited in the EMDB

under the following IDs: EMD-61088, EMD-61091, EMD-6092, and EMD-6193. The

raw tilt images, prior to 3D reconstruction, as well as the segmentation data will be

deposited to EMPIAR.

**Acknowledgments**

The authors are grateful to Dr. Florian Schur (Institute of Science and Technology (IST),

Austria) for helpful discussion about analyzing actin network and 3D printing and Yuriko

Sakamaki (the Research Core Center, TMDU) for technical help. We thank Drs T. Ishii

and T. Asano (TMDU), S. Kikkawa (Kobe University) for the helpful discussion, and S.

Nakamura (TMDU), K. Chin, Y. Sakihama, T. Shimizu (Kobe University), N. Makuta

(Osaka University), and M. Nakamura (Mie University) for assistance, and other

colleagues in Nakata lab and Nitta lab, the Research Core Center (TMDU) for usage of

the transmission electron microscopy and the carbon coater. This work was partly

supported by the Nanotechnology Platform Program and "Advanced Research

Infrastructure for Materials and Nanotechnology in Japan (ARIM)" of the Ministry of

Education, Culture, Sports, Science and Technology (MEXT, Japan). Proposal Numbers

JPMX09A18OS0055, JPMX09A19OS0046, JPMX09A20OS0002, JPMX09A21OS0020, and JPMX12220S0015. This research was also partially supported by the Research Support Project for Life Science and Drug Discovery (Basis for Supporting Innovative Drug Discovery and Life Science (BINDS)) from AMED under Grant Number JP22am121001.

**Funding**

This work was supported in part by JSPS KAKENHI (18K15002, 20K16105 and 22K06219 to HI, 18K19328 to HI and TN, 22K06809 to TI, 22K20680ZA and 23K14178ZA to SY, 23K23823 to TK and HT, 23K06674 to HG, 21H05254 to RN and TN), by AMED-CREST from the Japan Agency for Medical Research and Development (AMED; JP21gm161003 to TI), by JST [FOREST (JPMJFR214K to TI), Moonshot R&D (JPMJMS2024 to RN)], by Research Grants from the Takeda Science Foundation (to HI, TI and TN), the Uehara Memorial Foundation (to HI).

**Author contributions**

Conceptualization: H.I., T.I., R.N., T.N.; Methodology: H.I., T.I., A.K., H.T., T.K., K.M., R.T., T.K.; Validation: H.I., T.I., R.N., T.N.; Investigation: H.I., T.I., A.K., S.Y., R.N.,

T.N.; Resources: H.I., T.I., T.K., H.G., K.M, R.N., T.N.; Data curation: H.I., T.I., K.A.,

H.T., T.K., M.R., R.N., T.N.; Writing – original draft: H.I.; Writing -review & editing: T.I.,

S.Y., H.T., R.N., T.N; Visualization: H.I., T.I.; Supervision: R.N., T.N.; Project

administration: H.I., T.I., H.G., R.N., T.N.; Funding acquisition: H.I., T.I., S.Y., H.Y., H.T.,

T.K., H.G., R.N., T.N.

**Conflict of interests**

Kazuhiro Aoyama is affiliated with Thermo Fisher Scientific. The author has no financial

interests to declare. Other authors have no competing interests declared.

**Abbreviations**

Cryo-EM: cryogenic transmission microscopy

Cryo-ET: cryogenic electron tomography

CLEM: correlative light and electron microscopy

ECM: extracellular matrix

EM: electron microscopy

PA-Rac1: photoactivatable Rac1

LM: light microscopy

1 LLPS: liquid-liquid phase separation

2 MTs: microtubules

## References

1.  Small, J. V., Stradal, T., Vignal, E. & Rottner, K. The lamellipodium: Where motility begins. *Trends in Cell Biology* vol. 12 Preprint at https://doi.org/10.1016/S0962-8924(01)02237-1 (2002).

2.  Innocenti, M. New insights into the formation and the function of lamellipodia and ruffles in mesenchymal cell migration. *Cell Adh Migr* **12**, (2018).

3.  Nakata, T. & Hirokawa, N. Cytoskeletal reorganization of human platelets after stimulation revealed by the quick-freeze deep-etch technique. *J Cell Biol* **105**, (1987).

4.  Rottner, K., Faix, J., Bogdan, S., Linder, S. & Kerkhoff, E. Actin assembly mechanisms at a glance. *J Cell Sci* **130**, (2017).

5.  Svitkina, T. The actin cytoskeleton and actin-based motility. *Cold Spring Harb Perspect Biol* **10**, (2018).

6.  Luster, A. D., Alon, R. & von Andrian, U. H. Immune cell migration in inflammation: Present and future therapeutic targets. *Nature Immunology* vol. 6 Preprint at https://doi.org/10.1038/ni1275 (2005).

7.  Charpentier, J. C. & King, P. D. Mechanisms and functions of endocytosis in T cells. *Cell Communication and Signaling* vol. 19 Preprint at https://doi.org/10.1186/s12964-021-00766-3 (2021).

8.  Friedl, P. & Gilmour, D. Collective cell migration in morphogenesis, regeneration and cancer. *Nature Reviews Molecular Cell Biology* vol. 10 Preprint at https://doi.org/10.1038/nrm2720 (2009).

9.  Machesky, L. M. Lamellipodia and filopodia in metastasis and invasion. *FEBS Letters* vol. 582 Preprint at https://doi.org/10.1016/j.febslet.2008.03.039 (2008).

10. Ridley, A. J., Paterson, H. F., Johnston, C. L., Diekmann, D. & Hall, A. The small GTP-binding protein rac regulates growth factor-induced membrane ruffling. *Cell* **70**, (1992).

11. Nobes, C. D. & Hall, A. Rho, Rac, and Cdc42 GTPases regulate the assembly of multimolecular focal complexes associated with actin stress fibers, lamellipodia, and filopodia. *Cell* **81**, (1995).

12. Miki, H., Suetsugu, S. & Takenawa, T. WAVE, a novel WASP-family protein involved in actin reorganization induced by Rac. *EMBO Journal* **17**, (1998).

13. Machesky, L. M. *et al.* Scar, a WASp-related protein, activates nucleation of actin filaments by the Arp2/3 complex. *Proc Natl Acad Sci U S A* **96**, (1999).

14. Mullins, R. D., Heuser, J. A. & Pollard, T. D. The interaction of Arp2/3 complex with actin: Nucleation, high affinity pointed end capping, and formation of branching networks of filaments. *Proc Natl Acad Sci U S A* **95**, (1998).

15. Mattila, P. K. & Lappalainen, P. Filopodia: Molecular architecture and cellular functions. *Nature Reviews Molecular Cell Biology* vol. 9 Preprint at https://doi.org/10.1038/nrm2406 (2008).

16. Watanabe, N. & Mitchison, T. J. Single-molecule speckle analysis of actin filament turnover in lamellipodia. *Science (1979)* **295**, (2002).

17. Yamashiro, S. *et al.* New single-molecule speckle microscopy reveals modification of the retrograde actin flow by focal adhesions at nanometer scales. *Mol Biol Cell* **25**, (2014).

18. Mehidi, A. *et al.* Forces generated by lamellipodial actin filament elongation regulate the WAVE complex during cell migration. *Nat Cell Biol* **23**, (2021).

19. Rimoli, C. V., Valades-Cruz, C. A., Curcio, V., Mavrakis, M. & Brasselet, S. 4polar-STORM polarized super-resolution imaging of actin filament organization in cells. *Nat Commun* **13**, (2022).

20. Wu, C. *et al.* Arp2/3 is critical for lamellipodia and response to extracellular matrix cues but is dispensable for chemotaxis. *Cell* **148**, (2012).

21. Ponti, A. *et al.* Periodic patterns of actin turnover in lamellipodia and lamellae of migrating epithelial cells analyzed by quantitative fluorescent speckle microscopy. *Biophys J* **89**, (2005).

22. Koestler, S. A., Auinger, S., Vinzenz, M., Rottner, K. & Small, J. V. Differentially oriented populations of actin filaments generated in lamellipodia collaborate in pushing and pausing at the cell front. *Nat Cell Biol* **10**, (2008).

23. Svitkina, T. M. & Borisy, G. G. Arp2/3 complex and actin depolymerizing factor/cofilin in dendritic organization and treadmilling of actin filament array in lamellipodia. *Journal of Cell Biology* **145**, (1999).

24. Svitkina, T. M., Verkhovsky, A. B., McQuade, K. M. & Borisy, G. G. Analysis of the actin-myosin II system in fish epidermal keratocytes: Mechanism of cell body translocation. *Journal of Cell Biology* **139**, (1997).

25. Schneider, J. & Jasnin, M. Capturing actin assemblies in cells using in situ cryo-electron tomography. *Eur J Cell Biol* **101**, (2022).

26. Young, L. N. & Villa, E. Bringing Structure to Cell Biology with Cryo-Electron Tomography. *Annual Review of Biophysics* vol. 52 Preprint at https://doi.org/10.1146/annurev-biophys-111622-091327 (2023).

27. Fäßler, F., Javoor, M. G. & Schur, F. K. M. Deciphering the molecular mechanisms of actin cytoskeleton regulation in cell migration using cryo-EM. *Biochemical Society Transactions* vol. 51 Preprint at https://doi.org/10.1042/BST20220221 (2023).

28. Vinzenz, M. *et al.* Actin branching in the initiation and maintenance of lamellipodia. *J Cell Sci* **125**, (2012).

29. Mueller, J. *et al.* Load Adaptation of Lamellipodial Actin Networks. *Cell* **171**, (2017).

30. Fäßler, F., Dimchev, G., Hodirnau, V. V., Wan, W. & Schur, F. K. M. Cryo-electron tomography structure of Arp2/3 complex in cells reveals new insights into the branch junction. *Nat Commun* **11**, (2020).

31. Chung, W. L. *et al.* A network of mixed actin polarity in the leading edge of spreading cells. *Commun Biol* **5**, (2022).

32. Wu, Y. I. *et al.* A genetically encoded photoactivatable Rac controls the motility of living cells. *Nature* **461**, 104–108 (2009).

33. Riedl, J. *et al.* Lifeact: A versatile marker to visualize F-actin. *Nat Methods* **5**, (2008).

34. Nedozralova, H. *et al.* In situ cryo-electron tomography reveals local cellular machineries for axon branch development. *Journal of Cell Biology* **221**, (2022).

35. Aramaki, S., Mayanagi, K., Jin, M., Aoyama, K. & Yasunaga, T. Filopodia formation by crosslinking of F-actin with fascin in two different binding manners. *Cytoskeleton* **73**, (2016).

36. Dimchev, G., Amiri, B., Fäßler, F., Falcke, M. & Schur, F. K. Computational toolbox for ultrastructural quantitative analysis of filament networks in cryo-ET data. *J Struct Biol* **213**, (2021).

37. Jansen, S. *et al.* Mechanism of actin filament bundling by fascin. *Journal of Biological Chemistry* **286**, (2011).

38. Yang, S. *et al.* Molecular mechanism of fascin function in filopodial formation. *Journal of Biological Chemistry* **288**, (2013).

39. Langanger, G. *et al.* Ultrastructural localization of α-actinin and filamin in cultured cells with the immunogold staining (IGS) method. *Journal of Cell Biology* **99**, (1984).

40. Salmon, W. C., Adams, M. C. & Waterman-Storer, C. M. Dual-wavelength fluorescent speckle microscopy reveals coupling of microtubule and actin movements in migrating cells. *Journal of Cell Biology* **158**, (2002).

41. Gupton, S. L., Salmon, W. C. & Waterman-Storer, C. M. Converging populations of f-actin promote breakage of associated microtubules to spatially regulate microtubule turnover in migrating cells. *Current Biology* **12**, (2002).

42. Küppers, M., Albrecht, D., Kashkanova, A. D., Lühr, J. & Sandoghdar, V. Confocal interferometric scattering microscopy reveals 3D nanoscopic structure and dynamics in live cells. *Nat Commun* **14**, (2023).

43. Urban, E., Jacob, S., Nemethova, M., Resch, G. P. & Small, J. V. Electron tomography reveals unbranched networks of actin filaments in lamellipodia. *Nat Cell Biol* **12**, (2010).

44. Boujemaa-Paterski, R. *et al.* Network heterogeneity regulates steering in actin-based motility. *Nat Commun* **8**, (2017).

45. Suraneni, P. *et al.* The Arp2/3 complex is required for lamellipodia extension and directional fibroblast cell migration. *Journal of Cell Biology* **197**, (2012).

46. Case, L. B., Zhang, X., Ditlev, J. A. & Rosen, M. K. Stoichiometry controls activity of phase-separated clusters of actin signaling proteins. *Science (1979)* **363**, (2019).

47. Yang, S. *et al.* Self-construction of actin networks through phase separation-induced abLIM1 condensates. *Proc Natl Acad Sci U S A* **119**, (2022).

48. Graham, K. *et al.* Liquid-like VASP condensates drive actin polymerization and dynamic bundling. *Nat Phys* **19**, (2023).

49. Faix, J. & Rottner, K. Ena/VASP proteins in cell edge protrusion, migration and adhesion. *Journal of Cell Science* vol. 135 Preprint at https://doi.org/10.1242/jcs.259226 (2022).

50. Harker, A. J. *et al.* Ena/VASP processive elongation is modulated by avidity on actin filaments bundled by the filopodia cross-linker fascin. *Mol Biol Cell* **30**, (2019).

51. Yamashiro, S. & Watanabe, N. A new link between the retrograde actin flow and focal adhesions. *Journal of Biochemistry* vol. 156 Preprint at https://doi.org/10.1093/jb/mvu053 (2014).

52. Peskin, C. S., Odell, G. M. & Oster, G. F. Cellular motions and thermal fluctuations: the Brownian ratchet. *Biophys J* **65**, (1993).

53. Vallotton, P., Gupton, S. L., Waterman-Storert, C. M. & Danuser, G. Simultaneous mapping of filamentous actin flow and turnover in migrating cells by quantitative fluorescent speckle microscopy. *Proc Natl Acad Sci U S A* **101**, (2004).

54. Yamashiro, S. *et al.* Convection-Induced Biased Distribution of Actin Probes in Live Cells. *Biophys J* **116**, (2019).

55. Shaaya, M., Fauser, J. & Karginov, A. V. Optogenetics: The art of illuminating complex signaling pathways. *Physiology* **36**, (2021).

56. Tischer, D. & Weiner, O. D. Illuminating cell signalling with optogenetic tools. *Nat Rev Mol Cell Biol* **15**, 551–558 (2014).

57. Chakrabarti, R. *et al.* Optogenetics and electron tomography for structure-function analysis of cochlear ribbon synapses. *Elife* **11**, (2022).

58. Watanabe, S. Flash-and-freeze: Coordinating optogenetic stimulation with rapid freezing to visualize membrane dynamics at synapses with millisecond resolution. *Front Synaptic Neurosci* **8**, (2016).

59. Romero, A. M., Bonin, C. & Twomey, E. C. Time-Resolved Cryo-EM Specimen Preparation with Single Millisecond Precision. *bioRxiv* 2023.08.24.554704 (2023) doi:10.1101/2023.08.24.554704.

60. Fäßler, F., Zens, B., Hauschild, R. & Schur, F. K. M. 3D printed cell culture grid holders for improved cellular specimen preparation in cryo-electron microscopy. *J Struct Biol* **212**,

(2020).

61.  Kakumoto, T. & Nakata, T. Optogenetic Control of PIP3: PIP3 Is Sufficient to Induce the Actin-Based Active Part of Growth Cones and Is Regulated via Endocytosis. *PLoS One* **8**, 1–17 (2013).

62.  Inaba, H., Miao, Q. & Nakata, T. Optogenetic control of small GTPases reveals RhoA mediates intracellular calcium signaling. *Journal of Biological Chemistry* 100290 (2021) doi:10.1016/j.jbc.2021.100290.

63.  Schindelin, J. *et al.* Fiji: An open-source platform for biological-image analysis. *Nature Methods* Preprint at https://doi.org/10.1038/nmeth.2019 (2012).

64.  Kremer, J. R., Mastronarde, D. N. & McIntosh, J. R. Computer visualization of three-dimensional image data using IMOD. *J Struct Biol* **116**, (1996).

65.  Rigort, A. *et al.* Automated segmentation of electron tomograms for a quantitative description of actin filament networks. *J Struct Biol* **177**, (2012).

66.  Watanabe, R. *et al.* The In Situ Structure of Parkinson's Disease-Linked LRRK2. *Cell* **182**, (2020).

  **Figure legends**

**Fig. 1. PA-Rac1-induced lamellipodia formation on EM grids.** (**a**) Schematic
representation of subdomains in lamellipodia. (**b**) Lifeact-mCherry image of COS-7
cells, highlighting the crescent-shaped cell edges in red. Scale bar = 20 μm. **(c–e)** COS-
7 cells expressing PA-Rac1 and Lifeact-mCherry on EM grids. Images were captured
every 10 sec from -5 min, with PA-Rac1 activated by scanning with a single 458 nm
laser within 0–10 min at 10-sec intervals. **(c, d)** Representative time-lapse images. A
merged image of DIC and fluorescence of Lifeact-mCherry (yellow) **(c)**, and
fluorescence images of Lifeact-mCherry **(d)** are shown. A yellow arrowhead indicates
dotted actin structures within the interior of cells Scale bar = 20 μm. **(e)** Quantification
of cell area changes. Data are presented as the means ± SD from 6 cells. Blue
background indicates the activation period; red dotted line marks the timing of freezing.

**Fig. 2. Experimental workflow from sample preparation to tomogram acquisition.**
Laminin-coated EM grids were placed on the 3D-printed grid holders in 4 well dishes,
and cells were seeded on the EM grids. At the same time, plasmids encoded mVenus-
PA-Rac1 and Lifeact-mCherry were transfected into the cells. The next day, the cells
were rapidly frozen in a plunge freezer after inducing lamellipodia formation by
activating PA-Rac1 with blue light irradiation, controlling the timing of the freezing.
The frozen cells were observed using cryo-fluorescence microscopy to identify the cells
that had formed lamellipodia. By correlating the fluorescence images with low-
magnification electron microscopy images, the regions for tomography were
determined. Subsequently, continuous tilt series images were acquired by cryo-electron
microscopy from -70 degrees to +70 degrees, and tomography was reconstructed.

**Fig. 3. The organization of the actin cytoskeleton in PA-Rac1 induced lamellipodia**
**(a, b)** The averaged height **(a)** and the filament density **(b)** in lamellipodia across
individual cryo-ET. **(c)** The orientation of the actin filaments relative to the leading edge
(L.E.) in lamellipodia **(d)** Orientation distributions of actin filaments by length: 0–200
nm (left), 200–400 nm (center), >400 nm (right). **(e, f)** Actin branches within lamellipodia.
(**e**) Representative images of branched actin filaments. Scale bar = 25 nm. (**f**) The
frequency of actin branches in lamellipodia across individual cryo-ET.

**Fig. 4. Cryo-ET of PA-Rac1-induced lamellipodia in the early stage of formation.**

(a–c) A cryo-EM image (a) and corresponding cryo-CLEM images (b,c) of a PA-Rac1-induced lamellipodium in Cell 1 (Fig. S2) extended approximately 0.5 μm. (c) An enlargement of the area within the yellow rectangles in b. Scale bar = 10 μm in b and 2 μm in c, respectively. (d, e) A cross-sectional (x-y) slice from cryo-ET (d) and a corresponding segmented image (e) obtained from the area marked by a white box in b and c. The plasma membrane is represented in cyan. Actin filaments are color-coded to indicate their orientation relative to the leading edge (L.E.) of the cell. Here, 0 degrees represents perpendicular orientation relative to the leading edge, while 90 degrees indicates parallel orientation. Branching points of actin filaments are marked with black dots. Scale bar = 200 nm. (f, g) Enlarged images of the protrusive structure at the leading edge shown in the black dotted box in d and e. Scale bar = 50 nm.

**Fig. 5**. **Cryo-ET of PA-Rac1-induced lamellipodia in the middle stage of formation.** (a–c) A cryo-EM image (a) and corresponding cryo-CLEM images (b,c) of a PA-Rac1-induced lamellipodium in Cell 1 (Fig. S2) extended approximately 1.5 μm. (c) An enlargement of the area within the yellow rectangles in b. Scale bar = 10 μm in b and 2 μm in c, respectively. (d, e) A cross-sectional (x-y) slice from cryo-ET (d) and a corresponding segmented image (e) obtained from the area marked by a white box in b and c. Scale bar = 200 nm. (f, g) Enlarged images of protrusive structure at the leading edge shown in the black dotted box in d and e. Yellow dots mark branched points of actin filaments; red arrows point locations where actin filaments attach to the plasma membrane; an orange asterisk denote regions where the plasma membrane shows high intensity. Scale bar = 50 nm. (h) Enlarged images of the actin filaments running parallel in g, with three filaments indicated by red arrows running parallel to each other. Scale bar = 10 nm.

**Fig. 6**. **Cryo-ET of PA-Rac1-induced lamellipodia in the late stage of formation. (a–c)** A cryo-EM image (a) and corresponding cryo-CLEM images (b, c) of a PA-Rac1-induced lamellipodium in Cell 2 (Fig. S2) extended over 3 μm. (c) An enlargement of the area within the yellow rectangles in b. Scale bar = 10 μm in b and 2 μm in c, respectively. (d–g) Cross-sectional (x-y) slices from cryo-ET (d, f) and corresponding segmented images (e, g) obtained from the area marked by white boxes in b and c. Scale bar = 200 nm. (h-j) Enlarged images of actin bundles in filopodia (dotted boxes in d and e), and actin bundles running parallel to the L.E. within lamellipodia (dotted boxes in f and g). Scale bar = 10 nm.

**Fig. 7. The orientation and the density of actin filaments relative to the leading edge**
**(L.E.)** (**a**) The orientation of actin filaments relative to the L.E. Tomography numbers
designated as "anterior region" are marked in red, while those identified as "posterior
region" are noted in green. (**b**) The orientation distributions of actin filaments in anterior
regions (left: approximately 1.5 μm from the L.E.) and posterior region (right: more than
1.5 μm from L.E.). (**c**) The density of actin filaments is relative to the L.E., where straight
lines were drawn at 50 nm intervals from the L.E., and the number of intersecting actin
filaments was counted. The data obtained from each tomography were plotted on a graph.
9
**Fig. 8. Cryo-ET of the rare region of lamellipodia and cell cortical region before**
**lamellipodia formation.** (**a**) Cryo-CLEM image of the same cell shown in Fig. 2, cryo-
ET area in white box. Scale bar = 10 μm. (**b**) Segmented cryo-ET of white box area in
**a**. Blue: ER and inner cellular membrane structure, Green: Microtubules, Red: actin
filaments. Scale bar = 200 nm. (**c, d**) Magnified view of the area in B, focusing on the
parallel arrangement of tubular ER and microtubules. Includes a cryo-ET slice (**c**) and
its segmentation (**d**). Scale bar = 200 nm.
17
**Fig. 9. A model of actin cytoskeleton reorganization during formation of**
**lamellipodia.** This figure illustrates a model for the reorganization of the actin
cytoskeleton during lamellipodia formation, comprising three stages: **Early Stage**:
Minor protrusions form, with unbundled actin filaments growing towards the leading
edge, **Middle Stage**: mini filopodia emerge, with some actin filaments becoming cross-
linked. **Late Stage**: matured filopodia are present, with cross-linked actin filaments
running nearly parallel to the leading edge throughout the lamellipodia. See Discussion
for details. Illustration created with BioRender.

**a** Filopodium
Microspike
anterior
Lamellipodium
posterior
Cortical actin bundle
Convergence zone

**b**

**c**

**e**
458-nm light
Normalized cell area
1.15
1.10
1.05
1.00
0.95
-5    0    5    10
min

**d**
0:00
0:30
1:00
1:30
2:00
5:00

**Cell culture on EM grids**

3D printed grid holder

4 well dish

EM gird

**Transfection**
· **mVenus-PA-Rac1**
*Rac1 photoswitch*
· **Lifeact-mCherry**
*F-actin marker*

**Blue-light irradiation & plung freezing**

Plung freezer

tweezer

grid

480 nm LED

filter paper

**Time control**

liquid ethane

**Cryo-fluorescence microscopy**

**Cryo-electron tomography**

electron beam

tilt

sample

image

**Inaba et al., Fig. 3**

a

b

Lifeact-mCherry

↻ 90°

c

d

g

f

e

f

g

angle to the L.E. dircetion (deg)

**a**

**b**

**c**

**Early Stage**

Leading edge

cortical actin bundle

→ **cross-linked actin filaments**

**Middle Stage**

**mini filopodia**

**Late Stage**

**filopodia**