Behavior- and circuit-specific cortico-striatal decoupling during the early development of Parkinson's disease-like syndrome

Xu-Ran Yao[1,2], Yang Liu[1,2], Wei-Tong Zheng[1,2], Kai-Wen He[1,2,3*]

[1]*Interdisciplinary Research Center on Biology and Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 100 Haike Rd, Shanghai 201203, China*

[2]*University of Chinese Academy of Sciences, Beijing 100049, China*

[3] *Shanghai Key Laboratory of Aging Studies, Shanghai 201210, China*

[*] Correspondence: kwhe@sioc.ac.cn

ABSTRACT

Despite that cortico-striatal decoupling has been widely reported in individuals diagnosed with Parkinson's Disease (PD), its onset, circuit specificity and underlying mechanism remain largely unclear. To investigate these questions, dual fiber photometry is established for the first time to evaluate cortico-striatal coupling during varied motor behaviors, whose cell-type resolution was provided by the usage of Cre transgenic mouse lines. Contralateral turning, digging and licking show distinct coupling patterns, among which digging induces the strongest coupling. Striatal D1R-expressed medium spiny neurons (dMSNs) and D2R-expressed MSNs (iMSNs) similarly contribute to the cortical-striatal coupling during turning and licking but not digging, with much tighter coupling between the dMSNs and the M1 cortex. In PD-like mouse model induced via intra-striatal injection of synthetic mouse wildtype α-synuclein preformed fibril (PFF), digging-associated cortical-striatal decoupling emerges as early as 1-month post induction (Mpi), which becomes significant since 2 Mpi and correlates with later-onset behavioral deficit. Notably, impaired dMSNs but not iMSNs mediate this decoupling, which can be rescued by activation of D1 but not D2 receptor. Mechanistically, we found an inverted U-shape decline in striatal dopamine level along the disease development in PFF-injected mice. Supplement with L-DOPA alleviates the decoupling and motor deficit, suggesting that early dopamine deficiency directly contributes to the cortical-striatal decoupling and the associated motor deficit. These findings provide new insights into the temporal profile and mechanisms underlying the PD-associated cortico-striatal decoupling, which has been implicated as functional biomarker for early diagnosis of PD.

Keywords: Parkinson's disease; α-synuclein; fiber photometry; cortico-striatal coupling; striatal circuit specificity, dopamine.

INTRODUCTION

Parkinson's Disease (PD) is one of the most prevalent neurodegenerative disorders that primarily affects patients' motor function(1, 2). Various brain regions within the basal ganglia loop collaborate to regulate motor behavior(3, 4), with the functional coupling between the primary motor cortex (M1) and the striatum (STR) playing a crucial role in motor planning, execution, and learning(5-7). Functional magnetic resonance imaging (fMRI) revealed the reduced cortico-striatal connectivity, or cortico-striatal decoupling in PD patients(8-10). Furthermore, this PD-associated decoupling is correlated with both disease duration and behavioral impairment of patients(11, 12). Cortico-striatal decoupling is also associated with motor deficits in other disorders such as dystonia(13) and Huntington's disease(14). Together, these findings underscore the importance of understanding the mechanism of cortico-striatal coupling and characterizing the cortico-striatal decoupling associated with diseases.

D1 receptor- and D2 receptor-expressed medium spiny neurons (MSN) are the two main types of neurons in the striatum(15, 16). Both of them receive glutamatergic inputs from the cortex and are considered to play distinct yet debated roles in motor control via activation of the direct and indirect pathway, respectively(17-20). How these neurons contribute to the cortico-striatal coupling during motor behavior requires further investigation. In dopamine-depleted PD rodent models, impaired activities of D1R-expressed MSN (dMSN) and/or D2R-expressed MSN (iMSN) have been reported(21). While some studies found similar changes in both dMSN and iMSN firing(22) or increased similarity between their discharge frequency(23) after dopamine deletion, many others suggest the imbalanced activity between them might be the main cause of motor deficit(24-27). Whether and how dMSN and iMSN contribute to the cortico-striatal decoupling associated with parkinsonian remains to be answered.

The intrastriatal injection of α-Syn preformed fibril (PFF) effectively induces aggregation of α-Syn and

dopaminergic (DA) neuron cell loss in the substantia nigra (SN), two pathological hallmarks of PD(28)

along with motor dysfunction that gradually advances(29-31). This PD-like mouse model allows the

monitoring of potential alterations in cortico-striatal coupling preceding the onset of motor deficits.

In the present study, we first established a dual fiber photometry system to examine the cortico-striatal

coupling with behavioral and pathway specificity. Using the PFF-induced PD-like mouse model, we then

analyzed the characteristic as well as the underlying cellular and circuitry mechanism of cortico-striatal

decoupling during disease progression. Our findings validate a new method to study functional coupling

with cell type and circuit resolution, and for the first time unveil the temporal profile of M1 cortex-dorsal

striatum decoupling with the PFF-induced PD-like mouse model, uncover the distinct contributions of

dMSN and iMSN.

RESULTS

Dual fiber photometry measures coupling of cortical and striatal activity during different motor

behaviors

Several techniques, including electroencephalography (EEG), local field potential (LFP), and fMRI,

have been utilized to assess the coupling between neural activities in various brain subregions(32-34).

However, these approaches are limited in their capacity to elucidate the circuitry mechanisms

underlying such coupling. Considering that striatal neurons predominantly comprise dMSNs and iMSNs,

there is a pressing need for a novel method to explore the cell type-specific cortico-striatal coupling

during motor behavior in both healthy and diseased brains.

To address this challenge, we employed fiber photometry, which allows for the concurrent monitoring of

activity across different cell types in multiple brain regions in behaving mice(35). Two calcium indicators,

jRGECO1a and GCaMp7s, were respectively expressed in neurons in the deeper layers of the primary

motor cortex (M1) and dorsal striatum (STR) of the same hemisphere, followed by the implantation of

two optical fibers (Fig 1A-B; see Methods). Utilizing this dual fiber photometry approach, we successfully

monitored neuronal activities of M1 and the dorsal striatum without discernible crosstalk of fluorescence

signals (Fig 1B, Supplementary Figure 1A). Three naturalistic mouse motor behaviors, namely contra-

lateral turning, licking, and digging, were chosen to investigate cortico-striatal coupling (Fig 1C, 1G, 1K).

The coupling index is determined by calculating the Pearson's correlation coefficient between the

activities of the two channels within a time window around behavior initiation. Additionally, the phase

lag is estimated to assess the temporal alignment of the two responses (see Methods). Contralateral

turning has been demonstrated to elicit a striatal response(36). Fibers in the striatum consistently

detected robust neuronal activity in response to turning initiation (Fig 1D, right), validating the temporal

resolution of fiber photometry for monitoring fast motor behavior-evoked neuronal activity. Interestingly,

M1 neuronal activity was suppressed mildly during contralateral turning (Fig 1D, left), resulting in a

negative coupling between M1 and the striatum (Fig 1E; see Methods). In contrast, both M1 and the

striatum were activated during licking, with the striatal response exhibiting slightly earlier onset and

being more phasic compared to M1's response (Fig 1H-I). The distinct dynamics of these responses

consequently resulted in a coupling index close to 0 for licking behavior (Fig 1J). During digging, where

mice retrieved food pellets buried underneath the bedding (see Methods), both M1 and the dorsal

striatum exhibited robust activation upon initiation (Fig 1L). The digging-induced responses from both

channels were temporally aligned (Supplementary Figure 1C) and exhibited similar kinetics (Fig 1M),

reflected by a coupling index close to 1 (Fig 1N).

Thus, we have developed a dual fiber photometry approach to investigate the activity coupling between

M1 and the striatum. Our findings reveal distinct coupling patterns, implying that complex rules govern

cortico-striatal communication during various motor behaviors.

dMSNs and iMSNs distinctively contribute to the cortico-striatal coupling in different motor

behaviors

To investigate the cell type-specific cortico-striatal coupling, GCaMp7s was selectively expressed in

either striatal dMSNs or iMSNs by using D1R-cre or D2R-cre mice (Fig 2A-B, see Methods). The dual

fiber photometry system recorded clear calcium activities from M1 neurons and striatal dMSNs or iMSNs

concurrently (Supplementary Figure 2A-B). Both striatal dMSNs and iMSNs were similarly activated

upon contralateral turning and licking (Fig 2C, E), and their coupling with M1 activity remained

consistent across cell types (Fig 2D, F). These findings suggest that both cell types in the striatum

participate similarly in these two behaviors, consistent with previous studies (36-38). During digging

behavior, striatal dMSNs exhibited digging-evoked activity closely resembling that of total striatal

neurons and tightly coupled with M1 activity (Fig 2G-H). Notably, although iMSNs exhibited regular

activity during free moving (Supplementary Figure 2B), contralateral turning and licking, they displayed

a much weaker response during digging (Fig 2G) and significantly reduced coupling with M1 activity

compared to that of dMSN-M1 (Fig 2H). These findings further underscore the importance of

investigating cell type-specific M1-striatal coupling in motor behavior.

Intrastriatal injection of α-syn PFF induces PD-like pathology in wildtype mice

Cortico-striatal decoupling has been widely reported in patients with mild to severe Parkinson's disease

(PD), correlating positively with motor impairment(11, 39, 40). To investigate the onset and mechanism

underlying this PD-associated cortico-striatal decoupling, we adapted an inducible PD-like mouse

model established previously(41) by bilaterally inoculating α-synuclein (α-Syn) preformed fibrils (PFF)

into dorsal striatum of wildtype mice (Fig 3A). Behavioral and pathological phenotypes were examined

at both one- and three-month post PFF inoculation (Fig 3B). One-month post-injection (1Mpi), PFF led

to a slight reduction in rotarod learning ability (Fig $3C_1$) without affecting the rotarod test performance

(Fig $3C_2$) or the general locomotor activity (Fig $3C_3$). By 3-month post-injection, mice receiving PFF

displayed notable motor deficits (Fig $3D_1$, D_2), indicating a progressive PD-like motor impairment.

Concurrently, we assessed two pathological hallmarks of PD, α-Syn aggregates, and dopaminergic

neuron loss, via immunohistochemistry. PFF inoculation in striatum induced significant increase in the

phosphorylated α-Syn immunoreactivity (p-α-Syn), a common marker for synucleiopathy, both 1- and

3-month post injection in both the striatum (Fig 3E-F, L-M) and the substantia nigra (SN) (Fig 3H-I, O-

P). The rapid formation and spreading of synucleiopathy coincided with a progressive reduction in

striatal dopamine transporter (DAT) signal intensity in the striatum (Fig 3G, N) and a decreased in

dopaminergic neuronal cell count in SN (Fig 3J, Q). These findings confirm that PFF can initiate the

formation of α-Syn aggregates, which can further spread to the SN and trigger the pathology of

dopaminergic neurons in a time-dependent manner.

Behavior-specific cortico-striatal decoupling emerges early in the PFF-injected mice

Utilizing the PFF-induced PD-like mouse model, we aimed to investigate the onset timing and

mechanisms underlying cortico-striatal decoupling during disease progression by using the dual fiber

photometry method. PFF inoculation had no discernible effect on the fiber-based calcium imaging

(Supplementary Figure 3A-C). Given its robustness and circuitry dependency, we compared digging-

evoked M1-dorsal striatum coupling between PFF and PBS-injected mice. At one month post-injection,

cortico-striatal coupling largely persisted (Fig 4B-C), albeit with a moderate increase in trial-by-trial

phase lags among PFF mice compared to PBS counterparts (Fig 4B, C_3), indicating the emergence of

early temporal decoupling. By the second month post-injection, PFF mice exhibited a further increase

in phase lag relative to PBS controls (Fig 3D, E_2, E_3), accompanied by a declining trend in the cortico-

striatal coupling index (Fig $4E_1$) while cortical and striatal response amplitudes remained unchanged

(Supplementary Figure 3E). Following three to four months of PFF inoculation, decoupling in PFF mice

worsened, contrasting with sustained tight coupling observed in PBS mice between M1 and the striatum

during digging (Fig 4F, G_1). This decoupling in PFF mice were attributed to both elevated phase lag (Fig

$4G_2$, G_3) and a significant reduction in striatal response amplitude (Fig 4H). Temporal profiles of the

coupling index (Fig 4I, left), phase lag (Fig 4I, right), and response amplitude (Supplementary Figure

3F-G) collectively suggest an initial impairment in temporal precision of cortico-striatal communication

in this PD-like mouse model, with cortico-striatal decoupling exacerbating as pathology progresses. We

subsequently investigated whether the cortico-striatal decoupling in digging affect the digging behavior.

Digging duration and frequency were quantified and compared between groups (See Methods). It was

found that until three months post-injection PFF mice exhibited a shortened digging duration (Fig 5A)

and increased frequency (Fig 5B) relative to the PBS controls. Furthermore, the phase lag and coupling

index, but not the striatal response amplitude (Supplementary Figure 4C-D), strongly correlated

positively (Fig. 5C) and negatively (Fig. 5E) with digging frequency. Conversely, they exhibited opposite

correlations with digging duration (Fig 5D, F). These findings suggest that the functional decoupling

between M1 and dorsal striatum emerges early, predicting the later-onset behavioral deficit.

Of note, although the cortico-striatal coupling during digging was compromised in PFF mice, it remained

unaltered during licking even after 4 months of PFF injection (Fig 4K-M, Supplementary Figure 3H), at

which point PFF mice displayed normal licking behavior (Supplementary Figure 4E-H). This result

suggests that the PD-associated decoupling and motor deficit at early stages is behavior-specific.

Defects in dMSN neuronal activity mediates the digging-associated cortico-striatal decoupling

in PFF-injected mice

A functional imbalance between the dMSNs and iMSNs has been proposed to underlie the bradykinesia

in PD patients[16]. Therefore, we explored whether there is a cell-type effect in the observed early-

onset cortico-striatal decoupling in the PFF-induced PD-like mice. To investigate this, we employed a

methodology akin to that depicted in Figure 2A (Fig 6A). Additionally, D2R-Cre mice were also utilized

to examine the M1-dMSN coupling by expressing CreOFF-GCaMP in the striatum (See Method), which

could effectively label dMSNs (Supplementary Figure 5A) and generate reliable dMSNs responses

during various behaviors (Supplementary Figure 2C-F, Supplementary Figure 5B). We then tracked

cortico-striatal coupling in a pathway-specific manner beginning at 2 months post-injection, when

decoupling becomes apparent in PFF mice. The findings revealed that digging-related cortical and

striatal dMSN responses were markedly decoupled in PFF mice as early as 2 months post-injection

(Fig 6B), as evidenced by a significantly decreased coupling index (Fig $6C_1$) and an increased phase

lag (Fig $6C_2$). This M1-dMSN decoupling exacerbated further at 3~4 months post-PFF injection,

coinciding with a notably diminished striatal dMSN response compared to PBS mice (Fig 6D-E). In

contrast to the M1-dMSN decoupling, coupling between M1 and striatal iMSN during digging remained

largely unaffected in the PFF mice throughout the experimental time course (Fig 6F-J), indicating a cell

type-selective impairment in dMSNs contributing to the cortico-striatal decoupling in this PD-like mouse

model. Moreover, PFF inoculation had no discernible effect on the licking-evoked coupling between M1

and dMSN or iMSN (Supplementary Figure 5G-H), suggesting that cell-type-specific cortico-striatal

decoupling in PD is also behavior-specific.

Altered locomotion-associated striatal dopamine level in the PFF-injected mice

Insufficient striatal dopamine released from SN dopaminergic terminals is the primary cause of the

disturbed basal ganglia circuit that mediates the motor deficits in PD(16). Recent studies have indicated

a correlation between dopamine depletion and abnormal functional connectivity in both rodent models

and PD patients(42-45). Consequently, we hypothesized that abnormal dopamine level might underlie

the early-onset, cell-type-specific cortico-striatal decoupling observed in PFF-induced PD mice. To

monitor extracellular dopamine level at various time points, we expressed the dopamine sensor DA4.4

in striatal neurons (Fig 7A-B). In PBS mice, the fluorescence signal remained stable throughout the 3-

month recording period (Supplementary Figure 6A). The extracellular dopamine level was first

examined when PFF and PBS mice freely explored new cages (Fig 7C). PFF mice exhibited normal

general activity (Supplementary Figure 6B) and unaltered dopamine signals (Fig 7D-E), measured as

the summation of the fluorescent intensity during the exploration (see Methods). Since exploration of

new environments primarily triggers novelty-seeking behavior driven by dopamine release from the

ventral tegmental area (VTA)(46, 47), these findings suggest minimal impact of PFF inoculation on VTA-

mediated dopamine release up to 3 months post-injection. Dopaminergic neurons in the substantia

nigra compacta (SNc) are pivotal for voluntary movement(40, 48). To assess whether dopamine release

by SNc dopaminergic neurons is affected, mice were trained to run at a constant speed on a treadmill

(Fig 7F). Both PBS and PFF mice maintained their performance up to 3-month post-injection

(Supplementary Figure 6C). Interestingly, dopamine signal during constant speed running exhibited a

inverted U-shaped pattern of changes in PFF mice compared to PBS mice, with significant reductions

at 3-4 weeks and 10-14 weeks post-PFF injection, but a transient recovery at 6-7 weeks (Fig 7G-H).

These findings suggest impaired SNc-mediated dopamine release in PFF mice.

Activation of D1R but not D2R rescues the cortico-striatal decoupling and behavioral defects of

the PFF-injected mice

To investigate whether the impaired dopamine release is responsible for the dMSN-mediated cortico-

striatal decoupling and related behavioral deficit, PFF mice at 4-month post-injection received

intraperitoneal (i.p.) administration of L-dopa, SKF-81297 (SKF, a D1R receptor agonist), quinpirole

(Quin, a D2R receptor agonist), or saline (SAL) (Fig 8A). Consistent with our previous findings, both L-

dopa and SKF, but not Quin, rescued the digging-associated decoupling phenotypes, reflected by

decreasing phase lag (Fig 8C) and a trend of increasing coupling index in PFF mice (Fig 8D). Besides,

SKF but not Quin enhanced the striatal response amplitude in them (Fig 8E). These results further

support the notion that at this stage of PD development, deficient striatal dopamine leads to cortico-

striatal decoupling via impairing dMSN neuronal activation during digging.

Subsequently, the behavioral effects of these drugs were tested. Both L-dopa and SKF improved the

digging performance of the PFF mice (Fig 8F-G) without changing their general locomotion

(Supplementary Figure 7C). Moreover, SKF significantly increased the success rate, further supporting

that targeting dMSNs alone is more beneficial for the digging behavior (FigureS7D). In contrast, Quin

did not rescue the digging performance (Fig 8F, Supplementary Figure 7D), consistent with the absence

of digging-associated M1-iMSN decoupling in PFF mice.

DISCUSSION

Many clinical studies have revealed abnormal cortico-striatal functional coupling in patients diagnosed

with Parkinson's disease(8-11), among which the decoupling between M1 and striatum has been found

to tightly correlate with parkinsonian(49-51) even in the early stages of the disease(51). Therefore,

abnormal brain connectivity has been proposed as a functional biomarker for early diagnosis of

neurodegenerative diseases including PD(52, 53) and Alzheimer's disease(54, 55). The phenotypic

characterization and the underlying mechanistic investigation of such decoupling in PD animal model

may provide insight to facilitate not only the early diagnosis but also intervention.

Despite the development of various methods to detect inter-region brain coupling, such as LFP, EEG,

and fMRI(56), none of these techniques offer cell type-specific precision. Our current study establishes

a dual fiber photometry imaging method to investigate M1-striatal coupling with circuitry resolution by

using cell type-specific cre mouse lines. Although photometry imaging has lower temporal resolution

compared to LFP and EEG(57, 58), our method demonstrates its capability of differentiating distinct

M1-striatal coupling patterns associated with different motor behaviors (Fig 1). Among the three

behaviors, spontaneous contralateral turning as a reflexive behavior(59) evokes striatal but no obvious

M1 response, resulting in the weakest coupling reflected by the negative coupling index. Both tone-

cued licking and food pellet-driven digging evoke strong M1 and striatal activities, which are tightly

coupled with each other in particularly during digging. Interestingly, we found that only coupling

associated with digging, but not the other two motor behaviors with relatively weak coupling, displays

early abnormality in the PD-like mice. These findings suggests that motor tasks requiring effective

cortico-striatal co-activation might facilitate the early detection of PD-like pathological decoupling, which

is supported by neuroimaging studies showing task-related fMRI such as finger tapping(50, 60, 61)

detects abnormal brain activity and/or connectivity in early-stage PD(62).

Our current method further allows us to determine the contribution of dMSNs and iMSNs to the cortico-

striatal coupling. Consistent with previous studies(37, 63), we found that both types of MSNs were

activated during contralateral turning and licking, validating the accuracy of our method. Such co-

activation results in comparable contributions of dMSNs and iMSNs to the cortico-striatal coupling

during these behaviors. In contrast, our results reveal that M1 couples much more strongly with dMSNs

than iMSNs during digging behavior, mainly due to the greater and more timely activation of dMSNs.

Our findings are in line with previous studies using methods such as in vivo two-photon imaging (64) or

cell type specific optogenetic manipulation(19) that show distinct activation patterns of the dMSNs and

iMSNs during motor behaviors such as lever pressing(19, 64). Therefore, by simultaneously monitoring

M1 and striatal dMSNs or iMSNs activities, we provide evidences for the first time that the M1-striatal

coupling is regulated in a behavior- and circuit-specific manner.

Among all three motor behaviors we tested, only cortico-striatal coupling associated with digging is

impaired even before the behavioral deficit manifested. Digging is also the only behavior evoking

unequal activity of dMSNs and iMSNs. which may render it more susceptible to PD pathology. Striatal

MSNs form functional units, whose activity is responsible for driving specific behavior and is tightly

regulated by the complex lateral inhibitory network(65). Indeed, we found the significantly impaired

digging-evoked dMSN but not iMSN activity correlates with the behavioral deficit. It is noteworthy that

the phase lag, a parameter describing the relative timing of the M1 and striatal responses, shows the

strongest correlation with the digging performance (Fig 5C-F). This suggest that the effective

connectivity between the M1 and the striatum also, if not more, potently regulates specific behaviors.

Weakening of dMSNs while strengthening of iMSNs is observed in PD rodent models mimicking late

stages of PD, which lead to imbalance activity of the direct and indirect pathways and are considered

to mediate the motor dysfunction in PD(16, 66). Here we found that the activity of dMSNs but not iMSNs

is impaired during digging early in the PFF-induced PD-like mice. A previous study also found a selective

disconnection of dMSNs but not iMSN with the M1 in a 6-OHDA-induced partial dopamine lesion mouse

model(67). These findings together suggest that dMSNs might be more vulnerable at the early stages

of PD development. Interestingly, recent research has demonstrated that gene therapy selectively

activating striatal dMSNs can effectively reverse PD-like symptoms in both rodent and primate models

of PD(68). Here we also showed that administration of D1R agonist but not D2R agonist, rescued both

the digging-associated decoupling and behavioral deficits. Therefore, dMSNs might be better targets

for early PD intervention.

The lack of dopamine (DA) from SN dopaminergic terminals in the striatum can cause dysfunction in

the basal ganglia circuit and consequently lead to motor defects in PD patients(16, 69-71). Research

has revealed abnormal coordination between cortical and striatal ensembles in acutely dopamine-

depleted mice(72). Moreover, a clinical study reported an increase in M1-putamen connectivity after L-

DOPA intake in some PD patients(73). These findings collectively link the cortico-striatal coupling with

DA level in the PD state. In fibril-induced PD-like mice, a reduction in dopaminergic neurons in SNpc

begins as early as 4 weeks post inoculation, which is consistent with a significant reduction in the striatal

DA level during constant speed running. Interestingly, DA release transiently recovers 6-7 weeks post

inoculation, potentially due to a compensatory increase in spiking of the remaining dopaminergic

neurons in the SNpc(74). Therefore, the impaired DA level during motor behavior may mediate the

decoupling and behavioral deficit observed in the PD-like mice. Indeed, supplement of L-dopa almost

fully rescues these defects. Accumulating evidence suggests that abnormal axonal dopamine release

could occur before dopaminergic cell lost(71, 75), further supporting the notion that dopamine-

dependent, motor task-specific cortical-striatal decoupling might serve as an functional biomarker for

early PD.

Another mechanism regulating the cortico-striatal coupling might involve striatal fast-spiking

interneurons (FSIs), who is well known to dictate the temporal precision of neural information

processing(76-78). Within the striatal functional units that drive specific motor behavior, potent lateral

inhibitions are provided by the soma-targeting FSIs in addition to the MSN collaterals(65). In addition,

these FSIs are found to be activated faster with lower thresholds compared to MSNs(79, 80), making

them ideal for regulating the timing and dynamic of the striatal response. Moreover, FSI activity is

sensitive to DA(81) and its alteration has been shown to exacerbate striatal imbalance(24). It remains

to be determined whether the impaired DA level in the early stage of PD affects cortico-striatal coupling

by altering FSI activity.

CONCLUSION

Our study establishes a methodology utilizing fiber photometry to detect motor task-driven inter-regional

brain coupling with cell-type resolution. By employing this approach, we reveal the behavior-specific

M1-striatum decoupling preceding the behavioral deficit in the α-Syn PFF-induced PD-like mice,

supporting the use of motor task-specific cortico-striatal decoupling as a functional biomarker of early

PD. Additionally, we found that the decoupling associated with digging is selectively mediated by

impaired dMSNs in a dopamine-dependent manner, unveiling for the first time the circuitry mechanism

mediated task-specific decoupling in the PD state.

Acknowledgement

We thank Dr. Cong Liu and his students for their help in preparing the PFF. We also thank the staff

members of the animal facility at the National Facility for Protein Science in Shanghai (NFPS), Shanghai

Advanced Research Institute, Chinese Academy of Sciences, China for excellent support. This work is

supported by NSFC (Grant No. 32271004, 32070963), and Shanghai Municipal Science and

Technology Major Project (Grant No. 2019SHZDZX02) to K.-W.H.

Author contributions

349   Conceptualization, K.-W.H.; Primary Investigation, X.-R.Y.; Assistant Investigation, Y.L., W.-T.Z.; Data

350   Analysis, X.-R.Y., Y.L.; Writing, X.-R.Y., Y.L. and K.-W.H.; Funding Acquisition, K.-W.H; Supervision, K.-

351   W.H.

List of abbreviations

PD   Parkinson's disease

M1   Primary motor cortex

Str   Striatum

SN   Substantia nigra

DA   Dopamine

DAT   Dopamine transporter

D1R   Type 1 dopamine receptor

D2R   Type 2 dopamine receptor

MSN   Medium spiny neuron

dMSN   direct pathway MSN

iMSN   indirect pathway MSN

α-Syn   α-Synuclein

PFF   Preformed fibril

Mpi   Month(s) post injection

METHODS

Animals

All procedures were approved by the Institutional Animal Care and Use Committees at the

Interdisciplinary Research Center on Biology and Chemistry (IRCBC), Chinese Academy of Science.

C57BL/6J mice were purchased from Shanghai Lingchang Biotechnology Co., Ltd.. The transgenic

mice (D1R-Cre: MMRRC_034258-UCD; D2R-Cre: MMRRC_034258-UCD) were bred in-house. Mice

were group housed with a 12 h:12 h light:dark cycle and ad libitum access to food and water until

experiments. Male mice of 2-8 months of age were used with exception stated specifically. All

experiments were carried out at ZT (Zeitgeber time) 15~19.

Preparation and characterization of mouse wildtype α-Syn PFF

The procedures followed the same protocol as described previously(31). In brief, pET22 vector plasmid

containing α-syn gene was transfected into BL21 (DE3) cells. α-syn expression was induced by IPTG

(isopropyl-1-thio-D-galactopyranoside). Cells were then harvested and sonicated followed by boiling,

streptomycin (20 mg/mL) treatment, pH adjustment, and dialysis overnight in turn. The α-syn monomer

was then incubated in solution containing 100 μM in 50 mM Tris, pH 7.5, 150 mM KCl, and 0.05% NaN3

buffer at 37 °C with constant agitation (900 rpm) in ThermoMixer (Eppendorf) for one week to form fibril

structure. The fibril was quantified by subtracting the residual soluble α-syn from the total amount of α-

syn monomer and the fibril pellet was suspended with PBS to 2 µg/µL. The obtained fibril was sonicated

(20% power 15 times (1 s on, 1 s off) on ice, JY92-IIN sonicator) and the structure was confirmed by

TEM (FEI Company, USA).

Stereotaxic injection

Craniotomy was performed as described previously(31, 76). Briefly, mice were anesthetized with

isoflurane vapor (1~2% in air) and head-fixed in a stereotaxic frame (RWD Life Science, China) with

heating pad (~37 ℃) underneath to maintain body temperature. The scalp was removed, and

virus/fibril/PBS was delivered at a rate of 80 nl/min to either M1 (bregma/midline/depth: 1.77/1.5/-0.67

mm) or dorsal lateral striatum (dlSTR. For GCaMP7s: -0.11/2.5/2.3 mm; for PFF/PBS, DA sensor,

0.2/2.0/2.6 mm) by using a glass pipette connected to a micro syringe pump (Stoelting, USA). For

photometry calcium imaging, 150 nl of AAV-hSyn-NES-jGRECO1a was injected into M1 layer V; 350 nl

of AAV-hSyn-GCaMP7s (or 400nL AAV-hSyn-DIO-GCaMP7s for Cre mice) was injected into dlSTR. To

measure the striatal extra-cellular dopamine level, 400nl of AAV-hSyn-GRabeen-DA4.4 was injected

into dlSTR. All viruses were diluted to a titer of 10^{12} vg./ml. For establishing the α-Syn PFF model, PFF

or PBS (0.1 µl/g body weight, 2.5 µl at most) was injected at a speed of 0.5 µl/min for the first 0.2 µL

and 0.2 µL/min for the rest.

Fiber photometry surgery and imaging

Optical fibers (NA: 0.37; diameter: 200µm; Inper, China) were implanted using the same M1 and dlSTR

coordinates except 0.05 mm less in depth right after the virus injection. Fiber was secured to the skull

using dental cement (C&M Super Bond, Japan). Mice were then housed individually and allowed to

recover for at least 2 weeks before imaging.

Photometry recording during behavioral paradigms was conducted during the dark cycle utilizing a

commercial multichannel fiber photometry system (Inper, China). Excitation light at wavelengths of 470

nm, 561 nm, and 410 nm was used to excite GCaMP7s/DA-4.4, RGECO1a, and Ca^{2+}-independent

internal channel control, respectively. Fluorescence signal was collected at 40 Hz using InperStudio

(MultiColorEVAL15) software, which automatically synchronized the calcium signal with video footage

from a camera positioned above the arena. Throughout the imaging, the patch cord attached to the

optical fiber was suspended by a helium-filled balloon to minimize its weight impact on the mouse

behavior. All behavioral trials lasted for less than 30 min.

Immunofluorescence (IF) staining

IF was performed as previously describe(31). Briefly, mice were anesthetized with isoflurane vapor and

perfused with PBS followed by about 40 ml of 4% paraformaldehyde (PFA). Subsequently, whole brain

was removed and fixed overnight in PFA at 4°C. The fixed brains were then saturated in PBS

containing 20% sucrose for overnight to ensure the mice brains sinking bottom, and 30%

sucrose for 2 d. Afterward, the brains were embedded in tissue freezing medium (OCT, Leica) and

rapidly frozen and stored at -80°C. Next, 20 μm coronal slices were sectioned using the cryostat (Leica,

CM3050s-1-1-1). These sections were incubated in blocking solution (PBS containing 5% normal goat

serum (Sigma) and 0.1% Triton X-100) for 2 hours at room temperature, followed by overnight

incubation at 4°C with the primary antibody diluted in the blocking solution. After incubation, the primary

antibody solution was washed out three times with PBST (0.1% Tween-20 in PBS) and once with PBS,

followed by incubation with the fluorophore-conjugated secondary antibodies for 2 hours at room

temperature. For MSN type-specific labeling, DAPI (diluted at 1:500) was then used to stain the nuclei

for 10 minutes at room temperature. Information of all antibodies used were summarized in

Supplementary Table 1.

To validate the expression and location of the calcium and dopamine sensor, PFA-fixed brains were

sectioned into 60 μm coronal slices using a vibratome (Leica VT 1000S). Slices with optical fiber track

were selected for subsequent fluorescence imaging. Slices were further stained for either D1R or D2R

for cell-type specific colocalization.

Confocal Imaging

Fluorescence images were captured using a spinning disk confocal microscope (Andor, UK). Different

lasers with wavelengths of 630 nm, 561 nm, 488 nm, and 405 nm were sequentially utilized to excite

the Fluor 647, Fluor 568 or RGECO1a, Fluor 488 or GCaMP7s or DA4.4, and DAPI respectively.

Imaging was performed using a 20× air (NA = 0.75), or 40× water (NA = 1.25) objective, with consistent

settings maintained for each set of experiments. The image resolution is 0.6 μm:0.6 μm/pixel (x:y) for

20× and 0.15 μm:0.15 μm/pixel (x:y) for 40× objective. Z-stack images were acquired for all brain

regions and then projected in the z-direction to obtain the maximum intensity for subsequent analysis

in ImageJ. Step size (μm) and thickness (μm) of each brain region were as followed: SN: 0.5, 1; Striatum:

1, 5 (for characterization of the PFF-induced pathology); Others: 2, 10. The whole view of STR was

analyzed, while the SN region was manually selected based on the mouse brain atlas. The average

intensity of phosphorylated α-synuclein (P-α-syn) and dopamine transporter (DAT) was measured using

the same background subtraction and contrast transformation applied to each dataset, and then

normalized to the control (PBS group). DAT-positive dopamine neurons in the Substantia Nigra pars

compacta (SNpc) were automatically selected and counted using the Spot module of Imaris (Oxford

Instruments), with settings consistent with previous descriptions(41).

Behavioral task or test

a.   **Digging test**

Mice were food deprived for 24h before behavioral digging test. During the deprivation, two food

pallets (25mg, sucrose and milk powder in a 1:1 ratio. Xietong Bio-engineering Co., Ltd., China) were

placed in its home cage to introduce novel food. The test environment consisted of a lidless,

semitransparent box (FENGSHI group, China), identical in size and texture to the mouse's home

cage (l:w:h: 29 cm×18 cm×13 cm), serving as the arena for the digging test. The bottom of the

arena was covered with approximately 3cm-thick clean bedding, and twelve food pallets were evenly

buried in the bedding. Additionally, a solitary food pallet was strategically positioned at the center of

the arena's surface on top of the bedding, easily accessible for the mice upon entering the arena,

thereby enticing them to dig for buried pallets. The mice's performance in the arena was recorded

using two cameras—one providing a top view and the other a side view—for subsequent post-hoc

analysis. After completing the test, the bedding was discarded, and the arena was cleaned with 75%

ethanol.

b. Sound-cued licking task

Mice were habituated and trained for three days prior the final experiment.

Day0. Completely water deprivation

Day1. Mice were required to lick water from the water outlet connected to a sensor. Once the mice

lick the outlet, a single dose of 5μl water was pumped out from the outlet. Each mouse repeated this

process 100 times.

Day2. A sound cue (3000Hz, 200ms) with random interval (10-20 s) was introduced, and mice had

to associate it with water. 5μl water was pumped out if mice licked the outlet within 3s after the cue;

otherwise, they received nothing. A 5-second interval allowed mice to consume the water before

moving to the next random time interval. Each mouse was allowed to complete 80 water trials

Day3. The random time interval from Day 2 was replaced with a 5-second non-licking interval. Mice

had to refrain from licking the outlet for 5s to trigger the sound cue. If they licked during this interval,

they had to wait another 5s for the sound cue. The time window for mice to respond to the cue and

to consume water remained the same as Day 2. Each mouse was allowed sufficient time to complete

120 watered trials.

Day4. Fiber photometry recording was conducted while mice performed the task, following the

paradigm of Day3. The timestamp of licks within each watered trial (see Supplementary Figure 4)

was recorded by a sensor controlled by customized Arduino code. The initiation of licking was defined

as the first lick detected by the sensor after the sound cue.

c. Rotarod

The procedures followed the protocol described in a previous report(31). Briefly, mice underwent

three days of training, consisting of two adaptive trials at a low rolling speed of 5 rpm/min, followed

by a third trial with accelerating speed ranging from 0 to 40 rpm/min. After a rest day on the fourth

day, mice were tested twice on the rotarod on the fifth day. Each testing session lasted for a total of

5 minutes. During the first 1.5 minutes, the speed of the rotarod accelerated from 0 to 40 rpm/min,

which was maintained for the remaining time. Mice were tested twice, and the maximum duration

they stayed on the rotarod was recorded for final analysis. Between each trial, the rotarod was

thoroughly cleaned with 75% alcohol.

d. Open field test

After a period of habituation, mice were introduced into the center of the open field chamber (40 × 40

cm) and given 10 minutes to freely explore its surroundings. After the test, the arena was cleaned

with 75% ethanol. Their activities were observed and recorded using a camera positioned on the

ceiling, and the data were subsequently analyzed using EthoVision XT 11.5 software (Noldus,

Netherlands).

e. Free exploring in a new cage

The procedures and settings for the open field test were replicated with the exception of utilizing a

lidless and empty cage resembling to the mouse's home cage (l:w:h: 29 cm×18 cm×13 cm) as the

arena.

f. Free moving

The procedures and settings were exactly same with free exploring but the arena covered with clean

bedding.

g. Treadmill running

Mouse was placed on the treadmill (Zhenghua Biologic apparatus, China) 10mins before testing for

habitation. During the test, the rolling velocity of conveyor was constantly 10m/min and the testing

session would last for 10mins. To prevent interaction between mice, each mouse was isolated by

baffles while running on the treadmill. After the test, conveyor and baffles were cleaned with 75%

ethanol. Both male and female mice were included in this experiment but they were separated by

gender during the test. Any mice that failed to keep up with rolling conveyor during the test was

excluded from the analysis.

Pharmacology

The experimental design followed a previous study that investigated the effects of dopaminergic drugs

on striatal MSNs using calcium imaging in parkinsonian mice(26). The drugs (all compounds from

Merck), including D1R agonist (SKF81297; 1mg/kg), D2R agonist (quinpirole; 0.5mg/kg), L-DOPA

(1mg/kg) together with benserazide (15mg/kg), were dissolved in 0.9% saline and filtered through a

PES membrane filter unit (0.22μm, Millex). These solutions were stored at 4℃ and used within a week.

The drugs were administered via intraperitoneal (i.p.) injection, with a volume determined according to

body weight (0.1 ml/10 g). Both behavioral and photometry experiments commenced 15 minutes after

i.p. except quinpirole (60 minutes post i.p.). Information of all chemicals used were summarized in

Supplementary Table 1.

Quantification and statistics

Behavioral analysis

Digging behavior.　Trials was manually picked from the video by the investigator. The initiation of

digging behavior was determined when the bedding beneath the paws of the mice were visibly displaced.

The termination of a digging trial was defined as either no bedding displacement for 1 second or the

mice engaging in any other behaviors such as rearing, grooming, or eating pallets. The average duration

of all digging trials within the first ten minutes after the mice acquired the pallet on the top of the bedding

was considered as the digging duration. Digging frequency was defined as the number of digging per

minute during this period. The success rate was calculated as the percentage of trials in which the mice

successfully obtained the pallet out of the total number of trials.

Licking behavior.　It was recorded using a touch sensor controlled by an Arduino chip, with each lick

automatically captured and timestamped. The data were then written into a text file. Customized R code

was used to analyze the data from the text file and calculate various metrics to characterize licking

behavior. A series of successive licks with intervals less than 1 s was considered as a bout. The duration

of a bout was defined as the bout length.

Qualitative analysis of photometry signal

    a. *Photometry signal processing*

The photometry data collected was pre-processed for bleaching and motion correction using

InperDataProcess v0.7.2 software. The onset of digging and contra-turning behavioral events was

manually marked based on the synchronized videos. For digging behavior, only trials having ≥ 10 s of

dig-less baseline and ≥ 3 s of digging were selected for final analysis. For turning behavior, only trials

with ≥ 4 s intertrial interval and ≥ 90 degree turning were chosen for final analysis.

All fiber signal was presented as dF/ F_0 using the equations:

$$dF/F_0 = \frac{F - F_0}{F_0}$$

F represents the fluorescence value of each timepoint. F_0 is the average of F of baseline.

For digging and licking behavior, the time window was -5 to 5 seconds (relative to behavior initiation),

during which the -4 to -2 s was used to calculate the F_0. For contra-turning, the time window was -1 to

1 second, and the baseline was -1 to 0 s.

The data were further transformed into z-score using the formula to better compare the dynamic of M1

and striatal responses:

$$Z = \frac{F - \overline{F}}{\sigma}$$

Where $\overline{F}$ is the population mean, and σ is the population standard deviation.

    b*. Coupling index & Phase lag.*

The coupling index represented the Pearson's correlation coefficient of time-aligned averaged M1 and

STR signals. It was calculated using the R cor function. Additionally, the ccf function in R was utilized

to determine the lag at which the maximum cross-correlation between the signals occurred. The lag.max

parameter was set to be twice the sampling rate.

c. *Plateau & peak amplitude*.

The plateau amplitude was defined as the mean of the normalized (dF/F_0) signal within the plateau

stage (1~2 s, relative to initiation) for digging behavior. For licking behavior, the peak amplitude was

that within 0~0.5s.

d. *Sum dFF activity*.

The data of each mouse collected during 10 minutes of treadmill running or free-moving was separated

into bins of equal length, typically 8 seconds each. Each bin of data was normalized to dF/F_0, where F_0

represented the 20th percentile of the data within that bin. In the meanwhile, we calculated the

Pearson's correlation coefficient between the sequence of dF/F_0 values and the raw F sequence. If this

correlation coefficient lower than 0.8, the mouse was excluded from the further analysis. The sum dFF

activity was then determined by summing the resultant dF/F_0 values across all bins.

Statistical analysis

Sample sizes are represented in all figures as the number of mice included in the analysis. Statistical

analysis was conducted using Prism V9.0 software (GraphPad Software, Inc.). For unpaired data, the

non-parametric Mann-Whitney (M-W) test was employed, while one-way balanced ANOVA followed by

Dunnet's multiple-comparison test was used for comparisons among multiple groups. Cumulative

distributions were compared using the Kolmogorov-Smirnov (KS) test. The significance level was set at

$P < 0.05$, denoted as $*P < 0.05$, $**P < 0.01$, and $***P < 0.001$. Additionally, when the P-value fell between

0.05 and 0.1, it was labeled accordingly. Error bars in all figures represent the standard error of the

mean (s.e.m).

References

1.   Kalia LV, Lang AE. Parkinson's disease. The Lancet. 2015;386(9996):896-912.

2.   Sveinbjornsdottir S. The clinical symptoms of Parkinson's disease. Journal of Neurochemistry.
2016;139(S1):318-24.

3.   Dudman JT, Krakauer JW. The basal ganglia: from motor commands to the control of vigor.
Current Opinion in Neurobiology. 2016;37:158-66.

4.   Turner RS, Desmurget M. Basal ganglia contributions to motor control: a vigorous tutor. Current
Opinion in Neurobiology. 2010;20(6):704-16.

5.   Lemke SM, Ramanathan DS, Darevksy D, Egert D, Berke JD, Ganguly K. Coupling between
motor cortex and striatum increases during sleep over long-term skill learning. Elife. 2021;10.

6.   Santos FJ, Oliveira RF, Jin X, Costa RM. Corticostriatal dynamics encode the refinement of
specific behavioral variability during skill learning. Elife. 2015;4:e09423.

7.   Lemke SM, Ramanathan DS, Guo L, Won SJ, Ganguly K. Emergent modular neural control
drives coordinated motor actions. Nat Neurosci. 2019;22(7):1122-31.

8.   Gao LL, Wu T. The study of brain functional connectivity in Parkinson's disease. Transl
Neurodegener. 2016;5:18.

9.   Tessitore A, Cirillo M, De Micco R. Functional Connectivity Signatures of Parkinson's Disease. J
Parkinsons Dis. 2019;9(4):637-52.

10. Filippi M, Sarasso E, Agosta F. Resting-state Functional MRI in Parkinsonian Syndromes. Mov
Disord Clin Pract. 2019;6(2):104-17.

11. Lang S, Yoon EJ, Kibreab M, Kathol I, Cheetham J, Hammer T, et al. Mild behavioral impairment
in Parkinson's disease is associated with altered corticostriatal connectivity. Neuroimage Clin.
2020;26:102252.

12. Mi TM, Zhang W, Li Y, Liu AP, Ren ZL, Chan P. Altered Functional Segregated Sensorimotor,
Associative, and Limbic Cortical-Striatal Connections in Parkinson's Disease: An fMRI Investigation.
Front Neurol. 2021;12:720293.

13. Rothkirch I, Granert O, Knutzen A, Wolff S, Gövert F, Pedersen A, et al. Dynamic causal
modeling revealed dysfunctional effective connectivity in both, the cortico-basal-ganglia and the
cerebello-cortical motor network in writers' cramp. Neuroimage Clin. 2018;18:149-59.

14. Unschuld PG, Joel SE, Liu X, Shanahan M, Margolis RL, Biglan KM, et al. Impaired cortico-
striatal functional connectivity in prodromal Huntington's Disease. Neurosci Lett. 2012;514(2):204-9.

15. DeLong MR. Primate models of movement disorders of basal ganglia origin. Trends Neurosci.
1990;13(7):281-5.

16. McGregor MM, Nelson AB. Circuit Mechanisms of Parkinson's Disease. Neuron.
2019;101(6):1042-56.

17.   Kravitz AV, Tye LD, Kreitzer AC. Distinct roles for direct and indirect pathway striatal neurons in
reinforcement. Nat Neurosci. 2012;15(6):816-8.

18.   Surmeier DJ, Ding J, Day M, Wang Z, Shen W. D1 and D2 dopamine-receptor modulation of
striatal glutamatergic signaling in striatal medium spiny neurons. Trends Neurosci. 2007;30(5):228-35.

19.   Tecuapetla F, Jin X, Lima SQ, Costa RM. Complementary Contributions of Striatal Projection
Pathways to Action Initiation and Execution. Cell. 2016;166(3):703-15.

20.   Jin X, Tecuapetla F, Costa RM. Basal ganglia subcircuits distinctively encode the parsing and
concatenation of action sequences. Nat Neurosci. 2014;17(3):423-30.

21.   Lee K, Masmanidis SC. Aberrant features of in vivo striatal dynamics in Parkinson's disease. J
Neurosci Res. 2019;97(12):1678-88.

22.   Sharott A, Vinciati F, Nakamura KC, Magill PJ. A Population of Indirect Pathway Striatal
Projection Neurons Is Selectively Entrained to Parkinsonian Beta Oscillations. J Neurosci.
2017;37(41):9977-98.

23.   Ketzef M, Spigolon G, Johansson Y, Bonito-Oliva A, Fisone G, Silberberg G. Dopamine Depletion
Impairs Bilateral Sensory Processing in the Striatum in a Pathway-Dependent Manner. Neuron.
2017;94(4):855-65 e5.

24.   Mallet N, Ballion B, Le Moine C, Gonon F. Cortical inputs and GABA interneurons imbalance
projection neurons in the striatum of parkinsonian rats. J Neurosci. 2006;26(14):3875-84.

25.   Parker PR, Lalive AL, Kreitzer AC. Pathway-Specific Remodeling of Thalamostriatal Synapses in
Parkinsonian Mice. Neuron. 2016;89(4):734-40.

26.   Parker JG, Marshall JD, Ahanonu B, Wu YW, Kim TH, Grewe BF, et al. Diametric neural
ensemble dynamics in parkinsonian and dyskinetic states. Nature. 2018;557(7704):177-82.

27.   Ryan MB, Bair-Marshall C, Nelson AB. Aberrant Striatal Activity in Parkinsonism and Levodopa-
Induced Dyskinesia. Cell Rep. 2018;23(12):3438-46.e5.

28.   Bloem BR, Okun MS, Klein C. Parkinson's disease. Lancet. 2021;397(10291):2284-303.

29.   Luk KC, Kehm V, Carroll J, Zhang B, O'Brien P, Trojanowski JQ, et al. Pathological α-synuclein
transmission initiates Parkinson-like neurodegeneration in nontransgenic mice. Science.
2012;338(6109):949-53.

30.   Paumier KL, Luk KC, Manfredsson FP, Kanaan NM, Lipton JW, Collier TJ, et al. Intrastriatal
injection of pre-formed mouse α-synuclein fibrils into rats triggers α-synuclein pathology and bilateral
nigrostriatal degeneration. Neurobiol Dis. 2015;82:185-99.

31.   Long H, Zheng W, Liu Y, Sun Y, Zhao K, Liu Z, et al. Wild-type α-synuclein inherits the structure
and exacerbated neuropathology of E46K mutant fibril strain by cross-seeding. Proceedings of the
National Academy of Sciences. 2021;118(20).

32.   Brismar T. The human EEG--physiological and clinical studies. Physiol Behav. 2007;92(1-2):141-
7.

33.   Thompson JA, Lanctin D, Ince NF, Abosch A. Clinical implications of local field potentials for
understanding and treating movement disorders. Stereotact Funct Neurosurg. 2014;92(4):251-63.

34.   Lee MH, Smyser CD, Shimony JS. Resting-state fMRI: a review of methods and clinical
applications. AJNR Am J Neuroradiol. 2013;34(10):1866-72.

35.   Zhou ZC, Gordon-Fennell A, Piantadosi SC, Ji N, Smith SL, Bruchas MR, et al. Deep-brain
optical recording of neural dynamics during behavior. Neuron. 2023;111(23):3716-38.

36.   Tecuapetla F, Matias S, Dugue GP, Mainen ZF, Costa RM. Balanced activity in basal ganglia
projection pathways is critical for contraversive movements. Nat Commun. 2014;5:4315.

37.   Cui G, Jun SB, Jin X, Pham MD, Vogel SS, Lovinger DM, et al. Concurrent activation of striatal
direct and indirect pathways during action initiation. Nature. 2013;494(7436):238-42.
38.   Shin JH, Kim D, Jung MW. Differential coding of reward and movement information in the
dorsomedial striatal direct and indirect pathways. Nat Commun. 2018;9(1):404.
39.   Bell PT, Gilat M, O'Callaghan C, Copland DA, Frank MJ, Lewis SJ, et al. Dopaminergic basis for
impairments in functional connectivity across subdivisions of the striatum in Parkinson's disease. Hum
Brain Mapp. 2015;36(4):1278-91.
40.   Hensel L, Hoffstaedter F, Caspers J, Michely J, Mathys C, Heller J, et al. Functional Connectivity
Changes of Key Regions for Motor Initiation in Parkinson's Disease. Cereb Cortex. 2019;29(1):383-
96.
41.   Long H, Zheng W, Liu Y, Sun Y, Zhao K, Liu Z, et al. Wild-type α-synuclein inherits the structure
and exacerbated neuropathology of E46K mutant fibril strain by cross-seeding. Proc Natl Acad Sci U
S A. 2021;118(20).
42.   Crompe B, Aristieta A, Leblois A, Elsherbiny S, Boraud T, Mallet NP. The globus pallidus
orchestrates abnormal network dynamics in a model of Parkinsonism. Nat Commun. 2020;11(1):1570.
43.   Perlbarg V, Lambert J, Butler B, Felfli M, Valabrègue R, Privat AL, et al. Alterations of the
nigrostriatal pathway in a 6-OHDA rat model of Parkinson's disease evaluated with multimodal MRI.
PLoS One. 2018;13(9):e0202597.
44.   Beck MH, Haumesser JK, Kuhn J, Altschuler J, Kuhn AA, van Riesen C. Short- and long-term
dopamine depletion causes enhanced beta oscillations in the cortico-basal ganglia loop of
parkinsonian rats. Exp Neurol. 2016;286:124-36.
45.   Evangelisti S, Pittau F, Testa C, Rizzo G, Gramegna LL, Ferri L, et al. L-Dopa Modulation of Brain
Connectivity in Parkinson's Disease Patients: A Pilot EEG-fMRI Study. Front Neurosci. 2019;13:611.
46.   Costa VD, Tran VL, Turchi J, Averbeck BB. Dopamine modulates novelty seeking behavior during
decision making. Behav Neurosci. 2014;128(5):556-66.
47.   Kesner AJ, Calva CB, Ikemoto S. Seeking motivation and reward: Roles of dopamine,
hippocampus, and supramammillo-septal pathway. Prog Neurobiol. 2022;212:102252.
48.   Chinta SJ, Andersen JK. Dopaminergic neurons. Int J Biochem Cell Biol. 2005;37(5):942-6.
49.   Wu T, Liu J, Zhang H, Hallett M, Zheng Z, Chan P. Attention to Automatic Movements in
Parkinson's Disease: Modified Automatic Mode in the Striatum. Cereb Cortex. 2015;25(10):3330-42.
50.   Wu T, Wang L, Hallett M, Chen Y, Li K, Chan P. Effective connectivity of brain networks during
self-initiated movement in Parkinson's disease. Neuroimage. 2011;55(1):204-15.
51.   Luo C, Song W, Chen Q, Zheng Z, Chen K, Cao B, et al. Reduced functional connectivity in
early-stage drug-naive Parkinson's disease: a resting-state fMRI study. Neurobiol Aging.
2014;35(2):431-41.
52.   Le W, Dong J, Li S, Korczyn AD. Can Biomarkers Help the Early Diagnosis of Parkinson's
Disease? Neurosci Bull. 2017;33(5):535-42.
53.   Ellmore TM, Castriotta RJ, Hendley KL, Aalbers BM, Furr-Stimming E, Hood AJ, et al. Altered
nigrostriatal and nigrocortical functional connectivity in rapid eye movement sleep behavior disorder.
Sleep. 2013;36(12):1885-92.
54.   Ereira S, Waters S, Razi A, Marshall CR. Early detection of dementia with default-mode network
effective connectivity. Nature Mental Health. 2024.

55. Rossini PM, Miraglia F, Alu F, Cotelli M, Ferreri F, Iorio RD, et al. Neurophysiological Hallmarks of
Neurodegenerative Cognitive Decline: The Study of Brain Connectivity as A Biomarker of Early
Dementia. J Pers Med. 2020;10(2).
56. Varela F, Lachaux JP, Rodriguez E, Martinerie J. The brainweb: phase synchronization and large-
scale integration. Nat Rev Neurosci. 2001;2(4):229-39.
57. Harris AZ, Gordon JA. Long-range neural synchrony in behavior. Annu Rev Neurosci.
2015;38:171-94.
58. Gatev P, Darbin O, Wichmann T. Oscillations in the basal ganglia under normal conditions and in
movement disorders. Mov Disord. 2006;21(10):1566-77.
59. Ashby FG, Turner BO, Horvitz JC. Cortical and basal ganglia contributions to habit learning and
automaticity. Trends in Cognitive Sciences. 2010;14(5):208-15.
60. Wurster CD, Graf H, Ackermann H, Groth K, Kassubek J, Riecker A. Neural correlates of rate-
dependent finger-tapping in Parkinson's disease. Brain Struct Funct. 2015;220(3):1637-48.
61. Bologna M, Leodori G, Stirpe P, Paparella G, Colella D, Belvisi D, et al. Bradykinesia in early and
advanced Parkinson's disease. J Neurol Sci. 2016;369:286-91.
62. Tessitore A, Giordano A, De Micco R, Russo A, Tedeschi G. Sensorimotor connectivity in
Parkinson's disease: the role of functional neuroimaging. Front Neurol. 2014;5:180.
63. Chen Z, Zhang ZY, Zhang W, Xie T, Li Y, Xu XH, et al. Direct and indirect pathway neurons in
ventrolateral striatum differentially regulate licking movement and nigral responses. Cell Rep.
2021;37(3):109847.
64. Sheng MJ, Lu D, Shen ZM, Poo MM. Emergence of stable striatal D1R and D2R neuronal
ensembles with distinct firing sequence during motor learning. Proc Natl Acad Sci U S A.
2019;116(22):11038-47.
65. Burke DA, Rotstein HG, Alvarez VA. Striatal Local Circuitry: A New Framework for Lateral
Inhibition. Neuron. 2017;96(2):267-84.
66. Obeso JA, Marin C, Rodriguez-Oroz C, Blesa J, Benitez-Temiño B, Mena-Segovia J, et al. The
basal ganglia in Parkinson's disease: current concepts and unexplained observations. Ann Neurol.
2008;64 Suppl 2:S30-46.
67. Escande MV, Taravini IR, Zold CL, Belforte JE, Murer MG. Loss of Homeostasis in the Direct
Pathway in a Mouse Model of Asymptomatic Parkinson's Disease. J Neurosci. 2016;36(21):5686-98.
68. Chen Y, Hong Z, Wang J, Liu K, Liu J, Lin J, et al. Circuit-specific gene therapy reverses core
symptoms in a primate Parkinson's disease model. Cell. 2023;186(24):5394-410 e18.
69. Obeso JA, Rodriguez-Oroz M, Marin C, Alonso F, Zamarbide I, Lanciego JL, et al. The origin of
motor fluctuations in Parkinson's disease: importance of dopaminergic innervation and basal ganglia
circuits. Neurology. 2004;62(1 Suppl 1):S17-30.
70. Zhai S, Tanimura A, Graves SM, Shen W, Surmeier DJ. Striatal synapses, circuits, and
Parkinson's disease. Curr Opin Neurobiol. 2018;48:9-16.
71. Cramb KML, Beccano-Kelly D, Cragg SJ, Wade-Martins R. Impaired dopamine release in
Parkinson's disease. Brain. 2023;146(8):3117-32.
72. Costa RM, Lin SC, Sotnikova TD, Cyr M, Gainetdinov RR, Caron MG, et al. Rapid alterations in
corticostriatal ensemble coordination during acute dopamine-dependent motor dysfunction. Neuron.
2006;52(2):359-69.

73. Herz DM, Haagensen BN, Christensen MS, Madsen KH, Rowe JB, Løkkegaard A, et al.
Abnormal dopaminergic modulation of striato-cortical networks underlies levodopa-induced
dyskinesias in humans. Brain. 2015;138(Pt 6):1658-66.
74. Tozzi A, Sciaccaluga M, Loffredo V, Megaro A, Ledonne A, Cardinale A, et al. Dopamine-
dependent early synaptic and motor dysfunctions induced by alpha-synuclein in the nigrostriatal
circuit. Brain. 2021;144(11):3477-91.
75. Burke RE, O'Malley K. Axon degeneration in Parkinson's disease. Exp Neurol. 2013;246:72-83.
76. Zong FJ, Min X, Zhang Y, Li YK, Zhang XT, Liu Y, et al. Circadian time- and sleep-dependent
modulation of cortical parvalbumin-positive inhibitory neurons. EMBO J. 2023;42(3):e111304.
77. Kato HK, Asinof SK, Isaacson JS. Network-Level Control of Frequency Tuning in Auditory Cortex.
Neuron. 2017;95(2):412-23 e4.
78. Hu H, Gan J, Jonas P. Interneurons. Fast-spiking, parvalbumin(+) GABAergic interneurons: from
cellular design to microcircuit function. Science. 2014;345(6196):1255263.
79. Mallet N, Le Moine C, Charpier S, Gonon F. Feedforward inhibition of projection neurons by fast-
spiking GABA interneurons in the rat striatum in vivo. J Neurosci. 2005;25(15):3857-69.
80. Gittis AH, Kreitzer AC. Striatal microcircuitry and movement disorders. Trends Neurosci.
2012;35(9):557-64.
81. Chen H, Lei H, Xu Q. Neuronal activity pattern defects in the striatum in awake mouse model of
Parkinson's disease. Behav Brain Res. 2018;341:135-45.

Figures and Legends

Figure 1. Detection of task-specific cortico-striatal coupling via dual fiber photometry imaging

A. Scheme of virus injection and optic fiber implantation.

B. Anatomical verification of virus expression and optic fiber localization. Scale bar: 200μm.

C-N. Examination of the cortico-striatal couplings during contralateral turning (C-F), licking (G-J), and

digging (K-N).

(C, G, K) Behavioral diagrams (top) and example Ca^{2+} activities recorded from the primary motor cortex

(M1, red) and the striatum (STR, green) with dotted lines denoted onsets of the specified behaviors.

Ca^{2+}-independent fluorescence was monitored using 405 nm light pulses for individual channel (purple,

bottom).

(D, H, L) Heatmaps showing all compiled trials (Top panels) and the averaged Ca^{2+} signal intensities

(dF/F_0, bottom panels) aligned (dotted lines) to the initiation of the contralateral turning (D), licking (H),

and digging (L). All left panels, M1 signal; all right panels, STR signals.

(E, I, M) Temporally aligned traces showing z-scored average signal intensities of both M1 (red) and

STR (green).

(F, J, N) Calculated cortico-striatal coupling indices for contralateral turning (F), licking (J), and digging

(N).

Figure 2. Task- and cell type-specific cortico-striatal coupling

A. Schematic diagram showing the experimental design for analyzing cell type-specific cortico-striatal

coupling.

B. Immunofluorescence images showing the colocalization between DRD1 (red) and DIO-GCaMP

(green) in D1R-Cre mice (top panels), and the colocalization between DRD2 (magenta) and DIO-

GCaMP (green) in the D2R-Cre mice (bottom panels). Scale bar: 10µm.

C, E, G. Z-scored average traces of M1 (red) and STR dMSN (light green) and iMSN (dark green) Ca^{2+}

activities temporally aligned to the initiation of contralateral turning (C), licking (E), and digging (G).

D, F, H. Comparison of cell type-specific cortico-striatal coupling indices for turning (D), licking (F), and

digging behavior (H, P = 0.0087). Mann-Whitney test. **P<0.01.

Figure 3. Pathological characterization of α-synuclein preformed fibril (PFF) induced PD-like

mouse model

A. Schematic diagram describing the bilateral inoculation of PFF into striatum of wildtype mice. Right

panel, transmitted electron microscope image of PFF. Scale bar: 100 nm.

B. Experimental timeline. IHC: Immunohistochemistry.

C-D. Behavioral analysis of the PFF- and PBS-injected mice 1 (C) and 3 (D) months post injection (Mpi).

(C_1) Learning curve of rotarod running.

(C_2) Latency to fall in the rotarod test.

(C_3) Total travel distance in the open field test.

(D_1) Latency to fall in the rotarod test 3Mpi (P = 0.0207).

(D_2) Total travel distance in the open field test 3Mpi (P = 0.0147).

E-K. Immunofluorescence analyses of pathological hallmarks in PFF- and PBS-injected mice 1Mpi. All

data normalized to PBS control.

(E-G) Comparison of the p-α-Syn intensity (F, P = 0.0079) and dopamine transporter (DAT) signal

intensities (G) in STR. E, Representative immunofluorescence images. Scale bars: 50 μm.

(H-K) Comparison of the p-α-Syn intensity (I, P = 0.0043), DA neuron number (J, P = 0.0022) and DAT

intensity (K) in substantia nigra (SN). H, Representative immunofluorescence images. Scale bars: 250

μm.

L-R. Immunofluorescence analyses of pathological hallmarks in PFF- and PBS-injected mice 3Mpi. All

data normalized to PBS control.

(L-N) Comparison of the p-α-Syn intensity (M, P = 0.0022) and dopamine transporter (DAT) signal

intensities (N, P =0.0411) in STR. L, Representative immunofluorescence images. Scale bars: 100 μm.

(O-R) Comparison of the p-α-Syn intensity (P, P =0.0022), DA neuron number (Q, P = 0.0079) and DAT

intensity (R) in SN. O, Representative immunofluorescence images. Scale bars: 250 μm.

Mann-Whitney test, unless otherwise noted. Two-way ANOVA with Sidak's multiple comparisons test

(C_1). *P<0.05, **P<0.01.

Figure 4. Task-specific cortico-striatal decoupling in PD-like mice

A. Experimental timeline.

B-I, Dual fiber photometry imaging of M1 and STR in PFF- and PBS-injected mice at 1 Mpi (B-C), 2 Mpi

(D-E), and 3-4 Mpi (F-I) during digging.

(B, D, F) Z-scored average traces of M1 (red) and STR (green) Ca^{2+} activities temporally aligned to the

initiation of digging. Left, PBS; Right, PFF.

(C, E, G) Quantitative analyses of the cortico-striatal coupling.

(C_1, E_1, G_1) Coupling indices (E_1, $P = 0.0649$. G_1, $P = 0.0037$).

(C_2, E_2, G_2) Phase lags (E_2, $P = 0.0281$. G_2, $P = 0.0082$).

(C_3, E_3, G_3) Cumulative distribution of phase lag per trial (C_3: PBS = 140 trials, PFF = 124 trials. E_3:

PBS = 109 trials, PFF = 104 trials; $P = 0.0041$. G_3: PBS = 102 trials, PFF = 118 trials; $P = 0.0009$.

Kolmogorov-Smirnov test). Inserts, Comparison of the trial average phase lag (In C_3, $P = 0.0188$. In E_3,

$P = 0.0072$. In G_3, $P = 0.0110$).

(H) Plateau amplitude of M1 (left) and STR (right) Ca^{2+} activities at 3~4Mpi.

(I) The temporal profiles of coupling index (left) and phase lag (right).

J-K. Dual fiber photometry imaging of M1 and STR at 4 Mpi during licking.

(J) Z-scored average traces of M1 (red) and STR (green) Ca^{2+} activities temporally aligned to the

initiation of licking.

(K) Comparison of the coupling indices.

(L) Comparison of the phase lags.

Mann-Whitney test, unless otherwise noted. *$P<0.05$. **$P<0.01$. ***$P<0.001$.

Figure 5. Digging-associated cortico-striatal decoupling correlated with specific behavioral

deficits

A. Duration of a single digging event. Left, the temporal profiles of average digging duration. Right,

comparison of the duration at 3~4Mpi (*P* = 0.0022).

B. Frequency of digging events. Left, temporal profiles of digging frequency. Right, comparison of the

frequency at 3~4Mpi (*P* = 0.0152).

C, D. Correlation analysis of phase lag between digging frequency (C) and duration (D) from PBS- and

PFF-injected mice at 3~4Mpi. r: Pearson's correlation coefficient. Each dot represents data from an

individual mouse (the same below).

E, F. Correlation analysis of coupling index between digging frequency (E) and duration (F) from PBS-

and PFF-injected mice at 3~4Mpi.

Mann-Whitney test. *P<0.05. **P<0.01.

Figure 6. Cell type-specific contribution of digging-related cortico-striatal decoupling in PFF-

injected mice

A-E. Dual fiber photometry imaging of M1 and striatal dMSN in PFF- and PBS-injected mice at 2 Mpi

(B-C), and 3-4 Mpi (D-E) during digging.

(A) Scheme of virus injection and optic fiber implantation. Note the CreON-GCaMP virus was injected

into the STR of D1R-Cre mice, while CreOFF-GCaMP virus was injected into the STR of D2R-Cre mice.

Right panel, experiment timeline.

(B, D) Z-scored average traces of M1 (red) and STR dMSN (aqua) Ca^{2+} activities temporally aligned to

the initiation of digging. Left, PBS; Right, PFF.

(C, E) Quantitative analyses of the M1-dMSN coupling and plateau amplitude of dMSN Ca^{2+} activities

at 2 Mpi (C) and 3-4 Mpi (E). The hollow signal represents D1R-Cre mice; the cross signal represents

D2-Cre mice.

(C_1, E_1) Coupling indices (C_1, P = 0.0043. E_1, P = 0.0062).

(C_2, E_2) Phase lags (E_2, P = 0.0043. G_2, P = 0.0062).

(C_3, E_3) Plateau amplitude of dMSN Ca^{2+} activities (E_3, P = 0.0451).

F-J. Dual fiber photometry imaging of M1 and striatal iMSN in PFF- and PBS-injected mice at 2 Mpi (G-

H), and 3-4 Mpi (I-J) during digging. Right panel, experiment timeline.

(F) Scheme of virus injection and optic fiber implantation.

(G, I) Z-scored average traces of M1 (red) and striatal iMSN (dark green) Ca^{2+} activities temporally

aligned to the initiation of digging. Left, PBS; Right, PFF.

(H, J) Quantitative analyses of the M1-iMSN coupling and plateau amplitude of iMSN Ca^{2+} activities at

2 Mpi (H) and 3-4 Mpi (J).

(H_1, J_1) Comparison of the coupling indices.

(H_2, J_2) Comparison of the phase lags.

(H_3, J_3) Comparison of the plateau amplitude of iMSN Ca^{2+} activities.

Mann-Whitney test. *P<0.05. **P<0.01.

Figure 7 Cortico-striatal decoupling in PFF-injected mice accompanied by deficiency of striatal

extracellular dopamine

A. Schematic diagram showing the experimental design for assessing striatal extracellular dopamine

(DA) level. DA4.4 is a florescence-based DA sensor. Right panel, anatomical verification of virus

expression and optic fiber localization. Scale bar: 200μm.

B. Experimental timeline.

C. Schematic diagram for free-exploring paradigm.

D. Representative magnitude heatmap of normalized (dF/F_0) of DA sensor activity 3~4 weeks post-

injection (wpi), 6~7wpi, 10wpi, and 13~14wpi from one PBS-injected mouse and one PFF-injected

mouse during 2-minute free-exploring.

E. Normalized sum activity of DA sensor during free-exploring from PBS- and PFF-injected mice 3~4wpi,

6~7wpi, 10wpi, 13~14wpi. Data batch-normalized to the PBS group (the same below).

F.Schematic diagram for treadmill running paradigm.

G. Representative magnitude heatmap of normalized (dF/F_0) of DA sensor activity 3~4wpi, 6~7wpi,

10wpi, and 13~14wpi from same two mice as in (D) during 2-minute treadmill running.

921   **H**. Normalized sum activity of DA sensor during free-exploring from PBS- and PFF-injected mice at

922   3~4wpi (P = 0.0007), 6~7wpi, 10wpi, 13~14wpi (P = 0.0043).

923   Mann-Whitney test. **P<0.01. ***P<0.001.

Figure 8 Selective activation to D1R rescues both digging-associated cortico-striatal decoupling

and digging behavioral defects

A. Schematic diagram for pharmacological experiments. i.p., intraperitoneal administration.

B-E. Dual fiber photometry imaging of M1 and STR in PFF- and PBS-injected mice at 4Mpi during

digging, after ip. SAL (saline), L-dopa, SKF (SKF-81297, a D1R selective agonist), or Quin (quinpirole,

a D2R selective agonist).

(B) Z-scored average traces of M1 (red) and STR (green) temporally aligned, digging-associated Ca^{2+}

activities after administration of drugs. Left, PBS; Right, PFF.

(C, D, E) Quantitative analyses of the digging-associated cortico-striatal coupling (C,D) and   amplitude

of striatal Ca^{2+} activities (E) after administration of drugs.

(C$_1$) Comparison of the phase lags (SAL-PBS *vs.* SAL-PFF: Mann-Whitney test, $P = 0.0152$. PFF

groups: SAL-PFF as control, $F_{(1.588, 9.529)} = 9.270$, SAL-PFF *vs.* Ldopa-PFF, $P = 0.0156$; SAL-PFF vs.

SKF-PFF, $P = 0.0328$).

(C$_2$) Comparison of the phase lags, after saline or quinpirole administration.

(D$_1$) Comparison of the coupling indices (SAL-PBS *vs.* SAL-PFF: Mann-Whitney test, $P = 0.0043$).

(D$_2$) Comparison of the coupling indices, after saline or quinpirole administration.

(E$_1$) Comparison of the plateau amplitude of striatal activity (SAL-PBS *vs.* SAL-PFF: Mann-Whitney test,

$P = 0.0424$. PFF groups: SAL-PFF as control, $F_{(1.453, 8.719)} = 6.152$, SAL-PFF vs. SKF-PFF, $P = 0.0459$).

(E$_2$) Comparison of the plateau amplitude of striatal activity, after saline or quinpirole administration.

F-G. Effect of dopaminergic drugs on digging behavior.

(F) Comparison of digging duration (PFF groups: SAL-PFF as control, $F_{(1.463, 7.317)} = 19.97$, SAL-PFF *vs.*

Ldopa-PFF, $P = 0.0018$; SAL-PFF *vs.* SKF-PFF, $P = 0.0388$).

(G) Comparison of digging frequency (PFF groups: SAL-PFF as control, $F_{(2.120, 10.60)} = 11.78$, SAL-PFF

vs. SKF-PFF, $P = 0.0143$; SAL-PFF *vs.* Quin-PFF, $P = 0.0192$).

One-way ANOVA followed by Dunnett's multiple comparisons test, unless otherwise noted. Wilcoxon

matched-pairs test (C$_2$, D$_2$, E$_2$). *$P<0.05$. **$P<0.01$.