Monsoonal climate and asymmetric heating facilitate the slope-aspect to influence

landscape-scale tree structure in the Western Ghats, India

Devi Maheshwori[1], Shreyas Managave[1*], Girish Jathar[2,3], Sham Davande[4]

[1] Indian Institute of Science Education and Research Pune, India

[2] Srushti Conservation Foundation, Mahatma Society, Pune, Maharashtra, India

[3] Advance Research Centre for Ecology and Conservation, DES Pune University, Pune,
Maharashtra, India

[4] GIS Consultant, West Vinod Nagar, Delhi, India

[*] Corresponding Author: shreyas@iiserpune.ac.in

Abstract

The regional variation of rainfall and length of the rainy season are identified to control
the vegetation distribution in the Western Ghats (WG) of India, one of the important
biodiversity hotspots. Here we demonstrate the existence of two new modes related to
monsoonal climate and asymmetric solar heating that influence the vegetation structure at the
landscape-scale in the WG: (i) the north-facing slopes have a higher tree cover (TC) and canopy
height (CH) than the south-facing slopes and (ii) higher TC and CH on the west-facing slopes
than the east-facing slopes. The asymmetry associated with these modes increases with the
slope angle. The interaction of these modes often leads northwest and southeast aspects to have
the highest and lowest TC (and CH), respectively. We thus demonstrate that even in low-
latitude regions, the slope-aspect plays an important role in determining the TC and CH if the
relief is higher. This fundamental driver of vegetation structure needs to be considered while
formulating and executing programs aimed at increasing tree cover and conserving
biodiversity, especially in the high relief areas in the tropics having seasonal rainfall, such as
in the WG.

Key words: tree cover, canopy height, aspect, vegetation, monsoon, sun angle, tropics

1. Introduction

The Western Ghats (WG) constitutes a sequence of hills running parallel to the west
coast of India for about 1600 km (from 8°N to ~21°N) and covering an area of ~160000 km^2
(Das et al., 2006). It is a world heritage site and one of the important biodiversity hotspots
having high species endemicity (Myers et al., 2000). Understanding the factors controlling
vegetation distribution in the WG is crucial for the conservation of biodiversity and endemic
species. The length of the rainy season and total rainfall has been attributed to control the
regional scale vegetation distribution in the WG (Champion and Seth 1968; Pascal 1988;
Gadgil, 1996; Gimaret and Carpentier, 2003; Nagendra and Ghate, 2003; Prasad et al., 2008;
Ramachandra et al., 2016). However, the factors influencing vegetation distribution at
landscape-scale within individual protected areas (PAs) are not appreciated.

The slope-aspect is one of the important drivers of vegetation structure at the landscape-
scale (Singh, 2018). The effect slope-aspect results in the equator-facing slopes having a
relatively warmer, drier microclimate and soil with lower moisture content than the pole-facing
slopes (Geroy et al., 2011; Gutiérrez-Jurado et al., 2013; Zhou et al., 2013). This in conjunction
with soil edaphic factors leads to the North-facing slopes having a higher vegetation cover than
the South-facing slopes in the Northern Hemisphere; the asymmetry being higher on steeper
slopes (Smith and Bookhagen, 2021). The north-south asymmetry in the vegetation cover thus
established is common to mid-latitude regions (e.g. Champion and Seth 1968; Holland and
Steyn, 1975; Badano et al., 2005; Singh, 2018; Smith and Bookhagen, 2021); its influence in
the WG, however, is not recognised.

The influence of slope-aspect on the vegetation distribution in the WG was realized
only in terms of regional west-east asymmetry in the vegetation distribution. This asymmetry
was attributed to the interaction of the west/south-westerly monsoonal winds with the WG
which causes higher rainfall in the windward western part (i.e. area lying west of the crest line
of the WG hills) than in the leeward eastern part (Fig. 1). This orographic effect controls the
regional distribution of the vegetation: evergreen forests on the western side and deciduous
forests on the eastern side of the WG (Champion and Seth 1968; Pascal 1988; Gadgil, 1996).
The existence of north-south and west-east asymmetries and their interaction in controlling
vegetation structure at the landscape-scale is not known. Here we test whether the slope-aspect
controls the tree structure, viz. tree cover (TC) and canopy height (CH), at the landscape-scale
in the WG.

Figure 1. The topography of the WG (based on ASTER Global Digital Elevation Model) (left panel). The arrows indicate the direction of the Low-level winds (925 hPa) (Viswanadhapalli et al., 2020). Distribution of rainfall (Pai et al., 2014) and the length of the rainy season (shown by contours) (Singh et al., 1986) in the WG (right panel). The regions highlighted in red colour (right panel) are protected areas (PAs) analysed for their TC and CH. Northern (NWG), Central (CWG) and Southern (SWG) parts of the WG are demarcated by horizontal dashed lines. The notations for the PAs: a- Tamhini-Lavasa-Raigad; b- Koyna; c-Chandoli; d-Vishalgad; e-'West of the WG Escarpment'; f-Radhangiri; g-Bhagwan Mahavir; h-Netravali; i-Anshi; j-Cotiago; k-Kudremukh; l-Pushpagiri; m-Talakaveri; n-New Brahmagiri; o-Brahmagiri; p-Aralam; q-New Amarambalam-Silent Valley; r-Peechi; s-Chimmony; t-Parambikulun; u-Indira Gandhi; v-Idukki; w-Shendurney; x-Kallakad Mundanthurai; y-Neyyar.

2. Material and Methods

2.1. Regional Setting

The WG acts as an orographic barrier to the south-westerly monsoonal winds during

southwest monsoon (SWM) season (June to September) which results in higher and lower

rainfall on the windward (western margin) and leeward (eastern margin) sides of the WG,

respectively (Gunnell, 1997) (Fig. 1). The southern part of WG receives additional rains during

the northeast monsoon (NEM) season (October to December) (Fig 1, Suppl. Fig. S1). This, in

conjunction with an earlier onset and late withdrawal of SWM, leads the southern part of the

WG to have a longer rainy season (~260 days) as compared to that in the northern part (~130

days) (Fig. 1) (Singh, 1986).

The seasonal amplitude of temperature is higher in the northern Western Ghats (NWG)

than in the central and southern Western Ghats (CWG and SWG, respectively) (Suppl. Fig.

S1). In the WG, December and January are the coldest months (~23ºC); April and May are the

warmest (~30ºC in the NWG, and ~27ºC in the CWG and SWG) (Suppl. Figure S1). The lapse

rate of average annual temperature observed in the southern part of WG varied from 0.5° to

0.7º/100 m (von Lengerke,1977). At high altitude PAs (2000-2500 m) the average monthly

temperature varies from 9.5 (January) to 18.7 (April) (von Lengerke,1977).

To assess the effect of aspect on TC and CH we studied vegetation in 25 protected areas

(PAs) distributed throughout the WG (Fig. 1). This study covers ~8400 km^2 of area; different

aspect categories cover 9 to 15% of it (Suppl. File 1 and Fig. S2). About ~62 % of the area in

these PAs exhibits a slope between 10° to 30°, followed by 0° to 10° (~26 %), 30° to 50° (~11

%) and 50° to 90° (~1 %). Within individual PAs the area covered by slope 10° to 30° is the

most abundant (Suppl. Fig. S3). The median elevation of the PAs varies from 143 to 1346 m;

only two PAs (indicated by letters *q* and *u* in Figure 1) have elevations higher than 2000 m.

The higher altitudes, especially in the CWG and SWG, were associated with steeper slopes

(Suppl. Fig. S4). The high-altitude regions often show a mosaic of trees and grasslands, the

shola landscape (Meher-Homji, 1967).

2.2. Data sets

To demonstrate the influence of slope-aspect on the tree structure in the WG, we used

the Hansen Global Forest Change v1.9 (2000-2021) (Hansen et al., 2013) and Global Forest

Canopy Height, GEDI_V27 (Potapov et al., 2021) data to get TC and CH, respectively. Hansen

et al., (2013) defined the tree as vegetation taller than 5m in height and tree cover is expressed

as a percentage per output grid cell of size 30 x 30 meters. Potapov et al., (2021), who

considered the tree as vegetation taller than 3m in height, integrated lidar and Landsat analysis-

ready data to produce CH data of 30m resolution. SRTM DEM data (30 m resolution) was used

to characterize topography.

2.3. Methodology

To avoid the artifacts of anthropogenic activities on the aspect-related asymmetry in

vegetation structure, we restricted our analysis to PAs. The boundaries of PAs were taken from

The Ministry of Environment, Forest and Climate Change, Government of India's Decision

Support System (MoEF & CC-DSS). The boundaries of PAs Tamhini-Lavasa-Raigad, 'West

of the WG Escarpment', New Brahmagiri, New Amarambalam-Silent Valley and Vishalgad

were selected manually. Further, an area below the altitude of 200m was not considered as it

was generally affected by anthropogenic activity. Areas under anthropogenic influences (e.g.

plantations and human settlements), deciphered through Google Earth images, were ignored.

We considered the median of the TC and CH on eight aspects i.e. North (N), Northeast

(NE), East (E), Southeast (SE), South (S), Southwest (SW), West (W) and Northwest (NW),

and three slope angle (0° to 10°, 10° to 30° and 30° to 50°) categories. We did not consider the

area having a slope from 50° to 90°, which was less than 1% of the total area. The effect of

altitude on TC and CH was assessed for the altitude range of 200-500m, 500-900m, 900-1500m

and 1500-2600m. The TC and CH data were overlain on the DEM image in Q-GIS to determine

the TC and CH under different aspect, slope and altitude categories.

3. Results

The regional distribution of TC and CH were characterised by lower TC and CH in the

NWG, and higher TC and CH in CWG and SWG (Suppl. Figure S5). Each PA shows aspect-

related variation in the median TC and CH (Figure 2). The range in the median TC and CH on

various aspects within individual PAs vary from 2 to 32% and 1 to 8 m, respectively. The

distribution of TC (and CH) on different aspects within individual PAs was statistically

different ($p < 0.05$, the two-sample Kolmogorov-Smirnov test).

Figure 2. The median tree cover (left panel) and canopy height (right panel) on different aspect categories in the studied PAs. The abbreviations used for the PAs are as in Figure. 1. NWG, CWG and SWG indicate northern, central and southern Western Ghats, respectively.

Even though, the tree was defined differently in TC and CH datasets (i.e. vegetation >5m in TC; >3m in CH) a positive correlation was observed between the median values of TC and CH (i) of different PAs (Suppl. Fig. S5c) and (ii) on different aspects within individual PA (Suppl. Fig. S6).

3.1. TC and CH asymmetry on N- and S-aspects

The median TC percent and CH on the N-aspect in the individual PAs is higher than that on the S-aspect (Fig. 3a, b), the N-S asymmetry. This is to say that TC is denser and canopy height taller on the N-aspects than on the S-aspects. Similarly, the median TC and CH showed a decreasing trend from the NE- to E- to SE-aspects (Suppl. Fig. S7) and from NW- to W- to SW-aspects (Suppl. Fig. S8). The N-S asymmetry in the TC (and CH) showed dependence on latitude; although this dependence was statistically significant only for TC (Suppl. Fig. S9).

The median TC percent on the N-aspect was similar to that on the W-aspect ($W_{aspect} - N_{aspect} = -1 \pm 5\%$) but the median TC percent on the S-aspect was always lower than on the W-aspect ($W_{aspect} - S_{aspect} = 9 \pm 6\%$) (Figure 3e). A similar pattern was also observed for the CH: $W_{aspect} - N_{aspect} = 1 \pm 1$ m and $W_{aspect} - S_{aspect} = 3 \pm 2$ m (Figure 3f).

Figure 3. The median tree cover (left panels) and canopy height (right panels) on the N-aspect
and S-aspect (a, b) and the W-aspect and E-aspect (c and d) in the studied PAs. The bottom
panels show the difference in the tree cover (e) and canopy height (f) on the W-aspect and N-
aspect, and on W-aspect and S-aspect. The abbreviations ('a' to 'y') used for the PAs are as in
Figure. 1.

3.2. TC and CH asymmetry on W- and E-aspects

We observed higher median TC percent and CH on W-aspect than on the E-aspect (Fig.

3 c and d) (i.e. W-E asymmetry) which was also manifested in decreasing trend in TC and CH

from NW- to N- to NE-aspects (Suppl. Fig. S10) and from SW- to S- to SE-aspects (Suppl.

Fig. S11). The correlations of W-E asymmetry in the TC and CH with latitude were statistically

insignificant (Suppl. Fig. S9 c and d).

3.3. Slope and altitude dependence of the asymmetries in TC and CH

The N-S and W-E asymmetries in the TC and CH increased with the slope angle within

individual PAs (Figure 4). The slope-dependent asymmetry is established by occurrence of

lower TC and CH selectively on the S- and E-aspects on the steeper slopes (Suppl. Fig. S12

and S13).

Figure 4. The slope-dependant variation in the difference between the median tree cover on

the N-aspect and S-aspect (N_{aspect}-S_{aspect}) (a) and the W-aspect and E-aspect (W_{aspect}-E_{aspect})

(b). The same for canopy height is shown in (c) and (d). The abbreviations used for the PAs

are as in Figure. 1. Note, the area covered by slope 50-90° is mostly less than 1%.

As compared to their variation with slope, the N-S and W-E asymmetries in TC and

CH did not show systematic variations with altitude (Suppl. Fig. S14). Some of the PAs show

high values of N-S and W-E asymmetries in TC and CH at 900-1500 m and 1500-2600 m (e.g.

PAs *k*, *l*, *n*, *o* and *v* in Suppl. Fig. S14). These altitude bins were associated with steeper slopes.

Further, the variations in the TC and CH on N-, S-, W- and E-aspects with altitude (Suppl.

Figs. S15 and S16) were more heterogeneous than that with the slope (Suppl. Figs. S12 and
S13).

4. Discussion

The higher median TC in the CWG and SWG than that in the NWG (Suppl. Fig. S5)
reported here is consistent with the earlier finding of the effect of rainfall and the length of the
rainy season on the vegetation distribution (Pascal 1988; Ghate et al., 1998; Nagendra and
Ghate, 2003; Pascal et al., 2004). Higher rainfall and the length of the rainy season facilitate
higher TC in CWG and SWG. The CH is influenced by various factors including hydraulic
limitations (Koch et al., 2004; Moles et al., 2009; Tao et al. 2016; Adrah et al., 2022) and
sunlight availability (Fricker et al., 2019). The drivers of the CH were found to be scale-
dependent: the climate (e.g. water deficit, precipitation) dominates CH at all scales (25 to 1000
m), the relative importance of topography-induced variables (e.g. slope and sunlight exposure)
becomes important at fine scale (25-100 m) (Fricker et al., 2019). Here, we observed a positive
correlation between TC and CH on coarser (inter-PAs, Suppl. Fig. S5c) and finer (intra-PA,
scales between median TC and CH on 8 aspect categories, Suppl. Fig. S6) suggesting similar
drivers of TC and CH in the WG. At a landscape scale, the competition for sunlight might have
also led the CH to increase where tree cover is higher.

4.1. N-S asymmetry: the role of monsoon and solar declination

The TC and CH on the N-aspect were similar to that on the W-aspect (Figure 3 e and
f). However, the TC and CH on the S-aspect were consistently less than that on the W-aspect
suggesting N-S mode is generated by the factors that selectively lower TC and CH on the S-
aspect. This asymmetry was also manifested in decreasing TC and CH from NE- to E- to SE
aspects and from NW- to W- to SW-aspects. We attribute the N-S asymmetry to the differential
heating of the south-facing and north-facing slopes. We illustrate this with the help of
conditions at the northernmost (i.e. Tamhini) and the southernmost (i.e. Neyyar) PAs (Figure
5); the explanation thus also holds for PAs at in-between latitudes.

The duration the Sun spends over the N- and S-aspects is different at these PAs. The
Sun is over the S-aspect for 288 days (~80% of a year, from the 30th of July to the 10th of April)
and 225 days (~62% of a year, from the 1st of September to the 10th of April) at Tamhini and
Neyyar, respectively. On the contrary, the Sun is over the N-aspect for a shorter duration at

Tamhini (~20% of a year i.e. for 78 days, from the 12[th] of May to the 29[th] of July) and Neyyar

(~38% of a year i.e. for 141days, from the 11[th] of April to 30[th] of August).

Figure 5. The angle the Sun makes at its highest position (declination) during summer (21st

June) and winter (21st December) solstice at Tamhini (the northernmost PA) and Neyyar (the

southernmost PA). The figure also shows the SWM and NEM season (shown by shades of blues)

vis-à-vis the maximum Sun angles at their onset and withdrawal; the non-rainy season, which

is generally cloud-free (India Meteorological Department, 2022) is represented by grey colour.

The Sun angle data from www.sunearthtools.com for year 2020.

In addition to the longer duration spend by the Sun over the S-aspect, the monsoonal

climate further facilitates higher heating of the S-aspect. Out of the total days the Sun spends

over the S-aspect, the non-rainy season (with clearer sky) constitutes 223 days (223 out of 288

days; the 1[st] of October to the 11[th] of May) at Tamhini and 100 days (100 out of 225 days; the

1[st] of January to the 10[th] of April) at Neyyar. The lesser length of the dry season at Neyyar is

due to the northeast monsoon (NEM) rainfall (Figure 5). In comparison, when the Sun is over

the N-aspect, only ~29 (out of 78) and ~47 (out of 141) days belong to a non-rainy season at

these PAs, respectively (Figure 5). Even during these non-rainy days when the Sun is over N-

aspect wet and cloudy conditions often prevail due to occurrence the of pre-monsoon showers

(India Meteorological Department, 2022). In summary, the south-facing slopes receive sunlight

for a significantly longer duration as compared to the north-facing slopes; the asymmetry being

higher at Tamhini in the north than at Neyyar located in the south.

The solar radiation load can establish asymmetry in the soil moisture content on north-

facing and south-facing slopes. When the Sun is over the southern slopes, the amount of solar

radiation received by the southern slopes is higher than the northern slopes (Zhou et al., 2013).

In moisture-limited conditions (such as dry season in monsoonal India), the soils on the N-

aspect, receiving lower solar heating, have lesser evaporation losses, and thus have more soil

moisture content than the soil on the S-aspects (Geroy et al., 2011; Gutiérrez-Jurado et al.,

2013; Zhou et al., 2013). The N-aspects also have the potential to retain water for a longer

duration (Geroy et al., 2011). The S-aspect, on the contrary, is more soil-moisture deficient

(Champion and Seth, 1968; Guan et al., 2015; Lewis et al., 2011).

The vegetation productivity in monsoonal India is mainly controlled by the soil

moisture availability (Dubey and Ghosh, 2023), which is the maximum during the rainy season

and decreases rapidly afterward; the lowest soil moisture is observed prior to the onset of

monsoon (Venkates et al., 2011). Direct sunlight exposure of the southern slopes during the

(cloud-free) dry season likely leads to rapid loss of soil moisture on the southern side and can

explain the observed N-S asymmetry in the TC and CH in the WG.

4.2. W-E asymmetry: an effect of orographic rainfall alone?

The orographic rainfall leads to the evergreen and deciduous forests on the western and

eastern slopes of the WG, respectively (Champion and Seth 1968; Pascal 1988; Gadgil, 1996).

Thus the conditions are more conducive for tree growth on the western slopes than the eastern

slopes. The W-E asymmetry in the TC and CH observed here (Fig. 3, 4) also suggested the

influence of the orographic precipitation, especially in PAs that cover the western and eastern

slopes of the WG.

The intense orographic rainfall is restricted till 800 m height on the windward side of

the WG (Tawade and Singh, 2014); a higher amount of rain was observed on the western slope

of the WG and coast plains (Shige et al., 2017; Subrahmanyam and Kumar, 2022). Therefore,

the region lying on the western part of the WG crest line is not expected to show W-E

asymmetry in the orographic precipitation and hence in TC and CH. However, we observed

the W-E asymmetry in TC and CH even over the individual ridges (as low as 400 m in height)

lying in the western part of the WG, for example, in 'West of the WG Escarpment' and

Vishalgad (Figure 1, 2, 6). This alludes to the plausible role of additional factors (such as sea

breeze, the persistence of morning dew, and timing of the peak cloud cover vis-à-vis solar

inclination) (Smith and Bookhagen, 2021) in establishing W-E asymmetry in the vegetation.

In the WG region a higher cloud cover was observed in the afternoon (when the Sun is on the

western side) than in the morning (when the Sun is on the eastern side) during non-rainy days

(from October to May) (Nikumb et al., 2019). This might facilitate the protection of the soil

moisture for a longer duration on the W-aspect post-cessation of the monsoon and lead to the

W-aspect having more TC and CH than the E-aspect.

4.3. The role of slope, latitude and altitude

The N-S asymmetry in the solar insolation increases with the slope angle and latitude

(Smith and Bookhagen, 2021). The slope-dependence of N-S asymmetries in TC and CH (Fig.

4) suggested the role of asymmetric heating of North and South facing aspects in establishing

N-S asymmetry in the vegetation structure in the WG. The increase in the N-S asymmetries in

TC (and to some extent CH) with latitude (Suppl. Fig. 9a, b) further supports this conclusion.

The orographic rainfall is restricted to the windward side of the WG. A long and gentle

rise in topography is associated with a higher orographic rainfall than with a steep rise (Tawade

and Singh, 2014). Therefore, an increase in W-E asymmetry in rainfall with slope is

unexpected, and less likely to be attributed to the slope-dependent increase in W-E asymmetry

in the TC and CH. The slope-dependence of W-E asymmetry in TC and CH is consistent with

the effect of cloud-mediated asymmetric heating of W- and E- aspects (explained earlier). The

W-E asymmetry is established by lower TC and CH selectively on the E-aspect (Suppl. Fig.

S12, 13); steeper E-aspects get heated more in the morning (which are generally cloud-free),

leading to higher soil moisture loss and resulting in its lower TC and CH. Persistence of dew

for a longer duration (due to lower heating) on the western aspects (Smith, 1978) might also

help in retaining higher soil moisture on the W-aspect and support higher TC and CH. As (i)

there is no W-E asymmetry in solar insolation (i.e. radiation in the absence of cloud cover)

(Smith and Bookhagen, 2021) and (ii) cloud-mediated asymmetric heating of W- and E-aspects

is not known to vary with latitude, the W-E asymmetry in TC and CH did not show latitudinal

dependence (Suppl. Fig. 9c, d).

The vegetation in WG showed N-S and W-E asymmetry in TC and CH at all altitude

bins; the extent of asymmetry, however, did not change systematically with altitude (Suppl.

Fig. S14). It thus appears that lapse rate controlled temperature change may not be controlling
the magnitude of N-S and W-E asymmetry in TC and CH in the WG. At lower altitudes (<1500
m), the variation in N-S and W-E asymmetry could be due to the effect of precipitation and
slope. While the western side receives orographic precipitation, the precipitation decreases with
altitude on the eastern side (Figure 1). This makes it difficult to assess whether the variation in
the W-E asymmetry with altitude reported here is the effect of altitude or precipitation. In
addition, the slope increases with altitude in the CWG and SWG (Suppl. Fig. S4). As the N-S
and W-E asymmetry increase systematically with slope (Fig. 4), part of the altitude-induced
variation in vegetation asymmetry could also be due to the effect of slope. In the higher altitude
regions (>~1500 m), especially in the CWG, the vegetation is characterized by a mosaic of
grassland and shola (i.e. stunted evergreen forest) (Meher-Homji, 1967). The highest N-S and
W-E asymmetry in TC and CH in such areas (Suppl. Fig. S14) is due to this "bi-modal"
vegetation pattern.

4.4. Combined effect of N-S and W-E asymmetries in the TC and CH

Our work indicated two modes control the landscape-scale TC and CH in the WG: the
solar declination induced and monsoonal climate facilitated asymmetry in vegetation on N-
and S-aspects; and orographic rainfall, and plausibly solar heating, induced W-E asymmetry.
These asymmetries can also be observed within the small watersheds. For example, hill ranges
within the watershed at Vishalgad in NWG (area ~41 km^2) (Fig. 6). The combined effects of
the N-S and W-E decreasing trends in vegetation resulted generally in N- and NW-aspects
having the highest TC and CH and S- and SE-aspects, the least (Figure 2, 7). Further, as the N-
S and W-E asymmetries increase with slope angle, the vegetation on the steeper slopes on S-
and SE-aspects exhibit the lowest TC and CH.

Figure 6. Elevation (a), Google Earth Image (b), Tree cover (c) and Canopy Height (d) at Vishalgad watershed. The dotted lines with numbers indicate ridges.

Figure 7. Number of PAs showing the lowest and highest tree cover (a) and canopy height (b) in different aspect categories.

5. Conclusion

The WG is an ecologically sensitive zone with high biodiversity and species endemicity

(Myers et al., 2000). Understanding the factors controlling vegetation distribution is important

not only for academic pursuits but also for the effective implementation of afforestation

programs and the conservation of biodiversity. This work demonstrates the slope-aspect

influence TC and CH at the landscape-scale in the WG. There is a need to acknowledge the

tendency of the southern aspects to have a lower TC and CH as a natural phenomenon; lower

TC and CH on the southern aspects should not be considered as "degraded" and a higher TC

and CH on northern aspects as "pristine". The relatively less favourable conditions for tree

growth on the S- and SE-aspects should be considered while designing afforestation programs

in the WG. Biodiversity assessment studies in the WG should consider the slope-aspect effect

as an additional framework while assessing species distribution at the landscape scale. This

work shows that slope-aspect can influence the vegetation structure even in the low-latitude

regions. This fundamental property of vegetation control needs to be considered as the driver

of vegetation structure in the high-relief areas in the tropics, such as in the WG.

References

1. Adrah, E., Wan Mohd Jaafar, W. S., Omar, H., Bajaj, S., Leite, R. V., Mazlan, S. M.,

... & Mohan, M. (2022). Analyzing canopy height patterns and environmental

landscape drivers in tropical forests using NASA's GEDI spaceborne LiDAR. *Remote*

Sensing, *14*(13), 3172. https://doi.org/10.3390/rs14133172

2. Badano, E. I., Cavieres, L. A., Molina-Montenegro, M. A., & Quiroz, C. L. (2005).

Slope aspect influences plant association patterns in the Mediterranean matorral of

central Chile. *Journal of Arid Environments*, *62*(1), 93-108.

https://doi.org/10.1016/j.jaridenv.2004.10.012

3. Champion, H. G., & Seth, S. K. (1968). *A revised survey of the forest types of India*.

Manager of publications.

4. Das, A., Krishnaswamy, J., Bawa, K. S., Kiran, M. C., Srinivas, V., Kumar, N. S., &

Karanth, K. U. (2006). Prioritisation of conservation areas in the Western Ghats,

India. *Biological conservation*, *133*(1), 16-31.

https://doi.org/10.1016/j.biocon.2006.05.023

5. Dubey, N., & Ghosh, S. (2023). The relative role of soil moisture and vapor pressure deficit in affecting the Indian vegetation productivity. *Environmental Research Letters*, *18*(6), 064012. https://doi.org/10.1016/j.ecolind.2022.108698

6. Fricker, G. A., Synes, N. W., Serra-Diaz, J. M., North, M. P., Davis, F. W., & Franklin, J. (2019). More than climate? Predictors of tree canopy height vary with scale in complex terrain, Sierra Nevada, CA (USA). *Forest Ecology and Management*, *434*, 142-153. https://doi.org/10.1016/j.foreco.2018.12.006

7. Gadgil, M. (1996). Western Ghats: a lifescape. *Journal of the Indian Institute of Science*, *76*(3), 495-504.

8. Geroy, I. J., Gribb, M. M., Marshall, H. P., Chandler, D. G., Benner, S. G., & McNamara, J. P. (2011). Aspect influences on soil water retention and storage. *Hydrological Processes*, *25*(25), 3836-3842. https://doi.org/10.1002/hyp.8281

9. Ghate U. , Joshi, N. V., & Gadgil, M. (1998). On the patterns of tree diversity in the Western Ghats of India. *Current Science*, 594-603. https://www.jstor.org/stable/24100563

10. Gimaret-Carpentier, C., Dray, S., & Pascal, J. P. (2003). Broad-scale biodiversity pattern of the endemic tree flora of the Western Ghats (India) using canonical correlation analysis of herbarium records. *Ecography*, *26*(4), 429-444. https://doi.org/10.1034/j.1600-0587.2003.03356.x

11. Guan, K., Pan, M., Li, H., Wolf, A., Wu, J., Medvigy, D., ... & Lyapustin, A. I. (2015). Photosynthetic seasonality of global tropical forests constrained by hydroclimate. *Nature Geoscience*, *8*(4), 284-289. https://doi.org/10.1038/ngeo2382

12. Gunnell, Y. (1997). Relief and climate in South Asia: the influence of the Western Ghats on the current climate pattern of peninsular India. *International journal of climatology: a journal of the Royal Meteorological Society*, *17*(11), 1169-1182. https://doi.org/10.1002/(SICI)1097-0088(199709)17:11%3C1169::AID-JOC189%3E3.0.CO;2-W

13. Gutiérrez-Jurado, H. A., Vivoni, E. R., Cikoski, C., Harrison, J. B. J., Bras, R. L., & Istanbulluoglu, E. (2013). On the observed ecohydrologic dynamics of a semiarid basin

with aspect-delimited ecosystems. *Water Resources Research*, *49*(12), 8263-8284. https://doi.org/10.1002/2013WR014364

14. Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A., Tyukavina, A., ... & Townshend, J. R. (2013). High-resolution global maps of 21st-century forest cover change. *science*, *342*(6160), 850-853. https://doi.org/10.1126/science.1244693

15. Holland, P. G., & Steyn, D. G. (1975). Vegetational responses to latitudinal variations in slope angle and aspect. *Journal of Biogeography*, 179-183. https://doi.org/10.2307/3037989

16. India Meteorological Department. (2022). Climatological Tables of Observatories in India 1991-2020 = Bhāratīya Vedhaśālāom Kī Jalavāu Vijñāna Sambandhī Sāraṇiyām. *New Delhi: Director General of Meteorology* **9th ed.**

17. Koch, G. W., Sillett, S. C., Jennings, G. M., & Davis, S. D. (2004). The limits to tree height. *Nature*, *428*(6985), 851-854. https://doi.org/10.1038/nature02417

18. Lewis, S. L., Brando, P. M., Phillips, O. L., Van Der Heijden, G. M., & Nepstad, D. (2011). The 2010 amazon drought. *Science*, *331*(6017), 554-554. https://doi.org/10.1126/science.1200807

19. Meher-Homji, V. M. (1967). Phytogeography of the South Indian hill stations. *Bulletin of the Torrey Botanical Club*, 230-242.

20. Moles, A. T., Warton, D. I., Warman, L., Swenson, N. G., Laffan, S. W., Zanne, A. E., ... & Leishman, M. R. (2009). Global patterns in plant height. *Journal of ecology*, *97*(5), 923-932. https://doi.org/10.1111/j.1365-2745.2009.01526.x

21. Myers, N., Mittermeier, R. A., Mittermeier, C. G., Da Fonseca, G. A., & Kent, J. (2000). Biodiversity hotspots for conservation priorities. *Nature*, *403*(6772), 853-858. https://doi.org/10.1038/35002501

22. Nagendra, H., & Utkarsh, G. (2003). Landscape ecological planning through a multi-scale characterization of pattern: studies in the Western Ghats, South India. *Environmental Monitoring and Assessment*, *87*, 215-233. https://doi.org/10.1023/A:1024878925038

23. NASA JPL. NASA Shuttle Radar Topography Mission Global 1 arc second [Data set]. *NASA EOSDIS Land Processes Distributed Active Archive Center,* (2013). Accessed 2023-12-01 from https://doi.org/10.5067/MEaSUREs/SRTM/SRTMGL1.003

24. NASA/METI/AIST/Japan Spacesystems and U.S./Japan ASTER Science Team. ASTER DEM Product [Data set]. *NASA EOSDIS Land Processes Distributed Active Archive Center* (2001). Accessed 2023-11-05 from https://doi.org/10.5067/ASTER/AST14DEM.003

25. Nikumbh, A. C., Chakraborty, A., & Bhat, G. S. (2019). Recent spatial aggregation tendency of rainfall extremes over India. *Scientific reports*, *9*(1), 10321. https://doi.org/10.1038/s41598-019-46719-2

26. Potapov, P., Li, X., Hernandez-Serna, A., Tyukavina, A., Hansen, M. C., Kommareddy, A., ... & Hofton, M. (2021). Mapping global forest canopy height through integration of GEDI and Landsat data. *Remote Sensing of Environment*, *253*, 112165. https://doi.org/10.1016/j.rse.2020.112165

27. Pai, D. S., Rajeevan, M., Sreejith, O. P., Mukhopadhyay, B., & Satbha, N. S. (2014). Development of a new high spatial resolution (0.25× 0.25) long period (1901-2010) daily gridded rainfall data set over India and its comparison with existing data sets over the region. *Mausam*, *65*(1), 1-18. https://doi.org/10.54302/mausam.v65i1.851

28. Pascal, J.P., 1988. Wet evergreen forests of the Western Ghats of India: ecology, structure, floristic composition and succession. *Institut Français de Pondichéry, Pondicherry.*

29. Pascal, J. P., Ramesh, B. R., & Franceschi, D. D. (2004). Wet evergreen forest types of the southern Western Ghats, India. *Tropical Ecology*, *45*(2), 281-292.

30. Prasad, V. K., Badarinath, K. V. S., & Eaturu, A. (2008). Effects of precipitation, temperature and topographic parameters on evergreen vegetation greenery in the Western Ghats, India. *International Journal of Climatology: A Journal of the Royal Meteorological Society*, *28*(13), 1807-1819. https://doi.org/10.1002/joc.1662

31. QGIS.org, %Y. QGIS Geographic Information System. *QGIS Association*. http://www.qgis.org

32. Ramachandra, T. V., Kumar, U., & Dasgupta, A., (2016). Time-series MODIS NDVI based Vegetation Change Analysis with Land Surface Temperature and Rainfall in the Western Ghats, India. *ENVIS Technical Report 100*, Sahyadri Conservation Series 53, ETR 100.

33. Shige, S., Nakano, Y., & Yamamoto, M. K. (2017). Role of orography, diurnal cycle, and intraseasonal oscillation in summer monsoon rainfall over the Western Ghats and

Myanmar Coast. *Journal of Climate*, *30*(23), 9365-9381. https://doi.org/10.1175/JCLI-
D-16-0858.1

34. Singh, N. (1986). On the duration of the rainy season over different parts of
India. *Theoretical and Applied Climatology*, *37*, 51-62.
https://doi.org/10.1007/BF00866104

35. Singh, S. H. I. P. R. A. (2018). Understanding the role of slope aspect in shaping the
vegetation attributes and soil properties in Montane ecosystems. *Tropical
Ecology*, *59*(3), 417-430.

36. Smith, B. J. (1978). Aspect-related variations in slope angle near Béni Abbès, western
Algeria. *Geografiska Annaler: Series A, Physical Geography*, *60*(3-4), 175-180.
https://doi.org/10.1080/04353676.1978.11879972

37. Smith, T., & Bookhagen, B. (2021). Climatic and biotic controls on topographic
asymmetry at the global scale. *Journal of Geophysical Research: Earth
Surface*, *126*(1), e2020JF005692. https://doi.org/10.1029/2020JF005692

38. Subrahmanyam, K. V., & Kumar, K. K. (2022). Diurnal evolution of orographic
precipitating clouds over the southernmost part of the Western Ghats of India during
summer and winter monsoons. *International Journal of Climatology*, *42*(14), 7146-
7161. https://doi.org/10.1002/joc.7635

39. Tao, S., Guo, Q., Li, C., Wang, Z., & Fang, J. (2016). Global patterns and determinants
of forest canopy height. *Ecology*, *97*(12), 3265-3270. https://doi.org/10.1002/ecy.1580

40. Tawde, S. A., & Singh, C. (2015). Investigation of orographic features influencing
spatial distribution of rainfall over the Western Ghats of India using satellite
data. *International Journal of Climatology*, *35*(9). https://doi.org/10.1002/joc.4146

41. The Sun angle data. (2009). Calculation of sun's position in the sky for each location
on the earth at any time of
day. https://www.sunearthtools.com/dp/tools/pos_sun.php?lang=en#annual

42. Venkatesh, B., Lakshman, N., Purandara, B. K., & Reddy, V. B. (2011). Analysis of
observed soil moisture patterns under different land covers in Western Ghats,
India. *Journal of hydrology*, *397*(3-4), 281-294.
https://doi.org/10.1016/j.jhydrol.2010.12.006

43. Viswanadhapalli, Y., Dasari, H. P., Dwivedi, S., Madineni, V. R., Langodan, S., &
Hoteit, I. (2020). Variability of monsoon low-level jet and associated rainfall over

India. *International Journal of Climatology*, *40*(2), 1067-1089.
https://doi.org/10.1002/joc.6256

44. Von Lengerke HJ. 1977. The nilgiris: weather and climate of a mountain area in south
India. *Wiesbaden, Germany: Franz Steiner Verlag.*

45. Zhou, X., Istanbulluoglu, E., & Vivoni, E. R. (2013). Modeling the ecohydrological
role of aspect-controlled radiation on tree-grass-shrub coexistence in a semiarid
climate. *Water Resources Research*, *49*(5), 2872-2895.
https://doi.org/10.1002/wrcr.20259

Acknowledgement:

We thank Vishwas Kale and Deepak Barua for going through the manuscript and offering
suggestion. Help extended by Rahul Tak with programming is appreciated.

Author contributions

Conceptualization: Shreyas Managave

Methodology: Devi Maheshwori, Shreyas Managave, Girish Jathar, Sham Davande

Formal Analysis: Devi Maheshwori, Shreyas Managave, Girish Jathar, Sham Davande

Investigation: Devi Maheshwori, Shreyas Managave, Girish Jathar, Sham Davande

Resources: Shreyas Managave

Writing—Original Draft: Devi Maheshwori, Shreyas Managave

Writing—Review and Editing: Devi Maheshwori, Shreyas Managave, Girish Jathar, Sham Davande

Supervision: Shreyas Managave

Funding Acquisition: Shreyas Managave

Competing interests

The authors declare no competing interest

Materials and correspondence

Supplementary Information

Supplementary Figure S1. Climatology of monthly precipitation, temperature and relative

humidity at selected PAs (a) Tamhini, (b) Kudremukh, (c) Parambikulam, (d) Neyyar.

Tamhni is in the NWG, Kudremukh is in the CWG, Parambikulam and Neyyar are in the

SWG. NWG, CWG and SWG indicate northern, central and southern Western Ghats,

respectively.

Supplementary Figure S2. The percent area covered by different aspect categories in the

selected PAs. The flat area covers less than 0.05 %.

Supplementary Figure S3. The area covered by different slope categories in the selected PAs.
Note, the area covered by slope 50-90° is typically less than 1%. The notations for the PAs:
a- Tamhini-Lavasa-Raigad; b- Koyna; c-Chandoli; d-Vishalgad; e-'West of the WG
Escarpment'; f-Radhangiri; g-Bhagwan Mahavir; h-Netravali; i-Anshi; j-Cotiago; k-
Kudremukh; l-Pushpagiri; m-Talakaveri; n-New Brahmagiri; o-Brahmagiri; p-Aralam; q-
New Amarambalam-Silent Valley; r-Peechi; s-Chimmony; t-Parambikulun; u-Indira Gandhi;
v-Idukki; w-Shendurney; x-Kallakad Mundanthurai; y-Neyyar.

Supplementary Figure S4. Whisker plot showing variation in the slope in different altitude
bands in the NWG (a), CWG (b) and SWG (c).

Supplementary Figure S5. Median tree cover (TC) (a), canopy height (CH) (b) and variation
between them (c) in the PAs in the WG.

Supplementary Figure S6. The correlation between the median TC and CH on 8 aspects
within PAs. The abbreviations for the PAs are as in Supplementary Figure S3. The horizontal
line with 'P=0.05' indicates the correlation coefficient at p-value of 0.05.

Supplementary Figure S7. The median TC (left panel) and CH (right panel) on the northeast

(NE), east (E) and southeast (SE) aspects in the studied PAs. The abbreviations for the PAs are

as in Supplementary Figure S3.

Supplementary Figure S8. The median TC (left panel) and CH (right panel) on the northwest

(NW), west (W) and southwest (SW) aspects in the studied PAs. The abbreviations for the PAs

are as in Supplementary Figure S3.

Supplementary Figure S9. The latitude-dependence of the N-S asymmetry in the TC (a) and CH (b) and of W-E asymmetry in the TC (c) and CH (d).

Supplementary Figure S10. The median TC (left panel) and CH (right panel) on the northwest (NW), north (N) and northeast (NE) aspects in the studied PAs. The abbreviations for the PAs are as in Supplementary Figure S3.

Supplementary Figure S11. The median TC (left panel) and CH (right panel) on the southwest (SW), south (S) and southeast (SE) aspects in the studied PAs. The abbreviations for the PAs are as in Supplementary Figure S3.

Supplementary Figure S12. Variation in the TC on N- and S-aspects (a, b, c) and W- and E-aspects (d, e, f) with the slope in NWG (a, d), CWG (b, e) and SWG (c, f). The TC values of S- and E-aspects for the PA Kudremukh (indicated by letter 'k' in Suppl. Fig. S3) from CWG and Idukki (indicated by letter 'v' in Suppl. Fig. S3) from SWG are removed to reveal the variability in TC in other PAs.

566   Their values are as below

PAs		Aspect	0°-10°	10°-30°	30°-50°,
k	CWG	S	72	71	0
k	CWG	E	70	62	0
v	SWG	S	67	63	0
v	SWG	E	72	64	3

Supplementary Figure S13. Variation in the CH on N- and S-aspects (a, b, c) and W- and E-
aspects (d, e, f) with slope in NWG (a, d), CWG (b, e) and SWG (c, f). The CH values of S-
and E-aspects for the PA Kudremukh (indicated by letter 'k' in Suppl. Fig. S3) from CWG and
Idukki (indicated by letter 'v' in Suppl. Fig. S3) from SWG are removed to reveal the variability
in TC in other PAs.

Their values are as below

PAs		Aspect	0°-10°	10°-30°	30°-50°,
k	CWG	S	17	17	5
k	CWG	E	16	14	5
v	SWG	S	16	16	9
v	SWG	E	17	16	7

Supplementary Figure S14. The altitude-dependant variation in the difference between the

median tree cover on (a) the N-aspect and S-aspect and (b) the W-aspect and E-aspect. The

same for canopy height is shown in (c) and (d). The abbreviations for the PAs are as in

Supplementary Figure S3.

Supplementary Figure S15. Variation in the TC on N- and S-aspects (a, b, c) and W- and E-aspects (d, e, f) with altitude in NWG (a, d), CWG (b, e) and SWG (c, f).

Supplementary Figure S16. Variation in the CH on N- and S-aspects (a, b, c) and W- and E-aspects (d, e, f) with altitude in NWG (a, d), CWG (b, e) and SWG (c, f).