Growth rate-driven modelling reveals how phenotypic adaptation drives drug resistance in BRAFV600E-mutant melanoma

Sara Hamis[1⊛], Alexander P Browning[2], Adrianne L Jenner[3], Chiara Villa[4,5], Philip K Maini[2], Tyler Cassidy[6].

1 Department of Information Technology, Uppsala University, Uppsala, Sweden.
2 Mathematical Institute, University of Oxford, Oxford, United Kingdom.
3 School of Mathematical Sciences, Queensland University of Technology, Brisbane, Australia.
4 Sorbonne Université, CNRS, Université de Paris, Inria, Laboratoire Jacques-Louis Lions UMR, Paris, France.
5 Université Paris-Saclay, Inria, Centre Inria de Saclay, 91120, Palaiseau, France.
6 School of Mathematics, University of Leeds, Leeds, United Kingdom.

Author ORCiDs: **SH:** 0000-0002-1105-8078; **APB:** 0000-0002-8753-1538; **ALJ:** 0000-0001-9103-7092; **CV:** 0000-0003-3127-0532; **PKM:** 0000-0002-0146-9164; **TC:** 0000-0003-0757-0017.

⊛ **Corresponding author**: sara.hamis@it.uu.se.

Abstract

Phenotypic adaptation, the ability of cells to change phenotype in response to external pressures, has been identified as a driver of drug resistance in cancer. To quantify phenotypic adaptation in BRAFV600E-mutant melanoma, we develop a theoretical model that emerges from data analysis of WM239A-BRAFV600E cell growth rates in response to drug challenge with the BRAF-inhibitor encorafenib. Our model constitutes a cell population model in which each cell is individually described by one of multiple discrete and plastic phenotype states that are directly linked to drug-dependent net growth rates and, by extension, drug resistance. Data-matched simulations reveal that phenotypic adaptation in the cells is directed towards states of high net growth rates, which enables evasion of drug-effects. The model subsequently provides an explanation for when and why intermittent treatments outperform continuous treatments *in vitro*, and demonstrates the benefits of not only targeting, but also leveraging, phenotypic adaptation in treatment protocols.

Keywords: Cell plasticity, heterogeneity, treatment schedules, mathematical model, individual-based model.

Introduction

Over 50% of melanoma cases present with BRAFV600 mutations that result in constitutive activation of the Mitogen Activated Protein Kinase (MAPK).[1,2] Consequently, targeted small molecule therapies that selectively inhibit mutant BRAF signalling have been developed over the last decades and are now part of the standard of care for BRAF-mutant melanoma. While these targeted therapies have led to a remarkable increase in progression free survival, treatment resistance inevitably develops and limits long-term survival.[3] This drug resistance in BRAF-mutant melanoma, and other solid tumours, is increasingly understood as a result of intratumoural heterogeneity, which has long been viewed through the lens of inter-clonal differences.[4,5,6] However, a growing body of work implicates cell plasticity, i.e., phenotypic adaptation without associated genetic mutations, as a key component of therapy resistance in melanoma.[1,7,8]

Advances in transcriptomics and multi-omics have led to the identification of reversible and adaptive phenotype changes that confer treatment resistance and are mediated by a number of complex physiological factors on the single-cell level.[9,10,11] As systems-level experiments of the complex interactions driving these adaptations are currently intractable, computational models have become increasingly important to our understanding of how therapeutic selection pressure shapes the epigenetic evolution of malignant tumours.[12,13,14] Such evolution is driven by epigentic changes in cancer cells, i.e., changes in gene expression that do not involve changes in the cells' DNA sequence.[15]

Phosphorylated, i.e., activated, ERK is often considered to be the main output of interest in pathway-level studies of BRAF-mutated melanoma. This follows from the fact that activated ERK phosphorylates multiple proliferation-promoting substrates in the nucleus and cytoplasm, which can drive uncontrolled cell growth and division.[16] Consequently, many existing computational approaches informed by extensive perturbation experiments leverage precise mechanistic representations of the MAPK pathway to describe signalling dynamics culminating in phosphorylated ERK.[13,17] With this ERK-endpoint, such models are not explicitly linked with the cellular response. Further, due to the large number of chemical reactions being modelled, these models are computationally expensive to train and simulate. Here, we present a novel approach to understanding cell plasticity that is computationally inexpensive and directly bridges cellular phenotype with net growth rates. Specifically, we develop a computational model that is directly informed by growth rate analysis of BRAFV600E-WM239A melanoma cells exposed to the BRAF-inhibitor encorafenib (LGX818) in a recent study by Kavran et al.[9]

In our model, each individual cell is described by a plastic phenotype state that can take one of S discrete values, where S is an arbitrary integer larger than 0. Further, each phenotype state is directly related to drug dose-dependent growth rates. Modelled phenotype states are thus mathematical abstractions that link cells to drug resistance. To investigate phenotypic adaptation in response to drug exposure and removal, we propose four candidate cell-level phenotype adaptation strategies. Subsequent comparisons of model-generated cell populations and *in vitro* cell count data reveal that phenotypic adaptation is directed towards phenotype states with high net growth rates. As such, our individual-based model demonstrates the emergence of drug resistance at the population-level as a result of directed phenotypic adaptation on the individual cell-level. Further, our modelling captures the experimental observation that intermittent treatment strategies can outperform their continuous counterparts *in vitro*, despite halved cumulative drug doses.[9] Based on multiple-dose growth rate data from untreated and treated cells, our framework thus provides a mechanistic explanation for when and why intermittent treatments may be more effective than continuous treatments. Building on these results, we perform further simulation experiments to quantitatively assess the potential of not only targeting, but also leveraging, cell plasticity in melanoma towards improved clinical outcomes.

Results

Empirical growth rates are mapped to phenotype states and give rise to a theoretical model

Our theoretical model formulation emerges from data analysis of drug dose-dependent net growth rates of BRAFV600E-mutant melanoma cells exposed to the BRAF-inhibitor encorafenib. The growth rates are shown in Figure 1a and are extracted from a recent experimental study by Kavran et al.[9] in which cell count numbers of WM239A-BRAFV600E melanoma cells were measured at 0 and 72 hours in response to drug challenge following 0-4 weeks of 500 nM encorafenib incubation. Cell populations that have been continuously incubated with drug for 1-4 weeks present dose-dependent net growth rates that are not significantly different from each other. We therefore assume that cells reach a drug-adapted state within 1 week of encorafenib incubation and stratify the observed growth rates as belonging to one of two extreme phenotypes: drug-naive cells and drug-adapted cells, i.e, cells that have been incubated in 500 nM encorafenib for 0 and 1-4 weeks, respectively (Figure 1b). Further, upon drug removal, drug-adapted cells in the regarded *in vitro* experiment resumed growth rates similar to those of drug-naive cells[9] (results not shown) hence we allow for bi-directional phenotype updates in the model.

Building on our growth rate stratification, we formulate a discrete adaptive phenotype model in which each individual cell in a cell population is described by a discrete and plastic phenotype state that can be updated in response to phenotypic instability or external pressures, here encorafenib presence or absence. We use the variable x_i to describe the phenotype state of cell i, where a cell's phenotype state corresponds to its sensitivity to the BRAF-inhibitor such that, arbitrarily, cells in state $x = 0$ are the most drug-sensitive and cells in state $x = 1$ are the most drug-resistant. In our model, growth rates for these phenotypes correspond to those for drug-naive and drug-adapted cells in Figure 1b, respectively. We further assume that the cells can exist in n intermittent phenotype states between the extrema $x = 0$ and $x = 1$, where each state is associated with drug dose-dependent net growth rates that are estimated through linear interpolation of growth rate data and are mapped onto a fitness matrix (Figure 1c-d). The multi-step adaptation model is based on the observation that protein expression for L1CAM, a marker for drug resistance in melanoma, changes gradually in response to external pressures in the regarded experiments.[9] In the main body of this article, we use $n = 9$, but key results are shown in the Supplementary Material (SM1) for other choices of n, including $n = 0$ which results in a single-step adaptation model.

In our discrete adaptive phenotype model, positive values in the fitness matrix correspond to daily probabilities that cells divide, whereas negative values correspond to daily probabilities that cells die. As is discussed in the next sections, our presented growth rate-to-phenotype state mapping enables growth rate-driven phenotype characterisation of cells, which we use to elucidate the directionality of phenotypic adaptation, predict treatment responses, and investigate the possibility of both targeting and leveraging cell plasticity in treatments.

Phenotypic adaptation in BRAFV600E-mutant melanoma is directed towards high fitness

Curiously, it was observed that four weeks of intermittent (one week on/one week off) treatments with 500 nM encorafenib are more effective at suppressing WM239A-BRAFV600E cell counts than four weeks of continuous treatments with the same dose *in vitro*.[9] Here, we set out to elucidate the cell-level behaviour related to phenotypic adaptation that gives rise to this macroscopic, population-level result. To investigate the directionality of phenotypic adaptation in the regarded melanoma cells in response to encorafenib exposure and removal, we present four candidate phenotype update strategies for describing cell-level dynamics. These

are illustrated in Figure 2a and are mathematically defined in the Methods section.

With the *no update* strategy, cells never update their phenotype states. With the *unbiased* strategy, cells propose to update their phenotype states to adjacent states with equal probability, unless at a boundary, and subsequently move to the proposed state. With the *semi-biased* strategy, cells propose to update their phenotype states to adjacent states with equal probability, but only accept the move if it results in a higher fitness, i.e., net growth rate. This corresponds to the cells having a noisy measurement of the fitness gradient. Finally, with the *biased* strategy, cells update their phenotype states to states of higher fitness whenever possible. As such, phenotypic adaptation is non-directed in the first two (no update and unbiased) strategies, but directed in the last two (semi-biased and biased) strategies. Biologically, the unbiased PU strategy corresponds to phenotype instability, while the biased PU strategy corresponds to stress-induced phenotypic adaptation, in accordance with work by Chisholm et al.[18] Ultimately, the mechanisms responsible for the development of resistance are not yet fully known, and we therefore also explore the more general semi-biased PU strategy than encompasses both phenotypic instability and adaptation. This gamut of phenotype progression is highly studied in segments of the mathematical literature[19] but, to the best of our knowledge, has not been applied in the interpretation of experimental data to select between candidate phenotype update models. Note that, while the no update strategy suggests that the presence of drug-resistant cells that exist post treatment arise purely from pre-exisiting phenotypic heterogeneity, the other three strategies allow drug-resistant cells to arise through phenotypic plasticity during treatment.

The four update strategies are implemented through a simple computational algorithm (Figure 2b) and produce significantly different phenotype density distributions in response to 300 nM encorafenib exposure and removal following continuous and intermittent treatments (Figure 2c). In the implementation, the same update rule applies both on and off treatments, but the directionality of the updates in the semi-biased and biased phenotype update strategies is drug dose dependent. While the non-directed strategies yield phenotype density distributions that only moderately change over time in response to intermittent treatments, the directed strategies yield phenotype distributions that markedly move between the drug-sensitive ($x = 0$) and drug-resistant ($x = 1$) extrema. Simulation results pertaining to 0 and 300 nM treatment responses have been produced with dose-dependent net growth rates from the fitness matrix (Figure 1d) and ρ, which denotes the probability that a cell changes its phenotype state at a programmed update time, set to $\rho = 1$ (Figure 2a). Cells have a chance to update their phenotype state η_{off} and η_{on} times per day in drug absence and presence, respectively. In Figure 1, we set $\eta_{\text{on}} = \eta_{\text{off}} = 2$, ensuring that cells can traverse phenotype space within a week (following observations in Figure 1a). Comparing simulated cell counts in Figure 2d, our model predicts a distinct benefit in using 300 nM encorafenib intermittent over continuous treatments for the directed phenotype update strategies, but not for the non-directed strategies.

Further simulation results with 500 nM demonstrate that only the directed strategies are able to match *in vitro* data[9] and reproduce cell population dynamics such that 500 nM encorafenib intermittent treatments outperform continuous treatments, in terms of suppressing cell counts (Figure 2). Phenotype-dependent net growth rates in response to 500 nM encorafenib are fit to data through a least-squares parameter estimation for each update strategy, where net growth rates are bounded by those for the adjacent drug doses for which data is available i.e., 300 nM and 1000 nM. The number of cellular phenotype updates per day, η_{on} and η_{off} are also chosen to best match the data (details about the parameter fitting are available in the Methods section). For all update models, with the exception of the biased update model, cell counts in response to continuous and intermittent treatments are similar in response to 300 nM and 500 nM encorafenib. This result is further discussed and visualised in the Supplementary Material (Figure S1). The Supplementary Material (Figure S3) also includes simulated cell

counts in response treatments for different choices of phenotype granularity, i.e., n, showing that if we consider the cell population to be phenotypically non-binary, then the directed strategies better explain the observed data than do the non-directed strategies.

Taken together, our results suggest that phenotypic adaptation in the regarded WM239A-BRAFV600E cells is directed towards phenotype states of high fitness through semi-biased or biased update strategies (Figure 2e). With the used model and available data, we can not distinguish which of the two directed strategies best describe the model system. However, the conclusion that phenotypic adaptation is directed provides a growth rate data-driven explanation for why intermittent treatments outperform continuous treatments at 500 nM encorafenib over 4 weeks. Moreover, with directed phenotypic adaptation, our model predicts that death events spike once treatments resume after a drug holiday, when the cells have acquired drug-sensitive phenotype states, in the intermittent treatments (Figure 2f). This simulation result is in qualitative agreement with Kavran et al.'s[9] experimental observation that the percentage of propidium iodide (PI) positive cells peak during drug re-challenge, where PI is used as a cell death marker, giving further credence to our model formulation and our simplifying model assumption that positive and negative net growth rates, respectively, can be approximated to yield birth and death events on the cell-level.

Simulations demonstrate the benefits of targeting and leveraging cell plasticity in treatments

Our results raise the question: Given directed phenotypic adaptation, when do intermittent treatments outperform their continuous counterparts? To address this question, we perform a series of simulation experiments in which output dynamics are quantified in response to variations of two model-inputs: the number of times per day that a cell can update its phenotype state (η), and the duration of treatment on/off intervals (t_{on}/t_{off}). Next, to enable biologically interpretable and mathematically tractable comparisons of continuous and intermittent treatments, we introduce the effective net growth rate λ_{eff} as the average (per cell and day) net growth rate that cells experience over the course of a simulation. Using the biased update strategy, we derive analytical approximations of effective net growth rates λ_{eff} obtained in the limit of large total treatment times (see the Methods section for details). For 300 nM and 500 nM encorafenib and treatment on/off intervals such that $t_{on} = t_{off} = T$, both simulation results with instantaneous drug on/off switches and analytical expressions identify pairings of η and T values that outperform modelled continuous treatments, i.e., yield lower effective growth rates than those for fully drug-adapted cells in drug presence. These pairings are highlighted by yellow regions in Figure 3b,e and are separated by analytically derived curves from $\eta - T$ pairings that result in higher effective growth rates than continuous treatments. Note that in the 300 nM heatmaps, all visualised $\eta - T$ pairings outperform the continuous treatment strategy, hence no separating curve is shown. Throughout Figure 3, we have used phenotype-dependent net growth rates for 500 nM that are obtained through log-linear interpolation between net growth rates for 300 and 1000 nM in the fitness matrix (Figure 1a).

When all cells start in the most drug-sensitive phenotype state, the effective net growth rates increase with the number of cell updates per day for the regarded doses and simulation inputs (Figure 3a-c). In other words, intermittent treatment become more effective when η is decreased. As such, our simulations highlight the therapeutic benefit of targeting phenotype adaptation by slowing it down.

Next, motivated by clinical studies in which drug re-challenge shows anti-tumour activity in patients who have previously been treated with BRAF-inhibitors,[3,20,21] we investigate the impact of drug-holidays on treatment responses. We thus perform a simulation experiment in which all cells start in the most drug-resistant phenotype state (Figure 3d-f), as would follow from a period of continuous treatment. In such cases, our model produces non-monotonic

relationships between effective net growth rates and the number of daily cell updates. This follows from the fact that cells may not have time to traverse phenotype space into states that are susceptible to sudden changes in drug exposure or removal. In an effort to push drug-resistant cells to drug-sensitive states in such cases, we can schedule a drug holiday and delay the treatment restart time. The benefits of delaying intermittent treatments are shown in Figure 3e, where the most effective treatment schedules are those that allow time for cells to traverse phenotype space to reach drug-susceptible states. Our simulation results thereby highlight the benefit of leveraging phenotypic adaptation in treatments.

Figure 1: **Empirical growth rates are mapped to phenotype states and give rise to a theoretical model. (a)** Growth rates for BRAFV600E-mutant melanoma cells from the WM239A cell-line are shown in response to various doses of the BRAF-inhibitor encorafenib, as measured over 72 hours. The rates are extracted from quadruplicate cell count assays by Kavran et al.[9] Prior to drug challenge, the cells are incubated in 500 nM encorafenib for 0,1,2,3, or 4 weeks (wks), with four replicates per incubation time. **(b)** Mean growth rates for drug-naive phenotypes (0 wks in a), and drug-adapted phenotypes (1-4 wks in a) are plotted over drug doses. Between the two extreme phenotype states, n intermediate states are introduced (here, $n = 9$). Marker colours and sizes correspond to phenotype state values and drug doses, respectively. **(c)** Net growth rates in (b) are mapped onto a discrete phenotype-dose space, as is indicated by marker colours and sizes. **(d)** Points in phenotype-dose space in (c) are mapped onto a fitness matrix (FM), in which the values of the left-most and right-most columns correspond to the growth rates in for drug-naive and drug-adapted phenotype states, respectively. Other values in the fitness matrix are obtained via linear interpolation between $\mathrm{FM}(x = 0, d_j)$ and $\mathrm{FM}(x = 1, d_j)$ for each of the j doses. Negative net growth rates are underlined.

Figure 2: **Phenotypic adaptation in BRAFV600E-mutant melanoma cells is directed towards states of high fitness.** **(a)** We propose four phenotype update strategies. The figure shows probabilities that a cell in state x_0 at time t updates to states x_-, x_0 and x_+ at time $t + 1$ whenever possible, i.e., ensuring that the phenotype states lie between the extrema $x = 0$ and $x = 1$. Net growth rates increase with decreasing x-values in the top row, and with increasing x-values in the bottom row. The directionality of the net growth rates, i.e., whether the top or bottom row applies, depends on the applied drug dose and is inferred from growth rate data. **(b)** A simple algorithm is used to implement our discrete adaptive phenotype model with the update strategies in (a). Each simulated day, cells update their phenotype states η times either before (with probability 0.5) or after (with probability 0.5) the resolution of cell division and death. **(c)** The histograms show how cell population phenotype density distributions change over time in response to no, continuous and intermittent 300 nM BRAF-inhibitor treatments for the update strategies in (a). Densities are shown with 100 simulations layered over each other (shaded regions) and means (solid bars). 28 days before the experiments start, cells are seeded with uniformly distributed phenotype states. **(d)** The plots show normalised cell counts over time in response to 300 and 500 nM continuous and intermittent treatments. Simulations start with 100 cells at day 1, with initial phenotype distributions matching day 1 in panel (c). The yellow box highlights that only the directed strategies (semi-biased and biased updates) are able to capture *in vitro* cell count dynamics. Mean (triangles) and standard errors for 6 replicate experiments are shown. **(e)** The highlighted result in panel (d), together with panels (a,c), suggest that phenotypic adaptation is directed towards states of high fitness, where this direction depends on the applied drug dose. Drug doses included in the top and bottom arrows respectively induce phenotypic adaptation in the direction of increasing and decreasing x-values. **(e)** The plots show the number of death events per cell over time for the simulation experiments in (d) with directed update strategies in response to 500 nM encorafenib. In (d,f) mean counts (solid/dashed lines) and standard deviations (shaded bands) are shown for 100 simulation runs.

Figure 3: **Simulations demonstrate the benefits of targeting and leveraging cell plasticity in treatments.**
(a) We consider an initial condition in which all cells are in the most drug-sensitive phenotype state ($x = 0$) in
(a-c). (b) Effective growth rates in response to 8 week intermittent treatments with 300 nM (left) and 500 nM (right)
encorafenib are obtained trough simulations with the biased phenotype update strategy. Based on mean values
from 100 simulations, heatmap bins show effective growth rates in response to different phenotype updates per
day (η), and on/off treatment intervals (T). Effective growth rates for 300 nM (left) and 500 nM (right) continuous
treatments are highlighted in the colour legends. In the 300 nM heatmap, all shown $\eta - T$ pairings yield lower
effective growth rates than the continuous treatment. In the 500 nM heatmap, the green, dashed curve separates
the $\eta - T$ pairings into two groups: pairings that yield lower effective growth rates than continuous treatments,
and pairings that do not. The curve is obtained through mathematical analysis (see Methods). (c) The plots show
simulated cell count dynamics in response to 1 and 4 cell updates per day, and treatment intervals of 1 and 4
days. These results demonstrate the benefit of targeting phenotypic adaptation in treatments by decreasing the
number of cell updates per day. (d) We consider an initial condition in which all cells are in the most drug-resistant
phenotype state ($x = 1$) in (d-e). (e) The experiments in (b) are repeated with the initial condition in (d). (f) The
plots show simulated cell count dynamics in response to 1 and 4 cell updates per day, and treatment intervals of 1
day. Treatments commence after a delay period of 1 (left), 4 (middle), or 10 (right) days. The results demonstrate
the benefit of leveraging phenotypic adaptation in treatments by delaying treatments and allowing drug-resistant
cells to desensitize by traversing phenotype space.

Discussion

Cell plasticity is not listed as one of the original *Hallmarks of Cancer* which were famously collated in 2000.[22] However, based on the last two decades' research advances, an updated version of the work identifies cancer cells' ability to unlock phenotypic adaptation as an emerging hallmark, intended to stimulate debate amongst researchers and inspire investigations that will improve our understanding of cancer.[8] The work presented in this article directly contributes to this understanding by evidencing that phenotypic adaptation in BRAFV600E-mutant melanoma is directed to states of high fitness, and demonstrating that this adaptation enables evasion of drug effects.

Our model emerges from growth rate data analysis of BRAFV600E-mutant melanoma cells exposed to the BRAF-inhibitor encorafenib. Our conclusion that phenotypic adaptation in the regarded cells is directed towards state of high fitness is obtained through a model-selection procedure, in which a gamut of phenotype update strategies on the cell-level are evaluated against dynamic cell count data in response to intermittent and continuous encorafenib treatments. One notable consequence of our results is that dynamic phenotypic adaptation alone suffices to explain why intermittent treatments can outperform continuous treatments. This explanation offers an alternative to variations of the more common inter-clonal competition-models in which two or more distinct subpopulations of cells, such as a drug-resistant and drug-sensitive clones, interact with each other and compete for resources.[23] This ecological perspective has served as the theoretical underpinning of *adaptive therapy*, where treatment is designed to maintain a drug-sensitive cancer cell population that competitively excludes drug-resistant clones.[24,25] This existing framework has historically relied on an assumed cost-of-resistance, where drug-resistant clones are less fit than drug-sensitive clones in the absence of treatment. However, recent theoretical work has demonstrated that this assumption is not necessary for adaptive therapies to outperform conventional maximum-tolerated dose strategies.[26,27] Here, in contrast with much of the adaptive therapy literature,[28,29] we use a simple model to capture improved treatment responses to intermittent therapy without explicitly modelling competition between clones.

The model developed in this paper is simple by design to enable direct mappings between phenotype states and growth rates. This simplicity, together with the informative results produced by the model, inspires and informs the design of further model-*in vitro* integrated research towards more detailed quantitative insight into phenotypic adaptation. For instance, data that clearly distinguish cell division and death events could be used to refine our model to include growth and death rates, as opposed to net growth rates only. Further, our methodology to linearly interpolate between drug-naive and drug-adapted growth rates to assign fitnesses to intermediate phenotype states could be refined by further drug challenge experiments. These would chart out growth rates in phenotype-dose parameter space through cell counts measured at multiple time points after the pre-assay drug incubation, and ideally include pre-assay drug incubation performed at multiple doses. It would also be informative to have data revealing if and when drug-adapted cells can fully recover the growth rates of drug-naive cells. This would allow us to assess dose-dependent velocities of phenotypic adaptation. Moreover, such experiments would contribute the open and contemporary discussion on whether cell-level drug resistance is, or is not, reversible in various cancer types.[30]

Whilst our results clearly indicate that phenotypic adaptation is directed towards state of high fitness, we can not determine if the semi-biased or biased update strategy best describes the regarded cells. This follows from the fact that we only consider mean growth rates in the model and do not have access to data that reveal population-level phenotype distributions. The observation that cell count data alone can not be used to distinguish between semi-biased and biased phenotype adaptation opens up for a broader question: what type of experimental data is needed to identify phenotypic heterogeneity in cell populations and quantify how bi-

ased phenotypic adaptation is? We address this question in a subsequent study,[31] where we have moved from a discrete to a continuous phenotype population model in order to allow for more rigorous mathematical analysis revealing that although cell count data alone is not sufficient to distinguish between the semi-biased and biased PU models, cell count data supplemented with information about birth and death events are. While we build our model on 3-day growth rates and (up to) four week treatment data, Russo et al.[32] combined long-term (over ten weeks) monitoring of experimental growth rates with an ordinary differential equation model to demonstrate that clinically approved targeted therapies induce drug-sensitive-to-tolerant adaptations in colorectal cancer cells. Further evidence of drug-induced cell plasticity was found by other recent studies combining mathematical modelling and *in vitro* experiments.[33,34] Beyond growth curve focussed studies, methods to quantify phenotype adaptation have been developed for data sets in which individual cells can be experimentally categorised into phenotype groups by, e.g., fluorescent markers.[35,36]

Mathematical models, analysis and simulations, especially those integrated with data, have made significant contributions to our understanding of cancer dynamics and treatment responses.[37,38] To highlight a few contributions pertaining to BRAFV600E-mutant melanoma, Gerosa et al.[17] integrated proteomics and modelling to show that exposure to drugs that target the BRAFV600E-MEK-ERK pathway can cause non-genetic drug resistance by signal rewiring in the targeted pathway. Such rewiring was also identified as a mechanism of drug resistance by Fröhlich et al.,[13] in an extension of the Gerosa et al. model, together with proteomic, transcriptomic and imaging data, to uncover that drug resistance can be mediated by the co-existence of two functionally distinct channels upstream of ERK, one initiated BRAFV600E monomers and the other by RAS. Whilst these models describe subcellular mechanisms that drive phenotypic adaptation, our model describes how cell-level phenotype traits, in the form of drug-dependent growth rates, change in response to BRAF-inhibitor exposure and removal. Comparing all three models, which complement each other, ours reveals less detailed subcellular information but is, on the other hand, considerably less data-intensive and therefore more accessible. Other data-integrated mathematical models of melanoma have been used to identify dose combinations of BRAFV600E, MEK, and ERK inhibitors that yield synergistic treatment responses,[39,40] predict initial treatment responses to BRAFV600E-inhibitors in xenografts,[41] and demonstrate phenotypic plasticity and multi-stability.[42] Beyond melanoma, cell plasticity, and its impact on treatment responses and the evolution of therapy resistance, has been studied mathematically through ordinary differential equation models[32] and partial integro-differential equation models[43,44,45,46,18] which, notably, can be used to suggest model-informed treatment strategies that impede the evolution of resistance.

Treatment scheduling and responses were also investigated in our current study. Our model-generated simulation results suggest that phenotypic adaptation can be both targeted and leveraged in an effort to suppress *in vitro* cell counts. These results support the development of molecular inhibitors that target mechanisms that drive cell plasticity.[47] Furthermore, strategies that leverage phenotypic adaptation, and, in other words, use cancers' ability to adapt against itself, can implicitly or explicitly be implemented through intermittent treatments and adaptive treatments.[48,28] In addition to these treatment strategies to combat adaptive resistance, there has been recent interest in epigenetic treatments that directly target phenotypic adaptation in the context of BRAF inhibitior resistant melanoma.[49,50] A recent proof-of-concept clinical trial of combination epigenetic and MAPK inhibitors demonstrated durable clinical responses in a minority of participants.[51] We emphasize that the treatment-related results presented in this study are based on data-driven models of *in vitro* systems, and that appropriate model extensions, as well as *in vivo* or clinical data, are needed to draw any conclusions about the role of phenotypic adaptation in xenografts or clinical tumours. However, one of the major benefits of modelling *in vitro* systems to study cell plasticity is the access to temporal and easy-to-interpret data that reveal how cancer systems change over time. Such

data are paramount to understanding dynamical aspects of cancer. Importantly, our limited understanding of cancer dynamics has been proposed as one of the main factors that hinder the development of efficient targeted therapy protocols[52,53] and whilst most cancer treatments target genomic cancer aberrations without consideration for evolutionary tumour aspects, clinical trials that are informed by mathematical models and account for tumour dynamics have started to emerge.[54]

The growth rate-to-phenotype modelling pipeline presented in this work can be used to quantify phenotypic adaptation velocities, i.e., directions and rates, in any cell lines for which cell counts can be measured. Our pipeline can thus be used to estimate such velocities in cancerous, non-cancerous and mutated cell lines, and thereby quantitatively assess to what extent phenotypic adaptation stratifies cancer cells, mutated or not, from other cells. This functionality of our pipeline is of significant value to contemporary cancer research, as an emerging body of work has identified non-genetic resistance as a reason for drug resistance in multiple cancer types, including melanoma,[7] neuroblastoma[55] and lung cancer.[56] One well-studied example mechanism of non-genetic resistance is the Epithelial-Mesenchymal Transition (EMT),[8,47,57] although a multitude of other mechanisms have been reported.[1,9,58,59,60,61] Our growth rate-driven methodology is agnostic to these subcellular mechanisms and can therefore be used to directly examine phenotypic adaptation in its role as an emerging hallmark of cancer in melanoma and beyond.

Methods

Extracting growth rates from cell count data

Our growth rate data are extrapolated from cell count assays performed by Kavran et al.[9] In the experiments, cells are incubated with 500 nM encorafenib prior to a drug challenge with an encorafenib dose of 0, 0.1, 0.3, 1, 3, 10, 30, 100, 300, 1000, 3000, or 10000 nM encorafenib. The fold changes in cell counts after 72 hours are reported for four replicates for each dose, relative to the fold changes in 500 nM encorafenib. To estimate effective growth rates from the relative fold change data, we assume exponential population growth over the 72 h period. Given a relative fold change measurement of FC_{72}^{rel}, the per-day growth rate λ is given by

$$\lambda = \frac{\log(FC_{72}^{rel})}{3} + \lambda^*. \tag{1}$$

As the fold change data are reported as relative to a standard treatment, for which $FC_{72}^{rel} = 1$, we normalise by λ^* i.e., the estimated growth rate for cells on a standard treatment. We estimate λ^* using reported cell count data following a 7 day period of 500 nM treatment, applying linear regression on the log-scale under the assumption that growth is approximately exponential. The growth rate data are obtained using the code file `ProcessData.m` in the project's GitHub repository.

Designing the fitness matrix

In this study, we model the net growth rate of a cell as a function of two variables: (1) the cell's phenotype state, and (2) the applied drug dose. The growth rates for cells in the melanoma system[9] that we study in this work are shown in the fitness matrix (Figure 1d), in which matrix element $FM(x, d)$ describes the net growth rate for cells in phenotype state x that are exposed to drug dose d. In this subsection, we describe how the fitness matrix is designed.

Based on the growth rate data, we say that drug-naive cells (0 wks in Figure 1a) are in a drug-sensitive phenotype state that we arbitrarily denote by $x = 0$. Similarly, we say that cells

that have been continuously exposed to 500 nM encorafenib for 1-4 weeks (1-4 wks in Figure 1a) are in a drug-adapted phenotype state $x = 1$. Based on the data, we assume that cells can move from state $x = 0$ to state $x = 1$ within one week when exposed to 500 nM of the BRAF-inhibitor. We further make the simplifying modelling assumption that cells can revert from state $x = 1$ to state $x = 0$ following a drug holiday. This is an approximation that is based on experimental results for the studied melanoma system.[9]

Assumption 1: Cells can revert from phenotype state $x = 1$ to phenotype state $x = 0$ and, by doing so, recover the growth rates of drug-naive cells.

Assumption 1 means that we can study phenotypic adaptation along a one-dimensional phenotype axis x when modelling how cells adapt to exposure and removal of drug at dose d. We note that *Assumption 1* may not be appropriate for all cell systems, as cell-level drug resistance in cancer ranges from reversible to irreversible.[30] Next, we make the simplifying modelling assumption that phenotype adaptation in directions of both increasing and decreasing x-values occurs in $n + 1$ discrete steps, where n is the number of intermediate phenotype states between the extrema $x = 0$ and $x = 1$.

Assumption 2: Cells move between phenotype states $x = 0$ to $x = 1$, and back, in $n + 1$ discrete steps.

In the main body of our article, we choose to work with $n = 9$, in order to allow us to study density distributions of phenotype states, such as the ones depicted in Figure 2c. However, our model could, for example, be reduced to study two-phenotype systems (with $n = 0$), or be extended to model continuous x-values, so that the phenotype density distributions would be described by partial differential equations. In the Supplementary Material (Figure S3), we include simulation results for a range of different n.

The growth rates FM(x,d) for cells in states $x = 0$ and $x = 1$ are directly assigned from the growth rate data extrapolation described in the previous subsection. The process of assigning growth rates to intermediate states $x \in (0, 1)$ is described in the next subsection. Note that the net growth rate values FM(x, d) in the right-most column of the fitness matrix ($x = 1$) are based on pre-incubation with 500 nM encorafenib, followed by 3 days of incubation in dose d. As such, the net growth rates of cells that have adapted to concentrations higher than 500 nM may have higher values than what our fitness matrix suggests. Experimental designs in which growth rates are measured following incubation with more than one dose (here 500 nM) would allow for a more detailed mapping of the fitness matrix.

Using growth rates to parameterise phenotype and drug dose-dependent fitnesses

We linearly interpolate between the extrema $x = 0$ and $x = 1$ to assign dose-dependent growth rate values to intermediate phenotypes $x \in (0, 1)$ for all doses included in the fitness matrix (Figure 1d). Since we do not have access to three-day cell count fold changes for 500 nM encorafenib, growth rates at this dose are instead estimated through a least-squares parameter fit evaluated against fold change data for 500 nM continuous and intermittent treatments (Figure 2d). In the fit, four free model parameters are estimated:

- FM($0, 500$ nM): the growth rate for drug-naive cells in 500 nM encorafenib.

- FM($1, 500$ nM): the growth rate for drug-adapted cells in 500 nM encorafenib.

- η_{on}: the number of times per day that cells have a chance to phenotype updates their phenotype when the drug is on.

- η_{off}: the number of times per day that cells have a chance to phenotype updates their phenotype when the drug is off.

In biased update models, the directionality of the phenotype adaptation is dependent on the applied drug dose, as determined by the data. As such, cells move towards states of higher proliferative net fitness (Figure 2a,c). To fit the free parameters, we further assume that the same phenotype update strategy applies both on and off drugs, although the phenotype update frequency may vary between drug-on and drug-off days.

Assumption 3: The same phenotype update strategy applies both on and off drugs.

The parameter combination C_ℓ^* that minimises the sum of squared distances between mean data values and model simulations (for 100 simulation runs) for continuous and intermittent treatments, simultaneously, are obtained for all phenotype update strategies ℓ. The fit is bounded by growth rates for the fitness matrix-adjacent drug doses 300 and 1000 nM between which 10 linearly interpolated values are tested. Further, η_{on} and η_{off} can take values $1/h$ and k, for integers h between 2 and 10, and k between 0 and 10. An update frequency of $1/h$ means that phenotype updates occur every hth day, and an update frequency of k means that phenotype updates occur k times a day. The parameter combinations that best match the *in vitro* data

$$C_{\ell*} = \Big(\text{FM}(0, 500 \text{ nM}), \text{FM}(1, 500 \text{ nM}), \eta_{\text{on}}, \eta_{\text{off}} \Big)$$

are used to produce the simulation results in Figures 2d,f for the 500 nM sub-panels, where

$$C_1^* = \Big(-0.202 \text{ day}^{-1}, 0.171 \text{ day}^{-1}, 0 \text{ updates day}^{-1}, 0 \text{ updates day}^{-1} \Big),$$

$$C_2^* = \Big(-0.202 \text{ day}^{-1}, 0.171 \text{ day}^{-1}, 1 \text{ updates day}^{-1}, 3 \text{ updates day}^{-1} \Big),$$

$$C_3^* = \Big(-0.335 \text{ day}^{-1}, 0.146 \text{ day}^{-1}, 3 \text{ updates day}^{-1}, 4 \text{ updates day}^{-1} \Big),$$

$$C_4^* = \Big(-0.268 \text{ day}^{-1}, 0.171 \text{ day}^{-1}, 1 \text{ updates day}^{-1}, 3 \text{ updates day}^{-1} \Big).$$

As is shown in the Supplementary Material (Figures S1-S2), fitting η_{on} and η_{off} only, using M(0,500 nM) and FM(1, 500 nM) obtained from log-linear interpolation between 300 nM and 1000 nM growth rates, does not result in simultaneously data-matched simulations for continuous and intermittent treatments. For all simulations in this article, $\rho =$ is fixed at $\rho = 1$ and thus the values of η_{on} and η_{off} determine the speed of phenotype adaptation.

Implementing cell division, cell death and phenotype updates

The algorithm (Algorithm 1) used to implement phenotype updates, cell division and cell death is illustrated in Figure 2b and is listed below. After an initial cell seeding, we loop over all simulated time points $\hat{t}$ and cells j. At every simulated time point, the drug dose is set according to a treatment schedule. Next, it is decided whether cell division and death events should be resolved *before* (with probability 0.5) or *after* (with probability 0.5) the phenotype state update events. If a cell has a positive net growth rate, the rate corresponds to the probability per day that the cell divides and produces an identical daughter cell. If a cell has a negative net growth rate, the rate corresponds to the probability per day that the cell dies. Within a simulated time unit (in our case, day), all phenotype updates are performed consecutively.

Algorithm 1 Implementation of cell division, cell death and phenotype updates.

1: Seed cells
2: **for** $\hat{t}$ = 1 to T **do**
3:     set the drug dose $d(\hat{t})$
4:     randomise a value v between 0 and 1
5:     **if** v < 0.5 **then**
6:         call function A
7:         call function B
8:     **else**
9:         call function B
10:         call function A
11:     **end if**
12: **end for**

    function A: resolve cell division and cell death
13: **for** j = 1 to J **do**
14:     g($x(j), d(\hat{t})$) ← FM(x, d)
15:     **if** g($x(j), d(\hat{t})$) > 0 **then**
16:         produce a daughter cell with a probability corresponding to g($x(j), d(\hat{t})$)
17:     **else**
18:         kill the cell with a probability corresponding to abs(g($x(j), d(\hat{t})$))
19:     **end if**
20: **end for**

    function B: resolve phenotype updates
21: **for** j = 1 to J **do**
22:     **for** update = 1 to η **do**
23:         update $x(j)$ according to the applied phenotype update rule
24:     **end for**
25: **end for**

Modelling cell-level phenotypic adaptation with different update strategies

The four phenotype update (PU) strategies in Figure 2a are formalised in this subsection. In order, these are the no, unbiased, semi-biased and biased PU strategies. We let x_0 denote the current phenotype state of an arbitrary cell. The probabilities of updating to higher (x_+) and lower (x_-) states, if possible with the restriction $x \in [0, 1]$, are given below.

PU1: $p(x_+) := 0,$ $p(x_-) := 0.$ (2a)

PU2: $p(x_+) := \dfrac{\rho}{2},$ $p(x_-) := \dfrac{\rho}{2}.$ (2b)

PU3: $p(x_+) := \dfrac{\rho}{2}H(\text{FM}(x_+, d) - \text{FM}(x_0, d)),$ $p(x_-) := \dfrac{\rho}{2}H(\text{FM}(x_-, d) - \text{FM}(x_0, d)).$ (2c)

PU4: $p(x_+) := H(\text{FM}(x_+, d) - \text{FM}(x_0, d)),$ $p(x_-) := H(\text{FM}(x_-, d) - \text{FM}(x_0, d)).$ (2d)

Probabilities to remain at the current state x_0 are implied from the above equations through the relationship $p(x_+) + p(x_0) + p(x_-) = 1$. The value FM($x, d$) can be read from the fitness matrix in Figure 1d, and denotes the net growth rate of a cell in phenotype state x that is exposed to drug dose d. In Eqs. 2a-d, $H(\cdot)$ is the Heaviside function, and $\rho/2$ denotes the probability that a cell updates its phenotype state in a given direction at an update time (Figure 2a). Note that

the no update rule can be considered a special case of both the unbiased and semi-biased update rule with $\rho = 0$. Further, the biased update rule can be considered a special case of the semi-biased update rule with $\rho/2 = 1$.

Deriving analytical expressions to identify simulated intermittent treatment regimes that outperform their continuous counterparts

To predict when intermittent treatment schedules outperform continuous schedules, we derive analytical expressions for long-term effective growth rates under the biased phenotype update strategy. We let t_{on} denote the duration (per cycle) that the drug is *on*, and t_{off} denote the duration (per cycle) that the drug is *off*. ω is the phenotype adaptation velocity (in both directions) such that $\eta = \omega/\Delta$ gives the number of phenotype updates per day. In the Figure 1d example, $\Delta = 0.1$ is the step size arising from the choice $n = 9$. Implicit to the biased update rule is that, at any time in a simulation, there is no phenotypic heterogeneity between cells (Figure 2c). As such, all cells have an identical phenotype, denoted by $x(t)$, the dynamics of which are governed by

$$\frac{dx}{dt} = \begin{cases} \omega, & x < 1 \text{ and the drug is on,} \\ -\omega, & x > 0 \text{ and the drug is off,} \\ 0 & \text{otherwise,} \end{cases} \tag{3}$$

for drug doses 300 and 500 nM, and any other doses listed inside the top arrow of Figure 2e. Now, let's assume that we have a cell population in equilibrium such that $x(s) = x(0)$. Then, over a time period $\tau = t_{\text{on}} + t_{\text{off}}$, the cell count $N(s + \tau)$ for cells on intermittent treatment is given by

$$N(s + \tau) = N(s)\text{e}^{\lambda_{\text{eff}} \cdot \tau}, \tag{4}$$

where the effective growth rate, λ_{eff}, is

$$\lambda_{\text{eff}} = \frac{1}{\tau} \left(\int_0^{t_{\text{on}}} \text{FM}(x(t), d)\, dt + \int_{t_{\text{on}}}^{\tau} \text{FM}(x(t), 0)\, dt \right). \tag{5}$$

Solving Eq. 5 yields the effective growth rate for cells on intermittent treatment to be

$$\lambda_{\text{eff}} = \begin{cases} \dfrac{\Delta\text{FM}(0) - \Delta\text{FM}(D)}{2\tau\omega} + \dfrac{t_{\text{on}}\text{FM}(1, D) + t_{\text{off}}\text{FM}(0, 0)}{\tau}, & \text{if } t_{\text{on}}, t_{\text{off}} > \dfrac{1}{\omega}, \\[2ex] \dfrac{\omega t_{\text{on}}^2(\Delta\text{FM}(0) + \Delta\text{FM}(D))}{2\tau} + \dfrac{t_{\text{on}}\text{FM}(0, D) + t_{\text{off}}\text{FM}(0, 0)}{\tau}, & \text{if } t_{\text{on}} < t_{\text{off}} \leq \dfrac{1}{\omega}, \\[2ex] \dfrac{-\omega t_{\text{off}}^2(\Delta\text{FM}(0) + \Delta\text{FM}(1))}{2\tau} + \dfrac{t_{\text{on}}\text{FM}(1, D) + t_{\text{off}}\text{FM}(1, 0)}{\tau}, & \text{if } t_{\text{off}} < t_{\text{on}} \leq \dfrac{1}{\omega}, \\[2ex] \dfrac{\text{FM}(0, 0) + \text{FM}(0, D) + \text{FM}(1, 0) + \text{FM}(1, D)}{4}, & \text{if } t_{\text{on}} = t_{\text{off}} = \dfrac{1}{\omega}, \\[2ex] \dfrac{\omega t_{on}(\Delta\text{FM}(0) + \Delta\text{FM}(D))}{4} + \dfrac{\text{FM}(0, 0) + \text{FM}(0, D)}{2}, & \text{if } t_{\text{on}} = t_{\text{off}} < \dfrac{1}{\omega}, \\[2ex] \dfrac{\Delta\text{FM}(0) - \Delta\text{FM}(D)}{4t_{on}\omega} + \dfrac{\text{FM}(0, 0) + \text{FM}(1, D)}{2}, & \text{if } t_{\text{on}} = t_{\text{off}} > \dfrac{1}{\omega}, \end{cases} \tag{6}$$

where

$$\Delta\text{FM}(d) = \text{FM}(1, d) - \text{FM}(0, d), \tag{7}$$

and D is a chosen drug dose such that Eq. 3 holds. The expressions in Eq. 6 are derived as shown in the Supplementary Material (SM2), and allow a user to find λ_{eff} for customised values of $D, t_{\text{on}}, t_{\text{off}}, \omega$. To, next, draw fair comparisons between cells on continuous and intermittent treatments, we can compare the long-term effective growth rate of cells under both regimes. To avoid effects of the initial drug desensitization for cells undergoing continuous treatments,

we consider the effective growth rates under continuous treatment to be given by $\mathrm{FM}(1, D)$. Thus, we expect intermittent treatment to outperform continuous treatment if $\lambda_{\mathrm{eff}}(D, t_{\mathrm{on}}, t_{\mathrm{off}}, \omega)$ is lower than $\mathrm{FM}(1, D)$. In Figure 3, we consider a treatment regime with equal on and off drug intervals, such that $t_{\mathrm{on}} = t_{\mathrm{off}} = \tau/2 = T$. Results are produced with net growth rates obtained through log-linear interpolation between $\mathrm{FM}(x, 300\ \mathrm{nM})$ and $\mathrm{FM}(x, 1000\ \mathrm{nM})$ so that $\mathrm{FM}(0, 500\ \mathrm{nM}) = -0.287\ \mathrm{day}^{-1}$ and $\mathrm{FM}(1, 500\ \mathrm{nM}) = 0.074\ \mathrm{day}^{-1}$ (see Figure S1, Supplementary Material). Inserting this choice into Eq. 6 for $T\omega \geq 1$, we find that intermittent treatment will outperform continuous treatment if $T\omega < 6.75245$ or, equivalently, $T\eta < 67.5245$.

Implementing the discrete adaptive phenotype model and performing simulation experiments

The model is implemented in MATLAB. Information on how to access, run and modify the code files is available on the public GitHub repository `https://github.com/SJHamis/phenotype_adaptation`.

Acknowledgments

SH was funded by the Swedish Research Council (project 2024-05621), Wenner-Gren Stiftelserna/the Wenner-Gren Foundations (WGF2022-0044), and the Kjell och Märta Beijer Foundation. APB thanks the Mathematical Institute, Oxford for a Hooke Research Fellowship. ALJ thanks the London Mathematical Society. This project was partially supported by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 945298-ParisRegionFP (CV). CV is a Fellow of the Paris Region Fellowship Programme, supported by the Paris Region. This work was partially supported by a Heilbronn Institute for Mathematical Research Small Maths Grant to TC.

Author contributions

Conceptualization: All authors. Formal analysis: SH, APB, ALJ. Investigation: SH, APB. Software: SH, APB. Methodology: All authors. Visualization: SH. Writing – original draft: SH, APB, TC. Writing – editing: All authors.

Declaration of interests

The authors declare no competing interests.

References

[1] Diazzi, S., S. Tartare-Deckert, and M. Deckert (2023). "The mechanical phenotypic plasticity of melanoma cell: an emerging driver of therapy cross-resistance", *Oncogenesis* 12.1, p. 7. DOI: `10.1038/s41389-023-00452-8`.

[2] Dummer, R. et al. (2022). "COLUMBUS 5-Year Update: A Randomized, Open-Label, Phase III Trial of Encorafenib Plus Binimetinib Versus Vemurafenib or Encorafenib in Patients With BRAFV600E-Mutant Melanoma", *J. Clin. Oncol.* 40.36, pp. 4178–4188. DOI: `10.1200/JCO.21.02659`.

[3] Fukushima, H., Y. Iwata, K. Saito, and K. Sugiura (2021). "Successful rechallenge therapy for BRAF/MEK inhibitor-resistant multiple brain metastases of melanoma", *J. Dermatol* 48.8, pp. 1291–1295. DOI: `10.1111/1346-8138.15969`.

[4] Dagogo-Jack, I. and A. T. Shaw (2018). "Tumour heterogeneity and resistance to cancer therapies", *Nat. Rev. Clin. Oncol.* 15.2, pp. 81–94. DOI: `10.1038/nrclinonc.2017.166`.

[5] McGranahan, N. and C. Swanton (2017). "Clonal Heterogeneity and Tumor Evolution: Past, Present, and the Future", *Cell* 168.4, pp. 613–628. DOI: `10.1016/j.cell.2017.01.018`.

[6] Burkhardt, D. B., B. P. San Juan, J. G. Lock, S. Krishnaswamy, and C. L. Chaffer (2022). "Mapping Phenotypic Plasticity upon the Cancer Cell State Landscape Using Manifold Learning", *Cancer Discov.* 12.8, pp. 1847–1859. DOI: 10.1158/2159-8290.CD-21-0282.

[7] Arozarena, I. and C. Wellbrock (2019). "Phenotype plasticity as enabler of melanoma progression and therapy resistance", *Nat. Rev. Cancer* 19.7, pp. 377–391. DOI: 10.1038/s41568-019-0154-4.

[8] Hanahan, D. (2022). "Hallmarks of Cancer: New Dimensions", *Cancer Discov.* 12.1, pp. 31–46. DOI: 10.1158/2159-8290.CD-21-1059.

[9] Kavran, A. J., S. A. Stuart, K. R. Hayashi, J. M. Basken, B. J. Brandhuber, and N. G. Ahn (2022). "Intermittent treatment of BRAFV600E melanoma cells delays resistance by adaptive resensitization to drug rechallenge", *Proc. Natl. Acad. Sci. U. S. A.* 119.12. DOI: 10.1073/pnas.2113535119.

[10] Fennell, K. A. et al. (2022). "Non-genetic determinants of malignant clonal fitness at single-cell resolution", *Nature* 601.7891, pp. 125–131. DOI: 10.1038/s41586-021-04206-7.

[11] Chen, J.-Y. et al. (2023). "Multi-range ERK responses shape the proliferative trajectory of single cells following oncogene induction", *Cell Rep.* 42.3, p. 112252. DOI: 10.1016/j.celrep.2023.112252.

[12] Foo, J. et al. (2022). "Roadmap on plasticity and epigenetics in cancer", *Phys. Biol.* 19.3, p. 031501. DOI: 10.1088/1478-3975/ac4ee2.

[13] Fröhlich, F., L. Gerosa, J. Muhlich, and P. K. Sorger (2023). "Mechanistic model of MAPK signaling reveals how allostery and rewiring contribute to drug resistance", *Mol. Syst. Biol.* 19.2, pp. 1–26. DOI: 10.15252/msb.202210988.

[14] Schubert, M., B. Klinger, M. Klünemann, A. Sieber, F. Uhlitz, S. Sauer, M. J. Garnett, N. Blüthgen, and J. Saez-Rodriguez (2018). "Perturbation-response genes reveal signaling footprints in cancer gene expression", *Nat. Commun.* 9.1, p. 20. DOI: 10.1038/s41467-017-02391-6.

[15] Easwaran, H., H. C. Tsai, and S. B. Baylin (2014). *Cancer Epigenetics: Tumor Heterogeneity, Plasticity of Stem-like States, and Drug Resistance*. DOI: 10.1016/j.molcel.2014.05.015.

[16] Wellbrock, C. and A. Hurlstone (2010). *BRAF as therapeutic target in melanoma*. DOI: 10.1016/j.bcp.2010.03.019.

[17] Gerosa, L. et al. (2020). "Receptor-Driven ERK Pulses Reconfigure MAPK Signaling and Enable Persistence of Drug-Adapted BRAF-Mutant Melanoma Cells", *Cell Syst.* 11.5, pp. 478–494. DOI: 10.1016/j.cels.2020.10.002.

[18] Chisholm, R. H., T. Lorenzi, A. Lorz, A. K. Larsen, L. N. D. Almeida, A. Escargueil, and J. Clairambault (2015). "Emergence of Drug Tolerance in Cancer Cell Populations: An Evolutionary Outcome of Selection, Nongenetic Instability, and Stress-Induced Adaptation", *Cancer Research* 75 (6). ISSN: 15387445. DOI: 10.1158/0008-5472.CAN-14-2103.

[19] Lorenzi, T., R. H. Chisholm, L. Desvillettes, and B. D. Hughes (2015). "Dissecting the dynamics of epigenetic changes in phenotype-structured populations exposed to fluctuating environments", *J. Theor. Biol.* 386, pp. 166–176. DOI: 10.1016/j.jtbi.2015.08.031.

[20] Schreuer, M., Y. J. L. Jansen, S. Planken, I. Chevolet, T. Seremet, V. Kruse, and B. Neyns (2017). "Combination of dabrafenib plus trametinib for BRAF and MEK inhibitor pretreated patients with advanced BRAFV600-mutant melanoma: an open-label, single arm, dual-centre, phase 2 clinical trial", *Lancet Oncol.* 18.4, pp. 464–472. DOI: 10.1016/S1470-2045(17)30171-7.

[21] Reschke, R., J. C. Simon, and M. Ziemer (2019). "Rechallenge of targeted therapy in metastatic melanoma", *J. Dtsch. Dermatol. Ges.* 17. DOI: 10.1111/ddg.13766.

[22] Hanahan, D and R. A. Weinberg (Jan. 2000). "The hallmarks of cancer", *Cell* 100, pp. 57–70. DOI: 10.1016/s0092-8674(00)81683-9.

[23] Madan, E. et al. (Dec. 2022). "Cell Competition in Carcinogenesis", *Cancer Res.* 82.24, pp. 4487–4496. DOI: 10.1158/0008-5472.CAN-22-2217.

[24] Zhang, J., J. Cunningham, J. Brown, and R. Gatenby (2022). "Evolution-based mathematical models significantly prolong response to abiraterone in metastatic castrate-resistant prostate cancer and identify strategies to further improve outcomes", *Elife* 11, pp. 1–105. ISSN: 2050-084X. DOI: 10.7554/eLife.76284.

[25] West, J., L. You, J. Zhang, R. A. Gatenby, J. S. Brown, P. K. Newton, and A. R. Anderson (2020). "Towards Multidrug Adaptive Therapy", *Cancer Res.* 80.7, pp. 1578–1589. ISSN: 0008-5472. DOI: 10.1158/0008-5472.CAN-19-2669.

[26] Strobl, M. A. R., J. Gallaher, J. West, M. Robertson-Tessi, P. K. Maini, and A. R. A. Anderson (2022). "Spatial structure impacts adaptive therapy by shaping intra-tumoral competition", *Commun. Med.* 2.1, p. 46. ISSN: 2730-664X. DOI: 10.1038/s43856-022-00110-x.

[27] Viossat, Y. and R. Noble (2021). "A theoretical analysis of tumour containment", *Nat. Ecol. Evol.* 5.6, pp. 826–835. ISSN: 2397-334X. DOI: 10.1038/s41559-021-01428-w.

[28] Gatenby, R. A., A. S. Silva, R. J. Gillies, and B. R. Frieden (2009). "Adaptive therapy", *Cancer Res.* 69.11, pp. 4894–4903. DOI: 10.1158/0008-5472.CAN-08-3658.

[29] Gatenby, R. A. and J. S. Brown (2020). "Integrating evolutionary dynamics into cancer therapy", *Nat. Rev. Clin. Oncol.* ISSN: 1759-4774. DOI: 10.1038/s41571-020-0411-1.

[30] Bhat, G. R. et al. (2024). *Cancer cell plasticity: from cellular, molecular, and genetic mechanisms to tumor heterogeneity and drug resistance*. DOI: 10.1007/s10555-024-10172-z.

[31] Browning, A. P., R. M. Crossley, C. Villa, P. K. Maini, A. L. Jenner, T. Cassidy, and S. Hamis (2024). "Identifiability of heterogeneous phenotype adaptation from low-cell-count experiments and a stochastic model", *bioRxiv*. DOI: 10.1101/2024.08.19.608540.

[32] Russo, M. et al. (2022). "A modified fluctuation-test framework characterizes the population dynamics and mutation rate of colorectal cancer persister cells", *Nature Genetics* 54 (7). ISSN: 15461718. DOI: 10.1038/s41588-022-01105-z.

[33] Gevertz, J. L., J. M. Greene, S. Prosperi, N. Comandante-Lou, and E. D. Sontag (2024). "Understanding therapeutic tolerance through a mathematical model of drug-induced resistance", *bioRxiv*. DOI: 10.1101/2024.09.04.611211.

[34] Wu, C., E. B. Gunnarsson, J. Foo, and K. Leder (2024). *A statistical framework for detecting therapy-induced resistance from drug screens*. arXiv: 2409.18879 [q-bio.PE]. URL: https://arxiv.org/abs/2409.18879.

[35] Gupta, P. B., C. M. Fillmore, G. Jiang, S. D. Shapira, K. Tao, C. Kuperwasser, and E. S. Lander (2011). "Stochastic state transitions give rise to phenotypic equilibrium in populations of cancer cells", *Cell* 146 (4). ISSN: 10974172. DOI: 10.1016/j.cell.2011.07.026.

[36] Buder, T., A. Deutsch, M. Seifert, and A. Voss-Böhme (2017). "CellTrans: An R Package to Quantify Stochastic Cell State Transitions", *Bioinformatics and Biology Insights* 11. ISSN: 11779322. DOI: 10.1177/1177932217712241.

[37] Altrock, P. M., L. L. Liu, and F. Michor (2015). "The mathematics of cancer: integrating quantitative models", *Nat. Rev. Cancer* 15.12, pp. 730–745. DOI: 10.1038/nrc4029.

[38] Rockne, R. C. et al. (2019). "The 2019 mathematical oncology roadmap", *Phys. Biol.* 16 (4). DOI: 10.1088/1478-3975/ab1a09.

[39] Hamis, S. J., Y. Kapelyukh, A. McLaren, C. J. Henderson, C. Roland Wolf, and M. A. J. Chaplain (Nov. 2021). "Quantifying ERK activity in response to inhibition of the BRAFV600E-MEK-ERK cascade using mathematical modelling", *Br. J. Cancer* 125.11, pp. 1552–1560. DOI: 10.1038/s41416-021-01565-w.

[40] De Carli, A., Y. Kapelyukh, J. Kursawe, M. A. J. Chaplain, C. R. Wolf, and S. Hamis (May 2024). "Simulating BRAFV600E-MEK-ERK signalling dynamics in response to vertical inhibition treatment strategies", *npj Syst. Biol. and Appl.* 10.1. DOI: 10.1038/s41540-024-00379-9.

[41] Smalley, I. et al. (2019). "Leveraging transcriptional dynamics to improve BRAF inhibitor responses in melanoma", *EBioMedicine* 48, pp. 178–190. DOI: 10.1016/j.ebiom.2019.09.023.

[42] Pillai, M., Z. Chen, M. K. Jolly, and C. Li (2022). "Quantitative landscapes reveal trajectories of cell-state transitions associated with drug resistance in melanoma", *iScience* 25.12, p. 105499. DOI: 10.1016/j.isci.2022.105499.

[43] Clairambault, J. and C. Pouchol (2019). "A survey of adaptive cell population dynamics models of emergence of drug resistance in cancer, and open questions about evolution and cancer", *BIOMATH*. hal-02126727v3.

[44] Stace, R. E., T. Stiehl, M. A. J. Chaplain, A. K. Marciniak-Czochra, and T. Lorenzi (2020). "Discrete and continuum phenotype-structured models for the evolution of cancer cell populations under chemotherapy", *Math. Model. Nat. Phenom.* 15.14. DOI: 10.1051/mmnp/2019027.

[45] Chisholm, R. H., T. Lorenzi, and J. Clairambault (2016). "Cell population heterogeneity and evolution towards drug resistance in cancer: Biological and mathematical assessment, theoretical treatment optimisation", *Biochim Biophys Acta* 1860.11, Part B. DOI: 10.1016/j.bbagen.2016.06.009.

[46] Cassidy, T., D. Nichol, M. Robertson-Tessi, M. Craig, and A. R. A. Anderson (Aug. 2021). "The role of memory in non-genetic inheritance and its impact on cancer treatment resistance", *PLoS Comput. Biol.* 17.8, e1009348. DOI: 10.1371/journal.pcbi.1009348.

[47] Boumahdi, S. and F. J. de Sauvage (2020). "The great escape: tumour cell plasticity in resistance to targeted therapy", *Nat. Rev. Drug Discov.* 19.1, pp. 39–56. DOI: 10.1038/s41573-019-0044-1.

[48] Labrie, M., J. S. Brugge, G. B. Mills, and I. K. Zervantonakis (2022). "Therapy resistance: opportunities created by adaptive responses to targeted therapies in cancer", *Nat. Rev. Cancer* 22.6, pp. 323–339. DOI: 10.1038/s41568-022-00454-5.

[49] Emmons, M. F. et al. (2019). "HDAC8 Regulates a Stress Response Pathway in Melanoma to Mediate Escape from BRAF Inhibitor Therapy", *Cancer Res.* 79.11, pp. 2947–2961. ISSN: 0008-5472. DOI: 10.1158/0008-5472.CAN-19-0040. URL: https://aacrjournals.org/cancerres/article/79/11/2947/633118/HDAC8-Regulates-a-Stress-Response-Pathway-in.

[50] Gallagher, S. J. et al. (2018). "HDAC inhibitors restore BRAF☐inhibitor sensitivity by altering PI3K and survival signalling in a subset of melanoma", *Int. J. Cancer* 142.9, pp. 1926–1937. ISSN: 0020-7136. DOI: 10.1002/ijc.31199. URL: https://onlinelibrary.wiley.com/doi/10.1002/ijc.31199.

[51] Embaby, A. et al. (2024). "A Proof-of-Concept Study of Sequential Treatment with the HDAC Inhibitor Vorinostat following BRAF and MEK Inhibitors in BRAF V600 -Mutated Melanoma", *Clin. Cancer Res.* 30.15, pp. 3157–3166. ISSN: 1078-0432. DOI: 10.1158/1078-0432.CCR-23-3171.

URL: `https://aacrjournals.org/clincancerres/article/30/15/3157/746581/A-Proof-of-Concept-Study-of-Sequential-Treatment`.

[52] Gatenby, R. A., Y. Artzy-Randrup, T. Epstein, D. R. Reed, and J. S. Brown (Feb. 2020). "Eradicating Metastatic Cancer and the Eco-Evolutionary Dynamics of Anthropocene Extinctions", *Cancer Res.* 80.3, pp. 613–623. DOI: `10.1158/0008-5472.CAN-19-1941`.

[53] Wang, Z. and T. S. Deisboeck (2019). "Dynamic Targeting in Cancer Treatment", *Front. Physiol.* 10, p. 96. DOI: `10.3389/fphys.2019.00096`.

[54] Poels, K. E. et al. (June 2021). "Identification of optimal dosing schedules of dacomitinib and os-imertinib for a phase I/II trial in advanced EFGR-mutant non-small cell lung cancer", *Nat. Commun.* 12.1, p. 3697. DOI: `10.1038/s41467-021-23912-4`.

[55] Thirant, C. et al. (2023). "Reversible transitions between noradrenergic and mesenchymal tumor identities define cell plasticity in neuroblastoma", *Nat. Commun.* 14.1, p. 2575. DOI: `10.1038/s41467-023-38239-5`.

[56] Wang, F. and L. Zhu (2023). "Phenotypic plasticity promotes lymph nodes metastasis and drug resistance in lung squamous cell carcinomas", *Heliyon* 9.4, e14614. DOI: `10.1016/j.heliyon.2023.e14614`.

[57] Shi, Z.-D., K. Pang, Z.-X. Wu, Y. Dong, L. Hao, J.-X. Qin, W. Wang, Z.-S. Chen, and C.-H. Han (2023). "Tumor cell plasticity in targeted therapy-induced resistance: mechanisms and new strategies", *Signal Transduct. Target. Ther.* 8.1, p. 113. DOI: `10.1038/s41392-023-01383-x`.

[58] Sharma, S. V. et al. (2010). "A Chromatin-Mediated Reversible Drug-Tolerant State in Cancer Cell Subpopulations", *Cell* 141.1, pp. 69–80. DOI: `10.1016/j.cell.2010.02.027`.

[59] Goldman, A., B. Majumder, A. Dhawan, S. Ravi, D. Goldman, M. Kohandel, P. K. Majumder, and S. Sengupta (2015). "Temporally sequenced anticancer drugs overcome adaptive resistance by targeting a vulnerable chemotherapy-induced phenotypic transition", *Nat. Commun.* 6.1, p. 6139. DOI: `10.1038/ncomms7139`.

[60] Ramirez, M. et al. (2016). "Diverse drug-resistance mechanisms can emerge from drug-tolerant cancer persister cells", *Nat. Commun.* 7.1, p. 10690. DOI: `10.1038/ncomms10690`.

[61] Vander Velde, R. et al. (2020). "Resistance to targeted therapies as a multifactorial, gradual adaptation to inhibitor specific selective pressures", *Nat. Commun.* 11.1, p. 2393. DOI: `10.1038/s41467-020-16212-w`.