Distinct bacterial and protist plankton diversity dynamics uncovered through DNA-based monitoring in the Baltic Sea area

Authors:

Krzysztof T Jurdzinski[1], Meike AC Latz[1,2], Anders Torstensson[3], Sonia Brugel[4,5], Mikael Hedblom[3], Yue O O Hu[1], Markus Lindh[3], Agneta Andersson[4,5], Bengt Karlson[3], Anders F Andersson[1,*]

Affiliations:

1. *KTH Royal Institute of Technology, Department of Gene Technology, Science for Life Laboratory, Stockholm, Sweden*
2. *University of Copenhagen, Department of Plant and Environmental Sciences, Frederiksberg C, Denmark*
3. *Swedish Meteorological and Hydrological Institute, Oceanographic Services, Västra Frölunda, Sweden*
4. *Umeå University, Department of Ecology and Environmental Sciences, Umeå, Sweden*
5. *Umeå Marine Sciences Centre, Umeå University, SE-905 71, Hörnefors, Sweden*

*Corresponding author: Anders F Andersson (anders.andersson@scilifelab.se)

Abstract:

Planktonic microorganisms in coastal waters form the foundation of food webs and biogeochemical cycles while exposed to pronounced environmental gradients, especially brackish salinities. Yet, commonplace ecological assessment overlooks most of their diversity. Here, we analyzed the protist and bacterial diversity from new and publicly available DNA metabarcoding data collected alongside the Swedish marine monitoring program. We show that salinity, unlike other environmental factors, had a stronger effect on bacterial than protist community composition. The seasonality of protist but not bacterial families showed high geographic variation. Bacterial alpha diversity increased with dissolved inorganic nitrogen, while protist alpha diversity was highest in near-marine salinities. Microbial community composition patterns displayed interannual stability despite technical differences affecting the detection of rare taxa. Co-occurrence analysis identified clusters of potentially interdependent microorganisms.

Bayesian modeling showed that the same bacterial lineages were less likely than
protists to occur in both lower (<9 PSU) and higher (>15 PSU) brackish salinities. We
propose that protists are less ecologically sensitive to salinity due to the disconnection
of basic metabolic processes from the cell membrane through compartmentalization.
Ultimately, incorporating DNA metabarcoding into an environmental monitoring program
allowed us to connect ecological and biogeographic processes with understudied taxa
and biodiversity dynamics.

Introduction:

Microorganisms play indispensable roles in biogeochemical cycles[1]. Planktonic bacteria
and protists account for around half of primary production on Earth[2], with a
disproportionally high contribution from coastal ecosystems[3]. More than a third of the
human population lives near these ecosystems[4] and depends on them[5,6], while these
environments experience extreme anthropogenic pressures[5,7]. Finally, coastal zones
comprise multiple land-to-sea environmental gradients, including a full range of brackish
salinities, leading to an immense diversity of habitats[5,8].
Baltic Sea and adjacent Kattegat and Skagerrak (hence "Baltic Sea area," Fig.
1a) are among the best-studied coastal areas[9]. Recent formation (8000 years ago[10])
and pronounced environmental gradients and barriers make the Baltic Sea a model for
studying eco-evolutionary processes driven by dispersal and adaptation[11].
Simultaneously, its early exposure to anthropogenic pressures, especially
eutrophication, can foreshadow the processes awaiting other coastal regions[12]. Finally,
the Baltic Sea area has a long record of microbial research. With the earliest data
coming from the XIX century[13], regular microscopy-based phytoplankton monitoring has
been run by multiple countries in the region for decades[14] and since the 1980s[15] in
Sweden[15]. At the same time, the Baltic Sea area was one of the first brackish waters in
which bacteria and protists were surveyed using DNA metabarcoding[16–18] and
metagenomics[19,20]. Still, most of the unknown diversity in the Baltic Sea area is believed
to be microbial[21]. The research so far has mainly focused on groups of phytoplankton
and metazooplankton[21], while the roles of other microorganisms on ecosystem
functioning are less understood.
Globally, salinity is one of the strongest factors structuring bacterial[22–24] and
protist[25] communities. However, the importance of salinity has been, to a large extent,
inferred from the divide between freshwater and marine environments[26–28]. Our previous
research shows brackish waters host unique bacterial species[20,29]. Both microscopy-
and DNA-based surveys in the Baltic Sea area suggest pronounced shifts in surface
water microbial community composition (beta diversity) along the horizontal salinity
gradient[17,18,30–32]. Moreover, bacterial communities change more across the salinity

gradient than between seasons[33]. However, whether protists are as ecologically
sensitive to salinity as bacteria is less known.
The number of aquatic animal species tends to reach a minimum within the
brackish spectrum across salinity gradients[21,34]. For phytoplankton, patterns with a
maximum[30] and minimum[31] species richness (alpha diversity) within brackish waters
have been reported. DNA metabarcoding transect surveys of plankton showed no clear
patterns for bacteria[17] and major eukaryotic taxa[18]. At the same time, bacterial alpha
diversity has been observed to be lower in winter than in summer in the Baltic Sea[33,35],
consistently with patterns observed in marine environments[36–38]. In contrast,
microscopy-based analysis of phytoplankton alpha diversity has shown different
seasonal patterns in low and high salinities in the Baltic Sea area[32]. Little is known
about the factors driving the seasonal patterns or confounding diversity and community
composition changes along the salinity gradient.
Here, we present the first in-detail spatiotemporal analysis of bacterial and protist
diversity across the Baltic Sea area. We base the analysis on a recently published DNA
metabarcoding dataset[39], as well as new data released here, together corresponding to
398 samples We show how protists and bacteria differ in their ecological sensitivity to
salinity, compare seasonal patterns of closely related species in different regions, and
analyze drivers of taxonomic richness. We propose an explanation of the contrasting
bacterial and protist diversity patterns based on fundamental physiological differences
between the groups. Finally, we use co-occurrence data to identify core microbiomes
characteristic of overlapping conditions within the Baltic Sea area.

Results and discussion

Processing of metabarcoding data for ecologically relevant
metabarcoding-based analyses

We based the first part of our analyses on a publicly available 16S and 18S
metabarcoding dataset, covering stations across broad environmental gradients
regularly sampled throughout a year[39]. After filtering procedures, we used the data from
246 independent samples. They came from 18 locations, extending from Bothnian Bay
(minimal salinity = 2.00 PSU) to Skagerrak (maximal salinity = 33.92 PSU), sampled at
least six times between January 2019 and February 2020 (Fig. 1a).
From the 16S amplicon sequence variants (ASVs), we chose only those
classified as *Bacteria* and excluded plastids and mitochondria. From the 18S
metabarcoding results, we excluded ASVs classified as animals (*Metazoa*), *Fungi*, land
plants (*Embryophyceae*), and macroalgae (Rhodophyta, *Phaeophyceae*, *Cyanidiales*).
We thus treated the remaining 16S ASVs as bacteria and 18S ASVs as unicellular
eukaryotes, i.e., protists.

We clustered the ASVs by simultaneous genetic and distribution similarity, using
distribution-based operational taxonomic unit-calling (dbOTU[40]; we thus call those
taxonomic units dbOTUs). This method groups variants corresponding to different
marker gene copies within one organism and other ecologically indistinguishable intra-
species variation. Therefore, dbOTUs should correspond to ecologically coherent
lineages unless the sequenced regions are identical between organisms with distinct
ecologies.
In multiple analyses, we used family-level annotation to either group or
characterize dbOTUs. This choice was mainly motivated by family being the lowest
taxonomic level at which more than half of dbOTUs (and protist reads) were annotated
(Supplementary Fig. S1c-f). Additionally, in experimental conditions, bacterial
community assembly processes tend to converge at the family level but are more
stochastic at lower taxonomic levels, suggesting ecologically relevant functional
coherence[41].
Finally, a controlled amount of synthetic DNA amplifiable by the metabarcoding
primers, i.e., spike-in, was added to the samples. This approach aims to obtain absolute
values (copy number per volume sample) for the marker gene (16S/18S rRNA), which
better estimates absolute abundance than relative counts[42]. Thus, it can uncover
blooming and succession patterns obscured from relative abundance analysis[43]. At the
same time, the spike-in approach is highly sensitive to differences in sample preparation
and amplification[42]. This can lead to noisier results, especially in environmental
monitoring, as samples are collected by multiple researchers working in variable
conditions.

Regionally distinct seasonal dynamics of major bacterial and protist taxa

Major bacterial (Fig. 1b) and protist (Fig. 1c) clades displayed distinct seasonal
abundance dynamics. Both their abundance and seasonal patterns differed between
Skagerrak, Kattegat, and the three basins of the Baltic Sea included in this study (Baltic
Proper, Bothnian Sea, Bothnian Bay).
We observed an increased abundance of cyanobacteria as early as May in the
Baltic Proper (Fig. 1b), consistent with microscopic observations[44]. The increase in
cyanobacterial abundance was smaller and peaked later moving northwards within the
Baltic Sea. In the Baltic Proper, both dinoflagellates and *Gyrista* (the protist subdivision
including diatoms), peaked in abundance in March (Fig. 1c), as observed using
microscopic methods[44,45]. The dynamics of *Dinoflagellata* and *Gyrista* differed vastly
between basins. However, in all cases except for Bothnian Bay, those groups increased
in abundance in the first half of the year. Furthermore, in Kattegat and Skagerrak, there
was an additional dinoflagellate peak in September, corresponding to autumn blooms.
Haptophyta generally displayed increased abundance between May and August.

Cryptophyta were abundant in the Baltic Sea, with higher abundance in the second half
of the year than in the first.
We also observed geographic differences in abundance and seasonal dynamics
of major non-phytoplankton groups. *Bacteroidia* (Fig. 1b) peaked in abundance in May
across the Baltic Sea, consistently with previous reports[35,46], but displayed less clear
seasonal dynamics in Skagerrak and Kattegat. *Actinomycetia* and *Acidimicrobiia* were
abundant in the Baltic Sea but not in Kattegat/Skagerrak, consistent with *Actinobacteria*
abundance patterns from transect-based studies[17–19]. Additionally, *Acidimicrobiia*
decreased in abundance between May and July, a dynamic that was more pronounced
moving southwards in the Baltic Sea. Finally, contrary to previous results from a
coastline station[35,46], *Alphaproteobacteria* and *Gammaproteobacteria* comprised a
stable proportion of the whole community throughout the year in all basins but Kattegat
(Supplementary Fig. S2a). Still, consistently with metabarcoding[17] and metagenomic
transects[19], *Alphaproteobacteria* and *Gammaproteobacteria* were more abundant in
Skagerrak and Kattegat than in the Baltic Sea.
Overall, we observed geographic differences in both total abundance and
seasonal dynamics of major bacterial and protist taxa. There were some latitudinal
shifts, e.g., for cyanobacteria or *Acidimicrobiia*, and apparent differences between
Kattegat/Skagerrak and the Baltic Sea. Compared to relative abundance-based
analysis, spike-in correction added ecologically relevant information, as best exemplified
by the March dinoflagellate bloom in Baltic Proper (Fig. 1c) being indistinguishable
based on the relative abundance of *Dinoflagellata* (Supplementary Fig. S2b). Still, the
spike-in approach entailed substantial random noise, especially in less sampled basins
(Fig. 1b-c). Relative abundances better captured gradual succession patterns,
especially for bacteria (Supplementary Fig. S2a). These gradual changes likely reflect
sensitivity to factors other than the productivity of the system. Thus, the best
normalization approach depends on the goals of the analysis and suitable statistical
methods, at the very least until the consistency of absolute abundance quantification
improves.

176
Fig. 1 | Spatiotemporal changes in abundance of major microbial taxa, 2019-2020 a. A
map of 18 sampling locations labeled by the name of the station and size-coded by the number
of samples collected over a period of 13 months in 2019 and early 2020. **b-c.** The numbers of
rRNA marker gene copies per liter annotated to **b.** bacterial classes and **c.** protist subdivisions
averaged over each month and Baltic Sea area basin (as marked in **a.**). The values are based
on the number of reads relative to spike-in reads, and the amount of spike-in added per volume
of sample. Only the classes/subdivisions which, on average, corresponded to >0.01 of reads
across samples are shown. Above each bar, the number of samples collected in the respective
month and basin is given. Note: Two samples (January in Bothnian Bay and September in
Bothnian Sea) are removed for protists, due to spike-in detection failure. **d.** The distribution of
measurements of major physicochemical and geographic parameters for each basin. DIN -
dissolved inorganic nitrogen.

Salinity has a dominant impact on bacterial but not protist community composition

The dissimilarities of bacterial and protist communities between the same pairs of samples were roughly correlated ($R^2 = 0.58$). However, the protist communities were generally less similar to each other than bacterial ones (Supplementary Fig. S3a), as has been reported for picoeukaryotes and bacteria in ocean surface waters[47].

Principal Coordinate Analysis (PCoA) showed distinguishable geographic and seasonal structuring of bacterial and protist communities (Supplementary Fig. S4). To disentangle the environmental factors driving those structuring effects, we performed distance-based Redundancy Analysis (dbRDA). We focused on five fundamental environmental factors: salinity, temperature, day length (i.e., daytime length), dissolved inorganic nitrogen (DIN), and phosphate. These factors did not strongly correlate with each other ($R^2 < 0.4$, Supplementary Fig. S5). The dbRDA-based variation partitioning showed that salinity could explain almost half of the variation in bacterial community composition (Fig. 2a). The effect of salinity on protist community composition was much smaller, while temperature and nutrients were of greater importance than for bacteria (Fig. 2b). These results contradict the proposed rule that the distribution of smaller organisms is less limited by environmental factors than that of bigger organisms[48], and in particular that bacteria are less environmentally filtered than protists[49].

Nevertheless, for both bacterial and protist communities, there was a detectable salinity divide between the Baltic Sea and Kattegat/Skagerrak, and a more gradual latitudinal shift within the Baltic Sea (Fig 2c-d). The latter corresponded to decreasing salinity and phosphate-to-nitrogen ratios with increasing latitude (Fig. 1d). We connected those spatial changes in beta-diversity, calculated from ASV-based dissimilarities, with high relative abundance of specific microbial families (Fig 2c-d). For example, typically freshwater *Nanopelagicaceae* (acI *Actinobacteria*)[50] and *Limnocylindraceae* (*Chloroflexi*)[51] were associated with the lowest salinity samples, while commonly marine *Rhodobacteraceae*[52] and *Thioglobaceae*[53] were at the other end of the salinity spectrum (Fig 2c). Overall, many abundant and unevenly spatially distributed bacterial and protist families that we identified have unknown metabolic capabilities and preferred habitats. Our results suggest their potentially crucial ecological importance and point to candidate niche-defining factors.

**Fig. 2 | Environmental factors explaining geographic and seasonal community
structuring and the most correlated families of bacteria and protists a-b.** Variance
partitioning of bacterial (**a.**) and protist (**b.**) beta diversity, based on Bray-Curtis distances and
distance-based Redundancy Analysis (dbRDA). **c-d**. Partial dbRDA conditioned on seasonality
(sine and cosine of the day of the year), highlighting bacterial (**c.**) and protist (**d.**) geographic
community structuring. **e-f**. Partial dbRDA conditioned on salinity and latitude, highlighting
bacterial (**e.**) and protist (**f.**) seasonal community structuring. **c-f**. Weighted averages of families
deviating (Bonferroni corrected P < 0.05) were placed on the plot, with values rescaled by a
factor of 0.5. Percentages of variation explained by each dbRDA component are given in
brackets by the axes labels. DIN - dissolved inorganic nitrogen. Salinity is given in practical
salinity units (PSU).

Closely related bacteria but not protists follow analogous seasonal dynamics across the salinity gradient

Both bacterial and protist communities underwent gradual community composition changes throughout the year in all the analyzed basins of the Baltic Sea area (Fig. 2e-f, Supplementary Fig. S6). Diverse groups significantly contributed to the seasonal patterns in beta diversity, ranging from known seasonally blooming diatoms *Thalassiosiraceae* and *Skeletonemaceae*[54] to multiple uncharacterized *Bacteroides* families.

In all the basins, picocyanobacteria (*Cyanobiaceae*) were associated with high temperatures (Fig. 3a, Supplementary Fig. S6a). The rise in picocyanobacterial abundance was followed by blooms of filamentous heterocyst-forming cyanobacteria (*Nostocacaea*) in the Baltic Proper and the Bothnian Sea (Fig. 3a-b). Expectedly, *Nostocacaea* thrived in DIN-depleted conditions (Fig. 3b-c). The nitrogen fixed by filamentous cyanobacteria leaks to picocyanobacteria, which has been suggested to promote the growth of the latter group in DIN-deficient conditions[55]. However, we observed a high abundance of *Cyanobiaceae* in DIN-depleted Skagerrak and Kattegat, where no filamentous cyanobacteria blooms occurred (Fig. 3a-c). Thus, other mechanisms, potentially other diazotrophs or efficient recycling of organic nitrogen, can sustain picocyanobacterial populations under nitrogen-depleted conditions. Finally, only in Baltic Proper did we observe an increase in the abundance of *Cyanobiaceae* in winter. Therefore, previously identified winter increases of certain *Synechococcus* sp. in Baltic Proper[56] may be specific to this region.

Similar to cyanobacteria, the abundance of haptophyte families *Prymensiaceae* and *Chrysochromulinaceae* were associated with high temperatures (Fig. 2f). However, we could not connect any family to similar conditions across all the basins—not only among haptophytes but among protists in general (Supplementary Fig. S6b). This suggests that despite stronger geographic structuring of bacterial communities along the salinity gradient (Fig. 2a-d), protist seasonal succession patterns differ more between regions. The aforementioned haptophytes may best exemplify this heterogeneity. *Prymensiaceae* were by far most abundant in summer in Baltic Proper (Fig. 3d), while *Chrysochromulinaceae* bloomed at a similar time in Bothnian Bay (Fig. 3e). These families also increased in abundance around June in Skagerrak, largely explaining the increase in overall haptophyte counts (Fig. 1c). The same cannot be said about Kattegat, where haptophytes were most abundant in May (Fig. 1c). Both in Skagerrak and Kattegat, *Gephyrocapsa huxleyi* (formerly *Emiliania huxleyi*) is known to form summer/late spring blooms[57]. As *Gephyrocapsa oceanica* and *G. huxleyi* are indistinguishable using 18S metabarcoding[58], we investigated the abundance of *Noelaerhabdaceae*, a family containing these species. Expectedly[59], we observed this group to be abundant in nutrient-deficient conditions, with the inorganic N:P ratio much lower than the Redfield ratio of 16:1, and only in Kattegat/Skagerrak (Supplementary

Fig. S7a, d). Still, it did not explain Kattegat's high abundance of haptophytes in May.
We ultimately found a May bloom of *Phaeocystaceae* in Kattegat (Supplementary Fig.
S7e), which was short-lived, as is typical of *Phaeocystis*[57].

Overall, in each basin of the Baltic Sea area (except the Bothnian Sea), a
different set of haptophyte species bloomed under warm, nutrient-depleted conditions.
Ultimately, which taxa within *Haptophyta* flourish affects ecosystem functioning, as they
differ in their toxicity[57], as well as roles in carbon and sulfur cycles[60]. So is the case for
many other groups, including dinoflagellates and diatoms, which can be similarly
investigated using DNA monitoring data in focused studies.

BACL11, *Bacterioides* with streamlined genomes and low GC content[20], was the
only family other than *Cyanobiaceae* associated with seasonal changes in community
composition across all the studied regions (Supplementary Fig. S6a). However,
opposite to *Cyanobiaceae,* BACL11 thrived in cold waters (Fig. 2e, Supplementary Fig.
S6a). Our observations diverge from the previously suggested dinoflagellate-bloom
association of BACL11[20]. Complex seasonal patterns (Fig. 3g) of BACL11 suggest
dependence on a combination of factors or a factor not investigated here. On the other
hand, nitrite-oxidizing *Nitrospinaceae*[61], a winter-associated family (Fig. 2e,
Supplementary Fig. S6a), was clearly influenced by DIN dynamics (Fig. 3c, h). Still, their
abundance usually decreased before DIN depletion, as well as during the summer in
the Bothnian Bay, where nitrogen availability remained relatively high throughout the
year. A better understanding of the growth-limiting factors of *Nitrospinaceae* is needed
to explain these patterns.

Finally, *Flavobacteriaceae* showed seasonal patterns linearly uncorrelated with
the investigated environmental factors (Fig. 2e, Supplementary Fig. S6a). However, this
Bacteroides family rose in abundance at the periods corresponding to a rapid decrease
in phosphate availability (Fig. 3f, i), as did total *Bacteroidia* abundance (Fig. 1c).
Recently, it has been suggested that *Bacteroidia*, and in particular *Flavobacteria*, gain a
competitive advantage under limited phosphate availability by efficiently mobilizing
organic phosphorus 61[62,63]. Our results further support this notion and suggest that
phosphorus availability contributes to brackish *Flavobacteriaceae* seasonality. Still, this
adaptation strategy appears limited, as less *Flavobacteriaceae* was detected during
phosphate-minimum conditions in summer. Apart from phosphorus mobilization,
Flavobacteriaceae are adapted to efficiently utilize algal biomass[64], which might explain
why they are abundant under decreasing but not increasing phosphate availability.

307
Fig. 3 | Seasonal dynamics of selected taxa and nutrients. a-b., d-e., g-i. Numbers of rRNA
marker gene copies annotated to families associated with beta diversity seasonal dynamics
(see Fig. 2e-f). Families of cyanobacteria (**a-b**.), haptophytes (**d-e.**), *Bacteroidia* (**g-h**.), and
Nitrospinia (**i.**) are shown. The names of bacterial families are colored purple and of protist
families green. **c, d.** Concentration of inorganic nitrogen (**c.**) and phosphate (**f.**) across the
samples in 2019-2020. The lines correspond to the rolling mean for each basin, with values
averaged for measurements two weeks before and after each date. The same legend (**a.**)
applies to all the panels, with basins ordered from lowest to highest salinity (top to bottom). DIN
- dissolved inorganic nitrogen.

Inorganic nitrogen availability associated with bacterial alpha diversity

Bacterial alpha diversity declined and increased rapidly (Fig. 4a) at the beginnings and
ends of nitrogen-depletion periods (Fig. 3c). Consistently, DIN was the environmental
factor by far best-explaining variation in bacterial richness using a linear model (Fig. 4c).

Many bacteria use inorganic nitrogen as an energy source or an electron
acceptor in nitrification and denitrification processes. In the previous analyses, we found
the nitrate oxidizers *Nitrospinaceae* associated with high-DIN winter conditions in
multiple basins. Among dbOTUs preferentially found in high DIN (>0.5 μmol) conditions,
a few families other than *Nitrospinaceae* were overrepresented (Fig. 4e). Among them
was UBA2979 (corresponding to *Ca. Bathosphaeria*[65], *Chloroflexota*) and
Rhodospirillaceae, with representatives containing genes for nitrate reduction[51,66].
Indeed, the latter family, while comprising diverse metabolic strategies, includes major
marine denitrifiers[66,67]. Moreover, representatives of UBA8477 (*Marinisomatota*, 100%
barcode (ASV4915) sequence identity with clade SHBH1141[68] representatives
confirmed using BLAST[69]) express genes for nitrous oxide reduction[68]. While
denitrification is an anaerobic process, anoxic microenvironments can form in organic
particles suspended in surface waters[70] and sustain denitrification[71]. Indeed, such
oxygen-depleted microniches have been found during cyanobacterial blooms in the
Baltic Sea but with low denitrification rates connected to insufficient nitrate
concentrations in the surrounding water[71]. Still, a spatiotemporal metagenomic study is
needed to confirm the connection between the higher taxonomic diversity and the
nitrogen-related metabolic capabilities.
Inorganic nitrogen availability opens up niches for nitrifying and denitrifying, thus
potentially directly contributing to increased bacterial dbOTU richness in winter. Still,
other factors are likely also involved. It has previously been suggested that lower
bacterial alpha diversity in summer than in winter results from the effects of
cyanobacterial blooms on metabarcoding data[33]. Cyanobacteria would thus dominate
the 16S rRNA pool, decreasing the probability of detecting other taxa, even if those
taxa's abundances remain stable. However, bacterial dbOTU correlated much more
weakly with the relative abundance of cyanobacteria than with DIN in our dataset
(Supplementary Fig. S8). Neither could the higher alpha diversity easily be explained by
autumn water mixing, as previously suggested[33], as bacterial dbOTU richness remained
high throughout the winter (Fig. 4a). Also, day length, strongly correlated with bacterial
alpha diversity in a marine time series[36,38], had smaller explanatory power in our
spatiotemporal analysis, mainly overlapping with covarying factors (Fig. 4c). Finally,
nutrient depletion comes from and is maintained by subsequent blooms of different
protist and bacterial phytoplankton groups. Thus, DIN-associated low dbOTU richness
might also be indicative of the biological effects of these disrupted conditions. However,
we would then expect a similar DIN-dependency among protists, which is not the case
(Fig. 4b and d).

358
Fig. 4 | Seasonal dynamics of alpha diversity across regions a-b. Mean number of bacterial
(**a.**) and protist (**b.**) dbOTUs found across 100 refraction iterations (dbOTU richness). The lines
for the distinguished Baltic Sea area basins correspond to the rolling mean, with values
averaged for measurements two weeks before and after each date. **c-d.** Variance partitioning of
bacterial (**c.**) and protist (**d.**) dbOTU richness. **e.** Number of dbOTUs found significantly (FDR <
0.01 based on Wilcoxon test) more often in high DIN (> 0.5 μmol/l) and low DIN (≤ 0.5 μmol/l),
respectively, in the Baltic Proper. The differentially present dbOTUs are grouped by major
bacterial classes (classes with relative abundance in the whole dataset > 0.01). Families
overrepresented among the differentially present dbOTUs (FDR < 0.01 based on
hypergeometric test) have been named. If the family came from an unlabeled class, the name of
the class is given on the left-hand side of the family's name. **f.** Same as e., but comparing protist
dbOTUs more often found in high (>10 PSU) and low (≤ 10 PSU) salinity, grouped by major
protist subdivisions.

Higher protist alpha diversity in near-marine salinities

Protist dbOTU richness was strongly associated with salinity and was especially high in Kattegat and Skagerrak (Fig. 4b, d). Similarly, metazooplankton and fish alpha diversity in Skagerrak and Kattegat is higher than in the Baltic Sea, due to many marine species in higher brackish salinities, outweighing the presence of freshwater species in lower salinities[21]. The Baltic fauna comes mainly from a limited number of local adaptations to brackish conditions following the last glaciation period[11,21]. In contrast, for Baltic Sea bacteria, we recently showed evidence against widespread occurrence of such local adaptation, but rather colonization by a global brackish microbiota[29]. Consistently, both this (Fig. 4a, c) and previous studies[17,18,33] showed no evidence for decreased bacterial diversity in the Baltic Sea compared to Kattegat/Skagerrak. Whether the observed protist diversity pattern originates from a selected group of locally adapted species, as is the case for animals, is yet to be determined. Notably, unlike bacteria[72], marine protist species show a considerable dispersal limitation[28,73]. If the same is true in brackish waters, it would decrease the chances of colonization by brackish protists from distant locations, promoting local adaptation.

Among dbOTUs more often found in high salinity (>15 PSU), there were multiple families of *Syndiniales* ("Dino-Group" clades, Fig. 4f). These parasitic dinoflagellates are known to be more abundant and diverse in marine than freshwater environments[74]. Similarly, all described[75] MAST-3[76], *Florenciellales*[77,78], and *Phaeocystis*[79] species are marine (and all *Phaeocystaceae* we found were annotated to the genus *Phaeocystis*). As *Cyclostephanos*, *Stephanodiscus*, and *Discostella* are annotated to *Stephanodiscaceae* in the PR2 database[75], it leaves *Thalassiosiraceae* as a family with a majority of representatives from marine environments[80,81]. Systematic analyses with higher taxonomic resolution are needed to determine to what extent marine protists populate the higher brackish salinity ranges, as in Kattegat and Skagerrak.

Temperature was the second factor most strongly associated with protist alpha diversity (Fig. 4d), and in most basins, dbOTU richness increased from March to September (Fig. 4b, Supplementary Fig. S7b). This pattern might, however, reflect a slow recovery after spring blooms. Still, temperature was also the factor best explaining protist beta diversity (Fig. 2b), and our results therefore suggest that both common and rare protists might be more susceptible to the direct effects of climate change than bacteria.

Interannual stability of microbial community composition

To complement the 2019-2020 data analyzed above, we performed metabarcoding on stored DNA extracted from samples collected between 2015 and 2017 (Fig, 5a.b). The new dataset, corresponding to 143 samples, covered the southern range of the 2019-2020 dataset (Fig. 1a, Fig. 5a). Similarly to the 2019-2020 dataset, it mainly

corresponded to a 13-month long time-series, between February 2016 and March 2017,
with a few extra samples from 2015 from one station (Släggö, Fig. 5a). There were
minor differences in sample processing between the two datasets, most notable being:
DNA extraction kit, different Illumina sequencing systems, and lack of spike-in
sequences in the 2015-2017 dataset (see Methods for more details).
Overall, the community composition in the two datasets differed minimally,
following the spatial and seasonal environmental gradients in an analogous manner
(Fig. 5c-f). This apparent interannual stability of community composition is consistent
with previous observations from the Baltic Sea[46,82] and other waters[83]. Finally, the
similarity of communities between the two datasets shows that their technical
differences have little impact on the observed general community composition (Fig. 5e-
f).
Only a two-year break separated the datasets. The seasonal patterns likely
change over longer time scales, as, e.g., climate change has altered the phytoplankton
bloom dynamics in the Baltic Sea over decadal scales[84]. As climate change
accelerates[85] and nutrient inputs are being decreased in the Baltic Sea[86], the microbial
communities may shift in unforeseen directions. These changes may result not only
from shifting abiotic factors but also from biotic interactions, as, e.g. invasive species
introduction can alter microbial community dynamics[87]. A reproducible monitoring
scheme encompassing diverse taxa, achievable with DNA-based methods, is crucial for
understanding and informed responses to unforeseen changes.
In contrast to community composition, alpha diversity differed significantly ($P <$
10^{-14} in both cases) for both bacteria and protists (Fig. 5g-h). We doubt this increase,
amounting to hundreds of dbOTUs on average, represents an actual biological change.
Instead, it likely originates from the technical differences between the datasets. We
based our alpha-diversity estimates on rarefaction as it performs best in removing
sequencing depth biases[88] (expected due to different Illumina sequencing systems used
for the datasets). Moreover, in both datasets, rarefaction curves were saturated,
suggesting that extra reads would not change alpha diversity much (Supplementary Fig.
S9). Thus, another technical difference likely caused the dbOTU difference between
datasets. Its identification would require additional investigation. Ultimately, our results
suggest more caution is needed when comparing alpha diversity than beta diversity
across datasets.

Fig. 5 | Bacterial and protist diversity across datasets collected in 2015-2017 and in 2019-

2020. a. A map of 12 sampling locations labeled by the number of samples collected between

August 2015 and March 2017 (the 2015-2017 dataset), with all the 2015 samples coming from

Släggö in Skagerrak **b.** The distribution of measurements of major physicochemical and

geographic parameters for each basin in the 2015-2017 dataset. **c-d**. Principal Coordinatet

Analysis (PCoA) based on Bray-Curtis dissimilarities of bacterial (**c.**) and protist (**d.**)

communities across the 2015-2017 and 2019-2020 datasets. **e-f.** Variance partitioning of

bacterial (**e.**) and protist (**f.**) diversity, based on Bray-Curtis distances and distance-based

Redundancy Analysis (dbRDA). Dataset stands for a dummy variable differentiating the

samples collected in 2015-2017 and 2019-2020. **g-h.** dbOTU richness of bacteria (**g.**) and

protists (**h.**) in the two datasets. Only dbOTU richness values for stations present in both

datasets have been used. Moreover, samples were chosen in pairs, from the same station and

the same month of the year but different datasets. Sample pairs with the lowest distance in

terms of the day in a calendar year were chosen. P values obtained using pairwise Wilcoxon

signed-rank test are given above each plot. DIN - dissolved inorganic nitrogen. dbOTU -

distribution-based operational taxonomic unit.

Clusters of exclusively co-occurring microorganisms point to potential interdependencies

We performed a custom co-occurrence analysis identifying the pairs of dbOTUs most consistently found with but not without each other (FDR < 0.01 for both co-presence and co-absence), i.e., "exclusively co-occurring". These pairs correspond to either unprecedented examples of synchronized environmental filtering or dependence of at least one dbOTU on another. Thus, the clusters of co-occurring taxa we observed (Fig. 6a) represent either convergently evolved niche spaces or webs of interdependent organisms.

We found three clusters of exclusively co-occurring microorganisms (Fig. 6a). All the clusters contained bacteria with streamlined genomes and known metabolic dependencies. *Pelagibacteraceae* (SAR11), HIMB59 (formerly included in the SAR11 clade), D2472, and pelagic *Methylophilaceae* all have unusually small genomes missing essential functions[89–92]. A further, systematic, genome-based analysis is needed to assess whether the other members of the clusters metabolically complement the streamlined bacteria. At the very least, BACL14 (a genus to which *Methylophilaceae* from clusters 1 and 3 belong) and MB11C04 can provide vitamers that *Pelagibacteraceae* depend on[93]. Importantly, despite potentially providing the resources to *Pelagibacteraceae*, pelagic *Methylophilaceae* usually have streamlined genomes[92] (and all BACL14 representatives in GTDB release 09-RS220[94] have genomes smaller than 1.5 Mbp). Overall, the co-occurrence within the clusters might largely correspond to the dynamics of the "Strong Black Queen Hypothesis"[95]. That is, the clusters might represent metabolically interdependent taxa that need to disperse together to establish themselves in a new location. Consequently, the inability of some species to cross the environmental barriers might affect the whole cluster.

The distribution patterns of the exclusive co-occurrence clusters across the salinity barrier were even more pronounced based on abundance (Fig 6b-d) than the presence/absence of dbOTUs (Supplementary Fig. S10), despite the co-occurrence analysis being based on the latter data type. Clusters 1 and 3 showed strong preferences for high and low salinities, respectively. They also consisted almost entirely of bacterial dbOTUs. In contrast, cluster 2 spanned across the entire salinity gradient. However, it was less abundant and incompletely represented in the least saline samples from the northernmost location. The bacteria-to-protists ratio in cluster 2 was 2:1, on par with the ratio for the whole set of dbOTUs considered for this analysis (2.005:1.000). These results further support the notion of a stronger salinity limitation among bacteria than protists. However, if these microorganisms truly are interdependent, the effects might be amplified, as the salinity sensitivity of one dbOTU would affect the others.

The exclusive co-occurrence clusters were generally abundant and present together throughout the year (Fig 6b-d). However, in the warmest season, their geographic ranges decreased (Fig 6b-d). Some representatives were absent even in

the preferred salinities. The seasonal disappearance might be imposed by competition
for depleted nutrients or enhanced grazing on bacteria during the summer[96–98]. The
persistence of other members of the clusters might largely depend on cyanobacteria
providing the functions of the missing bacteria. For example, during the summer,
picocyanobacteria are the main producers of vitamers[93], on which other bacteria
depend.
Overall, our results suggest the existence of groups of potentially interdependent
microorganisms that tend to spread together. Their differential presence and abundance
highlight that the salinity gradient, limiting the ranges of some microorganisms, also
changes the biotic conditions for the rest of the community.
510

Fig. 6 | Co-occurring bacterial and protist dbOTUs. a. Grey lines link pairs of dbOTUs
occurring in and being absent from the same samples (FDR > 0.01 based on beta distribution,
corresponding to the less common dbOTU being present in at least 86% of the samples in
which the more common one was found and absent from all the samples where the more
common one was not found). Each dbOTU is labeled by annotation to bacteria or protists, a
class or subdivision (only taxa corresponding to >1% of all reads are labeled), and family (if
available). The three largest clusters are distinguished and ordered by the number of dbOTUs
belonging to them. Samples from both the 2015-2017 and 2019-2020 datasets were used to
build the network. Only dbOTUs present in at least ten samples were considered for the co-
occurrence analysis. **b-d.** Relative abundance based on spike-in normalized counts summed
over dbOTUs from each cluster across the 2019-2020 dataset. On the top and the right axes of
each plot are density plots based on mean spike-in normalized abundance per month (top) and
per unit salinity (right). The median given above each plot, unlike the means for the density
plots, is based only on samples in which at least one dbOTU from the cluster was present. The
same legend (**d.**) applies to panels b-d. Temperature is given in degrees Celsius (ºC). dbOTU -
distribution-based operational taxonomic unit.

Bacterial dbOTUs are less likely than protist dbOTUs to occur at both high and low brackish salinities

The salinity gradient in the Baltic Sea area contains a characteristic, rapid shift across the Danish straits. Consequently, our data contained geographically proximate samples from lower (<9 PSU) and higher (>15 PSU) brackish salinities (Fig. 1a, Fig. 5a). Our previous analyses showed the separation across this salinity barrier to be the strongest structuring effect on microbial communities, though less so for protists than for bacteria (Fig. 2c-d, Supplementary Fig. S4).

We further investigated whether bacteria are, in general, less likely to cross this salinity barrier between lower and higher salinities. First, the proportion of protist dbOTUs crossing the salinity barrier was significantly higher than the proportion of bacterial dbOTUs (Fig. 7a, Supplementary Fig. S11). Based on comparable subsets of samples, the proportion of both bacterial and protist dbOTUs crossing the salinity barrier was higher in the samples from 2016-2017 than from 2019-2020 (Supplementary Fig. S11b-c). This suggests that the above noted differences in alpha diversity (Fig. 4g-h) may be due to a higher sensitivity of the sequencing approach used for 2015-2017 data, not technical artifacts. Still, we based further analyses on the 2019-2020 dataset, since that allowed us to use spike-in normalized counts as an abundance metric comparable between bacteria and protists. Furthermore, to minimize the effects of other environmental filtering effects (especially climate), we chose the same number of stations and samples across the salinity barrier (see Methods for details).

The proportion of barrier-crossing bacteria and protists may be affected by detection biases coming from differences in relative abundance, length of vegetative season, and environmental filtering by factors other than salinity. Bacteria were on average more abundant in terms of rRNA gene marker copy numbers (Supplementary Fig. S12a). As protists tend to have higher rRNA copy numbers per cell[128], this difference likely translates to even larger difference in cell numbers, which is likely the most relevant abundance measure in the context of dispersal capabilities and detection probability. Meanwhile protists were present in more samples at one side of the salinity barrier (Supplementary Fig. S12b). Overall, being similarly widespread on one side of the salinity barrier, bacteria were more abundant (Supplementary Fig. S12c-f). Still, across the occupancy values, a higher proportion of protists crossed the salinity barrier (Fig. 7b). The same was true across the abundance values, apart from the most abundant outliers, which were barrier-crossing dbOTUs among bacteria but not protists (Fig. 7c). Generally, across different occupancy and abundance values, a higher proportion of protists crossed the salinity barrier (Fig. 7d, Supplementary Fig. 12g-h).

To confirm that protists are more likely to cross the salinity barrier, we build a Bayesian model using a Monte Carlo Markov Chain (MCMC) based on Just Another Gibbs Sampler[99] (JAGS). The model, based on the Bernoulli distribution, calculates the probability of crossing a barrier from logistic regression based on the maximal

occupancy on one side of the barrier, geometric mean abundance, and whether a
dbOTU is a protist or a bacterium. After model selection based on the Watanabe-Akaike
information criterion[100], we chose a version of the model assuming the probability
dependence on the other variables follows a different function for bacteria and for
protists (see Methods for details). The model predicted protists to be more likely to
cross the salinity barrier across the occupancy and abundance values, both on average
(Fig. 7e, Supplementary Fig. S12i-j) and in a majority of MCMC iterations (Fig. 7f). The
probability that protists are more likely to cross the salinity barrier was consistently close
to one except for dbOTUs with low occupancy and high abundance, in which case
detection biases might outweigh the biological signal. Finally, the model consistently
showed a positive relationship between occupancy and the probability of crossing the
salinity barrier (probability = 1 for both bacteria and protists). Meanwhile, an
independent effect of abundance was much less certain (probability of positive effect =
0.526 for protists and 0.614 for bacteria).
Finally, using a similar approach, we built a Bayesian model of total occupancy
as a function of the occupancy at the more occupied side of the salinity barrier, based
on the Poisson distribution. From this model we calculated the predicted occupancy on
the less occupied side of the salinity barrier, and compared it to the experimental values
(Fig. 7g). The model and the data showed that protists not only more often crossed the
salinity barrier, but also were, in general, more widespread in the less preferable salinity
regime.
589

Fig. 7 | Proportion and probability of bacteria and protists (being) found in both high (>15 PSU) and low (<9 PSU) brackish salinity. a. The number and proportion of bacterial and protists dbOTUs found in high (>15 PSU), low (<9 PSU) brackish salinity, and across the salinity barrier. b-c. The proportion of barrier crossing dbOTUs for each value of maximal occupancy at one side of the salinity barrier (b.) and across forty bins of mean abundance (c.). The proportion of all dbOTUs that lie within each bin is also given. The values are rolling means across +/- two bins. d. The difference in the proportion of protist and bacterial barrier crossing dbOTUs. Grey tiles correspond to the values across which either no protist or no bacterial dbOTUs were found. e-g. Results from Bayesian modeling. e. The mean difference in the predicted probability of a protist and a bacterial dbOTU crossing the salinity barrier. f. The probability (proportion of MCMC iterations) that protists are more likely to cross the salinity barrier. g. The occupancy at the less occupied side of the salinity barrier as a function of the occupancy at the more occupied side. The points correspond to means of the experimental values. The lines correspond to the values predicted from the model with 95% confidence intervals. The dashed black line is the identity line. For all the analyses, only the four stations with salinity >15 PSU, and four out of the most saline stations with salinity <9 PSU are chosen. The data for each station is downsampled to the same number of observations. During downsampling, samples closest in time to the samples from the least sampled station were chosen. Only observations from 2019-2020 are included. The abundance values correspond to numbers of rRNA copies per liter, based on the spike-in normalization, and are always given on a logarithmic scale. "Occupancy preferred side" and "occupancy other side" refer to the maximum and the minimum occupancy at one of the two sides of the salinity barrier, respectively.

Proposed explanation of the stronger ecological sensitivity to salinity among bacteria than protists

We propose that compartmentalization makes eukaryotic cells (thus, protists) more acclimatable than prokaryotic ones (thus, bacteria) to a different salinity regime (Fig. 8). In a compartmentalized cell, crucial metabolic processes happen across internal membranes. External ion concentrations and pH do not directly affect these processes. Notably, in eukaryotic cells, the proton motive force used for ATP synthesis is maintained along the inner membranes of mitochondria and plastids. In contrast, prokaryotes usually produce proton motive force across the cellular membrane directly exposed to the external environment. While eukaryotic cytosolic pH is kept at stable, neutral levels[101], marine and brackish waters are buffered and usually slightly alkaline (pH~8)[27,102]. Thus, in the presence of sea salt, some transferred protons are consumed in reactions with (bi-)carbonates or hydroxide. Consequently, many marine bacteria preferably use sodium motive force, i.e., Na^+-gradient across the membrane, for transmembrane transport, motility, and/or ATP production[19,27,103,104]. The competitive advantage of this strategy diminishes with decreasing salinity (in a logarithmic manner, assuming stable intracellular ion concentrations[105]), and so likely does the relative abundance of bacteria using it. Finally, many protists also have additional organelles with osmoregulating functions, especially the contractile vacuole[106].

Fig. 8 | The proposed mechanistic explanation for higher ecological sensitivity to salinity for bacteria than for protists. The cells and their components are drawn schematically and do not accurately represent sizes and shapes. All the cell components, except the protist nucleus and bacterial nucleoid, correspond to the machinery responsible for processes indicated by the arrows leading from them. Statements based on our results are written in italics. This figure was created using images from Togo Picture Gallery[107] Servier Medical Art (http://smart.servier.com).

Conclusions:

We show multiple lines of evidence that salinity, generally, has a stronger impact on the distribution of bacteria than protists. We propose that compartmentalization allows protists to acclimatize to a different salinity regime more easily (Fig. 8). Importantly, as further validation is needed, these considerations apply to withstanding different salinities in a competitive environment, and not necessarily to salinity ranges that isolated cultures can withstand. Nor does the proposed explanation apply to adaptation through long-term evolutionary changes. Finally, our theoretical model could be extended to pH, as we already discussed the bioenergetic aspects in this context (Fig.

8). Consistently, in soil, pH has a dominant effect on the community composition of
bacteria but not fungi[108].
We found that nitrogen-rich waters host higher numbers of bacterial taxa.
Furthermore, we connected this increase in alpha diversity to an overrepresentation of
nitrite oxidizers and denitrifiers. As eutrophication ultimately leads to an increase in
harmful algal blooms and subsequent nutrient depletion[109], the conclusion should not be
that adding nitrogen increases bacterial diversity. Instead, what requires more focus is
the nitrogen cycle-related roles of the rare taxa from nitrogen-rich waters. A monitoring
scheme sensitive to their presence is needed to understand their susceptibility to
environmental disruptions and subsequent impacts. DNA-based methods may well be
the only feasible option at present.
We show geographic differences not only in microbial community composition
but also in the seasonal dynamics of closely related taxa, especially protists. These
differences occur at both broad and more fine-grained taxonomic levels. Thus, time-
series results from multiple and diverse locations should be integrated for
generalizability and proper interpretation. We compared a publicly available DNA
metabarcoding dataset with newly sequenced samples from an earlier monitoring effort.
We observed interannual stability of community composition relative to the
environmental conditions. However, minor technical differences between the datasets
confounded alpha diversity analysis. Thus, a standardized DNA-based scheme is
needed to monitor rare microbial taxa.
Using DNA metabarcoding, we could connect bacterial and protist diversity to the
ecosystem processes in the Baltic Sea. Apart from detecting major already-known
dynamics, we highlight overlooked taxa that may have crucial roles in nutrient cycling in
brackish waters. Moreover, we show how salinity and inorganic nitrogen availability
affect microbial diversity, with conclusions generalizable beyond the coastal waters.
Still, DNA metabarcoding allows only the monitoring of phylogenetic diversity at a
limited resolution. Metagenomics and other omics methods can uncover full-scope
genetic diversity and its functional potential.
Ultimately, as Earth's environments undergo rapid change, microorganisms from
diverse, often understudied groups will be impacted and/or consequential for the future
trajectory of coastal and other ecosystems. DNA-based monitoring covers the
phylogenetic scope needed to understand and respond to those unprecedented
changes.

Methods:

Data collection and sequencing

The 2019-2020 data comes from Latz et al. 2024[39]. The collection and sequencing procedures are described in detail therein. In brief, samples were collected monthly or every second week, depending on the location, as a part of the Swedish National Marine Monitoring Program. The samples for metabarcoding were collected using a depth-integrating hose, covering 0-10m (except 0-5m at RÅNEÅ-1, and 0-20m at B1 and BY31). 500 ml of water from the hose sampling was filtered using 0.22 µm pore size. DNA was extracted using the Zymobiomics™ DNA miniprep kit following a modified protocol[110], and spike-in DNA was added. The DNA was sequenced using Illumina™ MiSeq flow cells. For 16S rRNA gene-based metabarcoding of prokaryotes, primers targeting the hypervariable V3-V4 regions[17], and for 18S metabarcoding of eukaryotes, primers targeting the V4 region[111], were used. Phased primers[112] were used for both 18S (on the forward primer) and 16S (on both primers) metabarcoding. Sequences of the primers and the details of the phasing strategy and sample indexing through a second PCR can be found in Latz et al. 2024[39]. Physico-chemical and geographic data has been obtained from the Swedish Meteorology and Hydrology Insitute's service SHARKweb: https://sharkweb.smhi.se/hamta-data/

      The 2015-2017 data was used for the first time for this study. The samples were stored at -80°C. DNA was extracted in the summer of 2017 and stored at -20°C until February 2023. The sampling and sequencing procedures were analogous to the one for 2019-2020 data, with a few exceptions:

1. Volumes varied from 200ml to 800ml. According to our previous tests run in the same setting, volumes ≥200ml should capture the microbial community structure equally well[39].
2. DNA extraction was performed using Qiagen™ DNeasy PowerWater Kit, and not ZymoBIOMICS™ DNA Miniprep Kit with a small modification[110] as for the 2019-2020 dataset.
3. No spike-in DNA was added.
4. Illumina™ NextSeq, rather than MiSeq system, was used.
5. For 18S, the initial use of 20 cycles for the first PCR amplification was insufficient, leading to many library preparation failures. It was then re-run with 24 cycles, and the results of both runs were merged after processing the tables by summing up the counts in the output count table.
6. For 18S, reverse primers were also phased, following a sequence (CTACGA)CTTTCGTTCTTGATYRR, with versions of the primer starting from any base in the brackets included in the primer mix.

720 Samples from other projects, which we did not analyze in this study, were sequenced
721 simultaneously and submitted to the European Nucleotide Archive (ENA) all together.

Processing of sequencing data

The data processing for 2019-2020 samples has been described in detail in Latz et al.
2024[39]. In brief, the phased primer sequences were removed from the obtained reads
using a modified version of a cutadapt[113]-based pipeline (original pipeline taken from:
https://github.com/biodiversitydata-se/amplicon-multi-cutadapt). DADA2[114] version
1.18.0 was used to denoise the data, infer amplicon sequence variants (ASVs), and
taxonomically annotate them. For taxonomic annotation of 16S ASVs, we used 16S
sequences from GTDB[115] version R06-RS202-1 corrected for mislabeled sequences
using SATIVA[116,117]. Annotation was additionally conducted using the SILVA
database[118] version 138.1 to identify plastids and mitochondria. The 18S ASVs were
annotated using the PR2 database[75] version 5.0.1.
The 2015-2017 data was analyzed the same way, with the only difference being
that the denoising and ASV inference with DADA2 was performed using the nf-
core/ampliseq pipeline[119] version 2.7.0 instead of an in-house R script. We kept the
DADA2 parameters as in Latz et al. 2024[39].
We further combined the 2015-2017 and 2019-2020 data. We matched ASVs
using exact matching. On the merged ASV-sequence list and count table, we ran an
additional chimera removal step using UCHIME v1[120] and clustered the remaining ASVs
based on their simultaneous genetic and distribution similarity using dbOTU3[40]. Both
these steps were performed using an in-house pipeline available at:
https://github.com/kjurdzinski/chimera_dbOTU_pipeline. Default parameters were used
for UCHIME, and for dbOTU#, only the maximum genetic distance for 16S has been
changed to 3% (instead of the default 10%). The procedure should minimize the
influence of intra-specific variation on our data and further reduce the effects of PCR
amplification and sequencing errors. We called thus obtained taxonomic units dbOTUs
(distance-based operational taxonomic units, as inferred using dbOTU3[40])
For both 16S and 18S, we removed dbOTUs with no annotation at the
phylum/supergroup taxonomic level. From 16S dbOTUs, *Archaea*, plastids, and
mitochondria were removed. Thus, the remaining ASVs should correspond to bacteria.
From 18S ASVs, animals (*Metazoa*), *Fungi*, land plants (*Embryophyceae*), and
macroalgae (Rhodophyta, *Phaeophyceae*, *Cyanidiales*) were removed. Thus, the
remaining ASVs were primarily protists.
We kept spike-in dbOTUs in a separate table and used them only to obtain spike-
in corrected abundance, i.e. the inferred number of rRNA copies per liter. The rRNA
copier per liter were calculated according to the equation:

rRNA copies per liter

$$758 \qquad = \frac{dbOTU\ counts}{spike-in\ counts}$$

$$759 \qquad \times \frac{amount\ of\ spike-in\ added\ [g]}{volume\ water\ sampled\ [l]\ \times\ spike-in\ molar\ mass\ (g/mol)} \times 6.022$$

$$760 \qquad \times 10^{23}$$

We have excluded over- and under-sequenced samples to remove the
amplification errors (including distorted relative abundances) and not to rarefy to too low
a read number during rarefaction. We have excluded three samples with less than
30,000 reads for bacteria, as well as one sample with more than 2,000,000 reads.
Moreover, we have excluded two samples from 2019-2020 with more than 200,000
protist reads (this number of reads was normal for 2015-2017 data since Illumina™
NextSeq was used instead of MiSeq, yielding overall higher numbers of reads).
Additionally, one sample (20191015_263953_1) with suspected contamination has
been excluded.
For comparability of results for protists and bacteria, we used only the samples
with data available from both 16S and 18S amplicon sequencing (excluding the under-
and over-sequenced samples) for all the further analyses. Additionally, in two samples
(station RÅNEÅ-2 on 2019-01-29 (sample ID: 20190129_263856_1) and B7 on 2019-
09-17 (sample ID: 20190917_264450_1)) 18S spike-in addition failed. Those samples
were excluded from analyses in which spike-in normalization was used, except for
bacteria-only focused analyses in Fig. 1b and Fig. 3a-b, g-i.

Abundance, alpha and beta diversity spatiotemporal patterns

For the spatiotemporal analysis of abundance, beta- and alpha diversity across the
samples from 2019-2020, we removed the samples from stations H4, BY29 / LL 19, and
SR3. Those stations were sampled at only a few unevenly distributed times, and their
exclusion/inclusion at certain seasons could obscure seasonal patterns.
All the statistical analyses and data visualization was performed using R version
4.2.2[121]. Inkscape[122] was used to assemble multi-panel figures and to make Fig. 8.
The multivariate analyses were performed using vegan R package[123], and all the
functions in this paragraph can be found there unless another package is explicitly
stated. Bray–Curtis distances between communities were obtained using rarefaction via
the avgdist function. Those distances were used for Principal Coordinate Analysis
(PCoA) and distance-based Redundancy Analysis (dbRDA), performed with,
respectively, capscale (without setting constraints) and dbrda functions. Variance
partitioning was performed using rdacca.hp R package[124]. Families were positioned on
the ordination plots by calculating weighted averages (sppscores function). Only the
families deviating (Bonferroni corrected P-value < 0.05) from normal distribution around

the center of the dbRDA coordinate system were shown on the plots. To find the
corresponding two-dimensional ellipsoid confidence intervals, aspace[125] package was
used. Whenever only samples were visualized, their ordination scores were scaled
using the eigenvalues, and symmetric scaling was used if both the families and the
samples were visualized (using, respectively, "sites" and "symmetric" scaling options of
the plotting functions respectively). For dbRDA and variance partitioning, we used a
subset of 228 (out of 246) samples with all the environmental variables of interest
available. For more informative visualization, to keep the sample structuring visible
despite very high family scores, we divided the family scores in half.
dbOTU richness was calculated as the mean number of dbOTUs detected in a
sample over 100 rarefaction iterations. Variance partitioning was performed using
rdacca.hp R package[124]. dbOTUs differentially present in high/low DIN/salinity were
identified using the Wilcoxon signed-rank test based on presence/absence data. The
families overrepresented among these dbOTUs were found using a hypergeometric test
(Fisher's exact test).

Co-occurrence analysis

810   dbOTUs found in at least ten samples across the 2015-2017 and 2019-2020
811   datasets were considered for the co-occurrence analysis. Their presence/absence data
812   was used to calculate the proportion of samples in which they were co-present relative
813   to the number of samples in which the more widespread dbOTU was found:

814
$$co\_presence_{i,j} = \frac{N_{i\,\&\,j}}{max(N_i, N_j)}$$

815   Where i and j stand for two different dbOTUs, $N_{i\,\&\,j}$ is the number of samples in which
816   both dbOTUs were found, and N_i and N_j are the total numbers of samples in which
817   each dbOTU was present.
818   Additionally, a co-absence value was calculated, also relative to the more
819   widespread dbOTU:

820
$$co\_absence_{i,j} = \frac{N_{\neg i\,\&\neg\,j}}{min(N_{\neg i}, N_{\neg j})}$$

where $N_{\neg i\,\&\neg\,j}$ is the number of samples in which neither dbOTUs was found, and $N_{\neg i}$
and $N_{\neg j}$ are the total numbers of samples in which each dbOTU was absent.
We used the beta-distribution-based test (pbeta in R) to identify the co-
occurrence and co-absence values significantly different from an average pair of
dbOTUs (FDR < 0.01 for each of the two values, in this case corresponding to the less
common dbOTU being present in at least 86% of the samples in which the more
common one was found and absent from all the samples where the more common one

was not found). The co-occurrence network was visualized using the igraph R
package[126].

Proportion salinity barrier crossing dbOTUs and Bayesian modelling

Any dbOTU found among included samples with salinity >15 PSU and among
samples with salinity <9 PSU was regarded as barrier crossing. For balancing out the
number of samples on both sides of the salinity barrier, apart from the four stations with
salinity >15 PSU (Å17, SLÄGGÖ), four out of twelve stations with salinity <9 PSU: BY2
ARKONA, BY5 BORNHOLMSDJ, BY15 GOTLANDSDJ, REF M1V1 (see Fig. 1a for
locations of the stations). First, we chose the four most saline stations from among
those with salinity <9 PSU. Then, we switched BCS III-10 to REF M1V1 (see Fig. 1),
since the latter had more samples, and, unlike other chosen stations from lower
salinities, was located close to the mainland coastline. For each station but the least
sampled (REF M1V1), the samples were downsampled, choosing samples closest in
time to the ones collected at the least sampled stations. Whenever available, those
samples should correspond to the same monitoring cruises.

The Bayesian models were built and implemented in JAGS [99], using the R
package rjags[127]. The parameters, probabilities, and predictions were calculated based
on every fifth out of 10000 MCMC iterations.

The model for probability of crossing the salinity barrier was optimized based on
Bernoulli distribution:

848
$$if\_crossed\_barrier_i \sim Bernoulli(p_i)$$

849 where for each dbOTU i, $if\_crossed\_barrier_i$ is a binary variable = 1 if it is a barrier
850 crossing dbOTU. p_i was calculated according to the formula

$$logit(p_i) = (1 - if\_protist_i) \times (\alpha_{bacterium} + \beta_{abundance\_bacterium} \times log(mean\_abundance_i) + \beta_{occurence\_bacterium} \times log(max\_occurence\_one\_side_i)) + if\_protist_i \times (\alpha_{protist} + \beta_{abundance\_protist} \times log(mean\_abundance_i) + \beta_{occurence\_protist} \times log(max\_occurence\_one\_side_i))$$

where $if\_protist_i$ is a binary variable = 1 if the dbOTU i is a protist. $mean\_abundance_i$ is
geometric mean of the abundance of the dbOTU i in the samples where it was present.
$max\_occurence\_one\_side_i$ is the maximum out of the numbers of samples in which the
dbOTU i was found at one side of the salinity barrier (in salinity <9 PSU or >15 PSU).
The other variables are the parameters of the model.

We compared this model with one assuming the same abundance and
occupancy dependence for bacteria and for protists (one $\beta_{abundance}$ and $\beta_{occupancy}$,
same for protists and bacteria, different $\alpha_{bacterium}$ and $\alpha_{protist}$). For both the model with
different and the same abundance and occupancy dependencies, we also compared
the versions without log-transforming either only the occupancy, or both abundance and

occupancy values. Based on the Watanabe-Akaike criterion[100], the model we used
performed the best.
The model for total occupancy was build based on the Poisson distribution

$$total\_occupancy_i \sim Poisson(\lambda_i)$$

870 where $total\_occupancy_i$ is the total number of samples in which dbOTU i was found. λ_i
871 was calculated according to the formula:

$$\lambda_i = max\_occurence\_one\_side_i + (1 - if\_protist_i) \times (\alpha_{bacterium} +$$
$$\beta_{bacterium} \times log(max\_occurence\_one\_side_i) + if\_protist_i \times (\alpha_{protist} +$$
$$\beta_{protist} \times max\_occurence\_one\_side_i)$$

875 with $if\_protist_i$ and $max\_occurence\_one\_side_i$ having the same meaning as described
876 above. The occupancy at the less occupied side of the salinity barrier was calculated by
877 subtracting $max\_occurence\_one\_side$ from the predicted λ.

Acknowledgements:

We would like to thank Jenny Lycken for the effort put into sampling and sample
preparation for the 2019-2020 dataset. National Genomics Infrastructure (NGI) Sweden,
and especially Elísabet Einarsdóttir, for seamless and adjustable sequencing service.
Karin Garefeld for help with preparing the 2015-2017 samples for sequencing.
Additionally, KTJ would like to thank Maliheh Mehrshad for hosting him in Uppsala
during the critical period of work on this manuscript, as well as suggesting courses
which taught him the skills necessary for the crucial analyses of this study. Ulf Gardin,
for help with the multivariate analyses. Mathew Low and Malin Aronsson for help with
Bayesian modeling.

Study funding information:

This work was funded by the Swedish Agency for Marine and Water Management and
the Swedish Environmental Protection Agency under the grant number NV-03728-17.
AFA was additionally funded by a research grant (2021-05563) from the Swedish
Research Council VR.
For this study, we used computational resources provided by the Swedish
National Infrastructure for Computing (SNIC) through the Uppsala Multidisciplinary
Center for Advanced Computational Science (UPPMAX).

Competing interests:

897 The authors declare no conflict of interest.

Author contributions:

KTJ analyzed the data under the supervision of AFA and with help from MACL and AT. YOOH prepared the DNA for the 2015-2017 dataset. KTJ wrote the first version of the manuscript under supervision of AFA. All authors contributed to conceptualization, interpretation of the data, revised the manuscript, and approved the final version.

Data and code availability:

This is a preliminary version of the manuscript. We are still working on making the data and code publicly available. The 2019-2020 data (reference) is available at the European Nucleotide Archive (ENA) under the study accession number https://identifiers.org/ena.embl:PRJEB55296.

Supplementary figures:

Supplementary Fig. S1 | Taxonomic annotation statistics a-b. Number of different bacterial
(**a.**) and protist (**b.**) taxa found at different taxonomic levels. **c-d**. Proportions of bacterial (**c.**) and
protist (**d.**) taxa annotated at different taxonomic levels. **e-f**. Proportions of bacterial (**e.**) and
protist (**f.**) reads annotated at different taxonomic levels. "Single species" corresponds to an
analysis that provides a species-level annotation only if the query sequence has identical
matches to database sequences of a single species. "Multiple species extra" is the extra
number/proportion of annotated species when exact matches to multiple species are allowed.

918
Supplementary Fig. S2 | Spatiotemporal changes in abundance of major microbial taxa,
2019-2020 a-b. The proportion of rRNA marker gene reads annotated to **a.** bacterial classes
and **b.** protist subdivisions averaged over each month and Baltic Sea area basin. Only the
classes/subdivisions which, on average, corresponded to >0.01 of reads across samples are
shown. Above each bar, the number of samples collected in the respective month and basin is
given.

Supplementary Fig. S3 | Dissimilarity of diversity metrics between bacteria and protist, 2019-2020 a. Beta diversity calculated as Bray-Curtis distances. The golden line represents the best linear fit, while the dashed black line is the identity line. **b.** Number of detected dbOTUs (alpha diversity).

Supplementary Fig. S4 | Principal coordinate analysis (PCoA) 2019-2020· The first four
ordination axes are based on Bray-Curtis distances between bacterial (**a-b**) and protist (**c-d**)
communities. Note: The axes are grouped and ordered by which ones, after visual examination,
correspond to geographic structuring (**a**, **c**) and which ones to seasonal effects (**b**, **d**). This
ordering corresponded to showing ordination axes with decreasing variation explained for
bacteria (**a-b**). However, for protists, the first plot (**c**) shows the second and the third
components, while the second plot (**d**) shows the first and the fourth components.

939
Supplementary Fig. S5 | Correlations between chosen environmental variables for the
2019-2020 dataset a. Pearson correlation coefficients (ρ) on an upper-triangle correlation plot.
b. Coefficients of determination (R^2) on a lower-triangle correlation plot.

943
Supplementary Fig. S6 | Seasonal changes in community composition within
distinguished basins of the Baltic Sea area. distance-based Redundancy Analysis (dbRDA)
based on Bray-Curtis distances between the bacterial (**a.**) and protist (**b.**) communities within
each of the distinguished basins of the Baltic Sea area. Weighted averages of families deviating
(Bonferroni corrected P < 0.05) were placed on the plot, with values rescaled by a factor of 0.5.
Percentages of variation explained by each dbRDA component are given in brackets by the
axes labels.

Supplementary Fig. S7 | The seasonal dynamics of inorganic N:P ratio, salinity, and
additional protist taxa. a. DIN (dissolved inorganic nitrogen) to phosphate concentration ratio
across the 2019-2020 samples. **b-c.** Temperature (**b.**) and salinity (**c.**) across the 2019-2020
samples. **d-e**. Spike-in normalized abundance of the haptophyte families *Phaeocystaceae* and
Noelaerhabdaceae. **f.** Spike-in normalized abundance of the dinoflagellate class *Sindiniales*.
The lines correspond to the rolling mean, with values averaged for measurements two weeks
before and after each date. The same legend (a.) applies to all the panels, with basins ordered
from lowest to highest salinity (top to bottom). PSU - practical salinity units

960
961 **Supplementary Fig S8 | Correlation of bacterial db OTU richness a.** with dissolved inorganic
962 nitrogen (DIN) **b.** relative abundance of cyanobacteria.

Supplementary Fig. S9 | Refraction curves a-b. Based on 16S filtered (only bacteria) read from the 2015-2017 (**a.**) and 2019-2020 (**b.**) datests. **c-d**. Based on 18S filtered (only protists) read from the 2015-2017 (**c.**) and 2019-2020 (**d.**) datests. dbOTU - distribution-based operational taxonomic unit.

968
Supplementary Fig. S10 | Number of dbOTUs from co-occurring clusters (Fig. 6a) found
in each sample. The samples in which all the dbOTUs from a cluster were present are outlined
in red. On the top and the right axes of each plot are density plots based on mean number of
dbOTUs per month (top) and per unit salinity (right). The same legend (**a.**) applies to all the
panels except for the size-coding corresponding to the number of dbOTUs from a cluster.

Supplementary Fig. S11 | The number and proportion of dbOTUs found only in high (>15
PSU), low (<9 PSU) brackish salinity, and across the salinity barrier. **a.** Proportions across
all samples. **b-c.** Proportions for the samples from 2016-2017 (**b.**) and 2019-2020 (**c.**). Included
samples from the two datasets were chosen in pairs, from the same station and the same
month of the year but different datasets. Sample pairs with the lowest distance in terms of the
day in a calendar year were chosen. **d.** Proportions across the two datasets with a balanced
number of stations and observations across the salinity barrier. Only the four stations with
salinity >15 PSU, and four out of the most saline stations with salinity <9 PSU were chosen. The
data for each station was downsampled to the same number of observations. Samples closest
in time were chosen. The same legend applies to all the panels.

985
Supplementary Fig. S12 | The fractions of all dbOTUs, the proportions of barrier-crossing
dbOTUs, and the probability of crossing the barrier across occupancy and abundance
values. a-b. The distribution of maximal occupancy at one side of the salinity barrier (**a.**) and
mean abundance (**b.**). **c.** Rolling mean (+/- two bins) of abundance across the occupancy
values. **d-f.** The fraction of all bacterial (**d.**) and protist (**e.**) dbOTUs, and the difference between
the fraction of protist and bacterial dbOTUs (**f.**). **g-h**. The proportion of barrier crossing bacterial
(**g.**) and protist (**h.**) dbOTUs. **d-h.** Grey tiles correspond to the values across which either no
protist or no bacterial dbOTUs were found. **I-j.** The predicted probabilities of a bacterial (**i.**) and
protist (**j.**) dbOTU crossing the salinity barrier. For all the analyses, only the four stations with
salinity >15 PSU, and four out of the most saline stations with salinity <9 PSU are chosen. The
data for each station is downsampled to the same number of reads. During downsampling,
observations closest in time to the samples from the least sampled station were chosen. Only
observations from 2019-2020 are included. The abundance values correspond to the numbers
of rRNA copies per liter, based on the spike-in normalization, and are always given on a
logarithmic scale. "Occupancy preferred side" is the maximum occupancy at one of the two
sides of the salinity barrier.

References:

1. Falkowski, P. G., Fenchel, T. & Delong, E. F. The microbial engines that drive Earth's biogeochemical cycles. *Science* **320**, 1034–1039 (2008).

2. Field, C. B., Behrenfeld, M. J., Randerson, J. T. & Falkowski, P. Primary production of the biosphere: integrating terrestrial and oceanic components. *Science* **281**, 237–240 (1998).

3. Mathis, M. *et al.* Enhanced CO2 uptake of the coastal ocean is dominated by biological carbon fixation. *Nat. Clim. Chang.* **14**, 373–379 (2024).

4. Cohen, J. E., Small, C., Mellinger, A., Gallup, J. & Sachs, J. Estimates of Coastal Populations. *Science* **278**, 1209–1213 (1997).

5. Williams, B. A. *et al.* Global rarity of intact coastal regions. *Conserv. Biol.* **36**, e13874 (2022).

6. De Valck, J. *et al.* Valuing ecosystem services in complex coastal settings: An extended ecosystem accounting framework for improved decision-making. *Mar. Policy* **155**, 105761 (2023).

7. O'Hara, C. C., Frazier, M. & Halpern, B. S. At-risk marine biodiversity faces extensive, expanding, and intensifying human impacts. *Science* **372**, 84–87 (2021).

8. Ray, G. C. Coastal-Zone Biodiversity Patterns: Principles of landscape ecology may help explain the processes underlying coastal diversity. *Bioscience* **41**, 490–498 (1991).

9. Kuliński, K. *et al.* Biogeochemical functioning of the Baltic Sea. *Earth Syst. Dyn.* **13**, 633–685 (2022).

10. Björck, S. A review of the history of the Baltic Sea, 13.0-8.0 ka BP. *Quat. Int.* **27**, 19–40 (1995).

11. Johannesson, K., Le Moan, A., Perini, S. & André, C. A Darwinian Laboratory of Multiple Contact Zones. *Trends Ecol. Evol.* **35**, 1021–1036 (2020).

12. Reusch, T. B. H. *et al.* The Baltic Sea as a time machine for the future coastal ocean. *Sci*

Adv **4**, eaar8195 (2018).

13. Lundsør, E. *et al.* Marine phytoplankton community data and corresponding environmental

properties from eastern Norway, 1896-2020. *Sci Data* **9**, 767 (2022).

14. Heiskanen, A. S. *et al.* Monitoring strategies for phytoplankton in the Baltic Sea coastal

waters. European Commission. *Institute for Environment and Sustainability Inland and*

Marine Waters. Unit I-21020. Ispra (VA), Italy (2005).

15. SHARK. SHARK - National marine environmental monitoring of Phytoplankton in Sweden

since 1983. The Swedish Meteorological and Hydrological Institute

https://doi.org/10.15468/S3YG8Z (2022).

16. Andersson, A. F., Riemann, L. & Bertilsson, S. Pyrosequencing reveals contrasting

seasonal dynamics of taxa within Baltic Sea bacterioplankton communities. *ISME J.* **4**,

171–181 (2010).

17. Herlemann, D. P. *et al.* Transitions in bacterial communities along the 2000 km salinity

gradient of the Baltic Sea. *ISME J.* **5**, 1571–1579 (2011).

18. Hu, Y. O. O., Karlson, B., Charvet, S. & Andersson, A. F. Diversity of Pico- to Mesoplankton

along the 2000 km Salinity Gradient of the Baltic Sea. *Front. Microbiol.* **7**, 679 (2016).

19. Dupont, C. L. *et al.* Functional tradeoffs underpin salinity-driven divergence in microbial

community composition. *PLoS One* **9**, e89549 (2014).

20. Hugerth, L. W. *et al.* Metagenome-assembled genomes uncover a global brackish

microbiome. *Genome Biol.* **16**, 279 (2015).

21. Snoeijs-Leijonmalm, P. Patterns of biodiversity. in *Biological Oceanography of the Baltic*

Sea (eds. Snoeijs-Leijonmalm, P., Schubert, H. & Radziejewska, T.) 123–191 (Springer

Netherlands, Dordrecht, 2017).

22. Wu, Q. L., Zwart, G., Schauer, M., Kamst-van Agterveld, M. P. & Hahn, M. W.

Bacterioplankton community composition along a salinity gradient of sixteen high-mountain

lakes located on the Tibetan Plateau, China. *Appl. Environ. Microbiol.* **72**, 5478–5485

(2006).

23. Lozupone, C. A. & Knight, R. Global patterns in bacterial diversity. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 11436–11440 (2007).

24. Caporaso, J. G. *et al.* Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proc. Natl. Acad. Sci. U. S. A.* **108 Suppl 1**, 4516–4522 (2011).

25. Singer, D. *et al.* Protist taxonomic and functional diversity in soil, freshwater and marine ecosystems. *Environ. Int.* **146**, 106262 (2021).

26. Logares, R. *et al.* Infrequent marine-freshwater transitions in the microbial world. *Trends Microbiol.* **17**, 414–422 (2009).

27. Walsh, D. A., Lafontaine, J. & Grossart, H.-P. On the Eco-Evolutionary Relationships of Fresh and Salt Water Bacteria and the Role of Gene Transfer in Their Adaptation. in *Lateral Gene Transfer in Evolution* (ed. Gophna, U.) 55–77 (Springer New York, New York, NY, 2013).

28. Burki, F., Sandin, M. M. & Jamy, M. Diversity and ecology of protists revealed by metabarcoding. *Curr. Biol.* **31**, R1267–R1280 (2021).

29. Jurdzinski, K. T. *et al.* Large-scale phylogenomics of aquatic bacteria reveal molecular mechanisms for adaptation to salinity. *Sci Adv* **9**, eadg2059 (2023).

30. Telesh, I. V., Schubert, H. & Skarlato, S. O. Revisiting Remane's concept: evidence for high plankton diversity and a protistan species maximum in the horohalinicum of the Baltic Sea. *Mar. Ecol. Prog. Ser.* **421**, 1–11 (2011).

31. Olli, K., Ptacnik, R., Klais, R. & Tamminen, T. Phytoplankton Species Richness along Coastal and Estuarine Salinity Continua. *Am. Nat.* **194**, E41–E51 (2019).

32. Olofsson, M., Hagan, J. G., Karlson, B. & Gamfeldt, L. Large seasonal and spatial variation in nano- and microphytoplankton diversity along a Baltic Sea-North Sea salinity gradient. *Sci. Rep.* **10**, 17666 (2020).

33. Herlemann, D. P. R., Lundin, D., Andersson, A. F., Labrenz, M. & Jürgens, K. Phylogenetic

Signals of Salinity and Season in Bacterial Community Composition Across the Salinity

Gradient of the Baltic Sea. *Front. Microbiol.* **7**, 1883 (2016).

34. Remane, A. Die Brackwasserfauna : mit besonderer Berücksichtigung der Ostsee. *Zool.*

Anz. **Suppl. 7**, (1934).

35. Lindh, M. V. *et al.* Disentangling seasonal bacterioplankton population dynamics by high-

frequency sampling. *Environ. Microbiol.* **17**, 2459–2476 (2015).

36. Gilbert, J. A. *et al.* Defining seasonal marine microbial community dynamics. *ISME J.* **6**,

298–308 (2012).

37. Ladau, J. *et al.* Global marine bacterial diversity peaks at high latitudes in winter. *ISME J.* **7**,

1669–1677 (2013).

38. Raes, E. J. *et al.* Seasonal patterns of microbial diversity across the world oceans. *Limnol.*

Oceanogr. Lett. (2024) doi:10.1002/lol2.10422.

39. Latz, M. A. C. *et al.* A comprehensive dataset on spatiotemporal variation of microbial

plankton communities in the Baltic Sea. *Sci Data* **11**, 18 (2024).

40. Olesen, S. W., Duvallet, C. & Alm, E. J. dbOTU3: A new implementation of distribution-

based OTU calling. *PLoS One* **12**, e0176335 (2017).

41. Goldford, J. E. *et al.* Emergent simplicity in microbial community assembly. *Science* **361**,

469–474 (2018).

42. Tourlousse, D. M. *et al.* Synthetic spike-in standards for high-throughput 16S rRNA gene

amplicon sequencing. *Nucleic Acids Res.* **45**, e23 (2017).

43. Lloréns-Rico, V., Vieira-Silva, S., Gonçalves, P. J., Falony, G. & Raes, J. Benchmarking

microbiome transformations favors experimental quantitative approaches to address

compositionality and sampling depth biases. *Nat. Commun.* **12**, 3562 (2021).

44. Wesslander, K., Viktorsson, L., Thor, P., Nilsson, M. & Skjevik, A.-T. *The Swedish National*

Marine Monitoring Programme 2019 : Hydrography Nutrients Phytoplankton. 59 (2020).

45. Zettler, M. L., Kremp, A. & Dutz, J. Biological assessment of the Baltic Sea 2019.

Meereswiss. Ber. , Warnemünde **115**, (2020).

46. Fridolfsson, E. *et al.* Multiyear analysis uncovers coordinated seasonality in stocks and

composition of the planktonic food web in the Baltic Sea proper. *Sci. Rep.* **13**, 11865

(2023).

47. Logares, R. *et al.* Disentangling the mechanisms shaping the surface ocean microbiota.

Microbiome **8**, 55 (2020).

48. Farjalla, V. F. *et al.* Ecological determinism increases with organism size. *Ecology* **93**,

1752–1759 (2012).

49. Wu, W. *et al.* Contrasting the relative importance of species sorting and dispersal limitation

in shaping marine bacterial versus protist communities. *ISME J.* **12**, 485–494 (2018).

50. Neuenschwander, S. M., Ghai, R., Pernthaler, J. & Salcher, M. M. Microdiversification in

genome-streamlined ubiquitous freshwater Actinobacteria. *ISME J.* **12**, 185–198 (2018).

51. Mehrshad, M. *et al.* Hidden in plain sight—highly abundant and diverse planktonic

freshwater Chloroflexi. *Microbiome* **6**, 176 (2018).

52. Pujalte, M. J., Lucena, T., Ruvira, M. A., Arahal, D. R. & Macián, M. C. The Family

Rhodobacteraceae. in *The Prokaryotes: Alphaproteobacteria and Betaproteobacteria* (eds.

Rosenberg, E., DeLong, E. F., Lory, S., Stackebrandt, E. & Thompson, F.) 439–512

(Springer Berlin Heidelberg, Berlin, Heidelberg, 2014).

53. Morris, R. M. & Spietz, R. L. The Physiology and Biogeochemistry of SUP05. *Ann. Rev.*

Mar. Sci. **14**, 261–275 (2022).

54. Spilling, K. *et al.* Shifting diatom—dinoflagellate dominance during spring bloom in the

Baltic sea and its potential effects on biogeochemical cycling. *Front. Mar. Sci.* **5**, (2018).

55. Karlson, A. M. L. *et al.* Nitrogen fixation by cyanobacteria stimulates production in Baltic

food webs. *Ambio* **44 Suppl 3**, 413–426 (2015).

56. Aguilera, A. *et al.* Ecophysiological analysis reveals distinct environmental preferences in

closely related Baltic Sea picocyanobacteria. *Environ. Microbiol.* **25**, 1674–1695 (2023).

57. Karlson, B. *et al.* Harmful algal blooms and their effects in coastal seas of Northern Europe.

Harmful Algae **102**, 101989 (2021).

58. Egge, E. S. *et al.* Seasonal diversity and dynamics of haptophytes in the Skagerrak,

Norway, explored by high-throughput sequencing. *Mol. Ecol.* **24**, 3026–3042 (2015).

59. Rost, B. & Riebesell, U. Coccolithophores and the biological pump: responses to

environmental changes. in *Coccolithophores: From Molecular Processes to Global Impact*

(eds. Thierstein, H. R. & Young, J. R.) 99–125 (Springer Berlin Heidelberg, Berlin,

Heidelberg, 2004).

60. Jordan, R. W. & Chamberlain, A. H. L. Biodiversity among haptophyte algae. *Biodiversity &*

Conservation **6**, 131–152 (1997).

61. Lücker, S. & Daims, H. The Family Nitrospinaceae. in *The Prokaryotes: Deltaproteobacteria*

and Epsilonproteobacteria (eds. Rosenberg, E., DeLong, E. F., Lory, S., Stackebrandt, E. &

Thompson, F.) 231–237 (Springer Berlin Heidelberg, Berlin, Heidelberg, 2014).

62. Lidbury, I. D. E. A. *et al.* Niche-adaptation in plant-associated Bacteroidetes favours

specialisation in organic phosphorus mineralisation. *ISME J.* **15**, 1040–1055 (2021).

63. Lidbury, I. D. E. A. *et al.* A widely distributed phosphate-insensitive phosphatase presents a

route for rapid organophosphorus remineralization in the biosphere. *Proc. Natl. Acad. Sci.*

U. S. A. **119**, (2022).

64. Teeling, H. *et al.* Substrate-controlled succession of marine bacterioplankton populations

induced by a phytoplankton bloom. *Science* **336**, 608–611 (2012).

65. Wiegand, S. *et al.* Taxonomic Re-Classification and Expansion of the Phylum Chloroflexota

Based on over 5000 Genomes and Metagenome-Assembled Genomes. *Microorganisms*

11, (2023).

66. Baldani, J. I. *et al.* The Family Rhodospirillaceae. in *The Prokaryotes: Alphaproteobacteria*

and Betaproteobacteria (eds. Rosenberg, E., DeLong, E. F., Lory, S., Stackebrandt, E. &

Thompson, F.) 533–618 (Springer Berlin Heidelberg, Berlin, Heidelberg, 2014).

67. Urios, L., Michotey, V., Intertaglia, L., Lesongeur, F. & Lebaron, P. Nisaea denitrificans gen.

nov., sp. nov. and Nisaea nitritireducens sp. nov., two novel members of the class

Alphaproteobacteria from the Mediterranean Sea. *Int. J. Syst. Evol. Microbiol.* **58**, 2336–

2341 (2008).

68. Hawley, A. K. *et al.* Diverse Marinimicrobia bacteria may mediate coupled biogeochemical

cycles along eco-thermodynamic gradients. *Nat. Commun.* **8**, 1507 (2017).

69. Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421

(2009).

70. Alldredge, A. L. & Cohen, Y. Can microscale chemical patches persist in the sea?

Microelectrode study of marine snow, fecal pellets. *Science* **235**, 689–691 (1987).

71. Klawonn, I., Bonaglia, S., Brüchert, V. & Ploug, H. Aerobic and anaerobic nitrogen

transformation processes in N2-fixing cyanobacterial aggregates. *ISME J.* **9**, 1456–1466

(2015).

72. Louca, S. The rates of global bacterial and archaeal dispersal. *ISME J.* **16**, 159–167 (2022).

73. de Vargas, C. *et al.* Ocean plankton. Eukaryotic plankton diversity in the sunlit ocean.

Science **348**, 1261605 (2015).

74. Guillou, L. *et al.* Widespread occurrence and genetic diversity of marine parasitoids

belonging to Syndiniales (Alveolata). *Environ. Microbiol.* **10**, 3349–3365 (2008).

75. Guillou, L. *et al.* The Protist Ribosomal Reference database (PR2): a catalog of unicellular

eukaryote small sub-unit rRNA sequences with curated taxonomy. *Nucleic Acids Res.* **41**,

D597–604 (2013).

76. Massana, R., del Campo, J., Sieracki, M. E., Audic, S. & Logares, R. Exploring the

uncultured microeukaryote majority in the oceans: reevaluation of ribogroups within

stramenopiles. *ISME J.* **8**, 854–866 (2014).

77. Eikrem, W. *et al.* Florenciella parvula gen. et sp. nov. (Dictyochophyceae,

Heterokontophyta), a small flagellate isolated from the English Channel. *Phycologia* **43**,

658–668 (2004).

78. Eckford-Soper, L. & Daugbjerg, N. The ichthyotoxic genus Pseudochattonella (Dictyochophyceae): Distribution, toxicity, enumeration, ecological impact, succession and life history - A review. *Harmful Algae* **58**, 51–58 (2016).

79. Schoemann, V., Becquevort, S., Stefels, J., Rousseau, V. & Lancelot, C. Phaeocystis blooms in the global ocean and their controlling mechanisms: a review. *J. Sea Res.* **53**, 43–66 (2005).

80. Kaczmarska, I., Beaton, M., Benoit, A. C. & Medlin, L. K. MOLECULAR PHYLOGENY OF SELECTED MEMBERS OF THE ORDER THALASSIOSIRALES (BACILLARIOPHYTA) AND EVOLUTION OF THE FULTOPORTULA[1]. *J. Phycol.* **42**, 121–138 (2006).

81. Alverson, A. J., Jansen, R. K. & Theriot, E. C. Bridging the Rubicon: phylogenetic analysis reveals repeated colonizations of marine and fresh waters by thalassiosiroid diatoms. *Mol. Phylogenet. Evol.* **45**, 193–210 (2007).

82. Olli, K. *et al.* Harmonizing large data sets reveals novel patterns in the Baltic Sea phytoplankton community structure. *Mar. Ecol. Prog. Ser.* **473**, 53–66 (2013).

83. Yeh, Y.-C. & Fuhrman, J. A. Contrasting diversity patterns of prokaryotes and protists over time and depth at the San-Pedro Ocean Time series. *ISME Commun* **2**, 36 (2022).

84. Hjerne, O., Hajdu, S., Larsson, U., Downing, A. S. & Winder, M. Climate driven changes in timing, composition and magnitude of the Baltic Sea phytoplankton spring bloom. *Front. Mar. Sci.* **6**, (2019).

85. Masson-Delmotte, V. *Climate Change 2021: The Physical Science Basis : Working Group I Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. (Cambridge University Press, 2021).

86. HELCOM. Baltic Sea Action Plan – 2021 update. Preprint at (2021).

87. Rohwer, R. R., Hale, R. J., Vander Zanden, M. J., Miller, T. R. & McMahon, K. D. Species invasions shift microbial phenology in a two-decade freshwater time series. *Proc. Natl.*

Acad. Sci. U. S. A. **120**, e2211796120 (2023).

88. Schloss, P. D. Rarefaction is currently the best approach to control for uneven sequencing

effort in amplicon sequence analyses. *mSphere* **9**, e0035423 (2024).

89. Giovannoni, S. J. *et al.* Genome streamlining in a cosmopolitan oceanic bacterium. *Science*

309, 1242–1245 (2005).

90. Grote, J. *et al.* Streamlining and core genome conservation among highly divergent

members of the SAR11 clade. *MBio* **3**, (2012).

91. Roda-Garcia, J. J., Haro-Moreno, J. M., Rodriguez-Valera, F., Almagro-Moreno, S. &

López-Pérez, M. Single-amplified genomes reveal most streamlined free-living marine

bacteria. *Environ. Microbiol.* **25**, 1136–1154 (2023).

92. Salcher, M. M., Schaefle, D., Kaspar, M., Neuenschwander, S. M. & Ghai, R. Evolution in

action: habitat transition from sediment to the pelagial leads to genome streamlining in

Methylophilaceae. *ISME J.* **13**, 2764–2777 (2019).

93. Bittner, M. J. *et al.* New chemical and microbial perspectives on vitamin B1 and vitamer

dynamics of a coastal system. *ISME Commun* **4**, ycad016 (2024).

94. Parks, D. H. *et al.* GTDB: an ongoing census of bacterial and archaeal diversity through a

phylogenetically consistent, rank normalized and complete genome-based taxonomy.

Nucleic Acids Res. **50**, D785–D794 (2022).

95. Fullmer, M. S., Soucy, S. M. & Gogarten, J. P. The pan-genome as a shared genomic

resource: mutual cheating, cooperation and the black queen hypothesis. *Front. Microbiol.* **6**,

728 (2015).

96. Kuosa, H. Picoplanktonic algae in the northern Baltic Sea: seasonal dynamics and

flagellate grazing. *Mar. Ecol. Prog. Ser.* **73**, 269–276 (1991).

97. Weitere, M., Dahlmann, J., Viergutz, C. & Arndt, H. Differential grazer-mediated effects of

high summer temperatures on pico- and nanoplankton communities. *Limnol. Oceanogr.* **53**,

477–486 (2008).

98. Novotny, A. *et al.* DNA metabarcoding highlights cyanobacteria as the main source of

primary production in a pelagic food web model. *Sci Adv* **9**, eadg1096 (2023).

99. Plummer, M. JAGS: A program for analysis of Bayesian graphical models using Gibbs

sampling. *3rd International Workshop on Distributed Statistical Computing (DSC 2003);*

Vienna, Austria (2003).

100.Watanabe, S. A widely applicable Bayesian information criterion. *J. Mach. Learn. Res.* **14**,

867–897 (2012).

101.Casey, J. R., Grinstein, S. & Orlowski, J. Sensors and regulators of intracellular pH. *Nat.*

Rev. Mol. Cell Biol. **11**, 50–61 (2010).

102.Schneider, B. *et al.* Biogeochemical cycles. in *Biological Oceanography of the Baltic Sea*

(eds. Snoeijs-Leijonmalm, P., Schubert, H. & Radziejewska, T.) 87–122 (Springer

Netherlands, Dordrecht, 2017).

103.Kogure, K. Bioenergetics of marine bacteria. *Curr. Opin. Biotechnol.* **9**, 278–282 (1998).

104.Zhang, H. *et al.* Repeated evolutionary transitions of flavobacteria from marine to non-

marine habitats. *Environ. Microbiol.* **21**, 648–666 (2019).

105.Benarroch, J. M. & Asally, M. The Microbiologist's Guide to Membrane Potential Dynamics.

Trends Microbiol. **28**, 304–314 (2020).

106.More, K. J., Kaur, H., Simpson, A. G. B., Spiegel, F. W. & Dacks, J. B. Contractile vacuoles:

a rapidly expanding (and occasionally diminishing?) understanding. *Eur. J. Protistol.* **94**,

126078 (2024).

107.Kawano, S., Ono, H., Takagi, T. & Bono, H. Tutorial videos of bioinformatics resources:

online distribution trial in Japan named TogoTV. *Brief. Bioinform.* **13**, 258–268 (2012).

108.Rousk, J. *et al.* Soil bacterial and fungal communities across a pH gradient in an arable soil.

ISME J. **4**, 1340–1351 (2010).

109.Heisler, J. *et al.* Eutrophication and Harmful Algal Blooms: A Scientific Consensus. *Harmful*

Algae **8**, 3–13 (2008).

110.Agneta, A. *et al.* DNA extraction protocol for DNA-metabarcoding of marine phytoplankton

using Zymobiomics DNA minprep kit (Zymo Research; D4300) v1. (2021)

doi:10.17504/protocols.io.bucjnsun.

111.Balzano, S., Abs, E. & Leterme, S. C. Protist diversity along a salinity gradient in a coastal

lagoon. *Aquat. Microb. Ecol.* **74**, 263–277 (2015).

112.Wu, L. *et al.* Phasing amplicon sequencing on Illumina Miseq for robust environmental

microbial community analysis. *BMC Microbiol.* **15**, 125 (2015).

113.Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal **17**, 10–12 (2011).

114.Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon data.

Nat. Methods **13**, 581–583 (2016).

115.Parks, D. H. *et al.* A standardized bacterial taxonomy based on genome phylogeny

substantially revises the tree of life. *Nat. Biotechnol.* **36**, 996–1004 (2018).

116.Kozlov, A. M., Zhang, J., Yilmaz, P., Glöckner, F. O. & Stamatakis, A. Phylogeny-aware

identification and correction of taxonomically mislabeled sequences. *Nucleic Acids Res.* **44**,

5022–5033 (2016).

117.Lundin, D. & Andersson, A. SBDI Sativa curated 16S GTDB database. *SciLifeLab*

https://doi. org/10. 17044/scilifelab **14869077**, v3 (2021).

118.Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data

processing and web-based tools. *Nucleic Acids Res.* **41**, D590–6 (2013).

119.Straub, D. *et al.* Interpretations of Environmental Microbial Community Studies Are Biased

by the Selected 16S rRNA (Gene) Amplicon Sequencing Pipeline. *Front. Microbiol.* **11**,

550420 (2020).

120.Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C. & Knight, R. UCHIME improves

sensitivity and speed of chimera detection. *Bioinformatics* **27**, 2194–2200 (2011).

121.R Core Team. R: A Language and Environment for Statistical Computing. Preprint at

https://www.R-project.org/ (2022).

122. Inkscape Project. *Inkscape*. (2020).

123. Oksanen, J. *et al.* vegan: Community Ecology Package. Preprint at https://CRAN.R-

project.org/package=vegan (2022).

124. Lai, J., Zou, Y., Zhang, J. & Peres-Neto, P. R. Generalizing hierarchical and variation

partitioning in multiple regression and canonical analyses using the rdacca.hp R package.

Methods Ecol. Evol. **13**, 782–788 (2022).

125. Buliung, R. N. & Remmel, T. K. Open source, spatial analysis, and activity-travel behaviour

research: capabilities of the aspace package. *J. Geogr. Syst.* **10**, 191–216 (2008).

126. Csardi, G. & Nepusz, T. The igraph software package for complex network research.

InterJournal vol. Complex Systems 1695 Preprint at https://igraph.org (2006).

127. Plummer, M. rjags: Bayesian Graphical Models using MCMC. Preprint at https://CRAN.R-

project.org/package=rjags (2023).

128. Lavrinienko, A., Jernfors, T., Koskimäki, J. J., Pirttilä, A. M. & Watts, P. C. Does

Intraspecific Variation in rDNA Copy Number Affect Analysis of Microbial Communities?

Trends Microbiol. 29, 19–27 (2021).