Opsin-based photoreception in Crinoids

Youri Nonclercq[1], Marjorie Lienard[2], Alexia Lourtie[1], Emilie Duthoo[1], Lise Vanespen[1], Igor Eeckhaut[1], Patrick Flammang[1], Jérôme Delroisse[1,3*]

[1] Laboratory of Biology of Marine Organisms and Biomimetics, Research Institute for Biosciences, University of Mons, Belgium

[2] Laboratory of Molecular Biology of Sensory Systems, GIGA, University of Liège, Belgium

[3] Laboratory of Cellular and Molecular Immunology, GIGA, University of Liège, Belgium

* **Corresponding author**: jerome.delroisse@umons.ac.be; youri.nonclercq@umons.ac.be
E-mail addresses : marjorie.lienard@uliege.be; igor.eeckhaut@umons.ac.be; patrick.flammang@umons.ac.be; alexia.lourtie@umons.ac.be; emilie.duthoo@umons.ac.be; lise.vanespen@student.umons.ac.be

Abstract

Opsin-mediated light perception has been investigated in many marine invertebrates including some clades of echinoderms such as sea stars, sea urchins and brittle stars. On the other hand, the understanding of potential light perception in crinoids, the basal lineage of the echinoderm phylum, remains largely unexplored. Only a few behavioural observations suggest that crinoids may be sensitive to light stimuli.

This study investigates the behavioural and molecular basis of opsin-based photoreception in *Antedon bifida,* a European crinoid species belonging to the Comatulid order. In this context, the behavioural response to different light wavelengths, the characterisation of opsin genes in the recent chromosome-scale genome of this species and the opsin immunolocalisation within the crinoid tissues have been investigated.

Behavioural tests pointed to a significant negative phototactic behaviour induced by a wide range of light wavelengths (463 to 630 nm) with maximum sensitivity to blue light (λmax = 463 nm). *In silico* genome analyses revealed the presence of only three rhabdomeric opsin genes located on chromosomes 4 and 6: *Abif-opsins 4.1*, *4.2* and *4.3.* All crinoid opsins are phylogenetically clustered as a sister-group of all other echinoderm rhabdomeric opsins, supporting their evolution via duplication of an ancestral gene in the crinoid lineage. The low opsin diversity contrasts with other echinoderms which are generally characterised by up to eight bilaterian opsin groups. Interestingly, *A. bifida* opsin sequences present typical amino acid residues of rhabdomeric opsins of other bilaterians, including two conserved cysteines (C110, C187), the probable ancestral E181 counterion, a NPxxY$(x)_6$F pattern, a highly conserved lysine potentially covalently bound to a chromophore, and the (D)RY motif, all supportive of photoreceptive functions. Finally, immunoreactivity to newly generated antibodies designed against sea star opsins was highlighted in several tissues associated with the ambulacral grooves of the calyx and the pinnules. Within these tissues, Abif-opsins (potentially the Abif-opsin 4.1) are expressed in the ectoneural basiepithelial nerve plexus-and the hyponeural nerve plexus. On the other hand, a different opsin type (potentially the Abif-opsin 4.2) is also expressed in the sensory papillae of tube feet. The localization of at least two opsin types in different sensory structures suggests the presence of a complex extraocular photoreception system exclusively based on rhabdomeric opsins in this crinoid species.

Keywords

Crinoidea – Opsin – Photoreception – Echinodermata – Negative phototaxis

Background

Light plays a major role in marine ecosystems. Most marine organisms have specific photosensitive organs, such as eyes, which contain opsins, the prototypic light-sensitive GPCR receptors of eumetazoans. Interestingly, echinoderm often lack specialised visual organs with the exception of optic cushions located at the tip of sea star arms [Takasu and Yoshida 1983; Johnsen 1997; Garm and Nilsson 2014; Petie et al. 2016] and putative eyespots in a few holothurian species [Berrill 1966, Yamamoto and Yoshida 1978]. However, numerous ethological observations confirm that most investigated echinoderms are light-sensitive [for review, Yoshida et al. 1984, Sumner-Rooney and Ullrich-Lüter 2023]. Most echinoderm species are then characterised by extraocular photoreception abilities, mediated by different classes of opsins expressed in different tissues, organs and structures intimately connected to the nervous system. Two types of opsins, ciliary and rhabdomeric, have been particularly investigated and identified in various adult anatomical structures, such as tube feet, integument, and spines of sea urchins [Ullrich-Lüter et al. 2011; 2013], sea stars [Delroisse et al. 2013, Ullrich-Lüter et al. 2013, Clarke et al. 2024] and brittle stars [Moore and Cobb, 1985; Cobb and Hendler, 1990; Delroisse et al. 2014; 2016; Sumner-Rooney et al. 2018; 2020; 2021]. In addition, sea stars (Asteroidea), the optic cushion is specifically expressing several opsin types [Ullrich-Lüter et al. 2011, Lowe et al. 2018]. A third type of opsin, a GO-opsin, has also been identified in two clusters of sensory cells on either side of the apical organ of pluteus sea urchin larvae [Valero-Gracia et al. 2016; Valencia et al. 2021; Cocurullo et al. 2023]. Extensive phylogenetic analyses of bilaterian opsins have shown that echinoderms are one of the phyla that have the largest diversity of ancestral opsin types with 7 opsin classes on a total of 9 predicted in the common ancestor of all bilaterians [D'Aniello et al. 2015; Ramirez et al. 2016]. Some of these ancestral opsin types, like bathyopsins and chaopsins, are currently almost exclusively restricted to echinoderms [Ramirez et al. 2016].

The crinoid group has been the least studied echinoderm class in terms of sensory perception in general, and photoreception in particular. However, given that Crinoidea currently represents the most basal clade of echinoderms, separated from the other orders since at least the Lower Ordovician period *i.e.* 480 million years ago, their phylogenetic position makes them an interesting group to better understand the evolution of photoreception not only within the whole phylum. It has long been known that some crinoid species are sensitive to light stimuli. Indeed, ethological observations have shown that several species of comatulid (an order of crinoids, most of which are non-pedunculated in the adult stage) living at shallow depths are predominantly nocturnal, extending their arms to filter feed at night [Magnus 1964, Rutman and Fishelson 1969, Meyer 1973, Vail 1987]. On the other hand, when these organisms are exposed to daylight, they tend to hide in reef crevices or under shaded underwater overhangs. Other observations point to rapid retraction of their arms when exposed to strong sunlight [Meyer 1973]. This nocturnal activity is however less and less systematically observed in individuals living at greater depths (beyond 10-20 m below the surface) where the luminosity of the environment is lower [Fishelson 1974]. Some studies have also reported a swimming escape behaviour in these feather stars, such as the European species *Antedon bifida,* when exposed to strong light stimuli [Dimelow 1958]. However, these isolated observations leave many unexplored areas, such as a deeper characterisation of the photoreceptive behaviours, or a determination of the spectral sensitivity of these animals. No photosensitive sensory structure has yet been formally demonstrated in extant crinoids. The repertoire of opsin genes also remains mostly unexplored, with only one rhabdomeric-type opsin so far identified in two transcriptome datasets from *Antedon mediterranea* and *Florometra serratissima* [D'Aniello et al. 2015].

The highly specialised lifestyle of crinoids and their phylogenetic position make them a prime
group to better understand the evolution of the large opsin diversity in echinoderms, as well as the
evolution of their extraocular photoreception.
This study sheds light on the mechanisms of crinoid opsin-based photoreception by focusing
on the phototactic behaviour of a model feather star species: *Antedon bifida*. This small common
species lives in shallow water, from a few centimetres below the surface down to a depth of 200 m
along the North Atlantic European coasts and is generally found during the day in shaded rocky
environments, sheltered from full light. Leveraging a recently available chromosome-scale genome,
we further annotate and describe the complete opsin gene repertoire in this crinoid species and assess
the precise anatomical location of several opsin-based photoreceptors. Altogether our combined
behavioural, *in silico*, morphological and expression approaches reveal novel types of candidate
photosensitive tissues, opening for molecular and functional studies in this under-studied class of
echinoderms.
104

Material and Methods
105

Collection of crinoid specimens
Specimens of *Antedon bifida* **(Fig 1 A)** were collected from two coastal harbours on the Crozon
peninsula in Brittany, France **(Fig 1B-D)**. The organisms were present in large amounts clinging to the
underside of boat mooring pontoons, just a few centimetres below the surface **(Fig 1E, F)**. Around
thirty individuals were brought back to the University of Mons (Belgium) and kept in an aquarium
containing around 400 L of seawater at 15°C (with a salinity of 34 PSU) in a closed circuit following a
12H/12H day/night cycle **(Fig 1G)**.
114

Figure 1. Distribution, habitat, and collection site of *Antedon bifida*. (A) Geographical distribution of *A. bifida* on the North Atlantic coasts of Europe. (B) Focus on the two collection sites (Morgat and Camaret-sur-Mer) located on the Crozon peninsula in Brittany (France). (C) Adult specimen of *Antedon bifida* clinging to a sabellid tubeworm. (D) Collecting site in the marina of Camaret-sur-mer. (E) The dense population of *A. bifida* hanging under the mooring pontoons. Scales: B. 15km, C. 20mm, E. 20mm.

Ethological experiments in *Antedon bifida*

Phototactic behaviour tests were carried out using a small elongated experimental aquarium (50 cm long and 7.5 cm wide) with three of its sides rendered opaque by black paper **(Fig 2)**. A monochromatic light source (LED bulb GU10 of 1 Watt) was placed against the still transparent glass end of the experimental device, creating a light gradient across the entire aquarium. The light intensity along this gradient was measured with a light metre (Digital Lux Meter AR823$^+$, Smart Sensor) and the brightness values varied from 46,000 Lux (6.7 µW/cm^2) next the light source to 3,600 Lux (0.5 µW/cm^2) at the opposite extremity. The mean value in the middle part of the aquarium was 10,000 Lux (1.5 µW/cm^2). The floor was covered with a rigid plastic mesh, allowing the crinoids to cling on and easily move in the aquarium. Three different light wavelengths were tested corresponding to blue (λ_{max}= 463 nm, half peak bandwidth HBW= 27 nm), green (λ_{max}= 512 nm, HBW= 33 nm) and red light (λ_{max}= 630 nm, HBW= 20 nm). The spectra of each LED lamp were precisely measured beforehand using a spectrometer (Ocean view, FLAM-S-UV-VIS spectrometer). One test was also carried out with polychromatic white light, and a negative control was used without a light source. Each of these five experimental conditions was carried out on 10 individuals of *A. bifida* during the same time of the day in a room completely immersed in darkness. Individuals were tested separately to avoid an aggregation phenomenon. Each crinoid specimen was first conditioned to darkness for around 5 minutes. Then, for each test, each individual was placed in the centre of the aquarium (25 cm from the illuminated end) and its displacement towards (positive displacement) or against (negative displacement) the light source was measured after 10 minutes. These displacement measurements were based on the position of the crinoid calyx centre relative to its initial position. First, statistical tests were used to confirm the homogeneity variance and the normal distribution of the displacement data with Bartlett and Quantil/Quantil tests, respectively. The mean values of the distance covered for each light wavelength tested were then compared to the mean distance travelled by individuals without illumination (negative control), using a one-way analysis of variance (ANOVA), followed by a Dunnett post hoc test with RStudio statistical software.

In silico* identification of opsin genes in *crinoids

The diversity of opsin genes in our model crinoid species, *A. bifida*, was determined through a homology-based approach. This method used several published opsin reference sequences as templates [D'Aniello et al. 2015, Ramirez et al. 2016], *i.e.* opsins from the model echinoderm species, the sea urchin *Strongylocentrotus purpuratus,* as well as opsins from additional representative bilaterians **(Suppl. Table S1)**.

Homology "tBLASTn" searches of predicted protein sequences were performed against the assembled *A. bifida* reference genome (ecAntBifi1.1) available on the NCBI database. The following tBLASTn parameters were used: *Matrix: Blosum62; gap costs: existence 11, extension 1*. We then applied a cut-off e-value < 1^{e-05} for putative opsin candidates showing a strong similarity to two highly conserved regions in functional opsins: the 7th transmembrane helix corresponding to the Schiff base binding zone of retinal (equivalent K296 lysine residue in the bovine rhodopsin), and the G-protein binding zone. Since the genome of *A. bifida* has not yet been annotated, a reciprocal BLAST of these

selected sequences was then performed on the entire NCBI database to validate their opsin
identification.
To obtain the most complete and accurate opsin gene predictions, we used several gene
prediction softwares like GENSCAN [Burge and Karlin 1997:
http://hollywood.mit.edu/GENSCAN.html], AUGUSTUS [Oliver et al. 2011: https://bioinf.uni-
greifswald.de/augustus/] and FGENESH [Solovyev et al. 2006:
http://www.softberry.com/berry.phtml?topic=index&group=programs&subgroup=gfind] using
tBLASTn on all chromosome portions of the *A. bifida* genome identified to contain opsin loci.
Then, we carried out a manual curation as follows: for each predicted opsin, only the longest
portion of the protein sequence which, using a MAFFT alignment (MAFFT version 7, Research Institute
for Microbial Diseases, Japan, https://mafft.cbrc.jp/alignment/server/index.html) [Katoh et al. 2019]
and maximising the high similarity to other echinoderm opsin reference sequences **(Suppl. Table S1)**
has been selected as the most accurate crinoid opsin sequence.
Next, a sequence similarity comparison between opsin sequences found in *Antedon bifida* and
the reference opsin repertoire of the sea urchin *S. purpuratus* (Sp-opsin 1; 2; 3.1; 3.2; 4; 5; 6; 7 and 8)
was performed using SIAS webtool (Sequence Identity And Similarity, Universidad Complutense de
Madrid) (http://imed.med.ucm.es/Tools/sias.html).
The chromosomal locations of *A. bifida* opsin genes within the genome and the number of
intron/exon regions were finally determined using internal tBLASTn search of the candidate protein
sequences identified above against the *A. bifida* genome. Gene positions within chromosomes were
visualised using IGV (Integrative Genomics Viewer) genome browser [Robinson et al 2023]. The
number and position of introns and exons were annotated by combining tBlastn, FGENESH in Softberry
[http://www.softberry.com], and the Kablammo software [Wintersinger and Wasmut 2014,
https://kablammo.wasmuthlab.org].
A similar procedure was carried out to determine all opsin sequences of two additional crinoid
species: *Anneissia japonica* (Comatulidae) and *Nesometra sesokonis* (Antedonidae like *A. bifida*),
whose reference genomes are also present in the NCBI genomic database (ASM1163010v1 and
ASM2563120v1, respectively).
191

Phylogenetic analyses of crinoid opsins

A phylogenetic analysis was carried out on opsin protein sequences previously discovered in the two
crinoid model species (*A. bifida*, *N. sesokonis*) to highlight their relationship with other opsins. For this
analysis, 111 different protein sequences from 21 species representing the 5 echinoderm classes were
used alongside opsin sequences of a few other representative groups of bilaterians. These sequences
were selected based on public metadata [Delroisse et al. 2014; D'Aniello et al. 2015; Ramirez et al.
2016; Lowe et al. 2018] and genomic databases such as NCBI to represent all nine ancestral opsin
lineages present in bilaterians (ciliary opsin, bathyopsin, Go-opsin, canonical and non-canonical
Rhabdomeric opsin, chaopsin, peropsin/RGR opsin, neuropsin and xenopsin) **(Suppl. Table S1)**. Eight
additional melatonin receptor protein sequences were added as outgroups. The sequences were then
aligned using MAFFT version 7 [Katoh et al. 2019]. The alignment was then trimmed using the TrimAl
software tool hosted on the NGPhylogeny.fr online platform
(https://ngphylogeny.fr/tools/tool/284/form). The tree was constructed using IQ-Tree
(http://iqtree.cibiv.univie.ac.at/) [Nguyen et al. 2015] based on maximum likelihood analysis with 1000
ultrafast bootstraps. The best-fit model recommended by ModelFinder [Kalyaanamoorthy et al. 2017]
to our unpartitioned dataset was LG+F+I+G4, and the number of parsimony informative sites was 726
on the 1583 amino-acid sites used in total. Tree visualisation was finally performed on the iTOL
Interactive Tree Of Life platform (https://itol.embl.de/).
210

Histology

Specimen preparation

Antedon bifida specimens intended for histological and immunohistochemical analysis were anaesthetised with a 3.5% magnesium chloride in sea water solution for 2 min before being placed 4 h in a non-acetic Bouin fixative fluid (75% of a picric acid saturated solution and 25% of formaldehyde commercial solution at 37%). After fixation, samples were decalcified for 2 days with a solution of 2% ascorbic acid and 0.3 M sodium chloride. After complete decalcification, samples were dehydrated for 24 h in graded ethanol 70° to 95° followed by a bath of butanol for 24 h at 60°C and finally embedded in paraffin (Paraplast Plus). Five μm-thick sections of arm and calyx samples were cut with a Microm HM340 E microtome and mounted on silane-coated glass slides.

Staining and imaging

Transverse sections of the calyx and longitudinal sections of the pinnules were stained with Masson's trichrome stain. Images were taken using a microscope (Orthoplan optical microscope, Leica) equipped with a high sensibility camera (DFT7000 T, Leica) using the Application Suite X software (LAS X, Leica).

Immunodetection of opsins on crinoid tissue sections

To detect opsins within the tissues of *A. bifida*, both immunofluorescence and immunohistochemical methods were carried out using two sets of purified polyclonal antibodies. These two primary antibodies, anti-Arub_ops1.1 and anti-Arub_ops4, produced in rabbits by Eurogentec, are directed respectively against ciliary opsin 1.1 (residues 159 to 174) and the c-terminal part of rhabdomeric opsin 4 (residues 447 to 461) (***Suppl. Fig. S2***) from the common European sea star species *Asterias rubens*. Histological paraffin sections of the calyx and pinnules of *A. bifida* specimens were used for opsin immunodetection.

Immunofluorescence

The dewaxed and rehydrated sample sections were rinsed with a solution of phosphate-buffered saline containing 0.5% Tween (PBS-T) and blocked 1h in PBS-T with 3% BSA (Bovine Serum Albumin). Samples were then incubated overnight at 4°C with either primary anti-opsin 1 antibodies or anti-opsin 4 antibodies at 1:200 dilution. Tissues were then rinsed in PBS-T and incubated 1h with commercial secondary antibodies (goat anti-rabbit IgG coupled to a Texas-Red fluorochrome; Alexa-fluorTM 594, Invitrogen A11012) at 1:100 dilution, revealing red fluorescence (peak at 618 nm) after exposure to an optimum excitation length of 590nm. After a final rinse with PBS-T, the slides were mounted with an aqueous mounting medium (Vectashield, Vector Laboratories) containing DAPI (4',6-diamidino-2-phenylindole) highlighting cell nuclei with blue fluorescence (450-490 nm). Opsin labelling was visualised by an Olympus FV1000D inverted confocal laser scanning microscope under three channels of fluorescence by laser excitation (peak excitation/emission): DAPI (358/461 nm), FITC (495/521 nm) and Texas Red (590/618 nm).

Immunohistochemistry

Sections used for immunohistochemistry were treated slightly differently from those used for immunofluorescence. After dewaxing and rehydration, slides were heated in a citrate buffer (0.01 M, pH 6.2) in the microwave (2X5 min, 900 W) to unmask specific antigenic epitopes present in the tissue. Tissues were then immersed in 0.06% hydrogen peroxide to quench endogenous peroxidases. The following steps of rinsing, blocking and incubation treatments with primary antibodies were similar to those described above for immunofluorescence. After a second rinse, the tissues were incubated 1 h with a horse anti-rabbit/peroxidase secondary antibody complex (ImmPressTMReagent kit, Vector

259   Laboratories). For opsin revelation, sections were incubated 1 to 3 minutes with a PBS-buffered
260   solution (pH 7.2) containing 4 mg/ml DAB (4-dimethylaminoazobenzene) and 0.02% H_2O_2. The sections
261   were counterstained with hemalun/Luxol Blue to enhance the contrast with the brown DAB-
262   immunostaining.
263
Controls
Various negative and positive controls were conducted. Negative controls included omitting
the primary antibody or using a pre-immune rabbit serum diluted at 1:2000 in place of the primary
antibody. In each case, all control sections were not labelled **(*Suppl. Fig. S3*)**. Sections from tissues of
Asterias rubens, including the optic cushions, tube feet and nervous system were used as positive
controls for the detection of opsins 1 and 4. In all cases, the immunostaining gave the expected results
as described in previous publications [Ullrich-Lüter et al. 2011, Clarke et al. 2024].
271
Scanning electron microscopy
Two crinoid specimens were prepared (calyx and arms separated) for precise anatomical observation
under a scanning electron microscope. After fixation with non-acetic Bouin's fluid, the specimens were
first progressively dehydrated with ethanol before being dried in a chamber using the CO_2 critical point
drying technique. The samples were then coated with a thin film of gold and palladium using a Jeol
JFC-1100E sputter coater (JEOL Company, Tokyo, Japan). Observations of the fine structure of the
pinnules and calyx of the feather star were carried out with a Jeol JSM-7200F scanning electron
microscope.
280

<u>Results</u>

***Antedon bifida* is negatively phototactic with a maximum response under blue light.**
Behavioural phototactic tests showed that adult crinoids of the species *A. bifida* were sensitive to white
light and moved significantly away from the light source (negatively oriented displacement) compared
to the control test without light ($p = 0.0099^{**}$) **(Fig. 2B)**. This negative phototactic behaviour was also
significant with the red-light ($p = 0.031^*$) and reached its maximum amplitude with blue light ($p = 0.0062^{**}$). This result points to a high sensitivity of crinoids to the blue part of the visible spectrum. In
0.0062^{**}). This result points to a high sensitivity of crinoids to the blue part of the visible spectrum. In
contrast, the results for the green-light tests showed no significant difference in displacement values
compared to the dark control values (green: $p = 0.167$).
Displacement values were also normalised by individual calyx size (distance travelled/calyx
diameter) to avoid any bias in displacement speed due to variation in individual size. However, we
found that there was no correlation between the size of the individuals and the travelled distance
during the test. After normalisation, significant differences were only observed for the blue and white
lights compared to the dark control.
Absolute displacement values were also considered (non-oriented displacements) to
determine the reactivity of crinoid individuals to light without considering the direction taken **(Fig. 2C)**.
As illustrated in Figure 2C, the distances travelled during the test, independently of the direction, did
not vary significantly between individuals exposed to the different wavelengths and the controls
remaining in the dark.
Thus, although light stimuli did not induce significant changes in the mobility of *A. bifida* (non-
directional displacement), white, red (630 nm) and blue (463 nm) light stimuli induced a significant
directional displacement of crinoids away from the light source (negative phototactic behaviour).
304

Figure 2. Phototactic behaviour in *Antedon bifida*: (A) Schematic representation of the experimental aquarium with a light gradient. (B) Graph showing the distance travelled by individuals within the experimental set-up over 10 minutes of time following different light stimuli (Black = no light; Blue = λmax of 463 nm; Green = λmax of 512 nm; Red = λmax of 630 nm and White = polychromatic white light). Negative displacement indicated a displacement away from the light source. One-factor ANOVA and Dunnett's post hoc statistical tests showed that crinoids exhibited a significant oriented displacement towards the darker region of the light gradient for white, blue and red light compared with the no-light control. (C) Graph showing the distances covered by the individuals in absolute values, showing the rate of movements of crinoids depending on the type of light stimulus. Dunnett's one-factor ANOVA and post hoc tests did not provide significant results between no-light controls versus different wavelengths tested.

***Antedon bifida* has a reduced opsin repertoire composed of only three rhabdomeric opsins**

The homology-based opsin search in the *A. bifida* genome enabled the discovery of three opsin genes: the *Abif-opsin 4.1* gene present on chromosome 4 [NCBI reference: OY727096.1], the *Abif-opsin 4.2* and *Abif-opsin 4.3* genes present on chromosome 6 [NCBI reference: OY727098.1] **(Fig. 3B)**. The three predicted *A. bifida* opsins contain, respectively 417, 356 and 408 amino acids (46.42, 41.17 and 45.63 kDa). A clue as to the identification of these three sequences was obtained through reciprocal blast searches and comparison of the sequence similarity with the set of reference opsins present in the sea

urchin *S. purpuratus* and three other non-echinoderm opsin types **(Fig. 3A)**. The results showed that the three *A. bifida* opsin sequences have higher similarity with the (non-canonical) rhabdomeric opsins of echinoderms (opsin 4). The same analyses carried out on the genome of another closely related crinoid species, *Nesometra sesokonis*, also revealed the presence of three similar opsin sequences, which also appeared to belong to the rhabdomeric opsin group. On the other hand, research carried out on the genome of the species *Anneissia japonica* (Comatulidae) revealed no opsin genes. It therefore seems that only one bilateral opsin lineage is still present in *A. bifida* and presumably in all Crinoids of the family Antedonidae.

All *Abif-opsin* genes share a conserved gene structure composed of one or two short exon sequences (80-270 bp) followed by a third, much longer, terminal exon region (800-960 bp) **(Fig. 3C)**. In addition, our genomic analysis of *A. bifida* opsin genes revealed two major structural differences compared to rhabdomeric opsin genes in other echinoderm classes, which lead to an overall shorter gene length due to a lower number of intronic regions (between 1 or 2 introns for *A. bifida* versus 3 for sea stars and sea urchins) and reduced intron sizes (between 540 and 1,800 bp maximum for crinoid opsins and from minimum 2,300 bp for the sea star *A. rubens* to maximum 11,400 bp for the sea urchin *S. purpuratus*). Genomic sequence alignments showed homology at the level of the first and last long exon for the three crinoid opsins, as well as for those of sea star and sea urchins. Finally, in the sea star *A. rubens* and the sea urchin *S. purpuratus* opsin 4 orthologs, the regions homologous to the third and fourth exon are fused into a single last exon in the three crinoid opsin sequences. Sequence homology analysis also shows that the first exon of *A. bifida* opsin 4.2 is composed of the assembly of the two first exons homologous to those present in the other two crinoid opsins. A short region of 57 base pairs is also found between these exons, and whose exonic nature recognised by gene prediction software is still uncertain.

Multiple alignment of the predicted opsins found in *A. bifida* and *Nesometra sesokonis* with other rhabdomeric opsins reveals the presence of highly conserved amino acid residues characteristic of light-sensitive opsins **(Fig. 3D)**. These opsin characteristics are mainly amino-acids present in the third and seventh transmembrane helix (Tm) as well as in the second extracellular loop (EC), which allow interactions with the conjugated chromophore molecule (retinal) and the G-protein. The most representative opsin residue is a lysine residue in the 7^{th} Tm, corresponding to K296 position in the rat rhodopsin, which performs the Schiff-based covalent binding with the retinal [Terakita 2005]. This lysine is conserved in all aligned predicted opsins of crinoids. For the stabilisation of this protonated Schiff-based binding, a counterion is necessary and this function is typically assumed by a glutamate residue [Terakita et al. 2004; Varma et al. 2019]. The probable ancestral position of this counterion is the equivalent of the glutamate E181 in the rat rhodopsin present in the second extracellular loop. This E_{181}, as the only counterion function, is typical of bistable opsins such as rhabdomeric opsins [Terakita et al. 2012; 2014]. The monostable opsins like vertebrate ciliary opsins possess another glutamate counterion in Tm III (corresponding to the E113 in the rat rhodopsin) closer to the retinal binding. This glutamate E_{113} is replaced by a tyrosine residue in rhabdomeric opsins. The three crinoid opsins present the same ancestral counterion glutamate E181, with the tyrosine residue also replacing E113 like in other echinoderm rhabdomeric opsins. These results point to the probable bistable character of crinoid opsins. Another typical amino acid pattern of all GPCR proteins like opsins is the $NPxxY(x)_6F$ pattern (corresponding to the sequence NPII(Y)306 in the rat rhodopsin) which could contribute, amongst other functions, to the binding between the opsin and the G-protein [Fritze et al. 2003]. This motif is also present in all the sequences including the Melatonin GPCR receptor. Another tripeptide also seems to participate in the association between the opsin and the G-protein, this is the (D)RY motif (corresponding to the ER(Y)136 in the rat rhodopsin). We can also mention the two opsin-typical cysteine residues (corresponding to C110 in the Tm III and the C187 in the EC II of rat rhodopsin) which allow the stabilisation of protein conformation owing to the formation of a disulfide bridge. The

three crinoid opsins possess all these key residues typical of functional opsins. We can also note the presence, in two of the three crinoid opsins (4.1 and 4.3), of the HxK pattern inside the motif NPxxY(x)6F, which is rather typical of rhabdomeric opsins and is homologous of the pattern NxQ found in ciliary opsins (corresponding to the sequence Nx(Q)312 in the rat rhodopsin) [Upton et al. 2021]. This typical-rhabdomeric pattern is only partly present in the crinoid-opsins 4.2 in which the histidine residue is replaced by a tyrosine. It is interesting to note that this is the only case in the echinoderm rhabdomeric opsins identified thus far.

Figure 3. *In silico* analysis of crinoid opsins: (A) Similarity comparison (%) between the three crinoid opsin protein sequences predicted from the *Antedon bifida* genome with all echinoderm opsin groups present in the model species of sea urchin (*Strongylocentrotus purpuratus*). This comparative study was also extended to two vertebrate canonic opsins present in *Rattus norvegicus* and one non-echinoderm opsin group in a brachiopod *Lingula anatina*. The three bold boxes highlight the best similarity percentage between the crinoid opsins and the non-canonical rhabdomeric opsin of *S. purpuratus*. (B) The chromosomal position of the three opsin genes in the *Antedon bifida* genome represented by a leading arrow indicating the reading direction of the gene. (C) The representation of intronic and exonic regions (and indication of homologous regions) in the three opsin genes of the studied species (*A. bifida*) compared to the sea star and sea urchin rhabdomeric opsin genes of the

model species (*A. rubens* and *S. purpuratus*). (D) Alignments in the third (Tm III) and seventh (Tm VII)
transmembrane helix domains as well as in the second extracellular loop (EC II) protein region of the
six opsin sequences evidenced in both crinoid species (Abif = *Antedon bifida*; Nses = *Nesometra*
sesokonis) versus 14 rhabdomeric opsins of other bilaterian metazoans (Arub = *Asterias rubens*; Apla
= *Acanthaster plancii*; Spur = *Strongylocentrotus purpuratus*; Lvar = *Lytechinus variegatus*; Afil =
Amphiura filiformis; Blan = *Branchiostoma lanceolatum*; Rnor = *Rattus norvegicus*; Ajap = *Apostichopus*
japonicus; Hgla = *Holothuria glaberrima*; Dmel = *Drosophila melanogaster*). The first and the last
supplementary sequences in these alignments are respectively the rat rhodopsin (a reference ciliary
opsin which gives the amino acid numeration) and the melatonin receptor of sea urchin *S. purpuratus*
(non-opsin GPCR protein) used as comparison models. Different amino acid patterns, characteristic of
typical opsins, are highlighted in colour in these alignments and are all present in the three crinoid
opsins.
404
Phylogenetic position of crinoid opsins
A first exploded global opsin phylogeny in different species of Echinoderms and other typical bilaterian
phyla allowed it to recover all nine ancestral bilaterian opsin lineages **(Fig. 4A, full phylogenetic tree**
presented in Supplementary Figure S1). In this way, we were able to confirm in *silico* analyses that all
three crinoid opsins belong to the rhabdomeric opsin group (opsins 4). A focus on rhabdomeric opsins
4 then allowed us to highlight their relationships between investigated crinoid species and other
echinoderms **(Fig. 4B)**. Crinoid and other echinoderm rhabdomeric opsins gathered into a well-
separated clade **(Fig. 4B)** that further branched apart from classical rhabdomeric opsins like vertebrate
melanopsins and *Drosophila* r-opsins, supporting that all rhabdomeric opsins in echinoderms
diversified from a unique common ancestor. This is consistent with the fact that all echinoderm opsins
4 have so far been identified as non-canonical rhabdomeric opsins and in this phylogeny, appear to be
the sister group to the canonical opsins of the chordates and protostomes. Other "non-conventional"
rhabdomeric opsins of brachiopods and polychaetes are not directly grouped with echinoderm opsins
and are represented as a sister group of the clade containing canonical opsins and non-canonical opsin
of echinoderms. It is also important to notice that the three opsin sequences discovered from the
genomes of both crinoid species (*Antedon bifida* and *Nesometra sesokonis*) are exclusively grouped
with two other published partial opsin sequences from the transcriptomes of two other feather star
species. This crinoid opsin clade is placed in the basal position of the other echinoderm rhabdomeric
opsins, in agreement with the phylogeny of extant echinoderms. The other echinoderm opsins 4 are
also grouped according to the different echinoderm classes, in perfect agreement with the well-
established phylogenetic classification of the phylum. The relationships between crinoid opsins point
to the existence of three rhabdomeric opsin lineages in the Antedonidae family, the most basal
position being occupied by crinoid opsin 4.2 orthologs.
428

A

Tree scale: 1

Melatonin receptors

Neuropsins

GO-opsins

Rhabdomeric opsin

Abif-opsin 4.2
Abif-opsin 4.3
Abif-opsin 4.1

RGR-opsins

Peropsins

Ciliary opsins

Chaopsins

Bathyopsins

Xenopsin

B

Melanopsin receptors
98

Ciliary opsins
97
25
Chaopsins
100
87
Xenopsins
94
92
Bathyopsins
20
37
GO-opsins
66
92
Neuropsins
98
89
RGR-opsins
64
99
Peropsins
90

Non-canonical rhabdomeric opsins
Canonical rhabdomeric opsins

Lana-opsin 4.2
Lana-opsin 4.1
100
Pdum-opsin 4.5
81
Pdum-opsin 4.4
99.5
Blan-opsin 4
Xlae-opsin 4
98
Rnor-opsin 4
100
Pdum-opsin 4.3
99
Pdum-opsin 4.2
94
Pdum-opsin 4.1
64
Soff-opsin 4
68.5
Dmel-opsin 4.6
49
Dmel-opsin 4.2
100
Dmel-opsin 4.1
99
Dmel-opsin 4.7
99.4
Dmel-opsin 4.5
96
Dmel-opsin 4.4
99
Dmel-opsin 4.3
99

Nses-opsin 4.2
96.5
Abif-opsin 4.2
Nses-opsin 4.1
95
Amed-opsin 4
100
100
Abif-opsin 4.1
94
Nses-opsin 4.3
Fser-opsin 4
97
Abif-opsin 4.3
24
Lvar-opsin 4
Crinoidea
100
Spur-opsin 4
Echinoidea
91
Hgla-opsin 4
Ajap-opsin 4.2
100
Holothuroidea
Ajap-opsin 4.1
99.5
Apla-opsin 4
70
Asteroidea
Arub-opsin 4
100
Afil-opsin 4.4
79
Afil-opsin 4.6
100
Afil-opsin 4.5
78
Afil-opsin 4.1
Ophiuroidea
92.5
Afil-opsin 4.3
83.5
Aaga-opsin 4
53
Afil-opsin 4.2
99
100

85
99

Tree scale: 1

Figure 4. Phylogenetic analysis of Crinoid opsins. (A) Global phylogenetic tree of the nine representative ancestral opsin lineages in bilaterian metazoans supported by maximum likelihood method. Bootstrap values are indicated at the root of each opsin cluster. The melatonin receptor cluster is used to root the phylogenetic tree and the branch length scale gives an insight into the relative rate of amino acid substitution per site. The three opsin sequences evidenced in the crinoid *Antedon bifida* (Abif) are indicated in bold into the rhabdomeric opsin cluster and associated with blue crinoid icons. (B) Focus on the rhabdomeric opsin cluster with a distinction between Canonical (in light blue) and non-canonical (in dark blue) R-opsins. The branch lengths and bootstrap values are indicated in this case respectively above and below the branches. There are in total 40 rhabdomeric opsin sequences present in this phylogeny belonging to 18 representative bilaterian species (Echinoderms: Abif = *Antedon bifida*; Amed = *Antedon mediterranea*; Nses = *Nesometra sesokonis*; Fser = *Florometra serratissima*; Arub = *Asterias rubens*; Apla = *Acanthaster plancii*; Spur = *Strongylocentrotus purpuratus*; Lvar = *Lytechinus variegatus*; Afil = *Amphiura filiformis*; Aaga = *Astrotomma agassizii* ; Ajap = *Apostichopus japonicus*; Hgla = *Holothuria glaberima*; Chordates: Blan = *Branchiostoma lanceolatum*; Rnor = *Rattus norvegicus*; Xlae = *Xenopus laevis*; Lophotrochozoans: Lana = *Lingula anatina*; Pdum = *Platynereis dumerilii*; Soff = *Sepia officinalis*; Arthropods: Dmel = *Drosophila melanogaster*). The three opsin sequences of the crinoid species (*A. bifida*) (Abif-opsin4.1, 4.2 and 4.3) are highlighted in bold at the base of the non-canonical rhabdomeric opsin cluster of Echinoderms. Accession numbers of all sequences are presented in the Supplementary Table S1. The Full circular phylogeny is presented in the Supplementary Figure S1.

Antedon bifida opsins are expressed in the tube feet and nervous systems

Opsin detection in the calyx area

A. bifida has a typical morphology of feather star crinoids (*Comatulida*) with a central calyx with mobile hook-like cirri on the aboral side for locomotion **(Fig. 5A)** and ten filtering arms connected to the oral side of the calyx. Each arm possesses numerous bilateral ramifications called pinnules **(Fig. 5B)**. A deep aboral nervous system, typical of crinoids, is localised in the deep part of the calyx and constituted by a complex bowl-shaped cluster whose branches form the brachial nerves **(Fig. 5F)**. Ramifications of brachial nerves extend into the aboral side of the pinnules. The mouth is in the middle of the calyx oral side (tegmen), right next to the anal tube that is slightly off-centre **(Fig. 5C)**. Five ambulacral grooves, that carry filtered food to the mouth, start from the mouth and dichotomise before extending to the arms. Each calyx groove is alternately bordered on both sides by spherical bodies called saccules (**Fig. 5D-F)** and by a unique series of tube feet **(Fig. 5E-F)**. Histological cross-sections of the ambulacral groove region point to the presence of two neuronal networks, on one hand, the ectoneural basiepithelial nerve plexus confined at the base of the groove epidermis and, on the other hand, a diffuse hyponeural nerve plexus present slightly down in the connective tissue **(Fig. 5F)**.

Figure 5. Morphological structure of the Calyx: (A) An adult specimen of *Antedon bifida*. (B) Oral view of the central calyx surrounded by the ten branched arms. (C) Scanning electron microscopy picture of the whole oral surface (tegmen) of the calyx with a focus on the ambulacral groove indicated by a box. (D) Lateral histological cross section on the global calyx with a focus on the boxed ambulacral groove region. (E) Micrography of the calycinal groove portion bordered by well-visible tube feet. (F) Transversal histological section (Masson's trichrome stain) of the ambulacral groove on the calyx. AG: Ambulacral Groove ; ANM: Aboral Nerve Mass; Ar: Arm; ARC : Ambulacral Radial Canal ; AT: Anal Tube; ATC: Ambulacral Tentacular Canal; BN: Brachial Nerve; Ca: Calyx; Ci : Cirri; CT : Connective Tissue; EBP: Ectoneural Basiepithelial Plexus ; HN : Hyponeural Nerve; Hy: Hydropores; In: Intestine; M: Mouth; My: Myocyte; OP: Oral Pinnule; Os: Ossicule; Pi: Pinnules; Pp: Papilla; S: Saccule; So: Somatocoele; TF: Tube Foot. Scales: A. 20 mm, B. 10 mm, C. 400 µm, D. 300 µm, E. 20 µm, F. 50 µm.

Both ecto- and hyponeuronal plexus exhibited a strong immunoreactivity after exposure to antibodies raised against opsin 4 of the sea star *Asterias rubens*, supporting the presence of opsins in

these nervous structures **(Fig. 6 A-E)**. The high sequence similarity between the antigen sequence of
A. rubens opsin 4 and the sequence of *A. bifida* opsin 4.1 **(*Suppl. Fig. S2*)** suggests that the
immunoreactivity observed both by immunofluorescence **(Fig. 6 A-C)** and by immunohistochemistry
(Fig. 6 B-E) correspond to the expression of the Abif-opsin 4.1 in these nervous structures.

Other structures showing the presence of opsins corresponded to tube feet, bordering the
ambulacral grooves **(Fig. 5 E-F)**. The epidermis of each tube foot presents several papillae with ciliary
sensory cells. The opsin immunoreactivity was observed more precisely in the epidermal cells at the
base and tip of these sensory papillae **(Fig. 6 F-G)**. These immunodetections were performed with
antibodies raised against *A. rubens* opsin 1.1. The similarity of antigenic sequence between *A. rubens*
opsin 1.1. and *A. bifida* opsin 4.2 **(*Suppl. Fig S2*)** suggests that the immunoreactivity observed in the
papillae of tube feet corresponds to the expression of the Abif-opsin 4.2. However, sequence
divergence for the target binding region in the available peptide antibodies does not rule out that opsin
4.3 may co-localise with opsin 4.2 or be expressed in additional sensory cells.

Figure 6. Immunostainings of two rhabdomeric opsins in the Calix ambulacral groove tissues of
Antedon bifida: (A) Localisation of opsin 4.1 by red immunofluorescence into the ectoneural
basiepithelial nerve plexus just under the ambulacral groove epithelium. (AI; AII) Focus on the two
basiepithelial spots of opsin labelling in red. (B) Opsin 4.1 is highlighted at the base of arm with a
localisation on the one hand in the ectoneural basiepithelial plexus (white head arrow) and the other
hand in the hyponeural plexus present deeper into the connective tissue (black arrow). (C) Localisation

of the same opsin 4.1 by immunocytochemistry in brown into the hyponeural nerve plexus inside the
connective tissue of the calyx ambulacral groove. (D) Focus on the hyponeural opsin showing the
localisation near a saccule. (E) Focus on the brown immunostaining of opsin 4.1 in the hyponeural
plexus of the base of the arm. (F) Labelling in red by immunofluorescence of opsin 4.2 at the top of
tube feet (in the sensory papillae) bordered the ambulacral groove of the calyx. (G)
Immunocytochemistry showing the opsin 4.2 expression at the top of tube feet (arrow). Scales: A.
50µm, A'-A''. 10µm, B. 25µm, C. 50µm, D. 20µm, E. 20µm, F. 50µm, G. 50µm.
511
Opsin detection in the pinnules
The pinnules of crinoids are lateral extensions of the arms **(Fig. 7 A)**. A pinnular ambulacral
groove occupies the middle oral side of each pinnule and is bordered by two rows of tube feet allowing
food capture and transport **(Fig. 7 B)**. The pinnular tube feet are grouped in triads arranged
alternatively on either side of the groove **(Fig. 7 C)**. These triads consist of one long primary tube foot
and a medium-sized secondary tube foot dedicated to trap food particles, and a short tertiary tube
foot whose function is to form the food bolus that is transported to the mouth [Lahaye and Jangoux
1985]. Similarly to the calyx tube feet, pinnular tube feet also have several epidermal sensory-secretory
outgrowths called papillae. Papillae are known to feature several cell types: secretory cells, peripheral
support cells with a long apical cilium, and central neurosensory cells with a shorter cilium [Flammang
and Jangoux 1991]. Scanning electron microscopy observations highlighted ciliary structures at the
extremity of the papillae **(Fig. 7 H)**. As for the case in calyx papillae, these pinnular papillae showed an
immunoreactivity with the anti-Arub_opsin 1.1 antibodies **(Fig. 7 E-G)**. These opsin labelling
corresponds more probably to the expression of Abif-opsin 4.2 as mentioned above (***Suppl. Fig. S2***).
The labelling was strong in the basal part of papillae but was also present occasionally in the apical
pole of papillae cells **(Fig. 7 E-G)**.
The three parts of the pinnular nervous system are extensions of those found in the arms and
calyx. The pinnular ambulacral grooves feature an ectoneural basiepithelial plexus and a deep aboral
nerve plexus in the centre of the pinnules. On the other hand, the hyponeural nerve plexus is
constituted by two lateral nerve cords that run longitudinally on either side of the pinnules and
innervate the tube feet and saccules **(Fig. 7C)**. As was the case in the calyx section, aboral, ectoneural
and hyponeural networks present in pinnules expressed opsin proteins **(Fig. 7D)**. As mentioned before
this immunostaining in nervous structures corresponds, more probably, to the presence of Abif-opsin
4.1 (*Suppl. Fig. S2*).
536

537
538

Figure 7. Anatomical structures of pinnules tissues and rhabdomeric opsin immunostainings in the pineal ambulacral groove in *Antedon bifida*: (A) Crinoid arm with a focus on a pinnule indicated by a red box. (B) Scanning electron microscopy picture at the extremity of the pinnule. (C) Longitudinal histological section (Masson's trichrome stain) in the lateral pinnule edge. (D) Localisation of opsin 4.1 in the ectoneural basiepithelial plexus (dark arrow) and the deep aboral nerve of the pinnule (white arrow). These histological structures become visible thanks to the further longitudinal section in the centre of the ambulacral groove. (E) Labelling of opsin 4.2 by immunofluorescence in red into the tube feet epidermis and the extremity of sensory papillae. (F) Immunostaining (brown precipitate) of opsin 4.2 in the lateral sensory papillae (dark arrow) of one secondary tube foot. (G) Similar opsin 4.2 immunostainings in brown at the base of sensory papillae (dark arrow) of one primary tube foot. (H) Micrography of the sensory papillae structures of a tube foot. AG: Ambulacral groove; Ar: Arm; ATC: Ambulacral Tentacular Canal; CC: Coelomic Canal; CCC: Covering ciliary cell; NCC: Neuronal Ciliary Cell; Pi: Pinnules; TF: Tube foot; TF'/TF''/TF''': primary/secondary/tertiary Tube Foot; HN: Hyponeural nerve; Pp: Papilla; PRC: Pinnule Radial Canal. Scales: A. 5mm, B. 200µm, C. 50µm, D. 20µm, E. 20µm, F. 20µm, G. 20µm, H. 3µm.

Discussion

Crinoid light behaviour

Multiple reef-dwelling Comatulid species feed mainly at night and hide in rocky crevices or fold their arms and pinnules during the day which suggests photophobic behaviours [Magnus 1962; Meyer 1973; Rutman and Fishelson 1969]. Our results evidenced a negative phototactic behaviour in *Antedon bifida* which is consistent with previous *in situ* observations of hiding behaviour made in other shallow water feather stars. One study also mentioned a positive phototactic behaviour for a small feather star species *Dorometra nana* that was attracted to a light source [Clark 1909]. This isolated observation could indicate a species-specific response, but the photic response could also be affected by different variables such as the light intensity. Some observations suggested that the nocturnal feeding activity and the diurnal shelter-seeking behaviour of many crinoids are not linked to the circadian cycle but would be well induced by direct ambient light detection. Indeed, some specimens belonging to the same species (*Heterometra savignii)* but living at greater depths, in an environment of constant low luminosity, tended rather to lose their nocturnal activity but kept their high photophobic reaction to an intense light stimulus [Fishelson 1974]. It is therefore not surprising that the studied population of *A. bifida* living just below the surface, clinging to the shade under pontoons of the marina, flees the polychromatic white light as it was previously reported in the literature based on observations in natural conditions [Dimelow 1958].

The ethological tests carried out on *Antedon bifida* showed that the negative phototactic behaviour reached its maximum amplitude for blue light pointing to the greater sensitivity of this species for the short light wavelength range (λmax = 463 nm). The greater sensitivity to blue light has also been demonstrated several times by phototactic behavioural tests in other echinoderm groups, such as the sea urchin *Strongylocentrotus purpuratus* [Ullrich-Lüter et al 2011], the brittle star *Amphiura filiformis* [Delroisse et al 2014] and, more recently, the sea cucumber *Holothuria leucospilota* [Wang et al 2023]. The blue part of the light spectrum is the wavelength range that penetrates the farthest into the water column and is, therefore, the light component, together with green light, mostly present in open water environments (Davis 1991). Electrophysiological studies in ocellary structures of some reef-dwelling sea star species attested a very narrow sensitivity peak for the blue wavelength (450 nm) suggesting the presence of a blue-sensitive opsin in ocellary spots of these sea stars [Garm and Nilsson 2014, Petie et al. 2016]. By contrast, *A. bifida* is sensitive to a much broader light spectrum.

Experiments performed in the present study demonstrated a significant negative phototactic
behaviour for red light (λmax= 630nm). This sensitivity to red light may be correlated with the fact that
this species lives at very shallow depths (less than 10m) and therefore is exposed to longer wavelengths
like red penetrating only a few metres below the sea surface, especially in the Atlantic and North Sea
where the light intensity declines quickly with depth (Davis 1991, Capuzzo et al. 2013, 2015). This large
variability in the range of perceived light wavelengths, together with specific expression patterns,
suggests that directional photoreception in this species of Comatulida may be mediated by different
opsins. Our study confirms the presence of three different opsin genes evidenced in the genome of *A.*
bifida, bearing all conserved characteristics of functional opsins. This suggests that *A. bifida* relies on
rhabdomeric opsins for extraocular light perception. Another clue pointing to the presence of a multi-
opsin photoreception in *A. bifida* is the expression of two different rhabdomeric opsins localised in
specific tissues such as the subepidermal ectoneural nerve plexus and the tips of tube feet. However,
further *in vitro* investigations of the bioactivity of the three different opsins are required to functionally
characterise the crinoid opsins.

A trend for *A. bifida* to move away from the light source was also observed for green light
(λmax = 512 nm), but this was not significant. This lack of significance is probably due either to a low
detection of green light or to a lack of phototactic behavioural response for this wavelength in this
Comatulid species. Green light is the wavelength range most present in the shallow coastal waters
where *A. bifida* populations are found. This ecologically relevant green wavelength could therefore
lead to distinct behavioural responses in contrast to other light wavelengths. A similar absence of
reactivity to green light has already been described in tadpoles of the amphibian species *Xenopus laevis*
living in green eutrophic water [Roberts et al. 2000]. Nevertheless, future behavioural tests with green
light wavelength could be carried out on a larger number of individuals to ensure that there has been
no statistical bias due to insufficient sampling.
610

Opsin evolution in feather stars
In-silico analyses performed on the genome of the crinoid species *A. bifida* and *N. sesokonis* have
revealed the presence of only three opsin genes. In the chromosome-scale genome of *A. bifida*, the
opsin genes are distributed on chromosomes 4 and 6. All three opsin genes were identified as
belonging to the same group of non-canonical echinoderm rhabdomeric opsins. This low opsin gene
diversity was quite unexpected given that the other four echinoderm classes possess a much more
varied opsin repertoire with between 9 and 15 different opsin genes currently known [D'Aniello et al.
2015]. For instance, sea urchins and brittle stars, still exhibit seven of the nine ancestral opsin lineages
inherited from the common ancestor of bilaterians: ciliary opsins, bathyopsins, Go-opsins, non-
canonical rhabdomeric opsins, chaopsins, peropsins/RGR-opsins and neuropsins [Ramirez et al. 2016].

Crinoids are considered as the basal lineage of echinoderms which generally justified their
importance in comparative evolutionary studies. Crinoidea has a long evolutionary history of at least
485 million years [Rouse et al. 2013; Oji and Twitchett 2015] and all crinoids living today originated
from the single branch of Articulata, the only crinoid lineage to have survived the great extinction of
the late Paleozoic [Baumiller et al. 2011]. Current feather stars therefore represent a relatively derived
group compared to the ancestral forms of the Paleozoic. The Crinoidea available genomes (i.e.,
Anneissia japonica [ASM1163010v1 in NCBI, 0.5896 Gb], *Nesometra sesokonis* [ASM2563120v1 in NCBI,
0.5971 Gb] and *Antedon bifida* [ecAntBifi1.1 in NCBI, 0.3196 Gb] the only chromosome-scale genome)
are comparable to sea stars (e.g., *Asterias rubens* (0.4176 Gb) and *Marthasterias glacialis* (0.52Gb)
[Parey et al. 2024]) but appear considerably smaller compared to other echinoderm genomes (e.g.,
Strongylocentrotus purpuratus (0.9218 Gb), *Holothuria leucospilota* (1.4 Gb), and *Amphiura filiformis*
(1.57 Gb) [Parey et al. 2024]). In *A. bifida*, specifically, a reduced number of chromosomes is observed
(11 chromosomes) compared to other chromosome-scale echinoderm genomes (i.e., 18 for

Paracentrotus lividus [Parey et al. 2024], 22 for *A. rubens*, 23 for *H. leucospilota* and 20 for *A. filiformis*
[Parey et al. 2024]). Our study further reveals a reduced opsin repertoire accompanying these genomic
characteristics. It is possible that, during their evolutionary history, these modern crinoids (many live
at relatively great depths in low luminosity) may have lost many of the ancestral opsin groups still
present in other echinoderms. The smaller crinoid genome could also explain the lower number of
intronic regions in their opsin genes, making crinoid opsin genes shorter than those found in other
echinoderm classes.
However, the presence of a single bilaterian opsin class may not be generalisable to all crinoid
groups but may just be specific to the Comatulida family Antedonidae. Indeed, the two species (*A.*
bifida and *N. sesokonis*), whose genomes have been investigated in the present study, belong to the
family Antedonidae. Research carried out on the only other available crinoid genome belonging to the
Comatulidae family species (*Anneissia japonica*) revealed no opsin gene. A more complete sampling
across Crinoidea will be crucial to have a more complete representation of opsin diversity within the
class.
The alignment comparisons with other metazoan rhabdomeric opsins have demonstrated that
all three crinoid protein sequences possess the set of amino acid residues characteristic of functional
opsins allowing interaction with both retinal and the G-protein. This suggests that these three
rhabdomeric opsins potentially play a direct role in light detection in *A. bifida*. Our behavioural tests
support the view that *A. bifida* individuals possess photosensitivity covering a wide part of the light
spectrum, potentially related to the activity of different opsins exhibiting distinct spectral absorptions.
The involvement of rhabdomeric opsins as the main photoreception molecular actor in the light
detection has been mentioned several times in other Echinoderms [Ullrich-Lüter et al 2011; Delroisse
et al 2014] notably for sea stars which possess a largely predominant expression of a rhabdomeric
opsin in their compound ocelli (optic cushion) at the end of arms [Lowe et al. 2018].
The phylogenetic analysis of rhabdomeric opsins in bilaterians places the three crinoid opsins
together at the base of the non-canonical rhabdomeric opsin clade of Echinoderms. Within the latter,
the relationship of opsins belonging to the different classes follows perfectly the classical phylogenetic
classification of Echinoderms, as highlighted in other studies [Delroisse et al. 2014; D'Aniello et al.
2015; Lowe et al. 2018]. The fact that the three crinoid opsins are clustered together suggests that
they come from a duplication of an ancestral rhabdomeric opsin gene in this group. The same more
extended gene polyplication was observed in the brittle star species *Amphiura filiformis*, which has 6
different rhabdomeric opsin genes [Delroisse et al 2014].
666

Potential photosensory structures in crinoids

Opsin immunostaining revealed the presence of at least two opsins in various tissue structures
associated with the ambulacral grooves of the calyx and pinnules. One opsin was associated with the
different nerve plexi whereas another opsin was expressed at the tips of the tube feet. These
histological structures containing opsins are ideally located on the surface of the body and face away
from the substratum, allowing a maximum detection of the sunlight rays.
The first opsin detected in the nervous plexus by antibodies raised against opsin 4 from the
sea star *Asterias rubens* very likely corresponds to opsin 4.1 of *A. bifida*. Indeed, as evidenced by the
alignments of the protein region recognised by the antibody, *A. bifida* opsin 4.1 shows the highest
percentage (46.66%) of similarity with the epitope sequence of *Asterias rubens* rhabdomeric opsin (see
the alignment in ***Suppl. Fig. S2***). The presence of rhabdomeric opsins directly linked to the nervous
system is relatively well described in other echinoderms, notably in the radial nerves and sensory
nerves innervating the tube feet in sea urchins and brittle stars [Ullrich-Lüter et al 2011; Delroisse et
al 2014]. It is well known that the ectoneural basiepithelial nerve plexus of echinoderms is involved in
the transmission of sensory information [Weber and Grosmann 1977; García-Arrarás et al 2001;

Mashanov et al 2023; Adameyko 2023], suggesting that this nervous plexus may also be involved in
transmitting photo-sensorial information. By contrast, a photosensory activity within the hyponeural
plexus is more intriguing. Indeed, this nervous plexus, located deeper in the connective tissue, is
generally considered to perform only motor functions in echinoderms [García-Arrarás et al 2001].
Nevertheless, some paleontological studies have demonstrated that the hyponeural nerves in
Paleozoic crinoids Camerata innervated directly some specific sensory organs in the calyx [Haugh
1978], demonstrating a possible dual sensory-motor role in crinoids, including a possible role for opsins
in these deeper nerve fibres.

The detection of opsins in hyponeural nerves could shed new light on the still-unknown
function of the saccules present below the surface of crinoid pinnules. These spherical bodies are
highly refractive [Mallefet et al 2023] and are quite abundant in some species of feather stars, such as
our study model *Antedon bifida*. They are regularly arranged under the epidermis on either side of the
ambulacral grooves in the calyx and pinnules. The exact function of these saccules is still unknown, but
numerous hypotheses about their physiological role have already been proposed, such as mucus
secretion [Carpenter 1884; Clark 1921; Mironov and Pawson 2010], as an energy reserve [Hyman 1955]
or playing a role in light production in bioluminescent species [Mallefet et al 2023]. Because of their
optical properties, a hypothesis has been formulated suggesting their possible role as refractive lenses
enabling light rays to converge on photoreceptors [Holland 1967]. The expression of rhabdomeric
opsins in the hyponeural plexus located under the saccules, and innervating them, seems to support
this hypothesis.

Earlier studies have also highlighted the presence of numerous pigments in the subepithelial
connective tissue of *A. bifida* ambulacral grooves, as well as in other feather stars [Carpenter 1876,
Dimelow 1958]. This pigmentation could potentially play a shading role at the level of ecto- and
hyponeural nerve *plexi* expressing opsins. This shading system would be necessary to allow a
directional photoreception, as demonstrated by phototactic tests in *A. bifida*.

The second opsin was specifically localised in the epidermis of the tube feet bordering the
ambulacral grooves of the calyx and pinnules. This opsin was immunodetected using a different rabbit
antibody raised against *Asterias rubens* opsin 1.1. Although this antibody recognises a ciliary opsin in
the sea star, the epitope sequence shows a relatively high similarity with the opsin 4.2 from *A. bifida*
(62.5%, **Suppl. Fig. S2**). Furthermore, a search for sequence homology has been performed against the
entire *A. bifida* genome. Our results indicate that Abif-opsin 4.2 has a significant similarity (62.5%)
with the sequence of the immunogenic peptide used to produce antibodies against sea star opsin 1.1.
In the future, the results will need to be confirmed using immunoblots analyses.

Observing rhabdomeric opsin expression in tube feet of *A. bifida* is noteworthy, given that r-
opsin localisation in tube feet has been previously demonstrated in other echinoderm classes such as
sea urchins and brittle stars [Ullrich-Lüter et al 2011, Delroisse et al 2014], yet not in crinoids thus far.
In contrast to other echinoderm, crinoids have their tube feet facing away from the substratum, which
exposes them more effectively to light in the water column. Additionally, ciliary neurosensory cells
have been found in various epidermal extensions such as "papillae" and "hillocks" of crinoid tube feet
[Flammang and Jangoux 1991]. The opsin immunoreactivity is precisely located both at the base and
occasionally at the tip of these sensory papillae, suggesting that these sensorial structures may be
involved in light perception. This potential tube foot photoreception is corroborated by *pax 6* gene
expression observed in the tube foot tips of the feather star species *Anneissia japonica* [Omori et al
2020], suggesting the probable presence of photoreceptors at the tips of tube feet in crinoids.
Expression of *pax 6* is generally recognised for its pivotal role in the embryonic development of light-
sensitive structures across bilaterian metazoans [Callaerts et al. 199f7; Gehring and Ikeo 1999; Arendt;
2003; Zuber et al. 2003; Martínez-Morales et al. 2004; Stierwald et al. 2004; Kozmik 2005].
Interestingly, *pax 6* has also been shown to be expressed in adult photoreceptor organs in various

echinoderms, including co-expression of *pax 6* and opsin genes in the sea urchin species *S. purpuratus* [Ullrich-Lüter et al 2011]. However, it will be necessary to, e.g., express and characterise the function of these opsins by heterologous expression and develop gene knock-out workflows in this new model system to fully demonstrate the role and molecular mechanisms underlying non-canonical rhabdomeric opsins photoreception.

Funding

YN and ED are supported by FRIA grants from the "Fonds de la Recherche Scientifique" of Belgium (F.R.S.-FNRS) and a PDR grant (T.0169.20). AL was supported by a FRIA grant from the F.R.S.-FNRS. JD is a postdoctoral researcher funded by a PDR project from the F.R.S.-FNRS (T.0071.23). PF is research director from the F.R.S.-FNRS.

Acknowledgements

This study contributes to the BioSciences Research Institute from the University of Mons and the "Centre Interuniversitaire de Biologie Marine" (CIBIM). The authors are grateful to the staff of the Marine Station of Concarneau for help and support.

Conflict of interest

The authors declare no conflicts of interest.

Author contributions

YN, IE, JD collected samples. YN, ML, AL, LV and JD performed the experiments. YN and JD analysed and interpreted the results. YN and JD prepared the first draft of the manuscript. JD and PF supervised the study. All authors read and approved the final manuscript.

Data availability

Data generated/analysed during the current study are available from the corresponding author upon reasonable request.

References

Adameyko I. (2023). Evolutionary origin of the neural tube in basal deuterostomes. Current Biology 33: R319–R331 (2023). https://doi.org/10.1016/j.cub.2023.03.045

Arendt D. (2003). Evolution of eyes and photoreceptor cell types. Int J Dev Biol. 47(7-8): 563-571 (2003). PMID: 14756332.

Baumiller TK, Salamon MA, Gorzelak P, Mooi R, Messing CG, Gahn FJ. (2010). Post-Paleozoic crinoid radiation in response to benthic predation preceded the Mesozoic marine revolution. Proc Natl Acad Sci U S A. Mar 30;107(13):5893-6 (2010). https://doi.org/10.1073/pnas.0914199107

Berrill M. (1966). The ethology of the Synaptid holothurian, *Opheodesoma spectabilis.* Canadian Journal of zoology, 44: 457-482 (1966). https://doi.org/10.1139/z66-046

Burge, C. and Karlin, S. (1997) Prediction of complete gene structures in human genomic DNA. J. Mol. Biol. 268, 78-94 (1997). https://doi.org/10.1006/jmbi.1997.0951

Callaerts P., Halder G., Gehring W.J. (1997). Pax-6 in development and evolution. Annu Rev Neuroscience 220: 483–532 (1997). https://doi.org/10.1146/annurev.neuro.20.1.483

Capuzzo E, Painting SJ, Forster RM, Greenwood N, Stephens DT, Mikkelsen O (2013) Variability in the sub-surface light climate at ecohydrodynamically distinct sites in the North Sea. Biogeochemistry, 113, 85–103 (2013). https://doi.org/10.1007/s10533-012-9772-6

Capuzzo, E., Stephens, D., Silva, T., Barry, J. and Forster, R.M. (2015), Decrease in water clarity of the
southern and central North Sea during the 20th century. Glob Change Biol, 21: 2206-2214 (2015).
https://doi.org/10.1111/gcb.12854

Carpenter P.H. and von Graff L. (1884). On three new species of Metacrinus. Transactions of the
Linnean Society of London, 2nd Series, Zoology, 2(14) : 435-446 (1884).
https://doi.org/10.1111/j.1096-3642.1885.tb00296.x

Carpenter, P.H. (1885) Report upon the Crinoidea collected during the voyage of H.M.S. Challenger
during the years 1873-1876. Part I - General morphology of the stalked crinoids. Report of the Scientific
Results of the Exploring Voyage H.M.S. Challenger, London, Zoology, 11 (32): 1-442 (1885).

Clark A. H. (1909). New genera and higher groups of unstalked crinoids. Proceedings of the Biological
Society of Washington 22: 173-177 (1909).

Clark A. H. (1921). A monograph of the existing crinoids, vol. 1 . The comatulids, pt. 2: [General].
Bulletin of the United States National Museum. 1-795 (1921).
https://doi.org/10.5479/si.03629236.82.2

Clarke D.N., Formery L. and Lowe C.J. (2024). See-Star: a versatile hydrogel-based protocol for clearing
large, opaque and calcified marine invertebrates. BMC EvoDevo 15:8 (2024).
https://doi.org/10.1186/s13227-024-00228-0

Cobb, J. L. S. and Hendler, G. (1990). Neurophysiological characterisation of the photoreceptor system
in a brittlestar, *Ophiocoma wendtii* (Echinodermata: Ophiuroidea). Comp. Biochem. Physiol. A 97, 329-
333 (1990). https://doi.org/10.1016/0300-9629(90)90619-4

D'Aniello S., Delroisse J., Valero-Gracia A., Lowe E.K., Byrne M., Cannon J.T., Halanych K.M., Elphick
M.R., Mallefet J., Kaul-Strehlow S., Lowe C.J., Flammang P., Ullrich-Lüter E., Wanninger A., Arnone M.I.
(2015). Opsin evolution in the Ambulacraria. Mar. Genomics, 24 (2) :177-183 (2015).
https://doi.org/10.1016/j.margen.2015.10.001 Epub 2015 Oct 23. PMID: 26472700

Davis R. A.(1991). In Oceanography: An Introduction to the Marine Environment, 2nd ed. Dubuque, IA:
Wm. C. Brown Publishers, 1991.

Delroisse J., Lanterbecq D., Eeckhaut I., Mallefet J. and Flammang P. (2013). Opsin detection in the sea
urchin Paracentrotus lividus and the sea star Asterias rubens. Cahiers de Biologie Marine, 54: 721-727
(2013). http://hdl.handle.net/2078.1/134949

Delroisse, J., Mallefet, J., Flammang, P. and Ullrich-Lüter, E. (2014). High opsin diversity in a non-visual
infaunal brittle star. BMC Genomics, 15: 1035 (2014). https://doi.org/10.1186/1471-2164-15-1035

Delroisse, J., Mallefet, J. and Flammang, P. (2016). De novo adult transcriptomes of two European
brittle stars: spotlight on opsin-based photoreception. PLoS ONE, 11(4): e0152988 (2016).
https://doi.org/10.1371/journal.pone.0152988

Dimelow E. (1958). Pigments Present in Arms and Pinnules of the Crinoid, Antedon bifida (Pennant).
Nature 182, 812 (1958). https://doi.org/10.1038/182812a0

Flammang P. and Jangoux M. (1991). The sensory-secretory structures of the podia of the comatulid
crinoid *Antedon bifida* (Echinodermata). In Echinoderm Research 1991, L. Scalera-Liaci & C.Canicatti
(eds) 1992 Balkema, Rotterdam. ISBN 90 5410 049 4.

Fishelson L. (1974). Ecology of northern Red Sea crinoids and their epi- and endozoic fauna. Marine
Biology 26:183–192 (1974). https://doi.org/10.1007/BF00388888

Fritze O., Filipek S., Kuksa V., Ernst O. (2003). Role of the conserved NPxxY(x)5,6F motif in the rhodopsin
ground state and during activation. PNAS, 100(5): 2290-2295 (2003).
https://doi.org/10.1073/pnas.0435715100

García-Arrarás J. E., Rojas-Soto M., Jiménez L. B., Díaz-Miranda L. (2001). The Enteric Nervous System
of Echinoderms: Unexpected Complexity Revealed by Neurochemical Analysis. J Exp Biol 204 (5): 865–
873 (2001). https://doi.org/10.1242/jeb.204.5.865

Garm A, Nilsson DE. (2014). Visual navigation in starfish: first evidence for the use of vision and eyes in
starfish. Proc. Biol. Sci. 281(1777):20133011 (2014). https://doi.org/10.1098/rspb.2013.3011 PMID:
24403344; PMCID: PMC3896028.

Gehring W.J. and Ikeo K. (1999). Pax 6: mastering eye morphogenesis and eye evolution. Trends Genet.
15(9): 371-377 (1999). https://doi.org/10.1016/s0168-9525(99)01776-x PMID: 10461206.

Gehring W. J. (2005). New Perspectives on Eye Development and the Evolution of Eyes and
Photoreceptors, Journal of Heredity, 96(3): 171–184 (2005). https://doi.org/10.1093/jhered/esi027

Haugh B. N. (1978). Biodynamic and phyletic paradigms for sensory organs in camerate crinoids.
Lethaia 11 (2) :145–173 (1978). https://doi.org/10.1111/j.1502-3931.1978.tb01300.x

Holland N. D. (1967). Some observations on the saccules of *Antedon mediterranea* (Echinodermata,
crinoidea). Pubblicazioni Stazione Zool. Napoli 35, 257–262.

Hyman L.H. (1955) The invertebrates, vol IV, Echinodermata. McGraw-Hill (eds), New York.

Johnsen S. (1997) Identification and Localization of a Possible Rhodopsin in the Echinoderms *Asterias*
forbesi (Asteroidea) and *Ophioderma brevispinum* (Ophiuroidea). Biological Bulletin. 193: 97-105
(1997). https://doi.org/10.2307/1542739

Katoh k., Rozewicki J., Yamada K.D. (2019). MAFFT online service: multiple sequence alignment,
interactive sequence choice and visualization. Briefings in Bioinformatics, 20(4): 1160–1166 (2019).
https://doi.org/10.1093/bib/bbx108

Kozmik Z. (2005). Pax genes in eye development and evolution. Curr. Opin. Genet. Dev. 15: 430–438
(2005). https://doi.org/10.1016/j.gde.2005.05.001

Lahaye M.C. and Jangoux M. (1985). Functional morphology of the podia and ambulacral grooves of
the comatulid crinoid A ntedon bifida (Echinodermata). Marine Biology 86, 307-318 (1985). Doi:
10.1007/bf00397517

Lam-Tung Nguyen, Heiko A. Schmidt, Arndt von Haeseler, and Bui Quang Minh (2015) IQ-TREE: A fast
and effective stochastic algorithm for estimating maximum likelihood phylogenies. Mol Biol Evol,
32:268-274 (2015). https://doi.org/10.1093/molbev/msu300

Lowe EK, Garm A.L., Ullrich-Lüter E., Cuomo C., Arnone M.I. (2018). The crowns have eyes: multiple
opsins found in the eyes of the crown-of-thorns starfish *Acanthaster planci*. BMC Evol. Biol., 18(1):168
PMID: 30419810 (2018). https://doi.org/10.1186/s12862-018-1276-0

Li Y.Y., Su F.J., Hsieh Y.J., Huang T.C. and Wang Y.S. (2021). Embryo Development and Behavior in Sea
Urchin (*Tripneustes gratilla*) Under Different Light Emitting Diodes Condition. Front. Mar. Sci. 8: 684330
(2021). https://doi.org/10.3389/fmars.2021.684330

Mallefet J., Martinez-Soares P., Eléaume M.,O'Hara T. and Duchatelet L. (2023). New insights on crinoid
(Echinodermata; Crinoidea) bioluminescence.Front. Mar. Sci. 10:1136138 (2023).
https://doi.org/10.3389/fmars.2023.1136138

Magnus, D. B. E. (1964) Gezeitenströmung und Nahrungsfiltration bei Ophiuren und Crinoiden.
Helgoländer wiss. Meeresunters. 10: 105-117.

Martínez-Morales J.R., Rodrigo I., and Bovolenta P. (2004). Eye development: a view from the retina
pigmented epithelium. BioEssays : news and reviews in molecular, cellular and developmental biology.
26(7):766-777 (2004). https://doi.org/10.1002/bies.20064

Mashanov V., Ademiluyi S., Jacob Machado D., Reid R. and Janies D. (2023) Echinoderm radial glia in
adult cell renewal, indeterminate growth,and regeneration.Front. Neural Circuits 17:1258370 (2023).
https://doi.org/10.3389/fncir.2023.1258370

Meyer D. L. (1973). Feeding Behavior and Ecology of Shallow-Water Unstalked Crinoids
(Echinodermata) in the Caribbean Sea. Marine Biology 22: 105-129 (1973).
https://doi.org/10.1007/BF00391776

Mironov A.N. and Pawson D.L. (2010). A new genus and species of Western Atlantic sea lily in the family Septocrinidae (Echinodermata: Crinoidea: Bourgueticrinida). Zootaxa 2449: 49–68 (2010). https://doi.org/10.11646/zootaxa.2449.1.2

Moore A. and Cobb J.L.S (1985) Neurophysiological studies on photic responses in Ophiura ophiura. Comparative Biochemistry and Physiology Part A: Physiology. 80 (1): 11-16 (1985). https://doi.org/10.1016/0300-9629(85)90669-3

Oji T. and Twitchett R.J. (2015). The oldest post-Palaeozoic Crinoid and Permian-Triassic origins of the Articulata (Echinodermata). Zoolog Sci. Apr;32(2):211-5 (2015). https://doi.org/10.2108/zs140240

Oliver Keller, Martin Kollmar, Mario Stanke, Stephan Waack (2011) A novel hybrid gene prediction method employing protein multiple sequence alignments. Bioinformatics, 27(6): 757–763 (2011). https://doi.org/10.1093/bioinformatics/btr010

Omori A., Shibata T.F. and Akasaka, K. (2020). Gene expression analysis of three homeobox genes throughout early and late development of a feather star Anneissia japonica. Dev Genes Evol 230, 305–314 (2020). https://doi.org/10.1007/s00427-020-00665-6

Parey E., Ortega-Martinez O., Delroisse J., Piovani L., Czarkwiani A., Dylus D., Arya S., Dupont S., Thorndyke M., Larsson T., Johannesson K., Buckley K., Martinez P., Oliveri P., and Marlétaz F. (2024). The brittle star genome illuminates the genetic basis of animal appendage regeneration. nature ecology & evolution (2024). https://doi.org/10.1038/s41559-024-02456-y

Petie, R., Hall, M. R., Hyldahl, M., & Garm, A. (2016). Visual orientation by the crown-of-thorns starfish (*Acanthaster planci*). Coral Reefs, 35(4): 1139–1150 (2016). https://doi.org/10.1007/s00338-016-1478-0

Ramirez M.D., Pairett A.N., Pankey M.S., Serb J.M., Speiser D.I., Swafford A.J., Oakley T.H. (2016). The Last Common Ancestor of Most Bilaterian Animals Possessed at Least Nine Opsins. Genome Biol. Evol. 8(12): 3640–3652 (2016). https://doi.org/10.1093/gbe/evw248

Robinson J., Thorvaldsdóttir H., Turner D., Mesirov J. (2023). igv.js: an embeddable JavaScript implementation of the Integrative Genomics Viewer (IGV). Bioinformatics 39(1), btac830 (2023). https://doi.org/10.1093/bioinformatics/btac830

Roberts, A., Hill, N. A. and Hicks, R. (2000). Simple mechanisms organise orientation of escape swimming in embryos and hatchling tadpoles of Xenopus laevis. J. Exp. Biol. 203, 1869–1885 (2000). https://doi.org/10.1242/jeb.203.12.1869

Rouse GW, Jermiin LS, Wilson NG, Eeckhaut I, Lanterbecq D, Oji T, Young CM, Browning T, Cisternas P, Helgen LE, Stuckey M, Messing CG. (2013). Fixed, free, and fixed: the fickle phylogeny of extant Crinoidea (Echinodermata) and their Permian-Triassic origin. Mol Phylogenet Evol. 66(1):161-81 (2013). https://doi.org/10.1016/j.ympev.2012.09.018

Rutman J. and Fishelson L. (1969). Food composition and feeding behavior of shallow-water crinoids at Eilat (Red Sea). Mar. Biol. 3: 46—57 (1969). https://doi.org/10.1007/BF00355592

Solovyev V, Kosarev P, Seledsov I, Vorobyev D. (2006) Automatic annotation of eukaryotic genes, pseudogenes and promoters. Genome Biol 7 Suppl 1: S10. 1-12 (2006). https://doi.org/10.1186/gb-2006-7-s1-s10

Stierwald M., Yanze N., Bamert R. P., Kammermeier L., and Schmid V. (2004). The Sine oculis/Six class family of homeobox genes in jellyfish with and without eyes: Development andregeneration. Developmental Biology 274: 70-81 (2004). https://doi.org/10.1016/j.ydbio.2004.06.018

Subha Kalyaanamoorthy, Bui Quang Minh, Thomas KF Wong, Arndt von Haeseler, and Lars S Jermiin (2017). ModelFinder: Fast model selection for accurate phylogenetic estimates. Nature Methods, 14:587–589 (2017). https://doi.org/10.1038/nmeth.4285

Sumner-Rooney L., Rahman I.A., Sigwart J.D., Ullrich-Lüter E. (2018). Whole body photoreceptor networks are independent of 'lenses' in brittle stars. Proceedings of the Royal Society B: Biological Sciences, 285: 20172590 (2018). https://doi.org/10.1098/rspb.2017.2590

Sumner-Rooney L., Kirwan J.D., Lowe E., Ullrich-Lüter E. (2020). Extraocular Vision in a Brittle Star Is
Mediated by Chromatophore Movement in Response to Ambient Light. Current Biology, 30: 319–327
(2020). https://doi.org/10.1016/j.cub.2019.11.042
Sumner-Rooney L., Kirwan J.D., Lüter C. and Ullrich-Lüter E. (2021) Run and hide: visual performance
in a brittle star. J. Exp. Biol. 224 (11): jeb236653 (2021). https://doi.org/10.1242/jeb.236653
Sumner-Rooney L. & Ullrich-Lüter J. (2023). Extraocular vision in echinoderms. In *Distributed Vision:*
From Simple Sensors to Sophisticated Combination Eyes (pp. 49-85). Cham: Springer International
Publishing. https://doi.org/10.1007/978-3-031-23216-9_3
Takasu N. and Yoshida M. (1983). Photic effects on photosensory microvilli in the seastar *Asterias*
amurensis. Zoomorphology 103: 135-148 (1983). https://doi.org/10.1007/BF00310473
Terakita, A., Koyanagi, M., Tsukamoto, H. *et al.* Counterion displacement in the molecular evolution of
the rhodopsin family. *Nat Struct Mol Biol* 11, 284–289 (2004). https://doi.org/10.1038/nsmb731
Terakita A. The opsins. *Genome Biol* 6, 213 (2005). https://doi.org/10.1186/gb-2005-6-3-213
Terakita A., Kawano-Yamashita E., Koyanagi M. (2012). Evolution and diversity of opsins. WIREs Membr
Transp Signal, 1:104–111 (2012). https://doi.org/10.1002/wmts.6
Terakita A. and Nagata T. (2014). Functional Properties of Opsins and their Contribution to Light-
Sensing Physiology. Zoological Science, 31(10): 653-659 (2014). https://doi.org/10.2108/zs140094
Ullrich-Lüter E.M., Dupont S., Arboleda E., Hausen H., Arnone M.I. (2011). Unique system of
photoreceptors   in   sea   urchin   tube   feet.   PNAS,   108   (20):   8367–8372   (2011).
https://doi.org/10.1073/pnas.1018495108
Ullrich-Lüter E.M., D'Aniello S., Arnone M.I. (2013). C-opsin Expressing Photoreceptors in Echinoderms.
Integrative and Comparative Biology, 53 (1): 27–38 (2013). https://doi.org/10.1093/icb/ict050
Upton B., Díaz N., Gordon S., Van Gelder R., Buhr E., Lang R. (2021). Evolutionary Constraint on Visual
and Nonvisual Mammalian Opsins. JOURNAL OF BIOLOGICAL RHYTHMS, 36(2): 109–126 (2021).
https://doi.org/10.1177/0748730421999870
Vail L. (1987). Diel patterns of emergence of crinoids (Echinodermata) from within a reef at Lizard
Island,   Great   Barrier   Reef,   Australia.   Marine   Biology   93:551–560   (1987).
https://doi.org/10.1007/BF00392793
Varma N., Mutt E., Mühle J., Panneels V., Terakita A., Deupi X., Noglye P., Schertlera G., Lesca E. (2019).
Crystal structure of jumping spider rhodopsin-1 as a light sensitive GPCR. PNAS, 116(29): 14547-14556
(2019). https://doi.org/10.1073/pnas.1902192116
Wang M.H., Hsieh Y.J., Chang H.H., Wang Y. S. (2024). Effects of different visible light spectrums on
phototaxis and bottom preference behavior of sea cucumber. Aquaculture 578, 740112 (2024).
https://doi.org/10.1016/j.aquaculture.2023.740112
Weber W. and Grosmann M. (1977). Ultrastructure of the basiepithelial nerve plexus of the sea urchin,
Centrostephanus   longispinus.   Cell   Tissue   Res.,   175:   551–562   (1977).
https://doi.org/10.1007/BF00222418
Wintersinger J.A., Wasmut J.D. (2014). Kablammo: an interactive, web-based blast results visualizer.
Bioinformatics, 31 (2014), pp. 1305-1306 (2014). https://doi.org/10.1093/bioinformatics/btu808
Yamamoto M., Yoshida M. (1978). Fine Structure of the Ocelli of a Synaptid Holothurian, Opheodesoma
spectabilis, and the Effects of Light and Darkness. Zoomorphologie, 90: 1 – 17 (1978).
https://doi.org/10.1007/BF00993740
Yoshida, M., Takasu, N., & Tamotsu, S. (1984). Photoreception in echinoderms. In Photoreception and
vision in invertebrates (pp. 743-771). Boston, MA: Springer US. https://doi.org/10.1007/978-1-4613-
2743-1_22
Zuber M.E., Gestri G., Viczian A.S., Barsacchi G. and Harris W.A. (2003). Specification of the vertebrate
eye by a network of eye field transcription factors. Development. 130(21): 5155-5167 (2003).
https://doi.org/10.1242/dev.00723 Epub 2003 Aug 27. PMID: 12944429.

Supplementary data:

Supplementary Table S1. List of different opsin sequences from bilaterian species used for the *in silico* and phylogenetic analysis.

Supplementary Figure S1. Phylogenetic analysis of Crinoid opsins. (A) Global phylogenetic tree of the nine representative ancestral opsin lineages in bilaterian metazoans supported by maximum likelihood method. Bootstrap values are indicated at the root of each opsin cluster and represented further in the branch by a circle size proportional to the bootstrap value. Methodological details are presented in the legend of Figure 4.

Supplementary Figure S2. Multiple alignment, identity and similarity values between the two immunogenic peptides of the sea star (*Asterias rubens*) opsin 1.1 and opsin 4 recognised by antibodies and the crinoid (*Antedon bifida*) opsin sequences. Other opsin sequences of the sea star *Asterias rubens* and the sea urchin *Strongylocentrotus purpuratus* were added as model comparisons.

Supplementary Figure S3. Control tests for immunolabeling opsins in the ambulacral groove of the *A. bifida* calyx**:** (A) immunofluorescence detection of opsin 4.1 in the basiepithelial nerve plexus (opsin labelling in red and DAPI labelling of cell nuclei in blue). As saccules are highly refractive structures, they show strong red autofluorescence. (B) immunofluorescence detection of opsin 4.2 (in red) at the tip of tube feet. (C) Immunohistochemistry detection of opsin 4.2 (in brown) at the tip of tube feet. (D) Negative control in immunofluorescence without antibodies. (E) Negative control in immunofluorescence with rabbit preimmune serum. (F) Negative control in immunohistochemistry without antibodies.