Territoriality and resource availability modulate the effect of conspecific encounters on the

foraging behaviours of a mammalian predator

Authors: Jeanne Clermont[1,2,5,*], Frédéric Dulude-de Broin[3,5], Marie-Pier Poulin[4] & Dominique

Berteaux[2,5]

Authors affiliations:

1. Université de Sherbrooke, Sherbrooke (Québec), Canada.

2. Canada Research Chair on Northern Biodiversity, Université du Québec à Rimouski,

Rimouski (Québec), Canada.

3. Université Laval, Québec (Québec), Canada.

4. University of Wyoming, Laramie (Wyoming), USA.

5. Center for Northern Studies and Quebec Center for Biodiversity Science.

*Corresponding author information: email: jeanne_clermb@hotmail.com, full address:

Université de Sherbrooke, 2500 boulevard de l'Université, Sherbrooke (Québec), J1K 2R1,

Canada.

Abstract

The probability of encountering conspecifics shapes multiple dimensions of animal behaviour. For example, territorial individuals increase vigilance, scent marking and alarm calling when approaching home range boundaries. Whether territorial predators modify their foraging behaviours with respect to the probability of encountering neighbouring territory owners is poorly understood. However, this could strongly influence the landscape of predation risk and therefore modulate predator-prey interactions. We studied the movements and behaviours of 23 resident Arctic foxes occupying neighbouring home ranges during two years of contrasting resource availability (abundant resources in 2019, scarce resources in 2022) on Bylot Island, Nunavut, Canada. First, based on simultaneous GPS tracking of individuals, we established which individuals used a territory (an exclusive area) by estimating the spatial distribution of the probability of encountering a neighbour within their home range. Second, using GPS and accelerometry data, we evaluated if the probability of encountering a neighbour influenced the spatial distribution of foraging behaviours, and whether this relationship differed between territorial and non-territorial individuals. When resources were abundant, only breeding individuals excluded other foxes from a part of their home range and were thus territorial. When resources were rare, none of the foxes reproduced and all but one were territorial. Non-territorial individuals in 2019 were less likely to cache prey in areas with a high probability of encounter, possibly reducing the risk of cache pilfering. We found no effect of the probability of encounter on the behaviours of the non-territorial individual in 2022. The probability to cache prey of territorial individuals however did not depend on the probability of encountering a neighbour, suggesting they have no benefit of modulating the distribution of their caches according to potential encounters in a partly exclusive area. Our results thus suggest that Arctic foxes use different tactics to secure resources based on their degree of territoriality and the availability of resources. We highlight how the presence of resident, but non-territorial predators, whose home ranges overlap those of their territorial neighbours, may influence the distribution of predation risk by creating zones where predator density is high, potentially influencing predator-prey interactions.

Keywords: foraging, home range, intraspecific interactions, movement, territoriality, *Vulpes lagopus*

Introduction

The ability of animals to manage encounters with other individuals is key to all social and interspecific interactions, thus directly influencing reproduction, survival, and ultimately fitness (Silk et al., 2003; Cresswell, 2008; Wey et al., 2008). Consequently, many animal behaviours have evolved to modulate the probability of encountering other individuals. One well known example is how predators and prey modify their habitat use to either find or avoid each other (Sih, 1984; Smith et al., 2019a). Similarly, individuals may avoid conspecifics to decrease resource competition. They for example secure limited resources by becoming territorial, given the benefits of territoriality outweigh the costs (Maher & Lott, 2000; Ord, 2021). A territory represents the area of a home range that is successfully defended by one or several individuals, resulting in the territory owners having exclusive access to the defended area (Brown & Orians, 1970; Powell, 2000; Hinsch & Komdeur, 2017). Territorial individuals will often signal space ownership through visual, auditory or olfactory signals and aggress intruders when encountered (Sillero-Zubiri & Macdonald, 1998; Giuggioli et al., 2011; Tórrez-Herrera et al., 2020).

The intensity of territoriality is often assessed by measuring the amount of overlapping area between neighbouring home ranges (Persson et al., 2010), or the frequency with which individuals adopt territorial behaviours such as scent marking (Fawcett et al., 2013). Furthermore, the degree to which individuals of a given species are territorial can vary among and within populations (Macdonald et al., 1999; McLoughlin et al., 2000; Eide et al., 2004). Ecological variables such as the abundance, predictability and distribution of food resources influence the costs and benefits of territoriality, and therefore partly explain the degree of territoriality (Maher & Lott, 2000; Sells & Mitchell, 2020). Notably, a cost-benefit model developed by Maher and Lott (2000) suggests that individuals should be territorial when food abundance and predictability are intermediate, but non-territorial when food abundance and predictability are below or above some thresholds, which was supported by McLoughlin et al. (2000) who studied variation in home range overlap across brown bear (*Ursus arctos*) populations. Costs and benefits of territoriality may also vary according to predator and competitor densities (Maher & Lott, 2000; Webber et al., 2023), individual characteristics such as sex (Rosell & Thomsen, 2006; Fawcett et al., 2013), or other variables such as the degree of relatedness among neighbours (Persson et al., 2010; Humphries et al., 2021).

The probability of encountering neighbours varies spatially in the home range of territorial
animals, being higher near boundaries. Where the probability of encountering a competitor is high,
individuals should modify their behaviour to avoid potentially costly physical encounters while
still signaling territory ownership more intensively (Schlägel et al., 2017). For example, when
located near boundaries, white-faced capuchins (*Cebus capucinus*), which defend group territories,
tend to socialise less with each other (Tórrez-Herrera et al., 2020) and travel at lower speed
(Noonan et al., 2021), suggesting more vigilance. Also, Ethiopian wolves (*Canis simensis*) and
grey wolves (*Canis lupus*) increase scent-marking near boundaries (Sillero-Zubiri & Macdonald,
1998; Zub et al., 2003), while red foxes (*Vulpes vulpes*) spend more time patrolling where artificial
scent marks are deposited (Arnold et al., 2011).

An increase in time spent vigilant or in signaling territory ownership when the probability of
encountering a competitor is high could further result in fewer time spent foraging (Laundré et al.,
2001). Furthermore, probability of encounter may directly affect specific foraging behaviour. For
example, in food caching animals that are territorial, such as many canids, felids and mustelids, it
may be beneficial to cache food away from competitors to reduce cache pilfering (van der Veen et
al., 2020). Better understanding how intraspecific interactions and territoriality shape foraging
behaviours could be particularly important for predators, as they generate predation risk landscapes
that influence the dynamics of prey populations and community structure (Lima, 2002; Gaynor et
al., 2019; Clermont et al., 2021a). For example, in systems where the predator is highly territorial
and avoids encountering conspecifics, buffer zones between territories may serve as refuge for
prey (Mech, 1977; Lewis & Murray, 1993). In support of this, Anderson et al. (2005) found that
reintroduced elks (*Cervus elaphus*) established their home ranges in the periphery of wolf
territories and selected areas within their home ranges that were further from wolf territory centers.
Prey may thus benefit from fewer predator encounter in places where territorial interference is high
and could be further relieved if predators also spend less time foraging in these areas.

Modern tracking techniques help to study predator behaviours as they can locate animals at high
frequencies while producing detailed behavioural classification (Nathan et al., 2012; Wilmers et
al., 2015). For example, activity level can be identified using GPS data (Patterson et al., 2017),
while scent marking (Bidder et al., 2020), killing of prey (Studd et al., 2021) and food caching
(Clermont et al., 2021b) may be identified using accelerometry. Estimating where encounters

between individuals occur is however challenging, limiting our ability to assess how probabilities

of encounter affect behaviours (Noonan et al., 2021). So far, most research has compared how

animals behave in and out of overlapping areas or with respect to the distance to territory borders

(assuming these variables are good surrogates for encounter probability), instead of directly

estimating the spatial distribution of encounter probabilities (Rosell & Thomsen, 2006; Tórrez-

Herrera et al., 2020). To address this methodological gap, Noonan et al. (2021) proposed a

statistical framework using tracking data and home range estimation to evaluate the spatial

distribution of the probability of encounter, termed the conditional distribution of encounters

(CDE). The CDE quantifies where in space encounters are likely to occur within an individual's

home range. Combining behavioural classifications to the CDE should allow to better understand

how interactions with neighbours affect behaviours of territorial animals, further improving our

understanding of important ecological processes.

We used GPS tracking and accelerometry to monitor the movements and behaviours of Arctic

foxes living in a large Greater snow goose (*Anser caerulescens atlanticus*) colony on Bylot Island

(Nunavut, Canada) (Clermont et al., 2021a, 2021b) during two years of contrasting resource

availability. At this site, foxes are socially monogamous, offer biparental care, and most home

ranges are occupied by a mated pair (Cameron et al., 2011). Food caching is an important

dimension of the foraging ecology of Arctic foxes, which cache the majority of the goose eggs

they collect (Careau et al., 2007). Our first objective was to quantify the degree of summer

territoriality of each studied individual by mapping the probability of encountering conspecifics

other than the mate within the home range, and to evaluate whether fox territoriality depends on

the availability of food resources. Following Maher and Lott's (2000) model predictions, we

predicted that fewer foxes would be territorial when resources are scarce (P1). Our second

objective was to test the hypothesis that the probability of encountering a neighbour influences the

spatial distribution of foraging behaviours within home ranges, and test whether territoriality

modulates this relationship. As Arctic foxes may attack other foxes foraging within their territory

and perform cache pilfering (Samelius & Alisauskas, 2000; Careau et al., 2007), we predicted (P2)

that foxes should be less likely to hunt and cache prey where the probability of encountering a

neighbour and the risk of cache pilfering is high. Lastly, we predicted (P3) that the probability of

encountering a neighbour should more strongly affect the foraging and caching behaviour of non-

territorial than territorial foxes, since the latter individuals might focus on excluding other foxes
from their territory rather than avoiding encounters.

Materials and methods

Study system

We worked in 2019 and 2022 in the southwest plain of Bylot Island (72°53′ N, 79°54′ W) in
Sirmilik National Park of Canada (Nunavut), where the Arctic fox is the main terrestrial predator.
Arctic foxes at this site are highly range resident (i.e., they use a stable home range), they usually
share a home range with their mating partner and most individuals show low home range overlap
with their neighbours (Grenier-Potvin et al., 2021). They bark and scent mark to indicate territory
ownership, and they chase intruders (Eberhardt et al., 1982). In addition to selecting habitats
suitable to their main prey, they avoid home range borders, potentially to minimise interactions
with their neighbours (Grenier-Potvin et al., 2021).

On Bylot, Arctic foxes rely mostly on small prey, such as lemmings (*Lemmus trimucronatus* and
Dicrostonyx groenlandicus). They also feed on greater snow goose eggs when their home range
overlaps Bylot's large colony of more than 20,000 nesting pairs (Bêty et al., 2001). In 2022
however, nesting goose density was unusually low throughout the whole colony (density estimated
from random plots: 128 nests/km^2 in 2022, 438 nests/km^2 in 2019 and an average of 248 nests/km^2
between 2000-2019; Cadieux et al., 2023). Foxes also opportunistically prey on the nests of other
ground nesting birds (Duchesne et al., 2021). Lemming density fluctuates cyclically (Gruyer et al.,
2008), and was moderate in 2019 and very low in 2022 as determined by capture-recapture
methods (Fauteux et al., 2015; Duchesne et al., 2021; Cadieux et al., 2023). The snow goose
incubation period lasts 23 days from mid-June to early July, during which foxes collect eggs for
later consumption (Samelius et al., 2007). In years of low to moderate lemming densities, goose
eggs represent the majority of prey collected by foxes during the goose incubation period (Careau
et al., 2007). Foxes cache up to 90% of the goose eggs they collect, and caches are located 85 m
(median) from the nest (Careau et al., 2007). They also recache ca. 60% of goose eggs, and
recovery rate (for recaching or consumption) is highest at the end and after the goose incubation
period (Careau et al., 2008).

Furthermore, access to the snow goose colony influences fox probability of reproduction
(Chevallier et al., 2020). Despite interindividual variations, the phenology of fox reproduction
largely coincides with that of geese, with cubs remaining in the den during the goose egg
incubation period, and progressively emerging and becoming independent from the den during
goose brooding (Grenier-Potvin et al., 2021). Due to low densities of both lemmings and goose
nests in 2022, none of the studied foxes reproduced that year, contrary to 2019 when many foxes
reproduced.

Arctic fox tracking

In May and June, we captured 13 foxes in 2019 and 10 foxes in 2022 in the snow goose colony
using Softcatch #1 padded leghold traps (Oneida Victor Inc. Ltd., Cleveland, OH, USA). We
determined the individual's sex from genital characteristics and fitted them with four coloured-ear
tags and a GPS-accelerometer collar (95 g, ca. 4% of body mass; Radio Tag-14, Milsar, Romania)
for subsequent identification and tracking. Arctic fox capture techniques and immobilization
procedures were approved by the UQAR Animal Care Committee (CPA-64-16-169 R3) and field
research was approved by the Joint Park Management Committee of Sirmilik National Park of
Canada (SIR-2018-28021).

In 2019, the 13 tracked foxes consisted of six neighbouring pairs plus one individual, and their
home ranges formed our 2019 study area (Figures 1A and 1B). In 2022, the 10 tracked foxes
included one pair for which both individuals were tracked, and eight pairs for which only one
individual of the pair was tracked, and their home ranges formed our 2022 study area (Figures 1C
and 1D). We determined reproductive status of individuals based on whether automated cameras
recorded cubs at individuals' den (Cameron et al., 2011). Daily observations and automated
cameras at fox dens confirmed that all foxes within our study area were tracked in 2019 and at
least one individual per pair in 2022. We collected one GPS location every 4 minutes and one 30-
sec burst of accelerometry (50 Hz) every 4.5 minutes in 2019 and every 4 minutes in 2022
(Clermont et al., 2021b). We divided the 30-sec bursts of accelerometry into ten 3-sec sequences
and, using behavioural observations and a random forest algorithm, we assigned to each sequence
one of four behaviours: running, walking, digging and motionless (see analytical details in
Clermont et al., 2021b). During the goose incubation period, digging behaviour of foxes is mainly
associated with goose egg caching, but may also occasionally reflect caching and handling of other

prey species as well as cache recoveries (Clermont et al., 2021b). As most captured prey are cached

(Careau et al., 2007), fox digging behaviour is more generally a proxy of foraging activity.

Figure 1. GPS locations (dots) of male and female Arctic foxes tracked from 11 June to 4 July
2019 when resources were abundant (n = 67,109, panels A and B) and from 19 June to 12 July

2022 when resources were scarce (n = 47,631, panels C and D) at a 4-min fix interval on Bylot
Island, Nunavut, Canada. Solid (males) and dashed (females) lines are 95% home range contours
obtained through auto-correlated kernel density estimation. For each year, the same colours are
used for members of a given fox pair. Home ranges are located near the coast of Bylot island
shown by the thick grey line.

Data analysis

1) Estimation of homes ranges

To define Arctic fox summer home ranges, we used GPS data collected from 11 June to 4 July

2019, and 19 June to 12 July 2022, corresponding to goose egg incubation periods. Prior to home

range estimation, we confirmed home range residency for each individual using variogram analysis

(Fleming et al., 2014), and excluded location data resulting from extra-territorial trips as it affects

home range estimation (Calabrese et al., 2016). Indeed, during the study period, some individuals

made a few extra-territorial trips going up to 25 km away from the center of the home range. These

"outliers" were identified by evaluating distances between points and home range core, and a cut-

off distance was determined visually for each individual (see Appendix A for further details). We

excluded 11%, 9%, 6% and 4% of datapoints for four individuals that performed extra-territorial

trips (three individuals in 2019 and one in 2022). The remaining individuals showed high range

residency and we excluded < 1% of their GPS locations.

We then fitted range-resident continuous-time movement models to the data of each individual

(model selection resulted in the Ornstein-Uhlenbeck Foraging (OUF) process used for each

individual), to control for autocorrelation in both speed and location (Fleming et al., 2014;

Calabrese et al., 2016). Then, we estimated home range areas using auto-correlated kernel density

estimation (AKDE; Fleming et al., 2015). AKDE is more accurate than other home range

estimators for autocorrelated location data (Noonan et al., 2019). The complete workflow we used

is detailed in Silva et al. (2021), and was performed using the package ctmm (v0.6.2, Calabrese et

al., 2016) in R 4.1.0 (R Development Team, 2021).

2) Estimation of probability of encounter among neighbours

Following Noonan et al. (2021), we estimated the conditional distribution of encounters (CDE) for

each year separately (package ctmm v0.6.2). The CDE estimates the spatial distribution of

encounter events in the environment as the normalized product of AKDE home range estimates.

The probability of encounter estimated by the CDE thus represents the probability that more than
one individual uses a specific area, but not necessarily at the same time. A CDE value of 1 at a
given location indicates there is a 100% chance that more than one individual use that location,
and thus that the probability of encounter is high, and a value of 0 that only one individual uses
that location, meaning individuals should never encounter each other at that location. Since we
were interested in encounters among neighbours, we excluded interactions between mates within
their shared home range. We then determined the degree of territoriality of each fox and compared
the proportion of territorial individuals between years (P1). A fox was considered territorial if at
least some part of its home range was outside of the 95% CDE (Noonan et al., 2021), excluding
areas within 500 m of boundaries of study areas. Indeed, some interactions inevitably occurred at
home range boundaries located at the edge of the 2019 and 2022 study areas (but not near the
coastline, Figures 1 and 2), where neighbouring foxes were not tracked, leading to a local
underestimation of CDE values. Using a 1 km buffer instead of a 500 m one would have led to the
same classification (territorial yes/no) of all individuals but one in 2022 (female BJOV; see
Appendix B).

3) Arctic fox prey caching behaviour

We used our accelerometry behavioural classification to determine whether individuals engaged
in digging (indicating prey caching, cache recovery events and more generally foraging activity)
during each 30-sec accelerometry burst. At least one out of the ten 3-sec accelerometry sequences
had to be assigned to the digging behaviour to consider that the fox engaged in digging in that
burst. Each 30-sec burst was then associated to the closest GPS location, provided the time stamp
of the GPS location was within 30 sec of the start or end of the burst. Bursts occurring at less than
50 m of a den were excluded as digging may then reflect den maintenance rather than prey caching.

We excluded from analyses all accelerometry bursts occurring within 500 m of the boundaries of
study areas, since interactions with unknown foxes may then occur (Appendix B). To verify
whether buffer size influences results, we 1) repeated the analyses described in the next section by
excluding the individual of 2022 that would have been considered as non-territorial when using a
1 km buffer (female BJOV), and 2) repeated our analyses by excluding locations falling within a
1 km buffer instead of a 500 m one (Appendix B). In both cases, we obtained similar results.

4) Effect of the probability of encountering a neighbour on fox behaviours

To test the influence of the likelihood of encountering a neighbour on the probability of engaging

in foraging behaviours (objective 2), we used generalized additive mixed models (GAMM) with a

binomial error distribution and logit link function ("gam" function of the R package mgcv, v1.9-

0, Wood, 2017) and modelled fox probability of engaging in digging (0 = no digging, 1= digging;

P2) with respect to the probability of encounter. As not all foxes experienced the same range of

probability of encounter, we centered the probability of encounter values to a mean of 0 within

each fox home range. We modeled 2019 and 2022 data separately. In both models, we included as

fixed effect a categorical variable representing whether the fox was territorial (yes/no), which we

included in interaction with the probability of encounter, to test whether the degree of territoriality

modulated the effect of the probability of encounter on the probability of caching (P3). In 2019,

we also controlled for nesting goose density (individual geese/ha), a proxy for goose nest density

estimated from detailed field surveys (Grenier-Potvin et al., 2021), as nesting goose density was

previously shown to affect fox probability to dig (Clermont et al., 2021b). We did not perform

detailed field surveys in 2022 as nest density was very low throughout the study area, therefore we

could not include nesting goose density as a covariate in the 2022 models. We included individual

sex as another fixed effect, as well as fox identity and Julian date as random intercepts. To address

temporal autocorrelation in our datasets, we also added to all models a cyclic cubic spline function

of the numeric time of the day and a thin plate regression spline function of the Julian date, with

10 basis dimensions for each function (Pedersen et al., 2019). All continuous covariates were

centered and standardised to facilitate the interpretation of model estimates (Schielzeth, 2010). We

verified model assumptions and independence of residuals using the R package DHARMa (v0.4.3,

Hartig, 2021), and observed significant deviations in models' residuals. We solved these issues by

using a subset of data that included 150 randomly chosen observations by individual for both

models, and by keeping all covariates in the models even when effects were not significant. Results

are expressed using evidence-based language (Muff et al., 2021) where p-value ≥ 0.1 suggests no

evidence, p-value =]0.05, 0.1] suggests weak evidence, p-value =]0.01, 0.05] suggests moderate

evidence, p-value =]0.01, 0.001] suggests strong evidence and p-value < 0.001 suggests very

strong evidence.

Results

Conditional distribution of encounters in fox home ranges

Home range areas averaged 7.93 ± 3.15 (SD) km^2 (Figure 1, n = 23). We found no evidence that

home range areas differed between sexes (t-test: t = 0.53, p = 0.600, average male home range area

= 8.27 ± 3.12 (SD) km^2, n = 12, average female home range area = 7.56 ± 3.28 (SD) km^2, n = 11).

However, there was weak evidence that home ranges were smaller in 2022 than in 2019 (t-test: t

= 2.07, p = 0.051, average home range area in 2019 = 9.03 ± 3.12 (SD) km^2, n = 13, average home

range area in 2022 = 6.51 ± 2.70 (SD) km^2, n = 10).

As expected, higher CDE values occurred where home ranges of tracked neighbours overlapped

(Figure 2). When resources were abundant (2019), three individuals (green female RMJJ, green

male BORR and grey male BBJO) were non-territorial as they had all of their home range located

within the 95% CDE contour if we ignore 500 m study area edges (Figures 2A and 3A). These

three foxes did not reproduce in 2019. In contrast, all other ten individuals were territorial, having

a considerable area of their home range outside the 95% CDE contour, where the probability of

encounter with a neighbour is null (Figures 2A and 3A). These ten individuals reproduced in 2019.

When resources were rare (2022), all individuals but one (dark orange male BOBB) were territorial

(Figures 2B and 3B). None of the foxes reproduced in 2022. We thus find no support for an effect

of resource abundance on territoriality (P1) as 10/13 and 9/10 individuals were territorial in 2019

(high resource abundance) and 2022 (low resource abundance), respectively.

Figure 2. Conditional distribution of encounters (CDE) among A) 13 Arctic fox neighbours in 2019 when resources were abundant and B) ten Arctic fox neighbours in 2022 when resources were scarce, tracked with GPS, excluding interactions between pair members. The thick black line is the 95% CDE contour and coloured lines are 95% AKDE contours. The grey shaded area represents the 500 m inner buffer of study area boundaries where interactions with other foxes inevitably occur. Males and females are represented by solid and dashed lines, respectively, and members of a given pair are identified by the same colour within each year.

Figure 3. CDE values (probability of encountering a neighbour) observed at each GPS location from Figure 1, for each fox in A) 2019 and B) 2022. Fox colour IDs reflect the unique combinations of colours of the four ear tags (e.g., OBBB: orange-blue-blue-blue). Boxplot colours used for each individual are the same as in Figures 1 and 2. Males and females are represented by solid and dashed lines, respectively, and each year's members of a given pair are identified by the same colour. Boxplots show first quartile, median, and third quartile. Lower and upper whiskers extend, respectively, to the lowest and highest values within the interquartile range multiplied by 1.5. Points represent values outside this range.

Effect of the probability of encountering a neighbour on fox caching behaviour

When resources were abundant (2019), non-territorial foxes were more likely to engage in digging (strong evidence) in places with low encounter probability (Table 1; Figure 4A). However, when resource were scarce (2022), we found no evidence that the probability of encounter influenced digging behaviour of the only non-territorial individual (Table 1; Figure 4B). As for territorial foxes, we did not find evidence that digging behaviour was influenced by the probability of encountering a neighbour in either year (Table 1; Figure 4). Thus, when resources were abundant, the probability of encountering a neighbour only affected the behaviours of non-territorial individuals, and all individuals were unaffected when resources were scarce (Figure 4), providing partial support for an effect of encounter probability on foraging behaviours (P2) and for a larger influence of encounter probability in non-territorial foxes (P3). We also found very strong evidence that nesting goose density has a positive effect on the propensity to dig in the year where

that variable could be included (2019; Table 1). There was no evidence of an effect of sex in both
years (Table 1). To ensure results were robust despite the low sample size, we ran our models on
different subsets of the dataset and observed slight changes in coefficient values, but no differences
in the direction and significance of effects.

Table 1. Estimates of generalized additive mixed models predicting the probability of fox digging
in A) 2019 when resources were abundant (n = 1950 accelerometry bursts) and B) 2022 when
resources were scarce (n = 1500 accelerometry bursts). Models had a binomial error distribution
with a logit-link function. We included as parametric terms (fixed effects) the probability of
encountering a neighbour, whether the fox was territorial, an interaction between the probability
of encountering a neighbour and whether the fox was territorial, the individual's sex and goose
nest density (2019 only). We included splines for numeric time of day (ToD) and Julian date (day),
and the random effects fox identity (Fox ID) and Julian date (day_factor).

A) Probability of fox digging when resources are abundant (2019)

Component	Term	Estimate	Std.Error	z.value	p.value
1. Parametric coefficients	(Intercept)	-0.91	0.30	-3.03	0.002
	Probability of encountering a neighbour	0.07	0.07	0.96	0.336
	Territorial yes/no [no]	-0.03	0.48	-0.07	0.946
	Goose nest density	0.37	0.06	6.17	<0.001
	Sex [male]	-0.09	0.40	-0.22	0.826
	Prob encounter: Territorial yes/no [no]	-0.45	0.15	-3.06	0.002

Component	Term	edf	Ref.df	Chi.sq	p.value
2. Smooth terms	s(ToD)	3.42	8.00	42.88	<0.001
	s(day)	2.76	3.46	6.43	0.159
	s(Fox ID)	9.15	10.00	104.81	<0.001
	s(day_factor)	0.22	21.00	0.22	0.404

B) Probability of fox digging when resources are scarce (2022)

Component	Term	Estimate	Std.Error	z.value	p.value
1. Parametric coefficients	(Intercept)	-0.37	0.23	-1.58	0.115
	Probability of encountering a neighbour	0.043	0.06	0.72	0.471
	Territorial yes/no [no]	0.346	0.57	0.60	0.546
	Sex [male]	-0.03	0.34	-0.10	0.920
	Prob encounter: Territorial yes/no [no]	-0.25	0.17	-1.48	0.138

Component	Term	edf	Ref.df	Chi.sq	p.value
2. Smooth terms	s(ToD)	4.20	8.00	84.67	<0.001
	s(day)	1.00	1.00	0.52	0.473
	s(Fox ID)	6.14	7.00	49.01	<0.001
	s(day_factor)	5.48	22.00	9.21	0.045

Figure 4. Predicted effects of the probability of encountering a neighbour on probability of fox digging (0 = no digging, 1= digging) in A) 2019 when resources were abundant (n = 1950 accelerometry bursts) and B) 2022 when resources were scarce (n = 1500 accelerometry bursts). The probability of encountering a neighbour was obtained from conditional distributions of encounters (Figure 2). Orange dots and lines refer to territorial individuals (n = 10 foxes in 2019 and 9 foxes in 2022), and blue dots and lines refer to non-territorial individuals (n = 3 foxes in 2019 and 1 fox in 2022). Shaded areas represent 95% confidence intervals.

Discussion

Using high throughput tracking technologies, we estimated the spatial distribution of the probability of encounter among Arctic foxes using neighbouring home ranges, during two years of contrasted resource abundance varying from high (2019) to low (2022). We benefited from a natural situation where most, but not all individuals were territorial, thus enabling us to evaluate the effect of territoriality on the spatial distribution of foraging behaviour. When resources were abundant, non-territorial individuals responded to the variation in the probability of encountering a neighbour by shifting the distribution of prey caching away from the places where encounters were most likely to occur. The only non-territorial individual monitored when resource abundance was low however did not respond to variation in the probability of encounter in its home range. The probability of encountering a neighbour also did not influence territorial individuals' prey caching behaviour. Despite modern technology, it remains highly challenging to simultaneously track the movements and foraging behaviours of many neighbouring vertebrate predators at high

resolution, in a natural setting and over several weeks. Our sample of 13 individuals in 2019 and

10 individuals in 2022, each tracked for more than three weeks, is therefore of high value. Our

study provides valuable insights into how intraspecific interactions may influence the spatial

distribution of predator foraging.

The degree of territoriality varies among foxes

Variation in the distribution, abundance and predictability of resources often explains among-

individual differences in territoriality (Maher & Lott, 2000; Sells & Mitchell, 2020). For example,

Eide et al. (2004) observed that in Arctic foxes of Svalbard, high home range overlap occurred

near the coast where seabird colonies are concentrated. On the other hand, little home range

overlap occurred inland where reindeer (*Rangifer tarandus*) carcasses are scattered and

unpredictable. In our study, we do not suspect that variation in prey distribution, abundance or

predictability explains among-individual differences in territoriality within years, as all studied

foxes lived near the center of the goose colony where nest abundance is usually high and habitats

favourable to lemmings. However, we used two contrasted years of data, with one year of moderate

lemming abundance and typical goose density for this study area (2019), and one year of low

lemming abundance and low goose density (2022). In 2019, although resource abundance was

high, not all foxes reproduced and used a territory. This suggests that using a territory may be

necessary to secure enough resources to ensure the survival of young, even when food is abundant

(Webb et al., 2012; López-Bao et al., 2014). On the other hand, for non-breeding individuals, the

costs of defending a territory may have been too high compared to the benefits when resources are

abundant, as predicted by Maher and Lott's (2000) cost-benefit model of territoriality.

Although resources were scarce in 2022, all foxes but one could exclude neighbours from part of

their home range, which seems to contradict Maher and Lott's (2000) model, that predicts no

territoriality when resources are very rare. This could however indicate that food abundance inside

the colony in 2022 was still not low enough for territoriality not to be beneficial. In fact, foxes in

the colony might rely on stored eggs from the previous year (Samelius et al., 2007), which could

attenuate the effects of low resources. However, although high abundance of resources may permit

non-breeding individuals to forego being territorial, defending a territory may remain beneficial

under low or very low resource abundance, especially when the abundance of resources varies

cyclically through time. Similarly, López-Bao et al. (2014) observed that home range overlap
among Iberian lynx *(Lynx pardinus)* remains low across a gradient of prey availability. This
obstinate strategy (sensu von Schantz, 1984), i.e., maintaining a territory even when resources are
rare, could be beneficial if it allowed to secure a home range in areas where resources are usually
high, such as the snow goose colony on Bylot Island. Therefore, it may remain beneficial for foxes
to defend a territory in the goose colony even during years of low goose nesting density, to
maintain territory ownership and maximise chances of future reproduction (López-Bao et al.,
2014).

Variable effect of the probability of encountering a neighbour on Arctic fox foraging behaviours

Foraging and caching prey in places where the probability of encountering a neighbour is high can
increase the risk of injury or cache pilfering (Samelius & Alisauskas, 2000; van der Veen et al.,
2020). Therefore, Arctic foxes should be more likely to forage and cache prey where they are less
likely to encounter neighbours. However, such behaviour can be influenced by the intrinsic and
environmental constraints faced by individuals. We found that resource abundance and the level
of territoriality influenced how Arctic foxes responded to variation in the probability of
encountering a neighbour. When resources were abundant, non-territorial foxes had a higher
probability to dig where the probability of encountering neighbours was low. For these individuals
without exclusive home range, foraging and caching prey where the risk of cache pilfering is low
may increase foraging efficiency. Although during goose incubation period, digging events should
mostly reflect prey caching, we cannot differentiate caching events from prey captures, cache
recoveries and recaching events (Clermont et al., 2021b). Therefore, our results may also indicate
that non-territorial individuals could favour recaching away from their neighbours, in safer sites.
Caching and recaching prey in "out of view" sites are in fact common cache protection strategies
in other species such as corvids (Dally et al., 2006). Careau et al. (2007) also showed that Arctic
fox recaches goose eggs away from were they were collected, potentially to secure eggs in sites
where chances of pilfering are lower (e.g., closer to the den). The non-territorial individual of 2022
however cached prey independently of its probability of encountering neighbours, suggesting that
when resources are scarce, non-territorial foxes may not afford to cache prey only in specific areas
of the landscape, but this result should be confirmed with a larger sample of individuals.

Territorial foxes did not adjust caching behaviours to the probability of encountering a neighbour
in either year of contrasted resource abundance. By defending a territory, these individuals had
null encounter probability in substantial parts of their home range. This likely reduced the need to
modulate foraging and caching behaviours, a potential benefit of establishing a territory. Still,
these individuals could have withheld from foraging and caching in the few parts of their home
range where encounter probability was not null. However, our results suggest that they did not,
potentially because focusing on maintaining and defending their territory is more important than
avoiding risk of injury or cache pilfering by conspecifics. Indeed, territorial individuals are likely
more dominant and better suited to win an encounter than non-territorial individuals (Stamps &
Krishnan, 2001). It might therefore be beneficial for these individuals to actively use all parts of
their home range at the risk of injury and cache pilfering rather than leaving some areas
undefended. Our results suggest that territoriality could be enough to secure the necessary
resources for survival and reproduction.

In addition to spatial adjustments of behaviours, Arctic foxes may modify their foraging
behaviours temporally to avoid encounters, which the CDE did not allow to evaluate (Noonan et
al., 2021). Similarly, prey species access resources in high-predation risk areas at times of the day
when predators are less active (Kohl et al., 2018; Smith et al., 2019b). Future studies should include
temporal partitioning when estimating the probability of encounter to help uncover the different
tactics territorial animals use to avoid intraspecific interactions while foraging.

Implications for predator-prey interactions

Arctic foxes have important top-down effects in the tundra and generate multiple predator-
mediated interactions among prey (Bêty et al., 2002; Legagneux et al., 2012; Duchesne et al.,
2021). For example, nest survival of American golden plover (*Pluvialis dominica*) is lower within
the snow goose colony, where foxes use smaller home ranges and are thus present at higher
densities (Dulude-de Broin et al., 2023). At fine spatial scales, Arctic fox movements generate a
predation risk landscape influencing the behaviour and nest distribution of several migratory birds
(Clermont et al., 2021a). It is therefore of strong interest to identify the factors explaining Arctic
fox density and where they choose to forage. In this study, we observed that the degree of
territoriality and food abundance influence fox response to intraspecific interactions. When

resources are abundant, non-territorial individuals avoid caching prey where the probability of
encountering a neighbour is high. As most captured prey are also cached (Careau et al., 2007), our
results suggest non-territorial individuals forage less intensively in areas likely used by other
individuals. However, there was no effect of the probability of encounter on the caching behaviour
of territorial individuals (i.e., most individuals), suggesting intraspecific interactions may not have
a strong effect on the spatial distribution of predation risk. Interestingly, although most Arctic
foxes were territorial, we observed no unused areas between adjacent home ranges, but instead
high probability of encounters in overlapping areas (Figures 1 and 2). As such, the spatial
configuration of fox home ranges should not lead to buffer zones of low predation risk between
territories as observed in some wolf-ungulate systems (Mech, 1977; Anderson et al., 2005).
Conversely, the presence of non-territorial individuals, whose home ranges overlap those of their
territorial neighbours, creates zones where fox density is high, which can increase the risk of
predation (Clermont et al., 2021a; Dulude-de Broin et al., 2023). As such, we hypothesise that the
greatest risk of predation should be located within the home range of non-territorial individuals,
while the lowest predation risk should be found in defended territories where the probability of
encounter among neighbouring foxes is null.

Conclusion

We found that not all Arctic foxes on Bylot Island used a territory. We highlight that because costs
and benefits of territoriality may differ among individuals in a population, alternative behavioural
tactics may emerge from non-territorial individuals to secure resources. Including the conditional
distribution of encounters into habitat selection analyses would further help to better understand
how different individuals deal with both the physical and social environments when using their
habitat, and how they compromise between the needs to acquire valuable resources and avoid
competitors. Finally, a spatial configuration in which the home ranges of non-territorial individual
predators overlap those of their territorial neighbours may influence the distribution of predation
risk, which could modulate the distribution and behaviour of prey and the structure of
communities.

Data Accessibility Statement

Arctic fox GPS and accelerometry data are available through the Movebank Data Repository
(Movebank Study ID 1241071371 and 2602389249).

Competing Interests Statement

The authors declare no competing interests.

Author Contributions

All authors conceived the study and collected the data. JC performed the analyses with help from
FDB and MPP. JC wrote the first version of the manuscript, which all authors commented and
revised. All authors approved the final version.

Acknowledgements

We thank A. Grenier-Potvin, R. Gravel, G. Roy, L.-P. Ouellet, M. Beaudoin and M. Archambault
for field work, M. Noonan for advice on spatial analyses, and the community of Mittimatalik for
support. Funding was received from (alphabetical order) Canada Foundation for Innovation,
Canada Research Chairs Program, Fonds de Recherche du Québec – Nature et technologies
(FRQNT), Natural Sciences and Engineering Research Council of Canada (NSERC), Network of
Centers of Excellence of Canada ArcticNet, Northern Scientific Training Program (Polar
Knowledge Canada), Parks Canada Agency, Polar Continental Shelf Program (Natural Resources
Canada), and Weston Family Foundation.

Literature cited

Anderson, D.P., Turner, M.G., Forester, J.D., Zhu, J., Boyce, M.S., Beyer, H., Stowell, L. 2005.
Scale-dependent summer resource selection by reintroduced elk in Wisconsin, USA.
Journal of Wildlife Management 69, 298–310.
Arnold, J., Soulsbury, C.D., Harris, S. 2011. Spatial and behavioral changes by red foxes (*Vulpes*
vulpes) in response to artificial territory intrusion. Canadian Journal of Zoology 89, 808–
815.
Bêty, J., Gauthier, G., Giroux, J.-F., Korpimäki, E. 2001. Are goose nesting success and
lemming cycles linked? Interplay between nest density and predators. Oikos 93, 388–400.

Bêty, J., Gauthier, G., Korpimaki, E., Giroux, J.-F. 2002. Shared predators and indirect trophic
interactions: lemming cycles and Arctic-nesting geese. Journal of Animal Ecology 71,
88–98.
Bidder, O.R., Virgilio, A. di, Hunter, J.S., McInturff, A., Gaynor, K.M., Smith, A.M., Dorcy, J.,
Rosell, F. 2020. Monitoring canid scent marking in space and time using a biologging
and machine learning approach. Scientific Reports 10, 588.
Brown, J.L., Orians, G.H. 1970. Spacing patterns in mobile animals. Annual Review of Ecology
and Systematics 1, 239–262.
Cadieux, M.-C., Gauthier, G., Lefebvre, J., Fauteux, D., Berteaux, D., Legagneux, P. 2023.
Population study of greater snow geese and its nesting habitat on Bylot Island, Nunavut
in 2022: A progress report. Technical Report.
Calabrese, J.M., Fleming, C.H., Gurarie, E. 2016. ctmm: an R package for analyzing animal
relocation data as a continuous-time stochastic process. Methods in Ecology and
Evolution 7, 1124–1132.
Cameron, C., Berteaux, D., Dufresne, F. 2011. Spatial variation in food availability predicts
extrapair paternity in the arctic fox. Behavioral Ecology 22, 1364–1373.
Careau, V., Giroux, J.-F., Berteaux, D. 2007. Cache and carry: hoarding behavior of Arctic fox.
Behavioral Ecology and Sociobiology 62, 87–96.
Careau, V., Lecomte, N., Bêty, J., Giroux, J.-F., Gauthier, G., Berteaux, D. 2008. Hoarding of
pulsed resources: temporal variations in egg-caching by arctic fox. Écoscience 15, 268–
276.
Chevallier, C., Gauthier, G., Lai, S., Berteaux, D. 2020. Pulsed food resources affect
reproduction but not adult apparent survival in arctic foxes. Oecologia 193, 557–569.
Clermont, J., Grenier-Potvin, A., Duchesne, É., Couchoux, C., Dulude-de Broin, F., Beardsell,
A., Bêty, J., Berteaux, D. 2021a. The predator activity landscape predicts the anti-
predator behavior and distribution of prey in a tundra community. Ecosphere 12, e03858.
Clermont, J., Woodward-Gagné, S., Berteaux, D. 2021b. Digging into the behaviour of an active
hunting predator: arctic fox prey caching events revealed by accelerometry. Movement
Ecology 9, 58.
Cresswell, W. 2008. Non-lethal effects of predation in birds. Ibis 150, 3–17.
Dally, J.M., Clayton, N.S., Emery, N.J. 2006. The behaviour and evolution of cache protection
and pilferage. Animal Behaviour 72, 13–23.
Duchesne, É., Lamarre, J.-F., Gauthier, G., Berteaux, D., Gravel, D., Bêty, J. 2021. Variable
strength of predator-mediated effects on species occurrence in an arctic terrestrial
vertebrate community. Ecography 44, 1236–1248.
Dulude-de Broin, F., Clermont, J., Beardsell, A., Ouellet, L.-P., Legagneux, P., Bêty, J.,
Berteaux, D. 2023. Predator home range size mediates indirect interactions between prey
species in an arctic vertebrate community. Journal of Animal Ecology 92, 2373–2385.
Eberhardt, L.E., Hanson, W.C., Bengtson, J.L., Garrott, R.A., Hanson, E.E. 1982. Arctic fox
home range characteristics in an oil-development area. The Journal of Wildlife
Management 46, 183.
Eide, N.E., Jepsen, J.U., Prestrud, P. 2004. Spatial organization of reproductive Arctic foxes
Alopex lagopus: responses to changes in spatial and temporal availability of prey. Journal
of Animal Ecology 73, 1056–1068.
Fauteux, D., Gauthier, G., Berteaux, D. 2015. Seasonal demography of a cyclic lemming
population in the Canadian Arctic. Journal of Animal Ecology 84, 1412–1422.

Fawcett, J.K., Fawcett, J.M., Soulsbury, C.D. 2013. Seasonal and sex differences in urine marking rates of wild red foxes Vulpes vulpes. Journal of Ethology 31, 41–47.

Fleming, C.H., Calabrese, J.M., Mueller, T., Olson, K.A., Leimgruber, P., Fagan, W.F. 2014. From fine-scale foraging to home ranges: a semivariance approach to identifying movement modes across spatiotemporal scales. The American Naturalist 183, E154–E167.

Fleming, C.H., Fagan, W.F., Mueller, T., Olson, K.A., Leimgruber, P., Calabrese, J.M. 2015. Rigorous home range estimation with movement data: a new autocorrelated kernel density estimator. Ecology 96, 1182–1188.

Gaynor, K.M., Brown, J.S., Middleton, A.D., Power, M.E., Brashares, J.S. 2019. Landscapes of fear: spatial patterns of risk perception and response. Trends in Ecology & Evolution 34, 355–368.

Giuggioli, L., Potts, J.R., Harris, S. 2011. Animal interactions and the emergence of territoriality. PLoS ONE 7, e1002008.

Grenier-Potvin, A., Clermont, J., Gauthier, G., Berteaux, D. 2021. Prey and habitat distribution are not enough to explain predator habitat selection: addressing intraspecific interactions, behavioural state and time. Movement Ecology 9, 12.

Gruyer, N., Gauthier, G., Berteaux, D. 2008. Cyclic dynamics of sympatric lemming populations on Bylot Island, Nunavut, Canada. Canadian Journal of Zoology 86, 910–917.

Hartig, F. 2021. DHARMa: residual diagnostics for hierarchical (multilevel/mixed) regression models. https://CRAN.R-project.org/ package=DHARMa.

Hinsch, M., Komdeur, J. 2017. What do territory owners defend against? Proceedings: Biological Sciences 284, 1–8.

Humphries, D.J., Nelson-Flower, M.J., Bell, M.B.V., Finch, F.M., Ridley, A.R. 2021. Kinship, dear enemies, and costly combat: the effects of relatedness on territorial overlap and aggression in a cooperative breeder. Ecology and Evolution 11, 17031–17042.

Kohl, M.T., Stahler, D.R., Metz, M.C., Forester, J.D., Kauffman, M.J., Varley, N., White, P.J., Smith, D.W., MacNulty, D.R. 2018. Diel predator activity drives a dynamic landscape of fear. Ecological Monographs 88, 638–652.

Laundré, J.W., Hernández, L., Altendorf, K.B. 2001. Wolves, elk, and bison: reestablishing the "landscape of fear" in Yellowstone National Park, U.S.A. Canadian Journal of Zoology 79, 1401–1409.

Legagneux, P., Gauthier, G., Berteaux, D., Bêty, J., Cadieux, M.-C., Bilodeau, F., Bolduc, E., McKinnon, L., Tarroux, A., Therrien, J.-F., Morissette, L., Krebs, C.J. 2012. Disentangling trophic relationships in a High Arctic tundra ecosystem through food web modeling. Ecology 93, 1707–1716.

Lewis, M.A., Murray, J.D. 1993. Modelling territoriality and wolf–deer interactions. Nature 366, 738–740.

Lima, S.L. 2002. Putting predators back into behavioral predator–prey interactions. Trends in Ecology & Evolution 17, 70–75.

López-Bao, J.V., Rodríguez, A., Delibes, M., Fedriani, J.M., Calzada, J., Ferreras, P., Palomares, F. 2014. Revisiting food-based models of territoriality in solitary predators. Journal of Animal Ecology 83, 934–942.

Macdonald, D.W., Courtenay, O., Forbes, S., Mathews, F. 1999. The red fox (Vulpes vulpes) in Saudi Arabia: loose-knit groupings in the absence of territoriality. Journal of Zoology 249, 383–391.

Maher, C.R., Lott, D.F. 2000. A review of ecological determinants of territoriality within
vertebrate species 143, 1–29.
McLoughlin, P.D., Ferguson, S.H., Messier, F. 2000. Intraspecific variation in home range
overlap with habitat quality: a comparison among brown bear populations. Evolutionary
Ecology 14, 39–60.
Mech, L.D. 1977. Wolf-pack buffer zones as prey reservoirs. Science 198, 320–321.
Muff, S., Nilsen, E.B., O'Hara, R.B., Nater, C.R. 2021. Rewriting results sections in the
language of evidence. Trends in Ecology & Evolution 37, 203–210.
Nathan, R., Spiegel, O., Fortmann-Roe, S., Harel, R., Wikelski, M., Getz, W.M. 2012. Using tri-
axial acceleration data to identify behavioral modes of free-ranging animals: general
concepts and tools illustrated for griffon vultures. Journal of Experimental Biology 215,
986–996.
Noonan, M.J., Martinez-Garcia, R., Davis, G.H., Crofoot, M.C., Kays, R., Hirsch, B.T., Caillaud,
D., Payne, E., Sih, A., Sinn, D.L., Spiegel, O., Fagan, W.F., Fleming, C.H., Calabrese,
J.M. 2021. Estimating encounter location distributions from animal tracking data.
Methods in Ecology and Evolution 12, 1158–1173.
Noonan, M.J., Tucker, M.A., Fleming, C.H., Akre, T.S., Alberts, S.C., Ali, A.H., Altmann, J.,
Antunes, P.C., Belant, J.L., Beyer, D., Blaum, N., Böhning-Gaese, K., Cullen, L., Paula,
R.C., Dekker, J., Drescher-Lehman, J., Farwig, N., Fichtel, C., Fischer, C., Ford, A.T.,
Goheen, J.R., Janssen, R., Jeltsch, F., Kauffman, M., Kappeler, P.M., Koch, F., LaPoint,
S., Markham, A.C., Medici, E.P., Morato, R.G., Nathan, R., Oliveira-Santos, L.G.R.,
Olson, K.A., Patterson, B.D., Paviolo, A., Ramalho, E.E., Rösner, S., Schabo, D.G.,
Selva, N., Sergiel, A., Xavier da Silva, M., Spiegel, O., Thompson, P., Ullmann, W.,
Zięba, F., Zwijacz-Kozica, T., Fagan, W.F., Mueller, T., Calabrese, J.M. 2019. A
comprehensive analysis of autocorrelation and bias in home range estimation. Ecological
Monographs 89, e01344.
Ord, T.J. 2021. Costs of territoriality: a review of hypotheses, meta-analysis, and field study.
Oecologia 197, 615–631.
Patterson, T.A., Parton, A., Langrock, R., Blackwell, P.G., Thomas, L., King, R. 2017. Statistical
modelling of individual animal movement: an overview of key methods and a discussion
of practical challenges. AStA Advances in Statistical Analysis 101, 399–438.
Pedersen, E.J., Miller, D.L., Simpson, G.L., Ross, N. 2019. Hierarchical generalized additive
models in ecology: an introduction with mgcv. PeerJ 7, e6876.
Persson, J., Wedholm, P., Segerström, P. 2010. Space use and territoriality of wolverines (Gulo
gulo) in northern Scandinavia. European Journal of Wildlife Research 56, 49–57.
Powell, R.A. 2000. Animal home ranges and territories and home range estimators, in: Boitani,
L., Fuller, T.F. (Eds.), Research Techniques in Animal Ecology. Columbia University
Press, New York, NY, pp. 65–110.
R Development Team 2021. R: a language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna.
Rosell, F., Thomsen, L.R. 2006. Sexual dimorphism in territorial scent marking by adult
Eurasian beavers (Castor fiber). Journal of Chemical Ecology 32, 1301–1315.
Samelius, G., Alisauskas, R.T. 2000. Foraging patterns of arctic foxes at a large arctic goose
colony. Arctic 53, 279–288.

Samelius, G., Alisauskas, R.T., Hobson, K.A., Larivière, S. 2007. Prolonging the arctic pulse:
long-term exploitation of cached eggs by arctic foxes when lemmings are scarce. Journal
of Animal Ecology 76, 873–880.
Schielzeth, H. 2010. Simple means to improve the interpretability of regression coefficients.
Methods in Ecology and Evolution 1, 103–113.
Schlägel, U.E., Merrill, E.H., Lewis, M.A. 2017. Territory surveillance and prey management:
wolves keep track of space and time. Ecology and Evolution 7, 8388–8405.
Sells, S.N., Mitchell, M.S. 2020. The economics of territory selection. Ecological Modelling 438,
109329.
Sih, A. 1984. The behavioral response race between predator and prey. The American Naturalist
123, 143–150.
Silk, J.B., Alberts, S.C., Altmann, J. 2003. Social bonds of female baboons enhance infant
survival. Science 302, 1231–1234.
Sillero-Zubiri, C., Macdonald, D.W. 1998. Scent-marking and territorial behaviour of Ethiopian
wolves Canis simensis. Journal of Zoology 245, 351–361.
Silva, I., Fleming, C.H., Noonan, M.J., Alston, J., Folta, C., Fagan, W.F., Calabrese, J.M. 2021.
Autocorrelation-informed home range estimation: A review and practical guide. Methods
in Ecology and Evolution 13, 534–544.
Smith, J.A., Donadio, E., Pauli, J.N., Sheriff, M.J., Bidder, O.R., Middleton, A.D. 2019a. Habitat
complexity mediates the predator–prey space race. Ecology 100, e02724.
Smith, J.A., Donadio, E., Pauli, J.N., Sheriff, M.J., Middleton, A.D. 2019b. Integrating temporal
refugia into landscapes of fear: prey exploit predator downtimes to forage in risky places.
Oecologia 189, 883–890.
Stamps, J.A., Krishnan, V.V. 2001. How territorial animals compete for divisible space: a
learning-based model with unequal competitors. The American Naturalist 157, 154–169.
Studd, E.K., Derbyshire, R.E., Menzies, A.K., Simms, J.F., Humphries, M.M., Murray, D.L.,
Boutin, S. 2021. The purr-fect catch: using accelerometers and audio recorders to
document kill rates and hunting behaviour of a small prey specialist. Methods in Ecology
and Evolution 12, 1277–1287.
Tórrez-Herrera, L.L., Davis, G.H., Crofoot, M.C. 2020. Do monkeys avoid areas of home range
overlap because they are dangerous? A test of the risk hypothesis in white-faced capuchin
monkeys (Cebus capucinus). International Journal of Primatology 41, 246–264.
van der Veen, B., Mattisson, J., Zimmermann, B., Odden, J., Persson, J. 2020. Refrigeration or
anti-theft? Food-caching behavior of wolverines (Gulo gulo) in Scandinavia. Behavioral
Ecology and Sociobiology 74, 52.
von Schantz, T. 1984. Spacing Strategies, kin Selection, and Population Regulation in Altricial
Vertebrates. Oikos 42, 48–58.
Webb, W.C., Marzluff, J.M., Hepinstall-Cymerman, J. 2012. Differences in Space Use by
Common Ravens in Relation to Sex, Breeding Status, and Kinship. The Condor 114,
584–594.
Webber, Q.M.R., Dantzer, B., Lane, J.E., Boutin, S., McAdam, A.G. 2023. Density-dependent
plasticity in territoriality revealed using social network analysis. Journal of Animal
Ecology 92, 207–221.
Wey, T., Blumstein, D.T., Shen, W., Jordán, F. 2008. Social network analysis of animal
behaviour: a promising tool for the study of sociality. Animal Behaviour 75, 333–344.

Wilmers, C.C., Nickel, B., Bryce, C.M., Smith, J.A., Wheat, R.E., Yovovich, V. 2015. The
701   golden age of bio-logging: how animal-borne sensors are advancing the frontiers of
702   ecology. Ecology 96, 1741–1753.
703   Wood, S.N. 2017. Generalized Additive Models: an introduction with R, 2nd ed. CRC Press,
704   Florida.
705   Zub, Kowalczyk, Jedrzejewska, Theuerkauf, Schmidt, Jedrzejewski 2003. Wolf pack territory
706   marking in the Bialowieza primeval forest (Poland). Behaviour 140, 635–648.
707

Appendix A: Workflow to exclude GPS locations associated to extra-territorial trips for home range estimation

Some individuals performed extra-territorial trips (Figure A1), which affected home range estimation. We considered these GPS locations associated to extra-territorial trips as outliers and identified them by calculating the distance between the core of the home range and each GPS location in the R package ctmm with the function 'outlie' (Calabrese et al., 2016). Figure A2 shows distance values for the individual BORR in 2019 (green male on Figure 1A of the main text) and VOVV in 2022 (dark purple male on Figure 1C of the main text). Zooming in on lower distance values helped to determine cut-offs for individuals travelling far from their home range core (Figure A3). For BORR, we removed GPS locations >1700 m from the home range core, which removed 9% of its original dataset. For VOVV, we removed GPS locations >2300 m from the home range core, which removed 11% of its original dataset. These thresholds were determined visually so that extra-territorial trips disappeared from the spatial representation of the individual's movements (Figure A4). We also removed 4% of GPS locations for RMJJ in 2019 (green female on Figure 1B), 6% for BBJO in 2019 (grey male on Figure 1A), and < 1% for all others.

Figure A1. GPS locations for A) individual BORR in 2019 and B) individual VOVV in 2022, that performed a few extra-territorial trips during the study period. X and Y axis show distance from home range core.

Figure A2. Distances between GPS locations and the core of the home range for A) individual BORR in 2019 and B) individual VOVV in 2022.

Figure A3. Zoom in on distances between GPS locations and the core of the home range for A) individual BORR in 2019 and B) individual VOVV in 2022. Red lines show the 1700 m and 2300 m cut-offs respectively.

736 **Figure A4.** GPS locations for A) individual BORR in 2019 and 2) VOVV in 2022 after the removal
737 of extra-territorial trips.

Appendix B: Exclusion of datapoints near study area boundaries

Interactions with foxes that were not tracked occurred on the outer margin of our study area, thus biasing negatively our CDE estimates. For that reason, when we determined whether foxes were territorial (i.e., whether they had a part of their home range located outside of the 95% CDE), we excluded areas within 500 m of study area boundaries (Figure 2). This led to three individuals considered as non-territorial in 2019 (green female RMJJ, green male BORR and grey male BBJO, Figure 2), and one in 2022 (dark orange male BOBB, Figure 2). Using a larger buffer size of 1 km would have led to the pink female BJOV in 2022 to also be considered as non-territorial, as the 1 km buffer overlaps all her home range (Figure C1).

To test the effect of the probability of encountering a neighbour on Arctic fox behaviour, we excluded from our dataset all locations falling within the 500 m inner buffer of the study area and considered that BJOV of 2022 was territorial (results in Table 1). Due to uncertainty in BJOV territorial degree, we repeated the analyses of 2022 while excluding this individual. Results are similar to those shown in Table 1 (Table C4).

We also verified whether our choice of buffer size influenced the results by repeating models, this time excluding locations falling within an inner buffer of 1 km instead of 500 m. This also resulted in excluding all locations of individual BJOV in 2022 as they were all within the 1 km buffer (Figure C1). Results are similar to those shown in Table 1 (Table C5).

Figure C1. Study area with 95% auto-correlated kernel density estimation contours for A) 7 male
and 6 female Arctic foxes in 2019 when resources were abundant and B) 5 male and 5 female
Arctic foxes in 2022 when resources were scarce. The light grey shaded area represents a 500 m
inner buffer of study area boundaries where observations were excluded, while the dark grey
shaded area indicates a 1 km buffer. The thick black line is the 95% CDE contour used to determine
whether foxes were territorial, and coloured lines are 95% AKDE contours. Males and females are
represented by solid and dashed lines, respectively, and members of a given pair are identified by
the same colour (within year).

766 **Table C4**. Estimates of generalized additive mixed models predicting the probability of fox
767 digging in 2022 (n = 1500 accelerometry bursts), while excluding female BJOV for which degree
768 of territoriality was uncertain. Model specifications are the same as those described in Table 1.

Component	Term	Estimate	Std.Error	z.value	p.value
1. Parametric coefficients	(Intercept)	-0.27	0.25	-1.07	0.285
	Probability of encountering a neighbour	0.070	0.06	1.14	0.254
	Territorial yes/no [no]	0.292	0.56	0.52	0.603
	Sex [male]	-0.14	0.35	-0.39	0.699
	Prob encounter: Territorial yes/no [no]	-0.08	0.17	-0.45	0.655

Component	Term	edf	Ref.df	Chi.sq	p.value
2. Smooth terms	s(ToD)	4.39	8.00	83.27	<0.001
	s(day)	1.00	1.00	3.21	0.073
	s(Fox ID)	5.23	6.00	40.30	<0.001
	s(day_factor)	4.64	22.00	6.26	0.163

Table C5. Estimates of generalized additive mixed models predicting the probability of fox digging in A) 2019 (n = 1950 accelerometry bursts) and B) 2022 (n = 1500 accelerometry bursts), while excluding locations falling within a 1 km inner buffer of study area boundaries. Model specifications are the same as those described in Table 1.

A) Probability of fox digging when resources are abundant (2019)

Component	Term	Estimate	Std.Error	z.value	p.value
1. Parametric coefficients	(Intercept)	-0.87	0.34	-2.56	0.011
	Probability of encountering a neighbour	-0.04	0.07	-0.50	0.619
	Territorial yes/no [no]	0.01	0.54	0.03	0.979
	Goose nest density	0.32	0.06	5.04	<0.001
	Sex [male]	-0.17	0.45	-0.38	0.700
	Prob encounter: Territorial yes/no [no]	-0.47	0.14	-3.29	0.001

Component	Term	edf	Ref.df	Chi.sq	p.value
2. Smooth terms	s(ToD)	2.89	8.00	33.98	<0.001
	s(day)	2.77	3.29	9.20	0.038
	s(Fox ID)	9.33	10.00	130.70	<0.001
	s(day_factor)	5.34	21.00	8.64	0.053

B) Probability of fox digging when resources are scarce (2022)

Component	Term	Estimate	Std.Error	z.value	p.value
1. Parametric coefficients	(Intercept)	-0.26	0.29	-0.90	0.371
	Probability of encountering a neighbour	-0.035	0.07	-0.51	0.608
	Territorial yes/no [no]	0.059	0.65	0.09	0.927
	Sex [male]	0.04	0.41	0.11	0.915
	Prob encounter: Territorial yes/no [no]	0.07	0.19	0.38	0.703

Component	Term	edf	Ref.df	Chi.sq	p.value
2. Smooth terms	s(ToD)	5.01	8.00	70.07	0.000
	s(day)	4.54	5.55	14.89	0.016
	s(Fox ID)	5.39	6.00	49.00	0.000
	s(day_factor)	0.00	22.00	0.00	0.817