Territoriality and resource availability modulate the effect of conspecific encounters on the

foraging behaviours of a mammalian predator

Authors: Jeanne Clermont[1,2,5,*], Frédéric Dulude-de Broin[3,5], Marie-Pier Poulin[4] & Dominique

Berteaux[2,5]

Authors affiliations:

1. Université de Sherbrooke, Sherbrooke (Québec), Canada.

2. Canada Research Chair on Northern Biodiversity, Université du Québec à Rimouski,

Rimouski (Québec), Canada.

3. Université Laval, Québec (Québec), Canada.

4. University of Wyoming, Laramie (Wyoming), USA.

5. Center for Northern Studies and Quebec Center for Biodiversity Science.

*Corresponding author information: email: jeanne_clermb@hotmail.com, full address:

Université de Sherbrooke, 2500 boulevard de l'Université, Sherbrooke (Québec), J1K 2R1,

Canada.

Abstract

The probability of encountering conspecifics shapes multiple dimensions of animal behaviour. For

example, territorial individuals increase vigilance, scent marking and alarm calling when

approaching home range boundaries. Whether territorial predators modify their foraging

behaviours with respect to the probability of encountering neighbouring territory owners is poorly

understood. However, this could strongly influence the landscape of predation risk and therefore

modulate predator-prey interactions. We studied the movements and behaviours of 23 resident

Arctic foxes occupying neighbouring home ranges during two years of contrasting resource

availability (abundant resources in 2019, scarce resources in 2022) on Bylot Island, Nunavut,

Canada. First, based on simultaneous GPS tracking of individuals, we established which

individuals used a territory (an exclusive area) by estimating the spatial distribution of the

probability of encountering a neighbour within their home range. Second, using GPS and

accelerometry data to identify prey searching and caching events, we evaluated if the probability

of encountering a neighbour influenced the spatial distribution of foraging behaviours, and whether

this relationship differed between territorial and non-territorial individuals. In 2019, when

resources were abundant, only breeding individuals excluded other foxes from a part of their home

range and could thus be considered territorial. In 2022, when resources were rare, none of the foxes

reproduced and all but one were territorial. Non-territorial individuals in 2019 were less active in

areas with a high probability of encounter, suggesting they searched prey less intensively and

engaged in more vigilance in these areas. They were also less likely to cache prey in areas with a

high probability of encounter, possibly reducing the risk of cache pilfering. We found no effect of

the probability of encounter on the behaviours of the non-territorial individual in 2022. Territorial

individuals in 2019 and 2022 were on the contrary more likely to be active in areas with a high

probability of encounter. Yet, their probability to cache prey did not depend on the probability of

encountering a neighbour, suggesting the increase in activity near borders may be related to an

increase in territory patrolling rather than in prey searching. Our results suggest that Arctic foxes

use different tactics to secure resources based on their degree of territoriality and the availability

of resources. We highlight how the presence of resident, but non-territorial predators, whose home

ranges overlap those of their territorial neighbours, may influence the distribution of predation risk

by creating zones where predator density is high, potentially influencing predator-prey
interactions.

Keywords: foraging, home range, intraspecific interactions, movement, territoriality, *Vulpes*
lagopus

Introduction

The ability of animals to manage encounters with other individuals is key to all social and
interspecific interactions, thus directly influencing reproduction, survival, and ultimately fitness
(Silk et al., 2003; Cresswell, 2008; Wey et al., 2008). Consequently, many animal behaviours have
evolved to modulate the probability of encountering other individuals. One well known example
is how predators and prey modify their habitat use to either find or avoid each other (Sih, 1984;
Smith et al., 2019a). Similarly, individuals may avoid conspecifics to decrease resource
competition. They for example secure limited resources by becoming territorial, given the benefits
of territoriality outweigh the costs (Maher & Lott, 2000; Ord, 2021). A territory represents the area
of a home range that is successfully defended by one or several individuals, resulting in the
territory owners having exclusive access to the defended area (Brown & Orians, 1970; Powell,
2000; Hinsch & Komdeur, 2017). Territorial individuals will often signal space ownership through
visual, auditory or olfactory signals and aggress intruders when encountered (Sillero-Zubiri &
Macdonald, 1998; Giuggioli et al., 2011; Tórrez-Herrera et al., 2020).

The intensity of territoriality is often assessed by measuring the amount of overlapping area
between neighbouring home ranges (Persson et al., 2010), or the frequency with which individuals
adopt territorial behaviours such as scent marking (Fawcett et al., 2013). Furthermore, the degree
to which individuals of a given species are territorial can vary among and within populations
(Macdonald et al., 1999; McLoughlin et al., 2000; Eide et al., 2004). Ecological variables such as
the abundance, predictability and distribution of food resources influence the costs and benefits of
territoriality, and therefore partly explain the degree of territoriality (Maher & Lott, 2000; Sells &
Mitchell, 2020). Notably, a cost-benefit model developed by Maher and Lott (2000) suggests that
individuals should be territorial when food abundance and predictability are intermediate, but non-
territorial when food abundance and predictability are below or above some thresholds, which was

supported by McLoughlin et al. (2000) who studied variation in home range overlap across brown

bear (*Ursus arctos*) populations. Costs and benefits of territoriality may also vary according to

predator and competitor densities (Maher & Lott, 2000; Webber et al., 2023), individual

characteristics such as sex (Rosell & Thomsen, 2006; Fawcett et al., 2013), or other variables such

as the degree of relatedness among neighbours (Persson et al., 2010; Humphries et al., 2021).

The probability of encountering neighbours varies spatially in the home range of territorial

animals, being higher near boundaries. Where the probability of encountering a competitor is high,

individuals should modify their behaviour to avoid potentially costly physical encounters while

still signaling territory ownership more intensively (Schlägel et al., 2017). For example, when

located near boundaries, white-faced capuchins (*Cebus capucinus*), which defend group territories,

tend to socialise less with each other (Tórrez-Herrera et al., 2020) and travel at lower speed

(Noonan et al., 2021), suggesting more vigilance. Also, Ethiopian wolves (*Canis simensis*) and

grey wolves (*Canis lupus*) increase scent-marking near boundaries (Sillero-Zubiri & Macdonald,

1998; Zub et al., 2003), while red foxes (*Vulpes vulpes*) spend more time patrolling where artificial

scent marks are deposited (Arnold et al., 2011).

An increase in time spent vigilant or in signaling territory ownership when the probability of

encountering a competitor is high could further result in fewer time spent foraging. Furthermore,

probability of encounter may directly affect specific foraging behaviour. For example, in food

caching animals that are territorial, such as many canids, felids and mustelids, it may be beneficial

to cache food away from competitors to reduce cache pilfering (van der Veen et al., 2020). Better

understanding how intraspecific interactions and territoriality shape foraging behaviours could be

particularly important for predators, as they generate predation risk landscapes that influence the

dynamics of prey populations and community structure (Lima, 2002; Gaynor et al., 2019;

Clermont et al., 2021a). For example, in systems where the predator is highly territorial and avoids

encountering conspecifics, buffer zones between territories may serve as refuge for prey (Mech,

1977; Lewis & Murray, 1993). In support of this, Anderson et al. (2005) found that reintroduced

elks (*Cervus elaphus*) established their home ranges in the periphery of wolf territories and

selected areas within their home ranges that were further from wolf territory centers. Prey may

thus benefit from fewer predator encounter in places where territorial interference is high and could

be further relieved if predators also spend less time foraging in these areas.

Modern tracking techniques help to study predator behaviours as they can locate animals at high
frequencies while producing detailed behavioural classification (Nathan et al., 2012; Wilmers et
al., 2015). For example, activity level can be identified using GPS data (Patterson et al., 2017),
while scent marking (Bidder et al., 2020), killing of prey (Studd et al., 2021) and food caching
(Clermont et al., 2021b) may be identified using accelerometry. Estimating where encounters
between individuals occur is however challenging, limiting our ability to assess how probabilities
of encounter affect behaviours (Noonan et al., 2021). So far, most research has compared how
animals behave in and out of overlapping areas or with respect to the distance to territory borders
(assuming these variables are good surrogates for encounter probability), instead of directly
estimating the spatial distribution of encounter probabilities (Rosell & Thomsen, 2006; Tórrez-
Herrera et al., 2020). To address this methodological gap, Noonan et al. (2021) proposed a
statistical framework using tracking data and home range estimation to evaluate the spatial
distribution of the probability of encounter, termed the conditional distribution of encounters
(CDE). The CDE quantifies where in space encounters are likely to occur within an individual's
home range. Combining behavioural classifications to the CDE should allow to better understand
how interactions with neighbours affect behaviours of territorial animals, further improving our
understanding of important ecological processes.

We used GPS tracking and accelerometry to monitor the movements and behaviours of Arctic
foxes living in a large Greater snow goose (*Anser caerulescens atlanticus*) colony on Bylot Island
(Nunavut, Canada) (Clermont et al., 2021a, 2021b). At this site, foxes are socially monogamous,
offer biparental care, and most home ranges are occupied by a mated pair (Cameron et al., 2011).
Our first objective was to quantify the degree of summer territoriality of each studied individual,
by mapping the probability of encountering conspecifics other than the mate within the home
range. Our second objective was to test the hypothesis that the probability of encountering a
neighbour influences the spatial distribution of foraging behaviours within home ranges, and test
whether territoriality modulates this relationship. We predicted (P1) that foxes should be less likely
to search prey when they are in areas associated to a high probability of encounter with neighbours,
as they should engage in more vigilance to avoid potentially costly encounters. As Arctic foxes
may perform cache pilfering (Samelius & Alisauskas, 2000; Careau et al., 2007), we also predicted
(P2) that foxes should be less likely to cache prey where the probability of encountering a
neighbour, and thus the risk of cache pilfering, is high. Lastly, we predicted (P3) that the

probability of encountering a neighbour should more strongly affect the foraging behaviours of
non-territorial than territorial foxes, since the latter individuals are free from encounters in some
fraction of their home range.

Materials and methods

Study system

We worked in 2019 and 2022 in the southwest plain of Bylot Island (72°53′ N, 79°54′ W) in
Sirmilik National Park of Canada (Nunavut), where the Arctic fox is the main terrestrial predator.
Arctic foxes at this site are highly range resident (i.e., they use a stable home range), they usually
share a home range with their mating partner and most individuals show low home range overlap
with their neighbours (Grenier-Potvin et al., 2021). They bark and scent mark to indicate territory
ownership, and they chase intruders (Eberhardt et al., 1982). In addition to selecting habitats
suitable to their main prey, they avoid home range borders, potentially to minimise interactions
with their neighbours (Grenier-Potvin et al., 2021).

On Bylot, Arctic foxes rely mostly on small prey, such as lemmings (*Lemmus trimucronatus* and
Dicrostonyx groenlandicus). They also feed on the eggs of the greater snow goose when their
home range overlaps Bylot's large colony of more than 20,000 nesting pairs (Bêty et al., 2001). In
2022 however, nesting goose density was unusually low throughout the whole colony (density
estimated from random plots: 72 nests/km^2 in 2022, 245 nests/km^2 in 2019 and an average of 150
nests/km^2 between 2000-2019). Foxes also opportunistically prey on the nests of other ground
nesting birds (Duchesne et al., 2021). Lemming density fluctuates cyclically (Gruyer et al., 2008),
and was moderate in 2019 and low in 2022 as determined by capture-recapture methods (Fauteux
et al., 2015; Duchesne et al., 2021; data unpublished for 2022). The snow goose incubation period
lasts 23 days from mid-June to early July, during which foxes collect eggs for later consumption
(Samelius et al., 2007). In years of low to moderate lemming densities, goose eggs represent the
majority of prey collected by foxes during the goose incubation period (Careau et al., 2007). Foxes
cache up to 90% of the goose eggs they collect, and caches are located 85 m (median) from the
nest (Careau et al., 2007). They also recache ca. 60% of the goose eggs, and recovery rate (for

recaching or consumption) is highest at the end and after the goose incubation period (Careau et
al., 2008).

Furthermore, access to the snow goose colony influences fox probability of reproduction
(Chevallier et al., 2020). Despite interindividual variation, the phenology of fox reproduction
largely coincides with that of geese, with cubs remaining in the den during the goose egg
incubation period, and progressively emerging and becoming independent from the den during
goose brooding (Grenier-Potvin et al., 2021). Due to low densities of both lemmings and goose
nests in 2022, none of the studied foxes reproduced that year, contrary to 2019 when many foxes
reproduced.

Arctic fox tracking

In May and June, we captured 13 foxes in 2019 and 10 foxes in 2022 in the snow goose colony
using Softcatch #1 padded leghold traps (Oneida Victor Inc. Ltd., Cleveland, OH, USA). We
determined the individual's sex from genital characteristics and fitted them with four coloured-ear
tags and a GPS-accelerometer collar (95 g, ca. 4% of body mass; Radio Tag-14, Milsar, Romania)
for subsequent identification and tracking. Arctic fox capture techniques and immobilization
procedures were approved by the UQAR Animal Care Committee (CPA-64-16-169 R3) and field
research was approved by the Joint Park Management Committee of Sirmilik National Park of
Canada (SIR-2018-28021).

In 2019, the 13 tracked foxes consisted of six neighbouring pairs plus one individual, and their
home ranges formed our 2019 study area (Figures 1A and 1B). In 2022, the 10 tracked foxes
included one pair for which both individuals were tracked, and eight pairs for which only one
individual of the pair was tracked, and their home ranges formed our 2022 study area (Figures 1C
and 1D). We determined reproductive status of individuals based on whether automated cameras
recorded cubs at individuals' den (Cameron et al., 2011). Daily observations and automated
cameras at fox dens showed that we tracked all foxes using our 2019 study area, whereas our 2022
study area was only used by tracked foxes and their pair mates. We collected one GPS location
every 4 minutes and one 30-sec burst of accelerometry (50 Hz) every 4.5 minutes in 2019 and
every 4 minutes in 2022 (Clermont et al., 2021b). We divided the 30-sec bursts of accelerometry
into ten 3-sec sequences and, using behavioural observations and a random forest algorithm, we

assigned to each sequence one of four behaviours: running, walking, digging and motionless (see

analytical details in Clermont et al., 2021b). During the goose incubation period, digging behaviour

of foxes is mainly associated with goose egg caching, but could also reflect caching or handling

of any prey as well as cache recoveries (Clermont et al., 2021b).

Figure 1. GPS locations (dots) of male and female Arctic foxes tracked from 11 June to 4 July
2019 (n = 67,109, panels A and B) and from 19 June to 12 July 2022 (n = 47,631, panels C and D)

at a 4-min fix interval on Bylot Island, Nunavut, Canada. Solid (males) and dashed (females) lines are 95% home range contours obtained through auto-correlated kernel density estimation. For each year, the same colours are used for members of a given fox pair. Home ranges are located near the coast of Bylot island shown by the thick grey line.

Data analysis

   1) Estimation of homes ranges

To define Arctic fox summer home ranges, we used GPS data collected from 11 June to 4 July 2019, and 19 June to 12 July 2022, corresponding to goose egg incubation periods. Prior to home range estimation, we confirmed home range residency for each individual using variogram analysis (Fleming et al., 2014), and excluded location data resulting from extra-territorial trips as it affects home range estimation (Calabrese et al., 2016). Indeed, during the study period, some individuals made a few extra-territorial trips going up to 25 km away from the center of the home range. These "outliers" were identified by evaluating distances between points and home range core, and a cut-off distance was determined visually for each individual (see Appendix A for further details). We excluded 11%, 9%, 6% and 4% of datapoints for four individuals that performed extra-territorial trips (three individuals in 2019 and one in 2022). The remaining individuals showed high range residency and we excluded < 1% of their GPS locations.

We then fitted range-resident continuous-time movement models to the data of each individual (model selection resulted in the Ornstein-Uhlenbeck Foraging (OUF) process used for each individual), to control for autocorrelation in both speed and location (Fleming et al., 2014; Calabrese et al., 2016). Then, we estimated home range areas using auto-correlated kernel density estimation (AKDE; Fleming et al., 2015). AKDE is more accurate than other home range estimators for autocorrelated location data (Noonan et al., 2019). The complete workflow we used is detailed in Silva et al. (2021), with R scripts provided, and was performed using the package ctmm (version 0.6.2, Calabrese et al., 2016) in R 4.1.0 (R Development Team, 2021).

   2) Estimation of probability of encounter among neighbours

Following Noonan et al. (2021), we then estimated the conditional distribution of encounters (CDE) from tracking data and AKDE home range estimates for each year separately (package ctmm v0.6.2). In short, the CDE estimates the spatial distribution of the probability of encounter between individuals based on their occurrence distribution. The probability of encounter estimated

by the CDE thus represents the probability that more than one individual uses a specific area, but
not necessarily at the same time. A CDE value of 1 at a given location indicates there is a 100%
chance that more than one individual use that location, and thus that the probability of encounter
is high, and a value of 0 that only one individual uses that location, meaning individuals should
never encounter each other at that location. Since we were interested in encounters among
neighbours, we excluded interactions between mates within their shared home range. We then
determined the degree of territoriality of each fox (objective 1) through a visual analysis of the
distribution of the probability of encounter within each home range. A fox was considered
territorial if at least some part of its home range was outside of the 95% CDE (Noonan et al., 2021),
excluding areas within 500 m of boundaries of study areas. Indeed, some interactions inevitably
occurred at home range boundaries located at the edge of the 2019 and 2022 study areas (but not
near the coastline, Figures 1 and 2), where neighbouring foxes were not tracked, leading to a local
underestimation of CDE values. Using a 1 km buffer instead of a 500 m one would have led to the
same classification (territorial yes/no) of all individuals but one in 2022 (female BJOV; see
Appendix C).

3) Arctic fox foraging behaviour: prey searching and prey caching

The Arctic fox is a cursorial, active hunting predator for which most active periods are associated
with prey searching (Schmitz et al., 2004; Poulin et al., 2021). We therefore determined whether
foxes were searching for prey by assigning to each GPS location an active or inactive state. To do
so we fitted a hidden Markov model (HMM) using the R package moveHMM (version 1.7,
Michelot et al., 2016). The active state is characterised by long step lengths and small turning
angles, and the inactive state by short step lengths and large turning angles. HMM construction
and model selection are detailed in Grenier-Potvin et al. (2021). HMM behavioural states were
well correlated with the accelerometry behavioural classification, where the inactive state was
almost exclusively associated to motionless sequences (Appendix B). The active state was
associated to various behaviours, the most common one being running (Appendix B).

We also used our accelerometry behavioural classification to determine whether individuals
engaged in digging (indicating prey caching or cache recovery events) during each 30-sec
accelerometry burst. At least one out of the ten 3-sec accelerometry sequences had to be assigned
to the digging behaviour to consider that the fox engaged in digging in that burst. Each 30-sec

burst was then associated to the closest GPS location, provided the time stamp of the GPS location

was within 30 sec of the start or end of the burst. Bursts occurring at less than 50 m of a den were

excluded as digging may then reflect den maintenance rather than prey caching.

We excluded from analyses all GPS locations and accelerometry bursts occurring within 500 m of

the boundaries of study areas, since interactions with unknown foxes may then occur (Appendix

C). To verify whether buffer size influences results, we 1) repeated our analyses by excluding the

individual of 2022 that would have been considered as non-territorial when using a 1 km buffer

(female BJOV), and 2) repeated our analyses by excluding locations falling within a 1 km buffer

instead of a 500 m one (Appendix C).

4) Effect of the probability of encountering a neighbour on fox behaviours

To test the influence of the likelihood of encountering a neighbour on the probability of engaging

in foraging behaviours (objective 2), we used generalized additive mixed models (GAMM) with a

binomial error distribution and logit link function ("gam" function of the R package mgcv, v1.9-

0, Wood, 2017) and modelled fox probability of being active (0 = inactive, 1 = active; P1) and of

engaging in digging (0 = no digging, 1= digging; P2) with respect to the probability of encounter.

As not all foxes experienced the same range of probability of encounter, we centered the

probability of encounter values to a mean of 0 within each fox home range. We modeled 2019 and

2022 data separately, which resulted in four models. In all models, we included as fixed effect a

categorical variable representing whether the fox was territorial (yes/no), which we included in

interaction with the probability of encounter, to test whether the degree of territoriality modulated

the effect of the probability of encounter on the probability of engaging in foraging behaviours

(P3). In 2019, we also controlled for nesting goose density (individual geese/ha), a proxy for goose

nest density estimated from detailed field surveys (Grenier-Potvin et al., 2021). Indeed, nesting

goose density was previously shown to affect fox habitat selection during active and inactive

periods (Grenier-Potvin et al., 2021) and fox probability to dig (Clermont et al., 2021b). We did

not perform detailed field surveys in 2022 as nest density was very low throughout the study area,

therefore we could not include nesting goose density as a covariate in the 2022 models. We

included individual sex as another fixed effect in all models, as well as fox identity and Julian date

as random intercepts. To address temporal autocorrelation in our datasets, we also added to all

models a cyclic cubic spline function of the numeric time of the day and a thin plate regression

spline function of the Julian date, with 10 basis dimensions for each function (Pedersen et al.,
2019). All continuous covariates were centered and standardised to facilitate the interpretation of
model estimates (Schielzeth, 2010). We verified model assumptions and independence of residuals
using the R package DHARMa (version 0.4.3, Hartig, 2021), and observed significant deviations
in all models' residuals. We solved these issues by using a subset of data that included 150
randomly chosen observations by individual for all models, and by keeping all covariates in the
models even when effects were not significant. Results are expressed using evidence-based
language (Muff et al., 2021) where p-value ≥ 0.1 suggests no evidence, p-value =]0.05, 0.1]
suggests weak evidence, p-value =]0.01, 0.05] suggests moderate evidence, p-value =]0.01,
0.001] suggests strong evidence and p-value < 0.001 suggests very strong evidence.

Results

Conditional distribution of encounters in fox home ranges

Home range areas averaged 7.93 ± 3.15 (SD) km^2 (Figure 1, n = 23). We found no evidence that
home range areas differed between sexes (t-test: t = 0.53, p = 0.600, average male home range area
= 8.27 ± 3.12 (SD) km^2, n = 12, average female home range area = 7.56 ± 3.28 (SD) km^2, n = 11).
However, there was weak evidence that home ranges were smaller in 2022 than in 2019 (t-test: t
= 2.07, p = 0.051, average home range area in 2019 = 9.03 ± 3.12 (SD) km^2, n = 13, average home
range area in 2022 = 6.51 ± 2.70 (SD) km^2, n = 10).

As expected, higher CDE values occur where home ranges of tracked neighbours overlap (Figure
2). The CDE differs among home ranges. In 2019, three individuals (green female RMJJ, green
male BORR and grey male BBJO) have all of their home range located within the 95% CDE
contour if we ignore 500 m study area edges, thus having no part of their home range with a null
probability of encountering a tracked neighbour (Figures 2A and 3A). As such, these three
individuals are considered as non-territorial. Furthermore, these three foxes did not reproduce in
2019. By contrast, all other ten individuals are territorial, having a considerable area of their home
range outside of the 95% CDE contour, where the probability of encounter with a neighbour is
null (Figures 2A and 3A). These ten individuals reproduced in 2019. In 2022, all individuals but
one (dark orange male BOBB) are territorial as they had an area of their home range located outside

of the 95% CDE contour where the probability of encounter with neighbours is null, ignoring 500

m study area edges (Figures 2B and 3B). None of the foxes reproduced in 2022.

Figure 2. Conditional distribution of encounters (CDE) among A) 13 Arctic fox neighbours in
2019 and B) ten Arctic fox neighbours in 2022, tracked with GPS, excluding interactions between
pair members. The thick black line is the 95% CDE contour and coloured lines are 95% AKDE
contours. The grey shaded area represents the 500 m inner buffer of study area boundaries where
interactions with other foxes inevitably occur. Males and females are represented by solid and
dashed lines, respectively, and members of a given pair are identified by the same colour within
each year.

Figure 3. CDE values (probability of encountering a neighbour) observed at each GPS location from Figure 1, for each fox in A) 2019 and B) 2022. Fox colour IDs reflect the unique combinations of colours of the four ear tags (e.g., OBBB: orange-blue-blue-blue). Boxplot colours used for each individual are the same as in Figures 1 and 2. Males and females are represented by solid and dashed lines, respectively, and each year's members of a given pair are identified by the same colour. Boxplots show first quartile, median, and third quartile. Lower and upper whiskers extend, respectively, to the lowest and highest values within the interquartile range multiplied by 1.5. Points represent values outside this range.

Effect of the probability of encountering a neighbour on fox behaviour

Non-territorial foxes of 2019 are more likely to be active (very strong evidence) and to engage in digging (strong evidence) where they are less likely to encounter a neighbour, thus supporting P1 and P2 (Table 1; Figure 4). However, we found no evidence of an effect of the probability of encounter on the behaviours of the only non-territorial individual of 2022 (Table 1; Figure 4). On the other hand, territorial foxes of 2019 and 2022 are more likely to be active where they are more likely to encounter a neighbour, with weak evidence for 2019 and moderate evidence for 2022, thus contradicting P1 (Table 1). However, when excluding individual BJOV in the 2022 model, or when excluding all locations falling within a 1 km inner buffer of study area boundaries, we find no evidence for an effect of the probability of encounter on the probability of being active (Appendix C). Furthermore, we found no evidence that the propensity to dig of territorial foxes (both years) varies according to the probability of encountering a neighbour, contradicting P2

(Table 1; Figure 4). In 2019, when resources were abundant, the probability of encountering a
neighbour more strongly affected the behaviours of non-territorial individuals compared to
territorial ones, but differences were minor or absent in 2022 when resources were scarce (Figure
4), providing partial support for P3. We also found very strong evidence that nesting goose density
has a positive effect on the level of activity and propensity to dig in 2019, but no evidence of an
effect of sex in both years (Table 1). Rerunning models on different subsets of the dataset sightly
changes coefficient values, but not the direction and significance of effects.

Table 1. Estimates of generalized additive mixed models predicting the probability of fox being
active in A) 2019 (n = 1950 GPS locations) and B) 2022 (n = 1500 GPS locations), and the
probability of fox digging in C) 2019 (n = 1950 accelerometry bursts) and D) 2022 (n = 1500
accelerometry bursts). All models had a binomial error distribution with a logit-link function. We
included as parametric terms (fixed effects) the probability of encountering a neighbour, whether
the fox was territorial, an interaction between the probability of encountering a neighbour and
whether the fox was territorial, the individual's sex and goose nest density (2019 only). We
included splines for numeric time of day (ToD) and Julian date (day), and the random effects fox
identity (Fox ID) and Julian date (day_factor).

A) Probability of fox being active in 2019

Component	Term	Estimate	Std.Error	z.value	p.value
1. Parametric coefficients	(Intercept)	0.13	0.18	0.71	0.477
	Probability of encountering a neighbour	0.12	0.07	1.72	0.086
	Territorial yes/no [no]	0.12	0.29	0.40	0.692
	Goose nest density	0.52	0.06	8.37	<0.001
	Sex [male]	-0.05	0.23	-0.20	0.840
	Prob encounter: Territorial yes/no [no]	-1.12	0.14	-7.84	<0.001

Component	Term	edf	Ref.df	Chi.sq	p.value
2. Smooth terms	s(ToD)	5.36	8.00	168.20	<0.001
	s(day)	1.00	1.00	0.09	0.771
	s(Fox ID)	7.96	10.00	39.45	<0.001
	s(day_factor)	2.70	21.00	3.21	0.265

B) Probability of fox being active in 2022

Component	Term	Estimate	Std.Error	z.value	p.value
1. Parametric coefficients	(Intercept)	0.65	0.10	6.32	<0.001
	Probability of encountering a neighbour	0.14	0.06	2.12	0.034
	Territorial yes/no [no]	-0.37	0.21	-1.81	0.070
	Sex [male]	0.00	0.13	-0.01	0.994
	Prob encounter: Territorial yes/no [no]	-0.22	0.18	-1.23	0.219

Component	Term	edf	Ref.df	Chi.sq	p.value
2. Smooth terms	s(ToD)	6.03	8.00	211.18	<0.001
	s(day)	1.00	1.00	0.20	0.658
	s(Fox ID)	0.01	7.00	0.01	0.465
	s(day_factor)	9.95	22.00	18.96	0.007

C) Probability of fox digging in 2019

Component	Term	Estimate	Std.Error	z.value	p.value
1. Parametric coefficients	(Intercept)	-0.91	0.30	-3.03	0.002
	Probability of encountering a neighbour	0.07	0.07	0.96	0.336
	Territorial yes/no [no]	-0.03	0.48	-0.07	0.946
	Goose nest density	0.37	0.06	6.17	<0.001
	Sex [male]	-0.09	0.40	-0.22	0.826
	Prob encounter: Territorial yes/no [no]	-0.45	0.15	-3.06	0.002

Component	Term	edf	Ref.df	Chi.sq	p.value
2. Smooth terms	s(ToD)	3.42	8.00	42.88	<0.001
	s(day)	2.76	3.46	6.43	0.159
	s(Fox ID)	9.15	10.00	104.81	<0.001
	s(day_factor)	0.22	21.00	0.22	0.404

D) Probability of fox digging in 2022

Component	Term	Estimate	Std.Error	z.value	p.value
1. Parametric coefficients	(Intercept)	-0.37	0.23	-1.58	0.115
	Probability of encountering a neighbour	0.043	0.06	0.72	0.471
	Territorial yes/no [no]	0.346	0.57	0.60	0.546
	Sex [male]	-0.03	0.34	-0.10	0.920
	Prob encounter: Territorial yes/no [no]	-0.25	0.17	-1.48	0.138

Component	Term	edf	Ref.df	Chi.sq	p.value
2. Smooth terms	s(ToD)	4.20	8.00	84.67	<0.001
	s(day)	1.00	1.00	0.52	0.473
	s(Fox ID)	6.14	7.00	49.01	<0.001
	s(day_factor)	5.48	22.00	9.21	0.045

Figure 4. Predicted effects of the probability of encountering a neighbour on probability of fox being active (0 = inactive, 1 = active) in A) 2019 (n = 1950 GPS locations) and B) 2022 (n = 1500 GPS locations) and on probability of fox digging (0 = no digging, 1= digging) in C) 2019 (n = 1950 accelerometry bursts) and D) 2022 (n = 1500 accelerometry bursts). The probability of encountering a neighbour was obtained from conditional distributions of encounters (Figure 2). Orange dots and lines refer to territorial individuals (n = 10 foxes in 2019 and 9 foxes in 2022), and blue dots and lines refer to non-territorial individuals (n = 3 foxes in 2019 and 1 fox in 2022). Shaded areas represent 95% confidence intervals.

Discussion

Using high throughput tracking technologies, we estimated the spatial distribution of the probability of encounter among Arctic foxes using neighbouring home ranges. We benefited from a natural situation where most, but not all individuals were territorial, thus enabling us to evaluate the effect of territoriality on the spatial distribution of foraging behaviour. We also used data from two contrasted years during which resource abundance varied from high (2019) to low (2022). Non-territorial foxes had some probability of encountering neighbours throughout their entire home range. Non-territorial individuals of 2019 responded to the variation in the probability of encountering a neighbour by shifting the distribution of prey searching and caching away from the places where encounters were most likely to occur. The only non-territorial individual of 2022 however did not respond to variation in the probability of encounter in its home range. Territorial individuals of 2019 and 2022 did the opposite by being slightly more likely to be active where they had a higher probability of encountering a neighbour (however strength of evidence varied), but the latter did not influence their tendency to cache prey.

Despite modern technology, it remains highly challenging to document the movements and foraging behaviours of many neighbouring vertebrate predators continuously and with high precision and frequency, in a natural setting over several weeks. Our sample of 13 individuals in 2019 and 10 individuals in 2022, each tracked for more than three weeks, is therefore of high value. Our study provides valuable insights into how intraspecific interactions may influence the spatial distribution of predator foraging. Specifically, our results suggest that in predators characterized by territoriality, when resources are abundant individuals that do not establish a territory strongly adjust their foraging behaviours based on the probability of encountering neighbours, which could represent an alternative tactic to secure resources.

The degree of territoriality varies among foxes

Variation in the distribution, abundance and predictability of resources often explains among-individual differences in territoriality (Maher & Lott, 2000; Sells & Mitchell, 2020). For example, Eide et al. (2004) observed that in Arctic foxes of Svalbard, high home range overlap occurred near the coast where seabird colonies are concentrated. On the other hand, little home range overlap occurred inland where reindeer (*Rangifer tarandus*) carcasses are scattered and

unpredictable. In our study, we do not suspect that variation in prey distribution, abundance or
predictability explains among-individual differences in territoriality within years, as all studied
foxes lived near the center of the goose colony where nest abundance is usually high and habitats
favourable to lemmings. However, we used two contrasted years of data, with one year of moderate
lemming abundance and typical goose density for this study area (2019), and one year of low
lemming abundance and low goose density (2022). In 2019, although resource abundance was
high, not all foxes reproduced and used a territory. This suggests that using a territory may be
necessary to secure enough resources to ensure the survival of young, even when food is abundant
(Webb et al., 2012; López-Bao et al., 2014). On the other hand, for non-breeding individuals, the
costs of defending a territory may have been too high compared to the benefits when resources are
abundant, as predicted by Maher and Lott's (2000) cost-benefit model of territoriality.

Although resources were scarce in 2022, all foxes but one had exclusive use (apart from pair
members) in some part of their home range, which seems to contradict Maher and Lott's (2000)
model, that predicts no territoriality when resources are very rare. This could however indicate
food abundance inside the colony in 2022 was still not low enough for territoriality not to be
beneficial. In fact, foxes in the colony may have relied on stored eggs from the previous year
(Samelius et al., 2007), which could have attenuated the effects of low resources. However,
although high abundance of resources may permit non-breeding individuals to forego being
territorial, defending a territory may remain beneficial under low or very low resource abundance,
especially when the abundance of resources varies cyclically through time. Similarly, López-Bao
et al. (2014) observed that home range overlap among Iberian lynx *(Lynx pardinus)* remains low
across a gradient of prey availability. This obstinate strategy (sensu von Schantz, 1984), i.e.,
maintaining a territory even when resources are rare, could be beneficial if it allows to secure a
home range in areas where resources are usually high, such as the snow goose colony on Bylot
Island. Indeed, having access to the colony allows for fox reproduction when lemming density is
low (Giroux et al., 2012; Chevallier et al., 2020). Therefore, it may remain beneficial to defend a
territory even during episodes of low food availability, to maximise chances of future reproduction
(López-Bao et al., 2014). Although results from 2019 suggest that pair members have very similar
home range boundaries, we did not monitor the activity of both pair members in 2022 (except for
one pair), which may have affected our ability to assess spatial variation in the probability of

encountering a neighbour. Our interpretations should be confirmed by monitoring the activity of
more neighbouring individuals.

Variable effect of the probability of encountering a neighbour on Arctic fox foraging behaviours

In 2019, being territorial or not determined how Arctic foxes responded to variation in the
probability of encountering a neighbour. The three non-territorial individuals were less active in
areas where the probability of encounter with neighbours was highest. As they had no or low
exclusive use in their home range, those Arctic foxes may have increased foraging efficiency by
searching prey more intensely where the risk of encounter with other foxes was the lowest, where
they do not need to be as vigilant. Indeed, inactive periods indicates that the fox was stationary,
which might reflect periods of resting or vigilance. A hypothesis is thus that non-territorial foxes
were more vigilant when using areas with a high probability of encounter, to avoid interactions
with dominant neighbours, as observed with white-faced capuchins (Tórrez-Herrera et al., 2020;
Noonan et al., 2021). In 2022, we found no effect of the probability of encounter on the activity of
the only non-territorial individual, suggesting that when resources are scarce, non-territorial foxes
may not afford to search prey only in specific areas of the landscape, but this result should be
confirmed with a larger sample of individuals.

Territorial individuals of 2019 and 2022 however were slightly more likely to be active where the
probability of encounter with neighbours was highest. However, the evidence was weak in 2019,
moderate in 2022, and absent when excluding individual BJOV in 2022 and when excluding
locations falling within a 1 km buffer of study area boundaries. There are therefore some
uncertainties in these findings and the following interpretations. As most active periods should
reflect prey searching in Arctic foxes, being more active where the probability of encountering a
neighbour is high could indicate that territorial individuals search prey more intensely where the
risk of encounter with other foxes is the highest. However, probability of encounter did not
influence their probability of caching prey (see next paragraph). Therefore, as most captured prey
are also cached (Careau et al., 2007), we believe that higher probability of being active near home
range boundaries for territorial individuals is more likely to reflect an increase in territory
patrolling or scent-marking, rather than prey searching. Indeed, locations of territorial boundaries
are not fixed in space and time and can for example vary according to the movements of animals
and the longevity of olfactory signals (Giuggioli et al., 2011). Therefore, even though territorial

individuals should overall avoid home range boundaries if encounters with other individuals are
risky, they should still revisit boundaries to refresh olfactory signals to maintain their exclusive
access (Schlägel et al., 2017).

Furthermore, greater probability of encounter with neighbours may generate a greater risk of cache
pilfering (Samelius & Alisauskas, 2000; van der Veen et al., 2020). Therefore, Arctic foxes should
be more likely to cache prey where they are less likely to encounter neighbours. As such we found
that non-territorial foxes of 2019 had a higher probability to dig where the probability of
encountering neighbours was lower. Although during the goose incubation period, digging events
should mostly reflect prey caching, we cannot differentiate caching events from prey captures,
cache recoveries and recaching events (Clermont et al., 2021b). Therefore, our results may also
indicate that non-territorial individuals could favour recaching away from their neighbours, in safer
sites. Caching and recaching prey in "out of view" sites are in fact common cache protection
strategies in other species such as corvids (Dally et al., 2006). Careau et al. (2007) also showed
that Arctic fox recaches of goose eggs were made further from the nest compared to initial caches,
potentially to secure eggs in sites where chances of pilfering are lower (e.g., closer to the den). All
other individuals however cached prey independently of their probability of encountering
neighbours. This is likely because territorial individuals have no benefit of modulating the
distribution of their caches according to potential encounters in an area that is in part used
exclusively, and under low resource abundance, non-territorial individuals may not afford to adjust
foraging behaviours to the probability of encounter.

In addition to spatial adjustments of behaviours, Arctic foxes may modify their foraging
behaviours temporally to avoid encounters, which the CDE did not allow to evaluate (Noonan et
al., 2021). Similarly, prey individuals use areas with abundant resources but high-predation risk
only at times of the day when predators are less active (Kohl et al., 2018; Smith et al., 2019b).
Future studies should incorporate time to their estimation of the probability of encounter to help
uncover the different tactics territorial animals use to avoid intraspecific interactions while
foraging.

Implications for predator-prey interactions

Arctic foxes have important top-down effects in the tundra and generate multiple predator-mediated interactions among prey (Bêty et al., 2002; Legagneux et al., 2012; Duchesne et al., 2021). For example, nest survival of American golden plover (*Pluvialis dominica*) is lower within the snow goose colony, where foxes use smaller home ranges and are thus present at higher densities (Dulude-de Broin et al., 2023). At fine spatial scales, Arctic fox movements generate a predation risk landscape influencing the behaviour and nest distribution of several migratory birds (Clermont et al., 2021a). It is therefore of strong interest to identify the factors explaining Arctic fox density and where they choose to forage. In this study, we observed that the degree of territoriality and food abundance influence fox response to intraspecific interactions. When resources are abundant, non-territorial individuals show the highest behavioural response by avoiding encounters with neighbours when prey searching and caching. However, the effect of the probability of encounter on the foraging behaviours of territorial individuals (i.e., most individuals), was either weak or absent, suggesting intraspecific interactions may not have a strong effect on the spatial distribution of predation risk. Interestingly, although most Arctic foxes are territorial, we observed no unused areas between home ranges, which overlap (Figures 1 and 2). As such, the spatial configuration of fox home ranges does not lead to buffer zones of low predation risk between territories as observed in some wolf-ungulate systems (Mech, 1977; Anderson et al., 2005). Conversely, the presence of non-territorial individuals, whose home ranges overlap those of their territorial neighbours, creates zones where fox density is high, which may increase the risk of predation (Clermont et al., 2021a; Dulude-de Broin et al., 2023). As such, we hypothesise that the greatest risk of predation should be located within the home range of non-territorial individuals, while the lowest predation risk should be found in defended territories where the probability of encounter among neighbouring foxes is null.

Conclusion

We found that not all Arctic foxes on Bylot Island used a territory. We highlight that because costs and benefits of territoriality may differ among individuals in a population, alternative behavioural tactics may emerge from non-territorial individuals to secure resources. Including the conditional distribution of encounters into habitat selection analyses would further help to better understand

how different individuals deal with both the physical and social environments when using their habitat, and how they compromise between the needs to acquire valuable resources and avoid competitors. Finally, a spatial configuration in which the home ranges of non-territorial individual predators overlap those of their territorial neighbours may influence the distribution of predation risk, with could modulate the distribution and behaviour of prey and the structure of communities.

Data Accessibility Statement

Arctic fox GPS and accelerometry data are available through the Movebank Data Repository (Movebank Study ID 1241071371 and 2602389249).

Competing Interests Statement

The authors declare no competing interests.

Author Contributions

All authors conceived the study and collected the data. JC performed the analyses with help from FDB and MPP. JC wrote the first version of the manuscript, which all authors commented and revised. All authors approved the final version.

Acknowledgements

We thank A. Grenier-Potvin, R. Gravel, G. Roy, L.-P. Ouellet, M. Beaudoin and M. Archambault for field work, M. Noonan for advice on spatial analyses, and the community of Mittimatalik for support. Funding was received from (alphabetical order) Canada Foundation for Innovation, Canada Research Chairs Program, Fonds de Recherche du Québec – Nature et technologies (FRQNT), Natural Sciences and Engineering Research Council of Canada (NSERC), Network of Centers of Excellence of Canada ArcticNet, Northern Scientific Training Program (Polar Knowledge Canada), Parks Canada Agency, Polar Continental Shelf Program (Natural Resources Canada), and Weston Family Foundation.

Literature cited

Anderson, D.P., Turner, M.G., Forester, J.D., Zhu, J., Boyce, M.S., Beyer, H., Stowell, L. 2005. Scale-dependent summer resource selection by reintroduced elk in Wisconsin, USA. Journal of Wildlife Management 69, 298–310.

Arnold, J., Soulsbury, C.D., Harris, S. 2011. Spatial and behavioral changes by red foxes (*Vulpes vulpes*) in response to artificial territory intrusion. Canadian Journal of Zoology 89, 808–815.

Bêty, J., Gauthier, G., Giroux, J.-F., Korpimäki, E. 2001. Are goose nesting success and lemming cycles linked? Interplay between nest density and predators. Oikos 93, 388–400.

Bêty, J., Gauthier, G., Korpimaki, E., Giroux, J.-F. 2002. Shared predators and indirect trophic interactions: lemming cycles and Arctic-nesting geese. Journal of Animal Ecology 71, 88–98.

Bidder, O.R., Virgilio, A. di, Hunter, J.S., McInturff, A., Gaynor, K.M., Smith, A.M., Dorcy, J., Rosell, F. 2020. Monitoring canid scent marking in space and time using a biologging and machine learning approach. Scientific Reports 10, 588.

Brown, J.L., Orians, G.H. 1970. Spacing patterns in mobile animals. Annual Review of Ecology and Systematics 1, 239–262.

Calabrese, J.M., Fleming, C.H., Gurarie, E. 2016. ctmm: an R package for analyzing animal relocation data as a continuous-time stochastic process. Methods in Ecology and Evolution 7, 1124–1132.

Cameron, C., Berteaux, D., Dufresne, F. 2011. Spatial variation in food availability predicts extrapair paternity in the arctic fox. Behavioral Ecology 22, 1364–1373.

Careau, V., Giroux, J.-F., Berteaux, D. 2007. Cache and carry: hoarding behavior of Arctic fox. Behavioral Ecology and Sociobiology 62, 87–96.

Careau, V., Lecomte, N., Bêty, J., Giroux, J.-F., Gauthier, G., Berteaux, D. 2008. Hoarding of pulsed resources: temporal variations in egg-caching by arctic fox. Écoscience 15, 268–276.

Chevallier, C., Gauthier, G., Lai, S., Berteaux, D. 2020. Pulsed food resources affect reproduction but not adult apparent survival in arctic foxes. Oecologia 193, 557–569.

Clermont, J., Grenier-Potvin, A., Duchesne, É., Couchoux, C., Dulude-de Broin, F., Beardsell, A., Bêty, J., Berteaux, D. 2021a. The predator activity landscape predicts the anti-predator behavior and distribution of prey in a tundra community. Ecosphere 12, e03858.

Clermont, J., Woodward-Gagné, S., Berteaux, D. 2021b. Digging into the behaviour of an active hunting predator: arctic fox prey caching events revealed by accelerometry. Movement Ecology 9, 58.

Cresswell, W. 2008. Non-lethal effects of predation in birds. Ibis 150, 3–17.

Dally, J.M., Clayton, N.S., Emery, N.J. 2006. The behaviour and evolution of cache protection and pilferage. Animal Behaviour 72, 13–23.

Duchesne, É., Lamarre, J.-F., Gauthier, G., Berteaux, D., Gravel, D., Bêty, J. 2021. Variable strength of predator-mediated effects on species occurrence in an arctic terrestrial vertebrate community. Ecography 44, 1236–1248.

Dulude-de Broin, F., Clermont, J., Beardsell, A., Ouellet, L.-P., Legagneux, P., Bêty, J., Berteaux, D. 2023. Predator home range size mediates indirect interactions between prey species in an arctic vertebrate community. Journal of Animal Ecology 92, 2373–2385.

Eberhardt, L.E., Hanson, W.C., Bengtson, J.L., Garrott, R.A., Hanson, E.E. 1982. Arctic fox
home range characteristics in an oil-development area. The Journal of Wildlife
Management 46, 183.
Eide, N.E., Jepsen, J.U., Prestrud, P. 2004. Spatial organization of reproductive Arctic foxes
Alopex lagopus: responses to changes in spatial and temporal availability of prey. Journal
of Animal Ecology 73, 1056–1068.
Fauteux, D., Gauthier, G., Berteaux, D. 2015. Seasonal demography of a cyclic lemming
population in the Canadian Arctic. Journal of Animal Ecology 84, 1412–1422.
Fawcett, J.K., Fawcett, J.M., Soulsbury, C.D. 2013. Seasonal and sex differences in urine
marking rates of wild red foxes Vulpes vulpes. Journal of Ethology 31, 41–47.
Fleming, C.H., Calabrese, J.M., Mueller, T., Olson, K.A., Leimgruber, P., Fagan, W.F. 2014.
From fine-scale foraging to home ranges: a semivariance approach to identifying
movement modes across spatiotemporal scales. The American Naturalist 183, E154–
E167.
Fleming, C.H., Fagan, W.F., Mueller, T., Olson, K.A., Leimgruber, P., Calabrese, J.M. 2015.
Rigorous home range estimation with movement data: a new autocorrelated kernel
density estimator. Ecology 96, 1182–1188.
Gaynor, K.M., Brown, J.S., Middleton, A.D., Power, M.E., Brashares, J.S. 2019. Landscapes of
fear: spatial patterns of risk perception and response. Trends in Ecology & Evolution 34,
355–368.
Giroux, M.-A., Berteaux, D., Lecomte, N., Gauthier, G., Szor, G., Bêty, J. 2012. Benefiting from
a migratory prey: spatio-temporal patterns in allochthonous subsidization of an arctic
predator. Journal of Animal Ecology 81, 533–542.
Giuggioli, L., Potts, J.R., Harris, S. 2011. Animal interactions and the emergence of territoriality.
PLoS ONE 7, e1002008.
Grenier-Potvin, A., Clermont, J., Gauthier, G., Berteaux, D. 2021. Prey and habitat distribution
are not enough to explain predator habitat selection: addressing intraspecific interactions,
behavioural state and time. Movement Ecology 9, 12.
Gruyer, N., Gauthier, G., Berteaux, D. 2008. Cyclic dynamics of sympatric lemming populations
on Bylot Island, Nunavut, Canada. Canadian Journal of Zoology 86, 910–917.
Hartig, F. 2021. DHARMa: residual diagnostics for hierarchical (multilevel/mixed) regression
models. https://CRAN.R-project.org/ package=DHARMa.
Hinsch, M., Komdeur, J. 2017. What do territory owners defend against? Proceedings:
Biological Sciences 284, 1–8.
Humphries, D.J., Nelson-Flower, M.J., Bell, M.B.V., Finch, F.M., Ridley, A.R. 2021. Kinship,
dear enemies, and costly combat: the effects of relatedness on territorial overlap and
aggression in a cooperative breeder. Ecology and Evolution 11, 17031–17042.
Kohl, M.T., Stahler, D.R., Metz, M.C., Forester, J.D., Kauffman, M.J., Varley, N., White, P.J.,
Smith, D.W., MacNulty, D.R. 2018. Diel predator activity drives a dynamic landscape of
fear. Ecological Monographs 88, 638–652.
Legagneux, P., Gauthier, G., Berteaux, D., Bêty, J., Cadieux, M.-C., Bilodeau, F., Bolduc, E.,
McKinnon, L., Tarroux, A., Therrien, J.-F., Morissette, L., Krebs, C.J. 2012.
Disentangling trophic relationships in a High Arctic tundra ecosystem through food web
modeling. Ecology 93, 1707–1716.
Lewis, M.A., Murray, J.D. 1993. Modelling territoriality and wolf–deer interactions. Nature 366,
738–740.

Lima, S.L. 2002. Putting predators back into behavioral predator–prey interactions. Trends in Ecology & Evolution 17, 70–75.

López-Bao, J.V., Rodríguez, A., Delibes, M., Fedriani, J.M., Calzada, J., Ferreras, P., Palomares, F. 2014. Revisiting food-based models of territoriality in solitary predators. Journal of Animal Ecology 83, 934–942.

Macdonald, D.W., Courtenay, O., Forbes, S., Mathews, F. 1999. The red fox (Vulpes vulpes) in Saudi Arabia: loose-knit groupings in the absence of territoriality. Journal of Zoology 249, 383–391.

Maher, C.R., Lott, D.F. 2000. A review of ecological determinants of territoriality within vertebrate species 143, 1–29.

McLoughlin, P.D., Ferguson, S.H., Messier, F. 2000. Intraspecific variation in home range overlap with habitat quality: a comparison among brown bear populations. Evolutionary Ecology 14, 39–60.

Mech, L.D. 1977. Wolf-pack buffer zones as prey reservoirs. Science 198, 320–321.

Michelot, T., Langrock, R., Patterson, T.A. 2016. moveHMM: an R package for the statistical modelling of animal movement data using hidden Markov models. Methods in Ecology and Evolution 7, 1308–1315.

Muff, S., Nilsen, E.B., O'Hara, R.B., Nater, C.R. 2021. Rewriting results sections in the language of evidence. Trends in Ecology & Evolution 37, 203–210.

Nathan, R., Spiegel, O., Fortmann-Roe, S., Harel, R., Wikelski, M., Getz, W.M. 2012. Using tri-axial acceleration data to identify behavioral modes of free-ranging animals: general concepts and tools illustrated for griffon vultures. Journal of Experimental Biology 215, 986–996.

Noonan, M.J., Martinez-Garcia, R., Davis, G.H., Crofoot, M.C., Kays, R., Hirsch, B.T., Caillaud, D., Payne, E., Sih, A., Sinn, D.L., Spiegel, O., Fagan, W.F., Fleming, C.H., Calabrese, J.M. 2021. Estimating encounter location distributions from animal tracking data. Methods in Ecology and Evolution 12, 1158–1173.

Noonan, M.J., Tucker, M.A., Fleming, C.H., Akre, T.S., Alberts, S.C., Ali, A.H., Altmann, J., Antunes, P.C., Belant, J.L., Beyer, D., Blaum, N., Böhning-Gaese, K., Cullen, L., Paula, R.C., Dekker, J., Drescher-Lehman, J., Farwig, N., Fichtel, C., Fischer, C., Ford, A.T., Goheen, J.R., Janssen, R., Jeltsch, F., Kauffman, M., Kappeler, P.M., Koch, F., LaPoint, S., Markham, A.C., Medici, E.P., Morato, R.G., Nathan, R., Oliveira-Santos, L.G.R., Olson, K.A., Patterson, B.D., Paviolo, A., Ramalho, E.E., Rösner, S., Schabo, D.G., Selva, N., Sergiel, A., Xavier da Silva, M., Spiegel, O., Thompson, P., Ullmann, W., Zięba, F., Zwijacz-Kozica, T., Fagan, W.F., Mueller, T., Calabrese, J.M. 2019. A comprehensive analysis of autocorrelation and bias in home range estimation. Ecological Monographs 89, e01344.

Ord, T.J. 2021. Costs of territoriality: a review of hypotheses, meta-analysis, and field study. Oecologia 197, 615–631.

Patterson, T.A., Parton, A., Langrock, R., Blackwell, P.G., Thomas, L., King, R. 2017. Statistical modelling of individual animal movement: an overview of key methods and a discussion of practical challenges. AStA Advances in Statistical Analysis 101, 399–438.

Pedersen, E.J., Miller, D.L., Simpson, G.L., Ross, N. 2019. Hierarchical generalized additive models in ecology: an introduction with mgcv. PeerJ 7, e6876.

Persson, J., Wedholm, P., Segerström, P. 2010. Space use and territoriality of wolverines (Gulo gulo) in northern Scandinavia. European Journal of Wildlife Research 56, 49–57.

688   Poulin, M., Clermont, J., Berteaux, D. 2021. Extensive daily movement rates measured in
689   territorial arctic foxes. Ecology and Evolution 11, 2503–2514.
690   Powell, R.A. 2000. Animal home ranges and territories and home range estimators, in: Boitani,
691   L., Fuller, T.F. (Eds.), Research Techniques in Animal Ecology. Columbia University
692   Press, New York, NY, pp. 65–110.
693   R Development Team 2021. R: a language and environment for statistical computing. R
694   Foundation for Statistical Computing, Vienna.
695   Rosell, F., Thomsen, L.R. 2006. Sexual dimorphism in territorial scent marking by adult
696   Eurasian beavers (Castor fiber). Journal of Chemical Ecology 32, 1301–1315.
697   Samelius, G., Alisauskas, R.T. 2000. Foraging patterns of arctic foxes at a large arctic goose
698   colony. Arctic 53, 279–288.
699   Samelius, G., Alisauskas, R.T., Hobson, K.A., Larivière, S. 2007. Prolonging the arctic pulse:
700   long-term exploitation of cached eggs by arctic foxes when lemmings are scarce. Journal
701   of Animal Ecology 76, 873–880.
702   Schielzeth, H. 2010. Simple means to improve the interpretability of regression coefficients.
703   Methods in Ecology and Evolution 1, 103–113.
704   Schlägel, U.E., Merrill, E.H., Lewis, M.A. 2017. Territory surveillance and prey management:
705   wolves keep track of space and time. Ecology and Evolution 7, 8388–8405.
706   Schmitz, O.J., Krivan, V., Ovadia, O. 2004. Trophic cascades: the primacy of trait-mediated
707   indirect interactions. Ecology Letters 7, 153–163.
708   Sells, S.N., Mitchell, M.S. 2020. The economics of territory selection. Ecological Modelling 438,
709   109329.
710   Sih, A. 1984. The behavioral response race between predator and prey. The American Naturalist
711   123, 143–150.
712   Silk, J.B., Alberts, S.C., Altmann, J. 2003. Social bonds of female baboons enhance infant
713   survival. Science 302, 1231–1234.
714   Sillero-Zubiri, C., Macdonald, D.W. 1998. Scent-marking and territorial behaviour of Ethiopian
715   wolves Canis simensis. Journal of Zoology 245, 351–361.
716   Silva, I., Fleming, C.H., Noonan, M.J., Alston, J., Folta, C., Fagan, W.F., Calabrese, J.M. 2021.
717   Autocorrelation-informed home range estimation: A review and practical guide. Methods
718   in Ecology and Evolution 13, 534–544.
719   Smith, J.A., Donadio, E., Pauli, J.N., Sheriff, M.J., Bidder, O.R., Middleton, A.D. 2019a. Habitat
720   complexity mediates the predator–prey space race. Ecology 100, e02724.
721   Smith, J.A., Donadio, E., Pauli, J.N., Sheriff, M.J., Middleton, A.D. 2019b. Integrating temporal
722   refugia into landscapes of fear: prey exploit predator downtimes to forage in risky places.
723   Oecologia 189, 883–890.
724   Studd, E.K., Derbyshire, R.E., Menzies, A.K., Simms, J.F., Humphries, M.M., Murray, D.L.,
725   Boutin, S. 2021. The purr-fect catch: using accelerometers and audio recorders to
726   document kill rates and hunting behaviour of a small prey specialist. Methods in Ecology
727   and Evolution 12, 1277–1287.
728   Tórrez-Herrera, L.L., Davis, G.H., Crofoot, M.C. 2020. Do monkeys avoid areas of home range
729   overlap because they are dangerous? A test of the risk hypothesis in white-faced capuchin
730   monkeys (Cebus capucinus). International Journal of Primatology 41, 246–264.
731   van der Veen, B., Mattisson, J., Zimmermann, B., Odden, J., Persson, J. 2020. Refrigeration or
732   anti-theft? Food-caching behavior of wolverines (Gulo gulo) in Scandinavia. Behavioral
733   Ecology and Sociobiology 74, 52.

von Schantz, T. 1984. Spacing Strategies, kin Selection, and Population Regulation in Altricial
Vertebrates. Oikos 42, 48–58.
Webb, W.C., Marzluff, J.M., Hepinstall-Cymerman, J. 2012. Differences in Space Use by
Common Ravens in Relation to Sex, Breeding Status, and Kinship. The Condor 114,
584–594.
Webber, Q.M.R., Dantzer, B., Lane, J.E., Boutin, S., McAdam, A.G. 2023. Density-dependent
plasticity in territoriality revealed using social network analysis. Journal of Animal
Ecology 92, 207–221.
Wey, T., Blumstein, D.T., Shen, W., Jordán, F. 2008. Social network analysis of animal
behaviour: a promising tool for the study of sociality. Animal Behaviour 75, 333–344.
Wilmers, C.C., Nickel, B., Bryce, C.M., Smith, J.A., Wheat, R.E., Yovovich, V. 2015. The
golden age of bio-logging: how animal-borne sensors are advancing the frontiers of
ecology. Ecology 96, 1741–1753.
Wood, S.N. 2017. Generalized Additive Models: an introduction with R, 2nd ed. CRC Press,
Florida.
Zub, Kowalczyk, Jedrzejewska, Theuerkauf, Schmidt, Jedrzejewski 2003. Wolf pack territory
marking in the Bialowieza primeval forest (Poland). Behaviour 140, 635–648.
751

Appendix A: Workflow to exclude GPS locations associated to extra-territorial trips for

home range estimation

Some individuals performed extra-territorial trips (Figure A1), which affected home range estimation. We considered these GPS locations associated to extra-territorial trips as outliers and identified them by calculating the distance between the core of the home range and each GPS location in the R package ctmm with the function 'outlie' (Calabrese et al., 2016). Figure A2 shows distance values for the individual BORR in 2019 (green male on Figure 1A of the main text) and VOVV in 2022 (dark purple male on Figure 1C of the main text). Zooming in on lower distance values helped to determine cut-offs for individuals travelling far from their home range core (Figure A3). For BORR, we removed GPS locations >1700 m from the home range core, which removed 9% of its original dataset. For VOVV, we removed GPS locations >2300 m from the home range core, which removed 11% of its original dataset. These thresholds were determined visually so that extra-territorial trips disappeared from the spatial representation of the individual's movements (Figure A4). We also removed 4% of GPS locations for RMJJ in 2019 (green female on Figure 1B), 6% for BBJO in 2019 (grey male on Figure 1A), and < 1% for all others.

Figure A1. GPS locations for A) individual BORR in 2019 and B) individual VOVV in 2022, that performed a few extra-territorial trips during the study period. X and Y axis show distance from home range core.

Figure A2. Distances between GPS locations and the core of the home range for A) individual BORR in 2019 and B) individual VOVV in 2022.

Figure A3. Zoom in on distances between GPS locations and the core of the home range for A) individual BORR in 2019 and B) individual VOVV in 2022. Red lines show the 1700 m and 2300 m cut-offs respectively.

Figure A4. GPS locations for A) individual BORR in 2019 and 2) VOVV in 2022 after the removal
of extra-territorial trips.

Appendix B: Relation between behavioural state and accelerometry classification

30-sec accelerometry bursts are composed of ten 3-sec sequences which were attributed to a behaviour (Motionless, Walking, Digging or Running) through a machine learning classification (see the main text and Clermont et al., 2021b). Behavioural states (active or inactive) were attributed to each GPS location through a hidden Markov model (HMM, see main text and Grenier-Potvin et al., 2021 for the details). To evaluate the relation between behavioural states and the accelerometry classification, we first associated each 30-sec burst of accelerometry data to the closest GPS location (provided the time stamp of the GPS location was within 30 sec of the start or end of the burst). We found Running was the most common behaviour for the active state, while Motionless was by far the most common behaviour for the inactive state (Figure B1).

Figure B1. Average number of 3-sec sequences attributed to each behaviour for both behavioural states.

Appendix C: Exclusion of datapoints near study area boundaries

Interactions with foxes that were not tracked occurred on the outer margin of our study area, thus biasing negatively our CDE estimates. For that reason, when we determined whether foxes were territorial (i.e., whether they had a part of their home range located outside of the 95% CDE), we excluded areas within 500 m of study area boundaries (Figure 2). This led to three individuals considered as non-territorial in 2019 (green female RMJJ, green male BORR and grey male BBJO, Figure 2), and one in 2022 (dark orange male BOBB, Figure 2). Using a larger buffer size of 1 km would have led to the pink female BJOV in 2022 to also be considered as non-territorial, as the 1 km buffer overlaps all her home range (Figure C1).

To test the effect of the probability of encountering a neighbour on Arctic fox behaviour, we excluded from our dataset all locations falling within the 500 m inner buffer of the study area and considered that BJOV of 2022 was territorial (results in Table 1). Due to uncertainty in BJOV territorial degree, we repeated the analyses of 2022 while excluding this individual (Table C4). We then find no evidence of an effect of the probability of encounter on the probability of being active for territorial individuals in 2022 (Table C4). Other results are similar to those shown in Table 1.

We also verified whether our choice of buffer size influenced the results by repeating models, this time excluding locations falling within an inner buffer of 1 km instead of 500 m. This also resulted in excluding all locations of individual BJOV in 2022 as they were all within the 1 km buffer (Figure C1). We find no evidence of an effect of the probability of encounter on the probability of being active for territorial individuals in 2019 and 2022 (Table C5). Other results are similar to those shown in Table 1.

Figure C1. Study area with 95% auto-correlated kernel density estimation contours for A) 7 male
and 6 female Arctic foxes and B) 5 male and 5 female Arctic foxes in 2022. The light grey shaded
area represents a 500 m inner buffer of study area boundaries where observations were excluded,
while the dark grey shaded area indicates a 1 km buffer. The thick black line is the 95% CDE
contour used to determine whether foxes were territorial, and coloured lines are 95% AKDE
contours. Males and females are represented by solid and dashed lines, respectively, and members
of a given pair are identified by the same colour (within year).

Table C4. Estimates of generalized additive mixed models predicting the probability of fox being
active in A) 2022 (n = 1500 GPS locations), and the probability of fox digging in B) 2022 (n =
1500 accelerometry bursts), while excluding female BJOV for which degree of territoriality was
uncertain. Model specifications are the same as those described in Table 1. Changes in significance
in comparison to Table 1 are highlighted.

A) Probability of fox being active in 2022

Component	Term	Estimate	Std.Error	z.value	p.value
1. Parametric coefficients	(Intercept)	0.73	0.18	4.19	<0.001
	Probability of encountering a neighbour	0.05	0.07	0.79	0.430
	Territorial yes/no [no]	0.18	0.37	0.47	0.640
	Sex [male]	-0.29	0.24	-1.21	0.226
	Prob encounter: Territorial yes/no [no]	-0.07	0.18	-0.38	0.701

Component	Term	edf	Ref.df	Chi.sq	p.value
2. Smooth terms	s(ToD)	6.48	8.00	220.58	<0.001
	s(day)	1.00	1.00	0.10	0.754
	s(Fox ID)	4.00	6.00	12.19	0.008
	s(day_factor)	6.70	22.00	9.57	0.110

B) Probability of fox digging in 2022

Component	Term	Estimate	Std.Error	z.value	p.value
1. Parametric coefficients	(Intercept)	-0.27	0.25	-1.07	0.285
	Probability of encountering a neighbour	0.070	0.06	1.14	0.254
	Territorial yes/no [no]	0.292	0.56	0.52	0.603
	Sex [male]	-0.14	0.35	-0.39	0.699
	Prob encounter: Territorial yes/no [no]	-0.08	0.17	-0.45	0.655

Component	Term	edf	Ref.df	Chi.sq	p.value
2. Smooth terms	s(ToD)	4.39	8.00	83.27	<0.001
	s(day)	1.00	1.00	3.21	0.073
	s(Fox ID)	5.23	6.00	40.30	<0.001
	s(day_factor)	4.64	22.00	6.26	0.163

Table C5. Estimates of generalized additive mixed models predicting the probability of fox being active in A) 2019 (n = 1950 GPS locations) and B) 2022 (n = 1500 GPS locations), and the probability of fox digging in C) 2019 (n = 1950 accelerometry bursts) and D) 2022 (n = 1500 accelerometry bursts), while excluding locations falling within a 1 km inner buffer of study area boundaries. Model specifications are the same as those described in Table 1. Changes in significance in comparison to Table 1 are highlighted.

A) Probability of fox being active in 2019

Component	Term	Estimate	Std.Error	z.value	p.value
1. Parametric coefficients	(Intercept)	0.18	0.24	0.76	0.448
	Probability of encountering a neighbour	0.01	0.07	0.18	0.856
	Territorial yes/no [no]	0.09	0.38	0.24	0.810
	Goose nest density	0.59	0.07	8.90	<0.001
	Sex [male]	0.27	0.31	0.86	0.388
	Prob encounter: Territorial yes/no [no]	-1.03	0.15	-6.86	<0.001

Component	Term	edf	Ref.df	Chi.sq	p.value
2. Smooth terms	s(ToD)	6.57	8.00	183.47	<0.001
	s(day)	2.75	3.24	5.25	0.194
	s(Fox ID)	8.70	10.00	59.33	<0.001
	s(day_factor)	5.56	21.00	8.44	0.067

B) Probability of fox being active in 2022

Component	Term	Estimate	Std.Error	z.value	p.value
1. Parametric coefficients	(Intercept)	0.72	0.32	2.29	0.022
	Probability of encountering a neighbour	-0.05	0.07	-0.61	0.540
	Territorial yes/no [no]	-0.65	0.69	-0.95	0.344
	Sex [male]	0.45	0.44	1.03	0.302
	Prob encounter: Territorial yes/no [no]	-0.04	0.21	-0.19	0.853

Component	Term	edf	Ref.df	Chi.sq	p.value
2. Smooth terms	s(ToD)	5.93	8.00	246.96	<0.001
	s(day)	1.00	1.00	1.93	0.165
	s(Fox ID)	5.26	6.00	35.75	<0.001
	s(day_factor)	9.60	22.00	19.42	0.009

C) Probability of fox digging in 2019

Component	Term	Estimate	Std.Error	z.value	p.value
1. Parametric coefficients	(Intercept)	-0.87	0.34	-2.56	0.011
	Probability of encountering a neighbour	-0.04	0.07	-0.50	0.619
	Territorial yes/no [no]	0.01	0.54	0.03	0.979
	Goose nest density	0.32	0.06	5.04	<0.001
	Sex [male]	-0.17	0.45	-0.38	0.700
	Prob encounter: Territorial yes/no [no]	-0.47	0.14	-3.29	0.001

Component	Term	edf	Ref.df	Chi.sq	p.value
2. Smooth terms	s(ToD)	2.89	8.00	33.98	<0.001
	s(day)	2.77	3.29	9.20	0.038
	s(Fox ID)	9.33	10.00	130.70	<0.001
	s(day_factor)	5.34	21.00	8.64	0.053

D) Probability of fox digging in 2022

Component	Term	Estimate	Std.Error	z.value	p.value
1. Parametric coefficients	(Intercept)	-0.26	0.29	-0.90	0.371
	Probability of encountering a neighbour	-0.035	0.07	-0.51	0.608
	Territorial yes/no [no]	0.059	0.65	0.09	0.927
	Sex [male]	0.04	0.41	0.11	0.915
	Prob encounter: Territorial yes/no [no]	0.07	0.19	0.38	0.703

Component	Term	edf	Ref.df	Chi.sq	p.value
2. Smooth terms	s(ToD)	5.01	8.00	70.07	0.000
	s(day)	4.54	5.55	14.89	0.016
	s(Fox ID)	5.39	6.00	49.00	0.000
	s(day_factor)	0.00	22.00	0.00	0.817