Differential induction of *Medicago truncatula* defence metabolites in response to rhizobial symbiosis and pea aphid infestation

Goodluck BENJAMIN[1], Marie PACOUD[1], Stéphanie BOUTET[2], Gilles CLEMENT[2], Renaud BROUQUISSE[1], Jean-Luc GATTI[1], Marylène POIRIÉ[1*], Pierre FRENDO[1*]

1   Université Côte d'Azur, INRAE, CNRS, Institut Sophia Agrobiotech, 06903 Sophia Antipolis, France
2   Université Paris-Saclay, INRAE, AgroParisTech, Institut Jean-Pierre Bourgin for Plant Sciences (IJPB), 78000 Versailles, France

*Both authors supervised this work.

Correspondence: pierre.frendo@univ-cotedazur.fr; goodluck.benjamin@etu.univ-cotedazur.fr

Figures: 6
Tables: 1
Total word count: 6250
Introduction: 680
Materials and Methods: 2039
Results: 1590
Discussion: 1941

Summary

• Legumes symbiosis with rhizobial nitrogen-fixing bacteria enable them to grow in

nitrate-depleted soils. Rhizobial symbioses also induces systemic plant defence against

bioagressors.

• We investigate how nitrogen-fixing symbiosis (NFS) in the legume *Medicago*

truncatula can prime plant defence against the pea aphid *Acyrthosiphon pisum*. We

analysed metabolite modification both by LC-MS and GC-MS and defence pathway

gene expression by qPCR in leaves of both NFS and nitrate-fed (non-inoculated; NI)

plants after aphid infestation (Amp).

• The accumulation of primary and secondary metabolites was modulated by both NFS

and aphid infestation. 62 defense-related metabolites such as salicylate, pipecolate,

gentisic acid and several soluble sugars were differentially regulated by aphid

infestation in both NFS and NI conditions. 19 metabolites, including triterpenoid

saponins, accumulated specifically in NFS_Amp conditions. Gene expression analysis

showed that aphid-infested plants exhibited significantly higher expression of *Chalcone*

isomerase, *flavonol synthase*, *hydroxyisoflavone-O-methyl transferase* and

Pterocarpan synthase, while *D-pinitol dehydrogenase* was only significantly induced

in NI infested leaves.

• Our data suggest that NFS, in addition to being a plant nitrogen provider, stimulates

specific legume defenses upon pest attack and should also be considered as a potential

tool in Integrated Pest Management strategy.

Key words: *Medicago truncatula*, *Sinorhizobium meliloti*, nitrogen-fixing symbiosis,

Acyrthosiphon pisum, metabolomics, plant defence priming.

Introduction

Plants are under constant threat from pathogens and insect pests such as sap-feeding aphids (National Research Council *et al.*, 1985). Aphids are deleterious agronomic pests, not only because they feed on phloem sap and therefore weaken the plant, but also because they are vectors for various plant viruses (Ng & Perry, 2004). More than 5000 species of aphids are known today, a diversity that is partly due to sympatric speciation initiated by individuals adapting to new host plants (Diehl & Bush, 1984; Drès & Mallet, 2002). During feeding, the aphid stylet injures plant cells, inject saliva, and sucks up tiny amounts of the cell content to decide on plant acceptance (Martin *et al.*, 1997; Lu *et al.*, 2016). Previous studies have revealed that the secreted proteins present in saliva trigger plant responses (Pitino & Hogenhout, 2013; Rodriguez & Bos, 2013). These saliva proteins act as herbivore-associated molecular patterns (HAMPs) (Kaloshian & Walling, 2016) that bind to host plant pattern recognition receptors (PRRs) and trigger an immune response associated with the production of reactive oxygen species (ROS) and defence hormones Salicylic Acid (SA) and Jasmonic Acid (JA) pathways (Wu & Baldwin, 2010; Herrera-Vásquez *et al.*, 2015). The increase of SA and JA regulate the accumulation of various primary and secondary metabolites that play a role in plant defence (Isah, 2019) as feeding deterrents or toxins that decrease food intake or food use efficiency, decrease survival and reproduction of the pest, or indirectly act as attractants for its natural enemies (War *et al.*, 2012).The secondary metabolites involved in this process include terpenoids, phenolics, cyanogenic glycosides, glucosinolates, and alkaloids (Maag *et al.*, 2015; Züst & Agrawal, 2016). For example, high levels of saponins and phenolic compounds increase aphid mortality and result in reduced pest population (Züst & Agrawal, 2016), flavonoid glycosides also reduce aphid fecundity, while nitrogen-containing compounds cause host plant rejection (Kordan *et al.*, 2012).

One property of legume plants is their ability to associate with rhizobia to perform nitrogen fixation symbiosis (NFS), i.e., the conversion of atmospheric N_2 into plant-usable ammonium. The microbial partner supplies assimilable nitrogen to the plant in exchange for carbon resources and a protective environment in the root nodules (Lee & Hirsch, 2006). NFS has also been reported to potentially provide defence priming to plants against bio-aggressors (Benjamin *et al.*, 2022). For example, in pea (*Pisum sativum*), rhizobium inoculation decreased *Didymella pinodes* disease severity and significantly reduced the seed infection level (Desalegn *et al.*, 2016; Ranjbar Sistani *et al.*, 2017). NFS has also been reported to modulate resistance to biotrophic pathogens in both *Medicago truncatula* and pea by reducing penetration and sporulation of the powdery mildew fungus *Erysiphe pisi* (Smigielski *et al.*, 2019).

We have previously demonstrated that NFS also influences the *Medicago truncatula*–pea aphid

interaction and general plant defence response (Pandharikar *et al.*, 2020). A detrimental effect

of rhizobia-inoculated plants on aphid development was observed, with lower adult weights

compared to aphids from nitrate-fed plants. *Pathogenesis Related 1* (*PR1*) gene expression was

upregulated in aphid-infested shoots, indicating the activation of SA-dependent defence.

Moreover, a significantly higher expression of *Proteinase Inhibitor (PI)* gene, a marker for the

JA transduction pathway, was observed in NFS plants compared to nitrate-fed plants

(Pandharikar *et al.*, 2020).

Due to the observations from our preliminary studies, we have analysed by metabolomics, the

impact of the nitrogen source (KNO_3 *vs* NFS) on *M. truncatula* infestation by pea aphid. To

this end we analysed leaf metabolite profiles of NFS plants with and without aphid infestation

and compared them to KNO_3-fed plants (Non Inoculated; NI) and to obtain a maximum

coverage of the leaf metabolites, untargeted metabolomic was done both by gas

chromatography/mass spectrometry (GC-MS) and liquid chromatography/mass spectrometry

(LC-MS). Our results showed that both primary and secondary metabolisms are significantly

modulated by nitrogen source and aphid infestation. Gene expression analysis of enzymes

involved in the secondary metabolite synthesis pathways showed that regulation of secondary

metabolism is partially mediated by the modulation of gene expression. We identified specific

metabolites which are differentially regulated in NFS plants compared to NI plants under aphid

infestation.

Materials and Methods

Biological materials and experimental design

The pea aphid *Acyrtosiphon pisum* clone, YR2-amp, further named YR2, is derived from a

clover biotype line collected in England that was freed from the secondary symbiont *Regiella*

insecticola by ampicillin treatment (Simon *et al.*, 2011). This aphid line is stable (more than

15 years old) and was maintained on fava bean, 20°C, 16:8h light/dark cycle. To synchronize

aphid, 20-40 apterous female adults were placed in a petri dish containing a fava bean leaf and

let to reproduce for 24hrs. Then, 10 nymphs (L1) were collected and used for infestation.

For one experiment, four batches of 5 pots containing each six *Medicago truncatula* A17 plants

were grown as previously described (Pandharikar *et al.*, 2020) (Fig. S1). After 12 days, two

groups were inoculated with the nitrogen-fixing bacteria *Sinorhizobium meliloti* 2011

(nitrogen-fixing symbiosis, NFS condition), and two were supplemented once with 10 ml of 5

mM KNO$_3$ solution (non-inoculated, NI condition). Seven days after (the time to NFS plants

to develop nodules), for both NFS and NI conditions, one batch of NI plants and one batch of

NFS plants were infested with aphids (10 L1/pots). The aphid nymphs were then let to develop

into adult during 12 days. All pots were individually isolated in a ventilated plastic box and

maintained at 20°C under 16:8h light/dark photoperiod. For each condition, control pots were

treated and maintained under the same conditions except that no aphids were introduced in the

plastic box. At the end of the 12 days, harvested leaves were used immediately or frozen in

liquid nitrogen and stored at -80°C.

Rhizobium inoculation was done with a streptomycin-resistant strain of *Sinorhizobium meliloti*

2011. It was cultured on Luria-Bertani medium supplemented with 2.5mM CaCl$_2$ and MgSO$_4$

(LBMC) and streptomycin at 200µg ml^{-1} for 3 days at 30°C, then transferred and grown in

LBMC liquid medium for 24h, pelleted at 5,000g, washed twice with sterile distilled water,

and resuspended in sterile distilled water to a final optical density of 0.05 (OD$_{600}$). Each NFS

plant was inoculated with 10 mL of this *S. meliloti* suspension.

A total of 4 biological replicates (i.e. 4 times 4 pots of six plants for each condition), produced

as described above were used for analysis carried out in this study.

Analysis of metabolites using gas chromatography/mass spectrometry (GC-MS)

Leaves were ground in liquid nitrogen to obtain a fine powder and 50 mg was resuspended in

1ml of cold (-20°C) Water:Acetonitrile:Isopropanol (2:3:3 in volume) containing Ribitol

4µg/ml as internal standard. After extraction under shaking (10 min, 4°C), insoluble material

was removed by centrifugation at 20,000g for 5min and 50µl were collected and dried

overnight (SpeedVac™) and used immediately or stored at -80°C. Three blank tubes underwent

the same treatments to estimate possible contamination. A quality control was made by pooling

an equal volume of each condition.

Samples were warmed 15 min before opening and dried again for 1.5h at 35°C before addition

of 10µl of 20 mg ml^{-1} methoxyamine in pyridine and the reaction was performed for 90 min at

28°C under continuous shaking. 90µl of N-methyl-N-trimethylsilyltrifluoroacetamide

(MSTFA) were then added and the reaction continued for 30 min at 37°C. After cooling, 45µl

were taken for injection. 1µl of derivatized sample was injected in splitless and split (1:30)

modes on a gas chromatograph (Agilent 7890A; Santa Clara, USA) coupled to a mass

spectrometer (Agilent 5977B) with a heated separation column (Rxi-5SilMS; Restek, Lisses,

France) (temperature ramp: 70°C for 7 min then 10°C/min to 330°C for 5 min; run length 38

min). Helium flow was constant at 0.7 mL/min. 5 scans per second were acquired spanning a

50 to 600Da range. Instrument was tuned with PFTBA with the *m/z* 69 and *m/z* 219 of equal

intensities. Samples were randomized. Three independent quality controls were injected at the

beginning, middle and end of the analysis for monitoring the derivatization stability. An alkane

mix (C10, C12, C15, C19, C22, C28, C32, C36) was injected during the run for external

calibration. Three independent derivatizations of the quality control were injected at the

beginning, in the middle and at the end of the series. A response coefficient was determined

for 4ng each of a set of 103 metabolites, respectively to the same amount of ribitol. This

compound was used to give an estimation of the absolute concentration of the metabolite in

what we may call a "one point calibration" (Fiehn, 2006, 2008).

Analysis of metabolites using liquid chromatography/mass spectrometry (LC-MS)

Metabolites were extracted from 6 mg of fresh weight ground sample using a protocol adapted

from the literature (Kim *et al.*, 2008). Briefly, 1.6ml of a mix of Methanol/H2O/Acetone/TFA

(40/32/28/0.05, v:v:v:v) and 300ng of Apigenin (used as internal standard) were added to each

sample, which was then stirred at 4°C for 30 minutes. After centrifugation (10 min, 20,000 g,

4°C), the supernatant was collected, and the pellet was extracted again by stirring with 1.6ml

of the previous solvent mix for 30 minutes. After centrifugation the two supernatants were

pooled, dried and resuspended in 200µl of water (ULC/MS grade)/Acetonitrile (90/10)

(Biosolve Chimie, Dieuze, France) and filtered (filter paper grade GF/A Whatman®).

Metabolomics data were acquired using a UHPLC system (Ultimate 3000, Thermo Scientific,

Waltham, USA) coupled to quadrupole time of flight mass spectrometer (Q-Tof Impact II

Bruker Daltonics, Bremen, Germany). A Nucleoshell RP18 plus reversed-phase column (2 x

100 mm, 2.7 µm; Macherey-Nagel, Hoerdt, France) was used for chromatographic separation.

The mobile phases used for the chromatographic separation were (A) 0.1% formic acid in H_2O

and (B) 0.1% formic acid in acetonitrile. The flow rate was of 400µl min^{-1} and the following

gradient was used: 95% of A for 1-min, followed by a linear gradient from 95% A to 80% A

from 1 to 3-min, then a linear gradient from 80% A to 75% A from 3 to 8-min, a linear gradient

from 75% A to 40% A from 8 to 20-min. 0% of A was hold until 24-min, followed by a linear

gradient from 0% A to 95% A from 24 to 27-min. Finally, the column was washed by 30% A

for 3.5min then re-equilibrated for 3.5-min (35-min total run time). Data-dependent acquisition

(DDA) methods were used for mass spectrometer data in positive and negative ESI modes

using the following parameters: capillary voltage, 4.5kV; nebuliser gas flow, 2.1 bar; dry gas

flow, 6L min^{-1}; drying gas in the heated electrospray source temperature, 140ºC. Samples were

analysed at 8Hz with a mass range of 100 to 1500 m/z. Stepping acquisition parameters were

created to improve the fragmentation profile with a collision RF from 200 to 700 Vpp, a transfer
time from 20 to 70 µs and collision energy from 20 to 40eV. Each cycle included an MS full-
scan and 5 MS/MS CID on the 5 main ions of the previous MS spectrum.
208
Soluble sugars and starch analysis
Soluble sugars (sucrose, glucose, and fructose) were extracted from 150 mg of frozen tissue
powder (both roots and leaves separately) with ethanol and water solution (800µl at 80% of
ethanol) by incubation in a water bath at 80°C for 15 min, with shaking every 5-minute. The
sample was centrifuged (5 min, 5,000 rpm) and the supernatant collected. The extraction was
repeated using the same conditions with 800µl of a 50% ethanol solution, 800µl of 100% water
and then 800µl 80% ethanol solution. All supernatants were mixed and then evaporated. The
dried sample was resuspended in 1ml of water and kept in the dark at -20°C until soluble sugar
analysis.
The residual pellet from sugar extraction was used for starch analysis. Immediately after
removal of the last supernatant, the pellet was resuspended in 3ml of thermostable α-amylase.
The tube was plunged in a boiling water bath and mixed every 2 mins. After 6 min, the sample
was transferred to 50°C bath and 0.1ml of amyloglucosidase (20U) was added, the tube was
mixed and incubated for 30 min. The entire content of the tube was then transferred to a larger
tube and volume adjusted to 20ml with distilled water, mixed thoroughly, and centrifuged at
5,000 rpm for 5 min. The clear, undiluted supernatant was used for the determination of glucose
issued from starch hydrolysis.
Sugar analyses were performed using Sucrose, D-Fructose and D-Glucose assay kit
(Megazyme K-SUFRG; Neogene, Mi, USA) and Total starch HK assay kit (Megazyme K-
TSHK), respectively, as described by the provider. Spectrophotometric measurements were
conducted at 340nm in cuvettes with a 1 cm optical path.
230
Gene Expression Analysis
For RNA extraction, plant material was ground in liquid nitrogen. Total RNAs from 100mg
was then isolated using RNAzol® RT (Sigma), quantified, and analysed on nanodrop and 1.5%
agarose gel electrophoresis to assess the purity. DNA digestion (RQ1 RNAse-free DNAse) and
reverse transcription (GoScript™ Reverse Transcription) were performed as described by the
manufacturer (Promega, Madison, USA). qPCR was performed (qPCR Master Mix plus CXR;
Promega) using cDNA template and each set of primers. *Pathogenesis Related 1* (*PR1*;
MtrunA17_Chr2g0295371) was used as SA defence gene marker and the *Proteinase Inhibitor*

(*PI*; PSI-1.2; MtrunA17_Chr4g0014461) as JA pathway activation gene (Pandharikar *et al.*,

2020). Other genes of interest analysed were *Chalcone Isomerase* (*CHI*;

MtrunA17_Chr1g0213011), *Flavanol Synthase/Flavanone 3-Hydroxylase* (*FLS/F3H*;

MtrunA17_Chr3g0092531), *Isoflavone 4'-O-methyltransferase* (*HI4'O-MT*;

MtrunA17_Chr4g0046341), *D-pinitol Dehydrogenase* (*OEPB*; MtrunA17_Chr6g0480011),

Phenylalanine Ammonia-lyase (*PAL*; MtrunA17_Chr1g0181091), *Pterocarpan Synthase 1*

(*PTS*; MtrunA17_Chr7g0259091) and *SAR Deficient 4* (*SARD4*; MtrunA17_Chr1g0202471) .

Real-time qPCR was performed with specific primers (Table S1) using 95°C for 3 min

followed by 40 cycles at 95°C for 3 sec and 60°C for 30 sec, and melting curves from 65°C to

95°C in increments of 0.5°C (AriaMx Real-time PCR machine, Agilent). Cycle threshold

values (Ct) were normalized to the average Ct of two housekeeping genes *MtC27*

(MtrunA17_Chr2g0295871) and *a38* (MtrunA17_Chr4g0061551) genes (Del Giudice *et al.*,

2011). The expression of these two genes was not affected by the treatments. The original Ct

obtained (Ariamix software; Agilent) (Dataset S1) were further used in the R qPCRBASE

package (Hilliou, 2013). For each gene, the expression level of the aphid infested plants was

compared with that of the non-infested control plants. The results of the qPCR analysis were

generated from four independent biological repeats.

Metabolomics Data Processing

For GC-MS, raw datafiles were converted in NetCDF format and analysed with AMDIS

software (http://chemdata.nist.gov/mass-spc/amdis/). A home retention indices/mass spectra

library built from the NIST, Golm (http://gmd.mpimp-golm.mpg.de/) and Fiehn databases

(https://fiehnlab.ucdavis.edu), and standard compounds was used for metabolites

identification. Peak areas were also determined with the Targetlynx software (Waters) after

conversion of the NetCDF file in masslynx format. AMDIS, Target Lynx in split-less and split

30 modes were compiled in one single Excel file for comparison. After blank mean subtraction

peak areas were normalized to Ribitol and leaves fresh weight.

For LC-MS, data files were converted and treated as previously described (Boutet *et al.*, 2022).

A first search was done in library with open-source software MZmine2 (Pluskal *et al.*, 2010)

with identification module and "custom database search" to begin the annotation with our

library, currently containing 159 annotations (RT and *m/z*) in positive mode and 61 in negative

mode, with RT tolerance of 0.3 min and m/z tolerance of 0.0025 Da or 6 ppm. Molecular

networks were generated with MetGem software (Olivon *et al.*, 2018)

(https://metgem.github.io) using the .mgf and .csv files obtained with MZmine2 analysis. ESI-

and ESI+ molecular networks were generated using cosine score thresholds of 0.8 in two
modes. Metabolites annotation was performed as described (Boutet *et al.*, 2022).
275
Statistical Analysis
Statistical analysis was performed with TMEV (https://sourceforge.net/projects/mev-tm4/) for
GC-MS. Univariate analyses by permutation (1-way-anova and 2-way anova) were firstly used
to select the significant metabolites (P-value < 0.01). Multivariate analyses (hierarchical
clustering an principal component analysis) were then made on both LC-MS and GC-MS data.
Mapman (http://www.gabipd.org/projects/MapMan/) was used for graphical representation of
the metabolic changes after Log2 transformation of the mean of the 3 replicates (Fiehn, 2006,
2008).
Tukey's multiple comparison tests were also performed on independent metabolites to identify
statistical differences between the various conditions for both primary and secondary
metabolites. Statistical analysis of individual metabolites was done using Prism v9.1.1
(GraphPad software, USA). All experimental data are expressed as mean ± standard error (SE).
Venn diagram was created based on results of Tukey's tests, clustering metabolites based on
statistical significance (P-value ≤ 0.05) using InteractiVenn (Heberle *et al.*, 2015).
290
Results
Aphid infestation, more than nitrogen source, significantly alters leaf metabolite profile.
M. trunculata leaves metabolites were analysed in four experimental conditions: nitrate-fed
plants (NI), nitrate-fed plants infested with aphids (NI_Amp), nitrogen-fixing symbiotic plants
(NFS) and nitrogen-fixing symbiotic plants infested with aphids (NFS_Amp) (Fig. S2). The
leave extracts were analysed by GC-MS, to obtain mainly primary metabolites, and by LC-MS,
to have access to the secondary metabolites. After GC-MS, 237 compounds were retained and
126 could be identified with confidence (Dataset S2), and among them the five main
quantitative compounds were sucrose, phosphate, malate, citrate, and glutamate (Fig. S3). By
LC-MS, 2627 compounds (in positive and negative modes) were obtained and 213 could be
identified (Dataset S2). We observed a strong effect of aphid infestation and nitrogen source in
the accumulation of metabolites and the hierarchical clustering of these compounds separates
both nitrogen source and plant infestation status (see Fig. 1A, S4 and S5 for GC-MS and Fig.
S6 for LC-MS, respectively). From the four experimental conditions, a total of 194 unique
metabolites, 126 from GC-MS and 69 from the LC-MS, were found significantly different

according in their accumulation (Dataset S2). Only sucrose was found significantly in both analysis. Among the 194 identified the most represented classes were flavonoids with 40 different compounds, followed by 30 carboxylic acids, 26 amino acids and 19 sugars (Fig. 1b).

Fig. 1: Aphid infestation, more than nitrogen source, significantly alters leaf metabolite profile. a) Heatmap from GC-MS analysis showing unsupervised hierarchical clustering of compounds separated by both infestation and nitrogen source. b) Class abundance of metabolites from both GC-MS and LC-MS analyses.

We then identified statistical differences for the 194 unique metabolites between the four experimental conditions using multiple comparison tests (Fig 2; Dataset S3). 86 compounds were not found significatively accumulated in one specific condition, including many of the amino acids. 108 compounds were found to be differential accumulated in the different conditions. Amongst them, aphid infestation increased significantly the accumulation of 62 metabolites. 18 metabolites accumulated in control NFS conditions and in infected conditions, suggesting they are linked to a general infection/infestation response. 2 compounds are specifically found in NFS conditions and 19 compounds were significantly accumulated in NFS_Amp conditions.

Fig. 2: Metabolite distribution according to nitrogen source and aphid infestation affects the metabolite profile of plants. Venn diagram showing distribution of statistically significant metabolites across the various experimental conditions. The name of the compounds and the statistic can be found in Dataset S3.

Aphid infestation triggers a greater accumulation of sugars than amino acids in both NFS and NI Plants.

Amino Acids

The majority of the amino acids (20) were not significantly accumulated in any specific condition. In contrast, tryptophan, cysteine, asparagine were significantly accumulated in NFS control, NFS and NI Amp. 5-Hydroxynorvaline was significantly accumulated in NFS control and NFS Amp compared to NI control and NI Amp. Finally, the putative asparagine or asparagine derivative (U1194.6/188[Asn?]) is significantly more accumulated in NFS and NI Amp than in NFS control and NI control (Dataset S3).

Sugars

Among the 62 metabolites significantly accumulated under the aphid-infested conditions, glucose, sucrose and fructose contents were increased between three to five-fold compared to control conditions in leaves (Fig. 3). To investigate whether this leaf increase in sugar was associated with either a reduction in sugar transport from leaf to root or the results of the mobilization of leaf starch, sugar and starch contents were measured using biochemical assays in the leaves and roots of plants under the different conditions (Fig. 4). The increase in glucose, sucrose and fructose in infested leaves was confirmed (Fig. 4), but no significant accumulation

of starch in leaves and no change in root sugar concentration were observed, whatever the

plant's growth conditions (Fig. 4), suggesting an alteration of the leaf sugar metabolism in

response to aphid infestation.

Fig. 3: Aphid infestation triggers accumulation of soluble sugars in both NFS and NI plant leaves. Bar graphs from

GCMS analysis showing the significant accumulation of glucose sucrose and fructose in NFS and NI plants upon aphid

infestation. Metabolites analysed and quantified in µg mg^{-1} of fresh weight (FW) of leaves of NFS and NI plants 12 days after

infestation by the aphids (control = no aphid infestation). Mean ± SE (n=4): different letters indicate a $p \leq 0.05$ (Tukey's test).

Fig. 4: Aphid infestation affects sugar concentration in leaves but not in roots. Biochemical analysis showing the

significant accumulation of glucose, sucrose, and fructose in leaves of NFS and NI plants. No significant difference was found

in roots upon aphid infestation. No statistical difference was observed in leaves starch, measured as glucose equivalent,

between the different conditions. Metabolites analysed and quantified in µmol g^{-1} of fresh weight (FW) of leaves and roots

from NFS and NI plants 12 days after infestation by the aphids (control indicates no aphids infestation). Mean± SE (n=4);

different letters indicate a $p \leq 0.05$ (Tukey's test).

Aphid infestation triggers accumulation of defence related metabolites in both NFS and NI plants.

Amongst the metabolites significantly accumulated during aphid infestation (Fig. S5; Dataset S3), many are involved in the regulation of plant defence pathways such as salicylates, pipecolate, an intermediate of the lysine catabolic pathway, pinitol, a cyclitol derived from myo-inositol, as well as secondary metabolites with known defence activity such as the putative daidzin, the glycoside form of the aglycone daidzein (Fig. 5).

Fig. 5: Aphid infestation triggers accumulation of defence-related metabolites in both NFS and NI particularly from the salicylic acid defence pathway. Graphs show accumulation of metabolites expressed as arbitrary unit (in mg apigenin equivalent mg^{-1}) of leaves fresh weight (FW) of NFS and NI plants 12 days after infestation by the aphids (control indicates no aphids infestation). Mean± SE (n=4); different letters indicate a $p \leq 0.05$ (Tukey's test).

Aphid infestation induces expression of genes involved in the flavonoid synthesis pathway.

In order to test whether pipecolate, pinitol, and diadzin accumulation resulted also from an increase in expression of the gene(s) involved in their synthesis pathways, we have measured the expression of *Chalcone isomerase (CHI), flavonol synthase/ flavonone 3β-hydroxylase (FLS/F3H), hydroxyisoflavone-O-methyl transferase* (*HI4'O-MT*) and *Pterocarpan synthase* (*PTS), Phenylalanine Ammonia Lyase (PAL), SAR-DEFICIENT4 (SARD4)* and (*OEPB*) by RT-qPCR (Table 1). Since we previously showed that *PR1,* a marker for SA defence pathway

and *PI*, a marker for jasmonic acid defence pathway were also differently induced in NFS and NI plants after aphid infestation (Pandharikar *et al.*, 2020), we test these genes as the plant condition controls. Here, RT-qPCR analysis showed that *PR1* expression increased 3.4 times more in NI_Amp condition than in NFS_Amp condition. In contrast *PI* was 2.3 times more expressed in NFS_Amp condition than in NI_Amp condition, in agreement with our previous results. The analysis of the expression of genes involved in secondary metabolism pathway showed that no significant change in *SARD4* and *PAL* expression was detected in aphid infected compared to the control plants. *OEPB* expression was significantly induced in NI_Amp leaves but not NSF_Amp leaves, although the amplitude of increase was similar. *CHI*, *FLS/F3H*, *HI4'O-MT* and *PTS* showed significant induction upon aphid infestation in both NFS and NI conditions; the induction being 2-fold for *FLS/F3H*, and reaching more than 50-fold for *PTS* in NSF_Amp.

Taken together these results showed that the accumulation of secondary metabolites is at least partially associated to an higher expression of enzymes involved in their synthesis pathways.

Table 1: Gene expression analysis of NFS and NI plants.

Gene	Pathway	Product	Fold Increase from Control (Mean ± SE)		*p* value	
			NFS_Amp/ NFS	NI_Amp/NI	NFS_Amp/ NFS	NI_Amp/NI
PR1	SA pathway	Salicylic acid	3.5 ± 0.32	11.99 ± 1.25	0.05	0.02
PI	JA pathway	Jasmonic acid	10.72 ± 0.26	4.73 ± 0.75	0.0006	0.002
PAL	Phenylpropanoid pathway	Phenylpropanoids	1.40 ± 0.02	1.35 ± 0.14	0.06	0.31
SARD4	Lysine Biosynthesis pathway	Pipecolate	1.16 ± 0.29	0.74 ± 0.09	0.73	0.88
OEPB	Pinitol Biosynthesis	Pinitol	1.75 ± 0.01	1.96 ± 0.31	0.07	0.01
FLS/F3H	Isoflavonoid pathway	Flavonoids	1.96 ± 0.05	2.5 ± 0.40	0.00002	0.01
HI4'O-MT	Isoflavonoid pathway	Formononetin	4.32 ± 0.10	3.37 ± 0.53	0.001	0.02
CHI	Isoflavonoid pathway	Flavonoids	4.58 ± 0.11	5.83 ± 0.93	0.006	0.004
PTS	Isoflavonoid pathway	Medicarpin	53.35 ± 1.27	24.66 ± 3.91	0.0001	0.001

Rescaled qPCR expression showing the level of gene induction upon aphid infestation in NFS and NI plants (NFS_Amp and NI_Amp, respectively) compared to their control (NFS and NI, respectively). Data are expressed as mean ± standard error (SE); t-test on all genes, P ≤ 0.05, significant.

NFS induces a differential defence mechanism upon aphid infestation

Nineteen metabolites (10% of the identified metabolites) were significantly accumulated in NSF_Amp plants compared to other conditions (Fig. S7; Dataset S3). Fifteen of them are

secondary metabolites from phenyl-propanoid (13) and terpene (2) synthesis pathways. For example, amongst these metabolites, putative triterpenoid saponin 3-Glu(1-3)Glu-28-Xyl(1-4)Rha(1-2)Ara zanhic acid is increased two-fold in NFS_Amp plants compared to NI-Amp plants, putative 3-(4'O-Malonyl)Rha(1-2)Gal(1-2)GluA-Soyasaponenol B is 2.8 time more accumulated in infested NFS-Amp plants than in the other three plant growth conditions, flavonoid putative tricin 5-glucoside is about 2 times more accumulated in NFS-Amp compared to NI-Amp while glycosylsalicylate is half a fold more in NFS-Amp than in NI-Amp (Fig. 6).

Fig. 6: NFS induces a differential defence mechanism upon aphid infestation. Graph showing significantly accumulated metabolites in NFS upon aphid infestation. Metabolites expressed as arbitrary unit (in mg apigenin equivalent mg^{-1}) of leaves fresh weight (FW) of NFS and NI plants 12 days after infestation by aphids (control indicates no aphids infestation). Mean± SE (n=4); different letters indicate a $p \leq 0.05$ (Tukey's test).

Discussion

There is a growing interest in better understanding the roles of symbiotic microbes in plant defence and in a broader sense how this could influence plant interactions with bioagressors. Beneficial microbes, in addition to enhancing plant growth and development have been reported to induce defence reaction and confer protection on their host plants (Gopalakrishnan *et al.*, 2015; Liu *et al.*, 2020; Benjamin *et al.*, 2022). Rhizobia are able to induce systemic

resistance in legumes such as pigeon pea against fusarial wilt as it was found that a combination
of rhizobial strains was better in inducing resistance (Dutta *et al.*, 2008).

Upon pea infection with the fungi Didymella, Turetschek and colleagues (2017) observed a
strong increase in sugars, sugar alcohols and glycolysis/TCA intermediates when studying
cultivars in symbiosis with rhizobia and mycorrhiza. They also observed the accumulation of
galactose, raffinose, maltose, threitol, melibiose, fructose and pyruvate in the pea Protecta
cultivar. Similar accumulation was also reported for amino acid pools, however, there was
significant depletion of phenylalanine in the pea cultivar Messire (Turetschek *et al.*, 2017).

Plants produce diverse primary and secondary metabolites that are involved in various
functions including development and defence. Previous work have demonstrated that aphids
are able to modify the overall metabolite profile of plants to establish feeding and that plants
may react by producing metabolites that have antifeeding or deterrent effects (Giordanengo *et*
al., 2010; Kumar, 2017; Jakobs *et al.*, 2019; Shih *et al.*, 2023). Few metabolomic studies have
focused on pea aphids and legumes interaction (Sanchez-Arcos *et al.*, 2019) and none have
addressed the role of rhizobia bacteria on the plant metabolomic response during aphid attack.
Based on our previous observation that nitrogen-fixing symbiosis is detrimental to aphid fitness
in *M. truncatula* in association with potential changes in hormonal balance between SA and JA
(Pandharikar *et al.*, 2020), we performed an untargeted metabolomic analysis in *M. truncatula*
leaves 12 days after aphid infestation (chronic effect) in both NFS and NI plants.

In our study, among amino acids (Dataset S3), we observed an increase in asparagine content
in NFS plant leaves compared to NI control plants that could be associated with asparagine
formation in root nodule which account for 60% of nodule amino acid content (Sulieman &
Schulze, 2010). Similarly, cysteine was found to be more accumulated in symbiotic *Lotus*
japonicus plants as the nodule is an important source of reduced sulphur in the plant (Kalloniati
et al., 2015). In contrast, tryptophan accumulation in the leaves of NFS plants has not yet been
observed, and a potential role in defence mechanism can be assumed (Hiruma *et al.*, 2013). In
NFS- and NI-infested plants, the increase of asparagine, cysteine and tryptophan content
compared to NI plants could be associated with the change of metabolism induced by aphids
for better nutritional content (Chiozza *et al.*, 2010; Tegeder, 2014) or by the plant through the
induction of defence reaction (Taylor & Ostrowsky, 2019). Citrate, fumarate, malate and
succinate accumulation were not significantly different among our four growth conditions,
suggesting that tricarboxylic acid (TCA) cycle is not changed by NFS or aphid infestation. In
contrast to the large number of metabolites that did accumulate differentially in the different
growth conditions, two (5-hydroxynorvaline and tartrate) were specifically accumulated in

NFS conditions. Whereas the significance of tartrate accumulation, which is and end point product of the catabolism of ascorbic acid, is more difficult to analyse (Burbidge *et al.*, 2021), unless it is used by the rhizobial bacteria (Ramachandran *et al.*, 2011), that of the 5-hydroxynorvaline, a non-protein amino acid, may be related to its defensive functions against insects (Huang *et al.*, 2011; Yan *et al.*, 2015). No metabolite was specifically more accumulated in NI- and infested NI-conditions.

Our data show that the sugar metabolism is particularly affected by the aphid infestation (Fig. 3). Amongst the primary metabolites, 12 are sugars and represent 70% of the number of primary metabolites significantly accumulated in infested plants (Dataset S3). In our experimental conditions, we were interested by the sugar transport to root to feed the root nodule involved in biological nitrogen fixation. Indeed, we previously showed that biological nitrogen fixation is affected in the infested conditions (Pandharikar *et al.*, 2020) and we hypothesized that the decrease in biological nitrogen fixation could be associated to a reduced sugar transport from leaves to roots. However we did not observed a significant modification in the accumulation of root glucose, fructose and sucrose between control and infested plants, indicating that the modification of sugar metabolism is not associated with the sugar transport to the root. We also observed that starch seemed more accumulated in infested leaves than in control ones but this difference was not significant. In conclusion, the modification of sugar metabolism does not seem to be associated to a modification of sugar storage through starch or sugar transport to roots. Moreover, the modification of sugar metabolism by aphids seems not dependent of the nitrogen source. The accumulation of sugars has been previously reported in plant-aphid (Ponzio *et al.*, 2017) as well as in plant-pathogen interactions (Kanwar & Jha, 2019) as being involved in the coordination of plant defence signalling (Yamada & Mine, 2024), this perhaps could be an explanation for our observation.

A large number of significantly accumulated metabolites (62 metabolites, 31% of the total metabolites) is present in NFS- and NI- infested plants compared to their control counterparts. This results show that Medicago plants respond significantly to aphid infestation and that the nitrogen source does not play a significant role in their accumulation. Amongst these secondary metabolites, 72% are from 3 families, flavonoids (30), phenolics (11) and glycosides (4). These different compounds are mainly associated with plant defence against pests such as biocide activity (i.e. acacetin, chrysoeriol, daidzin), feeding deterring activity and defence signalling activity (i.e. salicylate, pipecolate) (Stochmal *et al.*, 2001; Goławska *et al.*, 2012, 2024; Kim *et al.*, 2022; Pawłowska & Stepczyńska, 2022). Pinitol has also been shown to participate to the biological control of powdery mildew in cucumber (Chen *et al.*, 2014) and myoinositol

influences the plant bacterial colonization (O'Banion *et al.*, 2023). Thus, in parallel with secondary metabolism, the modification of inositol metabolism in NFS plant may be involved in the differential defence process observed in infested NFS-plants compared to compared to infested NI-plants. The significant differential accumulation of 3-Glu(1-3)Glu-28-Xyl(1-4)Rha(1-2)Ara zanhic acid and 3-(4'O-Malonyl)Rha(1-2)Gal(1-2)GluA-Soyasaponenol B, two terpene molecules are also a marker of both the symbiotic state and the infection by aphids. Finally, 3-(4'O-Malonyl)Rha(1-2)Gal(1-2)GluA-Soyasaponenol B maybe one of the most interesting and intriguing metabolite in our experiment. This triterpenoid saponin is a defensive compound against pathogenic microbes and herbivores (Osbourn, 1996; Kuzina *et al.*, 2009; Szakiel *et al.*, 2011) and may act as feeding deterrents for plant specialist herbivores (Cui *et al.*, 2019) causing a cytotoxic effect in hindgut and fat body of insects (Adel & Sammour, 2012). Production of triterpenoids have also been observed as an effect of rhizobia on pea seeds infected with Didymella. Accumulation of seed terpenoid Pisumoside B was observed in uninfected rhizobial treated seeds, meanwhile, Soyasapogenol C, Api_Dai_Kae_Flavon and 6-Hydroxyapigenin 7-[6"-(3-hydroxy-3-methylglutaryl)glucoside] were significantly enhanced in infected rhizobial treated seeds in Protecta cultivar (Ranjbar Sistani *et al.*, 2017). It may represent a very interesting biological marker of the symbiosis induced defence priming. The induction of defence mechanisms is associated to reprograming of gene expression in plants (Aerts *et al.*, 2022). For example reciprocal interactions between a chewing herbivore, *Sitona lineatus* (pea leaf weevil) and *Pisum sativum* (pea) plants grown with or without rhizobia (*Rhizobium leguminosarum* biovar. *viciae)* revealed that plants grown with rhizobia had increased gene transcript expression associated with hormone-related defence (jasmonic acid, ethylene, abscisic acid) as well as physical and antioxidant-related defence, which may explain reduced feeding by *S. lineatus* (Basu *et al.*, 2022).

We have also looked at the expression of some genes involved in the secondary metabolism pathway. As expected from previous analysis, *PR1* and *PI* are differentially expressed in infested NFS plants and NI plants suggesting that different defence transcriptional reprograming occurs in the two conditions.

Looking at the genes involved in secondary metabolism synthesis pathway, transcript accumulation of *SARD4*, that encodes a key enzyme for pipecolic acid biosynthesis, and *PAL*, the first enzyme of the phenylpropanoid pathway, is not modified by aphid infestation neither in NFS-plants nor in NI-plants. This was surprising since *SARD4* was shown to be required for the establishment of systemic acquired resistance to pathogen infection in Arabidopsis (Ding *et al.*, 2016). However, *SARD4* deleted plants are still able to biosynthesize pipecolate,

this pathway may involve thus other enzyme(s). PAL is also a member of a multigenic family
and other members of this family may be induced. In contrast the isoflavonoid pathway genes
CHI, FLS/F3H, HI4'O-MT and *PTS* expression was significantly increased in infested plants:
while the induction of *CHI, FLS/F3H, HI4'O-MT* does not vary more than two fold in the two
infested growth conditions, the *PTS* expression is induced 53- and 25-fold times by aphids in
NFS and NI plants, respectively. *PTS* is involved in synthesis of pterocarpans that constitute
the second largest group of natural isoflavonoids and play an important role as phytoalexins.
In Medicago, the pterocarpans medicarpin was shown to protect the plant from the powdered
mildew *Erysiphe pisi* and activates the SA pathway (Gupta *et al.*, 2022). Medicarpin was also
shown to be accumulated in Medicago leaves upon long term or strong attack by pea aphids
(Stewart *et al.*, 2016) and also in response of infection with the fungal pathogen *Phoma
medicaginis* (Jasiński *et al.*, 2009) suggesting some large spectrum defensive role. In contrast,
medicarpin was shown to be an antagonists of nod gene expression necessary for rhizobia to
form their association with the plant roots (Zuanazzi *et al.*, 1998). Thus an increase in
medicarpin synthesis during aphid infestation could also explain in part the effect on the root
nodules previously observed (Pandharikar *et al.*, 2020). *OEPB*, a gene involved in pinitol
synthesis (Pupel *et al.*, 2019), was significantly induced in infested-NI plants and not in
infested-NFS plants. Pinitol has been shown to prolonged the pea aphid probing behaviour but
did not prevent them from feeding (Campbell & Binder, 1984; Kordan *et al.*, 2011). Pinitol has
also been involved in maintenance of the nodule osmotic balance during development (Poole
& Ledermann, 2022) and *Sinorhizobium meliloti*, may catabolize pinitol to form nodule
(Kennedy-Mendez, 2018). Thus why *OEPB* increases only in NI plant remains to be
understood.

In aphid infestation, signal transduction pathway is activated to induce related transcription
factors that regulate the expression of downstream disease resistance genes (Gao *et al.*, 2021).
Taken together, our results show that, in our conditions, aphids are able to trigger accumulation
of various metabolites and expression of defence genes responsible for SAR and SA defence
pathway as well as modify the metabolism of primary sugars in legumes. Under infestation by
pea aphids, legumes in symbiosis are able to produce a differential defence reaction by
activating the JA defence pathway and production of defence metabolites such as triterpenoid
saponins (soyasapogenol). Here, we have chosen one aphid line that develops well on the A17
Medicago truncatula cultivar used, however we know that the metabolic profiles of plants
infested with their native or non-native aphid host race could vary (Sanchez-Arcos *et al.*, 2019).
Meanwhile in these interactions the potential change induced by the presence of the rhizobium

562   is not questioned since the symbiotic status of the plant used is generally unknown. Thus our
563   results provide a foundation for the development and inclusion of symbiosis in such studies
564   and in a more applied role as a potential stimulant of plant defence against herbivores that may
565   be a new form of biocontrol in integrated pest management.
566

Acknowledgements
We are highly grateful to S. Tares-Amichot and L. Arthaud for help in aphids manipulation
and rearing and for M. Bosseno for help in Medicago plants seedling. GB is supported by a
doctoral fellowship from the department "Santé des Plantes et Environnement" of Institut
National de Recherche pour l'Agriculture, l'Alimentation et l'Environnement (INRAE) and
the Université Côte d'Azur (UniCA). The IJPB benefits from the support of Saclay Plant
Sciences-SPS (ANR-17-EUR-0007). This work has benefited from the support of IJPB's Plant
Observatory platform PO-Chem. This work was supported by UniCA, the SPE department of
INRAE and the French Government (National Research Agency, ANR) through the
"Investments for the Future" programs LABEX SIGNALIFE ANR-11-LABX-0028-01 and
IDEX UCAJedi ANR-15-IDEX-01.

Competing Interests
None declared.
581

Author contributions
GB, PF, MP, and JLG conceived the ideas. GB, PF, MP, and JLG planned and designed the
research. SB and GC performed the metabolomics analysis and analysed the data. GB and MP
performed qPCR analysis. GB and RB performed sugar and starch analysis. GB analysed all
other data. GB wrote the manuscript with inputs from PF, MP, and JLG. All authors contributed
and agreed to the final revisions.
588

ORCIDs
Goodluck Benjamin https://orcid.org/0000-0003-4849-804X
Marie Pacoud
Stéphanie Boutet https://orcid.org/0009-0002-5482-8308
Gilles Clement https://orcid.org/0000-0001-9899-2738
Renaud Brouquisse https://orcid.org/0000-0002-7818-9662

Jean-Luc Gatti https://orcid.org/0000-0001-7683-718X

Marylène Poirié https://orcid.org/0000-0002-3349-6925

Pierre Frendo https://orcid.org/0000-0002-4578-3366

Data Availability

References

Adel MM, Sammour EA. **2012**. Effect of sub-lethal dose of natural compound of Medicago sativa (L, Leguminaceae) on the hind gut and fat body of Spodoptera littoralis (Lepidoptera, Noctuidae). *Journal of Applied Sciences Research* **8**: 1398–1408.

Aerts N, Chhillar H, Ding P, Van Wees SC. **2022**. Transcriptional regulation of plant innate immunity. *Essays in Biochemistry* **66**: 607–620.

Basu S, Lee BW, Clark RE, Bera S, Casteel CL, Crowder DW. **2022**. Legume plant defenses and nutrients mediate indirect interactions between soil rhizobia and chewing herbivores. *Basic and Applied Ecology* **64**: 57–67.

Benjamin G, Pandharikar G, Frendo P. **2022**. Salicylic acid in plant symbioses: Beyond plant pathogen interactions. *Biology* **11**: 861.

Boutet S, Barreda L, Perreau F, Totozafy J, Mauve C, Gakière B, Delannoy E, Martin-Magniette M, Monti A, Lepiniec L. **2022**. Untargeted metabolomic analyses reveal the diversity and plasticity of the specialized metabolome in seeds of different Camelina sativa genotypes. *The Plant Journal* **110**: 147–165.

Burbidge CA, Ford CM, Melino VJ, Wong DCJ, Jia Y, Jenkins CLD, Soole KL, Castellarin SD, Darriet P, Rienth M. **2021**. Biosynthesis and cellular functions of tartaric acid in grapevines. *Frontiers in Plant Science* **12**: 643024.

Campbell BC, Binder RG. **1984**. Alfalfa cyclitols in the honeydew of an aphid. *Phytochemistry* **23**: 1786–1787.

Chen J, Fernandez D, Wang DD, Chen YJ, Dai GH. **2014**. Biological control mechanisms
of D-pinitol against powdery mildew in cucumber. *Physiological and molecular plant*
pathology **88**: 52–60.

Chiozza MV, O'Neal ME, MacIntosh GC. **2010**. Constitutive and induced differential
accumulation of amino acid in leaves of susceptible and resistant soybean plants in response to
the soybean aphid (Hemiptera: Aphididae). *Environmental Entomology* **39**: 856–864.

Cui C, Yang Y, Zhao T, Zou K, Peng C, Cai H, Wan X, Hou R. **2019**. Insecticidal activity
and insecticidal mechanism of total saponins from Camellia oleifera. *Molecules* **24**: 4518.

Del Giudice J, Cam Y, Damiani I, Fung-Chat F, Meilhoc E, Bruand C, Brouquisse R,
Puppo A, Boscari A. **2011**. Nitric oxide is required for an optimal establishment of the
Medicago truncatula–Sinorhizobium meliloti symbiosis. *New Phytologist* **191**: 405–417.

Desalegn G, Turetschek R, Kaul H-P, Wienkoop S. **2016**. Microbial symbionts affect Pisum
sativum proteome and metabolome under Didymella pinodes infection. *Journal of Proteomics*
143: 173–187.

Diehl S, Bush G. **1984**. An evolutionary and applied perspective of insect biotypes. *Annual*
review of entomology **29**: 471–504.

Ding P, Rekhter D, Ding Y, Feussner K, Busta L, Haroth S, Xu S, Li X, Jetter R, Feussner
I. **2016**. Characterization of a pipecolic acid biosynthesis pathway required for systemic
acquired resistance. *The Plant Cell* **28**: 2603–2615.

Drès M, Mallet J. **2002**. Host races in plant–feeding insects and their importance in sympatric
speciation. *Philosophical Transactions of the Royal Society of London. Series B: Biological*
Sciences **357**: 471–492.

Dutta S, Mishra A, Kumar BD. **2008**. Induction of systemic resistance against fusarial wilt
in pigeon pea through interaction of plant growth promoting rhizobacteria and rhizobia. *Soil*
Biology and Biochemistry **40**: 452–461.

Fiehn O. **2006**. Metabolite profiling in Arabidopsis. *Arabidopsis protocols*: 439–447.

Fiehn O. **2008**. Extending the breadth of metabolite profiling by gas chromatography coupled
to mass spectrometry. *TrAC Trends in Analytical Chemistry* **27**: 261–269.

Gao P, Zhang H, Yan H, Zhou N, Yan B, Fan Y, Tang K, Qiu X. **2021**. Transcriptomic and
metabolomic changes triggered by Macrosiphum rosivorum in rose (Rosa longicuspis). *BMC*
genomics **22**: 1–13.

Giordanengo P, Brunissen L, Rusterucci C, Vincent C, van Bel A, Dinant S, Girousse C,
Faucher M, Bonnemain J-L. **2010**. Compatible plant-aphid interactions: how aphids
manipulate plant responses. *Comptes Rendus. Biologies* **333**: 516–523.

Goławska S, Łukasik I, Chojnacki A. **2024**. Luteolin and quercetin affect aphid feeding
behavior. *The European Zoological Journal* **91**: 318–331.

Goławska S, Łukasik I, Kapusta I, Janda B. **2012**. Do the contents of luteolin, tricin, and
chrysoeriol glycosides in alfalfa (Medicago sativa L.) affect the behavior of pea aphid
(Acyrthosiphon pisum)? *Polish Journal of Environmental Studies* **21**.

Gopalakrishnan S, Sathya A, Vijayabharathi R, Varshney RK, Gowda CL,
Krishnamurthy L. **2015**. Plant growth promoting rhizobia: challenges and opportunities. *3*
Biotech **5**: 355–377.

Gupta A, Awasthi P, Sharma N, Parveen S, Vats RP, Singh N, Kumar Y, Goel A,
Chandran D. **2022**. Medicarpin confers powdery mildew resistance in Medicago truncatula
and activates the salicylic acid signalling pathway. *Molecular Plant Pathology* **23**: 966–983.

Heberle H, Meirelles GV, da Silva FR, Telles GP, Minghim R. **2015**. InteractiVenn: a web-
based tool for the analysis of sets through Venn diagrams. *BMC bioinformatics* **16**: 1–7.

Herrera-Vásquez A, Salinas P, Holuigue L. **2015**. Salicylic acid and reactive oxygen species
interplay in the transcriptional control of defense genes expression. *Frontiers in plant science*
6: 131929.

Hilliou F. **2013**. RqPCRAnalysis: analysis of quantitative real-time PCR data. In:
SCITEPRESS, 202–211.

Hiruma K, Fukunaga S, Bednarek P, Piślewska-Bednarek M, Watanabe S, Narusaka Y,
Shirasu K, Takano Y. **2013**. Glutathione and tryptophan metabolism are required for
Arabidopsis immunity during the hypersensitive response to hemibiotrophs. *Proceedings of the*
National Academy of Sciences **110**: 9589–9594.

681 **Huang T, Jander G, de Vos M**. **2011**. Non-protein amino acids in plant defense against insect
682 herbivores: representative cases and opportunities for further functional analysis.
683 *Phytochemistry* **72**: 1531–1537.

684 **Isah T**. **2019**. Stress and defense responses in plant secondary metabolites production.
685 *Biological research* **52**.

686 **Jakobs R, Schweiger R, Müller C**. **2019**. Aphid infestation leads to plant part-specific
687 changes in phloem sap chemistry, which may indicate niche construction. *New Phytologist*
688 **221**: 503–514.

689 **Jasiński M, Kachlicki P, Rodziewicz P, Figlerowicz M, Stobiecki M**. **2009**. Changes in the
690 profile of flavonoid accumulation in Medicago truncatula leaves during infection with fungal
691 pathogen Phoma medicaginis. *Plant Physiology and Biochemistry* **47**: 847–853.

692 **Kalloniati C, Krompas P, Karalias G, Udvardi MK, Rennenberg H, Herschbach C,**
693 **Flemetakis E**. **2015**. Nitrogen-fixing nodules are an important source of reduced sulfur, which
694 triggers global changes in sulfur metabolism in Lotus japonicus. *The Plant Cell* **27**: 2384–2400.

695 **Kaloshian I, Walling LL**. **2016**. Hemipteran and dipteran pests: effectors and plant host
696 immune regulators. *Journal of Integrative Plant Biology* **58**: 350–361.

697 **Kanwar P, Jha G**. **2019**. Alterations in plant sugar metabolism: signatory of pathogen attack.
698 *Planta* **249**: 305–318.

699 **Kennedy-Mendez AI**. **2018**. Characterization of Pinitol Catabolism in Sinorhizobium Meliloti
700 and its Role in Nodule Occupancy.

701 **Kim JH, Cheon YM, Kim B-G, Ahn J-H**. **2008**. Analysis of flavonoids and characterization
702 of the OsFNS gene involved in flavone biosynthesis in rice. *Journal of Plant Biology* **51**: 97–
703 101.

704 **Kim E-G, Yun S, Park J-R, Jang Y-H, Farooq M, Yun B-J, Kim K-M**. **2022**. Bio-efficacy
705 of chrysoeriol7, a natural chemical and repellent, against brown planthopper in rice.
706 *International Journal of Molecular Sciences* **23**: 1540.

Kordan B, Dancewicz K, Wróblewska A, Gabryś B. **2012**. Intraspecific variation in alkaloid
profile of four lupine species with implications for the pea aphid probing behaviour.
Phytochemistry Letters **5**: 71–77.

Kordan B, Lahuta L, Dancewicz K, Sądej W, Gabryś B. **2011**. Effect of lupin cyclitols on
pea aphid probing behaviour. *Journal of Plant Protection Research* **51**.

Kumar S. **2017**. Plant secondary metabolites (PSMs) of Brassicaceae and their role in plant
defense against insect herbivores–A review. *Journal of Applied and Natural Science* **9**: 508–
519.

Kuzina V, Ekstrøm CT, Andersen SB, Nielsen JK, Olsen CE, Bak S. **2009**. Identification
of defense compounds in Barbarea vulgaris against the herbivore Phyllotreta nemorum by an
ecometabolomic approach. *Plant physiology* **151**: 1977–1990.

Lee A, Hirsch AM. **2006**. Signals and responses: Choreographing the complex interaction
between legumes and α-and β-rhizobia. *Plant signaling & behavior* **1**: 161–168.

Liu H, Brettell LE, Qiu Z, Singh BK. **2020**. Microbiome-mediated stress resistance in plants.
Trends in plant science **25**: 733–743.

Lu H, Yang P, Xu Y, Luo L, Zhu J, Cui N, Kang L, Cui F. **2016**. Performances of survival,
feeding behavior and gene expression in aphids reveal their different fitness to host alteration.
Scientific Reports **6**: 19344.

Maag D, Erb M, Köllner TG, Gershenzon J. **2015**. Defensive weapons and defense signals
in plants: some metabolites serve both roles. *BioEssays* **37**: 167–174.

Martin B, Collar J, Tjallingii W, Fereres A. **1997**. Intracellular ingestion and salivation by
aphids may cause the acquisition and inoculation of non-persistently transmitted plant viruses.
Journal of General Virology **78**: 2701–2705.

National Research Council, Research in Agriculture, Committee on Biosciences. **1985**.
New directions for biosciences research in agriculture: High-reward opportunities.

Ng JC, Perry KL. **2004**. Transmission of plant viruses by aphid vectors. *Molecular plant*
pathology **5**: 505–511.

O'Banion BS, Jones P, Demetros AA, Kelley BR, Knoor LH, Wagner AS, Chen J-G,
Muchero W, Reynolds TB, Jacobson D. **2023**. Plant myo-inositol transport influences
bacterial colonization phenotypes. *Current Biology* **33**: 3111–3124.

Olivon F, Elie N, Grelier G, Roussi F, Litaudon M, Touboul D. **2018**. MetGem software
for the generation of molecular networks based on the t-SNE algorithm. *Analytical chemistry*
90: 13900–13908.

Osbourn AE. **1996**. Preformed antimicrobial compounds and plant defense against fungal
attack. *The plant cell* **8**: 1821.

Pandharikar G, Gatti J-L, Simon J-C, Frendo P, Poirié M. **2020**. Aphid infestation
differently affects the defences of nitrate-fed and nitrogen-fixing Medicago truncatula and
alters symbiotic nitrogen fixation. *Proceedings of the Royal Society B* **287**: 20201493.

Pawłowska A, Stepczyńska M. **2022**. Natural biocidal compounds of plant origin as
biodegradable materials modifiers. *Journal of Polymers and the Environment* **30**: 1683–1708.

Pitino M, Hogenhout SA. **2013**. Aphid protein effectors promote aphid colonization in a plant
species-specific manner. *Molecular Plant-Microbe Interactions* **26**: 130–139.

Pluskal T, Castillo S, Villar-Briones A, Orešič M. **2010**. MZmine 2: modular framework for
processing, visualizing, and analyzing mass spectrometry-based molecular profile data. *BMC*
bioinformatics **11**: 1–11.

Ponzio C, Papazian S, Albrectsen BR, Dicke M, Gols R. **2017**. Dual herbivore attack and
herbivore density affect metabolic profiles of Brassica nigra leaves. *Plant, Cell & Environment*
40: 1356–1367.

Poole PS, Ledermann R. **2022**. Maintaining osmotic balance in legume nodules. *Journal of*
Experimental Botany **73**: 8–10.

Pupel P, Szablińska-Piernik J, Lahuta LB. **2019**. Two-step d-ononitol epimerization
pathway in Medicago truncatula. *The Plant Journal* **100**: 237–250.

Ramachandran VK, East AK, Karunakaran R, Downie JA, Poole PS. **2011**. Adaptation of
Rhizobium leguminosarum to pea, alfalfa and sugar beet rhizospheres investigated by
comparative transcriptomics. *Genome biology* **12**: 1–12.

Ranjbar Sistani N, Kaul H-P, Desalegn G, Wienkoop S. **2017**. Rhizobium impacts on seed
productivity, quality, and protection of Pisum sativum upon disease stress caused by Didymella
pinodes: phenotypic, proteomic, and metabolomic traits. *Frontiers in plant Science* **8**: 1961.

Rodriguez PA, Bos JI. **2013**. Toward understanding the role of aphid effectors in plant
infestation. *Molecular Plant-Microbe Interactions* **26**: 25–30.

Sanchez-Arcos C, Kai M, Svatoš A, Gershenzon J, Kunert G. **2019**. Untargeted
metabolomics approach reveals differences in host plant chemistry before and after infestation
with different pea aphid host races. *Frontiers in Plant Science* **10**: 373118.

Shih P-Y, Sugio A, Simon J-C. **2023**. Molecular mechanisms underlying host plant specificity
in aphids. *Annual Review of Entomology* **68**: 431–450.

Simon J-C, Boutin S, Tsuchida T, Koga R, Le Gallic J-F, Frantz A, Outreman Y, Fukatsu
T. **2011**. Facultative symbiont infections affect aphid reproduction. *PLoS One* **6**: e21831.

Smigielski L, Laubach E-M, Pesch L, Glock JML, Albrecht F, Slusarenko A, Panstruga
R, Kuhn H. **2019**. Nodulation induces systemic resistance of Medicago truncatula and Pisum
sativum against Erysiphe pisi and primes for powdery mildew-triggered salicylic acid
accumulation. *Molecular Plant-Microbe Interactions* **32**: 1243–1255.

Stewart SA, Hodge S, Bennett M, Mansfield JW, Powell G. **2016**. Aphid induction of
phytohormones in Medicago truncatula is dependent upon time post-infestation, aphid density
and the genotypes of both plant and insect. *Arthropod-plant interactions* **10**: 41–53.

Stochmal A, Simonet AM, Macias FA, Oleszek W. **2001**. Alfalfa (Medicago sativa L.)
flavonoids. 2. Tricin and chrysoeriol glycosides from aerial parts. *Journal of agricultural and*
food chemistry **49**: 5310–5314.

Sulieman S, Schulze J. **2010**. The efficiency of nitrogen fixation of the model legume
Medicago truncatula (Jemalong A17) is low compared to Medicago sativa. *Journal of Plant*
Physiology **167**: 683–692.

Szakiel A, Pączkowski C, Henry M. **2011**. Influence of environmental abiotic factors on the
content of saponins in plants. *Phytochemistry Reviews* **10**: 471–491.

Taylor BN, Ostrowsky LR. **2019**. Nitrogen-fixing and non-fixing trees differ in leaf chemistry
and defence but not herbivory in a lowland Costa Rican rain forest. *Journal of Tropical Ecology*
35: 270–279.

Tegeder M. **2014**. Transporters involved in source to sink partitioning of amino acids and
ureides: opportunities for crop improvement. *Journal of Experimental Botany* **65**: 1865–1878.

Turetschek R, Desalegn G, Epple T, Kaul H-P, Wienkoop S. **2017**. Key metabolic traits of
Pisum sativum maintain cell vitality during Didymella pinodes infection: cultivar resistance
and the microsymbionts' influence. *Journal of Proteomics* **169**: 189–201.

War AR, Paulraj MG, Ahmad T, Buhroo AA, Hussain B, Ignacimuthu S, Sharma HC.
2012. Mechanisms of plant defense against insect herbivores. *Plant signaling & behavior* **7**:
1306–1320.

Wu J, Baldwin IT. **2010**. New insights into plant responses to the attack from insect
herbivores. *Annual review of genetics* **44**: 1–24.

Yamada K, Mine A. **2024**. Sugar coordinates plant defense signaling. *Science Advances* **10**:
eadk4131.

Yan J, Lipka AE, Schmelz EA, Buckler ES, Jander G. **2015**. Accumulation of 5-
hydroxynorvaline in maize (Zea mays) leaves is induced by insect feeding and abiotic stress.
Journal of Experimental Botany **66**: 593–602.

Zuanazzi JAS, Clergeot PH, Quirion J-C, Husson H-P, Kondorosi A, Ratet P. **1998**.
Production of Sinorhizobium meliloti nod gene activator and repressor flavonoids from
Medicago sativa roots. *Molecular plant-microbe interactions* **11**: 784–794.

Züst T, Agrawal AA. **2016**. Mechanisms and evolution of plant resistance to aphids. *Nature*
plants **2**: 1–9.

List of Figures
Fig. 1: Aphid infestation, more than nitrogen source, significantly alters leaf metabolite
profile. a) Heatmap from GC-MS analysis showing unsupervised hierarchical clustering of

compounds separated by both infestation and nitrogen source. b) Class abundance of
metabolites from both GC-MS and LC-MS analyses.

Fig. 2: Metabolite distribution according to nitrogen source and aphid infestation affects
the metabolite profile of plants. Venn diagram showing distribution of statistically significant
metabolites across the various experimental conditions. The name of the compounds and the
statistic can be found in Dataset S3.

Fig. 3: Aphid infestation triggers accumulation of soluble sugars in both NFS and NI
plant leaves. Bar graphs from GCMS analysis showing the significant accumulation of glucose
sucrose and fructose in NFS and NI plants upon aphid infestation. Metabolites analysed and
quantified in µg mg^{-1} of fresh weight (FW) of leaves of NFS and NI plants 12 days after
infestation by the aphids (control = no aphid infestation). Mean ± SE (n=4): different letters
indicate a $p \leq 0.05$ (Tukey's test).

Fig. 4: Aphid infestation affects sugar concentration in leaves but not in roots.
Biochemical analysis showing the significant accumulation of glucose, sucrose, and fructose
in leaves of NFS and NI plants. No significant difference was found in roots upon aphid
infestation. No statistical difference was observed in leaves starch, measured as glucose
equivalent, between the different conditions. Metabolites analysed and quantified in µmol g^{-1}
of fresh weight (FW) of leaves and roots from NFS and NI plants 12 days after infestation by
the aphids (control indicates no aphids infestation). Mean± SE (n=4); different letters indicate
a $p \leq 0.05$ (Tukey's test).

Fig. 5: Aphid infestation triggers accumulation of defence-related metabolites in both
NFS and NI particularly from the salicylic acid defence pathway. Graphs show
accumulation of metabolites expressed as arbitrary unit (in mg apigenin equivalent mg^{-1}) of
leaves fresh weight (FW) of NFS and NI plants 12 days after infestation by the aphids (control
indicates no aphids infestation). Mean± SE (n=4); different letters indicate a $p \leq 0.05$ (Tukey's
test).

Fig. 6: NFS induces a differential defence mechanism upon aphid infestation. Graph
showing significantly accumulated metabolites in NFS upon aphid infestation. Metabolites
expressed as arbitrary unit (in mg apigenin equivalent mg^{-1}) of leaves fresh weight (FW) of

NFS and NI plants 12 days after infestation by aphids (control indicates no aphids infestation).

Mean± SE (n=4); different letters indicate a $p \leq 0.05$ (Tukey's test).

List of Tables

Table 1: Gene expression analysis of NFS and NI plants. Rescaled qPCR expression

showing the level of gene induction upon aphid infestation in NFS and NI plants (NFS_Amp

and NI_Amp, respectively) compared to their control (NFS and NI, respectively). Data are

expressed as mean ± standard error (SE); t-test on all genes, $P \leq 0.05$, significant.

Supporting Information

Datasets available as: Benjamin, Goodluck, 2024, "Supplementary Data for Differential

induction of Medicago truncatula defence metabolites in response to rhizobial symbiosis and

pea aphid infestation", https://doi.org/10.57745/PCDG8H, Recherche Data Gouv, V1,

UNF:6:A0SbGGbxQocqa6JsmMuoMA== [fileUNF]

-Dataset S1: File containing qPCR Data (qPCR CT Data.xlsx)

-Dataset S2: File containing GCMS and LCMS metabolomics data (Metabolomics_data.xlsx)

-Dataset S3: File containing Significant accumulated metabolites analysed by Anova and Venn

Analysis (Significant metabolites_Venn.xlsx)

Fig. S1: Experimental setup showing timeline leading up to assay.

Fig. S2: Experiment treatment classification showing NFS_Control, NFS_Amp, NI_Control

(KNO$_3$) and NI_Amp (KNO$_3$).

Fig. S3: Abundance of quantified metabolites from GCMS analysis showing accuracy by

comparing the mean obtained from the quality controls (QC, blue bars) and mean obtained

from the samples (Samples, red bars).

Fig. S4: Heatmap showing top 100 features from LCMS analysis.

Table S1: List of genes and oligos sequences used for RT-qPCR analysis.

Differential induction of *Medicago truncatula* defence metabolites in response to rhizobial symbiosis and pea aphid infestation

Goodluck BENJAMIN[1], Marie PACOUD[1], Stéphanie BOUTET[2], Gilles CLEMENT[2], Renaud BROUQUISSE[1], Jean-Luc GATTI[1], Marylène POIRIÉ[1*], Pierre FRENDO[1*]

1   Université Côte d'Azur, INRAE, CNRS, Institut Sophia Agrobiotech, 06903 Sophia Antipolis, France
2   Université Paris-Saclay, INRAE, AgroParisTech, Institut Jean-Pierre Bourgin for Plant Sciences (IJPB), 78000 Versailles, France

*Both authors supervised this work.

 Correspondence: pierre.frendo@univ-cotedazur.fr; goodluck.benjamin@etu.univ-cotedazur.fr

930
931

Supplementary Data

Datasets available as: Benjamin, Goodluck, 2024, "Supplementary Data for Differential induction of Medicago truncatula defence metabolites in response to rhizobial symbiosis and pea aphid infestation", https://doi.org/10.57745/PCDG8H, Recherche Data Gouv, V1, UNF:6:A0SbGGbxQocqa6JsmMuoMA== [fileUNF]

Dataset S1: File containing qPCR Data (qPCR CT Data.xlsx)

Dataset S2: File containing GC-MS and LC-MS metabolomics data (Metabolomics_data.xlsx)

Dataset S3: File containing Significant accumulated metabolites analysed by Anova and Venn Analysis (Significant metabolites_Venn.xlsx)

973 **Fig. S1: Experimental setup.** Diagram showing timeline leading up to assay with key date

974 points for bacteria inoculation and aphid infestation marked up. Each pot contained six plants

975 and 4 pots were made for each condition per biological replicate. Assay was performed using

976 4 biological replicates produced using the same timeline.

1000
1001
1002
1003

1004

1005 **Fig. S2: Experiment treatment classification.** Diagram showing the classification of studied

1006 treatments into 4 groups, NFS_Control, NFS_Amp, NI_Control (KNO_3) and NI_Amp (KNO_3).

1007

1008

1009

1010

1011

1034
1035
1036 **Fig. S3: Abundance of quantified metabolites.** Graph of quantified metabolites from GC-
1037 MS analysis showing accuracy of abundance by comparing the mean obtained from the quality
1038 controls (QC, blue bars) and mean obtained from the samples (Samples, red bars).

Fig. S4: Nitrogen source significantly modifies the leaf metabolite profile. Heatmap of 2-way Anova analysis from GC-MS showing unsupervised hierarchical clustering of compounds separated by nitrogen source. NFS = Nitrogen fixing symbiosis, KNO3 = Non inoculated, Amp = aphid infested, no aphids = Control.

1068

Fig. S5: Aphid infestation significantly modifies the leaf metabolite profile. Heatmap of 2-

way Anova analysis from GC-MS showing unsupervised hierarchical clustering of compounds

separated by aphid infestation. NFS = Nitrogen fixing symbiosis, KNO3 = Non inoculated,

Amp = aphid infested, no aphids = Control.

1074

1075

1076

Fig. S6: Unsupervised clustering of LC-MS compounds. Heatmap showing the top 100

features from LC-MS analysis.

1077

1078

1079

1080

1081

1082

1083

1084

1085 **Fig. S7: NFS and Aphid infestation significantly modifies the leaf metabolite profile.**

1086 Heatmap of 2-way Anova analysis from GC-MS showing unsupervised hierarchical clustering

1087 of compounds separated by the interaction of NFS and aphid infestation. NFS = Nitrogen fixing

1088 symbiosis, KNO3 = Non inoculated, Amp = aphid infested, no aphids = Control.

1089
1090
1091
1092

1093   **Table S1: List of genes and oligos sequences used for RT-qPCR analysis.**

Description	Name	Genomic ID	Forward primer	Reverse primer	References
Chalcone Isomerase	*CHI*	MtrunA17_Chr1g0213011	CCTGAAAAGGAGGCTGCACT	ACAGCGCTTAAGATCAGGGG	This work
Flavanol Synthase/Flavanone 3-Hydroxylase	*FLS/F3H*	MtrunA17_Chr3g0092531	CACAATTCACCAAAGAGATTGGGG	ATCCTCTGGCCTAGAAGGTGG	This work
Isoflavone 4'-O-methyltransferase	*HI4'O-MT*	MtrunA17_Chr4g0046341	GCAACAGGGGAGAGTTTTTGG	AGACTCCAAACCCTCGAAAACG	This work
D-pinitol Dehydrogenase	*OEPB*	MtrunA17_Chr6g0480011	GCACTCAGATGGGTGTACCA	GCACTCAGATGGGTGTACCA	This work
Phenylalanine Ammonia-lyase	*PAL*	MtrunA17_Chr1g0181091	AGCGCTTATGTTAAAGCCGC	AGCGCTTATGTTAAAGCCGC	This work
Pterocarpan Synthase 1	*PTS*	MtrunA17_Chr7g0259091	AAGGAGATGCCATTGTGGAG	CCTATTCGTACTATAGGTGAGAGTG	This work
SAR Deficient 4	*SARD4*	MtrunA17_Chr1g0202471	CCCCAATTCGCCAACACTAC	CAGGGAAATGGGTCACGAGT	This work
Pathogenesis Related Protein-1	PR1	MtrunA17_Chr2g0295371	TTCGGGTTGGATGTGCTAAG	GGTTGAAGCTCAATGGCACT	Pandharikar *et al.*, 2020
Proteinase Inhibitor	*PI*	MtrunA17_Chr4g0014461	TGTGGTGCAATTCTTTCAGG	ATTTTGGGGTGAGGTGTTGA	Pandharikar *et al.*, 2020
Housekeeping gene	*MtC27*	MtrunA17_Chr2g0295871	TGAGGGAGCAACCAAATACC	GCGAAAACCAAGCTACCATC	Del Guidice *et al.*, 2011
Housekeeping gene	*a38*	MtrunA17_Chr4g0061551	TCGTGGTGGTGGTTATCAAA	TTCAGACCTTCCCATTGACA	Del Guidice *et al.*, 2011

1094