Large protein databases reveal structural complementarity and functional locality

Paweł Szczerbiak[1,2], Lukasz M. Szydlowski[1,2], Witold Wydmański[2,3],

P. Douglas Renfrew[4], Julia Koehler Leman[5], Tomasz Kosciolek[1*]

1. Sano Centre for Computational Medicine, Kraków, 30-054, Poland

2. Małopolska Centre of Biotechnology, Jagiellonian University, Kraków, 30-387, Poland

3. Faculty of Mathematics and Computer Science, Jagiellonian University, Kraków, 30-348, Poland

4. Center for Computational Biology, Flatiron Institute, New York, NY 10010, USA

5. Open Molecular Software Foundation, Davis, CA 95616, USA

*Corresponding author: t.kosciolek@sanoscience.org

Abstract

Recent breakthroughs in protein structure prediction have led to an unprecedented surge in high-quality 3D models, highlighting the need for efficient computational solutions to manage and analyze this wealth of structural data. In our work, we comprehensively examine the structural clusters obtained from the AlphaFold Protein Structure Database (AFDB), a high-quality subset of ESMAtlas, and the Microbiome Immunity Project (MIP). We create a single cohesive low-dimensional representation of the resulting protein space. Our results show that, while each database occupies distinct regions within the protein structure space, they collectively exhibit significant overlap in their functional profiles. High-level biological functions tend to cluster in particular regions, revealing a shared functional landscape despite the diverse sources of data. To facilitate exploration and improve access to our data, we developed an open-access web server. Our findings lay the groundwork for more in-depth studies concerning protein sequence-structure-function relationships, where various biological questions can be asked about taxonomic assignments, environmental factors, or functional specificity.

Introduction

The tectonic shift that has occurred in structural and molecular biology thanks to the abundance of high-quality protein 3D structure models calls for a new outlook on protein structures as a source of information about biology and a tool to understand it further. Thanks to decades of work on protein sequences in the field, efficient methods to store, search, cluster or annotate sequences exist and are well-documented (1–6). Since the release of AlphaFold2 (7) and accompanying AlphaFold Protein Structure Database (AFDB), the number of readily available protein structures grew 1,000-fold from ~200k to ~200M (8, 9). These data found its way into UniProt and other repositories, where they are accessible through text-based interfaces. Still, just as protein sequence information became ubiquitous and purpose-made methods to leverage this source of biological data became necessary, there is a need for purpose-built tools and a better understanding of the protein structure space. So far, we have seen a rapid development of efficient tools for protein structure processing, e.g. clustering (10, 11), compression (12, 13) or representation (14–16). Subsequently, the ESMAtlas repository (17), featuring over 600 million predictions, further expanded the available structural models, bringing the total, including those from the Protein Data Bank (18), to nearly 1 billion structures or models. Much effort has been put so far into building a deeper understanding of the AFDB structure space ranging from novel fold search (19, 20), domain identification (21), clustering (22), and structure space exploration (23). But, we still lack a comprehensive and complementary understanding of the protein structure space and how different sources of structure data – obtained using different methods, each utilizing various underlying sequence databases – contribute to this space.

In this study, we analyze three large protein structure datasets originating from distinct sources. The AlphaFold Protein Structure Database (AFDB) (9), based on UniProt (24), includes protein models from a wide range of organisms, with a significant representation of eukaryotes. The ESMAtlas (17), derived from MGnify (25), focuses exclusively on proteins from metagenomic studies, and therefore predominantly contains prokaryotic data. The Microbiome Immunity Project (MIP) database (26), based on The Genomic Encyclopedia of Bacteria and Archaea (GEBA) (27), primarily consists of short, single-domain proteins ranging from 40 to 200 residues, representing bacterial genomes, and may thus have similarities with ESMAtlas. In contrast, both the AFDB and ESMAtlas datasets include single- and multi-domain proteins, distinguishing them from the MIP dataset. This diversity allows us to explore various regions of the structural space more thoroughly, such as the localization of high- and low-quality (including intrinsically disordered proteins, IDP) predictions or single- and multi-domain proteins (see Methods for further discussion). The representative sequences and structures are then projected onto a lower-dimensional space according to their structural features and functionally annotated using an updated version of a structure-based function prediction method deepFRI (28).

In this work, we present a comprehensive analysis of the current protein structure space, integrating data from the most extensive sources available. We show that the protein structure databases are complementary and exhaustive in terms of folds and biological function. We examine the unique characteristics of each database, analyze the distribution of protein functions within this space, and introduce a web server that allows for customized exploration of these structural datasets (https://protein-structure-landscape.sano.science/). This work aims to provide a deeper understanding and contextualization of the current protein structure landscape, enabling further insights into protein function and evolution.

Results

Datasets occupy complementary regions in the structure space

Our workflow is outlined in Fig. 1A. To create a comprehensive dataset of protein structures we focused on non-redundant sequences from the major protein structure databases: AFDB50, highquality_clust30 (a high-quality, representative subset of ESMAtlas), and MIP (see Methods for details). In the first stage, we eliminated structural redundancy with Foldseek, optimizing parameters for each database individually (see section "Structural clustering" in the Supplement). For AFDB, we relied on the results from reference (22), which categorized the structural models into light (mapping onto Pfam) and dark clusters (without hits to Pfam). These clusters include both low- and high-quality models, whereas the highquality_clust30 dataset from ESMAtlas contains only high-quality predictions. Subsequently, all representative structures (excluding AFDB and ESMAtlas singletons – see Methods) were gathered into one set and clustered again with Foldseek to remove structural redundancy between databases (see Figs. S1-S18 and Table S1 in the Supplement for details as well as Table S2 for a summary). Next, we functionally annotate the dataset using deepFRI and generate Geometricus representations to explore structural features (14, 28). deepFRI has proven to be a reliable method for generating functional annotations, while Geometricus embeds protein structures into fixed-length 'shape-mer' vectors that can be successfully used in downstream tasks (29–31). Shape-mer features are then reduced to a two-dimensional structure space using PaCMAP (32). To assess cluster heterogeneity, we define heterogeneous clusters as those containing models from at least two distinct databases. Finally, all information is integrated into a web server.

Fig. 1B highlights the key features of the structure space, with all structural representatives (clusters and singletons) as inputs. For more details on the construction, refer to Figs. S19-S25. We also embedded CATH superfamilies (33) into this space to better understand its organization. Because our dataset includes all proteins from AFDB and ESMAtlas (with only MIP containing mostly single-domain structures), the CATH superfamilies only partially cover this space. Despite this, a clear separation between CATH classes is evident, with the highest density in the alpha/beta region. Except for two small areas in the structure space, we did not observe a gradient in structure length. Additionally, we found a strong correlation between structure length and mean pLDDT (as shown in Fig. 1B, AFDB pLDDT panel; see also Fig. S26 for dispersion measures).

Fig. 1C compares the fractions of structures from different databases. For a fair comparison with ESMAtlas, we included only high-quality AFDB models. We observed clear complementarity between AFDB, ESMAtlas, and MIP clusters in their coverage of the structure space (we call this *structural complementarity*). The separation between structures from different databases cannot be explained by structure length alone, as most AFDB and ESMAtlas structures are of similar size (see Fig. S17). Interestingly, ESMAtlas and AFDB light proteins largely occupy the same region of the structure space (see Fig. S27 for absolute values and Fig. S28 for all AFDB models), suggesting that both databases extensively cover the known structural landscape. Additionally, we see significant overlap between AFDB light and dark clusters.

Figure 1: Pipeline and structure space. (A): The diagram illustrates the pipeline from top to bottom. We begin from three sets of proteins that are non-redundant at the sequence level and aim to cluster them at the structure level. Red asterisk indicates the part that has been done in reference (22). Using a similar methodology, we clustered the remaining two databases and, in the second stage, we

clustered the cluster representatives (green box) to eliminate structural redundancy between databases. All structures have been annotated with deepFRI and Geometricus to elucidate its functional potential and visualize the structure space, respectively. Dimensional reduction into 2 dimensions has been performed with PaCMAP. Subsequently, we analyzed cluster heterogeneity. **(B):** CATH class (left), number of structures (middle left), protein length (middle right), and pLDDT for AFDB structures (right) distributed across the structure space (we considered all ~1.5 million cluster representatives + singletons). (**C**): Fraction of structures coming from a given database (we included only high-quality models i.e. ~3 million structures - see Table S2). Panels **B** and **C** generated using unnormalized Geometricus representations may be found in the Supplement (see Figs. S29-S32).

Protein conformations display gradual shifts across the structural landscape

We selected several representative models from different regions of our dataset to illustrate the structural landscape (Figure 2). Selected models primarily have high mean pLDDT scores (except for a few, such as A0A498NE77 with a mean pLDDT of 69, and five proteins marked in blue, coming from clusters with long representative structures (14–16)compare Fig. 5E).

The spatial distribution of the structures highlighted in red indicates the presence of two primary branches that emerge from a central area, predominantly composed of alpha-beta structures. These branches extend into beta-only structures on the left side of the figure and alpha-only structures, which are mainly located on the right and lower-right sections. The long representative structures, shown in blue and primarily consisting of low-quality AFDB models that may include IDPs, are located in the bottom-left corner, where significant divergence and continuity are also observed. However, their *in silico* characterization poses difficulties for structure-based annotation tools due to the lack of such information regarding IDPs. Overall, the structural landscape exhibits a high degree of coherence, with gradual and incremental variations in structural motifs throughout the space, although some outliers are present.

A particularly interesting region is located in the top left (green dots), characterized by fibril proteins with varying cross-sections. These include square (MGYP000417476362, A0A3G3K1M9), flat (A0A0F6LHQ3, A0A498NE77), irregular (MGYP001600570585, A0A7T9AEB2), and even triangular (P16466) cross-sections, which may be responsible for diverse functional roles. The proteins clustered in this region, often involved in pore-forming activity and transmembrane transport, are of particular medical interest due to their association with pathogenicity and virulence, such as hemolysin (P16466). Other fibrillar proteins in this area are associated with translation (A0A3G3K1M9), signaling and defense responses (A0A0F6LHQ3), macromolecule interactions (A0A498NE77), and kinase activity in response to stimuli (A0A7T9AEB2).

Figure 2: Structural landscape. Example structures within the protein structure space are represented by red, green, and blue dots. Red dots illustrate structures that are distributed relatively evenly across the space. Green dots, located in the upper left corner with corresponding structures shown from two different perspectives at the top, represent fibril proteins with various cross-sections; these examples were selected with high predicted alignment error to ensure confidence in their overall geometry. Blue dots, with structures displayed at the bottom, represent clusters with long representative structures (see also Fig. 5E, where structures are colored by pLDDT). For easier navigation, blue and green dots are labeled with letters.

High-level functional categories are localized in specific regions of the structure space

We performed functional analysis using deepFRI versions 1.0 and 1.1 (see Fig. S33) (28). While both versions share the same architecture, version 1.1 is trained on AFDB structures. This results in a higher number of predictions compared to version 1.0 but also increases the likelihood of false positives (see "Functional annotations / deepFRI v1.1" subsection in the Supplement). To map Gene Ontology (GO) predictions from deepFRI onto Clusters of Orthologous Groups (COG), we employed a customized GO-to-COG mapping (see Methods for details) (34, 35). We further combined COGs into three main functional groups: "Cellular processes and signaling" (COGs: D,M,N,O,U,W,X,Y,Z), "Metabolism" (COGs: C,E,F,G,H,I,P,Q) and "Information storage and processing" (COGs: A,B,J,K,L). Functions that were too generic and did not fall into any other category, were classified as "General Function" (COG: R). Validation on the *E.coli* proteome helped establish optimal deepFRI score thresholds (Figs. S34-S35).

The combined functional groups, hereby termed as superCOGs (superCOG 1: "Cellular processing and signaling", superCOG 2: "Metabolism", superCOG 3: "Information storage and processing") had functionally annotated proteins from five datasets from Fig. 1C and MIP novel folds as a subset of the entire MIP dataset into superCOG categories, which reveals similar superCOG distributions across the datasets, including smaller sets like MIP novel folds (Fig. 3A). Despite representing previously unseen folds, the MIP novel folds dataset does not consist of unknown functions, as most of these proteins are annotatable. Similar observations were made in AFDB dark clusters, where 67-80% of proteins were annotated. SuperCOG 3 functions (related to information storage and processing, including replication and gene expression) are less abundant in these datasets compared to others. This is expected since these processes are well-conserved and studied across living organisms (36, 37).

In Fig. 3B, we present the percentage of proteins annotated with deepFRI v1.0 for various SuperCOG functional categories. The data reveal that proteins belonging to distinct high-level functional groups are concentrated in specific regions of the structural space, a phenomenon we refer to as *functional locality*. This non-random distribution of normalized functional categories is supported by Moran's I statistic, which indicates significant spatial autocorrelation (p-value close to zero; see Fig. 3B).

DeepFRI v1.1 achieves 100% annotation coverage of all proteins, but due to being trained on the Uniprot dataset, it predicts superCOG 3 functions with less accuracy than deepFRI v1.0, which was trained on Swiss-Prot. In contrast, superCOG 2 and general function categories show strong agreement between versions 1.0 and 1.1 (compare Fig. 3B and S36). We can notice complementarity in superCOG 1+3 annotation whereas regions not annotated by v1.0 are mostly predicted by v1.1 as superCOG 1+2. Regions representing the general function category, often associated with low-quality predictions or disordered proteins (compare Fig. 1B), are located in the same corner, suggesting that generally this cluster is hard to predict, regardless of the annotation tool. Overlaying functions with CATH classes, we observed that the alpha/beta CATH class primarily overlaps with metabolic functions (superCOG 2). This is consistent with the higher structural complexity of enzymatic proteins, which are often determined by their substrates (38, 39). The largest COG categories for each dataset are shown in Fig. S43.

Figure 3: Distribution of functional categories within datasets and the structure space. (A): deepFRI annotation of high-quality structural models in each database based on superCOG aggregation. For deepFRI v1.0 and v1.1, annotations with score ≥ 0.3 and ≥ 0.5 were considered, respectively. **(B):** Normalized contribution of each superCOG category for deepFRI v1.0 (see Fig. S36 for deepFRI V1.1 predictions). All panels have been generated using normalized Geometricus vectors for all ~1.5 million cluster representatives + singletons (versions showing absolute values as well as based on unnormalized Gemetricus representations can be found in the Supplement – Figs. S37-S38

and Figs. S39-S42 respectively). Moran's I coefficient and its corresponding p-value are displayed in each panel.

Exploring functional diversity in novel protein folds

In gene ontology (GO) annotation database, the functional hierarchy is structured as an ontology tree, where Shannon Information Content (IC) serves as a proxy for measuring the specificity of annotations (40). In our analysis of MIP novel fold clusters, we identified several proteins with annotations characterized by high IC values, indicating they are associated with more specific GO terms. A higher IC reflects greater information entropy within the GO database, meaning these proteins are annotated with terms that offer more detailed functional insight (see also (29)). This suggests that novel folds can still correspond to known functions (Fig. 4). We hereby selected three examples of such proteins that – despite high IC – do not have structural homologs: proteins with tetrahydromethanopterin S-methyltransferase activity (GO:0030269), intramembrane lipid transporter activity (GO:0140303), and pectinesterase inhibitor activity (GO:0046910). The specificity of these GO terms suggests new structural alternatives for the variety of functions, including core metabolism (GO:0030269 and 00140303) and cellular processing (GO:0046910).

Tetrahydromethanopterin S-methyltransferase (Fig. 4A) forms catalytic complexes with cobalamin (vitamin B12) and transfers methyl groups from tetrahydromethanopterin (H4MPT) to coenzyme M. The gene encoding the MtrA subunit of this complex, mtrA, exhibits significant phylogenetic diversity (41, 42). The ability of these proteins to bind complex substrates, such as H4MPT, and form complexes with other proteins likely drives their structural diversity (41–44). The protein we identified was derived from the genome of *Blautia* sp., a known methanogen, which strengthens our confidence in this prediction. A Foldseek search revealed the highest similarity to another *Blautia* protein in the AFDB database (TM-score = 0.99, Fig. 4A), while alignments with other known tetrahydromethanopterin S-methyltransferases yielded poor matches (TM-score = 0.47).

Another example (Fig. 4B) involves intramembrane lipid transporters, a diverse group of proteins that move lipids within the membrane. These proteins exhibit structural diversity, with transmembrane domains ranging from a few to over a dozen, and they may function as flippases, floppases, or scramblases (45, 46). This diversity is further influenced by the unique lipid profiles of different cell membranes. The MIP_00055961 sequence was derived from the genome of *Ectobacillus panaciterrae* DSM 19096, a recently classified species with a distinct cellular fatty acid profile compared to other *Bacillales*. Its closest structural match (TM-score > 0.8, despite only 20% sequence similarity) is a RRXRR domain-containing protein from the cyanobacterial organism *Symploca* sp.

The third example (Fig. 4C) concerns pectinesterase inhibitors (PMEIs), extracellular enzymes that inhibit pectin methylesterases (PMEs) and regulate cell wall modification. PMEIs can act against endogenous PMEs or target enzymes from other species in symbiotic or host-pathogen interactions, leading to significant structural variability (47–49). MIP_00248745, derived from *Gordonia bronchialis* DSM 43247, a known opportunistic human pathogen, might function as a PMEI, potentially influencing glycoprotein formation in human lung mucus. This protein shows high similarity to a protein from *Gordonia insulae* (TM-score = 0.69, with 48% sequence similarity), as well as proteins from *Nocardia brasiliensis* and *Mycobacterium tuberculosis*, to which *G. bronchialis* is related.

	A	**B**	**C**
	MIP_00230606	MIP_00055961	MIP_00248745
Organism:	*Blautia* sp.	*Ectobacillus panaciterrae*	*Gordonia bronchialis*
deepFRI score:	**0.80**	**0.56**	**0.99**
Information content:	13.8	13.1	12.1
Max TM-score to PDB:	**0.47**	**0.49**	**0.48**
Max TM-score to AFDB:	**0.99**	**0.80**	**0.69**
GO-term:	GO:0030269	GO:0140303	GO:0046910
GO-name:	tetrahydromethanopterin S-methyltransferase activity	intramembrane lipid transporter activity	pectinesterase inhibitor activity
COG:	**H** (vitamin & cofactor metabolism)	**I** (lipid metabolism & transport)	**T** (signal transduction)
superCOG:	**S2**	**S2**	**S1**

Figure 4: MIP novel fold case study. Functional overview of MIP novel folds. deepFRI v1.1 has been used for function annotations.

Protein space is dominated by heterogeneous structural clusters

Figure 5 provides an overview of cluster heterogeneity across different databases. Panel A shows the total number of models in each database, categorized into singletons, homogeneous clusters, and heterogeneous clusters. A significant portion of AFDB clusters is of low or moderate quality (pLDDT ≤ 70), with 38% for AFDB light and 53% for AFDB dark (red bars). This can be attributed to the structural redundancy within AFDB, which was reduced using Foldseek, leaving behind high-quality protein building blocks and many poorly predicted or intrinsically disordered proteins (see also Fig. S18). As a result, these lower-quality models overshadow the high-quality AlphaFold models in the complete AFDB set.

Interestingly, applying a different procedure to AFDB – filtering after clustering instead of clustering after filtering, as with ESMAtlas – resulted in a similar proportion of singletons, homogeneous clusters, and heterogeneous clusters for both AFDB and ESMAtlas. Moreover, most AFDB and ESMAtlas structures combine to form heterogeneous clusters (Fig. 5A). Among all high-quality AFDB structures that cluster with ESMAtlas (or with both ESMAtlas and MIP), only 13% (or 11%) are annotated as dark. In contrast, 31% of all clustered AFDB structures are dark, dropping to 26% when considering only models with a mean pLDDT > 70. These observations quantitatively support the conclusions from the structure space analysis in Fig. 1C, indicating that there is a greater overlap between light AFDB and ESMAtlas compared to the overlap between dark AFDB and ESMAtlas.

Panel B shows that in heterogeneous clusters, the percentage of superCOG 3 proteins decreases with cluster size, while the percentages of superCOG 1 and 2 proteins increase. This diversity can be explained by the fact that superCOG 1 and 2 proteins are subject to fewer constraints than superCOG 3 proteins, whose structures are largely determined by

nucleic acid and ribosome conformation. Conversely, the increase in superCOG 3-related proteins in homogeneous ESMAtlas data (Fig. S46) is likely due to the presence of phage/viral proteins derived from metagenomic samples. Additionally, the percentage of unannotated proteins decreases with cluster size, indicating that larger clusters are better characterized functionally. The percentage of general functions also varies between AFDB + ESMAtlas clusters and AFDB + ESMAtlas + MIP clusters, partially influenced by mixed superCOG categories (not shown in the plot) that may express more general annotations. In this analysis, only cluster centroids were compared, so we also checked if the trends hold for functionally homogeneous clusters – and they do (see Fig. S44). Similar behavior is observed in homogeneous clusters (see Figs. S45-S46), though changes in superCOG 2 are less pronounced. These observations suggest that while all databases are large enough to display similar functional potential at a general level, there is still considerable diversity within specific clusters.

In Panel C, we see that for AFDB homogeneous clusters, there is a positive correlation between cluster size and representative structure length. This trend is not observed in other databases, although a weaker correlation exists for heterogeneous clusters containing AFDB structures (see Figs. S47-S50 for details). Panel D shows that the percentage of singletons rapidly decreases with increasing structure size, regardless of the database.In AFDB, a significant portion of long protein tails (>1300 amino acids) fail to cluster with others, likely due to a higher density of low-quality predictions and intrinsically disordered proteins, as well as statistical bias (e.g., the scarcity of long structures in ESMAtlas and AFDB). Finally, Panel E illustrates the number of structures in clusters composed of long proteins, which almost exclusively come from AFDB. Most of these clusters are small, and only 31% have centroids with a mean pLDDT higher than 70. The reason for the larger proteins in AlphaFold clusters could be the variation in organisms across datasets: ESMAtlas (and MIP) consist of prokaryotic proteins, whereas AFDB, based on UniProt, also includes eukaryotic proteins. As prokaryotic proteins are generally shorter than the eukaryotic counterparts across all functions, the higher proportion of eukaryotic structures in the AFDB could result in such bias (50).

3D structure visualizations of the largest representatives, forming both homogeneous and heterogeneous clusters, along with functional annotation coverage of different cluster types, are presented in Figs. S53-S54, accompanied by a detailed discussion. An interactive version of Fig. S54 is available on the paper's GitHub repository (see Data Availability section).

Figure 5: Cluster heterogeneity. (A): Number of structures (clusters + singletons) stratified by database type and cluster group (singletons, homogeneous and heterogeneous clusters respectively). AFDB structures have been split into low- and high-quality groups. **(B):** SuperCOG category of representative structures vs cluster size for the two largest heterogeneous cluster groups i.e. AFDB + ESMAtlas and AFDB + ESMAtlas + MIP (corresponding plots for homogeneous clusters as well as for

all structures building highly functionally homogeneous clusters may be found in the Supplement – Figs. S44-S46). **(C):** Dependance between representative structure length and cluster size for homogeneous clusters (corresponding plots for heterogeneous clusters may be found in the Supplement – Figs. S47-S50). Number of clusters with a given cluster size is plotted as lines on the right vertical axis. **(D):** Number of singletons for each database type versus structure length. Ratio between the number of singletons and the total number of structures is depicted in red (see right vertical axis). **(E):** Distribution of clusters (mostly coming from the AFDB with two exceptions i.e. ESMAtlas structures depicted in orange) with long representative structures (≥ 1300 residues). Examples of AFDB structures are also displayed (see also Fig. 2). The number next to each structure represents the mean pLDDT score: values ≤ 50 are shown in red, and those >50 in green. In **B – D** only high-quality structures have been considered. In **B** and **C** only structure representatives shorter than 1300 residues have been included.

Web server

Figure 6 illustrates the core features of the interactive web server. Users can search for structures by ID or click on them directly. Once a structure is selected, detailed information is provided, including SuperCOG predictions from deepFRI v1.0 and v1.1, protein length, and database origin, along with a 3D visualization. The server supports filtering by database origin, SuperCOG category, protein length, and AFDB pLDDT, which is particularly helpful for identifying disordered proteins. For instance, in Figure 6, structures from all databases were selected, but only high-quality AFDB models (pLDDT > 70) with a length < 1000 aa, and SuperCOG 1 annotations from deepFRI v1.0, were shown.

Figure 6: Web server interface. The central section (1) displays protein structures, with different colors representing distinct datasets (gray: AFDB dark clusters, blue: AFDB light clusters, green: ESMAtlas clusters, red: MIP clusters, light red: MIP singletons). Users can search for structures by name (2) or select them via mouse click, triggering the display of an information panel (3) along with a PyMOL 3D visualization (4). AFDB structures can be filtered by pLDDT score (5), while all structures can be filtered by length (6), superCOG category (7), and database origin (8).

Discussion

It has been already demonstrated that the protein structure space is continuous and that the Protein Data Bank contains sufficient structural information to understand the protein structure landscape (18, 51–54). This statement is supported by the fact that AlphaFold, trained exclusively on PDB structures, succeeded in predicting structures for the vast majority of the UniProt proteins. In this work, we show that when considering clusters of structures derived from gene-coding proteins in the largest available protein structure databases, the structure space is highly diverse yet protein conformations display gradual shifts across the space, resulting in a continuous structural landscape (Fig. 1). We also observe clear complementarity between these databases: AFDB and ESMAtlas clusters, although greatly overlap, cover different locations of the structure space on a similar level (Fig. 1C). MIP domains (clusters and singletons) complement ESMAtlas cluster representatives but we might speculate if inclusion of ESMAtlas and AFDB singletons would change the picture. At this point, we decide not to include singletons in the analysis due to computational and analytical constraints, though this remains a promising direction for future work. Another potential avenue is to split all input structures into domains, evolutionary conserved structural building blocks, before clustering. This approach would provide a different perspective on the protein structure space, highlighting similarities between AFDB and ESMAtlas at the domain level (19, 21).

Functional analysis revealed that all considered datasets possess similar high-level functional profiles (on superCOG level – Fig. 3A). Nevertheless, we can observe certain differences on a cluster level (see Fig. 5B). It appears that novel folds do not necessarily indicate unknown functions (Fig. 4). To accurately determine their functional capabilities, it is essential to utilize advanced annotation tools that do not rely solely on homology. deepFRI offers a great advantage to homology-independent functional annotation, which was demonstrated in a number of papers (22, 23, 29, 30, 55, 56). Furthermore, the differences between deepFRI v1.0 and v1.1 create a complementary advantage in analyzing large databases. While version 1.0 appears more reliable and produces more balanced predictions, version 1.1 proves particularly valuable for studying novel folds (Fig. 4).

Heterogeneity analysis shows that most of the high-quality AFDB and ESMAtlas structures build heterogeneous clusters (with at least one representative from each database), meaning that there is a large structural similarity between them (Fig. 5A). Nonetheless, we can still identify a substantial number of AFDB and ESMAtlas singletons and homogeneous clusters – their nature (including taxonomic assignment) is yet to be fully explored.

By design, we included all AFDB clusters created in reference (22). This approach contrasts with the highquality_clust30 definition but enables us to estimate where low-pLDDT AlphaFold predictions – either low-quality models or intrinsically disordered proteins (IDP) – may reside in the structure space. As expected, deepFRI primarily annotates these as having a general function (Fig. 3B). However, considering that many of them are likely disordered, this highlights the need for improved functional annotation of intrinsically disordered proteins – IDPs usually fall into the superCOG 1 group, given their function as molecular chaperones (57, 58). Given that deepFRI is a structure-based tool, the annotation of IDPs would require a development of novel approach, e.g. based on dynamic modeling of IDPs under specific physiological conditions.

In conclusion, this study provides a thorough analysis of the protein structure landscape by identifying representative structures from three large resources: the AlphaFold database (AFDB), a high-quality subset of ESMFold (called highquality_clust30) and the Microbiome Immunity Project (MIP) database. Our findings indicate that those databases are complementary in their coverage of protein folds (*structural complementarity*), although their high-level functional annotations (split into three large functional families called SuperCOGs) display similar profiles. Constructing a low dimensional representation of cluster representatives from all databases, we offer a detailed look into how proteins are distributed across the structure space. We show that proteins representing all three high-level functional groups, SuperCOGs, are localized in specific regions of the structure space (*functional locality*). We also provide a detailed analysis of cluster heterogeneity, showing quantitatively to what extent the databases are similar. Additionally, we have developed a web server for personalized exploration of the clustered dataset, facilitating more tailored analyses (Fig. 6). Ultimately, this work enhances our understanding of the current state of protein structures and their evolutionary and functional implications, offering valuable insights for future research.

Methods

Datasets

We considered three large protein structure datasets: AFDB50 i.e. AlphaFold-database clustered at 50% sequence identity level (22), highquality_clust30 i.e. high quality subset of ESMAtlas (see details here https://github.com/mosamdabhi/esm_meta_protein/blob/main/scripts/atlas/README.md) with pTM and pLDDT > 0.7 clustered at 30% sequence similarity level, and Microbiome Immunity Project (MIP) (which already consists of structures clustered at 30% sequence identity level) (26). They comprise predictions from AlphaFold (7), ESMFold (17), and Rosetta (59) respectively. By default, we did not filter the AFDB structures by pLDDT as we wanted to observe the distribution of structures with low pLDDT within the structure space (note that some of them may be intrinsically disordered proteins). However, quantitative conclusions are drawn for AFDB structures with mean pLDDT > 0.7 (to be in line with highquality_clust30 construction) i.e. Fig. 1C, 3A, and Fig. 5. We also provide additional plots in the Supplement where the aforementioned constraint is imposed (e.g. the whole heterogeneity analysis section). Similarly to reference (22) we exclude highquality_clust30 singletons. We did include MIP singletons since they may contain important but rare domains from the tree of life coming from a curated set of genomes. See the Supplement for more details.

Structural clustering

Structural clustering has been performed using Foldseek (11) version e99047b2214b20b7c93f36fc0d54394c843c046b. We followed a similar procedure as the one described in reference (22). Structure comparison that outputs TM-score has been done using US-align (60) version 20230609. Parameter tuning and more details may be found in the "Structure clustering" section in the Supplement.

Structure space construction

Structure embeddings have been generated using Geometricus (14, 31) v0.5.0 with the `ShapemerLearn` class that outputs 1024-dimensional shape-mer representation (see https://turtletools.github.io/geometricus/getting_started.html for details). We computed shape-mers for all representative structures in the final clustered database comprising 1,505,141 proteins (see Fig. 1 and Table S2). The Geometricus representation has been reduced into two dimensions using the Pairwise Controlled Manifold Approximation Projection (PaCMAP) (32) being the current state-of-the-art technique for this purpose. We performed an extensive grid search to tune PaCMAP parameters (see the "Structure space" section in the Supplement for details). By default, in order to reduce the structure length bias we normalize Geometricus representations, but we also provide results for unnormalized vectors (see Figs. S29-S32, S39-S42). 3D protein structure visualizations have been generated with PyMOL 2.5.0 Open-Source, 2022-11-16.

Function prediction

Function prediction has been done using deepFRI (28) with release v1.0 (https://github.com/flatironinstitute/DeepFRI) and release v1.1 (https://github.com/bioinf-mcb/DeepFRI). deepFRI v1.1 has been trained using exactly the same architecture and training procedure as v1.0 but with a different training and validation sets (see the "Functional annotation" section in the Supplement for details).

go2cog mapping

Mapping of Gene Ontology (GO) terms into Clusters of Orthologous Groups of proteins (COG) has been performed manually with the use of 1,227 GO-terms, 26 COGs and 3 superCOGs (34, 35). Thanks to GO-term propagation with GO graph (`go.obo` file available at https://geneontology.org/docs/download-ontology/, access 20 January 2024) we were able to map 4,504 GO-terms for deepFRI v1.0 (95% coverage of all outputted GO-terms), 8,466 GO-terms for deepFRI v1.1 (91% coverage of all outputted GO-terms), and 9,427 GO-terms in total. The mapping is freely available at https://github.com/Tomasz-Lab/go2cog. deepFRI predictions have been mapped to superCOG categories using the following scheme: 1) Use only high-quality deepFRI predictions (threshold put on deepFRI score equals 0.3 and 0.5 for v1.0, and v1.1 respectively – see the Supplement for details). 2) Group predictions by protein. 3) If only R (general function) annotations are present, annotate a given protein as R, but If more than one superCOG category appears, assign them all equally, ignoring R (if present). 4) If all 3 superCOG categories are present, assign them to the R category. For COG categories, deepFRI predictions have been mapped using the same scheme but in point 4) we relaxed the number of different categories from three to at least five.

Web server

The web server enables users to interactively explore the protein structure landscape. Features include filtering by database, superCOG, protein length, and AFDB pLDDT. The frontend has been created using React (https://react.dev/), while backend is based on Python's FastAPI library (https://fastapi.tiangolo.com/). We have visualized the 2D projections of the embeddings using SciChart (https://www.scichart.com/). The backbone of the visualization consists of 10,000 points randomly sampled from the whole database. Each change in filters or screen location prompts a request to the backend, which responds with

1000 points matching the criteria and displays them on the frontend. The code for both parts is available at https://github.com/wwydmanski/2dPointVis. The web server is accessible at https://protein-structure-landscape.sano.science/ and will be continuously improved.

Data availability

Lists of clusters and singletons obtained from Foldseek, along with all relevant metadata (including structure space coordinates, functional annotations, and number of structures coming from each dataset) as well as Fig. S54 in HTML format, can be found in https://dx.doi.org/10.6084/m9.figshare.27203073. The code and data are also available on GitHub: https://github.com/Tomasz-Lab/protein-structure-landscape.

Supplementary data

The supplement is available as a separate PDF file.

Author contributions

P.S. T.K. P.D.R. and J.K.L. designed the research. P.S. L.S. and T.K. performed analysis. L.S. created the go2cog mapping. P.S. and W.W. retrained deepFRI v.1.1. W.W. developed the web server. P.S. T.K. L.S. J.K.L. and P.D.R. wrote the manuscript.

Acknowledgements

The computations reported in this paper were (in part) performed using resources made available by the Flatiron Institute. The Flatiron Institute is a division of the Simons Foundation. We kindly acknowledge the support of World Community Grid team and the community of volunteers who donated their computational resources to the creation of the MIP database. We gratefully acknowledge Polish high-performance computing infrastructure PLGrid (HPC Center: ACK Cyfronet AGH) for providing computer facilities and support within computational grant no. PLG/2022/015689. Special thanks also go to the high-performance cluster at the Małopolska Centre of Biotechnology of Jagiellonian University, and to its dedicated system administrator, Wojciech Pilch, for his invaluable support in our computational endeavors.

Funding

This work was supported by the National Science Centre, Poland 2023/05/Y/NZ2/00080 to T.K. and P.S.

Conflict of interest statement

None declared.

References

1. Altschul,S.F., Gish,W., Miller,W., Myers,E.W. and Lipman,D.J. (1990) Basic local alignment search tool. *J. Mol. Biol.*, **215**.

2. Hunter,S., Apweiler,R., Attwood,T.K., Bairoch,A., Bateman,A., Binns,D., Bork,P., Das,U., Daugherty,L., Duquenne,L., *et al.* (2009) InterPro: the integrative protein signature database. *Nucleic Acids Res.*, **37**, D211.

3. Hauser,M., Steinegger,M. and Söding,J. (2016) MMseqs software suite for fast and deep clustering and searching of large protein sequence sets. *Bioinformatics*, **32**, 1323–1330.

4. The UniProt Consortium, Bateman,A., Martin,M.J., O'Donovan,C., Magrane,M., Alpi,E., Antunes,R., Bely,B., Bingley,M., Bonilla,C., *et al.* (2016) UniProt: the universal protein knowledgebase. *Nucleic Acids Res.*, **45**, D158–D169.

5. Li,W. and Godzik,A. (2006) Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics*, **22**.

6. Steinegger,M., Meier,M., Mirdita,M., Vöhringer,H., Haunsberger,S.J. and Söding,J. (2019) HH-suite3 for fast remote homology detection and deep protein annotation. *BMC Bioinformatics*, **20**, 1–15.

7. Jumper,J., Evans,R., Pritzel,A., Green,T., Figurnov,M., Ronneberger,O., Tunyasuvunakool,K., Bates,R., Žídek,A., Potapenko,A., *et al.* (2021) Highly accurate protein structure prediction with AlphaFold. *Nature*, **596**, 583–589.

8. Varadi,M., Bertoni,D., Magana,P., Paramval,U., Pidruchna,I., Radhakrishnan,M., Tsenkov,M., Nair,S., Mirdita,M., Yeo,J., *et al.* (2024) AlphaFold Protein Structure Database in 2024: providing structure coverage for over 214 million protein sequences. *Nucleic Acids Res.*, **52**, D368–D375.

9. Varadi,M., Anyango,S., Deshpande,M., Nair,S., Natassia,C., Yordanova,G., Yuan,D., Stroe,O., Wood,G., Laydon,A., *et al.* (2022) AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Res.*, **50**, D439–D444.

10. Elofsson,A. (2023) Progress at protein structure prediction, as seen in CASP15. *Curr. Opin. Struct. Biol.*, **80**, 102594.

11. van Kempen,M., Kim,S.S., Tumescheit,C., Mirdita,M., Lee,J., Gilchrist,C.L.M., Söding,J. and Steinegger,M. (2023) Fast and accurate protein structure search with Foldseek. *Nat. Biotechnol.*, **42**, 243–246.

12. Kim,H., Mirdita,M. and Steinegger,M. (2023) Foldcomp: a library and format for compressing and indexing large protein structure sets. *Bioinformatics*, **39**, btad153.

13. Deorowicz,S. and Gudyś,A. (2023) Efficient protein structure archiving using ProteStAr. *bioRxiv*, 10.1101/2023.07.20.549913.

14. Durairaj,J., Akdel,M., de Ridder,D. and van Dijk,A.D.J. (2020) Geometricus represents protein structures as shape-mers derived from moment invariants. *Bioinformatics*, **36**, i718–i725.

15. Wang,D., Abbas,U.L., Shao,Q., Chen,J. and Xu,D. (2023) S-PLM: Structure-aware Protein Language Model via Contrastive Learning between Sequence and Structure. *bioRxiv*, 10.1101/2023.08.06.552203.

16. Derry,A. and Altman,R.B. (2023) COLLAPSE: A representation learning framework for identification and characterization of protein structural sites. *Protein Sci.*, **32**, e4541.

17. Lin,Z., Akin,H., Rao,R., Hie,B., Zhu,Z., Lu,W., Smetanin,N., Verkuil,R., Kabeli,O., Shmueli,Y., *et al.* (2023) Evolutionary-scale prediction of atomic-level protein structure with a language model. *Science*, 10.1126/science.ade2574.

18. Berman,H.M., Westbrook,J., Feng,Z., Gilliland,G., Bhat,T.N., Weissig,H., Shindyalov,I.N. and Bourne,P.E. (2000) The Protein Data Bank. *Nucleic Acids Res.*, **28**, 235.

19. Bordin,N., Sillitoe,I., Nallapareddy,V., Rauer,C., Lam,S.D., Waman,V.P., Sen,N., Heinzinger,M., Littmann,M., Kim,S., *et al.* (2023) AlphaFold2 reveals commonalities and novelties in protein structure space for 21 model organisms. *Communications Biology*, **6**, 1–12.

20. Waman,V.P., Bordin,N., Alcraft,R., Vickerstaff,R., Rauer,C., Chan,Q., Sillitoe,I., Yamamori,H. and Orengo,C. (2024) CATH 2024: CATH-AlphaFlow Doubles the Number of Structures in CATH and Reveals Nearly 200 New Folds. *J. Mol. Biol.*

21. Lau,A.M., Bordin,N., Kandathil,S.M., Sillitoe,I., Waman,V.P., Wells,J., Orengo,C.A. and Jones,D.T. (2024) Exploring structural diversity across the protein universe with The Encyclopedia of Domains. *bioRxiv*, 10.1101/2024.03.18.585509.

22. Barrio-Hernandez,I., Yeo,J., Jänes,J., Mirdita,M., Gilchrist,C.L.M., Wein,T., Varadi,M., Velankar,S., Beltrao,P. and Steinegger,M. (2023) Clustering predicted structures at the scale of the known protein universe. *Nature*, **622**, 637–645.

23. Durairaj,J., Waterhouse,A.M., Mets,T., Brodiazhenko,T., Abdullah,M., Studer,G., Tauriello,G., Akdel,M., Andreeva,A., Bateman,A., *et al.* (2023) Uncovering new families and folds in the natural protein universe. *Nature*, **622**, 646–653.

24. The UniProt Consortium, Bateman,A., Martin,M.-J., Orchard,S., Magrane,M., Ahmad,S., Alpi,E., Bowler-Barnett,E.H., Britto,R., Bye-A-Jee,H., *et al.* (2022) UniProt: the Universal Protein Knowledgebase in 2023. *Nucleic Acids Res.*, **51**, D523–D531.

25. Mitchell,A.L., Almeida,A., Beracochea,M., Boland,M., Burgin,J., Cochrane,G., Crusoe,M.R., Kale,V., Potter,S.C., Richardson,L.J., *et al.* (2020) MGnify: the microbiome analysis resource in 2020. *Nucleic Acids Res.*, **48**.

26. Koehler Leman,J., Szczerbiak,P., Renfrew,P.D., Gligorijevic,V., Berenberg,D., Vatanen,T., Taylor,B.C., Chandler,C., Janssen,S., Pataki,A., *et al.* (2023) Sequence-structure-function relationships in the microbial protein universe. *Nat. Commun.*, **14**, 1–11.

27. Mukherjee,S., Seshadri,R., Varghese,N.J., Eloe-Fadrosh,E.A., Meier-Kolthoff,J.P., Göker,M., Coates,R.C., Hadjithomas,M., Pavlopoulos,G.A., Paez-Espino,D., *et al.* (2017) 1,003 reference genomes of bacterial and archaeal isolates expand coverage of the tree of life. *Nat. Biotechnol.*, **35**, 676–683.

28. Gligorijević,V., Renfrew,P.D., Kosciolek,T., Leman,J.K., Berenberg,D., Vatanen,T., Chandler,C., Taylor,B.C., Fisk,I.M., Vlamakis,H., *et al.* (2021) Structure-based protein function prediction using graph convolutional networks. *Nat. Commun.*, **12**, 1–14.

29. Maranga,M., Szczerbiak,P., Bezshapkin,V., Gligorijevic,V., Chandler,C., Bonneau,R., Xavier,R.J., Vatanen,T. and Kosciolek,T. (2023) Comprehensive Functional Annotation of Metagenomes and Microbial Genomes Using a Deep Learning-Based Method. *mSystems*, 10.1128/msystems.01178-22.

30. Szydlowski,L., Ehlich,J., Szczerbiak,P., Shibata,N. and Goryanin,I. (2022) Novel species identification and deep functional annotation of electrogenic biofilms, selectively enriched in a microbial fuel cell array. *Front. Microbiol.*, **13**, 951044.

31. Durairaj,J., Akdel,M., de Ridder,D. and van Dijk,A.D.J. (2021) Fast and adaptive protein structure representations for machine learning. *bioRxiv*, 10.1101/2021.04.07.438777.

32. Wang,Y., Huang,H., Rudin,C. and Shaposhnik,Y. (2021) Understanding How Dimension Reduction Tools Work: An Empirical Approach to Deciphering t-SNE, UMAP, TriMap, and PaCMAP for Data Visualization. *J. Mach. Learn. Res.*, **22**, 1–73.

33. Orengo,C.A., Michie,A.D., Jones,S., Jones,D.T., Swindells,M.B. and Thornton,J.M. (1997) CATH--a hierarchic classification of protein domain structures. *Structure*, **5**.

34. The Gene Ontology Consortium, Ashburner,M., Ball,C.A., Blake,J.A., Botstein,D., Butler,H., Michael Cherry,J., Davis,A.P., Dolinski,K., Dwight,S.S., *et al.* (2000) Gene Ontology: tool for the unification of biology. *Nat. Genet.*, **25**, 25.

35. Tatusov,R.L., Galperin,M.Y., Natale,D.A. and Koonin,E.V. (2000) The COG database: a tool for genome-scale analysis of protein functions and evolution. *Nucleic Acids Res.*, **28**, 33.

36. Fullmer,M.S., Ouellette,M., Louyakis,A.S., Papke,R.T. and Gogarten,J.P. (2019) The Patchy Distribution of Restriction–Modification System Genes and the Conservation of Orphan Methyltransferases in Halobacteria. *Genes* , **10**, 233.

37. Hübscher,U. (1983) DNA polymerases in prokaryotes and eukaryotes: Mode of action and biological implications. *Experientia*, **39**, 1–25.

38. Botticelli,S., La Penna,G., Minicozzi,V., Stellato,F., Morante,S., Rossi,G. and Faraloni,C. (2024) Predicting the Structure of Enzymes with Metal Cofactors: The Example of [FeFe] Hydrogenases. *Int. J. Mol. Sci.*, **25**, 3663.

39. Smith,S.P., Bayer,E.A. and Czjzek,M. (2017) Continually emerging mechanistic complexity of the multi-enzyme cellulosome complex. *Curr. Opin. Struct. Biol.*, **44**.

40. The Reference Genome Group of the Gene Ontology Consortium (2009) The Gene Ontology's Reference Genome Project: A Unified Framework for Functional Annotation across Species. *PLoS Comput. Biol.*, **5**, e1000431.

41. Wagner,T., Ermler,U. and Shima,S. (2016) MtrA of the sodium ion pumping methyltransferase binds cobalamin in a unique mode. *Sci. Rep.*, **6**, 1–10.

42. Ragsdale,S.W. ed. (2008) Catalysis of Methyl Group Transfers Involving Tetrahydrofolate and B12. In *Vitamins & Hormones*. Academic Press, Vol. 79, pp. 293–324.

43. Nguyen,T.-Q. and Nicolet,Y. (2022) Structure and Catalytic Mechanism of Radical SAM Methylases. *Life*, **12**, 1732.

44. Wang,T., Leibrock,N., Plugge,C.M., Smidt,H. and Zoetendal,E.G. (2023) In vitro interactions between Blautia hydrogenotrophica, Desulfovibrio piger and Methanobrevibacter smithii under hydrogenotrophic conditions. *Gut Microbes*, 10.1080/19490976.2023.2261784.

45. Levental,I. and Lyman,E. (2022) Regulation of membrane protein structure and function by their lipid nano-environment. *Nat. Rev. Mol. Cell Biol.*, **24**, 107–122.

46. Harayama,T. and Riezman,H. (2018) Understanding the diversity of membrane lipid composition. *Nat. Rev. Mol. Cell Biol.*, **19**, 281–296.

47. Jolie,R.P., Duvetter,T., Van Loey,A.M. and Hendrickx,M.E. (2010) Pectin methylesterase and its proteinaceous inhibitor: a review. *Carbohydr. Res.*, **345**, 2583–2595.

48. Wu,H.-C., Bulgakov,V.P. and Jinn,T.-L. (2018) Pectin Methylesterases: Cell Wall Remodeling Proteins Are Required for Plant Response to Heat Stress. *Front. Plant Sci.*, **9**, 412022.

49. Li,Z., Wu,L., Wang,C., Wang,Y., He,L., Wang,Z., Ma,X., Bai,F., Feng,G., Liu,J., *et al.* (2022) Characterization of pectin methylesterase gene family and its possible role in juice sac granulation in navel orange (Citrus sinensis Osbeck). *BMC Genomics*, **23**, 1–18.

50. Brocchieri,L. and Karlin,S. (2005) Protein length in eukaryotic and prokaryotic proteomes. *Nucleic Acids Res.*, **33**.

51. Lupas,A.N., Ponting,C.P. and Russell,R.B. (2001) On the Evolution of Protein Folds: Are Similar Motifs in Different Protein Folds the Result of Convergence, Insertion, or Relics of an Ancient Peptide World? *J. Struct. Biol.*, **134**, 191–203.

52. Hou,J., Sims,G.E., Zhang,C. and Kim,S.-H. (2003) A global representation of the protein fold space. *Proceedings of the National Academy of Sciences*, **100**, 2386–2390.

53. Hou,J., Jun,S.-R., Zhang,C. and Kim,S.-H. (2005) Global mapping of the protein structure space and application in structure-based inference of protein function. *Proceedings of the National Academy of Sciences*, **102**, 3651–3656.

54. Osadchy,M. and Kolodny,R. (2011) Maps of protein structure space reveal a fundamental relationship between protein structure and function. *Proceedings of the National Academy of Sciences*, **108**, 12301–12306.

55. Duller,S., Vrbancic,S., Szydłowski,Ł., Mahnert,A., Blohs,M., Predl,M., Kumpitsch,C., Zrim,V., Högenauer,C., Kosciolek,T., *et al.* (2024) Expanding the cultivated human archaeome by targeted isolation of novel Methanobrevibacter strains from fecal samples. *bioRxiv*, 10.1101/2024.04.10.588852.

56. Szydlowski,L.M., Bulbul,A.A., Simpson,A.C., Kaya,D.E., Singh,N.K., Sezerman,U.O., Łabaj,P.P., Kosciolek,T. and Venkateswaran,K.J. (2023) Adaptation to space conditions of novel bacterial species isolated from the International Space Station revealed by functional gene annotations and comparative genome analysis. *bioRxiv*, 10.1101/2023.09.28.559980.

57. Hand,S.C., Menze,M.A., Toner,M., Boswell,L. and Moore,D. (2011) LEA Proteins During Water Stress: Not Just for Plants Anymore. *Annu. Rev. Physiol.*, **73**, 115–134.

58. Chakrabortee,S., Boschetti,C., Walton,L.J., Sarkar,S., Rubinsztein,D.C. and Tunnacliffe,A. (2007) Hydrophilic protein associated with desiccation tolerance exhibits broad protein stabilization function. *Proceedings of the National Academy of Sciences*, **104**, 18073–18078.

59. Leman,J.K., Weitzner,B.D., Lewis,S.M., Adolf-Bryfogle,J., Alam,N., Alford,R.F., Aprahamian,M., Baker,D., Barlow,K.A., Barth,P., *et al.* (2020) Macromolecular modeling and design in Rosetta: recent methods and frameworks. *Nat. Methods*, **17**, 665–680.

60. Zhang,C., Shine,M., Pyle,A.M. and Zhang,Y. (2022) US-align: universal structure alignments of proteins, nucleic acids, and macromolecular complexes. *Nat. Methods*, **19**, 1109–1115.