Modelling medulloblastoma pathogenesis and treatment in human cerebellar

organoids.

Thomas Willott[1]*, James G Nicholson[1]*, Yunchen Xiao[1], Olumide Ogunbiyi[2],

Benjamin Draper[3], Talisa Mistry[2], Laura Donovan[3], Nicolae Radu Zabet[1], Ashirwad

Merve[2], Sara Badodi[1+] & Silvia Marino[1,4+]

[1] Brain Tumour Research Centre, Blizard Institute, Queen Mary University of

London, London (UK)

[2] Department of Histopathology, Great Ormond Street Hospital for Children NHS

Foundation Trust, Great Ormond Street, London (UK)

[3] UCL Great Ormond Street Institute of Child Health, London (UK)

[4] Barts Brain Tumour Centre, Queen Mary University of London and Barts Health,

London (UK)

*shared first authors

+shared senior authors: s.badodi@qmul.ac.uk and s.marino@qmul.ac.uk

Abstract

Faithful genetically engineered *in vivo* models of medulloblastoma (MB) are currently
available only for some molecular subgroups, in keeping with recent studies showing
the unique role of human-specific progenitors in the development of G3 and G4 MB
subgroups. We generated human cerebellar organoids (CbO) from expanded potential
stem cells (EPSC) and characterised their epigenetic and transcriptomic profile
compared to the developing human cerebellum. We show the presence of sub
compartment-specific cerebellar lineages linked to MB formation, including
populations expressing signature genes of putative G3 and G4 MB cells-of-origin. We
show that these lineages can be genetically engineered to model MB tumour onset.
Moreover, we demonstrate that CbO sustain proliferation and invasion of G3/4 MB
cells in a 3D co-culture model (CbO-MB) while preserving their molecular identity, and
that treatment of CbO-MB with anti-tumour compounds recapitulate the efficacy of *in*
vivo drug testing in xenograft models.

Introduction

Medulloblastoma (MB) is a cerebellar tumour accounting for the largest proportion of brain cancer diagnoses in children. Molecular subgrouping based on transcriptomic, epigenomic and genomic features has collectively identified 4 MB subgroups: SHH (Sonic Hedgehog), WNT (Wingless-related Integration Site), Group 3 (G3) and Group 4 (G4), each further divided into subtypes with distinct prognosis and responses to therapy[1-4]. Despite these advances in classification, MB standard of care remains unchanged, with patients undergoing surgical resection followed by chemotherapy and/or radiotherapy[5], which, albeit effective in a majority of patients, is associated with significant side effects.

The origins and molecular alterations for SHH and WNT MB are well characterised, leading to a surge in the development of molecularly targeted and/or immunotherapies in *in vitro* and *in vivo* models translating into early-phase clinical trials for patients in recent years[6]. The unclear origin and lack of universal pathway aberrations in G3/G4 MB subgroups along with the consequent lack of faithful pre-clinical models has hampered the identification of new treatment options for these tumours[7]. In particular, *in vitro* models for G4 MB, the most common MB subgroup, are highly underrepresented compared to other subgroups due to difficulties establishing 2D cell culture, whilst currently available G3 MB cell lines predominantly harbour *MYC* amplifications without encompassing the heterogeneity seen in these tumours[8,9]. The maintenance of primary MB cells as patient-derived orthotopic xenograft (PDX) models has overcome some of these issues[10-12], however, the cost and resources required for these models limit their broader applications and scalability for drug testing.

MB formation is closely linked to deregulated cerebellar development[13]. Mapping lineage trajectories from MB single cell RNA sequencing (scRNA-Seq) data to human foetal cerebellar development has recently identified spatiotemporally restricted mitotic glutamatergic neuron progenitors in the rhombic lip subventricular zone (RL^{SVZ}) harbouring photoreceptor (PRC) and unipolar brush cell (UBC) gene signatures as putative G3 and G4 MB cells of origin respectively[14,15]. These progenitors are halted in development prior to cell cycle exit leading to malignancy. Importantly, immunohistochemical and pseudo-time comparisons of relevant stages of human and murine cerebellar development showed significantly fewer of these proliferative

progenitors in mice, raising the possibility that mouse models may not allow the study

of key features of G3/G4 MB, including their onset[14,15]. Additionally, a human-specific

transitional cell progenitor (TCP) population has been identified in the developing

foetal cerebellum at 12-14 PCW, which is maintained or enriched in G3/4 MB

tumours[16]. More recently, neural stem cells in the RL^{VZ} harbouring a

$PRTG^+;SOX2^+;MYC^{high};NESTIN^{low}$ signature have been linked to G3 MB onset[17].

These studies suggest that the failed maturation of early cerebellar cell lineages at

specific sub-compartment location and developmental stage is linked to the ontogeny

of G3/4 MB tumours.

Advances in organoid research have enabled the generation of new tools to study

various aspects of cancer biology, including tumour initiation[18], metastasis[19] and as a

co-culture model for drug testing[20], with important implications for personalised

medicine[21]. Methods to generate cerebellar organoids from induced pluripotent stem

cells (iPSC) have been developed[22,23] and characterised at single cell transcriptome

level, achieving neuronal and glial differentiation with cerebellar sub-compartment

identity[24-26]. In the context of MB modelling, current applications of cerebellar

organoids are restricted to studies of neoplastic transformation in bulk genetically

engineered organoids which lacked specificity for their respective cell-of-origin[27,28], or

in co-culture with established MB cell lines belonging to the SHH subgroup[29].

Here, we have generated human cerebellar organoids (CbOs) from expanded

potential stem cells (EPSC) and assessed their suitability to model G3/G4 MB. We

have shown that they contain the putative MB cell-of-origin at stages of differentiation

comparable to foetal cerebellar development and demonstrated that these cells can

be genetically engineered to model early events in tumour initiation. In addition, we

have analysed the suitability of CbOs as a co-culture model of G3/G4 MB providing

insight into MB transcriptional state heterogeneity and tumour-microenvironment

interactions. Finally, we show the value of the model as a pre-clinical drug testing tool.

Results

CbO display a cellular composition recapitulating human cerebellar development.

We generated human CbO from two EPSC lines (CbO19 and CbO61) obtained from reprogramming of dura mater fibroblasts and extensively previously characterised[21,30]. Organoid differentiation was followed for 35 days, as outlined in previously published iPSC-derived protocols (Figure 1A and S1A)[22,23]. We observed a significant increase in organoid size throughout differentiation (Figure S1B) and the formation of characteristic flat-oval neural rosettes at Day 35 (D35) of culture (Figure 1B). Expression of the stemness marker *NANOG* showed a significant decrease by D7, confirming the commitment of the EPSC to differentiate (Figure S1C). Additionally, the neural stem cell (NSC) markers *NESTIN* and *SOX1* demonstrated early expression from D7-14, as well as the cerebellar homeobox and developmental genes *EN2* and *OTX2,* involved in early isthmic organiser signalling[62] (Figure 1C and Figure S1D). Early progenitor markers *KIRREL2* and *PAX6* were expressed at D7-14 followed by markers of more mature precursors including *OLIG2/SKOR2* and *ATOH1/BARHL1* at D21-28, indicating the presence of GABAergic and glutamatergic lineages respectively (Figure 1C and Figure S1E, F). Interestingly, the transcription factor *EOMES,* expressed in glutamatergic progenitors and later in UBC, was expressed both in early (D7) and late (D28-35) stages of CbO maturation (Figure 1C and Figure S1F). The presence of GABAergic and glutamatergic granule neurons (GN) was confirmed by expression of CALBINDIN and Neu-N proteins respectively, proximal to SOX2-positive neural rosettes at D35 (Figure 1D and 1E). We observed similar patterns of expression throughout the differentiation of CbO19 and CbO61 (Figure 1C), in keeping with the conclusion that these CbOs reproducibly differentiate into cerebellar cell lineages.

To characterise in depth the cellular composition of CbO we performed scRNA-seq at D7, D35 and D49 timepoints to capture the earliest and more mature stages of cerebellar development and neurogenesis (Figure 1F). Initial data exploration revealed broad differences between D7 and the more heterogeneous D35 and D49 timepoints which were separated for downstream analysis (Fig S1G). Louvain clustering and Uniform Manifold Approximation and projections (UMAP) of D7 CbO cells identified three broad clusters corresponding to *PAX6*+ and *OTX2*+ glutamatergic progenitors, *PAX2*+, *PAX5*+ and *PKDCC*+ GABAergic progenitors and SOX9+ and S100B+ glial progenitors. We further identified a small population of nascent

NEUROD4⁺ and *NEUROG1*⁺ glutamatergic neurons, suggesting an initial lineage

commitment (Figure S1H and S1I). By D35 and D49 there was a greater degree of

cellular diversity, and we identified 17 clusters (Figure S1J), with none specific to either

timepoint (Figure S1K). Clusters were annotated based on their expression of marker

genes (Figure 1G, H) and correlations to endogenous cell types from reference

datasets of human developing cerebellum (Figure S1L)[16,42-44]. We identified non-

committed *MKI67*⁺ and *CCN1B*⁺ cycling progenitors as well as *OTX2*⁺ and *SOX2*⁺

progenitor clusters. Two NSC clusters were identified, a broad cluster of *SOX1*⁺,

SOX2⁺*, EN2*⁺ and *NES*⁺ NSCs and a more defined cluster expressing *SOX2*⁺*, PRTG*⁺

and *PAX6*⁺. We found several clusters associated with glutamatergic lineages

including two clusters of *BARHL1*⁺ and *RBFOX3*⁺ progenitors characterised by *PAX6*

or *ATOH1* expression (glut_progenitors and GN/UBC_progs respectively).

Additionally, we detected a *NEUROD1*⁺ GN/UBC cluster also expressing mature

glutamatergic neuronal markers (*LMO1* and *GABRA2*) and a *NEUROD6*⁺ and *MEIS2*⁺

glutamatergic progenitor cell (GPC) cluster. We further identified two mixed clusters

of maturing cerebellar neurons (matCN1 and matCN2) positive for *MEIS2, SLC17A6*

and *NR4A2* which are expressed by neurons located in the neural transitory zone

(NTZ) during cerebellar development[44]. We further observed a neuronal cluster of

GABAergic *SKOR2*⁺, *LHX1*⁺ and *GAD2*⁺ Purkinje cells as well as *FABP7*⁺ and

SLC1A3⁺ Bergmann glial cells (Berg_Glia). In addition to cerebellar lineages, we also

identified other hindbrain clusters containing cell types contributing to cerebellar

development, including *FOLR1*⁺ and *RSPH1*⁺ ependymal/choroid plexus, *POST*⁺ and

COL1A1⁺ meninges and *SLIT2*⁺ and *RSPO3*⁺ roof plate (Figure 1G, H).

Our data show that the cellular composition of CbO recapitulates the main cell types

found during cerebellar development.

154   Maturation of CbO follows conserved epigenetically-regulated central nervous

155   system developmental pathways.

Epigenetic regulation of gene expression, in particular DNA methylation, plays a

fundamental role during central nervous system (CNS) development as it controls the

spatio-temporal activity of cell lineage-specific genes[63]. Hence, we analysed the DNA

methylation profile of CbOs during differentiation and maturation (Figure 2A). We

identified genes showing the most variable time-dependent methylation status (VMGs)

during CbO development and grouped them into time-dependent methylation
programs by k-means clustering. Among genes exhibiting an increasing methylation
status during CbO maturation (clusters 1 and 2) we found transcription factors linked
to early cerebellum development including *PAX2*, *PAX5* and *PAX6* and to cell lineage
specification such as *LMX1A* and *OTX2*. On the contrary, genes involved in neuronal
differentiation and axon-guidance including *MAP2*, *MAP4*, *MAP7* and *GRIP1* were
progressively hypomethylated during CbO differentiation (cluster 4 and 5) (Figure 2B).
Gene Ontology (GO) pathway analysis of the identified VMGs revealed that early
hypomethylated clusters were enriched for biological processes related to
embryogenesis, fate specification, and organ development. Conversely, genes
displaying later hypomethylation patterns were associated with neuronal
differentiation, signal transduction, and structural functions such as synaptic junctions
and neuron projections (Figure 2C). Analysis of human cerebellar samples (Figure
S2A) revealed similar clustering of VMGs with distinct methylation patterns through
human development (Figure S2B). GO analysis of human VMG clusters demonstrated
shared earlier pathways with CbOs including those involved in proliferation and tissue
development, followed by later pathways associated with signal transduction, neuron
projection and catalytic activity (Figure S2C). Notably, we identified 1086 common
VMGs demonstrating similar changes in methylation status between human and CbO
samples (Figure 2D) suggesting that CbOs recapitulate epigenetic regulation of the
human foetal cerebellum. Finally, we integrated DNA methylation and RNA expression
analysis to identify genes with concordant methylation status and expression between
D7 and D35 (i.e. hypomethylated and upregulated or hypermethylated and
downregulated, Figure 2E). We found a significant enrichment for pathways involved
in tissue morphogenesis, proliferation and receptor signalling for
hypermethylated/downregulated genes expressed during early CbO development and
enrichment of neuronal synapse pathways for hypomethylated/upregulated genes
expressed later in CbO differentiation (Figure S2D).

Taken together, our data show that maturation of CbOs is governed by epigenetic
regulation of key developmental pathways similar to the human cerebellum.

CbOs reproduce the spatiotemporal cellular complexity required for
medulloblastoma initiation.

Next, we compared DNA methylation profiles of CbO and human foetal cerebellum

samples to stage the development achieved. We observed the highest correlation of

D35 CbO with 12-17 PCW cerebellum by comparing the methylation profiles of the top

1% of CpG probes with a linear correlation of methylation status across human foetal

cerebellar development (Figure 3A and Figure S3A). Similarly, the application of an

epigenetic clock algorithm trained on independent foetal brain datasets[57] showed a

positive correlation between differentiation of CbO and their predicted epigenetic age

(R-squared = 0.8), confirming the staging of D35 CbO at 14 PCW and progressive

maturation of the organoid (Figure S3B).

Moreover, we analysed pseudo-bulk CbO RNA-seq data against the average

expression of reference datasets[16,42-44] from developing cerebellum and observed that

D35 CbO show a high correlation in expression of age-related genes with 9-15 PCW

(Figure 3B). Interestingly, our D35/49 CbOs acquire a similar expression profile to

other cerebellar organoids cultured for longer time periods[24,25,43] (Figure 3B).

Putative G3/4 MB cells-of-origin were first described in the RL^{SVZ}, and we indeed saw

an enrichment of an RL^{SVZ} gene signature score in the glutamatergic cerebellar

neuronal and progenitor clusters, which was depleted in less differentiated cycling

progenitors and NSC (Figure 3C). Correlation of CbO methylation levels to

microdissected RL and EGL samples[14] revealed an increased correlation through CbO

development with both regions, reflecting the expanding populations of predominantly

glutamatergic neuronal progenitors in the CbO, and showing high correlation of D35

CbO specifically with RL (Figure 3D)..

Previous studies showed enrichment of photoreceptor and UBC expression signatures

in RL^{SVZ} (PRC.RL^{SVZ} or UBC.RL^{SVZ}), with a high expression in G3 and G4 MB putative

cells-of-origin respectively. Quantification of these signatures within CbOs confirmed

their enrichment in discrete clusters of glut_progenitors or GN/UBC_progs

respectively, with the UBC.RL^{SVZ}-enriched GN/UBC_prog cluster also highly

correlated to TCP cells (Figure S1L), a progenitor population previously associated

with G3/4 MB[16]. Importantly, both of these progenitor clusters were characterised by

EOMES expression (Figure 3E, F and Figure S3C) and we observed EOMES$^+$Ki67$^+$

cells located predominantly within the neural rosettes in CbO, reminiscent of the

human-specific G3 and G4 MB-susceptible progenitors (Figure 3G, H).

Notably, the early_RL/NSC cluster, which RNA velocity analysis predicts to give rise

to PRC.RL^{SVZ}-enriched glut_progenitors, presents high expression of *PRTG, SOX2*

and *MYC* and low expression of *NESTIN* (Figure 1H, S3D), mirroring the gene

signature found in a $PRTG^+;SOX2^+;MYC^{high};NESTIN^{low}$ stem cell population in the

RL^{VZ} recently proposed as a putative G3 cells-of-origin[17]. Immunohistochemical

staining revealed PRTG^{High};SOX2$^+$ cells outside of CbO neural rosettes in areas

presenting concomitant low expression of NESTIN (Figure 3I, J), suggesting a spatial

complexity to glutamatergic NSC/progenitor cells in CbOs with differing MB cell-of-

origin signatures.

Given that pre-malignant G3/4 MB cells-of-origin are thought to persist in the RL, likely

due to their stalled differentiation[15], we performed pseudotime and RNA velocity

analysis through the D35/49 UMAP (Figure S3D,E). Plotting G3 and G4 trajectories

through each UMAP demonstrated sequential expression of glutamatergic neuronal

genes associated with early lineage commitment, progenitor expansion and more

mature neurons (Figure S3F). We observed an earlier high expression of the G3-

associated genes *PRTG, OTX2* and *MYC* during sustained *SOX2* expression in the

G3 trajectory in line with G3 cells-of-origin harbouring stem and early progenitor

phenotypes (Figure 3K). Whereas, in the G4 trajectory, MB-associated genes *EOMES*

and *PAX6* exhibited high expression relatively later in pseudotime, after *SOX2* loss

and before upregulation of the mature glutamatergic marker *SLC17A6*, highlighting a

more committed progenitor cell population (Figure 3K).

Taken together, our data show that D35 CbOs display an epigenetic and

transcriptional landscape mirroring the cerebellar developmental stage and cellular

profiles of G3/G4 MB initiation.

251 Genetic editing of MB cell-of-origin populations in CbO models tumour onset.

Leveraging the observation that CbO mirrors the cellular complexity of the human

cerebellum at the time of G3/G4 MB formation, we sought to model tumour initiation

in CbOs by specifically targeting their relevant cell-of-origin.

We set out to overexpress the GFP-tagged *c-MYC* oncogene in G3/G4 progenitors

and to achieve selective spatio-temporal targeting of the cells we used a Tet-inducible

system driven by the promoter of *PAX6* or *EOMES*, transcription factors known to have

a role in MB[64,65] and expressed in MB cell-of-origin clusters at D35 of CbO

differentiation (Figure 1H, S3C, 3K). EPSCs transduced with either *pEOMES;MYC-*

GFP or *pPAX6;MYC-GFP* were used to generate CbOs and c-*MYC* overexpression

was induced at D35 of maturation (Figure 4A). A *BFP* tag was added to p*EOMES* and
p*PAX6* constructs to allow purification of the engineered cells from non-induced CbOs
and we confirmed retention of early stem cell markers expression in transduced EPSC
(Figure S4A). Foci of c-MYC-GFP[+] cells were observed as early as 4 days after
doxycycline treatment in both editing systems, with their expansion observed upon
maturation (Figure S4B). D63 CbO showed clusters of highly proliferative GFP[+]Ki67[+]
cells, which corresponded to areas of densely packed hyperchromatic and mitotically
active cells (Figure 4B).

To investigate the transcriptional consequences of *c-MYC* activation in the putative
G3/4 cells-of-origin, we next performed scRNAseq on edited CbOs, transferring cell
type labels from our previously annotated D35/49 dataset. c-*MYC* expression was
particularly high in a cluster made up almost exclusively of cells from *pEOMES;MYC-*
GFP and *pPAX6;MYC-GFP* CbO samples (cluster 12) that we therefore labelled as
MYC overexpressing (MYC[OE]) cluster (Figure 4C, D and Figure S4C, D). Gene
signature scoring confirmed the proliferative status of the MYC[OE] cluster as well as its
similarity to G3 MB[47] (Figure S4E, F). Gene set enrichment analysis of the MYC[OE]
cluster confirmed enrichment of downstream MYC targets (Figure S4G) and GO
enrichment of upregulated cluster markers highlighted terms consistent with
epigenetic dysregulation (protein-DNA complex disassembly, chromatin remodelling,
regulation of DNA methylation) and signalling pathways linked to MB (Notch signalling
pathway, inositol phosphate-mediated signalling) (Figure 4E)[35,66]. Interestingly, we
found enrichment of oncogene-induced senescence, a result in apparent contrast with
the relatively proliferative signature scoring for this cluster but likely reflective of the
countervailing forces of tumour suppression and oncogenic signalling as these cells
undergo transformation.

Transformation of MB cells-of-origin has been linked to their stalled differentiation[15],
we therefore projected MYC[OE] cells onto the D39/45 CbO UMAP (Fig 4F). As
expected, the MYC[OE] cluster from *pEOMES;MYC-GFP* CbO samples (p*EOMES-*
MYC[OE]) overlaid the G4 lineage whereas those from the *pPAX6;MYC-GFP* CbOs
(p*PAX6-* MYC[OE]) were more equally distributed between the G4 and G3 lineages.
Notably, G4 lineage and p*EOMES-* MYC[OE] cells express *ATOH1*, while the G3 lineage
and p*PAX6-* MYC[OE] express higher levels of *PRTG* and *OTX2* (Fig 4G), supporting
our choice of *PAX6*/*EOMES* promoters to target specific progenitor clusters.

We next scored cells for patient-derived MB gene signatures, confirming that both our
G3/4 lineage and MYC[OE] clusters were more similar to G3/4 subgroup (Figure 4H).
Importantly, we observed a significantly higher G3/4 MB signature score (Figure 4I) in
engineered MYC[OE] cells compared to their G3/4 lineage cells-of-origin alongside with
significant depletion of gene sets associated with neuronal differentiation and
enrichment of a gene set corresponding to poor MB prognosis (Figure 4J and S4H).

We further investigated the methylation profiles of sorted GFP[+]-MYC[OE] cells as
compared to BFP[+] lineage-of-origin controls and found that they clustered with G3 MB
patient tumours, as assessed by a MB subgroup-specific CpG probe set[1,59] (Figure
S4I). Furthermore, when focusing the analysis on DMPs between MYC[OE] and lineage-
of-origin we observed a stronger correlation to G3 MBs for the *pPAX6*-MYC[OE]
compared to *pEOMES*-MYC[OE] model (Figure S4J). Interestingly, analysis of DMGs
revealed model-specific deregulation of genes with *pPAX6*-MYC[OE] demonstrating
specific hypomethylation of *OTX2* and *GFI1*, known drivers of G3 MB onset[27], whilst
pEOMES-MYC[OE] showed hypomethylation of genes with mutually exclusive roles to
OTX2 in G3/4 MB pathogenesis and cell fate decisions including *PAX3* and
CBFA2T2[15,64] (Figure 4K). Integration of DNA methylation and gene expression data
further showed a concordant positive regulation of several key genes (*HMGA1*,
SLC7A5 and *FABP3*) and negative regulation of genes associated with neuronal
differentiation (*SALL2*, *NEUROD2* and *GRIA2*) (Figure 4L).

Taken together, we show that using CbO we can model features of MB onset in G3/4
lineages-of-origin through *c-MYC* overexpression including increased proliferation,
stalled development and transcriptional changes consistent with epigenetic
dysregulation and poor MB outcome.

CbOs sustain co-cultured medulloblastoma cells proliferation and invasion
while preserving their molecular identity.

As well as utilising CbO to model early events in MB cell-of-origin transformation, we
also set out to assess whether CbO could provide a physiologically relevant 3D
environment for sustaining growth of human MB cells. We co-cultured GFP-tagged
G3/4 MB cell lines (CHLA-01-Med, CHLA-01R-Med and ICb1299) as well as G3 MB
primary cells (Med211-FH) maintained as a PDX with D35 CbO when cerebellar
lineages are clearly defined (Figure 5A). All MB lines successfully attached and

infiltrated the CbO with considerable cell growth after 14 days of co-culture (Figure 5A

and S5A-C). Morphological analysis of CbO bearing MB cells (CbO-MB) showed a

highly cellular and proliferative pleomorphic embryonic tumour, in keeping with MB,

diffusely infiltrating the adjacent CbO tissue (Figure S5D-F). We observed positivity

for the proliferation marker Ki67 in around 30% of GFP[+] MB cells (Figure 5B), thus

confirming that CbOs can sustain MB cells proliferation and infiltration.

To explore the effect of MB cells on CbO, we performed scRNA-Seq on pooled CbO-

MB samples (CHLA-01-Med or Med211-FH) after 14 days of co-culture (Figure 5A),

as well as control CbOs co-cultured with cerebellar NSCs (NSC73.5, CbO-NSC).

Louvain clustering and inferred copy number analysis identified CHLA-01-Med and

Med211-FH MB cells (Fig 5C, S5G-H) expressing tumour-specific markers

(*SIX3*/*APOE* and *NEUROD1*/*BCAT1* respectively) as well as shared MB-specific

markers we previously observed in our modelled MYC[OE] cells (*HMGA1*, *MYC*, *YBX3*

and *TRAP1*). NSC73.5 cells could be distinguished by their separation in the UMAP,

and expression of stemness markers (*SOX2/9*), doublecortin (*DCX*) and transcription

factors governing interneuron development (*DLX1/2*) (Figure S5I). In keeping with the

highly proliferative properties of tumour cells, we identified MB cells representing more

than 80% of the sequenced CbO-MB cells as compared to NSC73.5 forming around

20% of CbO-NSC (Figure 5C).

To investigate the degree to which co-cultured CbO-MB cells retain their subtype

specificity, we computationally isolated co-cultured MB cells and scored them for

patient-derived MB gene signatures[45]. We observed a significant increase in average

G3/G4 metaprogram scores when compared to WNT or SHH scores (Figure 5D),

indicating that both MB cell lines retained their molecular subtype in the CbO-MB

model. We next correlated pseudobulk transcriptional profiles with the same reference

cohort of patient MB tumours[45]. As expected, CHLA-01-Med fell within an intermediate

subcluster containing both G3 and G4 primary tumours, whilst Med211-FH clustered

with G3 tumours only (Figure 5E). Label transfer of G3 and G4 tumour scRNA-seq

profiles onto co-cultured MB clusters further confirmed these subgroup identities

(Figure S5J). Studies of glioblastoma have shown that tumour cells more closely

recapitulate their parental tumours when co-cultured in brain organoids as compared

to 2D models[21]. We therefore correlated expression profiles of each MB line co-

cultured with CbO or in 2D with patient tumour samples[35,45]. Importantly, we observed

a significantly higher correlation in all co-cultured MB cells compared to their 2D

cultured counterparts (Figure 5F), suggesting that CbO-MB better phenocopy *in vivo*

primary tumours.

These data show that CbO represents a novel 3D model for G3 and G4 MB cell growth

and expansion which is superior at preserving features of patients' tumour samples as

compared to 2D cultures.

Group 3 MB cells harbour a subset of cells with myogenic differentiation

We next analysed the MB fraction of CbO-MB co-cultures, performing Louvain

clustering and correlating cluster marker signature to each other and to references of

the developing cerebellum (Figure 6A-C and Figure S5K). In line with previous studies,

both cell lines were predominantly composed of clusters corresponding to cycling and

non-cycling GCP-like cells (clusters 1,7, and 2 for Med211-FH; clusters 3 and 0 for

CHLA-01-Med). Overall, Med211-FH clusters correlated more closely with developing

cerebellar cell types than those from the CHLA-01-Med cell line and expressed genes

consistent with a more neural phenotype (*SYT1, NEUROD1*). We also identified two

pairs of smaller cell states that were common to both MB lines, clusters 8 and 5 and

clusters 4 and 6 from Med211-FH and CHLA-01-Med, respectively (Figure 6D).

Clusters 8 and 5 correlated to glial progenitor/NSC developing cerebellar cell types

(Figure S5K) and specifically expressed stemness markers (*SOX2, NTRK2*) (Figure

6C). GO enrichment of the 44 shared marker genes highlighted terms associated with

neuronal development and radial glia differentiation (Figure 6E). Clusters 6 and 4

showed a phenotype more consistent with immune reference cells and expressing

immune-associated markers such as *IL17B* (Figure 6C and Figure S5K). Further

analysis of the 127 shared marker genes between clusters 6 and 4 revealed a strong

enrichment of GO terms associated with myogenic development and differentiation

(Figure 6F). This included expression of key transcription factors governing myogenic

differentiation (*MYOG, MYOD1* and *HLHL41*) as well as components of the myosin

light chain (*MYL4*) and troponins (*TNNT1*) (Figure 6C) with MYOGENIN[+] MB cells

confirmed by IHC in CbO-MB co-cultures (Figure 6G). Previous histological studies

have reported aberrant myogenic differentiation of MB cells[67,68], but the transcriptional

state associated with such phenotypes has not previously been described. By scoring

reference MB samples[45] for our CbO-MB derived myogenic state signature, we

observed the highest score in the WNT MB subtype (Figure 6H) while analysis of

Group 3 MB revealed a subset of cells scored highly for our signature suggesting
heterogenous expression of this metaprogramme (Figure 6H). Indeed, upon sub-
setting, and re-clustering Group 3 MB samples we observed two clusters, 4 and 6,
which expressed this myogenic signature strongly (Figure 6I). These myogenic cells
were found in 11/16 samples ranging from 0.3-70% of total cells (Figure 6J). Scoring
of the myogenic signature in a larger cohort of 763 MB samples[1] revealed a similar
distribution across MB groups (Figure 6K), with an enrichment in the poor prognosis
MYC-amplified Group3_gamma subtype (Figure 6L) and a significant reduction of
survival in samples scoring highly (> median) for the myogenic signature (Figure 6M).

Overall, we identify a subset of MB cells in CbO-MB exhibiting myogenic differentiation
which is recapitulated in patient tumours and linked to poorer prognosis.

406   CbO-MB co-cultures reveal a differential impact of medulloblastoma
407   engraftment on cells of the tumour microenvironment.

Little is known about the MB tumour microenvironment (TME), both in terms of the
crosstalk between MB cells and non-neoplastic neuronal lineages and the molecular
regulators of this interaction, or about the contribution of the MB-TME interplay to
treatment response and recurrence in patients[69]. To investigate how MB cells affect
CbO cellular complexity, we identified CbO-specific cell types in CbO-MB, annotating
clusters based on label transfer using our non-co-cultured organoid as reference
(Figure 7A and Figure S6A-C). The most striking difference was a depletion of mature
cerebellar neurons (MatCN) which was evident in all CbO-MB co-cultures but absent
in the CbO-NSC co-culture control (Figure 7B and Figure S6C,D). Selective negative
impacts on neuronal populations have been shown in other brain tumour organoid co-
cultures[70,71], we therefore reasoned that MB co-culture was most likely causing
neuronal death. Using IHC, we validated the presence of apoptotic neurons (cCASP3$^+$
and MAP2$^+$/TubIII$^+$) proximal to GFP$^+$ MB cells which were not present in CbO-NSC
controls (Figure 7C and Figure S6E).

To characterise the molecular events mediating MB-TME crosstalk we modelled
receptor-ligand (R/L) interactions in CbO co-cultures using CellphoneDB. Overall, we
found that both CHLA-01-Mef and Med211-FH cells had fewer R/L interactions with
CbO cells than NSC73.5 controls (Figure S6F), in keeping with MBs demonstrating
limited TME reactivity compared to other brain tumours [69]. Among the 27 cancer-

specific R/L interactions shared between MB cells but absent in NSC73.5 (Figure 7D),

we observed interactions mediated by the cell adhesion protein CADM3, the NOTCH

signalling pathway, the receptor NRP2 as well as glutamate signalling via SLC1A7

and SLC17A7 (Figure 7E). We validated the predicted NOTCH1/DLL3 interaction

between MB and cycling progenitor clusters using immunohistochemistry, revealing

NOTCH1[+]SOX2[+] CbO cells proximal to GFP[+] MB cells, as well as co-localisation of

NOTCH1 and DLL3 specifically on MB cells surface (Figure 7F).

To gain further insights into the effect of MB cells on the surrounding CNS tissue, we

investigated cell type-specific transcriptional modulation caused by the presence of

MB cells and identified differentially expressed genes (DEGs) (Figure S6G). GO

analysis showed a downregulation of synapse and cytoskeletal pathways in GPCs

whilst Ependymal/ChP cells demonstrated downregulation of cilia-related pathways

and cholesterol metabolism, suggesting changes to the normal development of these

populations within CbO-MB (Figure S6H). Altered neurogenesis, development and

apoptosis pathways in stem/progenitor populations further highlight negative changes

elicited by MB on CbO populations, suggesting a more stressed cell state.

Interestingly, meningeal cells exhibited a downregulation in apoptosis and

upregulation in metabolic pathways, perhaps indicating these cells better tolerate the

tumour burden in CbO-MB co-cultures (Figure S6H).

Our data show that CbO-MB is a 3D model suitable for studying cell-cell interactions

and molecular events/cascades induced by MB cell engraftment and identifies specific

populations of CNS tissue that are more exposed to detrimental stimuli during MB

formation and expansion.

CbO-MB are an effective pre-clinical drug testing tool.

Following on from the observations that CbO-MB maintain tumour cells in a molecular

state closer to human samples whilst simultaneously providing insights into the

functional impact of MB growth on the surrounding cerebellum, we sought to

investigate whether CbO-MB could be suitable as a drug testing tool. We previously

demonstrated that combination therapies extend the survival of mice xenografted with

MB cells recapitulating a specific molecular signature found in a proportion of MB

tumours, whereby the polycomb group gene BMI1 and the chromatin remodelling

factor CHD7 are overexpressed and downregulated, respectively (BMI1^{High};CHD7^{Low}).

Treatment with BMI1 (PTC-209) and MAPK (PD325901) inhibitors demonstrated a

synergistic, cytotoxic effect on BMI1^{High};CHD7^{Low} MB cells whilst the phospho-inositol

metabolic modulator IP6 exerted cytostatic anti-tumour efficacy[35,72]. Here, we co-

cultured GFP-tagged BMI1^{High};CHD7^{Low} CHLA-01-Med cells with D35 CbO and treated

CbO-MB at 14 days, a timepoint when a large population of proliferative MB cells was

established (Figure 5A, B), with BMI1/MAPK inhibitors or IP6 for four days (Figure 7G).

We observed a significant decrease of GFP^{+} BMI1^{High};CHD7^{Low} MB cells upon both

treatments and concomitant increase in cCASP3 or decrease in Ki67 respectively, with

no significant impact on CbO cells (Figure 6H-K and Figure S6I, J). This confirms that

MB cells grown in 3D models recapitulate the treatment response as observed in a

widely accepted pre-clinical *in vivo* model, whilst also indicating negligible side effects

on CbO-MB non-neoplastic cells.

Taken together, these data show that CbO-MB are an effective 3D *in vitro* drug testing

tool which perform similarly to *in vivo* xenograft models.

Discussion

We show that CbO faithfully recapitulate the developmental stages of the human

cerebellum relevant for MB initiation and contain cell lineages which display gene

signature enrichment for G3 and G4 MB cells-of-origin. Genetic editing of these cells

models MB onset and provides the first proof of concept that they play a role in MB

pathogenesis. Moreover, co-culture of patient-derived MB cells within the CbO

generates a 3D *in vitro* model which sustains MB cells proliferation and invasion while

preserving their molecular identity and recapitulates the results obtained in pre-clinical

in vivo drug testing.

Recent studies have shown the fundamental role of human-specific stem and

progenitor cells in the development of G3 and G4 MB[14,15,17], clearly exposing the

limitations of mouse models and highlighting the need for new human-specific models.

Putative G3 and G4 MB progenitor cells-of-origin are generated in the RL^{SVZ} upon RL

compartmentalisation at 11 PCW and show a specific spatiotemporal pattern that

peaks at around 14 PCW. Previously published cerebellar organoid protocols

observed a correlation between two-month-old organoid cell populations and RL cell

clusters from human datasets[26] and a modest enrichment for UBC.RL^{SVZ} signature in

the GNP cluster of a three-month-old organoid[14]. Here, we leveraged comparative

epigenetic and transcriptomic analysis of our CbO model and human cerebellar tissue

samples to accurately characterise and date the development of our model and

provide the first compelling evidence that it faithfully recapitulates stages of human

development known to enable MB initiation. We observed a significant enrichment for

RL^{SVZ} signatures in discrete cell clusters at D35 of CbO maturation stage. Importantly,

we identified specific cell clusters showing enrichment for UBC.RL^{SVZ} and PRC.RL^{SVZ}

expression signatures, the latter never observed before in organoid models. TCP have

been reported as putative G3/4 cells-of-origin, and we found these cells to most

strongly correlate to our UBC.RL^{SVZ}-high G4 MB lineage cells. Recent work suggest

G3 MB originates from *PRTG*⁺;*SOX2*⁺;*MYC*^high;*NESTIN*^low cells. We observed this less

differentiated NSC population in our CbO, which immediately preceded the PRC.RL^{SVZ}

-high cluster in pseudotime/RNAvelocity trajectories. Thus, our CbO data help unify

these competing theories and suggest that both putative G3 cells-of-origin are partially

overlapping populations corresponding to subtly different stages along a shared G3

lineage-of-origin. Interestingly, we found a difference in distribution of EOMES⁺Ki67⁺

and PRTG^High SOX2⁺ cells in mature CbO with respect to neural rosettes, suggesting

a spatial contribution to their differing lineage commitments mimicking the events

linked to neoplastic transformation of MB cells-of-origin during human cerebellar

development. Therefore, our CbO model provides an unparalleled opportunity for

modelling the initiation of G3 and G4 MB by providing access to their putative human-

specific cells-of-origin in a 3D experimental system.

c-MYC amplification has been robustly demonstrated as an oncogenic driver in a

subset of aggressive G3 MB, as well as reported in a proportion of G4 MB[2,73]. Previous

studies have overexpressed *c-MYC* in combination with other G3 drivers *OTX2* or

GFI1 in bulk cerebellar organoids, observing an increase in proliferative cells[27].

Leveraging the in-depth developmental characterisation of our CbOs, we implemented

a refined MB-modelling approach, specifically overexpressing *c-MYC* oncogene in

G3/4 MB lineages-of-origin in a CbO at an MB-relevant developmental stage. This

spatiotemporal inducible system generated clusters of highly proliferative cells

overexpressing *c-MYC* showing an increased correlation to G3/4 MB tumours at both

transcription and DNA methylation levels. Strikingly, we also observed

hypomethylation of *OTX2* and *GFI1* specifically in the *pPAX6* targeted G3 lineage-of-

origin model, reflecting a susceptibility of specific cells in our CbO to MB

transformation as seen in patients. Genetically engineered mouse models of *MYC*-

driven G3 MB derived from *in utero* electroporation of neuronal progenitors or
orthotopic xenografts of cerebellar stem cells have relied on concomitant loss of *p53*
function to evade cell death and induce transformation[74,75], whilst in most human G3
MB tumours *p53* remains functional[1]. We did not observe this effect in our CbO upon
c-MYC overexpression alone, highlighting the importance of modelling the human-
specific cells-of-origin at the correct stage of cerebellar maturation. Overall, we
demonstrate changes in keeping with early onset MB formation upon lineage-specific
disease-defining genetic editing in human CbO, paving the way for future studies
leveraging this model to dissect the events involved in the progression to full blown
G3 and G4 tumours.

In vitro growth of MB cells isolated from patients tissue has proven difficult, particularly
for G3 and G4, thus the number of reliable MB cell lines for use in 2D *in vitro* cell
culture or PDX assays is limited[8,9]. Moreover, *in vitro* expansion can lead to shifts in
molecular subgrouping, an issue that is particularly relevant for G3/G4 MB[4,76]. We
successfully co-cultured both established and primary G3/G4 MB cell lines with CbO,
showing that our 3D co-culture model represents a valuable tool to grow and maintain
primary MB cell lines for which 2D culture has proven challenging by exposing tumour
cells to cell-extrinsic signals from the adjacent CNS tissue. Importantly, we observed
for the first time that MB cells in the 3D CbO-MB model retain an expression profile
closer to that of patient tumours as compared to 2D cultures suggesting that the CbO
environment is superior in supporting and preserving the molecular identity of MB
cells, as has been demonstrated in co-culture models for other brain tumours[77]. Of
note, despite G3/G4 MB typically manifesting post-natally, cases of congenital MB
have been reported[78,79], suggesting that the pre-natal cerebellum which we model in
CbO is a permissive ecosystem for MB formation, proliferation and invasion. While
multiple studies have reported the successful co-culture of primary adult glioblastoma
cells with cerebral organoids[20,21] or cerebellar organoids with established MB cell lines
belonging to other subgroups[29], this is the first time that this approach has been
successfully applied to G3/4 or any primary MB cells.

Analysis of MB cells upon 3D co-culture identified clusters exhibiting neuronal and
glial/stem-like phenotypes, as expected; however a cluster of MB cells displaying
myogenic differentiation was also observed. Historically, this lineage of differentiation
has been occasionally observed histologically in patients' samples[67,68], particularly in
a small proportion of tumours which do not classify within the four accepted WHO

subgroups[80]. Scoring of patient MBs for our CbO-MB derived myogenic signature
revealed the highest enrichment in a subset of G3 patients, with myogenic
differentiation being associated with poorer prognosis. Given the fidelity with which
CbO-MB recapitulate this myogenic subset of G3 MB cells, they may prove a useful
tool to further investigate its clinical relevance.

The characterisation of the MB TME remains limited compared to other brain tumours,
with little known about the interactions or effect of MB tumours on the surrounding
cerebellum[69], however co-culture models have been successfully applied to identify
TME interactions in other primary and metastatic brain tumours[81,82]. Upon MB co-
culture, we observed a specific depletion of more mature neuronal CbO populations.
Whether this effect is directly driven by trophic signalling of co-cultured MB cells or
due to selection pressure within the CbO-MB linked to the highly proliferative nature
of MB cells remains unclear. Receptor-ligand modelling predicted SLC1A7-mediated
glutamate receptor signalling, which had been previously identified specifically in G3/4
MB tumours[83]. It also predicted interactions between MB and various CbO populations
via the transmembrane protein NRP2 (Neuropilin-2) which had been linked to cell
proliferation in an MB mouse model[84]. Proteomic studies have identified upregulation
of proteins involved in TGFβ specifically in the MB TME[85], whilst in our CbO-MB
models we also saw predicted R/L interactions associated with this pathway including
TGFB3/TGFBR3 and INHBE/ACVR2A[86]. We further identified and validated the
interaction of NOTCH1/DLL3 between MB and CbO NSC populations, previously
described to be involved in MB's prognosis, tumour microenvironment and
metastasis[87,88]. Overall, we demonstrate our CbO-MB co-culture is a useful model to
dissect potential TME mediators linked to MB progression and metastasis.

Finally, our CbO-MB model recapitulates the therapeutic response upon treatment
with BMI1/MAPK inhibitors and IP6 as previously demonstrated in xenograft models,
hence supporting its potential value as a system for testing MB subgroup specific and
signature-matched therapeutic approaches. Notably, CbO-MB represent a faster,
scalable and more cost-effective model system as compared to expansion of MB cells
in immunodeficient mice, whilst also offering the possibility to evaluate toxicity on
cerebellar cells induced by anti-tumour therapy, hence could be used in precision
medicine modelling to identify new treatment options that are still lacking for G3/G4
patients.

In conclusion, we have established a new human cerebellar organoid model which
enabled access to human G3/G4 MB cells-of-origin and provided proof of concept that
genetic editing of these cell lineages leads to changes defining MB onset. Moreover,
CbO sustained G3/G4 MB cell growth in a co-culture system, enabling the
identification of a myogenic signature linked to poor prognosis as well as modelling in
3D the tumour microenvironment. Finally, we show its potential value as a novel
scalable pre-clinical screening platform for precision medicine.

Material and Methods

Cell Culture

Full characterisation of the two EPSC lines used in this study has been reported previously[21]. Briefly, EPSC19 and EPSC61 were derived from fibroblasts isolated from the healthy dura mater of two female patients harbouring no somatic mutations and reprogrammed based on established protocols[30]. EPSCs were cultured as described previously[21] on plates coated in Geltrex (Life Sciences) in mTEsR plus basal media (StemCell Technologies). During routine passaging, cells were detached with 0.1 mM EDTA (Life Sciences) and re-plated with media supplemented with 10 µM Y-27632 Dihydrochloride (PeproTech, #1293823). CHLA-01-Med and CHLA-01R-Med MB cell lines purchased from ATCC, were cultured in suspension in Gibco DMEM-F12 media (Life Technologies) supplemented with 20 ng/mL human recombinant FGF and 20 ng/mL human recombinant EGF (Peprotech), 2% B-27 supplement (Invitrogen) and 1% Penicillin-Streptomycin. ICb1299 MB cells obtained from Dr Xiao-Nan Li, Baylor College of Medicine, Texas Children Cancer Centre, USA[11,31] were cultured semi-adherently in Gibco DMEM media (Life Technologies) supplemented with 10% foetal bovine serum (Gibco) and 1% Penicillin-Streptomycin (Life Sciences). Med211-FH cells tagged with GFP were cultured in Neurobasal Plus media (Life Sciences) supplemented with N2 (Bio-techne), B27 (Bio-techne), 75 µg/mL BSA, 2 µg/mL heparin (StemCell Technologies), 10 ng/mL human recombinant FGF and 10 ng/mL human recombinant EGF (Peprotech).

All cell lines were confirmed mycoplasma negative prior to experimental procedures and cultured in humidified incubators at 37°C with 5% CO_2 until ~80% confluent before passaging.

Cerebellar Organoid Culture

Cerebellar organoids were cultured based on a previously published protocol[22,23]. Briefly, EPSCs at 80% confluency were detached using gentle cell dissociation reagent (StemCell) and plated into 96-well ultra-low attachment V-bottomed plates (PHCBI) in mTEsR plus basal media (StemCell) supplemented with 10uM Y-27632 Dihydrochloride (PeproTech, #1293823) at 6,000 cells per well. On Day 1, media was replaced with mTEsR plus basal media. On Day 2, media was replaced with growth-factor free chemically defined media (gfCDM) containing: IMDM (Life Technologies)/Ham's F-12 (Life Technologies) 1:1, 1% v/v chemically defined lipid

concentrate (Life Technologies), 450 µM monothioglycerol (Sigma), 15 µg/ml apo-

transferrin (Sigma), 5 mg/ml crystallization-purified BSA (Sigma), 50 U/ml penicillin/50

µg/ml streptomycin and 7 µg/ml insulin (Sigma). On Day 2-7, 50 ng/ml human

recombinant FGF2 (PeproTech, #100-18B) and 10 µM SB431542 inhibitor (Life

Sciences, #S4317) were added to gfCDM. On Day 7, organoids were transferred to 6-

well ultra-low attachment plates (Corning) on an orbital shaker (90 rpm, Thermo

Scientific, #88881102) in a humidified incubator at 37°C with 5% CO2 where they were

maintained for the remainder of the protocol. For Day 7-14, 33.3 ng/ml human

recombinant FGF2 and 6.67 µM SB431542 inhibitor were added to gfCDM, then 100

ng/ml human recombinant FGF19 (PeproTech, #100-32) was added to gfCDM for Day

14-21 of culture. From Day 21, organoids were cultured in complete neurobasal media

containing: Neurobasal medium (Life Technologies), GlutaMax I (Life Technologies),

N2 supplement (Life Technologies), and 50 U/ml penicillin/50 µg/ml streptomycin. 300

ng/ml human recombinant SDF1 (PeproTech, #300-28A) was added to complete

neurobasal media from Day 28-35.

CbO editing strategy

To induce *c-MYC* expression in putative MB lineages-of-origin within developing

CbOs, three lentivirus constructs were designed utilising the doxycycline inducible Tet-

On system[32]. To target specific CbO lineages, the *rtTA* sequence element and *BFP*

tag were placed under the control of the *EOMES* promoter (p*EOMES*) or a *PAX6* mini

promoter (Ple260[33]) designed for CNS expression (p*PAX6*). In addition, each

construct contained the puromycin resistance gene under the control of a constitutively

expressed promoter (*mPGK)*, giving *pPAX6-rtTA-BFP_mPGK-PuroR* or *pEOMES-*

rtTA-BFP_mPGK-PuroR. The third construct contained the *c-MYC* sequence and *GFP*

tag under the control of the tetracycline responsive element (*TRE)* as well as

blasticidin resistance gene under the control of *mPGK,* giving *TRE-c-MYC-*

GFP_mPGK-BlastR. Lentiviral constructs were produced as described previously[34]

and titrated using Lenti-X GoStix Plus as per manufacturer instructions (Takara Bio).

EPSC19 were infected with *pPAX6-rtTA-BFP_mPGK-PuroR* or *pEOMES-rtTA-*

BFP_mPGK-PuroR (MOI = 1), negative selection for 24 hours performed with

puromycin (1.25 µg/mL) then co-infected with *TRE-c-MYC-GFP_mPGK-BlastR*

construct (MOI = 1) and Blasticidin (5 µg/mL) negative selection performed for 48

hours. Cerebellar organoid protocol was then begun as detailed above. From day 35,

doxycycline (2 µg/mL) was added to culture media and CbOs monitored for 4 weeks.

Two sets of controls were included, non-induced CbOs harbouring the full editing

system (no doxycycline control) as well as CbOs derived from EPSC infected with the

pPAX6-rtTA-BFP_mPGK-PuroR   or   *pEOMES-rtTA-BFP_mPGK-PuroR*   constructs

only and treated with doxycycline from day 35 onwards (doxycycline+ control). On day

63, edited CbO were taken for scRNA-seq analysis, IHC (both detailed below) or

dissociated, FACS sorted for BFP or GFP using BD FACSAria III Cell Sorter then DNA

extracted and submitted for EPICv2 array (outlined below).

CbO-MB co-culture

Prior to co-culture, G3/4 MB cell lines were infected with shCHD7 lentivirus containing

GFP as previously published[35]. On Day 35 of CbO culture, organoids were placed in

96-well ultra-low attachment U-bottomed plates (Corning) in a 1:1 mix of complete MB

cell media and complete neurobasal media with 100,000 MB cells. For Med-211-FH

cells previously cultured as PDX only, cells were thawed then underwent mouse cell

depletion using Miltenyi mouse cell depletion kit (#130-104-694) according to

manufacturer instructions then seeded onto CbO as described above. The following

day, half media from each well was replaced with complete neurobasal media. On Day

37, the entire contents of each well was transferred to a 24-well ultra-low attachment

plate (Corning) with 1ml complete neurobasal media. On Day 4, total media was

replaced with complete neurobasal media to remove non-infiltrating cells and

organoids placed back on the orbital shaker (90 rpm) in a humidified incubator at 37°C

with 5% CO2. Co-cultures were maintained for 14 days in total from MB cell seeding

before taking for drug assays or scRNA-Seq.

Human Cerebellar FFPE Samples Processing

12 human cases from Great Ormond Street Hospital (GOSH) at pre-natal

developmental timepoints (19-37 post-conception weeks) were selected as having

normal cerebellar histology and consent for research. Areas of formalin-fixed paraffin-

embedded (FFPE) tissue containing cerebellar material were identified and manually

macro dissected according to the marking shown in Figure S2A. Following dissection,

tissue was kept at -20°C until downstream processing. In addition, 11 samples of fresh

frozen cerebellar tissue from earlier developmental timepoints (Carnegie stage 22-17

post-conception weeks) were obtained from the Human Developmental Biology

Resource (HDBR).

DNA/RNA Extraction

For cell culture samples (CbO, EPSC and Med211-FH), DNA and RNA were extracted using the RNA/DNA/Protein Purification Plus kit (Norgen, #47700) following manufacturer protocol. For FFPE human cerebellar samples, initial deparaffinization and DNA extraction was performed using FFPE RNA/DNA Purification Plus Kit (Norgen, #54300). Samples were incubated with Proteinase K overnight at 37°C to increase yield, after which DNA extraction was performed as per manufacturer instructions. For fresh frozen human samples, extraction was performed using RNA/DNA Purification Plus Kit (Norgen, #47700) as per manufacturer instructions.

RT-qPCR

All cDNA synthesis was performed using the SuperScript III Reverse Transcriptase kit (Invitrogen) according to the manufacturer's protocol. Analysis of gene expression was carried out using the Applied Biosystems 7500 Real-Time PCR with SYBR Green PCR MasterMix (Applied Biosystems) according to standard protocols. All qPCRs were performed in triplicate with 10ng cDNA. Average Ct values for target genes were normalised to average Ct of ACTIN B and GAPDH housekeeping genes before calculating fold change. Primers used are listed in Table 1.

Immunohistochemistry (IHC)

Organoids were collected and fixed in 4% PFA in PBS, dehydrated in 30% sucrose in PBS and embedded in blocks of 7.5% gelatin, 10% sucrose in PBS. Blocks were then snap frozen in isopentane at -55°C before cryosectioning at 15 μm sections. Prior to staining, sections were de-gelatinised in warm water before permeabilisation in 4% normal donkey serum with 0.25% Triton-X in PBS for 2 hours. Sections were then washed in PBS before incubation with primary antibodies at 4°C overnight diluted in 4% normal donkey serum in PBS. A list of primary antibodies and dilutions are shown in Table 2. Sections were then washed 3 times in PBS followed by incubation for 3 hours at room temperature with corresponding AlexaFluor™ secondary antibodies at 1:500 dilution in PBS. After a final 3 PBS washes, sections were mounted with coverslips using ProLong™ Gold antifade reagent with DAPI (Thermo Fisher Scientific, P36931). Images were taken using Zeiss 800 Laser Scanning Confocal microscope with Zeiss ZEN software v2.3.

CbO-MB drug assays

On Day 0 of drug assays, CbO-MB were transferred to new 24-well ultra-low

attachment plates (Corning) containing 500 μL complete neurobasal media plus drug

(PTC-209/PD325901 inhibitors or IP6) or vehicle (DMSO or PBS respectively) then

placed back on orbital shaker where they remained for the duration of the assay. A

drug and media top up was performed on day 2.

On day 4, the entire content of each well was collected and the Neural Dissociation kit

(MACS Miltenyi Biotec) was used to dissociate treated CbO-MB co-cultures into single

cell suspension. Samples were then stained with the Zombie NIR™ Fixable Viability

Kit (Biolegend) at 1:300 dilution, after which all samples were analysed using the BD

LSRII flow cytometer. GFP-expressing cells were detected with the blue excitation

laser at 488nm, while the Zombie NIR positive cells were detected with the red

excitation at 633nm. Upon gating of single cells, Zombie NIR positive cells were gated

as dead, with GFP fluorescence distinguishing viable MB from CbO cells. All

subsequent FACS analysis was performed in FlowJo v10.6.2.

Dissociation of CbO and CbO-MB for scRNA-Seq

Pooled samples of CbO were taken at day 7 and day 35, as well as edited CbO at day

63 and CbO-MB at day 49. First, organoids were dissociated using the Neural

dissociation kit (MACS Miltenyi Biotec) into single cell suspension. Subsequent debris

removal was then performed using Debris removal solution according to manufacturer

protocol (Miltenyi Biotec, #130-109-398) to obtain a clean sample of highly viable cells.

Cells were resuspended in loading buffer (0.04% BSA in PBS), strained through a 70

μm cell strainer and counted using a hemacytometer with trypan blue staining. At least

10,000 cells per sample were taken for processing using 10X Genomics kits (detailed

below).

RNA sequencing

Med211-FH RNA from cells cultured *in vitro* was sequenced at Novogene with paired-

end, 150bp sequence reads. Library preparation (mRNA, poly-A enrichment) and

sequencing was performed on the NovaSeq6000 platform. Reads were quality

trimmed using trimgalore v0.6.5 and aligned to reference genome

GRCh38.release109 using STAR v2.7.9a[36]. Reads were counted using featurecounts

function and annotated with ENSEMBL Homo_sapiens.GRCh38.109.gtf.

scRNA-Seq Analysis

Library preparation was performed at the Single Cell Genomics Facility (UCL, Cancer

Institute) using the Chromium Next GEM Single Cell 3' Kit v3.1 (10X Genomics,

1000268) along with the Chromium Next GEM Chip G Single Cell Kit (10X Genomics,

1000120) according to the manufacturer's instructions. Pooled samples libraries were

sequenced with lane sequencing on Novaseq 6000 S4 (Novogene). The Cell Ranger

2.0.1 pipeline was used to align reads to the GRCh38 human reference genome and

produce count matrices for downstream pre-processing and analysis using the Seurat

v5.0.1 R package[37].

UMAP calculations and cell type annotations

For analysis of CbO samples we implemented stringent quality filtration, only cells with

fewer than 5×10^4 total reads, lower than 10% mitochondrial reads, more than 750 and

less than 6500 detected features were retained for analysis. Expression values were

library size corrected to 10,000 reads per cell and log1p transformed, with Principal

component analysis (PCA) performed on the scaled data for the top 2000 variable

genes. Batch correction between cell lines was performed on principal components

using Harmony[38]. Uniform Manifold Approximation Projection (UMAP) embeddings,

Nearest Neighbours and cell clusters were then calculated in harmony-corrected PCA

space using 35 dimensions, and cells were clustered using FindClusters(). For Day 7

COs, FindClusters() resolution was set at 0.2, and for Day 35-49, where the UMAP

indicated a greater degree of cell type complexity, we instead set a resolution of 0.5

and used FindSubCluster(resolution = 0.1, algorithm = 1) to separate cluster 7 into two

distinct subclusters 7_0 and 7_1. Cluster marker genes were calculated using a

Seurat's in-bult Wilcoxon Rank Sum test (logfc.threshold = 0.25, min.pct = 0.1,

only.pos = T), and differential expression analysis across experimental conditions was

performed using MAST[39]. Initial clustering identified two clusters 13 and 14 which

represented hypoxic/glycolytic cells, common to all human brain organoids[40]. As has

been done previously[41], these cells were excluded from analyses and a new UMAP

was calculated.

For analysis of engineered CbO and CbO co-cultures containing CHLA-01-Med and

Med211FH MB medulloblastoma (MB) cells or control NSC37.5 cells, we changed the

detected feature threshold greater than 500 and less than 7500, with no batch

correction was necessary. For engineered CbO UMAPs and Nearest neighbours were

calculated using 30 PCA dimensions and FindClusters() was run with a resolution of

0.5. For CbO co-cultures we used 15 PCA dimensions and a resolution of 0.35.

Malignant CHLA-01-Med and Med211FH MB cells were distinguished from co-
cultured CbO cells based on Leiden clustering and inferred copy number variants, with
co-cultured NSC73.5 identified based on clustering and marker gene expression
alone. InferCNV was run on CbO-CHLA-01-Med and CbO-Med211FH samples using
D49 CbO cells as a reference, the "subclusters" analysis mode, and setting the
following parameters: cutoff=0.1, k_obs_groups = 3, leiden_resolution = 1×10^{-5}. For
annotating engineered CbO and organoid cells from CbO-MB co-cultures we used the
processed R object from the D35-49 CbO (Fig1B) as a reference, and the default
settings of Seurat FindTransferAnchors(), TransferData() and AddMetaData()
functions. We then annotated new cluster names based on their composition of
transferred cell type labels Fig S2D-G.

Finally, for the analysis of MB cells alone, cells CHLA-01-Med and Med211FH
identified by InferCNV were subset, and reanalysed using the standard Seurat
pipeline; UMAPs and nearest neighbours were calculated using 20 PCA dimensions
and FindClusters() was run with resolution=0.225.

scRNA-Seq Analysis: Comparison to reference datasets
To approximate the transcriptomic age of our CbO we compared data from this study,
and three other cerebellar organoid studies: Nayler et al[25], Chen et al[26], and Atamian
et al[24] against four datasets derived from developing foetal cerebellum: Aldinger et
al[42], Luo et al[16], Zhong et al[43], and Sepp et al[44]. First, pseudo-bulk transcriptomic data
were generated from aggregated single cell counts for all samples, and expressed as
log(counts per million +1). Data was merged and batch correction between different
sequencing methodologies (scRNAseq vs snRNAseq) was performed using Combat()
from the sva.3.48 package. We next derived an 'ageing-signature' as the top 200
genes correlated with age across the foetal cerebellum samples. This signature was
then used to compute the average Pearson correlation between CbO and foetal
cerebellum samples at different weeks of development.

To aid with D35/49 CbO cell type annotations, we compared cluster marker signatures
against those from the four foetal cerebellum datasets. In each case we obtained
published count matrices, UMAP-embeddings and author-defined cluster meta data,
and used them to perform a Wilcoxon Rank Sum test with Seurat's FindAllMarkers()
command (logfc.threshold = 0.25, min.pct = 0.1, only.pos = T) to obtain the list of

marker genes for their cell types. Then, for each dataset and marker gene, cluster

specificity scores were computed (mean normalized counts per cluster/total mean

normalized counts) – with overlapping gene signature specificity scores compared

across studies by Pearson correlation. We performed the same analysis on CHLA-01-

Med and Med211FH cells to determine each MB cluster's closest analogues from

developing foetal cerebellum samples.

To compare CHLA-01-Med and Med211FH MB cells in co-cultures to patient tumours,

we generated pseudo-bulk transcriptomic data from aggregated single cells counts

from our co-cultured MB samples, and 36 patients in Hovestadt et al[45]. Combat batch

correction was performed between log(cpm+1) transformed data, and Pearson

correlation between samples were calculated across the top 2500 most variable genes

(ranked by standard deviation). Pearson correlation across the same gene list was

also used to query the relative similarity between patient tumours and CbO co-cultured

CHLA-01-Med cells profile here, and CHLA-01-Med cells we previously cultured *in*

vitro and profiled by RNAseq[35]. For Med211FH MB cells, we compared CbO co-

cultured to RNAseq data from an *in vitro* sample drawn from the literature[17] and a

second *in vitro* sample profiled as part of this study.

scRNA-Seq Analysis: Reference gene set/signature scoring

For reference signature scoring, average gene module expression was calculated for

each single cell, subtracted by the aggregated expression of a random control set of

features selected from the same average expression bins as the query genes[46] using

Seurat's AddModuleScore() function. G/S and G2M gene modules are included in the

Seurat v5.0.1 package; G3 RL.PRC and G4 RL.UBC Medulloblastoma cell of origin

signatures were taken from Smith et al[14]; Group 3 and 4 MB tumour signatures were

taken from Vladoiu et al[47]; WNT, SHH and G3/4 MB cell state signatures were taken

from Hovestad et al[45]. For gene set enrichment analysis, gene sets were obtained

from MolSigDB[48] and computations were performed using the R package fGSEA

v4.4[49]. Gene ontology enrichment analysis of was performed on upregulated

(p.adj<0.05, and avg_log2FC>2) genes in the MYC^{oe} cluster of engineered CbOs

using the topGO v2.52.0 and org.Hs.eg.db v3.17.0 R packages.

Processed cell count and metadata from patient-derived MB samples was accessed

from Hovestadt et al[45] and reanalysed in Seurat using default settings. The resulting

R object was then used as a reference, with the default settings of Seurat

FindTransferAnchors(), TransferData() and AddMetaData() functions to assign MB

Group3 and Group4 prediction scores to the co-cultured CHLA-01-Med and

Med211FH MB cells. To generate a Myogenic MB state signature, we took the

intersect of significant MB marker genes with avg_log2FC>1 from cluster 4 (CHLA-01-

Med) and cluster 6 (Med211FH). The processed reference R object from Hovestadt et

al[45] was then scored for the Myogenic MB state signature as detailed above using the

the AddModuleScore() function. Next, the Hovestadt et al[45] data was subset to group

3 samples, before Harmony[38] batch correction between samples was performed. New

UMAP embeddings, Nearest Neighbours were then calculated in harmony-corrected

PCA space using 15 dimensions, and clusters were identified using a resolution of 0.5.

We also inspected a large reference dataset of 763 bulk MB RNAseq samples (Cavelli

et al[1]) for expression of the Myogenic MB state signature using the GSVA 1.48.3 R

package[50]. Survival analysis was performed on patients with complete data (i.e. that

had died) using the survival_3.8-3 and survminer_0.5.0 R packages and splitting

patients into the Myogenic Signature High or Low groups based on the median

signature score.

Cell lineage trajectory analysis

For RNAvelocity analysis, velocyto[51] was used to calculate spliced and unspliced

transcriptomes, before subsequent analysis using scVelo[52], on Seurat-defined cell

types and UMAP embeddings, using default parameters. For single cell lineage

trajectory analysis, we imported the processed Seurat object into the Monocle3 R

package. The 'learn_graph' function was applied to infer single cell trajectories setting

close_loop = TRUE, minimal_branch_len to 10 and euclidean_distance_ratio to 1.5.

The 'order_cells' function was then used to set Progenitors and Cyc_Progenitors as

the root of states of pseudotime. Next we subset data to cell types involved in the G3

trajectory (early_RL/NSCs, glut_progenitors and matCN2) or G4 trajectory (NSCs,

GN/UBC_progs and GN/UBC)[53]. For each subset, the 'learn_graph' function was

applied again to infer single cell trajectories setting minimal_branch_len to 10 and the

'order_cells' function was used to select early_RL/NSCs and NSCs as the root states

of pseudotime. Differential gene expression was calculated across single cell

trajectories using Moran's I test. For each trajectory, the top 2500 most differentially

expressed genes over pseudotime were selected, and smoothened, alra-imputed[54]

expression over pseudotime was plotted.

Receptor Ligand Modelling

For receptor-ligand analysis, cell type meta data and count data for the top 15000 most highly variable genes in the CbO-MB dataset was input into CellphoneDB v5[55] and run using the statistical_analysis mode.

DNA methylation analysis

All CbO and human cerebellar DNA samples underwent standard BSC conversion at UCL Genomics facility (UCL, Great Ormond Street Institute of Child Health) and were submitted for Illumina Methylation EPIC v2 array following standard protocols. Quality control was performed and probe beta values were extracted from raw .idat files using wateRmelon R package[56] under reference genome hg38. Probes with detection p-values greater than 0.05 were removed using pfilter(), as well as probes associated with sex chromosomes, mitochondrial chromosomes and single nucleotide polymorphisms. Adjusted Dasen normalisation was then performed.

Cerebellar Organoid developmental staging (DNA methylation)

To stage the development of CbO compared with human cerebellar samples, an 'ageing signature' approach was applied, with the top 1% of probes whose beta values best correlated to age in human cerebellar samples selected for subsequent correlation with CbO probe betas. Combat batch correction was first performed between CbO and human samples before top 1% probe selection and correlation to CbO. To corroborate this approach, CbO methylation data adapted to EPIC v1 array probe annotation was also passed to the foetal brain clock (FBC), an R package specifically trained on foetal brain samples during development to estimate their epigenetic age[57].

Differentially and variably methylated probes and regions analysis

Differentially methylated probes (DMPs) and regions (DMRs) were calculated pairwise for CbO samples between comparison groups using the R package DMRcate[58] with FDR threshold of 0.01. Additional criteria for DMR identification were set such that DMRs must contain at least 6 CpG sites with <400 base pair separation. Variable methylated probes (VMPs) and regions (VMRs) across CbO and human development timepoints were identified within the DMRcate package using the same thresholds as stated above for DMR analysis and the top 5% VMP/VMRs defined as significant.

VMRs were then annotated to genes (VMGs), after which significant VMGs were

clustered based on z-scores of average beta-values of the probes contained within

them and taken for pathway analysis.

DNA methylation: Comparison to reference datasets

For human rhombic lip (RL) and external granular layer (EGL) correlation, raw .idat

files for micro-dissected RL and EGL samples were accessed from GSE207266[14] and

processed using the same raw data processing pipeline adapted for EPIC v1 array

annotation before downstream analysis. DMPs were identified between each

cerebellar compartment (n=7) as stated above with FDR threshold of 0.01. DMPs were

then subset to CbO average probe betas adapted to EPIC v1 array annotation and

correlated to each cerebellar compartment.

For CbO editing samples, reference 450K methylation data of 763 MB samples were

accessed from GSE85218[1]. CbO datasets were trimmed to probes present in both

450K and EPICv2 array and clustering of CbO and MB samples was performed based

on a published probe set shown to distinguish between MB subgroups[59] (34/48 CpG

probes present in QC filtered datasets). DMPs were calculated as described above on

the trimmed CbO dataset then correlated to patient MB samples.

scRNA-Seq/DNA methylation integration and pathway analysis

Differentially expressed genes (DEGs) between Day 7 and Day 35 CbO scRNA-Seq

data with adjusted p-value < 0.05 and absolute log2FC > 0.25 were integrated with

genes mapping to DMRs from methylation data at matched timepoints with p-value <

0.05 and greater than a 10% imposed threshold in methylation difference. Genes

demonstrating concordant hypermethylation and down-regulation or hypomethylation

and up-regulation were taken for subsequent pathway analysis. For pathway analysis,

gene lists were input into Gprofiler [60] and pathways calculated based on a Benjamini-

Hochberg FDR < 0.05. Pathway visualisation was performed using CytoScape [61]. For

scRNA-Seq upon MB co-culture pathway analysis, UMAP clusters were grouped

based on cell developmental state (un/differentiated) and effect of MB co-culture on

cluster percentage (increase/decrease). Common pathways from each group were

then identified and the top 100 most significant (adjusted-p<0.05) Reactome and

KEGG pathways were plotted. For VMR pathway analysis, the top 100 most significant

(adjusted-p<0.05) GO:Biological Process and GO:Molecular Function pathways were

plotted for human and CbO VMG clusters, whilst for overlapping pathways with

concordant DMR/DEG integration of Day 7 vs Day 35 CbO samples, GO:Cell
Compartments were also plotted.
988

989   **Ethics**
990   Ethical approval for the work on human cerebellar samples was obtained from Brain
991   UK, reference number 22/010.
992

993   *Table 1 Primers used for RT-qPCR reactions*

Gene	Forward Primer (5'-3')	Reverse Primer (5'-3')
ACTIN B	GCGAGAAGATGACCCAGATC	CCAGTGGTACGGCCAGAGG
GAPDH	CAATGACCCCTTCATTGACC	GACAAGCTTCCCGTTCTCAG
NESTIN	GAAACAGCCATAGAGGGCAAA	TGGTTTTCCAGAGTCTTCAGTGA
SOX1	TGCTTGTTCTGTTAACTCAC	AAAGAACCTCAGAGAGAGTC
EN2	CCGGCGTGGGTCTACTGTA	GGCCGCTTGTCCTCTTTGTT
OTX2	AGAGGACGACGTTCACTCG	TCGGGCAAGTTGATTTTCAGT
KIRREL2	ACAACTCTCCTTATGGTCATC	ATTCGCATCAGGTTCTTTTG
OLIG2	GACAAGCTAGGAGGCAGTGG	CGGCTCTGTCATTTGCTTCTTG
SKOR2	CCAGGTGTTAAAAGGAAACACA	GCTCCCTTTTCATCTGATCCT
PAX6	AGAGAATACCAACTCCATCAG	GATAATGGGTTCTCTCAAACTC
ATOH1	TGTTATCCCGTCGTTCAACAAC	TGGGCGTTTGTAGCAGCTC
BARHL1	GAGCGGCAGAAGTACCTGAG	GTAGAAATAAGGCGACGGGAAC
EOMES	CACATTGTAGTGGGCAGTGG	CGCCACCAAACTGAGATGAT
MYC	AGTGGAAAACCAGCAGCCTC	TTCTCCTCCTCGTCGCAGTA
GATA6	GCCAACTGTCACACCACAAC	TCATAGCAAGTGGTCTGGGC
NANOG	AGAAAAACAACTGGCCGAAGAAT	GTTGAATTGTTCCAGGTCTGGTT

996   *Table 2 Primary antibodies and dilutions used for immunohistochemistry*

Primary Antibody	Dilution
SOX2 (abcam, ab97595)	1:500
OLIG2 (Santa Cruz, sc-19967)	1:200
CALB (Swant, CB300)	1:200
NeuN (Chemicon, MAB377)	1:500
EOMES (Chemicon, AB15894)	1:200
KI67 (abcam, ab15580)	1:500
GFP (Chemicon, AB16901)	1:400

cCASP3 (Cell Signalling, 9664)	1:500
MAP2 (Sigma, M4403)	1:200
TubIII (abcam, ab7751)	1:200
Myogenin (abcam, ab1835)	1:100

Acknowledgements

This work is funded by grants from Brain Tumour Research (Centre of Excellence award to S.M.), Cancer Research UK (C23985/A29199 programme award to S.M.), Barts Charity (MGU0447 programme grant to S.M.). We thank Gary Warnes (Blizard Flow Cytometry Core Facility), Maeve McLaughlin (Blizard Advanced Light Microscopy Facility), Imran Uddin (Single Cell Genomics Facility, UCL, Cancer Institute), Myrianni Constantinou and Sara Lucchini for sharing their expertise. The collection and annotation of FFPE human tissue samples are funded by the NIHR GOSH BRC. The collection of fresh frozen human tissue samples is funded by the MRC-Wellcome Trust HDBR.

Author contribution

SB and SM conceived and supervised the project; TW, SB and SM designed the experiments; TW performed, analysed and interpreted all experiments and data; SB analysed and interpreted all experiments and data; JN analysed and interpreted scRNA-seq data; TW and YX analysed DNA methylation data; OO, TM and AM identified and provided suitable human foetal cerebellar samples; BD and LD shared G3 MB primary cells, datasets and expertise; NRZ contributed expertise for DNA methylation analysis; SM secured financial support; TW, SB, JN and SM wrote the paper with contributions from all authors.

Competing interests

The authors declare no competing interests for this study.

Data Availability

The authors declare that all the data supporting the findings of this study are available within the article and its supplementary files. The scRNA-seq and DNA methylation datasets generated in this study and processed data are available in the NCBI Gene Expression Omnibus database (GSE263652 and GSE263653 respectively) or are available from the corresponding authors upon reasonable request. Publicly available datasets used in the study were retrieved from relevant databases as outlined in materials and methods.

1030

1031 **Code Availability**

1032 The authors declare that no custom code was used in this study. A full description of

1033 analyses including functions and packages used are stated in the materials and

1034 methods. Specific code will be made available from the corresponding authors upon

1035 reasonable request.

1036

1037 **Figures and Legends**

1038

1039

1040

1041

Figure 1. CbO is composed of diverse cerebellar lineages and displays
characteristic structural organisation.

A. Schematic of CbO samples used to assess expression of cerebellar markers.

B. H&E staining of CbO_19 and CbO_61 showing formation of characteristic flat-
oval neural rosettes within each structure.

C. Heatmaps showing average expression of marker genes associated with
cerebellar differentiation up to Day 35 (D35) of development for CbO_19 and
CbO_61 derived from two independent EPSC lines (D0). Heatmaps show Z-
scores for average dCT values from n = 3 independent batches of CbO.

D. Representative immunofluorescence staining of OLIG2$^+$ GABAergic
progenitors (red), Calbindin$^+$ (CALB$^+$) mature neurons (pink) and SOX2$^+$
neural stem and progenitor cells (green) in D35 CbO_61.

E. Representative immunofluorescence staining of NeuN$^+$ glutamatergic
neuronal cells (red) and SOX2$^+$ neural stem and progenitor cells (green) in
D35 CbO_61.

F. Schematic of CbO samples used in single cell RNA-Seq (scRNA-Seq)
analysis.

G. Uniform manifold approximation and projection (UMAP) plot of scRNA-Seq
data from CbO_19 and CbO_61 at Day 35 and CbO_61 at Day 49 of
differentiation (left) and bar plots showing percentages of annotated cell
populations identified (right).

H. Dot plot showing key marker genes used to identify major cell types in CbO
(see also Figure S1M).

Scale bars = 150 µm (B) and 100 µm (D and E).

1066

1067

1068

1069

Figure 2. CbO development is epigenetically regulated similarly to human
cerebellum.

A. Schematic of CbO samples used to assess DNA methylation during
development.

B. Heatmap showing k-means clustering (n=5) of significant variable methylated
genes across 35 days of CbO maturation. Mean beta values were taken for
each timepoint from CbO_19 and CbO_61. Z-scores for average beta values
of probes contained within each gene are plotted, with key cerebellar
developmental and neuronal genes annotated.

C. Bubble plot showing top 100 significant (adjusted p-value < 0.05) GO
Biological Processes and Molecular Functions of genes from 5 clusters shown
in **B**. Bubbles are coloured based on each cluster and size is proportional to
number of genes in specific GO term.

D. Venn diagram showing 1086 common variably methylated genes (VMGs)
calculated through human (Carnegie stage 22 to 37 post conception weeks)
and CbO (CbO_19 and CbO_61, Day 7 to 35) development (Upper). Pearson
correlation of human and CbO samples on their beta values for 1086 common
VMGs.

E. Volcano plots of differentially expressed (DEG) and differentially methylated
(DMG) genes between D35 and D7 of CbO maturation. Orange and green
dots represent upregulated/hypomethylated and
downregulated/hypermethylated genes respectively. Genes found in VMG
clusters in B are highlighted.

1095

1096

1097

Figure 3. CbO displays spatiotemporal expression of G3/G4 MB cell-of-origin.

A. Heatmap showing correlation of methylation status of top 1% (6886) age-dependent CpG probes between human developing cerebellum samples and CbO (see also Figure S3A).

B. Heatmap showing correlation of top 1% (200) age-related genes between pseudo-bulk RNA-Seq of CbO and cerebellar organoids from previously published protocols[24-26] and human foetal cerebellar samples[16,42-44] calculated similarly to described above.

C. UMAP plot of scRNA-Seq data from CbO at Day 35-49 of differentiation showing signature score for rhombic lip subventricular zone (RL^{SVZ}) signature genes[15].

D. Heatmap showing correlation of average beta values of CbO to human microdissected rhombic lip (RL) and external granule layer (EGL) regions[15] for differentially methylated probe set calculated between RL and EGL samples.

E. UMAP plot of scRNA-Seq data from CbO at Day 35-49 of differentiation showing signature score for unipolar brush cell (UBC) RL^{SVZ} signature genes[15].

F. UMAP plot of scRNA-Seq data from CbO at Day 35-49 of differentiation showing signature score for photoreceptor RL^{SVZ} signature genes[15].

G. Representative immunohistochemistry pictures of D35 CbO_61 showing a neural rosette with cells positive for EOMES (green) and Ki67 (red). Nuclei are sown in grey.

H. Quantification of staining shown in G of Ki67+EOMES+ nuclei with respect to CbO neural rosette structures. students t-test ****p<0.0001. n = 5 regions of interest quantified.

I. Representative immunohistochemistry pictures of D35 CbO_61 showing a neural rosette with cells positive for PRTG (green), SOX2 (red) and NESTIN (light blue). Nuclei are shown in grey.

J. Quantification of staining shown in I of SOX2+PRTG^{High} nuclei with respect to CbO neural rosette structures. PRTG^{High} was defined as the top 50% of fluorescence signal for each image. students t-test ****p<0.0001. n = 5 regions of interest quantified.

K. Defined Group 3 (G3) and 4 (G4) medulloblastoma cell-of-origin trajectories selected for further analysis (G3 = early_RL/NSC, glut_progs, matCN2; G4 = NSCs, GN/UBC_progs, GN/UBC; see Figure 1H). Expression of genes associated with G3/4 medulloblastoma across each of the G3 and G4 trajectories.

Figure 4. Edited medulloblastoma lineages-of-origin display features of MB
dysregulation upon MYC overexpression

A. Schematic of cerebellar organoid (CbO) editing strategy (see Methods).

B. Representative immunohistochemistry staining for GFP (green) and Ki67
(red) (left) and haematoxylin and eosin staining (right) for CbO_19 collected
at Day 63 of differentiation for dox-induced *pEOMES;MYC-GFP* and
pPAX6;MYC-GFP organoids.

C. UMAP plot of scRNA-Seq data from *pEOMES;MYC-GFP, pPAX6;MYC-GFP*
and control CbOs at Day 63 of differentiation and colour by cell type.

D. UMAP shown in D with *MYC* expression plotted.

E. Gene ontology enrichment bar plot of MYC[OE] cluster marker genes.

F. UMAP plot of control D35/40 CbO scRNA-Seq data highlighting Group3 (red)
and Group4 (green) MB lineages of origins (Top). Projection of MYC[OE] cells
from D63 *pEOMES;MYC*-GFP (blue) *and pPAX6;MYC-GFP* (purple) CbOs
onto control UMAP embeddings.

G. Dot plot showing G3 and G4 lineage-of-origin marker and MB-associated
genes in G3/4 lineage genes compared to MYC[OE] cells from D63
pEOMES;MYC-GFP and pPAX6;MYC-GFP CbOs.

H-I. Violin plot showing mean signature scores for WNT, SHH or G3/G4 MB
subgroups[45] in G3/4 lineages-of-origin and MYC[OE] cells from D63
pEOMES;MYC-GFP and pPAX6;MYC-GFP cells CbOs.

J. Gene set enrichment plot of the
POMEROY_MEDULLOBLASTOMA_PROGNOSIS_DN gene set in the
MYC[OE] cell compared to G3/4 lineages-of-origin, NES=3.46, padj=2.09E-11

K. Heatmap showing z-scored average beta values of probes contained within
differentially methylated genes between MYC[OE] (*pEOMES;MYC-GFP,*
pPAX6;MYC-GFP) and control medulloblastoma lineage-of-origin samples
(*pEOMES*.ctrl, *pPAX6*.ctrl). Key cerebellar and medulloblastoma associated
genes are annotated.

L. Scatter plot of differentially expressed (DEG) and differentially methylated
(DMG) genes between MYC[OE] cells and G3/4 cell-of-origin. Orange and green
dots represent upregulated/hypomethylated and
downregulated/hypermethylated genes respectively.

1174

1175

1176

1177

Figure 5. CbO enables MB growth whilst also retaining MB transcriptional
profiles closer to those of patient tumour samples.

A. Schematic of protocol used to co-culture cerebellar organoids (CbO) with
medulloblastoma (MB) showing CbO-MB sample used for scRNA-Seq (left
panel). Representative immunofluorescence staining of GFP⁺ CHLA-01-Med
MB cells upon one, four, seven and fourteen days of co-culture with CbO_61,
scale bars = 100 µm (right panels).

B. Box and whisker plot showing percentage of MB cells (GFP⁺) expressing the
proliferative markers Ki67 upon 7 and 14 days of co-culture in CHLA-01-Med
and Med211FH cells. Graphs report mean ± SEM, unpaired t-test, n = 6
independent CbO-MB analysed, ns = not significant (left). Representative
immunofluorescence staining of GFP⁺ and KI67⁺ MB cells upon 14 days of co-
culture with CbO, scale bars = 100 µm (right).

C. UMAP plot of scRNA-Seq data from D49 control CbO_61, CbO-Med211-FH,
CbO-CHLA-01-Med and control CbO-NSC73_5 co-cultures, coloured by
genotype (left). Barplot showing proportion of cells from each genotype in the
different experimental models (right).

D. Violin plot showing signature scores for WNT, SHH or G3/G4 MB subgroups[45]
in CbO-MB. One-way ANOVA ****$p<0.0001$.

E. Heatmap showing correlation scores between expression of pseudo-bulk
RNA-Seq of MB fractions only from CbO-Med211-FH and CbO-CHLA-01-Med
and reference MB samples. Clustering based on expression profile has been
used to assess molecular subgroup of CbO-MB models.

F. Boxplots plot showing correlations between same MB cell line grown as 2D
(red) or 3D (blue) model and MB samples shown in D. One-way ANOVA
****$p<0.0001$.

1206

1207

1208

1209

Figure 6 CbO-MB uncovers a subset of Group3 MB cells with a myogenic
transcriptional state.

A. UMAP plot of scRNA-Seq data from MB cells only from D49 CbO-Med211-FH
and CbO-CHLA-01-Med CbO-MB co-cultures, coloured by genotype (top),
and Louvain clusters (bottom).

B. Barplot showing proportion of MB cells from each CbO-MB co-culture in each
cluster.

C. Dot plot showing key marker genes used to identify MB cell types in MB cells
only from D49 CbO-Med211-FH and CbO-CHLA-01-Med CbO-MB co-
cultures.

D. Heatmap showing Pearson correlation of MB cell clusters across the top 250
specific markers for each cluster.

E. Venn diagram showing the intersect of CbO-Med211-FH cluster 8 and CbO-
CHLA-01-Med cluster 5 marker genes (left). Gene ontology enrichment bar
plot of the 44 shared markers between clusters 8 and 5 (right).

F. Venn diagram showing the intersect of CbO-Med211-FH cluster 6 and CbO-
CHLA-01-Med cluster 4 marker genes (left). Gene ontology enrichment bar
plot of the 127 shared markers between clusters 6 and 4 (right).

G. Immunohistochemistry of CbO-Med211-FH co-cultures stained for GFP
(green) and myogenin (red). Scalebar = 10 µm.

H. Violin plot showing signature scores for the myogenic signature in different
MB subgroups of a reference cohort of patient MB scRNA-Seq data[45].

I. UMAP plot of Group 3 MB reference samples in H, coloured by myogenic
signature score, and labelled by their Louvain cluster.

J. Barplot showing the relative proportion of myogenic (clusters 4 and 6) vs non-
myogenic cells in Group 3 MB reference samples in H.

K. Violin plot showing signature scores for the myogenic signature in different
MB subgroups of a reference cohort of patient MB RNA-seq data[1].

L. Violin plot showing signature scores for the myogenic signature in different
MB subtypes of reference cohort of patient MB RNA-seq data in K

M. Survival curve showing survival data of MB patients shown in K, split by
myogenic high (> median, red) or myogenic low (< median, blue) signature
score.

1244

Figure 7. CbO-MB reveal cell-cell interactions and molecular events induced by MB growth and they represent a suitable model for drug screening.

A. UMAP plot of scRNA-Seq data from D49 control CbO_61, CbO-Med211-FH, CbO-CHLA-01-Med and control CbO-NSC73_5 co-cultures, coloured by celltype.

B. Barplot showing proportion of CbO cell types only present in control CbO_61, CbO-Med211-FH, CbO-CHLA-01-Med and control CbO-NSC73_5 co-cultures.

C. Immunohistochemistry images of cerebellar organoids co-cultured with Med211FH for 14 days stained for GFP (green), cCASP3 (white) and MAP2(red, left) or TubIII (red, right). Scale bars = 10 μm.

D. Venn diagram showing the shared significant receptor-ligand pairs modelled between co-cultured CbO-Med211-FH, CbO-CHLA-01-Med and CbO-NSC73_5 and CbO_61 cell types. Shared MB-specific receptor-ligand pairs are highlighted in bold and plotted in panel E.

E. Dot plot showing significant 27 shared MB-specific receptor-ligand pairs identified by CellPhoneDB analysis interaction are separated by cell line CbO-CHLA-01-Med (top) CbO-Med211-FH (bottom), and direction MB cells as a source (left) or as a target (right). Points size corresponds to the co-expression value of the receptor-ligand pair and point colour corresponds to the interacting CbO cell type.

F. Immunohistochemistry images of cerebellar organoids co-cultured with CHLA-01-Med MB cells for 14 days stained for predicted R/L interactions in E. Upper panel: NOTCH1 (green), SOX2 (red) and GFP (white); Lower panel: NOTCH1 (green), DLL3 (red) and GFP (white). Scale bars = 5 μm.

G. Schematic of combination treatment with PTC209 (BMI1i) and PD329501 (MAPKi) or IP6 of CbO-CHLA-01-Med co-cultures.

H. Viability assays of GFP$^+$ MB cells (green) and GFP$^-$ CbO cells (pink) upon 4 days of treatment with increasing concentrations of BMI1 and MAPK inhibitors. Measurement of Area Under Curve (AUC) to compare the overall response to treatment. n= 9 independent experiments. graphs report mean ± SEM, unpaired t-test ****$p<0.0001$

I. Quantification (left) and representative images (right) of vehicle and treated CbO-CHLA-01 co-cultures from H stained for GFP and cCASP3. graphs report mean ± SEM, paired t-test **$p<0.01$; ns = not significant. n = 6 regions of interest quantified. Scale bars = 10 μm.

J. Viability assays of GFP$^+$ MB cells (green) and GFP$^-$ CbO cells (pink) upon 4 days of treatment with increasing concentrations of IP6. Measurement of Area Under Curve (AUC) to compare the overall response to treatment. n= 5 independent experiments. graphs report mean ± SEM, unpaired t-test ****$p<0.0001$

K. Quantification (left) and representative images (right) of vehicle and treated CbO-CHLA-01-Med co-cultures from J stained for GFP and Ki67. graphs report mean ± SEM, paired t-test **$p<0.01$; ns = not significant. n = 5 regions of interest quantified. Scale bars = 10 µm.

References

1. Cavalli, F.M.G., Remke, M., Rampasek, L., Peacock, J., Shih, D.J.H., Luu, B., Garzia, L., Torchia, J., Nor, C., Morrissy, A.S., et al. (2017). Intertumoral Heterogeneity within Medulloblastoma Subgroups. Cancer Cell *31*, 737-754 e736. 10.1016/j.ccell.2017.05.005.

2. Northcott, P.A., Buchhalter, I., Morrissy, A.S., Hovestadt, V., Weischenfeldt, J., Ehrenberger, T., Grobner, S., Segura-Wang, M., Zichner, T., Rudneva, V.A., et al. (2017). The whole-genome landscape of medulloblastoma subtypes. Nature *547*, 311-317. 10.1038/nature22973.

3. Schwalbe, E.C., Lindsey, J.C., Nakjang, S., Crosier, S., Smith, A.J., Hicks, D., Rafiee, G., Hill, R.M., Iliasova, A., Stone, T., et al. (2017). Novel molecular subgroups for clinical classification and outcome prediction in childhood medulloblastoma: a cohort study. The Lancet Oncology *18*, 958-971. 10.1016/s1470-2045(17)30243-7.

4. Sharma, T., Schwalbe, E.C., Williamson, D., Sill, M., Hovestadt, V., Mynarek, M., Rutkowski, S., Robinson, G.W., Gajjar, A., Cavalli, F., et al. (2019). Second-generation molecular subgrouping of medulloblastoma: an international meta-analysis of Group 3 and Group 4 subtypes. Acta Neuropathol *138*, 309-326. 10.1007/s00401-019-02020-0.

5. Mueller, S., and Chang, S. (2009). Pediatric brain tumors: current treatment strategies and future therapeutic approaches. Neurotherapeutics *6*, 570-586.

6. Rechberger, J.S., Toll, S.A., Vanbilloen, W.J.F., Daniels, D.J., and Khatua, S. (2023). Exploring the Molecular Complexity of Medulloblastoma: Implications for Diagnosis and Treatment. Diagnostics (Basel) *13*. 10.3390/diagnostics13142398.

7. Thompson, E.M., Ashley, D., and Landi, D. (2020). Current medulloblastoma subgroup specific clinical trials. Transl Pediatr *9*, 157-162. 10.21037/tp.2020.03.03.

8. Badodi, S., Marino, S., and Guglielmi, L. (2019). Establishment and Culture of Patient-Derived Primary Medulloblastoma Cell Lines. In Brain Tumor Stem Cells, S. Singh, and C. Venugopal, eds. (Humana Press). https://doi.org/10.1007/978-1-4939-8805-1_3.

9. Ivanov, D.P., Coyle, B., Walker, D.A., and Grabowska, A.M. (2016). In vitro models of medulloblastoma: Choosing the right tool for the job. Journal of Biotechnology *236*, 10-25. 10.1016/j.jbiotec.2016.07.028.

10. Sanden, E., Dyberg, C., Krona, C., Gallo-Oller, G., Olsen, T.K., Enriquez Perez, J., Wickstrom, M., Estekizadeh, A., Kool, M., Visse, E., et al. (2017). Establishment and characterization of an orthotopic patient-derived Group 3 medulloblastoma model for preclinical drug evaluation. Sci Rep *7*, 46366. 10.1038/srep46366.

11. Shu, Q., Wong, K.K., Su, J.M., Adesina, A.M., Yu, L.T., Tsang, Y.T., Antalffy, B.C., Baxter, P., Perlaky, L., Yang, J., et al. (2008). Direct orthotopic transplantation of fresh surgical specimen preserves CD133+ tumor cells in clinically relevant mouse models of medulloblastoma and glioma. Stem Cells *26*, 1414-1424. 10.1634/stemcells.2007-1009.

12. Brabetz, S., Leary, S.E.S., Grobner, S.N., Nakamoto, M.W., Seker-Cin, H., Girard, E.J., Cole, B., Strand, A.D., Bloom, K.L., Hovestadt, V., et al. (2018). A biobank of patient-derived pediatric brain tumor models. Nat Med *24*, 1752-1761. 10.1038/s41591-018-0207-3.

13. Marino, S. (2005). Medulloblastoma: developmental mechanisms out of control. Trends Mol Med *11*, 17-22. 10.1016/j.molmed.2004.11.008.

14. Smith, K.S., Bihannic, L., Gudenas, B.L., Haldipur, P., Tao, R., Gao, Q., Li, Y., Aldinger, K.A., Iskusnykh, I.Y., Chizhikov, V.V., et al. (2022). Unified rhombic lip origins of group 3 and group 4 medulloblastoma. Nature *609*, 1012-1020. 10.1038/s41586-022-05208-9.

15. Hendrikse, L.D., Haldipur, P., Saulnier, O., Millman, J., Sjoboen, A.H., Erickson, A.W., Ong, W., Gordon, V., Coudiere-Morrison, L., Mercier, A.L., et al. (2022). Failure of human rhombic lip differentiation underlies medulloblastoma formation. Nature *609*, 1021-1028. 10.1038/s41586-022-05215-w.

16. Luo, Z., Xia, M., Shi, W., Zhao, C., Wang, J., Xin, D., Dong, X., Xiong, Y., Zhang, F., Berry, K., et al. (2022). Human fetal cerebellar cell atlas informs medulloblastoma origin and oncogenesis. Nature *612*, 787-794. 10.1038/s41586-022-05487-2.

17. Visvanathan, A., Saulnier, O., Chen, C., Haldipur, P., Orisme, W., Delaidelli, A., Shin, S., Millman, J., Bryant, A., Abeysundara, N., et al. (2024). Early rhombic lip Protogenin(+ve) stem cells in a human-specific neurovascular niche initiate and maintain group 3 medulloblastoma. Cell *187*, 4733-4750 e4726. 10.1016/j.cell.2024.06.011.

18. Bian, S., Repic, M., Guo, Z., Kavirayani, A., Burkard, T., Bagley, J.A., Krauditsch, C., and Knoblich, J.A. (2018). Genetically engineered cerebral organoids model brain tumor formation. Nat Methods *15*, 631-639. 10.1038/s41592-018-0070-7.

19. Choe, M.S., Kim, J.S., Yeo, H.C., Bae, C.M., Han, H.J., Baek, K., Chang, W., Lim, K.S., Yun, S.P., Shin, I.S., and Lee, M.Y. (2020). A simple metastatic brain cancer model using human embryonic stem cell-derived cerebral organoids. FASEB J *34*, 16464-16475. 10.1096/fj.202000372R.

20. Linkous, A., Balamatsias, D., Snuderl, M., Edwards, L., Miyaguchi, K., Milner, T., Reich, B., Cohen-Gould, L., Storaska, A., Nakayama, Y., et al. (2019). Modeling Patient-Derived Glioblastoma with Cerebral Organoids. Cell Rep *26*, 3203-3211 e3205. 10.1016/j.celrep.2019.02.063.

21. Vinel, C., Rosser, G., Guglielmi, L., Constantinou, M., Pomella, N., Zhang, X., Boot, J.R., Jones, T.A., Millner, T.O., Dumas, A.A., et al. (2021). Comparative epigenetic analysis of tumour initiating cells and syngeneic EPSC-derived neural stem cells in glioblastoma. Nat Commun *12*, 6130. 10.1038/s41467-021-26297-6.

22. Silva, T.P., Bekman, E.P., Fernandes, T.G., Vaz, S.H., Rodrigues, C.A.V., Diogo, M.M., Cabral, J.M.S., and Carmo-Fonseca, M. (2020). Maturation of Human Pluripotent Stem Cell-Derived Cerebellar Neurons in the Absence of Co-culture. Front Bioeng Biotechnol *8*, 70. 10.3389/fbioe.2020.00070.

23. Muguruma, K., Nishiyama, A., Kawakami, H., Hashimoto, K., and Sasai, Y. (2015). Self-organization of polarized cerebellar tissue in 3D culture of human pluripotent stem cells. Cell Rep *10*, 537-550. 10.1016/j.celrep.2014.12.051.

24. Atamian, A., Birtele, M., Hosseini, N., Nguyen, T., Seth, A., Del Dosso, A., Paul, S., Tedeschi, N., Taylor, R., Coba, M.P., et al. (2024). Human cerebellar organoids with functional Purkinje cells. Cell Stem Cell *31*, 39-51 e36. 10.1016/j.stem.2023.11.013.

25. Nayler, S., Agarwal, D., Curion, F., Bowden, R., and Becker, E.B.E. (2021). High-resolution transcriptional landscape of xeno-free human induced pluripotent stem cell-derived cerebellar organoids. Sci Rep *11*, 12959. 10.1038/s41598-021-91846-4.

26. Chen, Y., Bury, L.A., Chen, F., Aldinger, K.A., Miranda, H.C., and Wynshaw-Boris, A. (2023). Generation of advanced cerebellar organoids for

neurogenesis and neuronal network development. Hum Mol Genet *32*, 2832-2841. 10.1093/hmg/ddad110.

27. Ballabio, C., Anderle, M., Gianesello, M., Lago, C., Miele, E., Cardano, M., Aiello, G., Piazza, S., Caron, D., Gianno, F., et al. (2020). Modeling medulloblastoma in vivo and with human cerebellar organoids. Nat Commun *11*, 583. 10.1038/s41467-019-13989-3.

28. van Essen, M.J., Apsley, E.J., Riepsaame, J., Xu, R., Northcott, P.A., Cowley, S.A., Jacob, J., and Becker, E.B.E. (2024). PTCH1-mutant human cerebellar organoids exhibit altered neural development and recapitulate early medulloblastoma tumorigenesis. Dis Model Mech *17*. 10.1242/dmm.050323.

29. van Essen, M.J., Nicheperovich, A., Schuster-Bockler, B., Becker, E.B.E., and Jacob, J. (2025). Sonic hedgehog medulloblastoma cells in co-culture with cerebellar organoids converge towards in vivo malignant cell states. Neurooncol Adv *7*, vdae218. 10.1093/noajnl/vdae218.

30. Yang, J., Ryan, D.J., Wang, W., Tsang, J.C., Lan, G., Masaki, H., Gao, X., Antunes, L., Yu, Y., Zhu, Z., et al. (2017). Establishment of mouse expanded potential stem cells. Nature *550*, 393-397. 10.1038/nature24052.

31. Zhao, X., Liu, Z., Yu, L., Zhang, Y., Baxter, P., Voicu, H., Gurusiddappa, S., Luan, J., Su, J.M., Leung, H.C., and Li, X.N. (2012). Global gene expression profiling confirms the molecular fidelity of primary tumor-based orthotopic xenograft mouse models of medulloblastoma. Neuro Oncol *14*, 574-583. 10.1093/neuonc/nos061.

32. Gossen, M., Freundlieb, S., Bender, G., Müller, G., Hillen, W., and Bujard, H. (1995). Transcriptional activation by tetracyclines in mammalian cells. Science *268*, 1766-1769. 10.1126/science.7792603.

33. Hickmott, J.W., Chen, C.Y., Arenillas, D.J., Korecki, A.J., Lam, S.L., Molday, L.L., Bonaguro, R.J., Zhou, M., Chou, A.Y., Mathelier, A., et al. (2016). PAX6 MiniPromoters drive restricted expression from rAAV in the adult mouse retina. Mol Ther Methods Clin Dev *3*, 16051. 10.1038/mtm.2016.51.

34. Badodi, S., Dubuc, A., Zhang, X., Rosser, G., Da Cunha Jaeger, M., Kameda-Smith, M.M., Morrissy, A.S., Guilhamon, P., Suetterlin, P., Li, X.N., et al. (2017). Convergence of BMI1 and CHD7 on ERK Signaling in Medulloblastoma. Cell Rep *21*, 2772-2784. 10.1016/j.celrep.2017.11.021.

35. Badodi, S., Pomella, N., Zhang, X., Rosser, G., Whittingham, J., Niklison-Chirou, M.V., Lim, Y.M., Brandner, S., Morrison, G., Pollard, S.M., et al. (2021). Inositol treatment inhibits medulloblastoma through suppression of epigenetic-driven metabolic adaptation. Nat Commun *12*, 2148. 10.1038/s41467-021-22379-7.

36. Dobin, A., and Gingeras, T.R. (2015). Mapping RNA-seq Reads with STAR. Curr Protoc Bioinformatics *51*, 11 14 11-11 14 19. 10.1002/0471250953.bi1114s51.

37. Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., Zheng, S., Butler, A., Lee, M.J., Wilk, A.J., Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal single-cell data. Cell *184*, 3573-3587.e3529. https://doi.org/10.1016/j.cell.2021.04.048.

38. Korsunsky, I., Millard, N., Fan, J., Slowikowski, K., Zhang, F., Wei, K., Baglaenko, Y., Brenner, M., Loh, P.R., and Raychaudhuri, S. (2019). Fast, sensitive and accurate integration of single-cell data with Harmony. Nat Methods *16*, 1289-1296. 10.1038/s41592-019-0619-0.

39. Finak, G., McDavid, A., Yajima, M., Deng, J., Gersuk, V., Shalek, A.K., Slichter, C.K., Miller, H.W., McElrath, M.J., Prlic, M., et al. (2015). MAST: a flexible statistical framework for assessing transcriptional changes and characterizing

heterogeneity in single-cell RNA sequencing data. Genome Biology *16*, 278. 10.1186/s13059-015-0844-5.

40. Bhaduri, A., Andrews, M.G., Mancia Leon, W., Jung, D., Shin, D., Allen, D., Jung, D., Schmunk, G., Haeussler, M., Salma, J., et al. (2020). Cell stress in cortical organoids impairs molecular subtype specification. Nature *578*, 142-148. 10.1038/s41586-020-1962-0.

41. Kanton, S., Boyle, M.J., He, Z., Santel, M., Weigert, A., Sanchis-Calleja, F., Guijarro, P., Sidow, L., Fleck, J.S., Han, D., et al. (2019). Organoid single-cell genomic atlas uncovers human-specific features of brain development. Nature *574*, 418-422. 10.1038/s41586-019-1654-9.

42. Aldinger, K.A., Thomson, Z., Phelps, I.G., Haldipur, P., Deng, M., Timms, A.E., Hirano, M., Santpere, G., Roco, C., Rosenberg, A.B., et al. (2021). Spatial and cell type transcriptional landscape of human cerebellar development. Nat Neurosci *24*, 1163-1175. 10.1038/s41593-021-00872-y.

43. Zhong, S., Wang, M., Huang, L., Chen, Y., Ge, Y., Zhang, J., Shi, Y., Dong, H., Zhou, X., Wang, B., et al. (2023). Single-cell epigenomics and spatiotemporal transcriptomics reveal human cerebellar development. Nat Commun *14*, 7613. 10.1038/s41467-023-43568-6.

44. Sepp, M., Leiss, K., Murat, F., Okonechnikov, K., Joshi, P., Leushkin, E., Spanig, L., Mbengue, N., Schneider, C., Schmidt, J., et al. (2024). Cellular development and evolution of the mammalian cerebellum. Nature *625*, 788-796. 10.1038/s41586-023-06884-x.

45. Hovestadt, V., Smith, K.S., Bihannic, L., Filbin, M.G., Shaw, M.L., Baumgartner, A., DeWitt, J.C., Groves, A., Mayr, L., Weisman, H.R., et al. (2019). Resolving medulloblastoma cellular architecture by single-cell genomics. Nature *572*, 74-79. 10.1038/s41586-019-1434-6.

46. Venteicher, A.S., Tirosh, I., Hebert, C., Yizhak, K., Neftel, C., Filbin, M.G., Hovestadt, V., Escalante, L.E., Shaw, M.L., Rodman, C., et al. (2017). Decoupling genetics, lineages, and microenvironment in IDH-mutant gliomas by single-cell RNA-seq. Science *355*, eaai8478. doi:10.1126/science.aai8478.

47. Vladoiu, M.C., El-Hamamy, I., Donovan, L.K., Farooq, H., Holgado, B.L., Sundaravadanam, Y., Ramaswamy, V., Hendrikse, L.D., Kumar, S., Mack, S.C., et al. (2019). Childhood cerebellar tumours mirror conserved fetal transcriptional programs. Nature *572*, 67-73. 10.1038/s41586-019-1158-7.

48. Liberzon, A., Birger, C., Thorvaldsdottir, H., Ghandi, M., Mesirov, J.P., and Tamayo, P. (2015). The Molecular Signatures Database (MSigDB) hallmark gene set collection. Cell Syst *1*, 417-425. 10.1016/j.cels.2015.12.004.

49. Korotkevich, G., Sukhov, V., Budin, N., Shpak, B., Artyomov, M.N., and Sergushichev, A. (2016). Fast gene set enrichment analysis. bioRxiv. 10.1101/060012.

50. Hanzelmann, S., Castelo, R., and Guinney, J. (2013). GSVA: gene set variation analysis for microarray and RNA-seq data. BMC Bioinformatics *14*, 7. 10.1186/1471-2105-14-7.

51. La Manno, G., Soldatov, R., Zeisel, A., Braun, E., Hochgerner, H., Petukhov, V., Lidschreiber, K., Kastriti, M.E., Lönnerberg, P., Furlan, A., et al. (2018). RNA velocity of single cells. Nature *560*, 494-498. 10.1038/s41586-018-0414-6.

52. Bergen, V., Lange, M., Peidli, S., Wolf, F.A., and Theis, F.J. (2020). Generalizing RNA velocity to transient cell states through dynamical modeling. Nature Biotechnology *38*, 1408-1414. 10.1038/s41587-020-0591-3.

53. Cao, J., Spielmann, M., Qiu, X., Huang, X., Ibrahim, D.M., Hill, A.J., Zhang, F., Mundlos, S., Christiansen, L., Steemers, F.J., et al. (2019). The single-cell

transcriptional landscape of mammalian organogenesis. Nature *566*, 496-502. 10.1038/s41586-019-0969-x.

54. Linderman, G.C., Zhao, J., Roulis, M., Bielecki, P., Flavell, R.A., Nadler, B., and Kluger, Y. (2022). Zero-preserving imputation of single-cell RNA-seq data. Nat Commun *13*, 192. 10.1038/s41467-021-27729-z.

55. Efremova, M., Vento-Tormo, M., Teichmann, S.A., and Vento-Tormo, R. (2020). CellPhoneDB: inferring cell-cell communication from combined expression of multi-subunit ligand-receptor complexes. Nat Protoc *15*, 1484-1506. 10.1038/s41596-020-0292-x.

56. Pidsley, R., Wong, C.C.Y., Volta, M., Lunnon, K., Mill, J., and Schalkwyk, L.C. (2013). A data-driven approach to preprocessing Illumina 450K methylation array data. BMC Genomics *14*. https://doi.org/10.1186/1471-2164-14-293.

57. Steg, L.C., Shireby, G.L., Imm, J., Davies, J.P., Franklin, A., Flynn, R., Namboori, S.C., Bhinge, A., Jeffries, A.R., Burrage, J., et al. (2021). Novel epigenetic clock for fetal brain development predicts prenatal age for cellular stem cell models and derived neurons. Mol Brain *14*, 98. 10.1186/s13041-021-00810-w.

58. Peters, T.J., Buckley, M.J., Statham, A.L., Pidsley, R., Samaras, K., Lord, R.V., Clark, S.J., and Molloy, P.L. (2015). De novo identification of differentially methylated regions in the human genome. Epigenetics & Chromatin *8*. https://doi.org/10.1186/1756-8935-8-6.

59. Hovestadt, V., Remke, M., Kool, M., Pietsch, T., Northcott, P.A., Fischer, R., Cavalli, F.M., Ramaswamy, V., Zapatka, M., Reifenberger, G., et al. (2013). Robust molecular subgrouping and copy-number profiling of medulloblastoma from small amounts of archival tumour material using high-density DNA methylation arrays. Acta Neuropathol *125*, 913-916. 10.1007/s00401-013-1126-5.

60. Kolberg, L., Raudvere, U., Kuzmin, I., Adler, P., Vilo, J., and Peterson, H. (2023). g:Profiler-interoperable web service for functional enrichment analysis and gene identifier mapping (2023 update). Nucleic Acids Res *51*, W207-W212. 10.1093/nar/gkad347.

61. Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B., and Ideker, T. (2003). Cytoscape: a software environment for integrated models of biomolecular interaction networks. Genome Res *13*, 2498-2504. 10.1101/gr.1239303.

62. Frantz GD, W.J., Levin ME, McConnell SK (1994). Otx1 and Otx2 define layers and regions in developing cerebral cortex and cerebellum. J Neurosci *14*, 5725-5740. 10.1523/JNEUROSCI.14-10-05725.1994.

63. Badodi, S., and Marino, S. (2022). Epigenetic mechanisms in paediatric brain tumours: regulators lose control. Biochem Soc Trans *50*, 167-185. https://doi.org/10.1042/BST20201227.

64. Zagozewski, J., Shahriary, G.M., Morrison, L.C., Saulnier, O., Stromecki, M., Fresnoza, A., Palidwor, G., Porter, C.J., Forget, A., Ayrault, O., et al. (2020). An OTX2-PAX3 signaling axis regulates Group 3 medulloblastoma cell fate. Nat Commun *11*, 3627. 10.1038/s41467-020-17357-4.

65. Jones, D.T., Jager, N., Kool, M., Zichner, T., Hutter, B., Sultan, M., Cho, Y.J., Pugh, T.J., Hovestadt, V., Stutz, A.M., et al. (2012). Dissecting the genomic complexity underlying medulloblastoma. Nature *488*, 100-105. 10.1038/nature11284.

66. Kahn, S.A., Wang, X., Nitta, R.T., Gholamin, S., Theruvath, J., Hutter, G., Azad, T.D., Wadi, L., Bolin, S., Ramaswamy, V., et al. (2018). Notch1 regulates the

initiation of metastasis and self-renewal of Group 3 medulloblastoma. Nat Commun 9, 4121. 10.1038/s41467-018-06564-9.

67. Rattenberry, W., McDonough, C.H., Burger, P.C., and Cohen, K.J. (2011). Medulloblastoma with myogenic differentiation: long-term survival in a patient treated with aggressive combination therapy and autologous stem cell transplantation. J Neurooncol 105, 659-662. 10.1007/s11060-011-0626-1.

68. Crawford, J.R., and Levy, M.L. (2015). Medulloblastoma with myogenic differentiation: a rare medulloblastoma variant in a young child. BMJ Case Rep 2015. 10.1136/bcr-2015-212322.

69. van Bree, N.F.H.N., and Wilhelm, M. (2022). The Tumor Microenvironment of Medulloblastoma: An Intricate Multicellular Network with Therapeutic Potential. Cancers 14. 10.3390/cancers14205009.

70. Fan, Q., Wang, H., Gu, T., Liu, H., Deng, P., Li, B., Yang, H., Mao, Y., and Shao, Z. (2024). Modeling the precise interaction of glioblastoma with human brain region-specific organoids. iScience 27, 109111. 10.1016/j.isci.2024.109111.

71. Prior, V.G., Maksour, S., Miellet, S., Hulme, A.J., Chen, Y., Mirzaei, M., Wu, Y., Dottori, M., and O'Neill, G.M. (2024). Parsing the effect of co-culture with brain organoids on Diffuse Intrinsic Pontine Glioma (DIPG) using quantitative proteomics. Int J Biochem Cell Biol 174, 106617. 10.1016/j.biocel.2024.106617.

72. Badodi, S., Pomella, N., Lim, Y.M., Brandner, S., Morrison, G., Pollard, S.M., Zhang, X., Zabet, N.R., and Marino, S. (2022). Combination of BMI1 and MAPK/ERK inhibitors is effective in medulloblastoma. Neuro Oncol. 10.1093/neuonc/noac052.

73. Northcott, P.A., Shih, D.J., Peacock, J., Garzia, L., Morrissy, A.S., Zichner, T., Stutz, A.M., Korshunov, A., Reimand, J., Schumacher, S.E., et al. (2012). Subgroup-specific structural variation across 1,000 medulloblastoma genomes. Nature 488, 49-56. 10.1038/nature11327.

74. Kawauchi, D., Ogg, R.J., Liu, L., Shih, D.J.H., Finkelstein, D., Murphy, B.L., Rehg, J.E., Korshunov, A., Calabrese, C., Zindy, F., et al. (2017). Novel MYC-driven medulloblastoma models from multiple embryonic cerebellar cells. Oncogene 36, 5231-5242. 10.1038/onc.2017.110.

75. Pei, Y., Moore, C.E., Wang, J., Tewari, A.K., Eroshkin, A., Cho, Y.J., Witt, H., Korshunov, A., Read, T.A., Sun, J.L., et al. (2012). An animal model of MYC-driven medulloblastoma. Cancer Cell 21, 155-167. 10.1016/j.ccr.2011.12.021.

76. Williamson, D., Schwalbe, E.C., Hicks, D., Aldinger, K.A., Lindsey, J.C., Crosier, S., Richardson, S., Goddard, J., Hill, R.M., Castle, J., et al. (2022). Medulloblastoma group 3 and 4 tumors comprise a clinically and biologically significant expression continuum reflecting human cerebellar development. Cell Rep 40, 111162. 10.1016/j.celrep.2022.111162.

77. Pine, A.R., Cirigliano, S.M., Nicholson, J.G., Hu, Y., Linkous, A., Miyaguchi, K., Edwards, L., Singhania, R., Schwartz, T.H., Ramakrishna, R., et al. (2020). Tumor Microenvironment Is Critical for the Maintenance of Cellular States Found in Primary Glioblastomas. Cancer Discov 10, 964-979. 10.1158/2159-8290.CD-20-0057.

78. Yang, J., Kim, S.K., Wang, K.C., Kim, K.H., Park, S.H., and Phi, J.H. (2021). A boy with a congenital cerebellar mass. Childs Nerv Syst 37, 983-987. 10.1007/s00381-020-04767-4.

79. Moradi, B., Hamidian, Y., Soltanghoraee, H., and Katouli, F.S. (2023). Prenatally detected congenital medulloblastoma. J Clin Ultrasound 51, 432-435. 10.1002/jcu.23278.

80. Gupta, K., Jogunoori, S., Satapathy, A., Salunke, P., Kumar, N., Radotra, B.D., and Vasishta, R.K. (2018). Medulloblastoma with myogenic and/or melanotic differentiation does not align immunohistochemically with the genetically defined molecular subgroups. Hum Pathol *75*, 26-33. 10.1016/j.humpath.2018.01.014.

81. Nicholson, J.G., Cirigliano, S., Singhania, R., Haywood, C., Shahidi Dadras, M., Yoshimura, M., Vanderbilt, D., Liechty, B., and Fine, H.A. (2024). Chronic hypoxia remodels the tumor microenvironment to support glioma stem cell growth. Acta Neuropathol Commun *12*, 46. 10.1186/s40478-024-01755-6.

82. Qu, F., Brough, S.C., Michno, W., Madubata, C.J., Hartmann, G.G., Puno, A., Drainas, A.P., Bhattacharya, D., Tomasich, E., Lee, M.C., et al. (2023). Crosstalk between small-cell lung cancer cells and astrocytes mimics brain development to promote brain metastasis. Nat Cell Biol *25*, 1506-1519. 10.1038/s41556-023-01241-6.

83. Northcott, P.A., Korshunov, A., Witt, H., Hielscher, T., Eberhart, C.G., Mack, S., Bouffet, E., Clifford, S.C., Hawkins, C.E., French, P., et al. (2011). Medulloblastoma comprises four distinct molecular variants. J Clin Oncol *29*, 1408-1414. 10.1200/JCO.2009.27.4324.

84. Hayden Gephart, M.G., Su, Y.S., Bandara, S., Tsai, F.C., Hong, J., Conley, N., Rayburn, H., Milenkovic, L., Meyer, T., and Scott, M.P. (2013). Neuropilin-2 contributes to tumorigenicity in a mouse model of Hedgehog pathway medulloblastoma. J Neurooncol *115*, 161-168. 10.1007/s11060-013-1216-1.

85. Trombetta-Lima, M., Rosa-Fernandes, L., Angeli, C.B., Moretti, I.F., Franco, Y.M., Mousessian, A.S., Wakamatsu, A., Lerario, A.M., Oba-Shinjo, S.M., Pasqualucci, C.A., et al. (2021). Extracellular Matrix Proteome Remodeling in Human Glioblastoma and Medulloblastoma. J Proteome Res *20*, 4693-4707. 10.1021/acs.jproteome.1c00251.

86. Vestal, K.A., Kattamuri, C., Koyiloth, M., Ongaro, L., Howard, J.A., Deaton, A.M., Ticau, S., Dubey, A., Bernard, D.J., and Thompson, T.B. (2024). Activin E is a transforming growth factor beta ligand that signals specifically through activin receptor-like kinase 7. Biochem J *481*, 547-564. 10.1042/BCJ20230404.

87. Liu, H., Sun, Y., O'Brien, J.A., Franco-Barraza, J., Qi, X., Yuan, H., Jin, W., Zhang, J., Gu, C., Zhao, Z., et al. (2020). Necroptotic astrocytes contribute to maintaining stemness of disseminated medulloblastoma through CCL2 secretion. Neuro Oncol *22*, 625-638. 10.1093/neuonc/noz214.

88. Liang, K.H., Chang, C.C., Wu, K.S., Yu, A.L., Sung, S.Y., Lee, Y.Y., Liang, M.L., Chen, H.H., Fen, J.J., Chao, M.E., et al. (2021). Notch signaling and natural killer cell infiltration in tumor tissues underlie medulloblastoma prognosis. Sci Rep *11*, 23282. 10.1038/s41598-021-02651-y.

1641 **Supplementary Figures and Legends**

1642

1643

1644

Supplementary Figure 1. CbO displays sustained growth and maturation.

A. Schematic of protocol used to generate and differentiate CbO, (see Methods,
Y: Y-27632 Dihydrochloride and SB: SB431542) and representative brightfield
pictures of CbO_61 upon 7, 14, 21, 28 and 35 days of differentiation. Scale
bar = 1000 µm.

B. Diameter analysis of CbO obtained from two independent EPSC lines
(CbO_19 and CbO_61) across 35 days of maturation (left) and analysis of
relative growth comparing Day 14 to Day 7 or Day 35. Asterisks indicate p-
values between given time point and Day 7, two-way ANOVA, ****$p<0.0001$,
ns: not significant. n = 10 independent CbO average diameters per timepoint.

C. qPCR analysis showing quantification of *NANOG* expression in CbO_19 and
CbO_61 across 35 days of maturation.

D. qPCR analysis showing quantification of *NESTIN, SOX1, EN2* and *OTX2*
expression in CbO_19 and CbO_61 across 35 days of maturation.

E. qPCR analysis showing quantification of *KIRREL2* and *SKOR2* expression in
CbO_19 and CbO_61 across 35 days of maturation.

F. qPCR analysis showing quantification of *PAX6, ATOH1, BARHL1* and
EOMES expression in CbO_19 and CbO_61 across 35 days of maturation.

Relative RNA level was calculated as $2^{(-dCT)}$ values normalised to EPSC
baseline (Day 0). All graphs report mean ± SEM, Welch's unpaired t-test,
*$p<0.05$, **$p<0.01$, ***$p<0.001$. non-significant comparisons are not shown.
Data points show average values for n = 3 independent batches of CbO per
timepoint.

G. UMAP plot of scRNA-Seq data from all CbO samples submitted at Day 7
(CbO_19 and CbO_61), 35 (CbO_19 and CbO_61)and 49 (CbO_61) of
differentiation coloured by timepoint.

H. UMAP plot of scRNA-Seq data from CbO at Day 7 of differentiation coloured
by annotated Louvain clusters.

I. Dot plot showing key marker genes used to identify major cell types in CbO at
Day 7 of differentiation.

J. UMAP plot of scRNA-Seq data from CbO samples at Day 35/49 of
differentiation, identifying 17 clusters and sub clustering of cluster 7 into 7_0
and 7_1. Low quality hypoxic clusters 13 and 14 not shown.

K. UMAP plot of scRNA-Seq data shown in K, split by Day 35 (upper) and Day
49 (lower) timepoints.

L. Heatmap showing correlation of CbO D35/49 clusters identified in K to
clusters annotated by cell type of four independent scRNA-seq datasets of the
human developing foetal cerebellum[1-4].

1684

1685

1686

Supplementary Figure 2. Staging development of CbO comparatively to human
cerebellum reveals conserved epigenetically regulated genes and pathways.

A. H&E staining of FFPE samples from human foetal cerebellum. Non-cerebellar
tissue excluded from the microdissection is circled. Developmental stage in
post-conception weeks (PCW) and scale bars are shown.

B. Heatmap showing k-means clustering (n=5) of significant variable methylated
genes across human foetal cerebellar development from Carnegie stage (cs)
22 to 37 post-conception weeks (pcw). Z-scores for average beta values of
probes contained within each gene are plotted.

C. Bubble plot showing top 100 significant (adjusted p-value < 0.05) GO
Biological Processes and Molecular Functions of genes from 5 clusters shown
in B taken for pathway analysis in g:Profiler. Bubbles are coloured based on
each clusters and size is proportional to number of genes in specific GO term.

D. Venn diagram showing integrative analysis of significant GOs in most variable
methylated (Figure 2D) and concordant (Figure 2E) genes (left panel).
Bubbles plot showing common pathways between genes
upregulated/hypomethylated and in clusters 3 and 4 (middle panel) or genes
downregulated/hypermethylated and in clusters 1 and 2 (left panel). Bubbles
are coloured based on FDR and size is proportional to number of genes of
specific GO term.

1707

1708

1709

1710

Supplementary Figure 3. CbO clusters harbouring MB cell-of-origin are
enriched for distinct G3 and G4 MB associated gene signatures.

A. Heatmap showing correlation of methylation status of top 1% age-dependent
probes between human developing cerebellum samples, demonstrating a
linear correlation through development.

B. Correlation plot between days of maturation of CbO_19 and CbO_61
(chronological age) and the modelled epigenetic ages from the foetal brain
clock[5].

C. Uniform Manifold Approximation and projection (UMAP) of Day 35/49 CbO
showing alra imputed expression score for *EOMES.*

D. Day 35/49 CbO UMAP with RNA velocity streams shown, calculated in
velocyto and scVelo.

E. Pseudotime analysis of Day 35/49 CbO UMAP calculated in Monocle3, with
cycling progenitors cluster (see Figure 1G) selected as the root of
pseudotime.

F. Plots of z-scored gene expression through pseudotime within G3 and G4
trajectories defined in Figure 3K.

1728

1729

1730

1731

Supplementary Figure 4. Cerebellar organoid editing strategy displays
specificity to medulloblastoma lineages-of-origin and similar
methylation profiles

A. qPCR data of naïve expanded potential stem cells (EPSC19) and EPSC19
infected with *pPAX6;MYC-GFP* or *pEOMES;MYC-GFP* editing constructs
outlined in Figure 4A for pluripotency (*NANOG)* and differentiation (*GATA6)*
markers as well as c-*MYC.* Relative RNA level signifies dCT values
expressed relative to naïve EPSC19 (2^-ddCT). ns = not significant, one-way
ANOVA. n = 3 technical replicates.

B. Representative images of CbO19 at Day 36, 46 and 63 of culture infected with
different combinations of *pPAX6, pEOMES* or *MYC-GFP* editing constructs
(refer to Figure 4A), with or without the addition of doxycycline (DOX) to
organoid media from Day 35 of culture.

C. UMAP plot of scRNA-Seq data from D63 *pEOMES;MYC-GFP, pPAX6;MYC-*
GFP (*pEOMES*-MYC^{OE}, *pPAX6*-MYC^{OE}) and control (CTRL) CbOs coloured
by Louvain clusters (left). Barplot showing the relative proportion of cells
clusters in each sample (right).

D. Barplot showing the relative proportion of cell types in each cluster in **C**
(labels transferred from D35/49 control CbOs (Figure 1G).

E. UMAP plot shown in C, coloured by cell cycle phase.

F. UMAP plot shown in C, showing Group 3 MB signature score[6].

G. Gene set enrichment plot of the HALLMARKs MYC TARGETS V1 gene set in
the MYC^{OE} cluster markers (Figure 4C), NES=3.63, padj=2.24E-22.

H. Gene set enrichment plot showing depletion of gene sets associated with
neural differentiation in MYC^{OE} cells relative to D35/49 G3/4 lineage-of-origin
cells.

I. Clustered heatmap of medulloblastoma (MB) samples taken from Cavalli et
al.[7] and edited CbO samples based on their beta values of probes shown to
delineate MB subgroups[8].

J. Clustered heatmap of edited CbO sample correlation to group 3 and group 4
MBs from the Cavalli et al. dataset[7] on the beta values of differentially
methylated probes between *MYC* overexpressing (*pEOMES*-MYC^{OE}, *pPAX6*-
MYC^{OE}) and control medulloblastoma lineage-of-origin samples
(*pEOMES*.ctrl, *pPAX6*.ctrl).

1766

1767

1768

Supplementary Figure 5. Medulloblastoma (MB) cells display the morphology
of human MB upon co-culture with cerebellar organoids (CbO) and retain their
genetic features.

A. Representative immunofluorescence staining of GFP⁺ ICb1299 G3/4 MB cells
upon fourteen days of co-culture with CbO_61, scale bar = 100 µm.

B. Representative immunofluorescence staining of GFP⁺ CHLA-01R-Med
recurrent G3/4 MB cells upon fourteen days of co-culture with CbO_61, scale
bar = 100 µm.

C. Representative immunofluorescence staining of GFP⁺ Med211FH cells upon
fourteen days of co-culture with CbO_61, scale bar = 50 µm.

D. H&E staining of CbO-Med211FH co-cultures upon 14 days of co-culture with
MB cells, scale bars = 50 µm.

E. H&E staining of CbO-CHLA-01-Med co-cultures upon 14 days of co-culture
with MB cells, scale bars = 100 µm.

F. H&E staining of CbO-Med211FH co-cultures upon 14 days of co-culture with
MB cells, scale bars = 100 µm.

G. UMAP plot of scRNA-Seq data from D49 control CbO_61, CbO-Med211-FH,
CbO-CHLA-01-Med and control CbO-NSC73.5 co-cultures, coloured by
sample.

H. Hierarchical clustering of inferred copy number profiles separates co-cultured
CHLA-01-Med (suede) and CbO-Med211-FH (yellow) from CbO (cyan) cells.

I. Dot plot showing key marker genes that distinguish co-culture NSC73_5,
CHLA-01-Med and CbO-Med211-FH cell populations from CbO.

J. UMAP plot of scRNA-Seq data from MB cells only from D49 CbO-Med211-FH
and CbO-CHLA-01-Med CbO-MB co-cultures, coloured by label transfer
scores for Group3 (top), Group 4 (bottom).

K. Heatmap showing correlation of co-cultured Med211-FH and CHLA-01-Med
scRNA-Seq clusters to reference cell type from four independent scRNA-seq
datasets of the human developing foetal cerebellum[1-4].

1800

1801

1802

Supplementary Figure 6. Co-culture with medulloblastoma cells modulates
transcriptional profiles of specific cerebellar organoid lineages.

A. UMAP plot of scRNA-Seq data from D49 control CbO_61, CbO-Med211-FH,
CbO-CHLA-01-Med and control CbO-NSC73_5 co-cultures, coloured by
Louvain clusters.

B. Barplot showing proportion of each cluster made up of cells from CbO_61
control, CbO-Med211-FH, CbO-CHLA-01-Med or CbO-NSC73_5 co-culture
samples (left). Barplot showing proportion of each cluster made up of different
cell types from CbO_61 control, CbO-Med211-FH, CbO-CHLA-01-Med or
CbO-NSC73_5 co-culture samples (right). For CbO clusters cell labels were
computationally transferred from D35/49 controls (Fig 1G).

C. Barplot showing proportion of samples in A, B made up by the joint cell type
labels.

D. Uniform manifold approximation and projections (UMAPs) plot of scRNA-Seq
data from D49 control CbO, CbO-Med211-FH, CbO-CHLA-01-Med and
control CbO-NSC73_5 co-cultures. Cells are coloured by their assigned cell
type and UMAPs are split according to experimental model.

E. Immunohistochemistry images of cerebellar organoids co-cultured with
NSC73.5 cells for 14 days. Upper panel: GFP (green), cCASP3 (red) and
TubIII (white); Lower panel: GFP (green), cCASP3 (red) and MAP2 (white).
Scale bars = 10 μm.

F. Heatmap showing the total number of predicted significant receptor-ligand
interactions (regardless of directionality) between co-cultured CbO-Med211-
FH, CbO-CHLA-01-Med and control CbO-NSC73_5 cells and CbO_61 cell
populations.

G. Volcano plots showing significant differentially expressed genes (DEGs) upon
MB cells co-culture in indicated CbO_61 cell populations. Red and blue dots
represent significantly upregulated and downregulated genes respectively.

H. Bubble plot showing top 100 significant (adjusted p-value < 0.05) GO
Biological Processes and Molecular Functions of up- (red) and down- (blue)
regulated genes from 6 volcano plots shown in G taken for pathway analysis
in g:Profiler. Bubbles are coloured based on each cell population and size is
proportional to number of genes in specific GO term.

I. Schematic of CbO-CHLA-01-Med treatment with tables reporting the
concentrations of PTC209 (BMI1i), PD329501 (MAPKi) and IP6 used.

J. Viability assays of GFP$^+$ MB cells (green) and GFP$^-$ CbO cells (pink) in CbO-
MB upon 4 days of treatment with increasing concentrations of BMI1 and
MAPK inhibitors (left) or IP6 (right) as shown in Figure 5H and 5I respectively.
All graphs report mean ± SEM, number of independent CbO analyzed is
reported in each graph, one-way ANOVA *$p<0.05$, ***$p<0.001$,
****$p<0.0001$, ns = not significant.

1846

References

1. Zhong, S., Wang, M., Huang, L., Chen, Y., Ge, Y., Zhang, J., Shi, Y., Dong, H., Zhou, X., Wang, B., et al. (2023). Single-cell epigenomics and spatiotemporal transcriptomics reveal human cerebellar development. Nat Commun *14*, 7613. 10.1038/s41467-023-43568-6.

2. Aldinger, K.A., Thomson, Z., Phelps, I.G., Haldipur, P., Deng, M., Timms, A.E., Hirano, M., Santpere, G., Roco, C., Rosenberg, A.B., et al. (2021). Spatial and cell type transcriptional landscape of human cerebellar development. Nat Neurosci *24*, 1163-1175. 10.1038/s41593-021-00872-y.

3. Luo, Z., Xia, M., Shi, W., Zhao, C., Wang, J., Xin, D., Dong, X., Xiong, Y., Zhang, F., Berry, K., et al. (2022). Human fetal cerebellar cell atlas informs medulloblastoma origin and oncogenesis. Nature *612*, 787-794. 10.1038/s41586-022-05487-2.

4. Sepp, M., Leiss, K., Murat, F., Okonechnikov, K., Joshi, P., Leushkin, E., Spanig, L., Mbengue, N., Schneider, C., Schmidt, J., et al. (2024). Cellular development and evolution of the mammalian cerebellum. Nature *625*, 788-796. 10.1038/s41586-023-06884-x.

5. Steg, L.C., Shireby, G.L., Imm, J., Davies, J.P., Franklin, A., Flynn, R., Namboori, S.C., Bhinge, A., Jeffries, A.R., Burrage, J., et al. (2021). Novel epigenetic clock for fetal brain development predicts prenatal age for cellular stem cell models and derived neurons. Mol Brain *14*, 98. 10.1186/s13041-021-00810-w.

6. Vladoiu, M.C., El-Hamamy, I., Donovan, L.K., Farooq, H., Holgado, B.L., Sundaravadanam, Y., Ramaswamy, V., Hendrikse, L.D., Kumar, S., Mack, S.C., et al. (2019). Childhood cerebellar tumours mirror conserved fetal transcriptional programs. Nature *572*, 67-73. 10.1038/s41586-019-1158-7.

7. Cavalli, F.M.G., Remke, M., Rampasek, L., Peacock, J., Shih, D.J.H., Luu, B., Garzia, L., Torchia, J., Nor, C., Morrissy, A.S., et al. (2017). Intertumoral Heterogeneity within Medulloblastoma Subgroups. Cancer Cell *31*, 737-754 e736. 10.1016/j.ccell.2017.05.005.

8. Hovestadt, V., Remke, M., Kool, M., Pietsch, T., Northcott, P.A., Fischer, R., Cavalli, F.M., Ramaswamy, V., Zapatka, M., Reifenberger, G., et al. (2013). Robust molecular subgrouping and copy-number profiling of medulloblastoma from small amounts of archival tumour material using high-density DNA methylation arrays. Acta Neuropathol *125*, 913-916. 10.1007/s00401-013-1126-5.

1891