Failure to Resolve Inflammation Contributes to Juvenile-Onset Cardiomyopathy in a

Mouse Model of Duchenne Muscular Dystrophy

James S. Novak[1,2†], Amy Lischin[1,3], Prech Uapinyoying[1,4], Ravi Hindupur[1], Young Jae Moon[1,5],

Surajit Bhattacharya[1], Sarah Tiufekchiev[1,6], Victoria Barone[1,3], Davi A. G. Mázala[1,7],

Iteoluwakishi H. Gamu[1], Gabriela Walters[1], Karuna Panchapakesan[1], Jyoti K. Jaiswal[1,2†]

[1] Center for Genetic Medicine Research, Children's National Research Institute, Children's

National Research and Innovation Campus, Children's National Hospital, Washington, D.C.,

20012, USA

[2] Departments of Pediatrics and Genomics and Precision Medicine, The George Washington

University School of Medicine and Health Sciences, Washington, D.C., 20037, USA

[3] Columbian College of Arts and Sciences, The George Washington University, Washington,

D.C. 20052, USA.

[4] Neuromuscular and Neurogenetic Disorders of Childhood Section, National Institute of

Neurological Disorders and Stroke, National Institutes of Health, Bethesda, MD, USA.

[5] Department of Biochemistry and Orthopaedic Surgery, Jeonbuk National University Medical

School and Hospital, Jeonju, 54907, Republic of Korea

[6] Integrated Biomedical Sciences, The George Washington University School of Medicine and

Health Sciences, Washington, D.C., 20037, USA

[7] Department of Kinesiology, College of Health Professions, Towson University, Towson, MD,

21252, USA

† Author for correspondence

Corresponding Author Address: James S. Novak (jnovak@childrensnational.org) and Jyoti K.

Jaiswal (jkjaiswal@childrensnational.org); Center for Genetic Medicine Research, Children's

National Research Institute, Children's National Hospital, Washington, D.C., 20012, USA

Abstract

The absence of dystrophin protein causes cardiac dysfunction in boys with Duchenne Muscular

Dystrophy (DMD). However, the common mouse model of DMD (B10-*mdx*) does not manifest

cardiac deficits until late adulthood limiting our understanding of the mechanism and therapeutic

approaches to target the pediatric-onset cardiac pathology in DMD. We show the mdx mouse

model on the DBA/2J genetic background (D2-*mdx*) displays juvenile-onset cardiomyopathy.

Molecular and histological analysis revealed heightened leukocyte chemotactic signaling and

failure to resolve inflammation, leading to chronic inflammation and extracellular matrix (ECM)

fibrosis, causing cardiac pathology in juvenile D2-*mdx* mice. We show that pharmacologically

activating the N-formyl peptide receptor 2 (FPR2) - a receptor that physiologically resolves

acute inflammation, mitigated chronic cardiac inflammation and fibrosis, and prevented juvenile

onset cardiomyopathy in the D2-mdx mice. These studies offer insights into pediatric onset of

cardiac damage in DMD, a new therapeutic target, and identify a drug-based potential therapy.

Introduction

Duchenne Muscular Dystrophy (DMD) is a severe and progressive muscle disease caused by the absence of dystrophin protein[1-4]. Dystrophin plays a crucial role in maintaining the integrity of the sarcolemmal membrane by facilitating the assembly and function of the dystrophin-associated protein complex, thus its absence renders skeletal muscle cells more susceptible to mechanical damage, and increased muscle degeneration[5-7]. Dystrophin deficiency in cardiomyocytes also increases their vulnerability to sarcolemmal damage, cell death, chronic inflammation and cardiac fibrosis. These pathologies also manifest in patients and lead to the thinning of the left ventricle (LV) wall, causing their progressive dilation and dilated cardiomyopathy that results in heart failure[4,8].

Patients with DMD experience symptoms early in life, with cardiac deficits being a major contributor to premature mortality not only in DMD, but also in DMD carriers and in Becker Muscular Dystrophy (BMD) patients[9-11]. While the common B10-*mdx* model exhibits skeletal myopathy at an early age, the cardiac deficit is not evident until late adulthood[12,13]. The advent of the D2-*mdx* model identified greater disease severity and fibrosis as compared to B10-*mdx* even in younger mice[14-17]. Excess skeletal muscle fibrosis in the D2-*mdx* model, results from an increase in transforming growth factor beta (TGFβ) signaling, and cardiac deficit is reported as early as adulthood (16-weeks of age)[17-19].

While therapeutic approaches to address cardiac deficit in patients with dystrophin deficiency are a topic of active investigation, anti-inflammatory glucocorticoids (GCs), angiotensin-converting enzyme inhibitors (ACEi), and angiotensin receptor blockers (ARBs), are commonly used for these patients[11,20-22]. Anti-inflammatory therapy by GC has the longest history of use in DMD patients and has mixed reports of cardiac benefit and side effects in both DMD patients and mouse models[23-27]. An anti-inflammatory protein activated by GCs that mediates the GC efficacy is Annexin A1 (AnxA1)[28,29]. AnxA1 works similarly to the endogenous pro-resolving lipid mediators Lipoxin A4, and Resolvin D1 by helping resolve acute inflammation

by binding the Formyl Peptide Receptors (FPRs)[29,30]. This feature of AnxA1 has led to the

advent of natural and synthetic agonists of FPR2, that unlike GCs, reduces inflammation by

promoting resolution of chronic inflammation instead of suppressing the tissue's inflammatory

response[31].

Use of FPR2-agonists reduces acute cardiac damage in various tissue injury models

including myocardial infarction, where it restricts premature heart failure and restores tissue

function[32-36]. Just as endogenous FPR2 agonists, nanomolar dose of a synthetic agonist BMS-

986235 also resolves chronic inflammation in preclinical models, which has led to its progress to

clinical studies[36] (Clinical Trial NCT03335553). This drug activates macrophage transition to a

pro-resolving (M2-like) state by enhancing phagocytosis and neutrophil apoptosis to regulate

inflammatory cell chemotaxis, all of which help improve mouse survival, reduce scarring, and

preserve tissue degeneration[36,37]. These are desirable features of therapies to target cardiac

inflammation and fibrosis associated with the cardiac pathology observed in DMD patients.

To understand the early onset of cardiac dysfunction in the D2-*mdx* model, we investigated

the factors that distinguish the pediatric initiation of cardiac dysfunction as compared with the

late adult onset in B10-*mdx* model. This revealed onset of cardiac pathology in juvenile (< 6-

weeks old) D2-*mdx* mice, and showed this is associated with excessive immune infiltration,

fibrotic ECM replacement, and degenerative remodeling of cardiac ventricular walls. It identified

increased leukocyte chemotactic signaling and failure to resolve inflammation as major

contributors to the initiation of cardiac pathology. To address this deficit, we test a drug-based

pro-resolution therapy for a preclinical evaluation as a therapeutic approach to mitigate early-

onset cardiac damage in DMD mouse models and patients.

Methods.

Animals and Sex as a biological variable: The study involved use of mice of both sexes with

procedures used followed guidelines for the care and use of laboratory animals as approved by

the Institutional Animal Care and Use Committee (IACUC) of Children's National Research

Institute (CNRI). The C57BL/10ScSn-DMD^{mdx}/J (B10-*mdx*) and D2.B10-DMD^{mdx}/J mouse

models of DMD were utilized for all experiments and both harbor the same nonsense point

mutation in exon 23 of the dystrophin (*Dmd*) gene thereby abolishing dystrophin protein

expression [38,39]. The C57BL/10ScSn/J (B10-WT) and DBA2/J (D2-WT) mouse models were

used as age-matched, model-specific controls. Mice were obtained from the Jackson Laboratory

and were housed at the CNRI Comparative Medicine Unit where they were provided daily

monitoring, food, water and enrichment ad libitum, while being maintained under 12 h light/dark

cycles.

Tissue harvesting and sample collection. Mice were euthanized via CO_2 inhalation and

cervical dislocation at designated ages corresponding to specific stages of disease progression.

Muscles were surgically removed, mounted on cork with tragacanth gum, flash-frozen in liquid

nitrogen-chilled isopentane and stored at –80°C. For all assays, samples were collected from

matched regions of the same muscles by collecting cryosections (Leica CM1950 cryostat) for

RNA analyses or histology and immunostaining assays.

RNA extraction, RNA library preparation, RNA sequencing and bioinformatic analyses.

Total RNA was extracted using TRIzol RNA isolation (Life Technologies) from frozen muscle

samples. RNA was purified using RNeasy mini elute columns (Qiagen) and DNAse treated

using Turbo DNA-free kit (Invitrogen). Purified, DNAse-treated RNA was quantified by

NanoDrop and quality was assessed using Qbit RNA Assays (Thermofisher) and BioAnalyzer

nano chips (Agilent) (RIN>7.8, Average 8.3±0.32). RNAseq library preparation and sequencing

was performed using the TruSeq mRNA stranded kit (Illumina) and the Illumina HiSeq4000 Flow

Cell with an average coverage of 63.05 million read pairs per sample at 2x75 base pair read

length. The quality of the raw fastq reads from sequencer were evaluated using FastQC version

0.11.5 followed by adapter and quality trimming using Trimgalore. STAR 2.5.3a was used to

map the reads to the reference mouse genome (GRCm38-mm10). The mapped reads were

counted using HTSeq (version v0.11.0) with a reference genomic feature file (Gene transfer

format, GTF). Overall analysis summary reports were analyzed using MultiQC v1.6. Differential

gene expression analysis was performed using default parameters with Deseq2 version 1.26 (R

3.6). Visualization was performed by PCA, pheatmap, EnhancedVolcano and ggplot2 R

packages. We set a threshold for $\log_2$ fold change (Log_2FC) change of greater than an absolute

value of 0.6 to select for the genes with significant differential expression. Gene lists were

sorted by Log_2FC (highest to lowest) to obtain the ranked list of differentially expressed genes

and a p_{adj} value cutoff of 0.05 was used to assess statistical significance.

Gene Set Enrichment Analysis (GSEA). DESeq2 pairwise comparison results were filtered for

0.6 $\log_2\text{FC}$ and 0.05 adjusted p_{adj} value and exported as tab delimited rank files (.rnk) files for

upload into the stand-alone desktop version of GSEA (v4.1.0). The GSEA pre-ranked analysis

was used with most default parameters except the following: the Ranked list = pairwise

comparison ".rnk" file, Gene sets database = "c5.bp.v7.4.symbols" (Hallmark gene sets, GO

biological processes, and gene symbols) and the Chip platform = "Mouse_Gene_Symbol_Re

mapping_to_Human_Orth ologs_MSigDB.v7.4.chip".

Gene ontology analysis with Cytoscape and EnrichmentMap. The GSEA pairwise

comparison results were uploaded into Cytoscape (v3.9.0) and analyzed using the

EnrichmentMap pipeline collection plugins (v1.1.0). Comparisons were loaded into

EnrichmentMap and the AutoAnnotate function was used with the MCL Cluster Annotation

algorithm set to 5 words. The results are networks of related GO terms that are grouped

together into a named network based on the most common words in each GO term within.

Autogenerated names of networks were renamed to fix grammar and nodes arranged to

improve legibility. The leading-edge genes for each cluster (node) from the immune and

extracellular matrix networks were exported for further analysis.

Histology and histological analyses. Frozen muscles were removed from -80°C cryostorage

and sectioned at an 8 µm thickness using a Leica CM1950 cryostat chilled to -20°C, where

tissues were then mounted on slides and stained using Hematoxylin and Eosin (H&E), Alizarin

Red, Picosirius red, and Masson's Trichrome according to TREAT-NMD Standard Operating

Procedures (SOPs) [39]. Thresholding parameters were applied uniformly to whole cross-section

tiled images acquired on the Olympus VS120-S5 Virtual Slide Scanning System using CellSens

Version 1.13 and ImageJ FIJI Version 2.1./1.53c. For H&E-stained images, areas of damage

were selected using CellSens and quantified as percent damaged tissue area per total cross-

sectional muscle area. For Alizarin Red stained images, calcified areas were selected and

quantified using CellSens as percent calcified tissue area per total cross-sectional muscle area.

For Masson's Trichrome stained images, areas of fibrosis were calculated using FIJI (Image J)

and reported as percent fibrosis per total cross-sectional muscle area [39].

Immunofluorescence. Frozen muscles were removed from -80°C cryostorage and sectioned

at an 8 µm thickness and mounted on slides for immunostaining procedures. Muscle sections

were stained with anti-F4/80 (1:100, MCA497R, Bio-Rad), anti-COL1A1 (1:100, ab21286,

Abcam), and anti-GAL-3 (1:100, ab76245, Abcam). First, muscle sections were fixed in ice-cold

PFA for 10 min, washed in PBS (0.1% Tween-20), and blocked for 1 h in PBS supplemented

with 10% goat serum (GeneTex), 0.1% Tween-20 (Sigma-Aldrich), and 10 mg/mL BSA (Sigma-

Aldrich). Then sections were incubated with primary antibodies overnight at 4°C and

subsequently probed with Alexa Fluor secondary antibodies, including goat anti-rat (H+L) Alexa

Fluor 647 (1:500, A-21247, Thermo Fisher), goat anti-rabbit (H+L) Alexa Fluor 488 (1:500, A-

11008, Thermo Fisher), and goat anti-rabbit (H+L) Alexa Fluor 568 (1:500, A-11011, Thermo

Fisher). Sections were counterstained with wheat germ agglutinin (WGA) Alexafluor-647 (1:500,

W32466, Thermo Fisher) to delineate cardiomyocytes and ProLong Gold Antifade with DAPI

(P36935, Thermo Fisher) for nuclear staining.

Gene expression analysis. Hearts from juvenile and adult dystrophic mice were used to

perform gene expression analysis. In brief, total RNA was extracted from muscle samples by

standard TRIzol (Life Technologies) isolation. Purified RNA (1000ng) was reverse-transcribed

using Random Hexamers and High-Capacity cDNA Reverse Transcription Kit (Thermo Fisher

Scientific). The mRNAs were then quantified using individual TaqMan assays on an ABI

QuantStudio 7 Real-Time PCR machine (Applied Biosystems) using TaqMan Fast Advanced

Master Mix (Thermo Fisher Scientific). Specific mRNA transcript levels were quantified using

individual TaqMan assays (Thermo Fisher) specific for each mRNA target, including Ccl8

(Mm01297183_m1), Ccl3 (Mm00441259_g1), Ccl2 (Mm00441242_m1), Il1b

(Mm00434228_m1), Anxa1 (Mm00440225_m1), Lgals3 (Mm00802901_m1), Stab2

(Mm00454684_m1), Arg1 (Mm00475988_m1), Il7r (Mm00434295_m1), Adam8

(Mm01163449_g1), Trem2 (Mm04209424_g1), Fpr2 (Mm00484464_s1), Spp1

(Mm00436767_m1), Fn1 (Mm01256744_m1), Col1a1 (Mm00801666_g1), Itgax

(Mm00498701_m1), Mmp12 (Mm00500554_m1), and Timp1 (Mm01341361_m1).

Pro-resolution preclinical drug trial. D2-*mdx* mice (18-19 days-old, n=6, males and females)

were treated daily with pro-resolving drug BMS-986235 (6mg/kg, oral gavage; HY-131180,

MedChemExpress) for 3 weeks. Control D2-*mdx* mice were administered saline (18–19-day-old,

n=6, males and females) for 3 weeks. BMS-986235 was initially resuspended in 10% DMSO

(Sigma) and 90% corn oil (Sigma), and further diluted in cherry syrup (NDC-0395-2662-16,

Humco) for oral gavage. After treatment, hearts were harvested, imaged, flash-frozen in liquid

nitrogen-chilled isopentane and stored at –80°C for molecular and histopathological analyses.

Microscopy. We used Olympus VS120-S5 Virtual Slide Scanning System with UPlanSApo

40×/0.95 objective, Olympus XM10 monochrome camera, and Olympus VS-ASW FL 2.7

imaging software. Analysis was performed using Olympus CellSens 1.13 and ImageJ FIJI

Version 2.1./1.53c software (National Institutes of Health). Brightfield whole tissue imaging was

performed using Labomed Luxeo 6Z Digital HD Stereo Microscope with 10x/22mm objective

and camera system.

Statistics. GraphPad Prism 9.2.0 was used for all statistical analyses of data. Statistical

analysis was performed using non-parametric Mann-Whitney tests. Data normality was

assessed for all statistical comparisons. All p-values less than 0.05 were considered statistically

significant; $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, and $****p < 0.0001$. Data plots reported as scatter

plot with mean ± SD.

Results

D2-mdx model Exhibits Pediatric-Onset Cardiomyopathy

We have previously described the use of D2-*mdx* model of DMD to define mechanisms

contributing to the pediatric onset severe skeletal muscle degeneration[39,40]. Here we observed

extensive pericardial damage of the ventricular wall in juvenile (<6-week-old) D2-*mdx* hearts

(**Fig. 1A**). In contrast, the age-matched milder DMD model, B10-*mdx*, did not demonstrate any

conspicuous histopathology (**Fig. 1B**). This identified the utility of D2-*mdx* as a model for

'pediatric-onset' cardiomyopathy for DMD. The histopathological damage extended to both the

right and the left ventricular walls (**Fig. 1C**). Use of Sirius Red staining revealed extensive

fibrosis, which manifests at lower but notable levels in both endomysium and perimysium of the

ventricular wall (**Fig. 1D**). Whole tissue cross-sectional analysis of H&E and Sirius Red stained

hearts revealed increased cardiac wall damage and cardiomyocyte loss (*p<0.001*) (**Fig. 1C, E,**

F), and a concomitant increase in fibrosis (*p<0.05*) in the juvenile D2-*mdx* hearts compared to

age-matched B10-*mdx* hearts (**Fig. 1D, G, H**). The extent of damage in D2-*mdx* mice varied

between the mice reaching levels of nearly 20% of total cardiac muscle cross-sectional area

(**Fig. 1E, F**). Reminiscent of the skeletal muscle damage, cardiac muscle also exhibited

increased calcification in the damaged areas, and this too was observed in the juvenile D2-*mdx*

but not in B10-*mdx* mice (**Supplemental Fig. 1**). Overall, these histopathological analyses

reveal early onset spontaneous cardiac damage with significant fibro-calcification in the D2-

mdx, offering a model to investigate the mechanisms of pediatric onset cardiac damage and

accompanying endomysial fibrosis observed in DMD.

Dysregulated inflammatory response characterize cardiomyopathy in juvenile D2-*mdx*

To identify the molecular alterations associated with the pediatric- versus adult-onset

cardiac damage caused by dystrophin deficit, we performed a comparative transcriptomic

analysis of hearts from 6-week-old male B10-*mdx* and D2-*mdx* hearts. Use of bulk RNA

sequencing we identified a total of 5,344 differentially expressed genes (DEGs) - Log_2 Fold

change of 0.6 with adjusted p-value of 0.05. Principal component analysis (PCA) identified that

gene expression profiles of D2-*mdx* hearts were distinctly segregated from B10-*mdx* along both

PC1 (72% variance) and PC2 (15% variance) axes (**Fig. 2A**). Intra- and inter sample variability

of DEGs by heatmap analysis of the top 2,719 DEGs revealed consistent trend of altered

expression at disease onset, with 1,586 genes upregulated in D2-*mdx* and 1,133 genes

upregulated in B10-*mdx* (**Fig. 2B**).

To identify the functions of the DEGs and assess how they may contribute to the observed

histopathology and functional deficit in juvenile D2-*mdx* hearts, we performed Gene Set

Enrichment Analysis (GSEA) using our DESeq2 pairwise comparison using the same fold

change and adjusted p-values used above for DEG analysis and performed Gene Ontology

(GO) analysis using Cytoscape's EnrichmentMap pipeline (**Fig. 2C**). Outputs from GSEA

provided a comprehensive GO analysis of all upregulated and downregulated GOBP terms,

their respective normalized enrichment scores (NES), p-values, and enrichment plots identifying

alterations in inflammatory and extracellular matrix GOBPs have strongest positive enrichment

scores (ES) (**Supplemental Table 1**, **Supplemental Fig. 2**). The top 20 upregulated GOBP

terms identified dysregulation of the inflammatory response or extracellular matrix architecture

that included a total of 331 of the most upregulated DEGs common between them

(**Supplemental Table 2**). Specifically, GO analysis implicated regulation of the inflammatory

response, immune cell activation, leukocyte chemotaxis and migration, and cytokine signaling

(**Fig. 2C,** red clusters), as well as extracellular matrix organization and disassembly, osteoblast

differentiation, substrate adhesion and bone remodeling, and overall tissue homeostasis (**Fig.**

2C, blue clusters). This was validated by the quantification of the normalized expression for the

top 20 most upregulated DEGs specific to unique GOBPs such as inflammatory response

(GO:0006954) (**Fig. 2D**) and external encapsulating structure (extracellular matrix) organization

(GO:0045229) (**Fig. 2E**). Analysis of leading-edge genes identified increased expression of pro-

inflammatory chemokines, and components of fibrotic extracellular matrix remodeling as

potential drivers of overt pediatric-onset cardiomyopathy in the D2-*mdx* (**Figure 2,**

Supplementary Fig. 2).

To independently validate the role of aberrant acute inflammatory response of granulocyte

and leukocyte chemotaxis by way of chemokine/cytokine signaling pathways, we examined

these genes by qPCR in an expanded (D2-*mdx* n = 9 and B10-*mdx* n = 7) cohort of juvenile

mdx mice. This validated our findings from RNAseq analysis and confirmed significant

upregulation of macrophage-secreted pro-inflammatory C-C family chemokines, *Ccl3*

(macrophage inflammatory protein-1α) and *Ccl8* (monocyte chemoattractant protein-2), that

regulate neutrophil, monocyte and lymphocyte chemotaxis following acute tissue damage (**Fig.**

3A; *p<0.001*). Similarly, expression of *Il7r* (interleukin-7 receptor), that promotes neutrophil and

monocyte recruitment, was also found upregulated in D2-*mdx* hearts (**Fig. 3A**; *p<0.001*), while,

expression of *Stab2* (stabilin-2), a macrophage-expressed phosphatidylserine (PS) surface

receptor that mediates phagocytosis and extracellular matrix remodeling during inflammation [41],

was also upregulated in D2-*mdx* hearts (**Fig. 3A**; *p<0.001*). Expression of *Adam8* (a disintegrin

and metalloproteinase 8), that promotes release of pro-inflammatory cytokines and cell

adhesion molecules and degradation of extracellular matrix [42], was also highly overexpressed in

D2-*mdx* hearts in accordance with bulk RNAseq results (**Fig. 3A**; *p<0.001*). *Trem2* (triggering

receptor expressed on myeloid cells 2), which drives NF-kB signaling and production of pro-

inflammatory cytokines including IL-6 and TNFα [43], was also upregulated in juvenile D2-*mdx*

hearts (**Fig. 3A**; *p<0.01*).

With the abundant increase in inflammatory cell chemokines, we examined the abundance

of macrophages using the pan-macrophage marker F4/80. This confirmed extensive presence

of macrophages in the damaged regions in D2-*mdx* hearts, which is distinct from what is

observed in the B10-*mdx* hearts (**Fig. 3C, D;** *top panels*). To assess the attributes of infiltrating

macrophages in the D2-*mdx* hearts, we next profiled transcript levels of *Il-1b* (interleukin-1b), a

pro-inflammatory macrophage marker, *Arg1* (arginase-1), a pro-regenerative macrophage

marker, and *Lgals3* (galectin-3), a pathogenic macrophage marker[44,45]. While the inflammatory

and regenerative marker genes were increased by 3-, and 9-folds respectively in D2-*mdx*,

compared to B10-*mdx* hearts, the level of pathogenic marker (*Lgals3*) was elevated by ~45-fold

(*p<0.001*). To determine if the different genetic background may contribute to this dysregulation

we examined these transcripts in age-matched juvenile D2-WT and B10-WT mice. This showed

no influence of genetic background for *Adam8*, *Il1b*, *Lgals3*, and *Arg1* between strains, and

relatively small impact on the expression of *Ccl3* (*p<0.01*), *Ccl8* (*p<0.01*), *Trem2* (*p<0.01*), and

Il7r (*p<0.05*) (**Supplemental Fig. 3A**). As our previous investigations have revealed galectin-3

enriched macrophages as a driver of fibrotic degeneration of D2-*mdx* skeletal muscles[46], we

assessed the tissue localization and abundance galectin-3 protein (GAL-3) in D2-*mdx*. This

identified that GAL-3+ macrophages were highly abundant and nearly exclusively localized

within the damaged regions of D2-*mdx* hearts (**Fig. 3C**), while in accordance with qPCR results,

these pathogenic macrophages were absent in B10-*mdx* hearts (**Fig. 3B**, **D**).

Increased expression of chemokines observed in these tissues explains the excessive

infiltration of macrophages in D2-*mdx*. However, during acute injury, the inflammatory response

is controlled by a pro-resolving response that follows cytokine-mediated activation of

inflammation and involves activation of the G protein coupled receptors including Formyl peptide

receptors (FPRs). FPRs are activated by the endogenous ligands produced in response to

tissue damage, including lipids (Resolvin D1, Lipoxin A4) and protein (Annexin A1; AnxA1) that

bind FPR1/2[47-49] and serve as a master switch at the site of damage that helps resolve the

inflammation. They do so by promoting macrophage skewing from pro- to anti-inflammatory

fates and regulating signaling pathways that help clear immune cell infiltration by activating their

apoptosis and non-phlogistic clearance as well [49-53]. To assess if the excessive inflammatory

responses in D2-*mdx* hearts is due to reduced FPR signaling, we examined the expression of

FPRs (*Fpr1*, *Fpr2*) and its ligand *Anxa1*. This revealed over 4-fold upregulation of FPRs (both

Fpr1 and *Fpr2*) expression in D2-*mdx* relative to B10-*mdx* heart, but no change in the

expression of *Anxa1* between D2-*mdx* and B10-*mdx* (**Fig. 3B**). These results validated the

findings from the previous cohort used for bulk RNAseq analysis (**Fig. 2, Supplementary Table**

1). Together, they suggest reduced activation of FPR signaling hinders resolution of

inflammation and may be a driver of the excessive inflammation seen in juvenile D2-*mdx* hearts,

which in turn is linked to their fibro-degenerative state.

Fibrotic ECM remodeling drives early onset cardiac fibrosis in juvenile D2-*mdx*.

To evaluate the prominent upregulation of ECM remodeling pathways implicated by the

RNAseq analysis, we performed qPCR for multiple extracellular matrix-associated components

and remodeling enzymes identified by the analysis of our bulk RNAseq cohort. This validated

the observed upregulation of *Fn1 (*fibronectin; *p<0.01), Col1a1 (*collagen 1A; *p=0.0712), and*

Itgax (integrin alpha X; *p<0.001)* in D2-*mdx* hearts, relative to B10-*mdx* (**Fig. 4A**)*.* Assessment

of *Spp1* (Osteopontin), a DMD genetic modifier that links inflammation to extracellular matrix

assembly and fibrosis [54-58] and macrophage-expressed matrix remodeling enzyme *Mmp12*

*(*matrix metalloproteinase 12), both showed over 200-fold upregulation in D2-*mdx* hearts,

compared to B10-*mdx* (**Fig. 4A**). While other ECM regulator and structural components

including *Timp1 (*TIMP metallopeptidase inhibitor 1), Itgax (Integrin X), *Fn1* (fibronectin), and

Col1a1 (Collagen A1) showed between 4-40-fold upregulation in D2-*mdx* hearts (**Fig. 4A**).

These validate the findings from the bulk RNAseq cohort and implicate a nexus of inflammatory-

ECM dysregulation in pediatric-onset cardiac pathogenesis in the D2-*mdx*. To monitor the sites

of fibrosis in juvenile D2-*mdx* hearts, we immunostained heart cross-sections for COL1A1 and

co-stained with wheat germ agglutinin (WGA), which showed dense COL1A1 expression in D2-

mdx in the damaged regions along the RV wall and throughout the endomysium (**Fig. 4B, B'**),

while endomysial COL1A1 staining in B10-*mdx* counterparts was minimal in comparison (**Fig.**

4C, C'). Again, to address the influence of genetic background on the above findings, we

assessed expression in D2-WT and B10-WT hearts, which indicated no genotype-related

differences in the expression of *Fn1*, *Col1a1*, *Itgax*, or *Timp1*, and a comparatively modest

increases in the expression of *Spp1* (*p<0.01*) and *Mmp12* (*p<0.01*) in D2-WT hearts, relative to

B10-WT (**Supplemental Fig. 3B**), when compared to differences between mdx strains (**Fig.**

4A). This identified the site and composition of the fibrotic ECM in the D2-*mdx* heart. It also

highlighted the potential of targeting the aberrant pro-inflammatory response to attenuate fibrotic

cardiac degeneration of the D2-*mdx* dystrophic heart.

Activation of pro-resolving FPR signaling prevents cardiac damage in D2-*mdx* hearts.

The above findings link inflammation and cardiac fibrosis in D2-*mdx*. With mixed success of

corticosteroid use in treating this in heart by dampening inflammation, coupled with our findings

of poor activation of pro-resolving FPR2 signaling, we hypothesized that the use of FPR2-

targeting therapy may resolve the chronic inflammation, without blocking the acute inflammation

required for the reparative ability of the injured tissue[59,60] (**Fig. 5A**).

To assess the benefit of a pro-resolving FPR2-agonist therapy for pediatric-onset cardiac

pathology in D2-*mdx*, we tested use of synthetic FPR2 agonist BMS-986235 compared to saline

(n = 6 animals each/cohort). Mice were orally dosed with drug beginning at 3-weeks of age

(coinciding with onset of cardiac pathology) and were maintained on drug (or saline) until 6-

weeks of age when tissues were harvested for further analysis (**Fig. 5B**). Analysis of

histopathology of the cardiac tissue cross-section, revealed clear lack of fibro-calcified damaged

areas along the RV wall in drug-treated hearts, compared to saline controls (**Fig. 5C**).

Histological analyses performed by H&E staining, confirmed this observed therapeutic effect

with some of the drug-treated hearts showing small and discrete areas of damage within the RV

wall, and the rest lacked any signs of damage or inflammation altogether, while the saline

treated mice showed extensive cardiac damage, (**Fig. 5D**). To further assess this impact of pro-

resolving therapy on inflammation and extracellular matrix remodeling in D2-*mdx* hearts, we

immunostained tissue cross-sections for macrophages, which confirmed the reduction in

macrophage infiltration through the heart and within and surrounding any small sites of damage

that existed in our treated cohort (**Fig. 5E**). This was in stark contrast to the heightened

macrophage infiltration both within and surrounding sites of damage in saline control hearts

(**Fig. 5E**). Next, to more directly assess fibrotic gene expression we immunostained these

hearts for COL1A1 and found the drug treatment also significantly reduced the COL1A1

expression throughout the heart, when compared to the expression in the saline controls (**Fig.**

5F).

To assess the effect of acute BMS-986235 treatment on inflammation and extracellular

matrix remodeling pathways, we next performed targeted qPCR for both inflammatory- and

extracellular matrix remodeling-associated transcripts previously shown to be dysregulated in

D2-*mdx* hearts (**Fig. 2-4**). We found acute BMS-986235 treatment of juvenile D2-*mdx* mice

resulted in a significant reduction in the levels of inflammatory transcripts *Trem2*, *Lgals3*, and

Anxa1 (**Fig. 5G**), confirming the efficacy of this pro-resolving therapy to attenuate aberrant

inflammatory signaling via the FPR2-ANXA1 axis. Similarly, quantification of extracellular matrix

remodeling targets, *Spp1*, *Timp1,* and *Col1a1*, showed significant depletion of *Spp1* and *Timp1*

transcripts (**Fig. 5H**), and a trending reduction in *Col1a1* transcript levels (**Fig. 5H**) which aligns

with COL1A1 immunostaining results (**Fig. 5F**). Thus, future chronic studies will be required to

assess the full benefit of chronic pro-resolving therapy to delay the onset and lessen the

severity of fibrosis with disease progression in older D2-*mdx* mice.

Discussion

While there has been a long-standing recognition of early-onset cardiac deficit and its lethal

consequences for boys with DMD, our experimental understanding of the molecular deficits and

preclinical interventions have been based on the use of adult mouse models. This is due to the

late onset of cardiac damage in the *mdx* mouse model. Our studies here introduce D2-*mdx* as a

model that manifests pediatric-onset cardiomyopathy. Histopathological analyses reveal early

onset spontaneous cardiac damage with significant fibro-calcification in the D2-*mdx* model,

offering an opportunity to investigate the mechanisms of pediatric-onset cardiac damage and

accompanying endomysial fibrosis observed in DMD.

Our studies here identify dysregulation of inflammatory and ECM remodeling as two such

pathological pathways. Next, we focused on harnessing underlying molecular pathways that

distinguish the cardiac deficit in D2-*mdx* model as compared to the adult-onset cardiac deficit

B10-*mdx* model. This analysis allowed distinguishing the differences between mild and severe

cardiomyopathy independent of the presence/absence of dystrophin protein. Such differences

are reminiscent of the DMD patients, who manifest varying level and severity of cardiomyopathy

despite lacking dystrophin expression.

Our studies identify excessive inflammatory response that fails to resolve through FPR2-

mediated pro-resolving pathway prevents restoring the injured heart tissues to their uninflamed

state. While the infiltrating leukocytes are needed in damaged heart to clear the dead cells,

FPR2 and other mediators that repress inflammation are released leading to predominance of

anti-inflammatory cells - a response associated with activation of cardiac repair. We find that

this latter repressive response is poor, and that the excessive inflammatory signaling proceeds

via Spp1 and TGFβ pathway to activate downstream cardiac fibrosis and other degenerative

response. Activation of these profibrotic degenerative responses have long been recognized as

a feature of the D2-*mdx* model[61] and we previously identified that this pathway contributes to

excessive skeletal muscle pathology in the juvenile D2-*mdx* mice by the suppression of skeletal

muscle regeneration[39,40]. Unlike skeletal muscle, cardiac muscle does not undergo regeneration

and our comparative analysis of D2-*mdx* and B10-*mdx* identify cardiomyocyte degeneration due

to chronic inflammation and loss by way of fibro-calcified ECM. We observe the ventricular

pericardium as the region most affected by this damage, but this can progress to the atria as

well (**Fig. 1**). This variability in the affected region is in addition to the variability we observe in

the severity of cardiac damage between individual mice. This hints at a level of stochasticity in

the onset and progression of cardiac damage typical of patients with DMD. We believe this

reflects the level of initial damage to the affected heart, which is then amplified as the ensuing

inflammation becomes chronic.

We find chronic cardiac inflammation in D2-*mdx* is marked by higher level of activation of

inflammatory and immune response, in part by higher expression of chemokines that attract

these immune cells (**Fig. 2, 3**). This inflammatory response involves accumulation of

Spp1(OPN)/*Lgals3*(GAL-3) expressing pathogenic macrophages that we recently identified by

single cell RNAseq analysis of the skeletal muscles from *mdx* mice[45]. Osteopontin secreted by

these macrophages promote skeletal muscle fibrosis by activating the stromal progenitors and

we suggest a similar mechanism may be in place following accumulation of these macrophage

in the damaged heart (**Fig. 3**). In support of this mechanism, we find these GAL-3+

macrophages enriched in the same pericardial region that are enriched in COLA1 indicative of

active fibrosis (**Fig. 3**). The indication that this is an active phenomenon comes from the

concomitant enrichment of other ECM building (fibronectin) and degrading (TIMP/MMP)

components, along with leukocyte attracting (CCCL3/8) and resolving (FPR1/2) signaling (**Fig.**

2-4). This dynamics is indicative of a likely shift in the equilibrium of these two opposing -

inflammation building and resolving signals, rebalancing which could provide a likely beneficial

effect for the affected region. This is in line with the fact that while acute inflammation following

tissue damage is essential for cardiac repair, chronic inflammation can be disruptive[62]. To

address this ensuing imbalance of these two opposing inflammatory mechanisms, we made use

of the pro-resolving therapy that unlike GCs, precisely targets the resolution of inflammation by

activating FPR2 signaling, but not blocking activation of inflammation by NFκb or related pro-

inflammatory pathways. This approach showed excellent promise such that a short (3 week)

treatment of juvenile D2-*mdx* mice allowed full resolution of inflammation and prevented any

subsequent fibrotic cardiac damage detected histologically as well as by way of aberrant

molecular signature including *Gal3*/*Spp1* macrophages and *Col1A1* and *Fn1* expressing stromal

cells (**Fig. 5**).

In summary, our studies introduce the D2-*mdx* as a model that manifests pediatric-onset

cardiomyopathy. We identified dysregulation of inflammatory and ECM remodeling pathways as

key contributors. Specifically, our findings highlight that the excessive and unresolved

inflammatory response involving pathogenic macrophages, contributes to chronic inflammation

and fibrosis in the D2-*mdx* model. Finally, our use of FPR2-targeting therapy provides a

potential approach to prevent pediatric-onset cardiac pathologies in DMD.

Author Contributions. This study was conceived, and the experiments designed by JSN and

JKJ with inputs from all authors. RNAseq sample processing was conducted by JSN, RH, DAM,

and KP; bioinformatic analyses were performed by and AL, PU, SB, ST, and VB. AL conducted

molecular analyses with help from YJM. Histopathological analyses were conducted by AL, RH,

YJM, IHG, and GW. The manuscript was written by JSN, AL, and JKJ, and edited by all authors.

JKJ and JSN obtained funding and provided oversight for pursuit of the study.

Acknowledgements. This work was supported by funds from Children's National Research

Institute (JSN), the Foundation to Eradicate Duchenne (JKJ). JKJ and JSN acknowledge

support from Department of Defense DMDRP (W81XWH2110711; W81XWH2110680) for their

work on DMD. Microscopy was performed at the Cell and Tissue Microscopy Core supported by

CNRI and The National Institutes of Health NICHD (P50HD105328).

Competing interests. The authors have no competing or financial interests to declare.

References.

1.   Duan D, Goemans N, Takeda S, Mercuri E, Aartsma-Rus A. Duchenne muscular

dystrophy. Nat Rev Dis Primers. 2021;7(1):13.

2.   Hoffman EP, Brown RH, Jr., Kunkel LM. Dystrophin: the protein product of the

Duchenne muscular dystrophy locus. Cell. 1987;51(6):919-28.

3.   Koenig M, Hoffman EP, Bertelson CJ, Monaco AP, Feener C, Kunkel LM. Complete

cloning of the Duchenne muscular dystrophy (DMD) cDNA and preliminary genomic

organization of the DMD gene in normal and affected individuals. Cell. 1987;50(3):509-17.

4.   Haslett JN, Sanoudou D, Kho AT, Bennett RR, Greenberg SA, Kohane IS, et al. Gene

expression comparison of biopsies from Duchenne muscular dystrophy (DMD) and normal

skeletal muscle. Proceedings of the National Academy of Sciences of the United States of

America. 2002;99(23):15000-5.

5.   Meyers TA, Townsend D. Cardiac Pathophysiology and the Future of Cardiac Therapies

in Duchenne Muscular Dystrophy. Int J Mol Sci. 2019;20(17).

6.   Serrano AL, Munoz-Canoves P. Fibrosis development in early-onset muscular

dystrophies: Mechanisms and translational implications. Semin Cell Dev Biol. 2017;64:181-90.

7.   Shin J, Tajrishi MM, Ogura Y, Kumar A. Wasting mechanisms in muscular dystrophy.

The international journal of biochemistry & cell biology. 2013;45(10):2266-79.

8.   Mavrogeni S, Markousis-Mavrogenis G, Papavasiliou A, Kolovou G. Cardiac

involvement in Duchenne and Becker muscular dystrophy. World J Cardiol. 2015;7(7):410-4.

9.   Melacini P, Fanin M, Danieli GA, Villanova C, Martinello F, Miorin M, et al. Myocardial

involvement is very frequent among patients affected with subclinical Becker's muscular

dystrophy. Circulation. 1996;94(12):3168-75.

10.   Solheim TA, Fornander F, Raja AA, Mogelvang R, Poulsen NS, Duno M, et al. Cardiac

Involvement in Women With Pathogenic Dystrophin Gene Variants. Front Neurol.

2021;12:707838.

11.   Finsterer J, Stollberger C. The heart in human dystrophinopathies. Cardiology.

2003;99(1):1-19.

12.   Van Erp C, Loch D, Laws N, Trebbin A, Hoey AJ. Timeline of cardiac dystrophy in 3-18-

month-old MDX mice. Muscle & nerve. 2010;42(4):504-13.

13.   Li W, Liu W, Zhong J, Yu X. Early manifestation of alteration in cardiac function in

dystrophin deficient mdx mouse using 3D CMR tagging. J Cardiovasc Magn Reson.

2009;11(1):40.

14.   Gordish-Dressman H, Willmann R, Dalle Pazze L, Kreibich A, van Putten M,

Heydemann A, et al. "Of Mice and Measures": A Project to Improve How We Advance

Duchenne Muscular Dystrophy Therapies to the Clinic. J Neuromuscul Dis. 2018;5(4):407-17.

15.   Grady RM, Teng H, Nichol MC, Cunningham JC, Wilkinson RS, Sanes JR. Skeletal and

cardiac myopathies in mice lacking utrophin and dystrophin: a model for Duchenne muscular

dystrophy. Cell. 1997;90(4):729-38.

16.   Aartsma-Rus A, van Putten M, Mantuano P, De Luca A. On the use of D2.B10-

Dmdmdx/J (D2.mdx) Versus C57BL/10ScSn-Dmdmdx/J (mdx) Mouse Models for Preclinical

Studies on Duchenne Muscular Dystrophy: A Cautionary Note from Members of the TREAT-

NMD Advisory Committee on Therapeutics. J Neuromuscul Dis. 2023;10(1):155-8.

17.   De Giorgio D, Novelli D, Motta F, Cerrato M, Olivari D, Salama A, et al. Characterization

of the Cardiac Structure and Function of Conscious D2.B10-Dmd(mdx)/J (D2-mdx) mice from

16-17 to 24-25 Weeks of Age. Int J Mol Sci. 2023;24(14).

18.   Heydemann A, Ceco E, Lim JE, Hadhazy M, Ryder P, Moran JL, et al. Latent TGF-beta-

binding protein 4 modifies muscular dystrophy in mice. J Clin Invest. 2009;119(12):3703-12.

19.   Flanigan KM, Ceco E, Lamar KM, Kaminoh Y, Dunn DM, Mendell JR, et al. LTBP4

genotype predicts age of ambulatory loss in Duchenne muscular dystrophy. Annals of

neurology. 2013;73(4):481-8.

20.   Tandon A, Villa CR, Hor KN, Jefferies JL, Gao Z, Towbin JA, et al. Myocardial fibrosis

burden predicts left ventricular ejection fraction and is associated with age and steroid treatment

duration in duchenne muscular dystrophy. J Am Heart Assoc. 2015;4(4).

21.   Schram G, Fournier A, Leduc H, Dahdah N, Therien J, Vanasse M, et al. All-cause

mortality and cardiovascular outcomes with prophylactic steroid therapy in Duchenne muscular

dystrophy. J Am Coll Cardiol. 2013;61(9):948-54.

22.   Bourke J, Turner C, Bradlow W, Chikermane A, Coats C, Fenton M, et al. Cardiac care

of children with dystrophinopathy and females carrying DMD-gene variations. Open Heart.

2022;9(2).

23.   Janssen PM, Murray JD, Schill KE, Rastogi N, Schultz EJ, Tran T, et al. Prednisolone

attenuates improvement of cardiac and skeletal contractile function and histopathology by

lisinopril and spironolactone in the mdx mouse model of Duchenne muscular dystrophy. PloS

one. 2014;9(2):e88360.

24.   Heier CR, Yu Q, Fiorillo AA, Tully CB, Tucker A, Mazala DA, et al. Vamorolone targets

dual nuclear receptors to treat inflammation and dystrophic cardiomyopathy. Life Sci Alliance.

2019;2(1).

25.   Mavrogeni S, Giannakopoulou A, Papavasiliou A, Markousis-Mavrogenis G, Pons R,

Karanasios E, et al. Cardiac profile of asymptomatic children with Becker and Duchenne

muscular dystrophy under treatment with steroids and with/without perindopril. BMC Cardiovasc

Disord. 2017;17(1):197.

26.   Guglieri M, Bushby K, McDermott MP, Hart KA, Tawil R, Martens WB, et al. Effect of

Different Corticosteroid Dosing Regimens on Clinical Outcomes in Boys With Duchenne

Muscular Dystrophy: A Randomized Clinical Trial. JAMA. 2022;327(15):1456-68.

27.   Butterfield RJ, Kirkov S, Conway KM, Johnson N, Matthews D, Phan H, et al. Evaluation

of effects of continued corticosteroid treatment on cardiac and pulmonary function in non-

ambulatory males with Duchenne muscular dystrophy from MD STARnet. Muscle & nerve.

2022;66(1):15-23.

28.   Sinniah A, Yazid S, Flower RJ. From NSAIDs to Glucocorticoids and Beyond. Cells.

2021;10(12).

29.   Gobbetti T, Cooray SN. Annexin A1 and resolution of inflammation: tissue repairing

properties and signalling signature. Biol Chem. 2016;397(10):981-93.

30.   D'Acquisto F, Perretti M, Flower RJ. Annexin-A1: a pivotal regulator of the innate and

adaptive immune systems. Br J Pharmacol. 2008;155(2):152-69.

31.   Perretti M, Dalli J. Resolution Pharmacology: Focus on Pro-Resolving Annexin A1 and

Lipid Mediators for Therapeutic Innovation in Inflammation. Annu Rev Pharmacol Toxicol.

2023;63:449-69.

32.   Tourki B, Kain V, Pullen AB, Norris PC, Patel N, Arora P, et al. Lack of resolution sensor

drives age-related cardiometabolic and cardiorenal defects and impedes inflammation-

resolution in heart failure. Mol Metab. 2020;31:138-49.

33.   Qin C, Buxton KD, Pepe S, Cao AH, Venardos K, Love JE, et al. Reperfusion-induced

myocardial dysfunction is prevented by endogenous annexin-A1 and its N-terminal-derived

peptide Ac-ANX-A1(2-26). Br J Pharmacol. 2013;168(1):238-52.

34.   Kain V, Ingle KA, Colas RA, Dalli J, Prabhu SD, Serhan CN, et al. Resolvin D1 activates

the inflammation resolving response at splenic and ventricular site following myocardial

infarction leading to improved ventricular function. J Mol Cell Cardiol. 2015;84:24-35.

35.   Kain V, Liu F, Kozlovskaya V, Ingle KA, Bolisetty S, Agarwal A, et al. Resolution Agonist

15-epi-Lipoxin A(4) Programs Early Activation of Resolving Phase in Post-Myocardial Infarction

Healing. Scientific reports. 2017;7(1):9999.

36.   Garcia RA, Lupisella JA, Ito BR, Hsu MY, Fernando G, Carson NL, et al. Selective FPR2

Agonism Promotes a Proresolution Macrophage Phenotype and Improves Cardiac Structure-

Function Post Myocardial Infarction. JACC Basic Transl Sci. 2021;6(8):676-89.

37.   Lupisella J, St-Onge S, Carrier M, Cook EM, Wang T, Sum C, et al. Molecular

Mechanisms of Desensitization Underlying the Differential Effects of Formyl Peptide Receptor 2

Agonists on Cardiac Structure-Function Post Myocardial Infarction. ACS Pharmacol Transl Sci.

2022;5(10):892-906.

38.   Bulfield G, Siller WG, Wight PA, Moore KJ. X chromosome-linked muscular dystrophy

(mdx) in the mouse. Proceedings of the National Academy of Sciences of the United States of

America. 1984;81(4):1189-92.

39.   Mazala DA, Novak JS, Hogarth MW, Nearing M, Adusumalli P, Tully CB, et al. TGF-

beta-driven muscle degeneration and failed regeneration underlie disease onset in a DMD

mouse model. JCI Insight. 2020;5(6).

40.   Mazala DAG, Hindupur R, Moon YJ, Shaikh F, Gamu IH, Alladi D, et al. Altered muscle

niche contributes to myogenic deficit in the D2-mdx model of severe DMD. Cell Death Discov.

2023;9(1):224.

41.   Kayashima Y, Clanton CA, Lewis AM, Sun X, Hiller S, Huynh P, et al. Reduction of

Stabilin-2 Contributes to a Protection Against Atherosclerosis. Front Cardiovasc Med.

2022;9:818662.

42.   Nishimura D, Sakai H, Sato T, Sato F, Nishimura S, Toyama-Sorimachi N, et al. Roles of

ADAM8 in elimination of injured muscle fibers prior to skeletal muscle regeneration. Mech Dev.

2015;135:58-67.

43.   Wang M, Gao X, Zhao K, Chen H, Xu M, Wang K. Effect of TREM2 on Release of

Inflammatory Factor from LPS-stimulated Microglia and Its Possible Mechanism. Ann Clin Lab

Sci. 2019;49(2):249-56.

44.   Villalta SA, Nguyen HX, Deng B, Gotoh T, Tidball JG. Shifts in macrophage phenotypes

and macrophage competition for arginine metabolism affect the severity of muscle pathology in

muscular dystrophy. Human molecular genetics. 2009;18(3):482-96.

45.   Coulis G, Jaime D, Guerrero-Juarez C, Kastenschmidt JM, Farahat PK, Nguyen Q, et al.

Single-cell and spatial transcriptomics identify a macrophage population associated with skeletal

muscle fibrosis. bioRxiv. 2023.

46.   Coulis G, Jaime D, Guerrero-Juarez C, Kastenschmidt JM, Farahat PK, Nguyen Q, et al.

Single-cell and spatial transcriptomics identify a macrophage population associated with skeletal

muscle fibrosis. Sci Adv. 2023;9(27):eadd9984.

47.   Sanchez-Garcia S, Jaen RI, Fernandez-Velasco M, Delgado C, Bosca L, Prieto P.

Lipoxin-mediated signaling: ALX/FPR2 interaction and beyond. Pharmacol Res.

2023;197:106982.

48.   Yang M, Song XQ, Han M, Liu H. The role of Resolvin D1 in liver diseases.

Prostaglandins Other Lipid Mediat. 2022;160:106634.

49.   Perretti M, Godson C. Formyl peptide receptor type 2 agonists to kick-start resolution

pharmacology. Br J Pharmacol. 2020;177(20):4595-600.

50.   Dahlgren C, Gabl M, Holdfeldt A, Winther M, Forsman H. Basic characteristics of the

neutrophil receptors that recognize formylated peptides, a danger-associated molecular pattern

generated by bacteria and mitochondria. Biochem Pharmacol. 2016;114:22-39.

51.   Godson C, Mitchell S, Harvey K, Petasis NA, Hogg N, Brady HR. Cutting edge: lipoxins

rapidly stimulate nonphlogistic phagocytosis of apoptotic neutrophils by monocyte-derived

macrophages. Journal of immunology. 2000;164(4):1663-7.

52.   Fredman G, Kamaly N, Spolitu S, Milton J, Ghorpade D, Chiasson R, et al. Targeted

nanoparticles containing the proresolving peptide Ac2-26 protect against advanced

atherosclerosis in hypercholesterolemic mice. Science translational medicine.

2015;7(275):275ra20.

53.   Scannell M, Flanagan MB, deStefani A, Wynne KJ, Cagney G, Godson C, et al.

Annexin-1 and peptide derivatives are released by apoptotic cells and stimulate phagocytosis of

apoptotic neutrophils by macrophages. Journal of immunology. 2007;178(7):4595-605.

54.   Vetrone SA, Montecino-Rodriguez E, Kudryashova E, Kramerova I, Hoffman EP, Liu SD,

et al. Osteopontin promotes fibrosis in dystrophic mouse muscle by modulating immune cell

subsets and intramuscular TGF-beta. J Clin Invest. 2009;119(6):1583-94.

55.   Capote J, Kramerova I, Martinez L, Vetrone S, Barton ER, Sweeney HL, et al.

Osteopontin ablation ameliorates muscular dystrophy by shifting macrophages to a pro-

regenerative phenotype. J Cell Biol. 2016;213(2):275-88.

56.   Kramerova I, Kumagai-Cresse C, Ermolova N, Mokhonova E, Marinov M, Capote J, et

al. Spp1 (osteopontin) promotes TGFbeta processing in fibroblasts of dystrophin-deficient

muscles through matrix metalloproteinases. Human molecular genetics. 2019;28(20):3431-42.

57.   Hirata A, Masuda S, Tamura T, Kai K, Ojima K, Fukase A, et al. Expression profiling of

cytokines and related genes in regenerating skeletal muscle after cardiotoxin injection: a role for

osteopontin. The American journal of pathology. 2003;163(1):203-15.

58.   Bello L, Pegoraro E. The "Usual Suspects": Genes for Inflammation, Fibrosis,

Regeneration, and Muscle Strength Modify Duchenne Muscular Dystrophy. J Clin Med.

2019;8(5).

59.   Aurora AB, Porrello ER, Tan W, Mahmoud AI, Hill JA, Bassel-Duby R, et al.

Macrophages are required for neonatal heart regeneration. J Clin Invest. 2014;124(3):1382-92.

60.   Lavine KJ, Epelman S, Uchida K, Weber KJ, Nichols CG, Schilling JD, et al. Distinct

macrophage lineages contribute to disparate patterns of cardiac recovery and remodeling in the

neonatal and adult heart. Proceedings of the National Academy of Sciences of the United

States of America. 2014;111(45):16029-34.

61.   Heydemann A, Ceco E, Lim JE, Hadhazy M, Ryder P, Moran JL, et al. Latent TGF-β–

binding protein 4 modifies muscular dystrophy in mice. The Journal of clinical investigation.

2009;119(12):3703.

62.   Frangogiannis NG. Regulation of the inflammatory response in cardiac repair. Circ Res.

2012;110(1):159-73.

Figures and Legends.

Figure 1. Histopathology associated with disease onset in juvenile D2-*mdx* hearts.

Images show of hearts from juvenile (6 ± 0.5 wk) D2-*mdx* and B10-*mdx* mice at disease onset.

A-B. Whole tissue images of the matched orientation of D2-*mdx* (**A**) and B10-*mdx* (**B**) hearts

showing ventricular and atrial fibro-calcified damage (marked by arrowheads). **C-D.** Image

showing cross-section of juvenile D2-*mdx* and B10-*mdx* hearts through the ventricular lumen,

stained for histological features by H&E (**C**), and for fibrosis by Sirius Red (**D**). Arrowheads mark

areas of fibrosis. **E-F.** Image (**E**) and quantification (**F**) showing a portion of heart cross-section

from juvenile D2-*mdx* and B10-*mdx* hearts, showing damaged tissue areas characterized by the

presence of interstitial fibrosis, inflammatory cells and damaged cardiomyocytes. **G-H.** Image

(**G**) and quantification (**H**) showing a portion of heart cross-section from juvenile D2-*mdx* and

B10-*mdx* hearts labeled with Sirius Red to mark fibrotic tissue area in hearts from juvenile D2-

mdx and B10-*mdx* mice. Data represent mean ± SD from $n = 12$ hearts per cohort. Statistical

analyses performed using Mann Whitney U test; **p < 0.05*, ***p < 0.001*, ****p < 0.001*. For full

tissue images of E, G, refer to Supplementary Fig. 1.

Figure 2. RNAseq analysis at disease onset in D2-*mdx* and age-matched B10-*mdx* hearts.

Juvenile D2-*mdx* and B10-*mdx* hearts at disease onset (6 wk ± 0.5 wk) were analyzed by bulk

tissue RNAseq. **A.** Dimensionality reduction of whole transcriptomic data via PCA (n=3-4

hearts/genotype) to assess sample clustering and inter-/intra-sample variance. **B.** Differential

gene expression analysis depicted via heatmap plot of 2,719 DEGs observed between D2-*mdx*

(blue) and B10-*mdx* (red), with 1,586 genes upregulated for D2-*mdx* and 1,133 genes

upregulated for B10-*mdx*. Expression is z-score values of variance-stabilizing transformation

(VST) normalized data. **C.** Gene Ontology (GO) analysis performed using Cytoscape and

EnrichmentMap plugins to identify networks of related GO terms groups found upregulated (red

dots) in juvenile D2-*mdx* hearts relative to B10-*mdx.* Pink clusters refer to inflammatory-related

GO terms, while blue clusters refer to extracellular matrix-related GO terms. **D-E.** Boxplots

showing VST normalized gene expression levels for top 20 differentially expressed

inflammation-related (D; GOBP:Inflammatory Response) and extracellular matrix-related (E;

GOBP:External Encapsulating Structure) transcripts observed between juvenile D2-*mdx* and

B10-*mdx* hearts. Refer to Supplementary Fig. 3 for additional details.

Figure 3. Targeted analysis of inflammatory response at disease onset in D2-*mdx* B10-

***mdx* hearts. A.** qRT-PCR analysis of a distinct cohort of D2-*mdx* and B10-*mdx* hearts to assess

the expression of inflammatory genes identified by RNAseq analysis cohort to be differentially

expressed. Transcripts include top dysregulated genes involved in leukocyte activation,

migration and chemotaxis and regulation of inflammatory response and leukocyte-mediated

immunity (*Ccl3, Ccl8, Stab2, Adam8, Trem2, Il7r*). **B.** RT-PCR analysis of a distinct cohort of

D2-*mdx* and B10-*mdx* hearts to assess inflammatory genes that show broad dysregulation of

neutrophil and macrophage response in juvenile D2-*mdx* hearts. **C-D.** Images showing

immunostaining for pan-macrophage marker, F4/80 (green), and pro-inflammatory, pathogenic

macrophage marker, GAL-3 (red), in juvenile D2-*mdx* (**C**) and B10-*mdx* (**D**). Data represent

mean ± SD from n = 7-9 hearts per cohort. Statistical analyses performed using Mann Whitney U

test; *$p < 0.05$, **$p < 0.001$, ***$p < 0.001$. For age-matched WT controls, refer to Supplementary

Fig. 4.

Figure 4. Targeted analysis of extracellular matrix remodeling response at disease onset

in D2-*mdx* B10-*mdx* hearts. A. qRT-PCR analysis of a distinct cohort of D2-*mdx* and B10-*mdx*

hearts to assess the expression of extracellular matrix-associated genes involved in matrix

organization/re-organization (*Spp1, Fn1, Col1a1, Itgax, Mmp12, Timp1*) that are identified to be

dysregulated by the RNAseq cohort. **B-C.** Images showing extracellular matrix distribution

visualized using wheat germ agglutinin (WGA, pink) within, and surrounding, areas of cardiac

damage in juvenile D2-*mdx* (**B**), and B10-*mdx* (**C**). **B'-C'.** Zoom of the dotted area from whole

cross-sectional images showing immunostaining for COL1A1 within the extracellular matrix

shows increased COL1A1 expression in damaged D2-*mdx* hearts (**B'**), relative to B10-*mdx* (**C'**),

indicative of early-onset endomysial fibrosis. Data represent mean ± SD from $n = 7\text{-}9$ hearts per

cohort. Statistical analyses performed using Mann Whitney U test; *$p < 0.05$, **$p < 0.001$,

***$p < 0.001$. For age-matched WT controls, refer to Supplementary Fig. 4.

Figure 5. Pro-resolving therapy to mitigate cardiac disease onset in juvenile D2-*mdx*. A.

Schematic describing the inflammatory response following cardiac injury in health (dotted black

trace) or dystrophic (red trace), showing acute versus chronic inflammation respectively. Use of

anti-inflammatory drug (purple trace) lowers inflammatory response blunting inflammation,

instead use of pro-resolving therapy (dotted green trace) does not impact the onset of

inflammation but helps clear inflammation preventing the inflammation to become chronic. **B.**

Schematic detailing the pre-clinical testing of pro-resolving drug, BMS-986235 (6.0mg/kg, 3 wk

daily administration) in D2-*mdx* mice (n=6) just prior to disease onset (18-19 days old). **C.**

Whole tissue images of the matched orientation of hearts showing ventricular and atrial fibro-

calcified damage (marked by arrowheads) in saline or BMS-986235-treated D2-*mdx* mice. **D.**

Image showing cross-section of D2-*mdx* heart stained for histological features by H&E from

mice treated with saline or BMS-986235. **E.** Images showing cross-section of D2-*mdx* heart

immunostained for pan-macrophage marker, F4/80 (red) and counterstained with WGA (green)

and DAPI (blue) to mark the ECM and nuclei, respectively. **F.** Image showing heart cross-

section immunostained for COL1A1 and counterstained with DAPI (blue) to visualize nuclei. **G-**

H. qRT-PCR analysis of inflammatory (**G**) and (**H**) extracellular matrix genes to assess the

effect of drug treatment of D2-*mdx* mice (red triangles), as compared with saline-treated

controls (black triangles). Data represent mean ± SD from $n = 6$ hearts per cohort. Statistical

analyses performed using Mann Whitney U test; *$p < 0.05$, **$p < 0.001$, ***$p < 0.001$.

Failure to resolve inflammation contributes to juvenile onset cardiomyopathy in a mouse model of Duchenne Muscular Dystrophy.

James S. Novak[1,2†], Amy Lischin[1,3], Prech Uapinyoying[1,4], Ravi Hindupur[1], Young Jae Moon[1,5], Surajit Bhattacharya[1], Sarah Tiufekchiev[1,6], Victoria Barone[1,3], Davi A. G. Mázala[1,7], Iteoluwakishi H. Gamu[1], Gabriela Walters[1], Karuna Panchapakesan[1], Jyoti K. Jaiswal[1,2†]

Supplemental Material.

Supplemental Figure 1. Cardiac histopathology in juvenile D2-*mdx* model. A. Alizain Red staining of juvenile D2-*mdx* and B10-*mdx* hearts (whole cross-section) showing right ventricular (RV) heart damage and calcification in juvenile D2-*mdx* hearts. Scale bars indicate 200µm. **B-C.** High magnification images from panel A, showing Alizarin Red staining of juvenile D2-*mdx* and B10-*mdx* hearts, and corresponding quantification of calcified fiber area per total tissue area. of fibrosis, damage and calcification are highlighted by black arrowheads (**C**). Scale bars indicate 100µm. Statistical analyses performed using Mann Whitney U test; ***$p < 0.001$.

Supplemental Figure 2. GSEA enrichment plot analysis of top gene ontology (GO) hits from differential gene expression analysis in juvenile D2-*mdx* and B10-*mdx* hearts. A-F. Enrichment plot analysis of top immune-related (**A, C, E, F**) and extracellular matrix-related (**B, D**) GO biological process (BP) terms obtained using GSEA, including Inflammatory Response (**A**), Encapsulating Structure Organization (**B**), Myeloid Luekocyte Migration (**C**), Collagen Metabolic Process (**D**), Neutrophil Chemotaxis (**E**), and Granulocyte Chemotaxis (**F**).

Supplemental Figure 3. Gene expression analysis of D2-WT and B10-WT hearts for selected immune and extracellular matrix targets. A. Relative gene expression analysis for dysregulated immune-related genes between D2-*mdx* vs. B10-*mdx* (**Figure 3**) assessed in juvenile D2-WT and B10-WT hearts. **B.** Relative gene expression analysis for dysregulated immune-related genes between D2-*mdx* vs. B10-*mdx* (**Figure 4**) assessed in juvenile D2-WT and B10-WT hearts.