Title:

Pyruvate kinase modulates the link between β-cell fructose metabolism and insulin

secretion

Authors:

Naoya Murao[1,2*], Risa Morikawa[1], Yusuke Seino[1,2], Kenju Shimomura[3], Yuko Maejima[3],

Tamio Ohno[4], Norihide Yokoi[5], Yuichiro Yamada[1,2], Atsushi Suzuki[1]

Affiliations:

[1]   Department of Endocrinology, Diabetes and Metabolism, Fujita Health University, School

of Medicine, Toyoake, Japan

[2]   Yutaka Seino Distinguished Center for Diabetes Research, Kansai Electric Power Medical

Research Institute, Kyoto, Japan

[3]   Department of Bioregulation and Pharmacological Medicine, Fukushima Medical

University School of Medicine, Fukushima, Japan

[4]   Graduate School of Medicine Center for Research of Laboratory Animals and Medical

Research Engineering Division for Research of Laboratory Animals, Graduate School of

Medicine, Nagoya University, Nagoya, Japan

[5]   Laboratory of Animal Breeding and Genetics, Graduate School of Agriculture, Kyoto

University, Kyoto, Japan

*Corresponding author: Naoya Murao, M.D., PhD

Departments of Endocrinology, Diabetes and Metabolism, Fujita Health University,

School of Medicine, 1-98 Dengakugakubo, Toyoake, Aichi 4701192 Japan

Phone: +81-562-93-2699

Fax: +81-562-95-1879

E-mail: naoya.murao@fujita-hu.ac.jp

ORCID: 0000-0002-7424-4476

1   GRAPHICAL ABSTRACT

Pyruvate kinase modulates the link between β-cell fructose metabolism and insulin
secretion.

Naoya Murao[*], Risa Morikawa, Yusuke Seino, Kenju Shimomura, Yuko Maejima,
Tamio Ohno, Norihide Yokoi, Yuichiro Yamada, Atsushi Suzuki.

Left: Fructose-stimulated insulin secretion (FSIS) is driven by sweet taste receptor
(STR)-mediated PLC signaling in pancreatic β-cells. Meanwhile, fructose metabolism
does not promote FSIS because fructose causes accumulation of fructose 1-
phosphate (F1P), which suppresses pyruvate kinase M2 (PKM2), lowering the
ATP/ADP ratio.

Right: A small-molecule PK activator counteracted F1P-mediated PKM2 inhibition,
prevented ATP decrease, and substantially enhanced FSIS in normal and diabetic
mouse β-cells. Thus, PK has been identified as a key regulator linking β-cell fructose
metabolism and FSIS.

2   ABSTRACT

Glucose triggers insulin secretion from pancreatic β-cells through intracellular glucose
metabolism, ATP production, and closure of ATP-sensitive K$^+$ channels (K$_{ATP}$ channels).
Fructose also stimulates insulin secretion, but the underlying mechanisms remain unclear. This
study investigated the contribution of phospholipase C (PLC) signaling and fructose
metabolism to fructose-stimulated insulin secretion (FSIS) using MIN6-K8 clonal β-cells and
mouse islets.

Fructose-induced PLC activation, assessed by inositol 1-phosphate accumulation, was
reduced in fructose-unresponsive β-cell models, such as diabetic mouse islets and K$_{ATP}$
channel-deficient β-cells, suggesting that β-cell fructose responsiveness is primarily
determined by PLC signaling.

Although FSIS was dependent on K$_{ATP}$ channels and Ca^{2+} influx, the ATP/ADP ratio was
unexpectedly lowered by fructose, and suppression of intracellular fructose metabolism hardly
affected FSIS. Metabolic flux analysis revealed that the accumulation of fructose 1-phosphate
(F1P) suppressed pyruvate kinase (PK) activity, contributing to ATP depletion. Strikingly, a
small-molecule PK activator, TEPP-46, antagonized F1P-mediated PK suppression, prevented
the drop in the ATP/ADP ratio, and restored FSIS in MIN6-K8 cells, normal mouse islets, and
fructose-unresponsive diabetic mouse islets.

These findings revealed the metabolic effects of fructose in β-cells and identified PK as a
key regulator linking β-cell fructose metabolism and FSIS, thereby providing new insights into
the mechanisms of insulin secretion and potential therapeutic targets for fructose-associated
metabolic diseases.

3 ABBREVIATIONS

2PG, 2-phosphoglycerate
3PG, 3-phosphoglycerate
DHAP, dihydroxyacetone phosphate
FBP, fructose 1,6-bisphosphate; F1P, fructose 1-phosphate
FSIS, fructose-stimulated insulin secretion
G6P, glucose 6-phosphate; F6P, fructose 6-phosphate
GA3P, glyceraldehyde 3-phosphate
GCK, glucokinase
GCKR, glucokinase regulatory protein
GSIS, glucose-stimulated insulin secretion

IP1, inositol 1-phosphate

K_{ATP} channel, ATP-sensitive K^+ channel

KHK, Ketohexokinase

MASLD, metabolic dysfunction-associated steatotic liver disease

OAA, oxaloacetate

PEP, phosphoenolpyruvate

PK, pyruvate kinase

PLC, phospholipase C

4   KEYWORDS

Pancreatic beta cells, insulin secretion, fructose, pyruvate kinase, glycolysis, diabetes mellitus

5   INTRODUCTION

Dietary monosaccharides, such as glucose, fructose, and mannose, stimulate pancreatic β-cells

to increase insulin secretion (Grodsky et al., 1963; Ashcroft et al., 1972). The process of

glucose-stimulated insulin secretion (GSIS) is critical for glucose homeostasis, and numerous

studies have delineated its molecular mechanisms. It is well established that GSIS is regulated

by β-cell glucose metabolism through two signaling pathways: the triggering and amplifying

pathways (Henquin, 2000; Henquin, 2009). The triggering pathway refers to the increase in

intracellular $[Ca^{2+}]$ after a sequence of events: an increase in intracellular ATP levels, closure

of ATP-sensitive K^+ (K_{ATP}) channels, membrane depolarization, and opening of voltage-

dependent Ca^{2+} channels (VDCCs). The amplifying pathway augments the triggering pathway

only when the latter is already active through various glucose-related metabolites such as

adenine nucleotides, citrate, GTP, and glutamate (Prentki et al., 2013; Murao et al., 2017; Han

et al., 2021).

Fructose is the second most abundant monosaccharide in the diet after glucose (Larke et al.,

2023). It is widely found in fruits and certain sweeteners, making it a significant source of

energy for many people. Fructose has been known to stimulates insulin secretion at

physiological glucose levels (Grodsky et al., 1963; Ashcroft et al., 1972; Zawalich et al., 1977;

Grant et al., 1980). However, the mechanism of fructose-stimulated insulin secretion (FSIS) is

less well understood than that of glucose-stimulated insulin secretion (GSIS).

In various cell types, fructose is transported primarily through GLUT5 (SLC2A5) and is

converted to F1P by ketohexokinase (KHK), the first rate-limiting enzyme in fructolysis

(Hannou et al., 2018). It is then metabolized to glyceraldehyde 3-phosphate (GA3P) and

dihydroxyacetone phosphate (DHAP) and enters glycolytic/gluconeogenic metabolite pool,

part of which is used for ATP production (Hannou, 2018, see also Figure 2).

Thus, it is reasonable to assume that ATP production by fructose metabolism stimulates

FSIS in a pathway similar to that of glucose. However, this relationship has not yet been

thoroughly investigated. To address this, we examined β-cell fructose metabolism and its

functional relationship with insulin secretion using clonal β-cell lines and mouse islets as a

model. Unexpectedly, we found that fructose metabolism in β-cells did not contribute to FSIS.

Through detailed metabolic profiling, we discovered that fructose inhibits pyruvate kinase (PK)

and hinders glycolysis. Remarkably, we showed that PK activators counteracted this inhibition,

enhancing FSIS, which was also effective in fructose-unresponsive diabetic mouse islets.

6   MATERIALS AND METHODS

6.1. Animals

C57BL/6JJcl (RRID:IMSR_JCL:JCL:MIN-0003) mice were purchased from CLEA Japan

(Tokyo, Japan) and used for experiments at 19–20 weeks of age. NSY.B6-*Tyr+*, *A^y*/Hos mice

were a gift from Hoshino Laboratory Animals (Ibaraki, Japan) and were used for experiments

at 19–20 weeks of age. Mice were maintained under specific-pathogen-free conditions at 23 ±

2 °C and 55 ± 10% relative humidity with 12-hour light-dark cycles (8am–8pm), with free

access to water and standard chow CE-2 (CLEA Japan, Tokyo, Japan). The health status of the

mice was checked regularly. All experiments were performed using male mice. Body weight

and blood glucose levels were measured at 9am in the ad-lib-fed state. The number of mice

analyzed is indicated in the figure legends. All animal experiments were performed with the

approval APU22080 by the Institutional Animal Care and Use Committee of Fujita Health

University, complying with the Guidelines for Animal Experimentation at Fujita Health

University and current Japanese guidelines and regulations for scientific and ethical animal

experimentation (Kagiyama et al., 2006).

6.2. Cell lines

MIN6-K8 cells were generated as previously described (Iwasaki et al., 2010) and were kindly

provided by Professor Junichi Miyazaki (Osaka University). *Kcnj11$^{-/-}$* β-cells (clone *Kcnj11$^{-/-}$*

βCL1) are β-cells deficient in Kir6.2, a pore-forming subunit of the K$_{ATP}$ channel. They were

established by sub-cloning MIN6-K8 cells transfected with Cas9 nickase and guide RNA pairs

targeting mouse *Kcnj11*, as described previously (Oduori et al., 2020). All cells were cultured

in Dulbecco's modified Eagle's medium (DMEM) containing 4500 mg/L glucose (Sigma-

Aldrich, St. Louis, MO, USA, Cat# D5796) supplemented with 10% fetal bovine serum (FBS)

(BioWest, Nuaillé, France, Cat# S1400-500) and 5 ppm 2-mercaptoethanol. The cells were

maintained at 37 °C with 5% CO_2.

6.3. Plasmids

GW1-PercevalHR was a gift from Gary Yellen (Addgene plasmid # 49082;

http://n2t.net/addgene:49082; RRID: Addgene_49082) (Tantama et al., 2013).

6.4. Reagents

Krebs-Ringer bicarbonate buffer-HEPES (133.4 mM NaCl, 4.7 mM KCl, 1.2 mM KH_2PO_4,

1.2 mM $MgSO_4$, 2.5 mM $CaCl_2$, 5 mM $NaHCO_3$, 10 mM HEPES) supplemented with 0.1%

bovine-serum albumin (Sigma-Aldrich, St. Louis, MO, USA, Cat# A6003) and 2.8 mM

glucose (2.8G-KRBH) adjusted to pH 7.4 was used in in vitro experiments. Additional glucose

(final concentration: 8.8 mM), U73122 (CAS: 112648-68-7, Cat# 70740), and fructose were

added to KRBH during the stimulation period. Nifedipine (FUJIFILM Wako Pure Chemical,

Osaka, Japan, Cat# 14505781) and TEPP-46 (Selleck Chemicals, Houston, TX, USA, CAS:

1221186-53-3, Cat# S7302) were added during the pre-incubation and stimulation periods. The

reagents used for stimulation were stored as a 1000× concentrate in dimethyl sulfoxide

(DMSO) (FUJIFILM Wako Pure Chemical, Osaka, Japan, Cat# 041-29351) and diluted with

KRBH shortly before the experiment. An equal volume of DMSO was added to the vehicle

control.

6.5. Isolation of pancreatic islets from mice

Digesting solution was formulated by supplementing 0.1w/v% Collagenase from Clostridium

histolyticum (Sigma-Aldrich, St. Louis, MO, USA, Cat# C6885) to Hanks' balanced salt

solution (136.9 mM NaCl, 5.4 mM KCl, 0.8 mM $MgSO_4$, 0.3 mM Na_2HPO_4, 0.4 mM KH_2PO_4,

4.2 mM $NaHCO_3$, 10 mM HEPES, 1.3 mM $CaCl_2$, 2 mM glucose). The mice were euthanized

by isoflurane exposure. Pancreas was digested by 10-min incubation at 37 °C following

intraductal injection of digesting solution. Islets were hand-picked and separated from exocrine

tissues, transferred to 60-mm non-treated plates (AGC Techno Glass, Shizuoka, Japan, Cat#

1010-060), and cultured overnight in RPMI-1640 (Sigma-Aldrich, St. Louis, MO, USA, Cat#

R8758) supplemented with 10% FBS (BioWest, Nuaillé, France, Cat# S1400-500) and 1%

penicillin-streptomycin solution (FUJIFILM Wako Pure Chemical, Osaka, Japan, Cat# 168-

23191) at 37 °C with 5% CO_2 before the experiments.

6.6. Insulin secretion

Insulin secretion was measured using the static incubation method, as described previously

(Murao et al., 2022; Murao et al., 2024).

To measure insulin secretion from cell lines, cells were seeded in 24-well plates at a density

of 5×10^5 cells/well and cultured for 48 h. On the day of measurement, the cells were subjected

to three successive washes with 2.8G-KRBH, followed by a pre-incubation period of 30 min

with 300 μL/well of 2.8G-KRBH. Subsequently, the supernatant was replaced with 300

μL/well of fresh KRBH containing the specified stimulations and incubated for 30 min at 37 °C.

The reaction was terminated by cooling the plate on ice for ten minutes, after which the entire

supernatant was collected for the quantification of released insulin using the homogeneous

time-resolved fluorescence assay (HTRF) Insulin Ultrasensitive kit (Revvity, Waltham, MA,

USA, Cat# 62IN2PEH) in accordance with the manufacturer's instructions. Fluorescence was

measured using an Infinite F Nano+ microplate reader (Tecan, Zürich, Switzerland).

To measure insulin secretion from islets, overnight cultured islets were rinsed twice with

2.8G-KRBH, followed by pre-incubation with 2.8G-KRBH for 30 min at 37 °C. Size-matched

islets were hand-picked and dispensed in a 96-well plate (Corning, Glendale, AZ, USA, Cat#

353072) at 5 islets/well. KRBH (100 μL/well) containing the specified stimulations was added

and incubated for 30 min at 37 °C. The supernatant was subjected to insulin quantification as

described above.

6.7. Measurement of inositol 1-phosphate (IP1)

To measure IP1 content in cell lines, the cells were seeded in a 96-well plate (Corning, Glendale,

AZ, USA, Cat# 353072) at a density of 1.0×10^5 cells/well and cultured for 48h. On the day

of measurement, the cells were subjected to three successive washes with IP1 assay buffer (146

mM NaCl, 4.2 mM KCl, 50 mM LiCl, 1 mM $CaCl_2$, 0.5 mM $MgCl_2$, 10 mM HEPES, 0.1%

BSA, pH7.4) supplemented with 2.8 mM glucose (2.8G-IP1 assay buffer), followed by a pre-

incubation period of 30 min with 60 μL/well of 2.8G-IP1 assay buffer. Subsequently, 20

μL/well of IP1 assay buffer containing the specified stimulations at 4× concentration was added

to the cells and incubated for another 30 min at 37 °C. The cells were lysed by the addition of

20 μL/well lysis buffer from the HTRF IP-One Gq Detection Kit (Revvity, Waltham, MA,

USA, Cat# 62IPAPEB). 30 μL/well of the lysate were transferred to a 384-well low-volume

plate (Cat #3826; Corning, Glendale, AZ, USA) and used for IP1 quantification according to

the instructions of the HTRF IP-One Gq Detection Kit.

To measure IP1 content in mouse islets, overnight-cultured islets were rinsed twice with

2.8G-IP1 assay buffer, followed by pre-incubation with 2.8G-IP1 assay buffer for 30 min at

37 °C. Size-matched islets were hand-picked and dispensed in a 96-well plate 96-well plate

(Corning, Glendale, AZ, USA, Cat# 353072) at 5 islets/well. The islets were pre-incubated for

30 min with 45 µL/well of 2.8G-IP1 assay buffer. Subsequently, 15 µL/well of IP1 assay buffer

containing the specified stimulations at 4× concentration was added to the cells and incubated

for another 30 min at 37 °C. The cells were lysed by the addition of 15 µL/well of lysis buffer

supplied in the kit. The lysate (30 µL/well) was subjected to IP1 quantification, as described

above.

6.8.   Imaging of intracellular Ca^{2+}

Ca^{2+} imaging was performed as previously described (Murao et al., 2024). Briefly, cells were

seeded in a 35 mm glass-bottom dish (Matsunami Glass, Osaka, Japan, Cat# D11530H) at a

density of 1.28×10^5 cells/dish and cultured for 48 h. Subsequently, the cells were loaded with

1 µM Fluo-4 AM (Dojindo, Kumamoto, Japan, Cat# F312) in 2.8G-KRBH for 20 min at 37 °C

in room air. Following a brief washing, cells were loaded with 1 mL of fresh 2.8G-KRBH and

basal recordings were performed for 300 s (from time -300 to 0). Immediately after the addition

of 1 mL KRBH supplemented with stimulations at 2× concentration, recordings were resumed

for another 600 s (from time 0 to 600) with a time interval of 2 s.

Time-lapse images were obtained using a Zeiss LSM 980 Airyscan2 inverted confocal laser

scanning super-resolution microscope equipped with a Plan Apo 40×, 1.4 Oil DICII objective

lens (Carl Zeiss Microscopy, Jena, Germany). The cells were excited at 488 nm laser with 0.3%

output power, and fluorescence emission was measured at 508-579 nm. During observation,

the cells were maintained at 37 °C using an incubator XLmulti S2 DARK (Pecon, Erbach,

Germany).

Images were acquired in the frame mode at a rate of 2 frames per second and with an image

size of 212.2 × 212.2 µm (512 × 512 pixels). The obtained images were analyzed using the

ZEN 3.0 imaging software (Carl Zeiss Microscopy, Jena, Germany, RRID:SCR_021725).

Cells were randomly chosen for analysis for each stimulation, and the number of cells analyzed

is indicated in the figure legends. The fluorescence intensity of the entire cell body (F) was

monitored and normalized to the average fluorescence intensity between -300 and 0 s (F0).

6.9. Imaging of intracellular ATP/ADP ratio[+]

The Cells were seeded in a 35 mm glass-bottom dish (Matsunami Glass, Osaka, Japan, Cat#

D11530H) at a density of 6.0×10^4 cells/dish and cultured for 48 h. Subsequently, the cells

were transfected with 0.1 µg/dish of plasmid construct GW1-PercevalHR using Effectene

transfection reagent (Qiagen, Venlo, Netherlands, Cat# 301425) in accordance with the

manufacturer's instructions and were cultured for another 48 h. On the day of the measurement,

after a brief washing, the cells were loaded with 1 mL of fresh 2.8G-KRBH, and basal

recordings were performed for 300 s (from time -300 to 0). Immediately after the addition of

1 mL KRBH supplemented with stimulations at 2× concentration, recordings were resumed for

another 600 s (from time 0 to 600) with a time interval of 2 s.

The cells were excited at 445 nm and 488 nm laser with 0.8% and 0.3% output power,

respectively, and fluorescence emission was measured at 508-543 nm. The ratio (R) of the

fluorescence at 445 nm and 488 nm was calculated and normalized to the average fluorescence

intensity between -300 and 0 s (R0). The other microscope configurations were the same as

those used for Ca^{2+} imaging (Section 6.8).

6.10. Knockdown of Ketohexokinase (*Khk*) using small interfering RNA (siRNA)

siRNAs targeting *Khk* (Dharmacon, Lafayette, CO, USA, Cat# M-062217-01-0005) and non-

targeting siRNA (Dharmacon, Lafayette, CO, USA, Cat# D-001206-14-50) were reverse-

transfected using the DharmaFECT 2 transfection reagent (Dharmacon, Lafayette, CO, USA,

Cat# T-2002-03). Briefly, a complex of siRNA and DharmaFECT 2 was prepared in serum-

free DMEM (Sigma-Aldrich, St. Louis, MO, USA, Cat# D5796) at a volume of 100 µL/well

according to the manufacturer's instructions. Cells were resuspended in complete culture

media at 1.25×10^6 cells/mL. The cell suspension was then combined with

siRNA/DharmaFECT 2 complex. The final concentrations of siRNA and DharmaFECT 2 were

40 nM and 0.4%, respectively. For insulin secretion and RT-qPCR experiments, the cells were

seeded in 24-well plates (Corning, Glendale, AZ, USA, Cat# 353047) at 5×10^5 cells/500

µL/well. For immunoblotting, the cells were seeded in 12-well plates (Corning, Glendale, AZ,

USA, Cat# 353053) at 1×10^6 cells/mL/well. For metabolic flux analysis, cells were seeded in

6-well plates (Cat #353046; Corning, Glendale, AZ, USA) at a density of 2×10^6 cells/2

mL/well. The subsequent experiments were performed after a 48-hour culture.

6.11. Reverse transcription quantitative polymerase chain reaction (RT-qPCR)

cDNA was prepared using CellAmp Direct Lysis and RT set (Takara Bio, Shiga, Japan, Cat#
3737S/A) according to the manufacturer's instructions. Quantitative real-time PCR was
performed on a QuantStudio 7 Flex system (Thermo Fisher Scientific, Waltham, MA, USA,
RRID:SCR_020245) using TaqMan Universal Master Mix II with UNG (Thermo Fisher
Scientific, Waltham, MA, USA, Cat# 4440038) and Taqman probes: *Khk* (Cat#
Mm00434647_m1) and TATA-box binding protein (*Tbp*, Cat# Mm01277042_m1). Relative
gene expression of *Khk* was calculated using the $2^{-\Delta\Delta CT}$ method and normalized to *Tbp*.

6.12. Immunoblotting

The cells were lysed with 50 μL/well of RIPA buffer (50 mM Tris-HCl pH 7.4, 150 mM NaCl,
1% NP-40, 0.25% sodium deoxycholate, 1% SDS), and 1x complete protease inhibitor cocktail
(Sigma-Aldrich, St. Louis, MO, USA, Cat# 11697498001). The lysate was sonicated for 20 s
on ice and centrifuged at $15000 \times g$ at 4 °C for 10 min. The supernatant was collected and
separated on a 7.5% polyacrylamide-SDS gel and transferred to a PVDF membrane. The
membranes were blocked with 3% bovine serum albumin (BSA) (Sigma-Aldrich, St. Louis,
MO, USA, Cat# A7906) in Tris-Buffered Saline with Tween 20 (TBS-T) and incubated with
anti-KHK mouse monoclonal antibody (1:100) (Santa Cruz Biotechnology, Dallas, TX, USA,
Cat# sc-377411) in TBS-T supplemented with 3% BSA overnight at 4 °C. The membrane was
then incubated with HRP-conjugated anti-mouse immunoglobulins (1:2000) (Agilent, Santa
Clara, CA, USA, Cat# P0447, RRID:AB_2617137) in TBS-T supplemented with 1% BSA for
1 h at room temperature. Signals were visualized using ECL Prime detection reagent (Cytiva,
Buckinghamshire, UK, Cat# RPN2232). Images were taken using ImageQuant 800 (Cytiva,
Buckinghamshire, UK). Subsequently, the membrane was stripped using Restore Western Blot
Stripping Buffer (Thermo Fisher Scientific, Waltham, MA, USA, Cat# 21059) for 15 min and
re-probed with an anti-α-tubulin mouse monoclonal antibody (1:1000) (Thermo Fisher
Scientific, Waltham, MA, USA, Cat# A11126, RRID:AB_2534135). The images were
quantified using ImageJ (version 1.53k, https://imagej.nih.gov/ij/index.html,
RRID:SCR_003070).

6.13. Metabolic flux analysis

The cells were seeded in 6-well plates (Cat #353046; Corning, Glendale, AZ, USA) at 2×10^6
cells/2 mL/well and cultured for 48h. The cells were then rinsed three times with 2.8G-KRBH,
followed by pre-incubation with 2.8G-KRBH for 60 min at 37 °C. The supernatant was
replaced with KRBH supplemented with specified stimulations including [U-^{13}C]-glucose

(Sigma-Aldrich, St. Louis, MO, USA, CAS: 110187-42-3 Cat# 660663) and [U-^{13}C]-fructose

(Cambridge Isotope Laboratories, Tewksbury, MA, USA, CAS: 287100-63-4, Cat# CLM-

1553-1). The cells were incubated for 30 min at 37 °C and the supernatant was discarded. Cells

were quickly rinsed with ice-cold water, extracted by the addition of 500 μL ice-cold extraction

buffer (67.5% methanol, 7.5% chloroform, and 25% water), and the whole plate was snap-

frozen in liquid nitrogen. Cells were thawed on ice, scraped into 2 mL screw tubes along with

the supernatant, and stored at -80 °C until extraction of the metabolites.

For extraction of the metabolites, samples were supplemented with 80 μL of diluted (1:640)

internal standard (Human Metabolome Technologies, Yamagata, Japan, Cat# H3304-1002),

165 μL of methanol, and 465 μL of chloroform. The samples were then homogenized using a

pre-cooled bead crusher at 3200 rpm for 1 min and centrifuged at 15000 × g at 4 °C for 3 min.

The aqueous layer was transferred to pre-wetted ultrafiltration tubes (Human Metabolome

Technologies, Yamagata, Japan, Cat# UFC3LCCNB-HMT) and centrifuged at 9100 × g, 4 °C

until completely filtrated. The filtrate was freeze-dried, re-dissolved in 10 μL of water, and

subjected to mass spectrometry. The organic layer was evaporated by decompression at room

temperature and the residue was resuspended in lysis buffer (see Section 6.7). , which was then

subjected to a BCA protein assay (Thermo Fisher Scientific, Waltham, MA, USA, Cat# 23225).

cGMP was measured by G7100A capillary electrophoresis (Agilent Technologies, Santa

Clara, CA, USA) interfaced with a G6224A time-of-flight LC/MS mass spectrometer (Agilent

Technologies, Santa Clara, CA, USA). A G1310A isocratic pump (Agilent Technologies, Santa

Clara, CA, USA) equipped with a G1379B degasser (Agilent Technologies, Santa Clara, CA,

USA) was used to supply sheath liquid (Human Metabolome Technologies, Yamagata, Japan,

Cat# H3301-2020). The mass spectrometer was operated in the negative ionization mode. All

separations were performed on fused silica capillaries (Human Metabolome Technologies,

Yamagata, Japan, Cat# H3305-2002) at 25 °C using an anion analysis buffer (Human

Metabolome Technologies, Yamagata, Japan, Cat# H3302-2021) as the background electrolyte.

The applied voltage was set to 30 kV at 20 °C, together with a pressure of 15 mbar. Sheath

liquid was delivered to a nebulizer by an isocratic pump at 1 mL/min. Chromatograms and

mass spectra were analyzed by MassHunter qualitative analysis version 10.0 (Agilent

Technologies, Santa Clara, CA, USA, RRID:SCR_015040). Annotation and quantification of

chromatogram peaks were carried out using a standard mixture (Human Metabolome

Technologies, Yamagata, Japan, Cat# H3302-2021) as a reference.

6.14. Statistical Analysis

Sample sizes were estimated from the expected effect size based on previous experiments. No

randomization or blinding was used. For cell line experiments, including insulin secretion, IP1

measurement, RT-qPCR, immunoblotting, and metabolic tracing, *n* represents the number of

biological replicates of cells grown in different wells of the same multiwell plate. For islet

experiments, including insulin secretion and IP1 measurement, *n* represents the number of

wells, each containing five islets. For Ca^{2+} and ATP/ADP ratio measurements, *n* represents the

number of different single cells analyzed. Data are shown as the mean ± standard deviation

(SD) along with the plot of individual data points. For statistical comparisons between two

groups, a two-tailed unpaired Welch's unpaired *t*-test was used. For comparisons between three

groups, Welch's one-way analysis of variance (ANOVA) was followed by pairwise

comparisons corrected using Dunnett's method. Normality of the distribution was confirmed

by the Shapiro-Wilk test. P-values less than 0.05 were considered statistically significant and

are indicated in the figures. P-values more than 0.05 are not indicated in the figures. The

statistical analyses used are indicated in the figure legends. Statistical analyses were performed

using   GraphPad   Prism   10   (Graphpad   Software,   Boston,   MA,   USA,

https://www.graphpad.com; RRID:SCR_002798).

7   RESULTS

7.1 FSIS is dependent on K_{ATP} channels and calcium influx.

We first examined the responsiveness of B6 mouse islets and MIN6-K8 cells to various doses

of fructose. Fructose, at concentrations ranging from 2.5 mM to 12.5 mM and 0.3 mM to 10

mM, dose-dependently increased insulin secretion in B6 mouse islets (Figure 1A) and MIN6-

K8 cells (Figure 1B), respectively, when glucose levels were at a stimulatory level (8.8 mM).

However, fructose displayed no effect on insulin secretion at basal glucose level (2.8 mM) in

MIN6-K8 cells (Figure 1C), indicating that FSIS is glucose-dependent.

To explore whether intracellular fructose metabolism is coupled with insulin secretion in a

manner comparable to that of glucose, the involvement of K_{ATP} channels and intracellular Ca^{2+}

in FSIS was examined. We utilized *Kcnj11*[-/-] β-cells, which lack K_{ATP} channel activity and

exhibit continuous cell membrane depolarization regardless of glucose metabolism (Oduori et

al., 2020). Fructose did not significantly increase insulin secretion in *Kcnj11*[-/-] β-cells at 8.8

mM glucose (Figure 1B), suggesting that FSIS is dependent on K_{ATP} channel activity. This

finding is supported by the previous observation that FSIS was abolished by clamping the K_{ATP}

channel in the open state with diazoxide in MIN6-K8 cells (Seino et al., 2015).

In MIN6-K8 cells, fructose augmented the increase in intracellular Ca^{2+} ($[Ca^{2+}]i$) levels

induced by 8.8 mM glucose (Figure 1, D–E). In *Kcnj11*$^{-/-}$ β-cells, fructose did not increase

$[Ca^{2+}]i$ levels (Figure 1, F–G). These trends in $[Ca^{2+}]i$ responses were consistent with the

differential FSIS observed in Figure 1B. Furthermore, Nifedipine, an inhibitor of L-type

VDCCs, abolished FSIS in MIN6-K8 cells (Figure 1H). Overall, these findings indicate that

K_{ATP} channels and Ca^{2+} influx are necessary for FSIS.

Figure 1. FSIS is dependent on K_{ATP} channels and calcium influx.

(A) Dose-dependent effect of fructose on insulin secretion at 8.8 mM glucose in B6

mouse islets. n = 11–13.

(B) Dose-dependent effect of fructose on insulin secretion at 8.8 mM glucose in
parental MIN6-K8 and *Kcnj11*[-/-] cells. n = 4.

(C) Effect of fructose on insulin secretion at 2.8 mM glucose in MIN6-K8 cells. n = 4.

(D) –(G) Effect of fructose on intracellular Ca^{2+} levels measured using Fluo-4. The
time course of normalized fluorescence intensity at 508-579 nm is indicated in (D)
and (F). Black arrows represent the addition of the indicated stimulations at time
= 0. The magnitude of Ca^{2+} responses was quantified as iAUC in (E) and (G),
using F/F0 =1 as the baseline. (D)–(E): MIN6-K8 cells. 8.8 mM Glc: n = 14; 8.8
mM Glc + 10 mM Fru: n = 8. (F)–(G): *Kcnj11*[-/-] cells. 8.8 mM Glc: n = 12; 8.8 mM
Glc + 10 mM Fru: n = 11.

(H) Effect of nifedipine (10 µM) on FSIS in MIN6-K8 cells. n = 4.

Data are presented as the mean ± standard deviation (SD). Glc, glucose; Fru, fructose.
Statistical comparisons were performed using Welch's two-way ANOVA with
Dunnett's post-hoc test for (A) and (B), and Welch's unpaired two-tailed t-test for (E),
(G), and (H).

7.2 Characterization of β-cell fructose metabolism

According to our previously deposited RNA-seq datasets (Hashim et al., 2018), *Slc2a5*
(GLUT5) and *Khk* are expressed in both MIN6-K8 cells and B6 mouse islets (Supplementary
Table 1). To determine whether fructose is metabolized in β-cells, we traced the metabolic fate
of [13]C-labelled fructose. MIN6-K8 cells were stimulated for 30 min with 10 mM [U-13C]-
fructose, a stable isotopomer of fructose in which all six carbon atoms have been replaced with
[13]C, in the presence of 8.8 mM unlabeled glucose. For each intermediate, [13]C-labelled
isotopomers were quantified by capillary electrophoresis-time-of-flight mass spectrometry
(CE-TOFMS) (Figure 2). M–M6 represents the isotopomers in which 0–6 carbon atoms were
replaced with [13]C. Naturally occurring isotopomers are mostly M and M1, whereas
incorporation of [U-13C]-fructose-derived carbons generates M2–M6.

Upper glycolytic intermediates, such as G6P, F6P, and FBP, did not comprise M2–M6,
whereas F1P was found only as M6 under [U-13C]-fructose treatment. Lower glycolytic
intermediates such as 3PG, 2PG, PEP, glycerol 3P, and lactate were labelled as M3. Meanwhile,
TCA cycle intermediates such as citrate and malate were labelled as M2-M6. These results
indicate that [U-13C]-fructose enters lower glycolysis and the subsequent TCA cycle. We also
assessed labeling efficiency by [13]C fraction of the total carbon atoms in each intermediate

(Supplementary Figure 1A). ^{13}C Fraction was negligible at the basal state and increased up to

0.1–0.2 in the presence of [U-^{13}C]-fructose in glycolysis and TCA cycle intermediates.

Figure 2. [U-^{13}C]-fructose tracing in MIN6-K8 cells.

The total content and isotopic distribution of each intermediate are indicated by

stacked bar graphs alongside the schematic overview of metabolic fates of glucose

and [U-^{13}C]-fructose. n = 4. Data are presented as the mean ± standard deviation (SD).

Statistical comparisons were made between the total content using Welch's unpaired

two-tailed t-test.

Because FSIS was glucose-dependent (Figure 1, B–C), we sought to clarify whether fructose

affects glucose metabolism. To this end, MIN6-K8 cells were labelled with 10 mM unlabeled

fructose in the presence of 8.8 mM [U-^{13}C]-glucose (Figure 3A). Glucose-derived ^{13}C was

robustly incorporated into all intermediates, except for F1P. The upper and lower glycolytic

intermediates were labelled as M6 and M3, respectively. In those intermediates, the ^{13}C fraction

was 0.3–0.5, a value greater than that observed using [U-^{13}C]-fructose at a higher concentration

(10 mM), suggesting that the rate of fructose metabolism is lower than that of glucose in β-

cells (compare Figure 3B and Supplementary Figure 1A).

Notably, the content (total M–M6) and labeling of several intermediates were significantly

affected by fructose. The contents of G6P, 3PG, and 2PG were increased, while that of FBP

was decreased by fructose, a trend consistent regardless of the ^{13}C source (Figure 3A and Figure

2). Additionally, glucose-derived ^{13}C fraction was increased in FBP but was decreased in later

steps of glycolysis (Figure 3B). Previous studies have shown that high concentrations of

fructose inhibit phosphofructokinase (PFK) (Kreuzberg, 1978) and pyruvate kinase (PK)

(Mapungwana and Davies, 1982; Taylor et al., 2021). As both PFK and PK reactions are

irreversible (Chandel, 2021), it is reasonable to assume that the reduction in their activities

causes the accumulation of upstream intermediates and/or the depletion of downstream

intermediates. Therefore, our interpretation was that (1) increased content and labeling of G6P,

decreased content of FBP, and decreased labeling of intermediates after FBP (Figure 3, A–B)

reflect reduced PFK activity, and that (2) increased contents of 3PG and 2PG are due to reduced

PK activity. Overall, our metabolic tracing revealed interactions between fructose metabolism

and glycolysis, indicating that fructose enters lower glycolysis but simultaneously restricts

glycolysis by inhibiting PFK and PK.

Figure 3. [U-¹³C]-glucose tracing in MIN6-K8 cells.

(A) The total content and isotopic distribution of each intermediate are indicated as stacked bar graphs alongside a schematic overview of the metabolic fates of [U-¹³C]-glucose and fructose. n = 4. Statistical comparisons were made between the total content using Welch's unpaired two-tailed t-test.

(B) Fraction of ¹³C in each intermediate. Statistical comparisons were made using Welch's unpaired two-tailed t-test.

Data are presented as the mean ± standard deviation (SD).

7.3 β-cell fructose metabolism has little effect on insulin secretion

We next investigated the impact of intracellular fructose metabolism on cellular energy status in MIN6-K8 cells. Fructose did not significantly alter the levels of adenine nucleotides

(Supplementary Figure 1, B–D) or redox cofactors (NAD, NADH, NADP, and NADPH)

(Supplementary Figure 1, F–I). However, the ATP/ADP ratio tended to decrease

(Supplementary Figure 1E). Indeed, time-lapse imaging using PercevalHR, a genetically

encoded fluorescent biosensor (Tantama et al., 2013), revealed that fructose suppressed the

glucose-induced increase in the ATP/ADP ratio (Figure 4, A–B).

To clarify how intracellular fructose metabolism affects FSIS, we depleted KHK using

siRNA. KHK mRNA and protein levels were reduced by approximately 70% and 20%,

respectively (Supplementary Figure 2, A–C). Since the knockdown efficiency was moderate at

the protein level, we also validated the metabolic phenotypes using [U-^{13}C]-fructose tracing.

As expected, KHK knockdown decreased M6 F1P (Figure 4C). Although total content of other

intermediates remained unchanged (Supplementary Figure 3), ^{13}C fraction in later glycolysis

and TCA intermediates decreased (Figure 4D), confirming that KHK knockdown significantly

suppressed intracellular fructose metabolism. Knockdown of KHK resulted in a small decrease

in insulin secretion, regardless of the presence of fructose (Figure 4E). However, when

expressed as the fold-change over 0 mM fructose, KHK knockdown had no effect on fructose

responsiveness (Figure 4F). These results provide evidence against the hypothesis that β-cell

fructose metabolism fuels insulin secretion via ATP production.

Figure 4. Suppression of intracellular fructose metabolism does not affect
fructose responsiveness in MIN6-K8 cells.

(A) –(B) Effect of fructose on intracellular ATP/ADP ratio in MIN6-K8 cells as
visualized by Perceval-HR. 8.8 mM Glc: n = 16, 8.8 mM Glc + 10 mM Fru: n=15.
(A) Time course of normalized ratiometric signals (R/R0). (B) The magnitude of
the responses was quantified as iAUC using R/R0 = 1 as the baseline.

(E) –(D) [U-^{13}C]-fructose tracing under Khk knockdown in MIN6-K8 cells. The total
content and isotopic distribution of F1P are presented in (C). The fraction of ^{13}C
in each intermediate is presented in (D). n = 4. See Supplementary Figure 2 for
the total content of other metabolites.

(F) The effect of Khk knockdown on FSIS was expressed as fold change over 0 mM
fructose. See Supplementary Figure 3D for the original values.

All experiments were performed using MIN6-K8 cells. siNT, non-targeting siRNA. Data

are presented as the mean ± SD. Statistical comparisons were performed using

Welch's unpaired two-tailed t-test.

β-cell fructose responsiveness parallels PLC signaling

Given the limited role of fructose metabolism in FSIS, we investigated the molecular basis of

β-cell fructose responsiveness.

Fructose functions as a substrate for the sweet taste receptor (STR), a G protein-coupled

receptor complex composed of taste receptor type 1 member 2 (TAS1R2) and 3 (TAS1R3)

(Nelson et al., 2001; Zhao et al., 2003; Chandrashekar et al., 2006). STR signals through a

second messenger system that involves G-protein gustducin, phospholipase C β2 (PLCβ2),

inositol 1,4,5-trisphosphate (IP_3), and transient receptor potential M4/5 (TRPM4/5) (Huang et

al., 1999; Zhang et al., 2007). This pathway was originally implicated in taste transduction in

the tongue (Ahmad and Dalziel, 2020). Later studies have shown that STRs are expressed in

β-cells (Nakagawa et al., 2009) and regulate FSIS (Kyriazis et al., 2012). Indeed, U73122, an

inhibitor of PLCβ2 (Bleasdale et al., 1990), partially inhibited FSIS in MIN6-K8 cells (Figure

5A).

We sought to determine whether differential PLC activity can account for fructose

responsiveness in various β-cell models. To assess fructose-induced PLC activity, we measured

d-myo-inositol 1-phosphate (IP1) as a surrogate for IP_3. 0.5 mM and 3 mM fructose caused a

dose-dependent accumulation of IP1 in parental MIN6-K8 cells, but this increase was

diminished in *Kcnj11*[-/-] cells, suggesting that diminished PLC signaling underlies impaired

fructose responsiveness (Figure 5B).

The pathophysiological relevance of these findings was examined using diabetes mouse

models. NSY.B6-*Tyr*[+], *A*[y] is a spontaneous type 2 diabetes mouse strain that was recently

established by crossing the NSY and B6J- *A*[y] strains, thereby interrogating the obesity-related

agouti-yellow (*A*[y]) mutation in the agouti (*a*) gene into a diabetes-prone NSY background

(Ohno et al., 2022). Consistent with the observation by Ohno et al., Ay heterozygous male mice

(*A*[y]/*a*) displayed obesity and hyperglycemia after 13 weeks of age compared to wild-type male

mice (*a*/*a*) (Figure 5, C–D). Islets were isolated at 19–20 weeks of age and tested for fructose

responsiveness. Insulin secretion and IP1 levels in the a/a islets exhibited a significant increase

at 10 mM fructose (Figure 5, E–F). In contrast, insulin secretion and IP1 levels in *A*[y]/*a* islets

failed to respond to fructose. Hence, decreased fructose responsiveness in *Kcnj11*[-/-] cells and

diabetic islets correlated with impaired PLC activity. These findings suggest that β-cell fructose

responsiveness is not dictated by intracellular fructose metabolism, but rather by STR-mediated

PLC signaling.

Figure 5. Association between β-cell fructose responsiveness and PLC

signaling.

(A) Effect of U73122 (20 μM) on FSIS in MIN6-K8 cells. n = 4.

(B) Dose-dependent effects of fructose on intracellular IP1 levels in parental MIN6-K8
and *Kcnj11⁻/⁻* cells. The data are expressed as fold-change over 0 mM fructose. n
= 6. The two cell lines were compared for statistical significance. *$p < 0.05$.
(C) Age-dependent changes in body weight in NSY.B6 strains. The two strains were
compared for statistical significance. *a/a*: n = 6; *Aʸ/a*: n = 12. ****$p < 0.0001$.
(D) Age-dependent changes in ad-lib fed blood glucose levels in NSY.B6 strains. The
two strains were compared for statistical significance. *a/a*: n = 6; *Aʸ/a*: n = 12. *p
< 0.05, **$p < 0.01$, ***$p < 0.001$.
(E) FSIS in islets isolated from NSY.B6 strains. *a/a*: n = 9; *Aʸ/a*: n = 12.
(F) Effects of fructose on intracellular IP1 levels in islets isolated from NSY.B6 strains.
a/a: n = 10; *Aʸ/a*: n = 11.
Data are presented as the mean ± SD. Statistical comparisons were performed using
Welch's unpaired two-tailed t-test.
565
566 **7.4 PK activator counteracts fructose-mediated PK inhibition in β-cells**
567
Improving the responsiveness of β-cells to fructose is of therapeutic importance. We
hypothesized that neutralizing the inhibitory impact of fructose on glycolysis in β-cells would
allow for the effective utilization of fructose as a fuel source for insulin secretion.
It has recently been shown that F1P binds to and inhibits the M2 isoform of pyruvate kinase
(PKM2) in intestinal epithelial cells (Taylor et al., 2021). Notably, this inhibition can be
overcome by TEPP-46, a small-molecule PK activator (Jiang et al., 2010). Considering that
PKM2 (*Pkm*) is the primary PK isoform in MIN6-K8 cells and B6 mouse islets, as evidenced
by our previously deposited RNA-seq datasets (Supplementary Table 1) (Hashim et al., 2018),
this mechanism is also likely to be applicable to our study.
Motivated by these findings, we sought to determine whether TEPP-46 enhanced β-cell
fructose responsiveness. The effect of TEPP-46 on PK activity was evaluated using lysates
from MIN6-K8 cells. TEPP-46 significantly elevated PK activity, regardless of whether the
cells were pretreated with fructose (Figure 6A). However, fructose pretreatment caused only a
slight decrease in PK activity, implying that the binding of intracellular F1P to PK is disrupted
by cell lysis.
Therefore, to confirm TEPP-46's effect in situ, we conducted metabolic tracing experiments
using either [U-¹³C]-glucose or [U-¹³C]-fructose in the presence of TEPP-46. In both
experiments, fructose-induced accumulation of 2PG, 3PG, and PEP was substantially

attenuated by TEPP-46 (Figure 6, B–D, Supplementary Figures 4 and 5A), indicating effective

PK activation, as previously reported in hepatocytes (Abulizi et al., 2020).

In [U-^{13}C]-glucose tracing, ^{13}C fraction in 3PG and 2PG decreased, while that in later

intermediates (lactate, citrate, and malate) increased (Figure 6E). Similarly, in [U-^{13}C]-fructose

tracing, the ^{13}C fraction in citrate and malate slightly, but significantly, increased

(Supplementary Figure 5B). These results suggest that TEPP-46 facilitates the metabolic flux

of glucose and fructose after PK. Moreover, in the presence of TEPP-46, fructose did not alter

glucose-induced increases in the ATP/ADP ratio, as monitored by Perceval HR (Figure 6, F–

G). These results indicate that TEPP-46 was able to overcome fructose-mediated PK inhibition

and counteract fructose-induced ATP depletion.

Figure 6. PK activator counteracts fructose-mediated PK inhibition in β-cells

(A) Effect of TEPP-46 on PK activity in MIN6-K8 cell lysates. n = 3.

(B) –(E) [U-^{13}C]-glucose tracing after TEPP-46 treatment. The total content and isotopic distribution of the lower glycolytic intermediates are presented in (B)–(D). The fraction of ^{13}C in each intermediate is presented in (E). n = 4. See Supplementary Figure 4 for the total content and isotopic distribution of the other metabolites.

(F) –(G) Effect of fructose on intracellular ATP/ADP ratio in the presence of TEPP-46 as visualized by Perceval-HR. 8.8 mM Glc: n = 21; 8.8 mM Glc + 10 mM Fru: n =

606   19. (F) Time course of normalized ratiometric signals (R/R0). (G) The magnitude

607   of the responses was quantified as iAUC using R/R0 = 1 as the baseline.

All experiments were performed using MIN6-K8 cells. TEPP-46 was used at a

concentration of 10 µM. Data are presented as the mean ± SD. Statistical comparisons

were performed using Welch's one-way ANOVA with Dunnett's post-hoc test for (A)

and Welch's unpaired two-tailed t-test for (B) –(E).

7.5 PK activator enhances fructose responsiveness in β-cells

TEPP-46 had no effect on insulin secretion at 8.8 mM alone, but it significantly increased

insulin secretion when additional 10 mM fructose was present (Figure 7A). Consistently,

TEPP-46 did not significantly alter the Ca^{2+} response to glucose alone (Figure 7, B–C), but

augmented the increase in Ca^{2+} at 10 mM fructose (Figure 7, D–E). These findings indicate

that TEPP-46 requires fructose to exert insulinotropic effects. KHK knockdown markedly

suppressed fructose responsiveness in the presence of TEPP-46 (Figure 7F). Taken together,

these observations suggest that TEPP-46 enables intracellular fructose metabolism to

effectively fuel insulin secretion.

The effects of TEPP-46 were validated in both normal and diabetic islets. *a/a* mouse islets

failed to display significant FSIS to 3 mM fructose alone, but showed marked FSIS when

treated with TEPP-46 (Figure 7G). Similarly, A^y/a islets were responsive to 10 mM fructose in

the presence of TEPP-46 (Figure 7H). These results corroborate the effectiveness of PK

activation in enhancing fructose responsiveness in normal and diabetic β-cells.

Figure 7. PK activator enhances β-cell fructose responsiveness

(A) Effect of TEPP-46 on FSIS in MIN6-K8 cells. n = 4.

(B) (C) Effect of TEPP-46 on glucose-induced Ca^{2+} increase measured using Fluo-4. DMSO: n = 13; TEPP-46: n = 24. (B) The black arrow represents the addition of 8.8 mM glucose (8.8G) at time 0. (C) The magnitude of the responses was quantified as iAUC using F/F0 = 1 as the baseline.

(D) –(E) Effect of TEPP-46 on fructose-induced Ca^{2+} increase measured using Fluo-4. DMSO: n = 25; TEPP-46: n = 24. (D) The black arrow represents the addition of 8.8 mM glucose supplemented with 10 mM fructose (8.8G + 10F) at time 0. (E)

The magnitude of the responses was quantified as iAUC using F/F0 = 1 as the

baseline.

(F) Effect of Khk knockdown on FSIS in the presence of TEPP-46 in MIN6-K8 cells.

siNT, non-targeting siRNA. n = 4.

(G) Effect of TEPP-46 on FSIS in *a*/*a* mouse islets. n = 8.

(H) Effect of TEPP-46 on FSIS in *A^y*/*a* mouse islets. n = 6.

Experiments (A)–(F) were performed using MIN6-K8 cells. TEPP-46 was used at a

concentration of 10 µM. Data are presented as the mean ± SD. Statistical comparisons

were performed using Welch's unpaired two-tailed t-test for (A) –(E) and Welch's one-

way ANOVA with Dunnett's post-hoc test for (G) –(H).

7   DISCUSSION

The present study demonstrated that (1) the responsiveness of β-cells to fructose is not

regulated by intracellular fructose metabolism, but rather by PLC signaling; (2) fructose limits

glycolysis by inhibiting PK and decreasing the ATP/ADP ratio; and (3) a PK activator

overcomes the inhibition of PK caused by F1P, thereby enhancing β-cell responsiveness to

fructose. These findings highlight the distinction between FSIS and GSIS, most of which is

attributable to the unique metabolic property of fructose.

Fructose has been known to be metabolized in β-cells (Ashcroft et al., 1972; Zawalich et al.,

1977; Sener and Malaisse, 1988; Giroix et al., 1999). However, the extent to which β-cell

fructose metabolism contributes to insulin secretion has remained undetermined. In this study,

by employing ^{13}C-labelled metabolic tracing and Khk knockdown experiments, we sought to

revisit this question.

In accordance with previous studies (Ashcroft et al., 1970; Zawalich et al., 1977; Sener and

Malaisse, 1988; Giroix et al., 1999), we found that the rate of fructose metabolism was

significantly lower than that of glucose metabolism in β-cells. Nevertheless, some studies have

proposed that FSIS may be regulated by ATP produced through intracellular fructose

metabolism, as the rates of insulin secretion and fructose metabolism or ATP production have

been observed to be correlated (Zawalich et al., 1977; Sener and Malaisse, 1988). However,

these findings have been challenged by later studies using diabetic GK rats (Giroix et al., 1999)

and an INS-1E β-cell line (Bartley et al., 2019), which showed no impact of acute fructose on

mitochondrial respiration or ATP production, as well as by our observation that fructose

impedes the glucose-induced increase in the ATP/ADP ratio.

The relationship between intracellular fructose metabolism and ATP levels is complex and
involves multiple factors. This study identified two factors that likely contribute to fructose-
induced reduction in the ATP/ADP ratio: ATP consumption caused by KHK and fructose-
mediated inhibition of PFK and PK. Phosphorylation of fructose by KHK requires ATP
hydrolysis, which leads to a reduction in ATP levels in the liver and kidney cells (Van den
Berghe et al., 1977; Abdelmalek et al., 2012; Cirillo et al., 2009). Chronic fructose treatment
has also been shown to decrease ATP levels in INS-1E cells (Bartley et al., 2019).

In MIN6-K8 cells, M6 F1P accumulated substantially after [U-^{13}C]-fructose treatment
(approximately 40 times more than G6P), but ^{13}C incorporation in downstream intermediates
was relatively low. This might be due to the imbalance between the expression levels of
GLUT5 and KHK and that of aldorase B, which was very low (Supplementary Table 1). Hence,
we assume that ATP consumption by KHK cannot be balanced by subsequent ATP production
resulting from fructose catabolism, leading to a decrease in the ATP/ADP ratio. Consequently,
fructose catabolism and cellular energy status were not positively correlated.

Previous research has utilized radioisotope-labelled sugars to measure the utilization or
oxidation rates of the sugar in the entire islet (Ashcroft et al., 1970; Zawalich et al., 1977; Sener
and Malaisse, 1988; Giroix et al., 1999), reporting that glucose utilization rate was unaffected
by the addition of excess fructose (Zawalich et al., 1977). In contrast, the present study
observed an apparent decrease in glycolytic flux by fructose. These discrepant results may
imply that the whole-cell glucose utilization rate is largely determined by glucokinase activity
and does not necessarily reflect fluctuations in downstream steps. In this regard, the utilization
of mass spectrometry with ^{13}C-labelled sugars, which enabled direct tracing of carbon flux in
each intermediate, offered a significant advantage in revealing the interactions between
glycolysis and fructolysis.

In the liver, fructose promotes glucose uptake and phosphorylation by stimulating the release
of glucokinase (GCK) from glucokinase regulatory protein (GCKR) (Brown et al., 1997;
McGuinness and Cherrington, 2003). While the same mechanism may apply to β-cells, it is
less likely given the low expression of GCKR in MIN6-K8 cells (Supplementary Table 1).
Instead, the observed increase in G6P due to fructose was likely a result of diminished PFK
activity. Some studies have reported that fructose is converted to F6P and G6P, which are then
oxidized through the pentose phosphate pathway (Sener and Malaisse, 1988; Giroix et al.,
1999). However, our findings indicated that [U-^{13}C]-fructose was not converted to M6 F6P or
M6 G6P.   It is possible that differences in experimental conditions, such as sugar

concentrations and the species of animals tested, as well as methodological differences,
contributed to this discrepancy.

This study provides the first description of fructose-mediated PK inhibition in β-cells. The
effect of F1P on PK may be dependent on concentrations and cell types, because it is also
reported that liver PK is activated by F1P in liver homogenates (Eggleston and Woods, 1970).
Nevertheless, our results clearly demonstrated the inhibition of PKM2 by fructose in β-cells,
according to the significant accumulation of 2PG and 3PG, which was ameliorated by TEPP-
46.

Although TEPP-46 counteracts fructose-mediated PK inhibition, the reason why the
insulinotropic effect of TEPP-46 requires intracellular fructose metabolism remains unclear.
There was no evidence that ATP production from fructose directly stimulated FSIS, even when
TEPP-46 was present, as the whole-cell ATP/ADP ratio visualized by Perceval HR was similar
under fructose- and glucose-stimulated conditions. However, our results do not preclude the
possibility that local ATP production by PK near the K_{ATP} channel regulates FSIS. Recent
studies have proposed that membrane-associated PK directly provides ATP to β-cell K_{ATP}
channels and controls glucose-induced channel closure (Lewandowski et al., 2020; Foster et
al., 2022; Ho et al. 2023; Merrins and Kibbey, 2024), although this proposal remains
controversial (Corradi et al., 2023; Satin et al., 2024; Rutter and Sweet, 2024). A similar
mechanism may also participate in FSIS.

Furthermore, lipogenesis-related metabolites, such as diacylglycerol, free fatty acid, and
fatty acid CoA, are potential metabolic coupling factors (MCFs) in FSIS (Prentki et al., 2013).
Fructose-derived metabolites are preferentially used for de novo lipid synthesis (Herman and
Samuel, 2016; Silva et al., 2019). As our metabolic analysis did not cover these pathways, more
comprehensive metabolic profiling is required to elucidate the link between intracellular
fructose metabolism and FSIS.

Excessive fructose consumption has been linked to several health issues, including obesity,
metabolic dysfunction-associated steatotic liver disease (MASLD), and type 2 diabetes
(Hannou et al., 2018; Taskinen et al., 2019). It has been demonstrated that FSIS is impaired in
the islets of NSY.B6-Ay/a, a mouse model of obese diabetes, suggesting that the deleterious
effects of fructose on glucose tolerance are counterbalanced by increased insulin secretion.
Surprisingly, these diabetic islets became responsive to fructose when treated with TEPP-46.
Furthermore, a recent study showed that pyruvate kinase activators improve both insulin
sensitivity and insulin secretion in high-fat diet-fed mice in vivo (Abulizi et al. 2020). Given

these findings, additional research is necessary to evaluate TEPP-46's potential as a therapeutic

agent for fructose-induced metabolic disorders.

8   LIMITATIONS OF THE STUDY

In this study, we utilized fructose concentrations that are in line with recent investigations

(Kyriazis et al., 2012; Bartley et al., 2019), but are not confirmed to be physiologically relevant.

We recognize that insulin secretory response to ingested fructose is determined by blood

fructose levels in vivo. However, there are considerable variations in the reported blood

fructose levels, ranging from sub-millimolar values (Sugimoto et al., 2010; Patel et al., 2015)

to more than 10 millimolar (Hui et al., 2009), which is likely due to differences in the subjects

and quantification methods. Future studies should aim to establish dose-response relationships

for fructose-induced insulin secretion in vivo to better understand the physiological

implications of our findings.

9   AUTHOR CONTRIBUTIONS

Conceptualization, N.M.; Methodology, N.M..; Investigation, N.M. and R.M.; Resources –

T.O., and N.Y.; Writing – Original Draft: N.M.; Writing – Review & Editing: N.M., R.M.,

Y.S., K.S.., Y.M., T.O.., N.Y.., Y.Y., and A.S.; Data Curation: N.M.; Visualization: N.M.;

Supervision: K.S., Y.Y., and A.S.; Funding Acquisition: N.M. and A.S.

10   ACKNOWLEDGEMENTS

The authors extend their gratitude to President Yutaka Seino of Kansai Electric Power Hospital

for his general support in this research. The authors thank Yoshikazu Hoshino of Hoshino

Laboratory Animals, Inc. for his assistance with the NSY.B6 mice. The authors thank Shihomi

Sakai, Asami Yamaguchi, and Megumi Akiyama for their excellent technical assistance.

11   DATA AVAILABILITY

The RNA sequencing data used for Supplementary Table 1 are available from the DDBJ

Sequence Read Archive with the accession number DRA006332 (Hashim et al., 2018). Other

data supporting the findings of this study are available from the corresponding author upon

reasonable request.

12 FUNDING

776 This study was supported by JSPS KAKENHI Grant Numbers JP22K20869 and JP23K15401

777 for N.M. Research grants for N.M. were provided by the Japan Association for Diabetes

778 Education and Care, Daiwa Securities Foundation, Suzuken Memorial Foundation, Japan

779 Diabetes Foundation - Nippon Boehringer Ingelheim Co., Ltd., The Hori Sciences and Arts

780 Foundation, Manpei Suzuki Diabetes Foundation, and Fujita Health University.

13 CONFLICT OF INTEREST

784 The authors declare no conflict of interest.

14 REFERENCES

[1] Abdelmalek, M. F., Lazo, M., Horska, A., Bonekamp, S., Lipkin, E. W.,
Balasubramanyam, A., Bantle, J. P., Johnson, R. J., Diehl, A. M., and Clark, J. M. (2012)
Higher dietary fructose is associated with impaired hepatic adenosine triphosphate
homeostasis in obese individuals with type 2 diabetes. Hepatology 56, 952-960

[2] Abulizi, A., Cardone, R. L., Stark, R., Lewandowski, S. L., Zhao, X., Hillion, J., Ma, L.,
Sehgal, R., Alves, T. C., and Thomas, C. (2020) Multi-tissue acceleration of the
mitochondrial phosphoenolpyruvate cycle improves whole-body metabolic health. Cell
metabolism 32, 751-766. e711

[3] Ahmad, R., and Dalziel, J. E. (2020) G protein-coupled receptors in taste physiology and
pharmacology. Frontiers in pharmacology 11, 587664

[4] Ashcroft, S. J., Bassett, J. M., and Randle, P. J. (1972) Insulin secretion mechanisms and
glucose metabolism in isolated islets. Diabetes 21, 538-545

[5] Bartley, C., Brun, T., Oberhauser, L., Grimaldi, M., Molica, F., Kwak, B. R., Bosco, D.,
Chanson, M., and Maechler, P. (2019) Chronic fructose renders pancreatic β-cells hyper-
responsive to glucose-stimulated insulin secretion through extracellular ATP signaling.
American Journal of Physiology-Endocrinology and Metabolism 317, E25-E41

[6] Bleasdale, J. E., Thakur, N. R., Gremban, R. S., Bundy, G. L., Fitzpatrick, F. A., Smith,
R. J., and Bunting, S. (1990) Selective inhibition of receptor-coupled phospholipase C-
dependent processes in human platelets and polymorphonuclear neutrophils. Journal of
Pharmacology and Experimental Therapeutics 255, 756-768

[7] Brown, K. S., Kalinowski, S. S., Megill, J. R., Durham, S. K., and Mookhtiar, K. A. (1997)
Glucokinase regulatory protein may interact with glucokinase in the hepatocyte nucleus.
Diabetes 46, 179-186

[8] Chandel, N. S. (2021) Glycolysis. Cold Spring Harbor Perspectives in Biology 13,
a040535

[9] Chandrashekar, J., Hoon, M. A., Ryba, N. J., and Zuker, C. S. (2006) The receptors and
cells for mammalian taste. Nature 444, 288-294

[10] Cirillo, P., Gersch, M. S., Mu, W., Scherer, P. M., Kim, K. M., Gesualdo, L., Henderson,
G. N., Johnson, R. J., and Sautin, Y. Y. (2009) Ketohexokinase-dependent metabolism of
fructose induces proinflammatory mediators in proximal tubular cells. Journal of the
American Society of Nephrology 20, 545-553

[11] Corradi, J., Thompson, B., Fletcher, P. A., Bertram, R., Sherman, A. S., and Satin, L. S.
(2023) KATP channel activity and slow oscillations in pancreatic beta cells are regulated
by mitochondrial ATP production. The Journal of Physiology 601, 5655-5667

[12] Eggleston, L., and Woods, H. (1970) Activation of liver pyruvate kinase by fructose-1-
phosphate. FEBS letters 6, 43-45

[13] Foster, H. R., Ho, T., Potapenko, E., Sdao, S. M., Huang, S. M., Lewandowski, S. L.,
VanDeusen, H. R., Davidson, S. M., Cardone, R. L., and Prentki, M. (2022) β-cell deletion
of the PKm1 and PKm2 isoforms of pyruvate kinase in mice reveals their essential role as
nutrient sensors for the KATP channel. Elife 11, e79422

[14] Giroix, M.-H., Scruel, O., Ladriere, L., Sener, A., Portha, B., and Malaisse, W. J. (1999)
Metabolic and secretory interactions between D-glucose and D-fructose in islets from GK
rats. Endocrinology 140, 5556-5565

[15] Grant, A., Christie, M., and Ashcroft, S. (1980) Insulin release from human pancreatic
islets in vitro. Diabetologia 19, 114-117

[16] Grodsky, G. M., Batts, A. A., Bennett, L. L., Vcella, C., McWilliams, N. B., and Smith,
D. F. (1963) Effects of carbohydrates on secretion of insulin from isolated rat pancreas.
American Journal of Physiology-Legacy Content 205, 638-644

[17] Han, G., Takahashi, H., Murao, N., Gheni, G., Yokoi, N., Hamamoto, Y., Asahara, S. i.,
Seino, Y., Kido, Y., and Seino, S. (2021) Glutamate is an essential mediator in glutamine-
amplified insulin secretion. Journal of Diabetes Investigation 12, 920-930

[18] Hannou, S. A., Haslam, D. E., McKeown, N. M., and Herman, M. A. (2018) Fructose
metabolism and metabolic disease. The Journal of clinical investigation 128, 545-555

[19] Hashim, M., Yokoi, N., Takahashi, H., Gheni, G., Okechi, O. S., Hayami, T., Murao, N.,
Hidaka, S., Minami, K., and Mizoguchi, A. (2018) Inhibition of SNAT5 induces incretin-
responsive state from incretin-unresponsive state in pancreatic β-cells: study of β-cell
spheroid clusters as a model. Diabetes 67, 1795-1806

[20] Henquin, J.-C. (2000) Triggering and amplifying pathways of regulation of insulin
secretion by glucose. Diabetes 49, 1751-1760

[21] Henquin, J.-C. (2009) Regulation of insulin secretion: a matter of phase control and
amplitude modulation. Diabetologia 52, 739-751

[22] Herman, M. A., and Samuel, V. T. (2016) The sweet path to metabolic demise: fructose
and lipid synthesis. Trends in Endocrinology & Metabolism 27, 719-730

[23] Ho, T., Potapenko, E., Davis, D. B., and Merrins, M. J. (2023) A plasma membrane-
associated glycolytic metabolon is functionally coupled to KATP channels in pancreatic
α and β cells from humans and mice. Cell reports 42

[24] Huang, L., Shanker, Y. G., Dubauskaite, J., Zheng, J. Z., Yan, W., Rosenzweig, S.,
Spielman, A. I., Max, M., and Margolskee, R. F. (1999) Gγ13 colocalizes with gustducin
in taste receptor cells and mediates IP3 responses to bitter denatonium. Nature
neuroscience 2, 1055-1062

[25] Hui, H., Huang, D., McArthur, D., Nissen, N., Boros, L. G., and Heaney, A. P. (2009)
Direct spectrophotometric determination of serum fructose in pancreatic cancer patients.
Pancreas 38, 706-712

[26] Iwasaki, M., Minami, K., Shibasaki, T., Miki, T., Miyazaki, J. i., and Seino, S. (2010)
Establishment of new clonal pancreatic β-cell lines (MIN6-K) useful for study of
incretin/cyclic adenosine monophosphate signaling. Journal of Diabetes Investigation 1,
137-142

[27] Jiang, J.-k., Boxer, M. B., Vander Heiden, M. G., Shen, M., Skoumbourdis, A. P., Southall,
N., Veith, H., Leister, W., Austin, C. P., and Park, H. W. (2010) Evaluation of thieno [3,
2-b] pyrrole [3, 2-d] pyridazinones as activators of the tumor cell specific M2 isoform of
pyruvate kinase. Bioorganic & medicinal chemistry letters 20, 3387-3393

[28] Kagiyama, N., Ikeda, T., and Nomura, T. (2006) Japanese Guidelines and regulations for
scientific and ethical animal experimentation. In In Vivo Models of Inflammation: Volume
I pp. 187-191, Springer

[29] Kreuzberg, K. (1978) Interaction of D-fructose and fructose 1-phosphate with yeast
phosphofructokinase and its influence on glycolytic oscillations. Biochimica et
Biophysica Acta (BBA)-Enzymology 527, 229-238

[30] Kyriazis, G. A., Soundarapandian, M. M., and Tyrberg, B. (2012) Sweet taste receptor
signaling in beta cells mediates fructose-induced potentiation of glucose-stimulated
insulin secretion. Proceedings of the National Academy of Sciences 109, E524-E532

[31] Larke, J. A., Bacalzo, N., Castillo, J. J., Couture, G., Chen, Y., Xue, Z., Alkan, Z., Kable,
M. E., Lebrilla, C. B., and Stephensen, C. B. (2023) Dietary intake of monosaccharides
from foods is associated with characteristics of the gut microbiota and gastrointestinal
inflammation in healthy US adults. The Journal of Nutrition 153, 106-119

[32] Lewandowski, S. L., Cardone, R. L., Foster, H. R., Ho, T., Potapenko, E., Poudel, C.,
VanDeusen, H. R., Sdao, S. M., Alves, T. C., and Zhao, X. (2020) Pyruvate kinase controls
signal strength in the insulin secretory pathway. Cell metabolism 32, 736-750. e735

[33] Mapungwana, S. M., and Davies, D. R. (1982) The effect of fructose on pyruvate kinase
activity in isolated hepatocytes. Inhibition by allantoin and alanine. Biochemical Journal
208, 171-178

[34] McGuinness, O. P., and Cherrington, A. D. (2003) Effects of fructose on hepatic glucose
metabolism. Current Opinion in Clinical Nutrition & Metabolic Care 6, 441-448

[35] Merrins, M. J., and Kibbey, R. G. (2024) Glucose Regulation of β-Cell KATP Channels:
It Is Time for a New Model! Diabetes 73, 856-863

[36] Murao, N., Morikawa, R., Seino, Y., Shimomura, K., Maejima, Y., Yamada, Y., and
Suzuki, A. (2024) Sildenafil amplifies calcium influx and insulin secretion in pancreatic β
cells. Physiological Reports 12, e16091

[37] Murao, N., Yokoi, N., Honda, K., Han, G., Hayami, T., Gheni, G., Takahashi, H., Minami,
K., and Seino, S. (2017) Essential roles of aspartate aminotransferase 1 and vesicular

glutamate transporters in β-cell glutamate signaling for incretin-induced insulin secretion.
PLoS One 12, e0187213

[38] Murao, N., Yokoi, N., Takahashi, H., Hayami, T., Minami, Y., and Seino, S. (2022)
Increased glycolysis affects β-cell function and identity in aging and diabetes. Molecular
metabolism 55, 101414

[39] Nakagawa, Y., Nagasawa, M., Yamada, S., Hara, A., Mogami, H., Nikolaev, V. O., Lohse,
M. J., Shigemura, N., Ninomiya, Y., and Kojima, I. (2009) Sweet taste receptor expressed
in pancreatic β-cells activates the calcium and cyclic AMP signaling systems and
stimulates insulin secretion. PloS one 4, e5106

[40] Nelson, G., Hoon, M. A., Chandrashekar, J., Zhang, Y., Ryba, N. J., and Zuker, C. S.
(2001) Mammalian sweet taste receptors. Cell 106, 381-390

[41] Oduori, O. S., Murao, N., Shimomura, K., Takahashi, H., Zhang, Q., Dou, H., Sakai, S.,
Minami, K., Chanclon, B., and Guida, C. (2020) Gs/Gq signaling switch in β cells defines
incretin effectiveness in diabetes. The Journal of clinical investigation 130, 6639-6655

[42] Patel, C., Sugimoto, K., Douard, V., Shah, A., Inui, H., Yamanouchi, T., and Ferraris, R.
P. (2015) Effect of dietary fructose on portal and systemic serum fructose levels in rats
and in KHK−/− and GLUT5−/− mice. American Journal of Physiology-Gastrointestinal
and Liver Physiology 309, G779-G790

[43] Prentki, M., Matschinsky, F. M., and Madiraju, S. M. (2013) Metabolic signaling in fuel-
induced insulin secretion. Cell metabolism 18, 162-185

[44] Rutter, G. A., and Sweet, I. R. (2024) Glucose Regulation of β-Cell KATP Channels: Is a
New Model Needed? Diabetes 73, 849-855

[45] Satin, L. S., Corradi, J., and Sherman, A. S. (2024) Do We Need a New Hypothesis for
KATP Closure in β-Cells? Distinguishing the Baby From the Bathwater. Diabetes 73, 844-
848

[46] Seino, Y., Ogata, H., Maekawa, R., Izumoto, T., Iida, A., Harada, N., Miki, T., Seino, S.,
Inagaki, N., Tsunekawa, S., Oiso, Y., and Hamada, Y. (2015) Fructose induces glucose-
dependent insulinotropic polypeptide, glucagon-like peptide-1 and insulin secretion: Role
of adenosine triphosphate-sensitive K+ channels. Journal of Diabetes Investigation 6, 522-
526

[47] Sener, A., and Malaisse, W. J. (1988) Hexose metabolism in pancreatic islets: Metabolic
and secretory responses to d-fructose. Archives of biochemistry and biophysics 261, 16-
26

[48] Silva, J. C., Marques, C., Martins, F. O., Viegas, I., Tavares, L., Macedo, M. P., and Jones,
J. G. (2019) Determining contributions of exogenous glucose and fructose to de novo fatty
acid and glycerol synthesis in liver and adipose tissue. Metabolic engineering 56, 69-76

[49] Sugimoto, K., Hosotani, T., Kawasaki, T., Nakagawa, K., Hayashi, S., Nakano, Y., Inui,
H., and Yamanouchi, T. (2010) Eucalyptus leaf extract suppresses the postprandial
elevation of portal, cardiac and peripheral fructose concentrations after sucrose ingestion
in rats. Journal of clinical biochemistry and nutrition 46, 205-211

[50] Tantama, M., Martínez-François, J. R., Mongeon, R., and Yellen, G. (2013) Imaging
energy status in live cells with a fluorescent biosensor of the intracellular ATP-to-ADP
ratio. Nature communications 4, 2550

[51] Taskinen, M.-R., Packard, C. J., and Borén, J. (2019) Dietary fructose and the metabolic
syndrome. Nutrients 11, 1987

[52] Taylor, S. R., Ramsamooj, S., Liang, R. J., Katti, A., Pozovskiy, R., Vasan, N., Hwang,
S.-K., Nahiyaan, N., Francoeur, N. J., and Schatoff, E. M. (2021) Dietary fructose
improves intestinal cell survival and nutrient absorption. Nature 597, 263-267

[53] Van Den Berghe, G., Bronfman, M., Vanneste, R., and Hers, H. (1977) The mechanism
of adenosine triphosphate depletion in the liver after a load of fructose. A kinetic study of
liver adenylate deaminase. Biochemical Journal 162, 601-609

[54] Zawalich, W. S., Rognstad, R., Pagliara, A. S., and Matschinsky, F. M. (1977) A
comparison of the utilization rates and hormone-releasing actions of glucose, mannose,
and fructose in isolated pancreatic islets. J Biol Chem 252, 8519-8523

[55] Zhang, Z., Zhao, Z., Margolskee, R., and Liman, E. (2007) The transduction channel
TRPM5 is gated by intracellular calcium in taste cells. Journal of Neuroscience 27, 5777-
5786

[56] Zhao, G. Q., Zhang, Y., Hoon, M. A., Chandrashekar, J., Erlenbach, I., Ryba, N. J., and
Zuker, C. S. (2003) The receptors for mammalian sweet and umami taste. Cell 115, 255-
266

SUPPLEMENTARY MATERIALS

Supplementary Figure 1.

[U-[13]C]-fructose tracing in MIN6-K8 cells.

(A) Fraction of [13]C in each intermediate.

(B) –(I) Total metabolite content and their ratio.

All experiments were performed using MIN6-K8 cells. n = 4. Data are presented as the mean

± SD. Statistical comparisons were made using Welch's unpaired two-tailed t-test. See also

Figure 2 for the experimental details.

Supplementary Figure 2.

(A) Knockdown efficiency of *Khk* was assessed by RT-qPCR. mRNA levels were normalized to those in siNT (non-targeting siRNA)-treated cells. n = 4.

(B) –(C)Knockdown efficiency of KHK protein assessed by immunoblotting. (B) The membrane was first probed for KHK and subsequently re-probed for α-tubulin. The corresponding bands are indicated by black triangles. (C) The average intensity of KHK was normalized to that of α-tubulin and expressed as fold-change. n = 6.

All experiments were performed using MIN6-K8 cells. Data are presented as the mean ± SD. Statistical comparisons were made using Welch's unpaired two-tailed t-test.

Supplementary Figure 3.

Effect of Khk knockdown on [U-^{13}C]-fructose tracing in MIN6-K8 cells. The total content and isotopic distribution of each intermediate are presented as stacked bar graphs along with a schematic overview of the metabolic fates of glucose and [U-^{13}C]-fructose. n = 4. Statistical comparisons were made between the total content using Welch's unpaired two-tailed t-test.

986
Supplementary Figure 4.

988 Effect of TEPP-46 on [U-^{13}C]-glucose tracing in MIN6-K8 cells. The total content and isotopic

989 distribution of each intermediate are presented as stacked bar graphs, along with a schematic

990 overview of the metabolic fates of [U-^{13}C]-glucose and fructose. n = 4. Statistical comparisons

991 were made between the total content using Welch's unpaired two-tailed t-test.

Supplementary Figure 5.

Effect of TEPP-46 on [U-^{13}C]-fructose tracing in MIN6-K8 cells. n = 4.

(A) The total content and isotopic distribution in each intermediate are presented as stacked

bar graphs. Statistical comparisons were made between the total content.

(B) Fraction of ^{13}C in each intermediate.

Statistical comparisons were made using Welch's unpaired two-tailed t-test.

Supplementary Table 1

		FPKM					
		MIN6-K8			B6 islet		
GO:0006001 fructose catabolic process							
Aldh1a1	aldehyde dehydrogenase family 1, subfamily A1	0.00	0.00	0.00	2.12	1.13	5.87
Aldh1a7	aldehyde dehydrogenase family 1, subfamily A7	0.00	0.00	0.00	0.00	0.00	0.00
Aldob	aldolase B, fructose-bisphosphate	0.00	1.00	0.00	0.00	0.00	0.98
Glyctk	glycerate kinase	10.80	7.02	15.23	9.20	21.50	13.70
Khk	ketohexokinase	17.67	29.10	21.76	50.96	58.85	43.06
Tkfc	Triokinase/FMN cyclase	77.55	76.25	58.75	58.75	64.51	73.40
GO:0006007 glucose catabolic process							
Actn3	actinin alpha 3	0.98	1.00	3.26	2.12	3.40	1.96
Adcy10	adenylate cyclase 10	0.00	0.00	0.00	0.71	0.00	0.00
Aldoa	aldolase A, fructose-bisphosphate	1014.98	1010.31	1024.87	904.87	932.85	947.52
Bad	BCL2-associated agonist of cell death	22.58	35.12	44.61	40.34	37.35	46.00
Eno1	enolase 1, alpha non-neuron	61.84	62.20	54.40	55.91	41.87	46.97
Eno1b	eno1b enolase 1B, retrotransposed	0.00	0.00	0.00	0.00	0.00	0.00
Eno2	gamma-enolase isoform 2	171.78	152.50	147.96	150.76	122.23	128.20
Eno3	enolase 3, beta muscle	4.91	7.02	15.23	37.51	33.95	40.12
Foxk1	forkhead box K1	91.29	112.37	130.56	132.36	117.70	115.48
Foxk2	forkhead box K2	353.38	326.07	332.92	190.39	162.97	174.20
Gapdh	glyceraldehyde-3-phosphate dehydrogenase isoform 1	5.89	5.02	10.88	9.91	18.11	17.62
Gck	glucokinase isoform 2	112.88	137.45	141.44	292.31	291.99	112.88
Gpi1	glucose phosphate isomerase 1	245.40	229.75	242.62	263.29	264.83	282.83
Hk1	hexokinase 1	0.00	0.00	2.18	21.94	21.50	28.38
Ldha	lactate dehydrogenase A	0.98	3.01	2.18	27.60	30.56	24.47
Lrp5	low density lipoprotein receptor-related protein 5	25.52	28.09	20.67	105.46	95.07	118.42
Pfkfb2	6-phosphofructo-2-kinase/fructose-2,6-biphosphatase 2	317.06	367.20	349.24	369.12	331.21	322.54
Pfkl	phosphofructokinase, liver, B-type	228.71	224.74	213.24	242.06	222.95	238.79
Pfkm	phosphofructokinase, muscle	218.90	204.67	211.07	213.04	224.09	242.70
Pfkp	ATP-dependent 6-phosphofructokinase, platelet type isoform 2	175.71	167.55	166.46	134.48	113.18	159.52
Pgam2	phosphoglycerate mutase 2	0.00	0.00	0.00	0.00	0.00	0.00
Pgk1	phosphoglycerate kinase 1	57.91	74.24	64.19	39.64	40.74	35.23
Pkm	pyruvate kinase, muscle	1746.27	1669.47	1697.23	1573.39	1491.64	1533.53
Slc25a12	solute carrier family 25 (mitochondrial carrier, Aralar), member 12	214.97	251.83	221.95	186.15	202.58	194.75
Tigar	Trp53 induced glycolysis regulatory phosphatase	182.58	167.55	181.69	111.12	149.39	108.63
Tpi1	triosephosphate isomerase 1	565.40	565.85	562.48	392.82	374.61	407.12
Trp53	transformation related protein 53	75.58	59.19	60.93	43.17	35.08	46.00
GO:0015149 hexose transmembrane transporter activity							
Gm5134	predicted gene 5134	nd	nd	nd	nd	nd	nd
Mfsd4a	#N/A	nd	nd	nd	nd	nd	nd
Mfsd4b1	#N/A	nd	nd	nd	nd	nd	nd
Mfsd4b4	#N/A	nd	nd	nd	nd	nd	nd
Mfsd4b5	#N/A	nd	nd	nd	nd	nd	nd
Slc2a1	solute carrier family 2 (facilitated glucose transporter), member 1	55.95	57.19	53.31	42.47	39.61	51.87
Slc2a2	solute carrier family 2 (facilitated glucose transporter), member 2	330.80	325.07	352.50	1665.40	1972.63	1619.65
Slc2a3	solute carrier family 2 (facilitated glucose transporter), member 3	1.96	0.00	0.00	8.49	14.71	8.81
Slc2a4	solute carrier family 2 (facilitated glucose transporter), member 4	2.94	1.00	1.09	1.42	0.00	0.00
Slc2a5	solute carrier family 2 (facilitated glucose transporter), member 5	464.30	474.56	488.50	676.64	630.38	552.93
Slc2a6	solute carrier family 2 (facilitated glucose transporter), member 6	74.60	59.19	78.33	13.45	12.45	7.83
Slc2a7	solute carrier family 2 (facilitated glucose transporter), member 7	0.00	0.00	0.00	0.00	0.00	0.00
Slc2a8	solute carrier family 2, (facilitated glucose transporter), member 8	22.58	27.09	27.20	38.22	35.08	29.36
Slc2a9	solute carrier family 2 (facilitated glucose transporter), member 9	12.76	12.04	9.79	4.95	5.66	9.79
Slc2a10	solute carrier family 2 (facilitated glucose transporter), member 10	0.00	0.00	0.00	4.95	2.26	2.94
Slc2a12	solute carrier family 2 (facilitated glucose transporter), member 12	0.00	0.00	0.00	0.71	0.00	0.00
Slc5a1	solute carrier family 5 (sodium/glucose cotransporter), member 1	1.96	2.01	1.09	1.42	1.13	0.98
Slc5a2	solute carrier family 5 (sodium/glucose cotransporter), member 2	0.00	0.00	0.00	0.00	0.00	0.00
Slc5a3	solute carrier family 5 (inositol transporters), member 3	nd	nd	nd	nd	nd	nd
Slc5a4a	solute carrier family 5, member 4a	0.00	0.00	0.00	0.00	0.00	0.00
Slc5a4b	solute carrier family 5 (neutral amino acid transporters, system A), member 4b	0.00	0.00	0.00	0.00	0.00	0.00
Slc5a9	solute carrier family 5 (sodium/glucose cotransporter), member 9	1.96	4.01	1.09	1.42	0.00	5.87
Slc5a10	solute carrier family 5 (sodium/glucose cotransporter), member 10	1.96	1.00	0.00	4.25	1.13	0.00
Slc5a11	solute carrier family 5 (sodium/glucose cotransporter), member 11	0.00	0.00	0.00	0.00	0.00	0.00
Slc45a1	solute carrier family 45, member 1	85.40	62.20	62.01	0.00	0.00	0.00
Slc45a2	solute carrier family 45, member 2	0.00	0.00	0.00	0.00	0.00	0.00
Others							
Gckr	glucokinase regulatory protein	0.00	0.00	0.00	0.71	0.00	0.98
Pklr	pyruvate kinase liver and red blood cell	18.65	6.02	17.41	41.05	55.46	46.00

1000 **Supplementary Table 1.**

1001 Expression profiles of glucose/fructose catabolism-related genes in MIN6-K8 cells and B6
1002 mouse islets. Source data: DDBJ Sequence Read Archive (DRA) accession number
1003 DRA006332 (Hashim et al., 2018). Genes were grouped according to the indicated Gene
1004 Ontology (GO) terms. n = 3 for each. nd, not detected.