Bidirectional Optogenetic Modulations of Peripheral Sensory Nerve Activity: Induction

vs. Suppression through Channelrhodopsin and Halorhodopsin

Akito Kosugi[1], Ken-ichi Inoue[2], Masahiko Takada[2], and Kazuhiko Seki[1*]

[1]Department of Neurophysiology, National Institute of Neuroscience, National Center of

Neurology and Psychiatry, Kodaira, Tokyo 187-8502, Japan

[2]Systems Neuroscience Section, Department of Neuroscience, Primate Research Institute,

and Center for the Evolutionary Origins of Human Behavior, Kyoto University, Inuyama,

Aichi, 484-8506, Japan

*Corresponding author: Kazuhiko Seki, Ph.D.

Department of Neurophysiology, National Institute of Neuroscience, National Center of

Neurology and Psychiatry, Kodaira, Tokyo 187-8502, Japan

Phone: +81-42-346-1724

E-mail: seki@ncnp.go.jp

Summary

In this study, we investigated the potential of optogenetics for modulating activity of

peripheral sensory nerves, particularly tactile and proprioceptive afferents, which are vital for

movement control. Using adeno-associated virus serotype 9 vector, we selectively transduced

channelrhodopsin (ChR2) and halorhodopsin (eNpHR3.0) into large-diameter sciatic nerve

afferents of rats. Diverging from conventional dorsal root ganglion (DRG) approaches, we

applied optical stimulation at the distal portion of the afferent nerve. The intensity of optical

stimulation varied to modulate the extent of induction and suppression of afferent activity.

Then, the effect of optical stimulation was determined by the activity recorded in the dorsal

root of the same afferents. Our findings show successful induction and suppression of activity

in large-diameter afferents via optical stimulation. By increasing the intensity of blue (for

ChR2) and yellow (for eNpHR3.0) light stimulation, the activity of fast-conducting afferent

fibers was preferentially evoked or inhibited in an intensity-dependent manner. These data

indicate that the activity of large-diameter afferents can systematically be regulated by

optogenetics. The present innovative methodology for manipulating specific sensory

modalities at the nerve level offers a targeted and accessible alternative to DRG stimulation,

expanding the therapeutic scope of optogenetics for treating sensory disorders.

Keywords: adeno-associated virus, peripheral nerve, dorsal root ganglion neuron, touch,

proprioception

Introduction

Peripheral sensory nerve activity is integral in conveying sensory experiences to the central

nervous system (CNS). Dysregulation of the peripheral sensory nerves, manifesting hyper- or

hypoactivation, is a hallmark of various sensory disorders (1, 2). Conventional treatments for

regulating peripheral nerve activity, such as pharmacological and electrical interventions (3−

6), often lack specificity and may introduce adverse effects (4, 7, 8). The advent of

optogenetics – a technique that manipulates neural activity with light-sensitive proteins –

offers an alternative solution with unprecedented temporal and spatial precision that has

already been established in the CNS (9−15).

In the present study, we attempted the extension of this optogenetic technique to

peripheral sensory nerves, specifically targeting large-diameter afferents associated with

tactile and proprioceptive sensations. The dorsal root ganglion (DRG) is the standard site for

neural modulation (16−19), but is encumbered by its anatomical complexity and procedural

risks (20, 21). By contrast, peripheral nerves are more accessible targets with reduced

complexity, as evidenced by successful optogenetic manipulation of motor neuron axons (22−

25). The heterogeneity of peripheral sensory nerves, comprising afferents of varying

modalities and diameters (26−29), introduces specific challenges for optogenetic application.

Our previous work demonstrated the efficacy of adeno-associated virus serotype 9 (AAV9)

vector for gene transduction into DRG neurons linked to fast-conducting primary afferent

fibers (19, 30). In this study, we used the AAV9 vector to target selectively large-diameter

sensory fibers in the sciatic nerve of rats transduced with a light-activated ion channel,

channelrhodopsin (ChR2), and a light-driven chloride pump, halorhodopsin (eNpHR3.0). We

found that optical stimulation of the sciatic nerve evoked or suppressed action potentials in its

afferent fibers, with the magnitude of effect directly proportional to the light intensity. This

suggests the feasibility of optogenetic approaches to fine regulation of sensory signals

transmitted through peripheral nerve fibers.

Results

Transduction of ChR2 and eNpHR3.0 Using AAV9 Vector

We successfully transduced ChR2 ($n = 7$) or eNpHR3.0 ($n = 10$) into DRG neurons by intra-

nerve injection of AAV9 vector (Fig. 1A and B). Using an acute experimental setup, we

evaluated the effect of optical stimuli on the distal portion of the sciatic nerve (Fig. 1C and

E).

Optogenetic Induction and Suppression of Sensory Afferent Activity

In rats transduced with ChR2, the effect of blue light was examined by recording the

conducting afferent volleys at the dorsal root (Fig. 1C). For rats transduced with eNpHR3.0,

the nerve was pre-activated by electrical stimulation, and the effect of yellow light applied

proximally to the stimulation site was then assessed at the dorsal root (Fig. 1E).

Representative results are shown in Figure 1D and F. Specifically, in the ChR2-transduced

rats, 470-nm light produced long-lasting volleys in the dorsal root, while 593-nm light had no

modulatory effect (Fig. 1D). In the eNpHR3.0-transduced rats, the volley size evoked by

electrical stimulation (Fig. 1F, gray line) was significantly diminished by optical stimulation

at 593 nm (Fig. 1F, black line, and Fig. S1B). We found comparable induction or significant

suppression of action potentials in all ChR2- ($n = 7$) and eNpHR3.0-transduced rats ($n = 10$).

Since this is the first study to inhibit action potential conduction at the peripheral nerve level,

we further characterized the optogenetic suppression and confirmed a stable and reversible

manner for inhibiting action potential conduction (Fig. S1). These findings demonstrated that

light application could effectively evoke action potentials in sensory fibers of the ChR2-

transduced rats and suppress sensory fiber activity in the eNpHR3.0-transduced rats.

Modulations of Sensory Afferent Activity Dependent on Optical Stimulation Intensity

Next, we examined the impact of varying optical stimulation intensity on sensory afferent

activity. In the ChR2-transduced rats, increasing the intensity of blue light from 20.5 to 51.6

mW resulted in a gradual increase in the amplitude of the recruited volley (Fig. 2A and B;

significant main effect of light power analyzed by Friedman's test, $\chi^2_{(6,60)} = 63.9$, $p < 0.0001$).

In eNpHR3.0-transduced rats, on the other hand, increasing the power of yellow light (≥ 8.7

mW) gradually decreased the amplitude of the electrically evoked volley (Fig. 2D and E;

significant main effect of light power analyzed by Friedman's test, $\chi^2_{(9,81)} = 73.8$, $p < 0.0001$).

These observations confirmed that optical stimulation could regulate the amplitude of

recruited volleys by tuning the stimulation intensity. In addition, increasing the light power

resulted in a shorter onset latency of the recruited volley in the ChR2-transduced rats (Fig.

2C) and a longer onset latency of the electrically evoked volley in the eNpHR3.0-transduced

rats (Fig. 2F, latency suppression). The overall findings indicate that sensory afferent nerve

activity can be finely controlled by adjusting optical stimulation parameters.

Selective Modulation of Large-diameter Afferent Activity

Next, we examined the afferent selectivity of optogenetic modulation, given the tropism of

AAV9 vector for fast-conducting, large-diameter afferents (19, 30). In the ChR2-transduced

rats, the lower power of optical stimulation exclusively recruited early-onset volleys, while

its higher power additionally recruited late-onset volleys as well (Fig. 3A). Specifically,

early-onset volleys were not observed at 11.4 mW irradiation, but became apparent at around

20.5 mW irradiation (Fig. 3B). By increasing the light power, late-onset volleys were first

observed at around 64.7 mW (Fig. 3B). This outcome aligned with the expected large-

diameter selectivity of AAV9 vector. We further compared the maximal amplitude between

the early-onset (1.5–4 ms after stimulus onset; Fig.3A, red shading) and the late-onset (5–10

ms after stimulus onset; Fig.3A, blue shading) volleys. The earlier range corresponded with a

conduction velocity of 15–45 m/s, which well fitted within the expected range for fast-

conducting, large-diameter Aα/β fibers (> 14 m/s) (31). The results showed that the amplitude

was larger in volleys with shorter latencies (Fig. 3C; Wilcoxon signed-rank test, $p = 0.016$).

Therefore, we concluded that large-diameter afferents, potentially Aα/β fibers, were

preferential targets for optogenetic induction and modulation in the ChR2-transduced rats.

For the eNpHR3.0-transduced rats, we also confirmed preferential regulation of

large-diameter afferents. We increased the intensity of electrical stimulation to determine how

optical stimulation with the same parameters suppressed the size of electrically evoked

volleys. We found a marked effect on volleys evoked by a stimulus current just above the

recruitment threshold (1T in Fig. 3D). Notably, these lowest-threshold volleys were

eliminated (100% amplitude inhibition) by optical stimulation in five of the 10 animals. By

contrast, the optical stimulation did not eliminate the volley evoked at 2T, and the

suppressive effect was less marked at 5T. We quantified the relationship between the

suppressive effect and the electrical current in this animal (Fig. 3E). As reported earlier (32,

33), without optical stimulation, the volley size first increased rapidly and then slowly as a

function of electrical current. With optical stimulation, however, this relationship changed

from a non-linear to a more linear manner, and the size of the volley linearly increased as a

function of current. This suggests the possibility that the suppressive effect may be afferent

selective. For example, optogenetic suppression was more effective in the volley evoked at

2T (57%) than 5T (6%; ratio of gray line to black line in Fig. 3E). We calculated the efficacy

of inhibition of the volley amplitude in all animals (Fig. 3F) and found that it was indeed

larger for volleys evoked by a lower stimulus current. Maximal efficacy was obtained at

around 1T and then decreased quickly, though it was still obvious until at 2T. The inhibitory

effect was not statistically significant (Fig. 3F; one-tailed Wilcoxon signed-rank test, p =

0.063 without Holm–Bonferroni correction) despite that the stimulation intensity was

increased to five times as much as the threshold level. We also found a comparable change in

the onset latency of volleys. Indeed, the efficacy to generate a late-onset latency was

decreased in volleys evoked by a higher stimulus current (Fig. 3G and H). These findings

showed that optogenetic suppression had a greater effect on volleys evoked by lower-

intensity electrical stimulation, suggesting preferential suppression of large-diameter afferent

fibers. The overall results indicate that large-diameter sensory afferent activity can selectively

be evoked and suppressed by optogenetics.

Correlation Between Efficacy of Gene Expression and Extent of Optogenetic

Modulation

We examined the relationship between the gene expression levels and the optogenetic

modulation effects in both the ChR2- and the eNpHR3.0-transduced rats (Fig. 4). An example

in Figure 4A shows recruited volleys and yellow fluorescent protein (YFP) expression in

DRG neurons of two ChR2-transduced rats with varying expression levels. One animal

exhibited larger efficacy in optical stimulation and stronger YFP signal intensity. We found a

significant correlation coefficient between the maximal magnitude of recruited volleys and

the YFP signal intensity (Fig. 4B; $r = 0.99$, $p < 0.0001$). A similar comparison was done in the

eNpHR3.0-transduced rats (Fig. 4C). Again, a significant correlation coefficient was found

between the amplitude inhibition percentage and the YFP signal intensity (Fig. 4D; $r = 0.88$,

$p = 0.001$). It should be noted here that these significant correlations were maintained when

an animal with notably high YFP expression was excluded (ChR2: $r = 0.82$, $p = 0.047$;

eNpHR3.0: $r = 0.68$, $p = 0.043$). The results indicate the importance of consistent expression

efficacy across animals for reproducible optogenetic modulation of sensory afferent activity.

164   Overall, our findings lead to similar conclusions in both of the ChR2- and the

eNpHR3.0-transduced rats regarding the stimulation intensity-dependent modulation of large-

diameter sensory afferent fibers and the correlation between the optogenetic effects and the

gene expression efficacy. However, a notable difference was also observed between the two

rat models, as described below.

DRG vs. Nerve: Comparative Analysis of Light Irradiation Sites

Building on previous studies that reported the optogenetic modulation of sensory afferent

activity through light irradiation toward DRG neurons (7, 16−19), we next aimed to compare

the efficacy of modulation between optical stimulation of DRG neurons and peripheral nerve

fibers.

175   In the ChR2-transduced rats, we found that nerve-evoked volleys exhibited smaller

amplitudes compared with DRG-evoked volleys (Fig. 5A–C). Since we applied the same

intensity of optical stimulation in both conditions, this indicates that induction of sensory

afferent activity by nerve irradiation is less effective than by classical DRG stimulation (Fig.

5C; Wilcoxon signed-rank test, $p = 0.016$).

Contrastingly, in the eNpHR3.0-transduced rats, suppression of afferent fiber

activity through nerve irradiation displayed the superior efficacy to DRG irradiation. This

was demonstrated by more pronounced suppression of sensory afferent activity upon optical

stimulation of the nerve compared with the DRG stimulation (Fig. 5D and E; Wilcoxon

signed-rank test, $p = 0.031$).

These findings collectively suggest that while optical stimulation of sensory

afferent fibers and DRG are capable of both inducing and suppressing their activity (as shown

in Figs. 1–4), its effectiveness varies depending on the irradiation site. Specifically, in the

eNpHR3.0-transduced rats, nerve irradiation exhibited more prominent suppression of

activity compared with the classical DRG irradiation approach. Conversely, in the ChR2-

transduced rats, the nerve application proved to be less effective in inducing sensory afferent

activity. Such an efficacy difference highlights the significance of selecting the appropriate

irradiation site to achieve the desired outcome in optogenetic modulation of sensory afferents.

Notably, for induction of sensory afferent activity, the DRG may be a preferable choice,

despite the increased complexity and surgical challenges associated with access thereto. This

consideration is critical when weighing the trade-off between the efficacy and the procedural

complexity in optogenetic manipulation of sensory pathways.

Discussion

There are three points of novelty in the present study. First, we have succeeded in inducing

and suppressing peripheral sensory nerve activity by optical stimulation. Second, the activity

of fast-conducting, large-diameter afferent fibers was preferentially evoked or suppressed in a

stimulation intensity-dependent manner. Third, the efficacy of optical stimulation varied

depending on the irradiation site. In particular, the peripheral nerve was the preferred target

for optogenetic suppression.

Optogenetic Control of Peripheral Nerve Activity

In our study, we have found that optical stimulation can recruit primary afferent activity, with

their amplitude and onset latency modulated in an intensity-dependent manner. A previous

study reported that optical stimulation activated opsins expressed in axons at the node of

Ranvier and generated action potentials (34). When the estimated conduction velocity of

optically evoked volleys was investigated in the present study, it was consistently within the

range of fast-conducting, large-diameter Aα/β fibers (> 14 m/s) (31). This implies the

induction of Aα/β fiber activity by optical stimulation. In addition, our data confirmed

recruitment of high-threshold late component volleys upon optical stimulation at high

intensity with longer light-pulse durations, suggesting an off-target effect of optical

stimulation (19). Nevertheless, the amplitude for late component volleys was usually smaller

than for early component volleys (Fig. 3B and C). Thus, we can conclude that the optical

stimulation in this study preferentially induces the activity of Aα/β fibers.

We also effectively demonstrated optogenetic suppression of axonal action

potentials which were beforehand activated at the time of stimulation. When the action

potentials were generated by electrical stimulation near the threshold current, we observed

almost complete elimination of electrically evoked volleys. Electrical nerve stimulation is

known to recruit action potentials in accordance with axonal diameter, as axons with larger

diameters offer lower electrical resistance, making them more susceptible to depolarization in

response to smaller input signals (35, 36). The earliest component of the recorded volleys,

known as the "Aα" wave, reflects action potentials generated by group I muscle afferent

fibers, which possess the largest axonal diameter among various categories of primary

afferents (33, 37−39). Consequently, the complete elimination of volleys by optogenetic

suppression in this study (Fig. 3D) indicates the superior effectiveness of optical stimulation

in suppressing the activity of group I afferents. We also found the efficient optogenetic

suppression of axonal action potentials primarily evoked by electrical stimulation at

intensities twice as much as their recruitment threshold or below. It is well established that at

such stimulation intensities, activities of both group I and II afferents are introduced, whereas

finer afferents (like those of group III) are not activated (32). Therefore, we suggest that

optogenetic manipulation in this study indeed inhibits the activity of large-diameter primary

afferents, such as group I and II fibers.

Until recently, optogenetic suppression of primary afferent activity has exclusively

been demonstrated by targeting small-diameter DRG neurons that are responsible for

transmitting nociceptive signals (7, 17, 18, 40−42), with limited attention given to large-

diameter neurons. In an effort to fill this gap, we previously attempted to transduce genes by

intra-nerve injection of AAV9 vector (19, 30), known for its affinity toward medium-to-large-

sized DRG neurons that are involved in conveying mechanoreceptive and proprioceptive

signals (43). Here, we have further extended our approach using the AAV9 vector to achieve

not only excitation (19), but also inhibition of large-diameter primary afferents including

group I and II fibers.

DRG vs. Nerve

Theoretically, there are four possible locations for light irradiation to activate opsins

expressed in the peripheral sensory system: peripheral tissue, spinal cord, DRG, and

peripheral nerve (5, 44). Among these, targeting the peripheral tissue has limitations. Light

penetration is restricted to skin tissue when it is irradiated from the body surface (24), thereby

confining the modulatory effect to skin receptors. The impact on deep primary afferent

terminals, such as muscle spindles and Golgi tendon organs, is either impossible or

incomplete. Similarly, targeting the spinal cord (from its surface) is inefficient due to the

highly myelinated dorsal white matter which makes the light scattered (45, 46). Light

intensity is likely to be significantly diminished before reaching the intermediate layer of the

spinal cord where group I fibers terminate. In the present study, we targeted the remaining

two candidates, DRG and peripheral nerve, and revealed that targeting the DRG yielded a

superior efficacy for optogenetic induction of large-diameter primary sensory afferents,

whereas targeting the peripheral nerve had the advantage of their suppression.

These observed differences in the effectiveness of optical stimulation between the

DRG and the peripheral nerve suggest the involvement of distinct mechanisms. In the

peripheral nerve, light irradiation activates opsins expressed in axons and changes the intra-

axonal potential (34). By contrast, irradiation toward the DRG leads to activation of opsins

expressed not only in axons, but also in somata. It was our expectation that the effect of light

irradiation might be amplified by addition of the somal effect. This is because DRG neurons

have active ion channels in their somata and can generate action potentials (47, 48), although

these action potentials are generally believed to be rare. Also, hyperpolarization of the somata

can influence the membrane potential at the T-junction of DRG, where axonal bifurcation

occurs within DRG. Hyperpolarization at the T-junction may interfere with action potential

propagation due to the possible impedance mismatch (49, 50). It seemed surprising that we

obtained the expected result on optogenetic induction but the opposite result on optogenetic

suppression, indicating a lower efficacy of DRG irradiation. Here, we propose two potential

explanations for this outcome. First, the anticipated contribution of T-junction

hyperpolarization may have been less influential than axonal hyperpolarization. Second, the

effect of axonal hyperpolarization due to DRG irradiation may have been minor compared to

nerve irradiation, stemming from structural differences between the two. Specifically, somata

and satellite cells are located predominantly in the dorsomedial region of DRG, while passing

fibers are more evenly distributed throughout the rest of the ganglion (51). Such a location of

somata and satellite cells may result in scattering of the light when irradiated from the DRG

surface, diminishing its intensity by the time it reaches the axons.

The superior efficiency of optical stimulation targeting the nerve, compared with

targeting the DRG, is desirable for *in vivo* application of this technique. In the past, various

chronic optical stimulation tools have been developed, such as nerve cuffs with LED or

optical fibers (52−54), LED directly placed on the peripheral nerve (42), and even

transdermal light irradiation to the peripheral nerve (24, 55). Conversely, methods for chronic

optical DRG stimulation are limited *in vivo* (44) and are highly invasive, posing a risk of

damaging DRG neurons during exposure at the stimulation site (20). Optical stimulation,

especially for optogenetic suppression, at the peripheral nerve level opens new experimental

possibilities of studying the involvement of sensory afferents in a variety of motor behaviors.

Clinical implications

Damage to the CNS in the case of stroke and neurodegenerative diseases, including spinal

muscular atrophy (SMA), lead to maladaptive changes in the periphery. In stroke, one

common complication is spasticity, which results from hyperexcitability of the spinal reflex

(56, 57). While it is assumed that reducing proprioceptive input to the spinal cord could help

prevent the onset of spasticity, there has been no clinical treatment to date that offers selective

and reversible manipulation of proprioceptive afferents. Similarly, SMA is characterized by

motor neuron death and muscle atrophy, with reduced neuronal activity in spinal

sensorimotor circuits as a secondary change (58, 59). Increasing proprioceptive input to the

spinal cord may compensate for motor impairments in patients with SMA (60). In the present

study, we have shown the feasibility of selective and reversible induction and suppression of

large-diameter primary sensory afferent fiber activity using optogenetics. This novel

technique has a significant potential for advancing clinical research works on and therapeutic

approaches to conditioning spinal sensorimotor circuits, thereby offering new possibilities of

targeted interventions against CNS damage caused by stroke and neurodegenerative

disorders.

Methods

Ethical approval

All surgery was performed according to the institutional guidelines for animal experiments

and the National Institutes of Health Guide for the Care and Use of Laboratory Animals. All

experiments were approved by the experimental animal committee of the National Institute of

Neuroscience (approval number 2022020).

Experimental animals

Seventeen male Slc:Wistar rats (4 weeks old, 70–90 g) obtained from Japan SLC, Inc.

(Shizuoka, Japan) were used. All animals were housed under a normal light/dark cycle (12

h:12 h), with the light on from 08:00 h until 20:00 h, in a temperature-controlled environment

with food and water available ad libitum.

Production of viral vectors

AAV9-hSyn-ChR2(H134R)/EYFP (2.5×10^{13} genome copies/mL) (19) and AAV9-CMV-

eNpHR3.0/EYFP (1.8×10^{13} genome copies/mL) were produced using the helper-free triple

transfection procedure and purified by cesium chloride gradient. Viral titer was determined

by quantitative PCR using Taq-Man technology (Life Technologies, Gaithersburg, MD,

USA). Vector purity was assessed by 4–12% sodium dodecyl sulfate–polyacrylamide gel

electrophoresis and fluorescent staining (Oriole, Bio-Rad, Hercules, CA, USA). A transfer

plasmid (pAAV-CMV-eNpHR3.0-EYFP-WPRE) was constructed by inserting the

eNpHR3.0-EYFP fragment with a Woodchuck Hepatitis virus (WHV) posttranscriptional

regulatory element (WPRE) sequence into an AAV backbone plasmid (pAAV-CMV,

Stratagene, La Jolla, CA, USA).

Vector injections

ChR2 ($n = 7$) or eNpHR3.0 ($n = 10$) were transduced into DRG neurons by retrograde

transport following a sciatic nerve AAV injection (Fig. 1A). The protocol for AAV injection

was based on previous studies (19, 30). Briefly, animals were anesthetized by intraperitoneal

injection of medetomidine/midazolam/butorphanol anesthesia (0.2, 2.0, and 2.5 mg/kg,

respectively). Adequate depth of anesthesia was monitored frequently by monitoring pupil

size and flexion reflex to the paw pinch. After anesthesia, the animals were fixed in a prone

position, and an incision was made in the left gluteal muscle. The sciatic nerve was identified

below the biceps femoris muscle and isolated approximately 20 mm from the surrounding

tissue. A 31-gauge needle tube was inserted 10 mm into the isolated nerve. The needle tube

was connected by polyethylene tubing (JT- 10, EICOM, Kyoto, Japan) to a Hamilton syringe

(1702RN, GL Science, Tokyo, Japan) that was mounted into a microinjection pump

(LEGATO 180, KD Scientific, Holliston, MA, USA). The tubing, syringe, and needle tube

were filled with an electrically insulating stable fluorocarbon-based fluid (Fluorinert, 3M, St

Paul, MN, USA). Then 1% Fast Green (0.6 µL) was added to the viral vector solution to

visualize the injected solution. After insertion, the needle tube was left in place for 5 minutes

to allow sealing of the tissue around the needle tip. After, 3 µL of viral vector solution was

injected at a rate of 0.6 µL/min. The needle tube was removed at 10 minutes after the end of

the injection to ensure absorption of the solution. Two separate injections were performed

into the common peroneal and tibial branches of the sciatic nerve to ensure that the nerve was

filled uniformly (in total, 6 µL of viral vector solution was injected). Finally, the wound was

closed with a non-absorbable suture, and the animals were allowed to recover at 37°C.

Acute electrophysiological experiments

To test whether ChR2 and eNpHR3.0 could be used to recruit and inhibit activities of large-

diameter somatosensory afferents *in vivo*, an acute electrophysiological experiment was

performed 4–8 weeks after AAV injection. First, animals were anesthetized by intraperitoneal

injection of medetomidine/midazolam/butorphanol anesthesia (0.2, 2.0, and 2.5 mg/kg,

respectively). After a tracheotomy, anesthesia was maintained by inhalation of 0.5–1.0%

isoflurane throughout surgery. Electrocardiogram, oxygen saturation levels, and rectal

temperature were monitored and maintained within a physiological range.

To expose the lumbar spinal cord, a laminectomy was performed from the 1st

lumbar to the 1st sacral vertebrae. Then, the upper thoracic spine as well as the base of the

tail were immobilized using clamps. The dura was opened with a sharp needle, and the 4th

and 5th lumbar dorsal roots were separated from the other roots. For electrical and optical

stimulation, the left sciatic nerve was exposed and isolated from the surrounding tissue. The

skin and muscle around the exposed roots and nerves were raised and tied to form a pool,

which was filled with mineral oil to protect the exposed roots and nerves from drying out.

After surgery, anesthesia was maintained by thiopental sodium (10 mg/kg), and rocuronium

bromide (10 mg/kg) was administered intravenously to achieve neuromuscular blockade.

To apply electrical stimulation, a bipolar cuff electrode was mounted on the

isolated sciatic nerve. Stimulus current was generated using an isolator (SS 102L, Nihon

Kohden, Tokyo, Japan) coupled to an electrical stimulator (SEN7203, Nihon Kohden) with

various stimulus intensities (1–10-times threshold) at a fixed pulse width (100 μs). A

multimode optical fiber (960 μm in diameter, 0.63 numerical aperture; Doric Lenses, Quebec,

Canada) connected to a light source (LISER_LED470, Doric Lenses) was placed 10 mm

proximal to the cuff electrode to ensure the stimulus site was not directly irradiated (Fig. 1E).

To record stimulus-evoked volleys, a bipolar hook electrode was mounted on the

isolated dorsal root at 15–25 mm proximal to DRG. Recorded potentials were amplified (×

1000) and band-pass filtered (15–10 kHz) using an amplifier (MEG-6108M, Nihon Kohden),

and digitized at 50 kHz using an analog/digital converter (Digidata 1550, Molecular Devices,

LLC, San Jose, CA, USA). The average of 20 responses, consisting of a 10 ms pre-stimulus

baseline period followed by a 100 ms post-stimulus period, was obtained. Then, the peak-to-

peak amplitude and latency between onset of the stimulus and beginning of the response were

calculated. The beginning of the response was defined as a time that exceeded the value of

the mean ± 3× standard deviations of the background signal recorded 10 ms before the

stimulation.

To confirm the effect of optical facilitation, optical stimulation of blue light at 470

nm was applied to the sciatic nerve. The stimulation was applied 20-times with a fixed

stimulus duration (1 ms) at an interval of 0.5 s (2 Hz). To test whether the effect of optical

stimulation was caused by activation of ChR2, irradiation at a different wavelength (593 nm,

1 ms duration) was also performed. To characterize the response properties of optical

facilitation, intensity–response characteristics of the irradiated light power were examined by

changing the intensity of light power (11.4–127.2 mW) at a fixed stimulus duration (1 ms).

To test whether optical stimulation activated large-diameter afferents selectively, intensity–

response characteristics with longer light pulse duration (10 ms) were further examined

because longer and stronger optical stimulation can recruit high-threshold non-target fibers

(19). The light power was measured at the tip of the optical fiber using a power meter

(PM100D, Thorlabs, Newton, New Jersey, USA). To facilitate comparison across animals,

volley amplitudes were normalized to the maximum amplitude of electrically evoked volleys

(10-times threshold intensity).

To confirm the effect of optical inhibition, three experimental sessions were

performed: a baseline session ("Baseline", light-off condition), a session during irradiation of

yellow light at 593 nm ("Light", light-on condition), and a session immediately following

light deactivation ("Recovery", light-off condition). Each session consisted of 20 electrical

stimuli at an interval of 0.5 s (2 Hz) and proceeded consequently without a break. Yellow

light was continuously irradiated for 10 s in only the "Light" session. To facilitate

comparison across animals, volley amplitudes in the "Light" and "Recovery" sessions were

normalized against the averaged amplitude in the preceding "Baseline" session. Considering

reported side effects of light irradiation, such as heating (14, 61−63) and light-off rebound

effects (64), the reversibility of light irradiation was confirmed by comparing "Baseline" and

"Recovery" sessions. Because prolonged light irradiation can decrease the efficacy of

inhibition over time (63, 65, 66), the temporal stationarity of optical inhibition was confirmed

by calculating the correlation between the elapsed time of light irradiation for each stimulus

and the peak-to-peak amplitude of each volley in the "Light" session.

To characterize the response properties of optical inhibition, three sessions were

repeated under various conditions. First, intensity–response characteristics of irradiated light

power were examined by changing the light power intensity (8.7–107.3 mW) at a fixed

electrical stimulation (1.5-times threshold intensity). Then, intensity–response characteristics

of electrical stimulation were examined by changing the intensity of electrical stimulation (1–

5-times threshold) at a fixed light power (107.3 mW). The percentage of amplitude inhibition

was calculated as the difference in normalized amplitude between "Baseline" and "Light"

sessions. The latency difference was also calculated as the difference in latency between

"Baseline" and "Light" sessions and compared in each condition.

At the end of the experiments, small electrolytic lesions were made in the dorsal

root and sciatic nerve using a 40 µA direct current stimulation of the recording and

stimulating bipolar hook electrodes for 40 s. Then, the animals were deeply anesthetized by

intravenous injection of thiopental sodium (20 mg/kg) and perfused transcardially with

phosphate-buffered saline (pH 7.4), followed by 300 mL of 10% neutral buffered

formaldehyde. After perfusion, the lumbar region of the spinal cord together with the DRG

and sciatic nerve were removed.

Histological quantification

To check expression of ChR2 and eNpHR3.0, the fluorescent signal intensity of YFP was

examined. Fluorescent images of entire DRG neurons were acquired using an inverted

microscope (BZ-X710, Keyence, Osaka, Japan) using fixed detector gain values without

sectioning. To calculate the fluorescent intensity, the images were converted to grayscale, and

then mean gray values within the region of interest were determined manually.

Statistical analyses

The Wilcoxon signed-rank test was performed for all two group comparisons. For group

analysis, Friedman's test was performed with light power or recording sessions ("Baseline",

"Light", and "Recovery" sessions) as a within-subject factor. One-way ANOVA was

performed with three recording sessions only for single animal analysis in the optical

inhibition experiment. Post hoc analyses were performed using Tukey's or Dunn's tests for

multiple comparisons. When testing intensity–response characteristics of electrical

stimulation in the optical inhibition experiment, one-tailed Wilcoxon signed-rank test with

Holm–Bonferroni correction was performed to compare the percentage of amplitude

inhibition in each stimulus intensity to the hypothetical value (0% inhibition). For all

correlation analyses, Pearson's correlation coefficient was calculated. The level of

significance was set at $\alpha = 0.05$. All electrophysiological data were analyzed using MATLAB

2021b (MathWorks, Natick, MA, USA), all fluorescent images were analyzed using OpenCV

4.6.0 with python 3.8, and all statistical tests were performed using Prism 10 (GraphPad

Software, San Diego, CA, USA).

Acknowledgments:

We thank Dr. Hidemasa Furue for their helpful comments. Dr. Wupuer Sidikejiang, Dr. Shinji

Kubota, and Ms. Moeko Kudo for their technical assistance. This work was supported by a

Grant-in-Aid from the Japan Society for the Promotion of Science (JSPS) (grant numbers

JP19H05724, JP19H01092, JP26120003, and JP 23H05488 [to K.S.], JP22H04922 [to K. I.],

and JP21K17633 [to A.K.]), and research grants from the Japan Agency for Medical

Research and Development (AMED) (grant numbers JP21dm0207092, JP21dm0207066, and

JP19ek0109216 [to K.S.], and JP21dm0207077 [to K.S. and M.T.]).

Author contributions:

A.K. and K.S. designed the study. K.I. and M.T. developed the viral vectors. A.K. performed

the surgery and electrophysiological experiments. A.K. analyzed the data. A.K. and K.S.

wrote a draft of the manuscript. All authors approved the final version of the manuscript.

Declaration of interests: The authors declare they have no competing interests.

References

1. Woolf, C.J., and Salter, M.W. (2000). Neuronal plasticity: increasing the gain in pain. *Science* **288**, 1765–1769.

2. Navarro, X., Vivó, M., and Valero-Cabré, A. (2007). Neural plasticity after peripheral nerve injury and regeneration. *Prog. Neurobiol.* **82**, 163–201.

3. Navarro, X., Krueger, T.B., Lago, N., Micera, S., Stieglitz, T., and Dario, P. (2005). A critical review of interfaces with the peripheral nervous system for the control of neuroprostheses and hybrid bionic systems. *J. Peripher. Nerv. Syst.* **10**, 229–258.

4. Goins, W.F., Cohen, J.B., and Glorioso, J.C. (2012). Gene therapy for the treatment of chronic peripheral nervous system pain. *Neurobiol. Dis.* **48**, 255–270.

5. Chang, R.B. (2019). Optogenetic control of the peripheral nervous system. *Cold Spring Harb. Perspect Med.* **9**, a034397.

6. Graham, R.D., Sankarasubramanian, V., and Lempka, S.F. (2022). Dorsal Root Ganglion Stimulation for Chronic Pain: Hypothesized Mechanisms of Action. *J. Pain* **23**, 196–211.

7. Daou, I., Beaudry, H., Ase, A.R., Wieskopf, J.S., Ribeiro-da-silva, A., Mogil, J.S., and Séguéla, P. (2016). Optogenetic Silencing of $Na_v1.8$-Positive Afferents. *eNeuro* **3**, ENEURO.0140-15.2016.

8. Montgomery, K.L., Iyer, S.M., Christensen, A.J., Deisseroth, K. and Delp, S. L. (2016). Beyond the brain: Optogenetic control in the spinal cord and peripheral nervous system. *Sci. Trans. Med.* **8**, 337rv5.

9. Boyden, E.S., Zhang, F., Bamberg, E., Nagel, G., and Deisseroth, K. (2005). Millisecond-timescale, genetically targeted optical control of neural activity. *Nat. Neurosci.* **8**, 1263–1268.

10. Nagel, G., Brauner, M., Liewald, J.F., Adeishvili, N., and Bamberg, E. (2005). Light activation of Channelrhodopsin-2 in excitable cells of *caenorhabditis elegans* triggers rapid behavioral responses. *Curr. Biol.* **15**, 2279–2284.

11. Zhang, F., Wang, L.P., Brauner, M., Liewald, J.F., Kay, K., Watzke, N., Wood, P.G.

Bamberg, E., Nagel, G., Gottschalk, A. et al. (2007). Multimodal fast optical

interrogation of neural circuitry. *Nature* **446**, 633–639.

12. Deisseroth, K. (2011). Optogenetics. *Nat. Methods* **8**, 26–29.

13. Deisseroth, K. (2015). Optogenetics: 10 years of microbial opsins in neuroscience. *Nat.*

Neurosci. **18**, 1213–1225.

14. Yizhar, O., Fenno, L. E., Davidson, T. J., Mogri, M., and Deisseroth, K. (2011).

Optogenetics in Neural Systems. *Neuron* **71**, 9–34.

15. Piatkevich, K.D., and Boyden, E.S. (2023). Optogenetic control of neural activity: The

biophysics of microbial rhodopsins in neuroscience. *Q. Rev. Biophys.* **57**, e1.

16. Daou, I., Tuttle, A.H., Longo, G., Wieskopf, J.S., Bonin, R.P., Ase, A.R., Wood, J.N., de

Koninck, Y., Ribeiro-da-Silva, A., Mogil, J.S. et al. (2013). Remote optogenetic

activation and sensitization of pain pathways in freely moving mice. *J. Neurosci.* **33**,

18631–18640

17. Boada, M.D., Martin, T.J., Peters, C.M., Hayashida, K., Harris, M.H., Houle, T.T.,

Boyden, E.S., Eisenach, J.C., and Riri, D.G. (2014). Fast-conducting mechanoreceptors

contribute to withdrawal behavior in normal and nerve injured rats. *Pain* **155**, 2646–

2655.

18. Iyer, S.M., Montgomery, K.L., Towne, C., Lee, S.Y., Ramakrishnan, C., Deisseroth, K.,

and Delp, S.L. (2014). Virally mediated optogenetic excitation and inhibition of pain in

freely moving nontransgenic mice. *Nat. Biotechnol.* **32**, 274–278.

19. Kubota, S., Sidikejiang, W., Kudo, M., Inoue, K., Umeda, T., Takada, M., and Seki, K.

(2019). Optogenetic recruitment of spinal reflex pathways from large-diameter primary

afferents in non-transgenic rats transduced with AAV9/Channelrhodopsin 2. *J. Physiol.*

597, 5025–5040.

20. Fischer, G., Kostic, S., Nakai, H., Park, F., Sapunar, D., Yu, H., and Hogan, Q. (2011).

Direct injection into the dorsal root ganglion: Technical, behavioral, and histological

observations. *J. Neurosci. Methods* **199**, 43–55.

21. Yang, O.J., Robilotto, G.L., Alom, F., Alemán, K., Devulapally, K., Morris, A., and

Mickle A.D. (2023). Evaluating the transduction efficiency of systemically delivered

AAV vectors in the rat nervous system. *Front. Neurosci.* **17**, 1001007.

22. Llewellyn, M.E., Thompson, K.R., Deisseroth, K., and Delp, S.L. (2010). Orderly

recruitment of motor units under optical control in vivo. *Nat. Med.* **16**, 1161–1165.

23. Liske, H., Towne, C., Anikeeva, P., Zhao, S., Feng, G., Deisseroth, K., and Delp, S.L.

(2013). Optical inhibition of motor nerve and muscle activity in vivo. *Muscle and Nerve*

47, 916–921.

24. Maimon, B.E., Sparks, K., Srinivasan, S., Zorzos, A.N., and Herr, H.M. (2018).

Spectrally distinct channelrhodopsins for two-colour optogenetic peripheral nerve

stimulation. *Nat. Biomed. Eng.* **2**, 485–496.

25. Williams, J.J., Watson, A.M., Vazquez, A.L., and Schwartz, A.B. (2019). Viral-mediated

optogenetic stimulation of peripheral motor nerves in non-human primates. *Front. Genet.*

13, 759.

26. Stewart, J.D. (2003). Peripheral nerve fascicles: Anatomy and clinical relevance. *Muscle*

and Nerve **28**, 525–541.

27. Hochman, S. (2007). Spinal cord. *Curr. Biol.* **17**, R950-5.

28. Abraira, V.E., and Ginty, D.D. (2013). The sensory neurons of touch. *Neuron* **79**, 618–

639.

29. Gatto, G., Smith, K.M., Ross, S.E., and Goulding, M. (2019). Neuronal diversity in the

somatosensory system: bridging the gap between cell type and function. *Curr. Opin.*

Neurobiol. **56**, 167–174.

30. Kudo, M., Wupuer, S., Fujiwara, M., Saito, Y., Kubota, S., Inoue, K., Takada, M., and

Seki K. (2021). Specific gene expression in unmyelinated dorsal root ganglion neurons

in nonhuman primates by intra-nerve injection of adeno-associated virus 6 vector. *Mol.*

Ther. Methods Clin. Dev. **23**, 11–22.

31. Harper, A.A., and Lawson, S.N. (1985). Conduction velocity is related to morphological

cell type in rat dorsal root ganglion neurones. *J. Physiol.* **359**, 31-46.

32. Brock, L.G., Eccles, J.C., and Rall, W. (1951). Experimental investigations on the

afferent fibres in muscle nerves. *Proc. R. Soc. Lond. B Biol. Sci.* **138**, 453–475.

33. Bradley, K., and Eccles, J.C. (1953). Analysis of the fast afferent impulses from thigh

muscles. *J. Physiol.* **122**, 462–473.

34. Arlow, R.L., Foutz, T.J., and McIntyre, C.C. (2013). Theoretical principles underlying

optical stimulation of myelinated axons expressing channelrhodopsin-2. *Neuroscience*

248, 541–551.

35. McNeal, D.R. (1976). Analysis of a model for excitation of myelinated nerve. *IEEE*

Trans. Biomed. Eng. **23**, 329–337.

36. Lertmanorat, Z., and Durand, D.M. (2004). Extracellular voltage profile for reversing the

recruitment order of peripheral nerve stimulation: A simulation study. *J Neural. Eng.* **1**,

202–211.

37. Erlanger, J., and Gasser, H.S. (1937). Electrical Signs of Nervous Activity. *University*

Pennsylvania Press, Philadelphia.

38. Lloyd, D.P.C. (1946). Integrative pattern of excitation and inhibition in two-neuron

reflex arcs. *J. Neurophysiol.* **9**, 439–444.

39. Boyd, I.A., and Kalu. K.U. (1979). Scaling factor relating conduction velocity and

diameter for myelinated afferent nerve fibres in the cat hind limb. *J. Physiol.* **289**, 277–

297.

40. Iyer, S. M., Vesuna, S., Ramakrishnan, C., Huynh, K., Young, S., Berndt, A., Lee, S. Y.,

Gorini, C.J., Deisseroth, K., and Delp, S.L. (2016). Optogenetic and chemogenetic

strategies for sustained inhibition of pain. *Sci. Rep.* **6,** 30570.

41. Li, B., Yang, X.Y., Qian, F.P., Tang, M., Ma, C., and Chiang, L.Y. (2015). A novel

analgesic approach to optogenetically and specifically inhibit pain transmission using

TRPV1 promoter. *Brain Res.* **1609**, 12–20.

42. Mayer, P., Sivakumar, N., Pritz, M., Varga, M., Mehmann, A., Lee, S., Salvatore, A.

Magno, M., Pharr, M., Johannssen, H.C. et al. (2019). Flexible and lightweight devices

for wireless multi-color optogenetic experiments controllable via commercial cell

phones. *Front. Neurosci.* **13**, 819.

43. Schuster, D.J., Dykstra, J.A., Riedl, M.S., Kitto, K.F., Belur, L.R., McIvor, R.S., Elde,

R.P., Fairbanks, C.A., and Vulchanova, L. (2014). Biodistribution of adeno-associated

virus serotype 9 (AAV9) vector after intrathecal and intravenous delivery in mouse.

Front. Neuroanat. **8**, 42.

44.　Harding, E.K., Fung, S.W., and Bonin R.P. (2020). Insights Into Spinal Dorsal Horn

Circuit Function and Dysfunction Using Optical Approaches. *Front. Neural Circuits* **14**,

31.

45.　Park, S.I., Brenner, D.S., Shin, G., Morgan, C.D., Copits, B.A., Chung, H.U., Pullen,

M.Y., Noh, K.N., Davidson, S., Oh, S.J. et al. (2015). Soft, stretchable, fully implantable

miniaturized optoelectronic systems for wireless optogenetics. *Nat. Biotechnol.* **33**,

1280–1286.

46.　Mondello, S.E., Sunshine, M.D., Fischedick, A.E., Dreyer, S.J., Horwitz, X.G.D.,

Anikeeva, P., Horner, P.J., and Moritz, C.T. (2018). Optogenetic surface stimulation of

the rat cervical spinal cord. *J. Neurophysiol.* **120**, 795–811.

47.　Kent, A.R., Min, X., Hogan, Q.H., and Kramer, J.M. (2018). Mechanisms of Dorsal Root

Ganglion Stimulation in Pain Suppression: A Computational Modeling Analysis.

Neuromodulation **21**, 234–246.

48.　Graham, R.D., Bruns, T.M., Duan, B., and Lempka, S.F. (2019). Dorsal root ganglion

stimulation for chronic pain modulates Aβ-fiber activity but not C-fiber activity: A

computational modeling study. *Clin. Neurophysiol.* **130**, 941–951.

49.　Du, H., Hao, S., Gigout, D., Huang, Y., Yang, L., Li, C., Wang, D., Sundt, D.B. Jaffe, H.

Zhang, N. et al. (2014). Control of somatic membrane potential in nociceptive neurons

and its implications for peripheral nociceptive transmission. *Pain* **155**, 2306–2322.

50.　Sundt, D., Gamper, N., and Jaffe, D.B. (2015). Spike propagation through the dorsal root

ganglia in an unmyelinated sensory neuron: A modeling study. *J. Neurophysiol.* **114**,

3140–3153.

51.　Sperry, Z.J., Graham, R.D., Peck-Dimit, N., Lempka, S.F., and Bruns, T.M. (2020).

Spatial models of cell distribution in human lumbar dorsal root ganglia. *J. Comp. Neurol.*

528, 1644–1659.

52.　Towne, C., Montgomery, K.L., Iyer, S.M., Deisseroth, K., and Delp, S.L. (2013).

Optogenetic Control of Targeted Peripheral Axons in Freely Moving Animals. *PLoS*

ONE **8**, e72691.

53. Michoud, F., Sottas, L., Browne, L.E., Asboth, L., Latremoliere, A., Sakuma, M.,

Courtine, G., Woolf, C.J., and Lacour, S.P. (2018). Optical cuff for optogenetic control of

the peripheral nervous system. *J. Neural. Eng.* **15**, 015002.

54. Song, K.I., Park, S.E., Lee, S., Kim, H., Lee, S.H., and Youn, I. (2018). Compact Optical

Nerve Cuff Electrode for Simultaneous Neural Activity Monitoring and Optogenetic

Stimulation of Peripheral Nerves. *Sci. Rep.* **8**, 15630.

55. Maimon, B.E., Zorzos, A.N., Bendell, R., Harding, A., Fahmi, M., Srinivasan, S.

Calvaresi, P. and Herr H.M. (2017). Transdermal optogenetic peripheral nerve

stimulation. *J. Neural. Eng.* **14**, 34002.

56. Burke, D., Wissel, J., and Donnan, G.A. (2013). Pathophysiology of spasticity in stroke.

Neurology **80**, S20–26.

57. Li, S., and Francisco, G.E. (2015). New insights into the pathophysiology of post-stroke

spasticity. *Front. Hum. Neurosci.* **9**, 192.

58. Mentis, G.Z., Blivis, D., Liu, W., Drobac, E., Crowder, M.E., Kong, L., Alvarez, F. J.,

Sumner, C.J., and O'Donovan, M.J. (2011). Early functional impairment of sensory-

motor connectivity in a mouse model of spinal muscular atrophy. *Neuron* **69**, 453–467.

59. Fletcher, E.V., Simon, C.M., Pagiazitis, J.G., Chalif, J.I., Vukojicic, A., Drobac, E.,

Wang, X., and Mentis, G.Z. (2017). Reduced sensory synaptic excitation impairs motor

neuron function via Kv2.1 in spinal muscular atrophy. *Nat. Neurosci.* **20**, 905–916.

60. Simon, C.M., Blanco-Redondo, B., Buettner, J.M., Pagiazitis, J.G., Fletcher, E.V., Sime

Longang, J.K., and Mentis, G.Z. (2021). Chronic Pharmacological Increase of Neuronal

Activity Improves Sensory-Motor Dysfunction in Spinal Muscular Atrophy Mice. *J.*

Neurosci. **41**, 376–389.

61. Stujenske, J.M., Spellman, T., and Gordon, J.A. (2015). Modeling the Spatiotemporal

Dynamics of Light and Heat Propagation for In Vivo Optogenetics. *Cell Rep.* **12**, 525–

534.

62. Arias-Gil, G., Ohl, F.W., Takagaki, K., and Lippert, M.T. (2016). Measurement,

modeling, and prediction of temperature rise due to optogenetic brain stimulation.

Neurophotonics **3**, 045007.

63. Wiegert, J.S., Mahn, M., Prigge, M., Printz, Y., and Yizhar, O. (2017). Silencing

Neurons: Tools, Applications, and Experimental Constraints. *Neuron* **95**, 504–529.

64. Bansal, H., Gupta, N., and Roy, S. (2020). Comparison of low-power, high-frequency

and temporally precise optogenetic inhibition of spiking in NpHR, eNpHR3.0 and Jaws-

expressing neurons. *Biomed. Phys. Eng. Express* **6**, 045011.

65. Mattis, J., Tye, K.M., Ferenczi, E.A., Ramakrishnan, C., O'Shea, D.J., Prakash, R.,

Gunaydin, L.A., Hyun, M., Fenno, L.E., Gradinaru, V. et al. (2011). Principles for

applying optogenetic tools derived from direct comparative analysis of microbial opsins.

Nat. Methods **9**, 159–172.

66. Zhang, C., Yang, S., Flossmann, T., Gao, S., Witte, O.W., Nagel, G., Holthoff, K., and

Kirmse, K.(2019). Optimized photo-stimulation of halorhodopsin for long-term neuronal

inhibition. *BMC Biol.* **17**, 95.

Figure 1. Optogenetic manipulation of sensory afferent activity by adeno-associated

virus serotype 9 (AAV9)-mediated gene transduction into dorsal root ganglion (DRG)

neurons and optical stimulation of nerve fibers. (A) Schema of AAV injection. Retrograde

transport via sciatic nerve injection enabled gene transduction into DRG neurons. (B)

Expression of YFP in the 5th lumbar DRG neurons. Scale bar: 200 μm. (C, D) Results on

channelrhodopsin (ChR2) transduction. (**C**) Schema of the acute *in vivo* electrophysiological

experiment. (D) Examples of dorsal root potentials evoked by light stimulation with different

wavelengths. Waveforms represent the stimulus-triggered averaging of 20 responses. The bar

above each waveform indicates the timing of light irradiation. (E, F) Results on

670   halorhodopsin (eNpHR3.0) transduction. **(**E) Schema of the acute *in vivo* electrophysiological

671   experiment. (F) Examples of electrically evoked volleys with (black line) and without (gray

672   line) light irradiation. Waveforms represent the stimulus-triggered averaging of 20 responses.

673   The red square above the waveforms indicates the timing of electrical stimulation.

Figure 2. Effect of light power on optogenetic manipulation. (A−C) Results on

channelrhodopsin (ChR2) transduction. (A) Examples of recruited volleys using different

irradiated light powers. Waveforms represent the stimulus-triggered averaging of 20

responses. Blue bars above each waveform indicates the timing of light irradiation. (B, C)

Changes in the amplitude and latency of recruited volleys as a function of light power. Each

gray line represents one animal ($n = 7$), and thick black lines represent the median value of all

animals. (D−F) Results on halorhodopsin (eNpHR3.0) transduction. (D) Examples of

electrically evoked volleys using different irradiated light powers. Waveforms represent the

stimulus-triggered averaging of 20 responses. Black lines indicate electrically evoked volleys

with light irradiation and gray lines indicate volleys without irradiation. Red squares above

686   each waveform indicates the timing of electrical stimulation. (E, F) Changes in the

687   percentage of amplitude inhibition and latency of electrically evoked volleys as a function of

688   light power. Each gray line represents one animal ($n = 10$), and thick black lines represent the

689   median value of all animals.

Figure 3. Large-diameter afferent preference. (A−C) Results of channelrhodopsin (ChR2)

transduction. (A) Examples of recruited volleys using long light pulse durations. Waveforms

represent the stimulus-triggered averaging of 20 responses. Red shading indicates the range

of early-onset volleys (1.5−4 ms after stimulus onset), and blue shading indicates the range of

late-onset volleys (5−10 ms after stimulus onset). Blue bars above the waveforms indicate the

timing of light irradiation. Black arrows indicate individual peaks of late-onset volleys. (B)

Changes in normalized amplitude of early- and late-onset volleys as a function of light power.

The gray line indicates the early-onset component, and the black line indicates the late-onset

component. Note that each line is plotted only at the light power at which the volley was

observed. (C) Maximum normalized amplitude of early- and late-onset volleys evoked by

long light pulse duration. Each gray line represents one animal ($n = 7$), and thick black lines

represent the median value of all animals. *$p < 0.05$ (Wilcoxon signed-rank test). (D−H)

Results of halorhodopsin (eNpHR3.0) transduction. **(D)** Examples of electrically evoked

volleys using different intensities of electrical stimulation. Waveforms represent the stimulus-

triggered averaging of 20 responses. Black lines indicate electrically evoked volleys using

light irradiation and gray lines indicate volleys without irradiation. Red squares above each

waveform indicates the timing of electrical stimulation. (E) Example of intensity–response

curves of amplitudes with and without light irradiation from one animal. (F) Changes in the

percentage of amplitude inhibition as a function of electrical stimulation intensity. Each gray

line represents one animal ($n = 10$), and thick black lines represent the median value of all

animals. *$p < 0.05$; **$p < 0.01$ (one-tailed Wilcoxon signed-rank test with Holm–Bonferroni

correction compared to 0%). (G) Example of intensity–response curves of latency with and

without light irradiation from one animal. (H) Difference in latency as a function of electrical

stimulation intensity. Each gray line represents one animal ($n = 10$), and thick black lines

represent the median value of all animals.

ChR2

Figure 4. Correlation between optical efficacy and YFP intensity. (A, B) Results on

channelrhodopsin (ChR2) transduction. (A) Examples of recruited volleys and YFP

expression in dorsal root ganglion (DRG) neurons for two animals with different expression

levels. Waveforms represent the stimulus-triggered averaging of 20 responses. Blue bars

above each waveform indicates the timing of light irradiation. Scale bar: 200 μm. (B) Scatter

plots and linear correlation between the normalized amplitude and YFP signal intensity of all

animals ($n = 7$). (C, D) Results on halorhodopsin (eNpHR3.0) transduction. (C) Examples of

electrically evoked volleys with (black line) and without (gray line) light irradiation and YFP

expression in DRG neurons for two animals with different expression levels. Waveforms

represent stimulus-triggered averaging of 20 responses. Red squares above the waveforms

729   indicate the timing of electrical stimulation. Scale bar: 200 μm. (D) Scatter plots and linear

730   correlation between the amplitude inhibition percentage and YFP signal intensity of all

731   animals ($n = 10$).

Figure 5. Sites and effects of optical stimulation. (A−C) Results on channelrhodopsin

(ChR2) transduction. (A) Schema of the acute *in vivo* electrophysiological experiment. (B)

Examples of recruited volleys by optical stimulation to the sciatic nerve and dorsal root

ganglion (DRG). Waveforms represent the stimulus-triggered averaging of 20 responses.

Blue bars above the waveforms indicate the timing of light irradiation. (C) The effect of the

stimulus site on the amplitude. Each gray line represents one animal ($n = 7$), and thick black

lines represent the median value of all animals. (D, E) Results on halorhodopsin (eNpHR3.0)

transduction. (D) Schema of the acute *in vivo* electrophysiological experiment. (E) The effect

of the stimulus site on the percentage of amplitude inhibition. Each gray line represents one

animal ($n = 6$), and thick black lines represent the median value of all animals. *$p < 0.05$.

Figure S1. Characterization of optogenetic suppression. (A) Time course of peak-to-peak

amplitudes of electrically evoked volleys. Red dots indicate the timing of electrical

stimulation (2 Hz). The orange line and orange shaded area indicate the timing of light

irradiation (10 s). There was no significant correlation between the elapsed time of light

irradiation and the peak-to-peak amplitude in the "Light" session (r = -0.073, p = 0.76),

indicating no time trend in this session. (B, C) Effect of optical inhibition on the amplitude

from a representative animal (B) and for all animals (C). For representative animal, data are

represented mean ± standard deviation. For all animals, amplitudes in the "Light" and

"Recovery" sessions were normalized by the average amplitude in the preceding "Baseline"

session. Each gray line represents one animal (n = 10), and thick black lines represent the

median value of all animals. Significant differences were found between the "Baseline" and

the "Light" session in a representative animal (one-way analysis of variance [ANOVA], $F_{(2,57)}$

= 4365, $p < 0.0001$; Tukey's post hoc test, $p < 0.0001$), and all animals (Friedman's test,

$\chi^2_{(2,18)}$ = 15.2, $p < 0.0001$, Dunn's post hoc test, $p = 0.001$). No difference was seen between

the "Baseline" and the "Recovery" session ($p = 0.61$ in the representative animal and $p = 0.99$

in all animals), suggesting that light irradiation causes no side effects. (D, E) Effect of optical

inhibition on latency from a representative animal (D) and for all animals (E). For

representative animal, data are represented mean ± standard deviation. For all animals, each

gray line represents one animal ($n = 10$), and thick black lines represent the median value of

all animals. The onset latency of the volley in the "Light" session was significantly longer

than that in the "Baseline" session in a representative animal (one-way ANOVA, $F_{(2,57)}$ =

742.3, $p < 0.0001$; Tukey's post hoc test, $p < 0.0001$), and all animals (Friedman's test, $\chi^2_{(2,18)}$

= 10.6, $p = 0.002$, Dunn's post hoc test, $p = 0.011$). No difference was found between the

"Baseline" and the "Recovery" session ($p = 0.79$ in the representative animal and $p = 0.99$ in

all animals), suggesting the reversible nature of latency inhibition. *$p < 0.05$; **$p < 0.01$;

****$p < 0.0001$.