Title

Restraint of melanoma progression by cells in the local skin environment

Authors

Yilun Ma[1-3], Mohita Tagore[2], Miranda V. Hunter[2], Ting-Hsiang Huang[2], Emily Montal[2], Joshua M. Weiss[1-3], Richard M. White[2,4]

Affiliations

[1] Weill Cornell/Rockefeller/Sloan Kettering Tri-Institutional MD-PhD Program, New York, NY, USA

[2] Department of Cancer Biology and Genetics, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[3] Cell and Developmental Biology Program, Weill Cornell Graduate School of Medical Sciences, New York, NY, USA

[4] Nuffield Department of Medicine, Ludwig Cancer Research, University of Oxford, Oxford, UK

Abstract

Keratinocytes, the dominant cell type in the melanoma microenvironment during tumor initiation, exhibit diverse effects on melanoma progression. Using a zebrafish model of melanoma and human cell co-cultures, we observed that keratinocytes undergo an Epithelial–Mesenchymal Transition (EMT)-like transformation in the presence of melanoma, reminiscent of their behavior during wound healing. Surprisingly, overexpression of the EMT transcription factor Twist in keratinocytes led to improved overall survival in zebrafish melanoma models, despite no change in tumor initiation rates. This survival benefit was attributed to reduced melanoma invasion, as confirmed by human cell co-culture assays. Single-cell RNA-sequencing revealed a unique melanoma cell cluster in the Twist-overexpressing condition, exhibiting a more differentiated, less

invasive phenotype. Further analysis nominated homotypic jam3b-jam3b and pgrn-sort1a

interactions between Twist-overexpressing keratinocytes and melanoma cells as potential

mediators of the invasive restraint. Our findings suggest that EMT in the tumor microenvironment

(TME) may limit melanoma invasion through altered cell-cell interactions.

Introduction

The complex interplay between cancer cells and their microenvironment has emerged as a critical

determinant of tumor progression and therapeutic response. In melanoma, the tumor

microenvironment (TME) encompasses diverse cell types, including immune cells, fibroblasts,

and endothelial cells[1]. However, during melanoma initiation the dominant cell type in the TME is

the keratinocyte, an epithelial cell which makes up the majority of our skin surface. In normal

homeostasis, each melanocyte reciprocally interacts with 30-40 keratinocytes[2], and this

interaction is essential for skin and hair color[3,4]. Despite decades of research, our understanding

of keratinocytes in the context of melanoma remains incomplete.

Keratinocytes have been shown to both inhibit and promote melanoma initiation. They are tightly

adherent to melanocytes through homophilic interactions of the cell adhesion molecule E-

cadherin[5]. Through E-cadherin, keratinocytes can control melanocyte growth and behavior[6–9]. As

melanoma progresses, they undergo a cadherin subtype switch in which they downregulate E-

cadherin and upregulate N-cadherin, thereby escaping keratinocyte-mediated controls while

promoting migration and survival[10–12]. Beyond cadherins, invasive melanoma can downregulate

matricellular proteins such as CCN3, which usually facilitates melanoma attachment to the

basement membrane[13,14]. In addition to melanoma intrinsic changes, alterations in epidermal

keratinocytes such as the loss of PAR3 expression results in a local environment that facilitates

melanoma invasion and metastasis[15]. In contrast, keratinocytes can also promote tumor

development through secretion of growth factors such as endothelins or via GABAergic crosstalk

between the two cell types[16–18].

These conflicting data highlight that interactions between keratinocytes and nascent melanoma

cells are likely dynamic and change rapidly during tumor initiation. Studying the nature of these

interactions in human samples is challenging because biopsies are taken after the patient has

come to the clinic, meaning that the earliest interactions in tumor initiation will be missed. This

necessitates models which faithfully recapitulate the earliest stages of tumor initiation, yet have

the cellular resolution to measure interactions between melanoma cells and keratinocytes.

In this study, we utilized a zebrafish model of melanoma to investigate the earliest interactions

between melanoma cells and their neighboring keratinocytes[19]. Zebrafish have emerged as a

powerful tool for cancer research due to their genetic tractability, conserved biology, and the ability

to visualize tumor development and progression in real-time within the context of an intact

organism[20,21]. Using a combination of cell-type specific genetic manipulations, in vivo imaging,

and single-cell transcriptomics, we found that tumor-associated keratinocytes undergo changes

associated with EMT, similar to what is found in wounded skin. Unexpectedly, we found that this

keratinocyte EMT suppresses melanoma progression. This change in the keratinocytes occurs

shortly after melanoma initiation, and results in keratinocytes which are more adhesive to these

nascent tumor cells and prevents their movement out of the epidermis. Our data suggests that

melanoma initiation revises an evolutionarily conserved wounding response in the nearby skin

environment, which acts as a cell extrinsic tumor suppressor to prevent newly transformed cells

from becoming clinically meaningful.

Results

Melanoma initiation is associated with EMT in keratinocytes

To investigate the relationship between keratinocytes and melanoma cells *in vivo*, we created a

transgenic zebrafish line in which GFP is expressed under the *krt4* promoter[22]. This line faithfully

marks all adult keratinocytes present throughout the fish epidermis and scales, similar to previous

lines using this promoter (Figure 1A). We then initiated melanomas in this background using the

TEAZ method (Transgene Electroporation in Adult Zebrafish)[19,23], in which plasmids containing

oncogenes or sgRNAs against tumor suppressors can be introduced directly into the skin (Figure

1B). The major advantage of this method is that we can visualize melanoma initiation when the

tumor is in its early stages, consisting of a small number of cells. We initiated tumors with a

combination of BRAF^{V600E}, sgRNAs against PTEN (zebrafish *ptena/b)* and germline loss of p53,

using *mitfa*-Cas9 to ensure that *PTEN* were inactivated only in melanocytes and not in

surrounding skin cells (Figure 1C). To account for skin wounding from electroporation in

assessing changes in tumor-associated keratinocytes, we also performed TEAZ using a control

vector that labels melanocyte-precursors but does not induce melanoma formation. Fluorescent

imaging 8-weeks post-electroporation with the control vector demonstrated an injury-free

epidermis, in contrast to the pronounced melanoma development in zebrafish administered with

the oncogenic vectors (Figure 1D).

After initiating tumorigenesis, we examined the morphology of tumor-adjacent keratinocytes to

investigate whether the growing tumor could be influencing keratinocyte behavior. We performed

confocal microscopy on the scales of fish 8 weeks post-electroporation. This revealed a marked

disruption of keratinocyte morphology in the tumor bearing fish, which was not seen in control

fish. We specifically noted disrupted cell-cell junctions, a disorganized pattern of keratinocytes,

and loss of the normal hexagonal cell layer (Figure 1E). These changes were reminiscent of

keratinocyte EMT, which has been previously noted to occur in wounded epidermis[24-26]. To further

assess this possibility, we excised tissues from both tumor and control skin and used FACS to

isolate keratinocytes (GFP+) and melanoma cells (tdTomato+) and performed qPCR (Figure 1B).

As expected, we found enrichment of *mitfa* in melanoma cells and *krt4* in keratinocytes, validating

the successful cell-type isolation (Figure 1F). Comparative analyses between tumor-associated

keratinocytes (TAKs) and normal keratinocytes (NKCs) from tissue without melanoma revealed

upregulation of EMT markers vimentin and N-cadherin in TAKs, confirming the EMT-like

morphological changes of keratinocytes in our imaging results (Figure 1G).

We next asked whether these changes were also seen in human samples. To address this, we

performed co-culture experiments between keratinocytes and melanoma cells. We grew GFP-

labeled HaCaT keratinocytes either alone or with A375 melanoma cells for 21 days, followed by

isolation by FACS for bulk RNA-sequencing as published in Tagore et al.[18] (Figure 1H). Consistent

with our *in vivo* results in the fish, the top pathway altered in the co-cultured HaCaT cells was

enrichment of EMT (Figure 1I). Differential gene expression analysis showed notable upregulation

of the mesenchymal markers vimentin and N-cadherin in co-cultured keratinocytes compared to

monocultured control keratinocytes (Figure 1J), similar to what was found in vivo. Collectively, our

data indicate that melanoma cells induce morphological and molecular markers of EMT in nearby

keratinocytes.

EMT transcription factors are upregulated in tumor-associated keratinocytes

EMT is usually driven by upstream transcription factors, which then act on downstream targets to

repress adhesion molecules such as E-cadherin or activate other adhesion molecules such as N-

cadherin[26]. We next wanted to understand which of these transcription factors was responsible

for the EMT-like behavior in tumor-associated keratinocytes. To address this, we utilized an

existing scRNA-sequencing dataset of a BRAF^{V600E}-driven zebrafish melanoma[27] (Figure 2A).

Dimensionality reduction with UMAP and subsequent clustering revealed two keratinocyte

populations as indicated by module scoring for genes enriched in zebrafish keratinocyte

populations (Figure 2B). Subsequent differential gene expression and GSEA analysis of the two

keratinocyte clusters revealed one cluster with enrichment for EMT, similar to what we observed

in Figure 1 (Figure 2C-D). We refer to this EMT cluster as Tumor Associated Keratinocytes

(TAKs), and the other cluster as a Normal Keratinocyte Cluster (NKCs). We focused on three

EMT-transcription factors expressed by zebrafish keratinocytes in this dataset, *snai1a*, *snai2*,

twist1a, zebrafish homologs of human SNAIL, SLUG and TWIST. Differential expression showed

significant enrichment in *snai1a* and *twist1a* in TAK vs. NKC clusters (Figure 2F). The significant

enrichment of *twist1a* in the TAK cluster, coupled with its rare expression in NKCs, positioned

twist1a as a promising candidate for further investigation into its potential role in driving EMT-like

changes in keratinocytes and, consequently, its impact on melanoma progression.

Keratinocyte *TWIST* restrains melanoma invasion in zebrafish

Having identified Twist as a potential driver of EMT-like changes in tumor-associated

keratinocytes, we next asked how it affected melanoma phenotypes. To investigate the role of

Twist in keratinocytes, we created new transgenic zebrafish in which the keratinocyte-specific *krt4*

promoter was used to drive the two zebrafish *TWIST* paralogs (*twist1a* and *twist1b*) in the

presence of BRAF^{V600E}-driven melanomas. We injected the plasmids for melanoma initiation to

zebrafish embryos and then monitored the fish for tumor-free survival as well as overall survival

over the ensuing 26 weeks (Figure 3A). Interestingly, we found no difference in melanoma

initiation rate in the TWIST overexpression condition compared to empty vector (Figure 3B).

Unexpectedly, however, we noted that overall survival was improved in the transgenic animals

expressing TWIST in the keratinocytes (Figure 3B).

This discrepancy between melanoma-free and overall survival suggested that the tumors in the

TWIST condition should be phenotypically distinct from the control tumors. To assess this, we

performed immunohistochemistry on the tumors and surrounding tissues from the control and

TWIST conditions (Figure 3C). The oncogenic driver in this melanoma model is hBRAF^{V600E} and

serves as an IHC marker for the tumor cells. Comparison of hBRAF^{V600E} staining revealed

significant melanoma infiltration into the zebrafish body in the CTRL condition, as opposed to a

nearly non-invasive tumor in the TWIST condition (Figure 3D). The lack of melanoma invasion

was also observed when the melanoma developed in other anatomical locations (Supp. Figure

1). These findings suggest that *twist1a/twist1b* overexpression in keratinocytes does not impair

tumor initiation, but instead impairs melanoma invasion and improves survival when expressed in

the microenvironment.

To further validate this finding, we developed a cell culture-based invasion assay by pairing the

HaCaT keratinocyte cell line with multiple melanoma cell lines that were pre-cultured on coverslips

(Figure 4A). Due to the migratory nature of melanoma cells, they will migrate off the coverslip and

infiltrate the layer of keratinocytes, allowing us to assess relative differences between culture

conditions. HaCaT keratinocytes were transformed via lentiviral transduction to overexpress

human *TWIST1* (HaCaT-TWIST) or an empty vector control (HaCaT-CTRL) (Figure 4B). Western

blot analysis confirmed robust Twist protein overexpression in HaCaT-TWIST compared to

HaCaT-CTRL (Figure 4C), and immunofluorescence imaging revealed nuclear localization of

Twist (Figure 4D). Co-culture of HS294T melanoma cells with HaCaT-TWIST resulted in

significantly reduced melanoma cell invasion into keratinocytes compared to HaCaT-CTRL

(Figure 4E-F), similar to what we had observed *in vivo*. This finding was recapitulated using the

SKMEL2 human melanoma cell line, demonstrating the inhibitory effect of *TWIST1*

overexpression in keratinocytes on melanoma invasion across different cell lines (Figure 4G-H).

Collectively, our results demonstrate that induction of EMT in keratinocytes is associated with

reduced melanoma invasion and improvement in animal survival.

Keratinocyte EMT promotes aberrant adhesion to nascent melanoma cells

While EMT of tumor cells is well recognized to promote invasion, our data suggest that EMT in

the microenvironment paradoxically restrains tumor invasion. We wanted to better understand the

downstream mechanisms accounting for this result. We therefore analyzed our CTRL vs. TWIST

overexpression tumors and adjacent microenvironment using single-cell RNA sequencing, which

would allow us to understand potential mechanisms by which melanoma cells were interacting

with these keratinocytes. Melanoma and adjacent skin from three fish per condition (CTRL or

TWIST) were dissociated into single cell suspensions and FACS-sorted for eGFP+ keratinocytes

and tdTomato+ melanoma cells (Figure 5A and Supp Figure 2A). In the resultant scRNA-seq

dataset, we identified distinct clusters of eGFP+ keratinocytes and tdTomato+ melanoma cells

(Figure 5B). Of the keratinocyte clusters identified, two were present in both the CTRL and TWIST

conditions, while a third was only present in samples overexpressing TWIST in keratinocytes

(Figure 5C). To identify biological processes that may be active within the keratinocyte, we

performed GSEA, comparing differentially expressed genes between the two clusters present in

both conditions. GSEA identified an enrichment in the EMT pathway as seen in Figure 2, allowing

us to label the EMT-enriched cluster as TAK and other as NKC (Figure 5D and Supp Figure 2B).

As expected, we also identified a unique cluster of keratinocytes only present in the TWIST

dataset that highly expresses *twist1a/b*, the genes that we overexpressed in the TWIST condition,

and labeled this cluster Twist-High (Figure 5D and Supp Figure 2C).

To understand how the Twist-high keratinocytes may be restraining melanoma progression, we

compared gene expression in melanomas that arose in the presence of control vs. Twist

overexpression keratinocytes by comparing their gene signatures to published melanoma gene

signatures representing a range of differentiation states[28]. It is now widely recognized that

melanoma cells exist along a trajectory of differentiation, ranging from undifferentiated/invasive,

to neural crest, to intermediate, to melanocytic/proliferative[28]. Interestingly, we found a cluster of

melanoma   cells   that   developed   in   the   TWIST   condition   was   enriched   for   the

melanocytic/proliferative state but not undifferentiated/invasive gene markers (Figure 5E). This is

consistent with our in vivo observations that these melanomas are phenotypically less invasive.

We hypothesized that this change in cell state might be induced by physical interactions between

the Twist-high keratinocytes and the melanoma cells. To address this, we analyzed potential cell-

cell interactions using CellChat, a software tool that allows us to quantitatively characterize and

visualize cell-cell communications using a curated zebrafish ligand-receptor interaction

database[29] (Figure 5F). Two unique ligand-receptor pairings were identified that only occur

between Twist-high keratinocytes and the melanomas that arose in these animals: a homophilic

jam3b-jam3b interaction and a *pgrn-sort1a* (progranulin-sortilin) interaction (Figure 5G).

The *jam3b* interaction was of particular interest to us, as this protein has been recently identified

as one required for melanophore survival in zebrafish[30] and for human melanoma metastasis[31,32].

In normal human skin, JAM1 (or F11R) is expressed in keratinocytes of the superficial epidermis,

whereas its heterophilic partner JAM3 is exclusively found in basal keratinocytes[33,34]. This

distribution is significant because melanomas predominantly originate in the basal area of the

skin. Interestingly, basal keratinocytes, but not superficial keratinocytes, have been shown to

inhibit melanocyte growth[7]. This observation, combined with our findings, suggests that Twist

expression in the keratinocytes was leading to aberrant expression of jam3b, resulting in stronger

homophilic jam3b-jam3b attachments between these keratinocytes and melanoma cells,

potentially inhibiting melanoma invasion.

Discussion

In this study, we observed that keratinocytes in both zebrafish and human models of melanoma

undergo an EMT-like transformation in the presence of melanoma. This alteration is reminiscent

of keratinocyte behavior during wound healing, in which keratinocytes exhibit markers and

morphological changes associated with EMT in development[25,35]. Interestingly, we observed an

increase in N-cadherin expression in KC, which is usually attributed to melanoma as it becomes

more aggressive and invades into the dermis to associate with fibroblasts[10]. Our findings would

suggest a subpopulation of keratinocytes could maintain contact with melanoma through

upregulation of N-cadherin. Additionally, re-analysis of a published zebrafish melanoma scRNA-

sequencing dataset showed distinct populations of keratinocytes that expressed markers of

EMT, demonstrating the feasibility of studying this KC population using our zebrafish model and

nominating Twist1 as a potent EMT transcription factor in this cell type[27]. Twist expression has

been found to be upregulated at the edge of wounded skin upon treatment with bFGF, a well-

characterized growth factor produced by melanoma[35,36]. If melanoma acts as an open-wound in

the skin, then Twist1 might be upregulated in tumor-associated keratinocytes as an adaptive

response to close this wound.

Our zebrafish melanoma survival experiment showed improvements in overall survival in

zebrafish with keratinocytes overexpressing Twist compared to those that received an empty

vector, despite the fish forming tumors at the same rate. This survival improvement was shown

to be caused by a decrease in melanoma invasion, raising the possibility that Twist

overexpressing keratinocytes could restrain melanoma invasion. Human cell co-cultures with a

HaCaT cell line overexpressing Twist showed a similar finding to our in vivo zebrafish model,

with reduced melanoma cell infiltration into keratinocytes. To learn more about the dynamics of

melanoma and TME keratinocytes, we performed scRNA-sequencing on our zebrafish

melanoma model to account for both keratinocytes in contact with melanoma and those in the

periphery. Compared to the fish that received an empty vector control that had two

subpopulations of keratinocytes, fish with Twist overexpression contained an additional novel

keratinocyte subpopulation with overexpression of *twist1a/b*[27].

Interestingly, we also found a cluster of melanoma cells unique to the TWIST condition, which

shared gene signatures similar to that of the genes observed in both transitory and melanocytic

states as published by Tsoi et al.[28]. These cell states were defined to be more differentiated with

higher MITF expression and correlated to a more proliferative but less invasive cohort from

Hoek et al.[37]. The enrichment for specific melanoma cell states when surrounded by a Twist-

overexpressing keratinocyte TME could be responsible for the reduced overall melanoma

invasion.

Further analysis using CellChat nominated *jam3b-jam3b* and *pgrn-sort1a* as unique interactions

between Twist-High keratinocytes and TWIST-Melanoma cells. As previously described, jam3b

has been identified as a critical protein in zebrafish melanophore survival with known

involvement in melanoma metastasis[30]. The aberrant expression of jam3b on Twist-

overexpressing keratinocytes could indicate strong homophilic interactions with melanoma

jam3b that retains the melanoma in the epidermis. Whether jam3b acts in concert or

independent of cadherin-based adhesion remains to be determined in future studies. Although

the progranulin-sortilin interaction has not been characterized in melanoma, sortilin has been

identified as a key regulator of progranulin levels[38]. Progranulin is known to be constitutively

expressed by keratinocytes, which could be cleaved to epithelins that promote KC

proliferation[39,40]. Progranulin is also a potent mediator of the wound response produced by

dermal fibroblasts in addition to epidermal keratinocytes[41]. Perhaps the increase in progranulin

is cleared by melanoma cells through sortilin, resulting in the endocytosis and lysosomal

transport of sortilin[42]. The degradation of sortilin could be responsible for decreased cell

migration and invasion, as sortilin is required for the interaction of proNGF, a neurotrophin

produced by melanoma, with NGFR in promoting melanoma migration[43]. Further studies are

needed to elucidate the precise mechanisms underlying the nominated interactions, jam3b-

jam3b and pgrn-sort1a, and to explore their potential as therapeutic targets in melanoma.

Methods

Zebrafish husbandry

Zebrafish were maintained in a dedicated facility with controlled temperature (28.5 °C) and salinity. The fish were kept on a 14-hour light/10-hour dark cycle and fed a standard zebrafish diet consisting of brine shrimp followed by Zeigler pellets. Embryos were obtained through natural mating and incubated in E3 buffer (5 mM NaCl, 0.17 mM KCl, 0.33 mM CaCl$_2$, 0.33 mM MgSO$_4$) at 28.5 °C. For procedures requiring immobilization, zebrafish were anesthetized using Tricaine-S (MS-222, Syndel) prepared as a 4 g/L stock solution with a pH of 7.0. The stock solution was protected from light exposure and diluted to the appropriate concentration to achieve fish immobilization. All experimental procedures and animal protocols described in this manuscript were conducted in compliance with the Institutional Animal Care and Use Committee (IACUC) protocol #12-05-008, approved by the Memorial Sloan Kettering Cancer Center (MSKCC).

Generation of zebrafish line with fluorophore labeled keratinocytes

Embryos at the one-cell stage from the Casper Triple zebrafish line (*mitfa:BRAF^{V600E};p53-/-;mitfa-/-;mpv17-/-*)[19,44] were injected with a *krt4:eGFP* expression cassette in the 394 vector of the Tol2Kit[45] with tol2 mRNA. Larvae were sorted for positive GFP fluorescence at day 3 and raised to adult for breeding. F0 fish were in-crossed and resulting F1 were outcrossed with Casper Triple zebrafish for consistent GFP expression. Starting from F2, the *krt4:eGFP* zebrafish line was maintained by out-crossing with Casper Triple zebrafish and sorting for GFP expression.

Transgene Electroporation in Adult Zebrafish (TEAZ)

TEAZ was utilized to generate melanoma as previously described[19,46]. *Krt4:eGFP* zebrafish

(*krt4:eGFP* Casper Triple) were anesthetized with tricaine and injected with a plasmid solution

containing miniCoopR-tdT (250ng/µl), mitfa:Cas9 (250 ng/µl); *zU6:sgptena* (23 ng/µl),

zU6:sgptenb (23ng/µl), and the tol2 plasmid (55ng/µl). For control electroporation without

generating melanoma, zebrafish were injected with *mitfa:tdTomato* (250ng/µl), *mitfa:*Cas9 (250

ng/µl), *zU6:*non-targeting (46ng/µl) and tol2 plasmid (55ng/µl). All fish were injected on the left

flank below the dorsal fin and electroporated with the BTX ECM 830 electroporator using 3mm

platinum Tweezertrodes (BTX Harvard Apparatus; #45-0487). Electroporator settings used: LV

Mode, 40V, 5 pulses, 60ms pulse length, and 1s pulse interval. Electroporated zebrafish were

screened for successful electroporation 7 days post-electroporation by tdTomato expression

using fluorescence microscopy and melanoma tracked by imaging once per week. All live

zebrafish imaging were performed with the Zeiss AxioZoom V16 fluorescence microscope.

sgRNA sequences for TEAZ listed below:

Nontargeting: 5'-AACCTACGGGCTACGATACG-3'

ptena: 5'-GAATAAGCGGAGGTACCAGG-3'

ptenb: 5'-GAGACAGTGCCTATGTTCAG-3'

Confocal imaging of zebrafish epidermis

Zebrafish with or without melanoma were anesthetized in tricaine (MS-222, Syndel) as

described above. Site of injection is visually identified by the presence of melanoma or the area

below the dorsal fin. Scales were removed with tweezers and fixed in 4% PFA in PBS (Santa

Cruz 281692) in a 96-well plate for 15 minutes. Fixed scales were washed three times with PBS

and permeabilized with 0.1% Triton-X 100 (Thermo Scientific 85111) in PBS, then blocked with

10% goat serum (Thermo Fisher 50062Z). Scales were incubated with 1:250 GFP polyclonal

antibody, Alexa Fluor 488 (Thermo Fisher A21311) overnight at 4°C. Next day, scales were

washed three times with PBS, incubated with 1:1000 Hoechst 33342 (Thermo Fisher H3570) for

1 hour and mounted onto slides with VECTASHIELD Vibrance Antifade Mounting Media (Vector

Laboratories H-1700). Samples were imaged on the Zeiss LSM880 inverted confocal

microscope and images were processed using FIJI v1.53.

Flow cytometry of adult zebrafish cells

Zebrafish were euthanized using ice-cold water. Melanoma and adjacent skin were dissected

from fish with melanoma and skin alone was dissected from below the dorsal fin. Subsequently,

samples were cut into 1mm strips using a clean scalpel and placed into 15 mL conical tubes

(Falcon 352099) with 3 ml of DPBS (Gibco 14190250) and 187.5µl of 2.5 mg/ml Liberase TL

(Roche 5401020001). Samples were incubated in dissociation solution at room temperature for

30 minutes on a shaker with gentle movement to prevent tissue from settling at the bottom of

the tube. At 15 minutes of incubation, a wide bore p1000 pipette tip (Thermo Scientific 2079G)

was used to gently pipette the sample up and down for 90 seconds. After 30 minutes, 250 µl of

FBS (Gemini Bio) was added to stop the enzymatic activity of Liberase TL and samples were

pipetted up and down using a wide bore p1000 pipette tip for 90 seconds. Dissociated cells

were then filtered through a 70 µm cell strainer (Falcon 352350) into a 50 mL conical tube

(Falcon 352098) placed on ice. Samples were centrifuged at 500g at 4°C for 5 minutes and

supernatant was removed by pipetting. The cell pellet was resuspended in 500 µl of PBS with

5% FBS and filtered again through 40 µm tip filters (Bel-Art H136800040) into 5 ml

polypropylene tubes (Falcon 352063). For subsequent FACS analysis, 0.5 µl of 1000x DAPI

(Sigma-Aldrich D9542) was added to each sample. Samples were FACS sorted (BD FACSAria)

at 4°C for GFP-positive keratinocytes and tdTomato-positive melanoma gated using

fluorophore-negative zebrafish controls.

Zebrafish tissue RNA extraction and real-time quantitative PCR (RT-qPCR)

FACS sorted zebrafish cells were deposited directly into 750 µl TRIzol LS Reagent (Invitrogen

10296010) in Eppendorf DNA LoBind Tubes (Eppendorf 022431021). After collection, samples

were snap-frozen using dry ice and stored at -80°C. RNA extraction was performed per TRIzol

LS manufacturer protocols. For precipitation of RNA, 10ug supplemental glycogen (Roche

10901393001) was used per sample to account for low cell numbers. Resulting RNA was

resuspended in Nuclease-free water (Fisher Scientific AM9937). 25ng RNA per sample was

transcribed to cDNA using Superscript III First-Strand Synthesis System (Invitrogen 18080051).

cDNA mix was diluted 1:10 with Nuclease free water for RT-qPCR using Power SYBR Green

PCR Master Mix (Applied Biosystems 4368708) and the Bio-Rad CFX384 Touch Real-Time

PCR System (Bio-Rad 1855484). Resulting Cq values were normalized to *hatn10* as previously

described.

qPCR primer sequences:

hatn10 fwd: 5'- TGAAGACAGCAGAAGTCAATG-3′

hatn10 rev: 5′-CAGTAAACATGTCAGGCTAAATAA-3′

mitfa fwd: 5'-GGCACCATCAGCTACAATGA-3'

mitfa rev: 5'-GAGACAGGGTGTTGTCCATAAG-3'

krt4 fwd: 5'-GGAGGTGTTTCCTCTGGTTATG-3'

krt4 rev: 5'-GAACCGAATCCTGATCCACTAC-3'

vim fwd: 5'-GGATATTGAGATCGCCACCTAC-3'

vim rev: 5'-GACTCTCGCAGGCTTAATGAT-3'

cdh2 fwd: 5'-GAGCCATCATCGCCATACTT-3'

cdh2 rev: 5'-CTTGGCCTGTCTCTCTTTATCC-3'

Re-analysis of zebrafish scRNA-sequencing data from Hunter et al.

Zebrafish scRNA-seq data from ref[27] was re-analyzed using R 4.2.0. and Seurat 4.3.0[47,48].

Cluster identities were maintained as published. Keratinocyte Module Scores were calculated

using the AddModuleScore function with default parameters using published gene lists.

Differential Gene Expression (DGE) analyses between clusters were performed using

FindMarkers. Differentially expressed gene lists were converted from zebrafish genes to human

orthologs using DIOPT as previously described[49,50]. GSEA analysis on differentially expressed

genes between keratinocyte clusters was performed using fgsea 1.22.0 and the Hallmark

pathways set from MSigDB[51,52].

Twist overexpression in zebrafish keratinocytes

Twist1a (ENSDART00000043595.5) and *twist1b* (ENSDART00000052927.7) were TOPO

cloned into the attL1-L2 Gateway pME vector and LR cloned into the pDestTol2pA2 vector

(Tol2Kit 394) with p5E-*krt4* promoter and p3E-polyA (Tol2Kit 302). To generate the zebrafish

melanoma model as previously described, one-cell stage Casper Triple zebrafish embryos

(*mitfa:BRAF^{V600E};p53-/-;mitfa-/-;mpv17-/-*) were injected with miniCoopR-tdTomato, *krt4-eGFP*,

either *krt4-twist1a* and *krt4-twist1b* for Twist overexpression condition or empty vector for control

condition, tol2 mRNA and phenyl red. Injections were performed three times on different days

with parents from the same clutch. Embryos were grown at standard conditions and sorted at 5

days post-injection for eGFP and tdTomato expression using the Zeiss AxioZoom V16

fluorescence microscope. eGFP+/tdTomato+ fish in the CTRL (n=135) and TWIST (n=118)

conditions were maintained to adulthood.

Zebrafish imaging and tumor-free survival tracking

Zebrafish were regularly monitored for melanoma formation and survival every 4 weeks,

beginning at 10 weeks post-fertilization. Melanoma formation was screened visually using the

Zeiss AxioZoom V16 fluorescence microscope under 20X magnification. Kaplan-Meier curves

and corresponding statistics were generated using GraphPad Prism 9. Statistical differences in

survival between conditions were determined by the Mantel-Cox log-rank test.

Histology of zebrafish samples

Zebrafish were euthanized in tricaine (MS222, Syndel). Each fish was dissected in three

sections consisting of head, body, and tail. Samples were placed in 4% PFA in PBS (Santa

Cruz 281692) for 72h on a shaker at 4°C, then paraffin embedded. Histology was performed by

HistoWiz Inc. (histowiz.com). Samples were processed, embedded in paraffin, and sectioned at

5μm. Immunohistochemistry was performed on a Bond Rx autostainer (Leica Biosystems) with

enzyme treatment (1:1000) using standard protocols. Sections were stained with H&E or IHC

with antibodies including BRAFV600E (ab228461) and GFP (ab183734). Bond Polymer Refine

Detection (Leica Biosystems) was used according to the manufacturer's protocol. After staining,

sections were dehydrated and film coverslipped using a TissueTek-Prisma and Coverslipper

(Sakura). Whole slide scanning (40x) was performed on an Aperio AT2 (Leica Biosystems).

Cell culture

Human melanoma lines A375, HS294T, and SKMEL2 were obtained from ATCC. Human

keratinocyte line HaCaT was obtained from AddexBio. All cells were routinely tested and

confirmed to be free from mycoplasma. Cells were maintained in a humidified incubator at 37°C

and 5% CO2. Cells were maintained in DMEM (Gibco 11965) supplemented with 10% FBS

(Gemini Bio) and split when confluent, approximately 2-3 times per week.

Twist overexpression in HaCaT

The HaCaT cell line was labeled with eBFP to allow for identification during co-culture. 293T

(ATCC) was transfected with the pLV-Azurite plasmid (Addgene 36086) with pMD2.5 (Addgene

12259) and psPAX2 (Addgene 12260) using Invitrogen Lipofectamine 3000 Transfection

Reagent (Invitrogen L3000015) according to manufacturer protocol. HaCaTs were infected with

lentivirus containing *CMV:eBFP* and selected for eBFP positivity using ampicillin and FACS.

Subsequently, HaCaT-eBFP was infected with lentivirus containing *CMV:TWIST1* (Horizon

Precision LentiORF Human TWIST1 OHS5898-202622685) or *CMV:*empty control created by

removing ORF of *CMV:TWIST1* plasmid. HaCaT line overexpressing *TWIST1* was labeled as

HaCaT-TWIST and HaCaT line with empty vector was labeled as HaCaT-CTRL. HaCaT lines

were subsequently sorted for nuclear eGFP expression present in the plasmid as part of the

Precision LentiORF system. HaCaT-CTRL and HaCaT-TWIST were cultured as previously

described.

Western blot

Cells were washed with DPBS (Gibco 14190250) and lysed in RIPA buffer (Thermo Scientific

89901) with the addition of protease and phosphatase inhibitors (Thermo Scientific 78440).

Lysates were centrifuged at 13,000g at 4 °C and quantified using the Pierce BCA Protein Assay

Kit (Pierce 23227). Samples were reduced with the laemmli SDS-sample buffer (Boston

BioProducts BP111R) and boiled for 10 minutes. For gel electrophoresis, samples were loaded

into 4-15% precast protein gels (Bio-Rad 4561084), then transferred to 0.2µm nitrocellulose

membranes (Bio-Rad 1704158). Membranes were washed in TBST and blocked with 5% milk in

TBST (Boston BioProducts P1400) for 1 hour at RT. Membranes were washed and incubated

with primary antibodies overnight. Antibodies used includes Twist (Abcam ab50887) and beta-

actin (CST 3700S). On the next day, membranes were washed with TBST and incubated with

appropriate secondary antibodies for 1 hour. Blots were incubated with the Immobilon Western

Chemiluminescent HRP Substrate (Millipore WBKLS0500) and imaged with the Amersham

ImageQuant 800.

Immunofluorescence

Cells were cultured on chamber slides (Thermo Scientific 154739) overnight at standard cell

culture conditions. Culture media was washed with DPBS (Gibco 14190250) and fixed with 2%

PFA in PBS (Santa Cruz 281692) for 15 minutes at RT. Cells were washed with DPBS and

permeabilized with 0.1% Triton-X 100 (Thermo Scientific 85111) in PBS, then blocked with 10%

goat serum (Thermo Fisher 50062Z) for 1 hour at RT. Primary antibodies used include Twist

(Abcam ab50887). Cells were incubated with primary antibody in 10% goat serum overnight at

4 °C and washed with DPBS the next day, before incubation with 1:1000 Hoechst 33342

(Thermo Fisher H3570) for 1 hour and mounted with VECTASHIELD PLUS Antifade Mounting

Medium (Vector Laboratories H1900). Slides were imaged on the Zeiss LSM880 inverted

confocal microscope and images were processed using FIJI v1.53.

Melanoma infiltration assay

RFP-labeled melanoma cell lines, including A375, SKMEL2, HS294T, were plated on poly-l-

lysine coated round glass coverslips (Corning 354085) placed in 24-well plates at 150-200k

cells per coverslip. HaCaT cell lines were plated in 6-well plates at 250-300k cells per well. Cells

were allowed to attach overnight and the coverslip containing melanoma cells is transferred to

6-wells containing HaCaT cell lines using tweezers. All coverslips were placed in the center of

the well. KC-melanoma co-cultures were incubated in standard cell culture conditions for 24

hours. Co-cultures were imaged by fluorescence microscopy at 4 locations of each coverslip:

top, right, bottom and left, to capture variations in melanoma cell infiltration into KC lines. FIJI

v1.53 was used to count the number of infiltrating melanoma cells per image and average

infiltrating melanoma cells were calculated per well. All experiments were performed in 3 sets,

with 3 replicates per set per condition. Average infiltrating cell numbers per well were

normalized to the average infiltrating cell number per well in the HaCaT-CTRL condition.

scRNA-sequencing analysis of zebrafish Melanoma

Six zebrafish, three each from CTRL and TWIST conditions at 26 weeks post-injection were

selected for scRNA-sequencing of melanoma tumors. To account for set differences, one fish

from each of three injection sets were chosen in each condition. Melanoma and adjacent skin

were dissected from the fish, then enzymatically and mechanically dissociated into single cell

solutions as described above. The samples were FACS sorted for GFP and RFP positivity,

corresponding to eGFP expressed by keratinocytes and tdTomato expressed by melanoma.

The sorted cells were placed in (Gibco 11965) supplemented with 10% FBS (Gemini Bio) and

1% penicillin-streptomycin-glutamine. To enrich for keratinocytes, sorted keratinocytes and

melanoma cells from each fish was recombined at a 7:3 KC:melanoma ratio. Sorted cells were

pelleted and resuspended in DPBS + 0.1% BSA. Samples were also combined based on their

genetic perturbation condition. Droplet-based scRNA-seq was performed using the Chromium

Single Cell 3′ Library and Gel Bead Kit v3 (10X Genomics) and Chromium Single Cell 3′ Chip G

(10X Genomics). 10,000 cells were targeted for encapsulation. GEM generation and library

preparation was performed according to kit instructions. Libraries were sequenced on a

NovaSeq S4 flow cell. Resulting reads were aligned to the GRCz11 reference genome with the

addition of eGFP and tdTomato sequences using CellRanger v5.0.1 (10x Genomics). scRNA-

sequencing analysis was performed as detailed above. In addition, melanoma cells were scored

using AddModuleScore to assess their enrichment of genes associated with the four main

melanoma cell states and intermediate states[28]. The highest scoring gene module for each cell

was annotated as its cell state. CellChat[29] was used to analyze cell-cell communication between

KC and melanoma clusters using its zebrafish L-R database.

Statistics and reproducibility

Statistical analysis and figures were generated by GraphPad Prism 9, R Studio 4.2.0 and

Biorender.com. Image processing were performed in FIJI v1.53. Statistical tests are described

in figure legends and methods. Experiments were repeated at least three times unless

otherwise noted. All animal and cell experiments were performed with a reasonable number of

replicates by power calculations or feasibility of the experimental method.

509   Acknowledgements

We thank the MSKCC Single Cell Analytics Innovation Lab for their consult with single cell RNA-

sequencing, the MSKCC Molecular Cytology Core Facility for assistance with imaging, the

MSKCC Aquatics Core Facility for zebrafish care and maintenance, and the MSKCC Flow

Cytometry Core Facility for assistance with FACS. We also thank the members of the White Lab

for helpful discussions. YM was supported by a Medical Scientist Training Program (MSTP)

grant from the National Institute of General Medical Sciences of the National Institutes of Health

(NIH) under award number T32GM152349 to the Weill Cornell/Rockefeller/Sloan Kettering Tri-

Institutional MD-PhD Program, the Kirschstein-NRSA predoctoral fellowship (F30) from the

National Cancer Institute (NCI) of the NIH under award number F30CA265124, and the Barbara

and Stephen Friedman Pre-doctoral Fellowship from MSKCC. MT was support by the Marie-

Josée Kravis Women in Science Endeavor Postdoctoral Fellowship and the Experimental

Immuno-Oncology Scholars Fellowship. MVH was funded by a K99/R00 Pathway to

Independence Award from the NCI(1K99CA266931), the Scholarship for the Next Generation of

Scientists from the Cancer Research Society, and a postdoctoral fellowship from the Canadian

Institutes of Health Research. EM was supported by the NIH Individual Predoctoral to

Postdoctoral Fellow Transition Award under award number 5K00CA223016. JMW was

supported by the F30 fellowship from the NCI of the NIH under award number F30CA236442, a

predoctoral fellowship (T32) by the Cell and Developmental Biology Program at Weill Cornell

Graduate School supported by the NIH under award number T32GM008539, and a MSTP grant

from the NIH under award number T32GM007739. RMW was funded through the NIH/NCI

Cancer Center Support Grant P30 CA008748, the Melanoma Research Alliance, NIH Research

Program Grants R01CA229215 and R01CA238317, NIH Director's New Innovator Award

DP2CA186572, The American Cancer Society, The Alan and Sandra Gerry Metastasis

Research Initiative at the Memorial Sloan Kettering Cancer Center, The Harry J. Lloyd

Foundation, and Oxford Ludwig Cancer Research.

References

1. Lee, J. T. & Herlyn, M. Microenvironmental influences in melanoma progression. *Journal of*
Cellular Biochemistry **101**, 862–872 (2007).

2. Fitzpatrick, T. B. & Breathnach, A. S. The Epidermal Melanin Unit System. *Dermatologische*
Wochenschrift **147**, 481–9 (1963).

3. Kunisada, T. *et al.* Transgene expression of steel factor in the basal layer of epidermis
promotes survival, proliferation, differentiation and migration of melanocyte precursors.
Development **125**, 2915–2923 (1998).

4. Tanimura, S. *et al.* Hair Follicle Stem Cells Provide a Functional Niche for Melanocyte Stem
Cells. *Cell Stem Cell* **8**, 177–187 (2011).

5. Tang, A. *et al.* E-cadherin is the major mediator of human melanocyte adhesion to
keratinocytes in vitro. *Journal of Cell Science* **107**, 983–992 (1994).

6. Imokawa, G., Yada, Y. & Miyagishi, M. Endothelins secreted from human keratinocytes are
intrinsic mitogens for human melanocytes. *Journal of Biological Chemistry* **267**, 24675–
24680 (1992).

7. Valyi-Nagy, I. T. *et al.* Undifferentiated keratinocytes control growth, morphology, and
antigen expression of normal melanocytes through cell-cell contact. *Laboratory Investigation*
69, 152–159 (1993).

8.   Shih, I. M., Elder, D. E., Hsu, M. Y. & Herlyn, M. Regulation of Mel-CAM/MUC18 expression

on melanocytes of different stages of tumor progression by normal keratinocytes. *Am J*

Pathol **145**, 837–845 (1994).

9.   Hara, M., Yaar, M. & Gilchrest, B. A. Endothelin-1 of Keratinocyte Origin Is a Mediator of

Melanocyte Dendricity. *Journal of Investigative Dermatology* **105**, 744–748 (1995).

10. Hsu, M. Y., Wheelock, M. J., Johnson, K. R. & Herlyn, M. Shifts in cadherin profiles between

human normal melanocytes and melanomas. *J Investig Dermatol Symp Proc* **1**, 188–194

(1996).

11. Hsu, M. Y. *et al.* E-cadherin expression in melanoma cells restores keratinocyte-mediated

growth control and down-regulates expression of invasion-related adhesion receptors.

American Journal of Pathology **156**, 1515–1525 (2000).

12. Li, G., Satyamoorthy, K. & Herlyn, M. N-cadherin-mediated intercellular interactions promote

survival and migration of melanoma cells. *Cancer Research* **61**, 3819–3825 (2001).

13. Fukunaga-Kalabis, M. *et al.* CCN3 controls 3D spatial localization of melanocytes in the

human skin through DDR1. *Journal of Cell Biology* **175**, 563–569 (2006).

14. Fukunaga-Kalabis, M. *et al.* Downregulation of CCN3 expression as a potential mechanism

for melanoma progression. *Oncogene* **27**, 2552–2560 (2008).

15. Mescher, M. *et al.* The epidermal polarity protein Par3 is a non-cell autonomous suppressor

of malignant melanoma. *Journal of Experimental Medicine* **214**, 339–358 (2017).

16. Jamal, S. & Schneider, R. J. UV-induction of keratinocyte endothelin-1 downregulates E-

cadherin in melanocytes and melanoma cells. *J Clin Invest* **110**, 443–452 (2002).

17. Kim, I. S. *et al.* Microenvironment-derived factors driving metastatic plasticity in melanoma.

Nature Communications **8**, 14343 (2017).

18. Tagore, M. *et al.* GABA Regulates Electrical Activity and Tumor Initiation in Melanoma.

Cancer Discovery **13**, 2270–2291 (2023).

19. Callahan, S. J. *et al.* Cancer modeling by Transgene Electroporation in Adult Zebrafish

(TEAZ). *Disease Models & Mechanisms* **11**, dmm034561 (2018).

20. Kaufman, C. K. *et al.* A zebrafish melanoma model reveals emergence of neural crest

identity during melanoma initiation. *Science* **351**, aad2197–aad2197 (2016).

21. Li, Q. & Uitto, J. Zebrafish as a Model System to Study Skin Biology and Pathology. *Journal*

of Investigative Dermatology **134**, 1–6 (2014).

22. Gong, Z. *et al.* Green fluorescent protein expression in germ-line transmitted transgenic

zebrafish under a stratified epithelial promoter from keratin8. *Developmental Dynamics* **223**,

204–215 (2002).

23. Montal, E., Suresh, S., Ma, Y., Tagore, M. M. & White, R. M. Cancer Modeling by

Transgene Electroporation in Adult Zebrafish (TEAZ). in *Zebrafish: Methods and Protocols*

(eds. Amatruda, J. F., Houart, C., Kawakami, K. & Poss, K. D.) 83–97 (Springer US, New

York, NY, 2024). doi:10.1007/978-1-0716-3401-1_5.

24. Haensel, D. & Dai, X. Epithelial-to-mesenchymal transition in cutaneous wound healing:

Where we are and where we are heading. *Developmental Dynamics* **247**, 473–480 (2018).

25. Leopold, P. L., Vincent, J. & Wang, H. A comparison of epithelial-to-mesenchymal transition

and re-epithelialization. *Seminars in Cancer Biology* **22**, 471–483 (2012).

26. Moreno-Bueno, G. *et al.* The morphological and molecular features of the epithelial-to-

mesenchymal transition. *Nature Protocols* **4**, 1591–1613 (2009).

27. Hunter, M. V., Moncada, R., Weiss, J. M., Yanai, I. & White, R. M. Spatially resolved

transcriptomics reveals the architecture of the tumor-microenvironment interface. *Nature*

Communications 2021 12:1 **12**, 1–16 (2021).

28. Tsoi, J. *et al.* Multi-stage Differentiation Defines Melanoma Subtypes with Differential

Vulnerability to Drug-Induced Iron-Dependent Oxidative Stress. *Cancer Cell* **33**, 890-904.e5

(2018).

29. Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nature*
communications **12**, 1088 (2021).

30. Eom, D. S., Patterson, L. B., Bostic, R. R. & Parichy, D. M. Immunoglobulin superfamily
receptor Junctional adhesion molecule 3 (Jam3) requirement for melanophore survival and
patterning during formation of zebrafish stripes. *Developmental Biology* **476**, 314–327
(2021).

31. Arcangeli, M. L. *et al.* The Junctional Adhesion Molecule-B regulates JAM-C-dependent
melanoma cell metastasis. *FEBS Letters* **586**, 4046–4051 (2012).

32. Langer, H. F. *et al.* A novel function of junctional adhesion molecule-C in mediating
melanoma cell metastasis. *Cancer Research* **71**, 4096–4105 (2011).

33. Tissue expression of F11R - Staining in skin. *The Human Protein Atlas*
https://www.proteinatlas.org/ENSG00000158769-F11R/tissue/skin.

34. Tissue expression of JAM3 - Staining in skin. *The Human Protein Atlas*
https://www.proteinatlas.org/ENSG00000166086-JAM3/tissue/skin.

35. Koike, Y., Yozaki, M., Utani, A. & Murota, H. Fibroblast growth factor 2 accelerates the
epithelial–mesenchymal transition in keratinocytes during wound healing process. *Sci Rep*
10, 18545 (2020).

36. Halaban, R., Kwon, B. S., Ghosh, S., Delli Bovi, P. & Baird, A. bFGF as an autocrine growth
factor for human melanomas. *Oncogene Res* **3**, 177–186 (1988).

37. Hoek, K. S. *et al.* Metastatic potential of melanomas defined by specific gene expression
profiles with no BRAF signature. *Pigment Cell Research* **19**, 290–302 (2006).

38. Hu, F. *et al.* Sortilin-Mediated Endocytosis Determines Levels of the Frontotemporal
Dementia Protein, Progranulin. *Neuron* **68**, 654–667 (2010).

39. Daniel, R., He, Z., Carmichael, K. P., Halper, J. & Bateman, A. Cellular Localization of Gene
Expression for Progranulin. *J Histochem Cytochem.* **48**, 999–1009 (2000).

40. Shoyab, M., McDonald, V. L., Byles, C., Todaro, G. J. & Plowman, G. D. Epithelins 1 and 2:

isolation and characterization of two cysteine-rich growth-modulating proteins. *Proceedings*

of the National Academy of Sciences **87**, 7912–7916 (1990).

41. He, Z., Ong, C. H. P., Halper, J. & Bateman, A. Progranulin is a mediator of the wound

response. *Nature Medicine* **9**, 225–229 (2003).

42. Tanimoto, R. *et al.* The perlecan-interacting growth factor progranulin regulates

ubiquitination, sorting, and lysosomal degradation of sortilin. *Matrix Biol* **64**, 27–39 (2017).

43. Truzzi, F. *et al.* Neurotrophins and Their Receptors Stimulate Melanoma Cell Proliferation

and Migration. *Journal of Investigative Dermatology* **128**, 2031–2040 (2008).

44. White, R. M. *et al.* Transparent Adult Zebrafish as a Tool for In Vivo Transplantation

Analysis. *Cell Stem Cell* **2**, 183–189 (2008).

45. Kwan, K. M. *et al.* The Tol2kit: A multisite gateway-based construction kit for Tol2

transposon transgenesis constructs. *Developmental Dynamics* **236**, 3088–3099 (2007).

46. Montal, E., Lumaquin, D., Ma, Y., Suresh, S. & White, R. M. Modeling the effects of genetic-

and diet-induced obesity on melanoma progression in zebrafish. *Disease Models \&*

Mechanisms **16**, dmm049671 (2023).

47. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell

transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol*

36, 411–420 (2018).

48. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29

(2021).

49. Hu, Y. *et al.* An integrative approach to ortholog prediction for disease-focused and other

functional studies. *BMC Bioinformatics* **12**, 357 (2011).

50. Campbell, N. R. *et al.* Cooperation between melanoma cell states promotes metastasis

through heterotypic cluster formation. *Developmental Cell* **56**, 2808-2825.e10 (2021).

51. Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for

interpreting genome-wide expression profiles. *Proceedings of the National Academy of*

Sciences of the United States of America **102**, 15545–15550 (2005).

52. Liberzon, A. *et al.* The Molecular Signatures Database Hallmark Gene Set Collection. *Cell*

Systems **1**, 417–425 (2015).

Figure 1

Figure 1. Keratinocytes in the melanoma microenvironment undergo EMT-like changes.

(A) Generation of transparent zebrafish with GFP-labeling of keratinocytes. Casper Triples (*mitfa-/-;mitfa:*BRAFV600E;*tp53-/-;mpv17-/-*) were injected with the Tol2Kit 394 vector containing a *krt4*:eGFP cassette and tol2 mRNA. Brightfield shows a transparent zebrafish while fluorescence imaging shows eGFP-labeling of keratinocytes. (Scale bar = 5mm)

(B) Schematic of TEAZ (Transgene Electroporation in Adult Zebrafish). Plasmid mix containing miniCoopR:tdTomato, *mitfa:*Cas9, *zU6:sgptena*, *zU6:sgptenb*, and the tol2 plasmid was injected superficially in the flank of the zebrafish. Electroporation of the injection site results in rescue of melanocyte precursors and the generation of a localized melanoma that could be analyzed by microscopy and FACS.

(C) Brightfield and immunofluorescence of zebrafish 8-weeks post-TEAZ with localized and fluorescently labeled melanoma. (Scale bar = 5mm)

(D) Immunofluorescence imaging of TEAZ region after 8-weeks, comparing empty vector control vs. miniCoopR:tdT conditions, with yellow dotted circles indicating general area of dissection for FACS. (Scale bar = 1mm)

(E) Confocal imaging of zebrafish epidermis. Normal epidermis of Tg(*krt4:*eGFP) Casper Triple post-TEAZ with empty vector control shows eGFP-labeled, polygonal shaped keratinocytes regularly connected while epidermis with melanoma generated with *miniCoopR*-driven melanocyte rescue shows disrupted epidermis and irregularly shaped keratinocytes. (Scale bar = 50um)

(F) qPCR of FACS sorted zebrafish epidermis with or without melanoma. tdTomato-labeled melanoma cells, eGFP-labeled keratinocytes and non-fluorescently labeled TME cells were isolated by dissection (as indicated in G) and FACS. Comparison of *mitfa* and *krt4* expression of samples normalized to non-fluorescent cells, either 'TME-Other' in tumor

samples or 'Other' in non-tumor samples, shows enrichment of *mitfa* in melanoma

sample and *krt4* in keratinocyte sample. ns is non-significant, * is p ≤ 0.05, *** is p ≤

0.001, **** is p ≤ 0.0001 by Tukey's multiple comparisons test.

(G) Comparison of the EMT-markers *vim* and *cdh2* shows enrichment in TME keratinocytes

vs. keratinocytes from epidermis without melanoma. * is p ≤ 0.05 by Welch's t-test.

(H) Schematic of keratinocyte-melanoma co-culture experiment. HaCaTs were cultured in

monoculture or co-culture with A375 melanoma cells in triplicates for 21 days, followed

by FACS isolation of keratinocytes for RNA-sequencing comparing co-culture vs.

monoculture keratinocytes.

(I) Top 5 enriched Hallmark pathways in HaCaTs co-cultured with A375 melanoma cells

compared with HaCaTs in monoculture.

(J) Normalized counts of EMT biomarkers vimentin (*VIM*) and N-cadherin (*CDH2*).

Figure 2

Figure 2. Zebrafish scRNA-sequencing shows upregulation of EMT-TFs in tumor-associated keratinocytes.

(A) Schematic of scRNA-sequencing experiment. Embryo injection with miniCoopR:eGFP plasmid and tol2 mRNA in Casper Triples (*mitfa-/-;mitfa:*BRAFV600E;*tp53-/-;mpv17-/-*) results in melanocyte rescue and subsequent melanoma formation. Melanoma was dissected and dissociated to single cell suspension for FACS isolation of eGFP+ melanoma cells and non-fluorescent TME cells for single cell RNA-sequencing.

(B) Dimensionality reduction and subsequent analysis with zebrafish keratinocyte gene module scoring highlights two keratinocyte clusters.

(C) Top 6 GSEA Hallmark pathways enriched in comparison between keratinocyte clusters.

(D) Hallmark EMT pathway enrichment in keratinocyte clusters.

(E) Expression of EMT-transcription factors Snail (*snai1a*), Slug (*snai2*), and Twist1 (*twist1a*) in TAK vs. NKC, Melanoma, and Other TME cells. ns is no significance, * is p ≤ 0.05, **** is p ≤ 0.0001.

Figure 3

Figure 3. Overexpression of twist1a/b results in improved survival of fish with melanoma.

(A) Schematic of zebrafish melanoma model with labeling and perturbation of keratinocytes. *Twist1a* and *twist1b* are overexpressed under the keratinocyte-specific *krt4* promoter in the TWIST condition and an empty vector control was used in the CTRL condition. Plasmid mix containing miniCoopR-tdTomato, *krt4*:eGFP, either empty vector or *krt4:twist1a/b*, with tol2 mRNA were injected into Casper Triples (*mitfa-/-;mitfa:*BRAFV600E;*tp53-/-;mpv17-/-*). Fish were sorted at 5 days for eGFP and tdTomato positivity as marker of successful keratinocyte labeling and melanocyte rescue.

(B) Tumor-free survival and overall survival of A. ns is no significance, ** is p ≤ 0.01 by Log-rank (Mantel-Cox) test.

(C) Sample images of zebrafish with melanoma at 26 weeks post-injection. Melanomas are pigmented in brightfield images. Keratinocytes are labeled by eGFP and melanoma are labeled by tdTomato in fluorescence images. Scale bar = 5mm.

(D) H&E and IHC of cross-sections through zebrafish body and melanoma. Dotted yellow line demarcates border of body. Percent of tumor in body is calculated as tumor area within body border divided by total tumor area. Scale bar = 500um.

Figure 4

Figure 4. Zebrafish findings are recapitulated in human cell lines.

(A) Schematic of coverslip cell infiltration assay. Melanoma cells are plated on a coverslip and allowed to attach overnight. The coverslip is then transferred into a well of keratinocytes to assess melanoma infiltration into keratinocytes.

(B) Generation of a HaCaT cell line overexpressing *TWIST1*. HaCaT-BFP was infected with lentivirus containing cassette with nuclear localized GFP and *CMV* driven *TWIST1* or no ORF. Infected cell lines were allowed to grow for a week before sorting for nuclear GFP as a marker of successful integration.

(C) Western blot for Twist expression in HaCaT-CTRL and HaCaT-TWIST.

(D) Immunofluorescence imaging for Twist localization in HaCaT-CTRL and HaCaT-TWIST. TWIST staining is pseudo-colored in magenta, DAPI in blue, phalloidin in white. Scale bar = 50um.

(E) Immunofluorescence imaging of coverslip cell infiltration assay after 20 hours with HS294T-tdT (orange) melanoma cells in co-culture with either HaCaT-CTRL or HaCaT-TWIST (cyan). Scale bar = 500um.

(F) Quantification of E. Infiltrating HS294T melanoma cells from each image were counted and averaged across four images per well. Resulting cell counts were normalized to average cell counts of HaCaT-CTRL from each set. N = 9, 3 sets, 3 replicates/wells per set. * is p ≤ 0.05 by t-test.

(G) Immunofluorescence imaging of coverslip cell infiltration assay after 20 hours with SKMEL2-tdT (orange) melanoma cells in co-culture with either HaCaT-CTRL or HaCaT-TWIST (cyan). Scale bar = 500um.

(H) Quantification of G. Infiltrating SKMEL2 melanoma cells from each image were counted and averaged across four images per well. Resulting cell counts were normalized to

average cell counts of HaCaT-CTRL from each set. N = 9, 3 sets, 3 replicates/wells per set. ** is p ≤ 0.01 by t-test.

Figure 5

Figure 5. scRNA-sequencing shows unique keratinocyte-melanoma communication with twist1a/b overexpression in keratinocytes.

(A) Schematic of scRNA-sequencing protocol. Melanoma and surrounding tissue were dissected from 26-weeks old zebrafish from either CTRL or TWIST conditions as shown in Figure 3. Samples were dissociated to single cell suspensions for FACS isolation of keratinocytes (GFP) and melanoma (tdTomato). Keratinocytes and melanoma were recombined per condition at a ratio of 7:3 for enrichment of keratinocytes for scRNA-sequencing.

(B) UMAP dimensional reduction and feature plots of scRNA-sequencing dataset. CTRL and TWIST samples were sequenced, and cell types were identified using eGFP+ for keratinocytes and tdTomato+ for melanoma, after which the datasets were integrated.

(C) UMAP showing cell type assignments of each cluster within either CTRL or TWIST conditions. Arrow indicates a TWIST-melanoma population clustering separately from other melanoma clusters.

(D) UMAP highlighting keratinocyte clusters, with a Tumor-Associated Keratinocyte (TAK) cluster, a Normal Keratinocyte Cluster (NKC), and a Twist-High cluster unique to the TWIST condition.

(E) Melanoma cell state analysis of the melanoma cluster unique to the Twist condition indicated by arrow in Figure 5C.

(F) Schematic overview of CellChat analysis. In CTRL condition, we analyzed Ligand-Receptor pairs with NKC as sender and CTRL-Melanoma as receiver. In TWIST condition, we analyzed L-R pairs with both NKC and Twist-High as sender and TWIST-Melanoma as receiver.

(G) CellChat analysis results. L-R pairs shown at p<0.01, with color scale indicating communication probability of L-R pair.