Calcium tunneling through the ER and transfer to other organelles for optimal signaling in *Toxoplasma gondii*.

Zhu-Hong Li[1], Beejan Asady[1,2], Le Chang[1], Miryam Andrea Hortua Triana[1], Catherine Li[1], Isabelle Coppens[2] and Silvia N.J. Moreno[1,3*]

[1]Center for Tropical and Emerging Global Diseases, Department of Computes Science, University of Georgia and [3]Department of Cellular Biology, University of Georgia, Athens, Georgia 30602
[2]Department of Molecular Microbiology and Immunology, Johns Hopkins Bloomberg School of Public Heath, Baltimore, MD 21205

Running title: Intracellular Ca^{2+} stores of *T. gondii*

[*]To whom correspondence should be addressed: Silvia N. J. Moreno, Department of Cellular Biology and Center for Tropical and Emerging Global Disease, 350A Paul D. Coverdell Center, University of Georgia, Athens, GA 30602. Tel.: 706-542-4736; Fax: 706-542-9493; E-mail: smoreno@uga.edu

Keywords: Calcium, endoplasmic reticulum, membrane contact sites, Toxoplasma gondii, SERCA-Ca^{2+}-ATPase, calcium tunneling

ABSTRACT

Ca^{2+} signaling in cells begins with the opening of Ca^{2+} channels in either the plasma membrane (PM) or the endoplasmic reticulum (ER) and results in a dramatic increase in the physiologically low (<100 nM) cytosolic Ca^{2+} level. The temporal and spatial Ca^{2+} levels are well regulated to enable precise and specific activation of critical biological processes. Ca^{2+} signaling regulates pathogenic features of apicomplexan parasites like *Toxoplasma gondii* which infects approximately one-third of the world's population. *T. gondii* relies on Ca^{2+} signals to stimulate traits of its infection cycle and several Ca^{2+} signaling elements play essential roles in its parasitic cycle. Active egress, an essential step for the infection cycle of *T. gondii* is preceded by a large increase in cytosolic Ca^{2+} most likely by release from intracellular stores. Intracellular parasites take up Ca^{2+} from the host cell during host Ca^{2+} signaling events to replenish intracellular stores. In this work, we investigated the mechanism by which intracellular stores are replenished with Ca^{2+} and demonstrated a central role for the SERCA-Ca^{2+}-ATPase to keep not only the ER filled with Ca^{2+} but also acidic stores. We also show mitochondrial Ca^{2+} uptake, by transfer of Ca^{2+} from the ER most likely through membrane contact sites. We propose a central role for the ER in tunneling of calcium from the extracellular milieu through the ER to other organelles.

HIGHLIGHTS

- The *T. gondii* ER efficiently takes up Ca^{2+} that enters the cytosol from the extracellular milieu.
- Filling of acidic stores in *T. gondii* appears to be dependent on the filling of the ER
- The mitochondrion of *T. gondii* has no direct access to extracellular calcium but is able to take up Ca^{2+} by transfer from the ER and/or acidic stores

INTRODUCTION

Toxoplasma gondii is an intracellular parasite from the Apicomplexan Phylum that infects
approximately one third of the world population (Weiss and Dubey, 2009). During the initial
infection with *T. gondii*, the parasite engages in multiple rounds of a lytic cycle, which consists of
invasion of host cells, replication inside a parasitophorous vacuole (PV), exit from the host cell
causing its lysis followed by invasion of a new host cell (Black and Boothroyd, 2000; Blader *et*
al., 2015). Cytosolic Ca^{2+} ($[Ca^{2+}]_c$) fluctuations precede the activation of several key steps of the
T. gondii lytic cycle like motility, attachment, invasion, and egress ($[Ca^{2+}]_c$) (Lourido and Moreno,
2015; Hortua Triana *et al.*, 2018). Egress from the host cell is an essential step for the infection
cycle of *T. gondii* (Bisio and Soldati-Favre, 2019) and it was shown that it is preceded by a
cytosolic Ca^{2+} increase (Endo *et al.*, 1982; Borges-Pereira *et al.*, 2015). Extracellular Ca^{2+} entry
was demonstrated in extracellular (Pace *et al.*, 2014; Hortua Triana *et al.*, 2024) and intracellular
replicating tachyzoites (Vella *et al.*, 2021). This activity was highly regulated and recent work
from our lab showed that a TRP-like activity was involved (Marquez-Nogueras *et al.*, 2021).

Ca^{2+} signaling is part of the signaling pathways that regulate a large number of cellular functions
(Clapham, 2007). All cells express a variety of channels, transporters, and Ca^{2+} pumps, located at
the plasma membrane (PM) and/or intracellular organelles (endoplasmic reticulum (ER), acidic
stores, and mitochondria) that regulate/control the concentration of cytosolic Ca^{2+} ($[Ca^{2+}]_c$). A
high cytosolic Ca^{2+} concentration for sustained lengths of time, is toxic to cells and may result in
their death (Bootman and Bultynck, 2020).

In *T. gondii*, both Ca^{2+} entry through the plasma membrane and influx from intracellular stores
like the ER may initiate a cascade of signaling events important for the stimulation of the biological
steps of the parasite lytic cycle (Lourido and Moreno, 2015; Hortua Triana *et al.*, 2018). Ca^{2+}
oscillations were observed in motile parasites loaded with fluorescent Ca^{2+} indicators, (Lovett and
Sibley, 2003) as well as expressing Genetically Encoded Calcium Indicators (GECIs) (Borges-
Pereira *et al.*, 2015). The significance of Ca^{2+} signals during all stages of the lytic cycle has been
demonstrated, but little is known about the mechanism by which intracellular stores contribute to
cytosolic Ca^{2+} signals and downstream regulation.

The ER, an exclusive organellar feature of eukaryotic cells, is the main store for Ca^{2+} in most
eukaryotic cells and it is widely accepted that: the ER is functionally heterogeneous with Ca^{2+}
binding proteins and Ca^{2+} pumps and channels having a nonuniform distribution resulting in the
presence of distinct subdomains within the organelle (Papp *et al.*, 2003); the ER facilitates Ca^{2+}
tunneling through its lumen as a mechanism of delivering Ca^{2+} to targeted sites without activating
inappropriate processes in the cell cytosol (Petersen *et al.*, 2017); and the ER is ubiquitously
distributed and is in close contact with all cellular organelles and the plasma membrane (PM)
(Spang, 2018). Over the past decade a new paradigm has emerged that seeks to decipher how
subcellular organelles communicate with each other in order to coordinate activities and efficiently
distribute ions and lipids within the cell. Numerous observations have highlighted the presence of
tight, stable and yet non-fusogenic associations between organellar membranes which have since
become known as membrane contact sites (MCSs) (Phillips and Voeltz, 2016).

The secretory pathway of *T. gondii* is organized in a highly polarized manner with the ER being an extension of the nuclear envelope (Hager *et al.*, 1999; Tomavo *et al.*, 2013). The ER at the apical surface of the nuclear envelope continues with the Golgi stacks and then the secretory organelles, micronemes and rhoptries, which are unique to the apicomplexan phylum (Hager *et al.*, 1999). These organelles perform important functions relevant to a successful lytic cycle like host cell attachment, invasion, and establishment of the parasitophorous vacuole (PV). Cytoplasmic Ca^{2+} increases, due to efflux from the ER, have been reported to initiate responses like microneme secretion (Carruthers and Sibley, 1999; Nagamune *et al.*, 2007b), conoid extrusion (Del Carmen *et al.*, 2009), invasion (Vieira and Moreno, 2000; Lovett and Sibley, 2003) and egress (Arrizabalaga and Boothroyd, 2004). These responses require precise spatiotemporal control of diverse targets and suggests the presence of distinct systems to deliver Ca^{2+} to specific locations rather than allowing global increases, which would activate unnecessary and potentially detrimental signaling events (Huet and Moreno, 2023).

In order to concentrate Ca^{2+} ions, the ER utilizes a SERCA-Ca^{2+}-ATPase, a transmembrane P-type ATPase, that couples ATP hydrolysis to the transport of ions across biological membranes and against a concentration gradient. SERCA pumps can be inhibited by various inhibitors including the very potent and highly specific thapsigargin (TG) (Thastrup *et al.*, 1990; Sagara and Inesi, 1991).

T. gondii expresses a SERCA Ca^{2+}-ATPase (TgSERCA), which possess SERCA conserved domains and Ca^{2+} binding sites and ATPase residues (Nagamune *et al.*, 2007a). The function of TgSERCA was determined by rescue experiments of yeast cells defective in Ca^{2+}-ATPases and by its specific inhibition by TG (Nagamune *et al.*, 2007a). TgSERCA was mainly localized to the ER of *T. gondii* but also showed a distinct distribution in extracellular parasites, where the protein was partially found in ER vesicles in the apical region near micronemes (Nagamune *et al.*, 2007a). This distribution pattern was different to the one obtained with the transient transfection of GFP-HDEL (an ER marker), which was retained near the nuclear envelope, suggesting an uneven distribution of ER markers in extracellular parasites. The authors suggested that this distribution to the apical end may be important for rapid release and effective recovery of cytosolic Ca^{2+}, events that likely govern both motility and microneme secretion (Nagamune *et al.*, 2007a; Nagamune *et al.*, 2007b).

In this work we characterize the mechanism by which the ER efficiently takes up Ca^{2+} promptly after the ion enters the cytosol from the extracellular milieu. We also characterize the role of the ER in transferring Ca^{2+} to other stores. We propose a model in which the ER is the distribution center for Ca^{2+} and due to the high Ca^{2+} affinity of the SERCA Ca^{2+}-ATPase the ER is highly efficient at storing Ca^{2+} and transferring it to other compartments.

RESULTS

Extracellular Ca^{2+} predominantly fills the endoplasmic reticulum and acidic stores

We first set up experiments to follow the destination of Ca^{2+} taken up by extracellular tachyzoites from the extracellular milieu into their cytosol. Tachyzoites were loaded with the ratiometric Ca^{2+}

indicator Fura-2 and incubated in Ringer buffer containing 100 µM EGTA to chelate extracellular Ca^{2+}, preventing its uptake, and investigated the release of intracellular Ca^{2+} pools by different inhibitors.

When thapsigargin (TG), which inhibits the activity of SERCA, was added, it prevented the re-uptake of Ca^{2+} into the ER and unmasked the leakage of Ca^{2+} from the ER into the cytosol, causing a Ca^{2+} cytosolic increase of approximately 160 nM **(Fig. 1B and S1A)**. We compared this with the effect of ionomycin (IO), a Ca^{2+}/H^+ ionophore which acts on neutral intracellular Ca^{2+} stores, inducing depletion of Ca^{2+}, mainly from the ER (Smith *et al.*, 1989). Exposure of *T. gondii* to 1 µM ionomycin caused a cytosolic increase of approximately 700-1100 nM Ca^{2+} **(Fig. S1A-B)**. We next tested another SERCA inhibitor, cyclopiazonic acid (CPA), with a different mode of action than TG. CPA caused a lower Ca^{2+} cytosolic increase than the one caused by TG, possibly implying less specificity toward TgSERCA **(Fig. S1B).** Exposure of parasites to TG before CPA ablated CPA's effect, but this did not occur when adding CPA prior to TG **(Fig. S1C-D)**. Although TG does a better job at depleting the ER of Ca^{2+}, the increase of cytosolic Ca^{2+} after adding TG is un-impressive when compared to the effect of IO **(Fig. S1A)**. This result could indicate the presence of other calcium stores not inhibited by TG or simply that TG was not very efficient at blocking TgSERCA.

Next, we tested the effect of Ca^{2+} entry in the filling of intracellular stores by measuring the cytosolic Ca^{2+} increases in response to inhibitors after pre-exposing the parasite suspension to extracellular Ca^{2+} **(Fig. 1C-F,** *dark traces*). Addition of 1.8 mM Ca^{2+} caused a cytosolic increase due to influx through a partially characterized plasma membrane pathway (**Fig. 1C**, *dark blue trace*) (Pace *et al.*, 2014). This Ca^{2+} influx appeared to be immediately taken up by the ER as the response to TG was much greater in parasites previously exposed to extracellular Ca^{2+} (compare light and dark blue traces) (**Fig. 1C**). We next tested the lysosomotropic agent glycyl-L-phenylalanine-naphthylamide (GPN), which mobilizes Ca^{2+} from acidic organelles (Haller *et al.*, 1996; Lloyd-Evans *et al.*, 2008; Miranda *et al.*, 2010; Yuan *et al.*, 2021) and observed a similar pattern (**Fig. 1D**). The cytosolic Ca^{2+} increase upon addition to GPN in parasites previously exposed to extracellular Ca^{2+} was much greater than in cells not exposed to extracellular Ca^{2+} (**Fig. 1D**, *compare light and dark blue traces*). A similar phenomenon was observed when adding nigericin (acting on acidic stores) or CCCP (most likely acting on mitochondria) (**Figs 1E-F**, *dark purple and orange traces, respectively*). These data indicate that the ER, mitochondrion, and PLVAC and other acidic stores, all appear to have the ability to access extracellular Ca^{2+}. The ER displayed the highest capacity to access a large portion of extracellular Ca^{2+}, with TG inducing close to ~500-900 nM of Ca^{2+} release after pre-exposure to Ca^{2+} and only 210 nM Ca^{2+} without Ca^{2+} pre-exposure **(Fig. 1C)**.

In summary, upon pre-exposure of *T. gondii* to physiological levels of extracellular Ca^{2+}, the ER, mitochondria, and acidic stores exhibited the capacity to release many times more Ca^{2+} than without pre-exposure, with the ER and GPN-sensitive stores displaying the greatest response **(Fig 1G).**

We next hypothesized that if the ER is the most efficient organelle to access extracellular Ca^{2+} and pass it to the other organelles, it should be possible to see an increased response to agonists like GPN and nigericin after inhibiting SERCA with TG permitting the buildup of Ca^{2+} on the cytosolic side of the ER membrane. For the next experiment we added Ca^{2+} to fill the stores, followed by TG to cause Ca^{2+} efflux from the ER, and then GPN (**Fig. 1H**). We compared the response to GPN between plus and minus the addition of TG and observed that the cytosolic increase caused by GPN when preceded by TG was significantly larger (**Fig. 1H**). We followed the same approach with nigericin and observed that the response increased after the addition of TG (**Fig. 1I**). These experiments support our hypothesis that extracellular Ca^{2+} is used to fill the ER and from the ER Ca^{2+} can be distributed to other organelles.

Ca^{2+} uptake by the SERCA-Ca^{2+} ATPase in permeabilized tachyzoites

The previous results highlighted an important role for the ER in taking up Ca^{2+} from the cytoplasm after an increase due to influx from the extracellular milieu. We believe that this ER uptake, essential for keeping stores filled with Ca^{2+}, is due to the high Ca^{2+} affinity of TgSERCA. The SERCA functions as an essential mechanism for the ER to maintain its Ca^{2+} levels, an especially important function considering the leakage of Ca^{2+} from the ER into the cytosol which occurs constitutively and passively (Camello *et al.*, 2002).

We developed a protocol that allowed us to measure directly Ca^{2+} uptake by the store where the SERCA is localized (ER), based in the compartmentation of a fluorescent dye in permeabilized cells. We used the low affinity indicator MagFluo4 which has a Kd for Ca^{2+} of 22 µM (Rossi and Taylor, 2020). The cytosolic concentration of Ca^{2+} in *T. gondii* is around ~70 nM (Moreno and Zhong, 1996), which is considerably below the detection limit of MagFluo4. We loaded parasites for an extended length of time and using high concentrations of MagFluo4-AM as the intention was for the indicator to compartmentalize. At the end of the incubation, parasites were washed and exposed to low concentration of digitonin, which permeabilized the plasma membrane leaving the organellar membranes intact. Under these conditions we had parasites loaded with a Ca^{2+} indicator inside organelles (**Fig. 2A**). We next determined the capacity of these permeabilized parasites to take up Ca^{2+}. SERCA activity is dependent on the presence of Mg-ATP (**Fig. 2B**) and we added this substrate under defined concentrations of Ca^{2+} (determined using the MaxChelator program). (Bers *et al.*, 1994). We observed that under these conditions a consistent and reproducible Ca^{2+} uptake was measured that was optimal at 220 nM Ca^{2+} and 500 µM MgATP (**Figs 2B-C**). Validation that this activity was due to SERCA is shown in the experiment in which we added TG highlighting the ER Ca^{2+} leakage (**Fig. 2D**). Note that the Ca^{2+} released by TG appears to be modest but the Kd of the indicator is 22 µM so a small decrease is highly significant when considering the amount of Ca^{2+} that it represents. Ionomycin can be more effective and larger amounts of Ca^{2+} were released from the ER and other stores as the minimum reached is lower than the starting level of Ca^{2+} (**Fig. 2E**).

In summary, these results showed that we can measure SERCA activity *in situ* in permeabilized parasites with the low affinity Ca^{2+} indicator MagFluo4. This activity is dependent on MgATP and

184 TG causes release of the accumulated Ca^{2+}. The better effect of IO could be due to Ca^{2+} release
185 from other compartments driven by other ATPases (TgA1).
186
TgSERCA and the *T. gondii* lytic cycle

To investigate the role of TgSERCA (TGGT1_230420) in the biology of *T. gondii* we created
conditional KOs (*iΔTgserca*) by inserting a tetracycline regulatable element at the 5' end of the
gene to control the expression of the gene with ATc (Sheiner *et al.*, 2011). In addition, we
endogenously tagged TgSERCA with HA. We isolated clonal cell lines of *iΔTgserca* and clones
expressing *iΔTgserca-HA* (**Fig. 3A**). Western blots (**Fig. 3B**) and immunofluorescence analysis
(**Fig. S2**) showed that the expression of TgSERCA is tightly regulated by ATc and it is absent after
1.5 days in culture. The growth of the *iΔTgserca* was severely retarded when cultured in the
presence of ATc as evaluated by plaque assays (**Fig. 3C-D**). In this assay, parasites engage in
repetitive cycles of invasion, replication, and egress causing host cell lysis and formation of
plaques in confluent monolayers infected with *T. gondii*. Replication was severely impacted after
down regulation of SERCA, and parasites were unable to proceed after one or two replication
cycles (**Fig 3E-F**). All (100%) PVs of the ATc cultured mutant contained 4 parasites or less (**Fig.**
3F, *bar graph on the right*). Invasion of host cells was also reduced in the *iΔTgserca* (+ATc)
mutant (**Fig. 3G**) and egress stimulated by ionomycin, or natural egress followed pre-incubation
with Compound 1, was significantly decreased. Interestingly, when testing egress after the addition
of saponin in the presence of extracellular Ca^{2+}, we observed that the tachyzoites egressed sooner
(**Fig. 3H**, *saponin egress*). This could be because as the SERCA is not active, the cytosolic Ca^{2+}
increase due to Ca^{2+} influx from the extracellular milieu after saponin permeabilization cannot be
compensated. In addition, the natural efflux from the ER would result in a higher cytosolic Ca^{2+}
increase that would cause a faster egress.
208
Ca^{2+} uptake by the SERCA-ATPase is essential for filling acidic Ca^{2+} stores

The *iΔTgserca* mutant showed a low response to TG (**Fig. 4A**), when measuring cytosolic calcium
(**Fig. 4A**) which was also the case when adding TG after extracellular calcium to fill the store (**Fig.**
4B). This was most likely the result of the lower activity of TgSERCA. Note that the change in
Ca^{2+} in Fig. 4A (*TatiΔku80* cells) is larger than in Fig. 1B (RH strain) and we believe that the
difference is related to the characteristics of each cell line (RH vs TatiΔku80). In addition, Mg-
ATP-driven Ca^{2+} uptake by permeabilized cells measured with MagFluo4 showed no TgSERCA
activity after 48 h of culture with ATc (**Fig. 4C**). This experiment validated the MagFluo4
approach for determination of SERCA activity. At 24 hours after culturing the parasites with ATc
there was still some SERCA activity (Fig. 4C). Ca^{2+} entry measured in Fura2 loaded *iΔTgserca*
(±ATc) parasites was not affected by the downregulation of TgSERCA (**Fig. 4D**). Interestingly,
the cytosolic resting Ca^{2+} concentration was not affected in the *iΔTgserca* (+ATc) mutant (Fig. 4
A,B, D-H) pointing to a critical role of the plasma membrane Ca^{2+} pump in the homeostasis of
cytosolic Ca^{2+}. The response to Zaprinast was diminished but was still present (**Fig 4E**) indicating

that Zaprinast acts by releasing Ca^{2+} from the ER and from an additional compartment. Adding
Zaprinast after calcium also resulted in a reduced response in the mutant pre-incubated with ATc
(**Fig 1F**). We next tested GPN, which acts mostly on acidic stores and the response was also
decreased (**Fig. 4G**). Adding GPN after replenishing the cells with Ca^{2+} resulted in an increased
response as we showed in Fig. 1, but this response was also reduced when the mutant was grown
with ATc (**Fig. 1H**). These results support a link between the stores on which GPN is acting upon
and the ER. Considering that downregulation of SERCA impacted the ER storage of Ca^{2+} but did
not affect cytosolic Ca^{2+} uptake, the diminished response to GPN, supports that the Ca^{2+}
released/leaked from the ER plays an important role in filling the GPN's targeted store.

233 The mitochondrion takes up Ca^{2+} from the ER

In mammalian cells, the high concentration of Ca^{2+} in the ER is important for the production of
mitochondrial ATP (Wenzel *et al.*, 2022). This is because of the close proximity between the ER
and mitochondria which allows for the directional flow of Ca^{2+} from the ER to the mitochondria
(Gincel *et al.*, 2001; Rapizzi *et al.*, 2002). With the aim of determining if the *T. gondii*
mitochondria can take up calcium, we introduced a genetic Ca^{2+} indicator in the mitochondrion of
T. gondii tachyzoites by attaching the *GCaMP6* gene (Chen *et al.*, 2013) to the mitochondrial
targeting signal of the *T. gondii* superoxide dismutase 2 (SOD2) gene (Pino *et al.*, 2007) and
isolated parasite clones (*Tgsod2gcamp6*) (Vella *et al.*, 2020). GCaMP6 was localized to the
mitochondria as detected by Fluorescence (**Fig 5A**). Direct Ca^{2+} uptake was observed in digitonin
permeabilized parasites incubated in the presence of increasing concentrations of Ca^{2+} (**Fig 5B-C**).
A measurable increase was observed but very high concentrations of Ca^{2+} were needed (**Fig. 5B-**
C) indicating that the *T. gondii* mitochondria can take up Ca^{2+} but with very low affinity and that
the concentrations of Ca^{2+} needed for the uptake were much higher than the concentrations of
cytosolic Ca^{2+} that are present in the cytosol of a live cell (**Fig 5B**).
We reasoned that the mitochondrion could be taking up Ca^{2+} through close membrane contacts
with the ER as it is possible that the concentration of Ca^{2+} in those microdomains could reach
much higher levels than in the cytosol and would allow for mitochondrial uptake. We next loaded
the *Tgsod2gcamp6* mutant with Fura2 and measured, first, if adding extracellular Ca^{2+} to the
parasite suspension led to an increase in GCaMP6 fluorescence. Parallel experiments looking at
cytosolic and mitochondrial Ca^{2+} showed that adding extracellular Ca^{2+} does result in an increase
in the cytosol, however mitochondrial Ca^{2+} was not affected (**Fig. 5D-E**). We next added TG
followed by Ca^{2+} and this led to an increase in cytosolic Ca^{2+} but only TG resulted in a measurable
increase in fluorescence of the mitochondrial indicator (**Fig 5G**). Our interpretation is that after
adding TG, there was accumulation of Ca^{2+} at the outer side of the ER membrane leading to an
increased concentration of the ion which activated mitochondrial uptake.
We next inserted the *SOD2GCaMP6* chimeric gene in the *iΔTgserca* mutant and isolated the
clonal mutant *iΔTgserca-sod2gcamp6.* We first looked at mitochondrial uptake after adding TG
and as shown previously, the parasites cultured without ATc showed a measurable and consistent
increase in GCaMP fluorescence after adding TG (**Fig. 5H**, *blue trace*). This was abolished in cells

cultured in the presence of ATc as the low expression of TgSERCA would result in the ER
depletion of Ca^{2+} (**Fig. 5H,** *pink trace*). Interestingly, we observed an increase in the GCaMP
fluorescence after adding GPN, ionomycin or Zaprinast (**Fig. 5H-J,** *blue traces*). In all cases the
fluorescence increase was muted in the *iΔTgserca* (+ATc) and the difference in the response was
always significant (**Fig. 5H-J,** *pink traces*).

In summary, we demonstrated that the mitochondrion of *T. gondii* can take up Ca^{2+} through its
transfer from the ER, a phenomenon that becomes evident after blocking the ER uptake with TG
as the concentration of Ca^{2+} needed to activate the uptake is very high and only reachable at
membrane contact sites between the ER and mitochondria. The *iΔTgserca* (+ATc) mutant, is
unable to keep the ER filled with Ca^{2+} and this impacts the capacity of the mitochondria to take up
Ca^{2+}. The mitochondria could be also taking up Ca^{2+} directly from acidic stores, which became
depleted when TgSERCA was downregulated.

Interaction between the ER and the mitochondrion and acidic compartments.

We next investigated if it was possible to see interaction between the ER with other organelles by
IFA and/or EM. We performed IFAs with ER and mitochondria markers and ER and PLVAC
markers (**Fig. 6**). We observed that in intracellular parasites the mitochondrion surrounds the ER
and multiple potential points of interaction are seen (**Fig. 6A** and supplemental video 1). The
mitochondrion of intracellular *T. gondii* tachyzoites surrounds the periphery of the cell in a lasso-
shape morphology. In extracellular parasites the mitochondrion changes drastically its shape
adopting either a sperm-like or a collapsed conformation (Ovciarikova *et al.*, 2017) (**Fig. 6B** and
supplemental video 2). Our hypothesis is that this retraction of the mitochondrion allows the ER
to expand in extracellular parasites into ER tubules, which extend toward the apical domain of the
parasite where Ca^{2+} is needed for secretion of micronemes and conoid extrusion.

The PLVAC also showed points of contact with the ER (**Fig. 6C** and supplemental video 3).
Multiple points of contact were also observed by EM between the ER and the PLVAC, the ER and
the apicoplast and the ER and the mitochondrion (**Fig. 6D**).

Interestingly these contacts were still present in the *iΔTgserca* (+ATc) mutant (**Fig. S3, A-D**),
as most likely TgSERCA would not be directly involved in the establishment of contacts. We
quantified the length of the limiting membrane of the organelle in contact with ER tubules at a
distance of less than 30 nM and found that after knockdown of TgSERCA, the contact was not
altered. However, Calcium transported from the ER into the mitochondrion after thapsigargin
treatment was significantly decreased which means that this phenotype is due to reduced ER
Calcium and not to lack of contacts (**Fig. S3E**). A similar result was seen when measuring contacts
between the ER and the apicoplast (**Fig. S3F**) and between the ER and the PLVAC (**Fig. S3G**).

In summary, this data showed that the ER form contacts with other organelles like the
mitochondrion, the PLVAC and the apicoplast and these contacts could be the ones allowing the
transfer of Calcium from the ER, the store with the highest amount of Calcium, to the other
organelles. The SERCA high affinity for Calcium is critical for the filling of the ER. However, the
molecular mechanisms involved in the uptake by the other organelles is less clear. Very little is

known about the structure and composition of Membrane Contact Sites in *T. gondii* (Huet and
Moreno, 2023). However, functional evidence for the transfer of Calcium has been demonstrated
for the ER-apicoplast (Li *et al.*, 2021) and in this work we presented evidence for the transfer of
Calcium between the ER and the mitochondrion and the ER and the PLVAC.
307
DISCUSSION

In this work we showed that the endoplasmic reticulum (ER) of *T. gondii* has an impressive
capacity for capturing Ca^{2+} that enters the cytosol from the extracellular milieu, and it
accomplishes this with a minimal increase in cytosolic Ca^{2+} levels. This is the result of a highly
efficient SERCA Ca^{2+}-ATPase (TgSERCA) which also showed high affinity for Ca^{2+}. The
TgSERCA activity combined with the plasma membrane Ca^{2+} pump (Luo *et al.*, 2001; Luo *et al.*,
2005) would not allow the cytosolic Ca^{2+} to increase above 200 nM (Hortua Triana *et al.*, 2024).
We also provide evidence of the ER's capacity for sharing Ca^{2+} with other organelles, which
originates from the ER unique ability to capture a sizable portion of extracellular Ca^{2+}. We showed
that extracellular Ca^{2+} can be efficiently taken up into the ER through the activity of SERCA and
from the ER it can be transferred to other compartments like the mitochondria and acidic stores.
These activities would ensure that Ca^{2+} is released at specific locations without flooding the whole
cell with Ca^{2+}, which may trigger unnecessary signals. Although initial data suggested a less
important role for the mitochondria and PLVAC in Ca^{2+} storage, our results showed that these
organelles may become more relevant Ca^{2+} reservoirs during ER calcium overload
The SERCA Ca^{2+}-ATPase is a P-type ATPase found in the membranes of the ER and the
secretory pathway (Kuhlbrandt, 2004). The mammalian genome encodes for three SERCA pumps:
SERCA1, SERCA2, and SERCA3, and seven different isoforms expressed from these genes,
SERCA1a/1b, SERCA2a/2b, and SERCA3a/3b/3c (Periasamy and Kalyanasundaram, 2007).
SERCA2b is the most widespread of all isoforms and phylogenetically likely the oldest and
considered the house-keeping isoform (Wuytack *et al.*, 2002). SERCA pumps translocate two Ca^{2+}
ions and hydrolyze one ATP for each catalytic turnover. This pumping activity serves several roles.
By translocating Ca^{2+} from the cytosol into the lumen of the ER, it restores the cytosolic Ca^{2+}
concentration to its low resting level (<100 nM). The stored Ca^{2+} can be released and activate
many cellular activities. In addition, SERCA maintains a sufficiently high (~500 µM) luminal Ca^{2+}
concentration that is necessary for enzyme activities important for proliferation, growth, and
differentiation (Wuytack *et al.*, 2002).
T. gondii appears to express a single SERCA protein (TgSERCA) (Nagamune and Sibley, 2006),
most likely playing a housekeeping role like the mammalian SERCA2. This was evident from the
strong phenotype of the *iΔTgserca* (+ATc) mutant, which could not replicate after 24 h and was
defective for all the steps of the tachyzoite lytic cycle. The activity of TgSERCA was highly
dependent on the presence of its substrate MgATP and it showed high affinity for Ca^{2+}.
Considering the low affinity of the indicator Magfluo4 (Kd: 22 µM), the measured uptake activity
and the accumulation of Ca^{2+} was impressive as it was evident even at 55 nM free calcium. This
result signified that TgSERCA is highly efficient at keeping the cytosol at the low 60-100 nM level

as well as efficiently filling the ER with Ca^{2+}.

Ca^{2+} signals are specifically contingent on their spatial, temporal and amplitude features.
Movement of Ca^{2+} in the cytosol of cells is severely limited due to the presence of high affinity
Ca^{2+} buffers. The concept of Ca^{2+} tunneling through the ER proposed that Ca^{2+} would move more
easily within the lumen of the ER as its binding capacity is almost 100 lower than the binding
capacity of the cytosol (Mogami *et al.*, 1997) and the high mobility of Ca^{2+} through the lumen of
the ER was directly demonstrated (Park *et al.*, 2000), and especially important in polarized cells.
In addition, the ER facilitates Ca^{2+} tunneling through its lumen as a mechanism of delivering Ca^{2+}
to targeted sites without activating inappropriate processes in the cell cytosol (Petersen *et al.*,
2017). In *T. gondii*, the Ca^{2+} binding capacity of the cytosol compared to the ER is less known as
the localization of all its predicted Ca^{2+} binding proteins have not been completely determined and
several calmodulin-like proteins showed localization to the conoid (Long *et al.*, 2017).

In *T. gondii* cytoplasmic Ca^{2+} increases, due to efflux from the ER, have been reported to initiate
essential parasite traits like microneme secretion (Carruthers and Sibley, 1999; Nagamune *et al.*,
2007b), conoid extrusion (Del Carmen *et al.*, 2009), invasion (Vieira and Moreno, 2000; Lovett
and Sibley, 2003) and egress (Arrizabalaga *et al.*, 2004). These responses require precise
spatiotemporal control of diverse targets and suggests the presence of distinct systems to deliver
Ca^{2+} to specific locations. Our hypothesis is that the ER plays a central role by distributing Ca^{2+} to
those locations at specific times for the initiation of parasite functions. The strong invasion,
replication and egress phenotypes of the *iΔTgserca* (+ATc) supports this hypothesis.

In mammalian cells Ca^{2+} ion is transferred from the ER to the mitochondrion through the outer
membrane voltage-dependent anion channel1 (VDAC1) (Gincel *et al.*, 2001; Rapizzi *et al.*, 2002)
and the inner membrane calcium uniporter (MCU1) (De Stefani *et al.*, 2011). A VDAC homologue
is present in *T. gondii*, which was shown to be essential for growth and for mitochondrial and ER
morphology (Mallo *et al.*, 2021). However, the molecular evidence for the presence of a Ca^{2+}
uniporter in the inner mitochondrial membrane driven by the electrochemical gradient generated
by the electron transport is not supported by the genome database. Calcium transport into the *T.*
gondii mitochondrion had not been previously shown and our data represents the first experimental
evidence showing this event although the mechanism is less clear. We showed that it has very low
affinity for Ca^{2+} and it was only possible to observe it in intact parasites after inhibiting SERCA,
which allowed the increase of the concentration of Ca^{2+} at the local ER membrane. Over the past
decade a new concept about the communication between subcellular organelles through tight,
stable, and yet non-fusogenic associations known as membrane contact sites (MCSs) has emerged
(Phillips and Voeltz, 2016). The ER forms an extensive network of MCSs with the PM,
mitochondria and endocytic vesicles for the exchange of Ca^{2+} (Burgoyne *et al.*, 2015).

In *T. gondii* the characterization of MCSs is only in its beginnings (Huet and Moreno, 2023) with
only a few evidences for their presence (Tomova *et al.*, 2009; Mallo *et al.*, 2021; Ovciarikova *et*
al., 2022) and function (Li *et al.*, 2021; Oliveira Souza *et al.*, 2022). When *T. gondii* is intracellular,
microscopy observations showed that its mitochondrion surrounds the periphery of the cell in a
lasso-shape conformation. On the other hand, in extracellular parasites the mitochondrion changes

its morphology and adopts a sperm-like or collapsed conformation (Ovciarikova *et al.*, 2017). Our
IFA analysis with ER and mitochondrial markers showed that the lasso-shaped mitochondrion
surrounds the ER with plenty of opportunities for contact between both organelles. We speculate
that at this stage the mitochondrion is buffering any leakage of Ca^{2+} from the ER to prevent
premature egress as a cytosolic increase would stimulate motility and possibly egress.

In this work, we showed that Ca^{2+} can be transferred from the ER to the mitochondrion, most
likely through MCSs as the low affinity of the mitochondrion would not permit uptake directly
from the cytosol. Down regulation of TgSERCA resulted in diminished response to acidic stores
triggers, which supported that the ER may also be responsible for their filling. The mitochondrion
may also be able to take up Ca^{2+} from acidic stores like the PLVAC as GPN was also able to
stimulate Ca^{2+} uptake and the response was diminished in the *iΔTgserca* (+ATc) mutant. Transfer
of Ca^{2+} from acidic stores to the mitochondrion was impacted by the ER depletion (Fig. 7).

In conclusion, this study showed how the ER of *T. gondii* replenish itself with calcium and how
it supplies Ca^{2+} to the cytosol for signaling and to acidic stores and the mitochondrion. *T. gondii*
is a protozoan parasite that causes disease by reiterating a lytic cycle that is driven by calcium
signaling. This work contributes to the understanding of how extracellular and intracellular Ca^{2+}
stores work together to sustain the pathologic features of *T. gondii*. Future studies would aim at
finding more about the role of Ca^{2+} in mitochondrial and acidic stores functions.

Methods

Cell Culture—*Toxoplasma gondii* tachyzoites (RH strain) were maintained in human telomerase
reverse transcriptase immortalized foreskin fibroblasts (hTERT) (Farwell *et al.*, 2000) grown in
Dulbecco's modified minimal essential media (DMEM) with 1% FBS.
406

Generation of SERCA mutants—A promoter insertion plasmid was generated by cloning three
PCR fragments into a modified pCR2.1-TOPO vector using Gibson Assembly Cloning Kit (NEB
#E5510). One fragment corresponding to the TgSERCA flanking region (predicted
promoter/5'UTR) was amplified with primers 1 and 2 (Table S1). The second fragment
corresponds to DHFR+T7S4 (Sheiner *et al.*, 2011) and was amplified with primers 3 and 4.
Another fragment corresponds to the 5' TgSERCA coding sequence beginning with start codon
and was amplified with primers 5 and 6. The vector pCR2.1-TOPO was used, which had only one
EcoRI site and it was cut with the enzyme NotI to use as a vector backbone. The promoter insertion
plasmid was transfected into the *TatiΔku80* cells and selected with 1 μM pyrimethamine using a
"ultra aggressive" screening method. Briefly, 200 μl of the suspension of transfected parasites was
added to 10 ml of media and then one to three drops (~65 μl per drop) were inoculated into three
24-well plates containing media. The clonal lines created after selection and subcloning were
termed *iΔSERCA*.

For *in situ* tagging, an approximately 2 kb fragment was amplified from the genomic locus (3'
region) of the TgSERCA gene using primers 7 and 8. The fragment was cloned into the pLic-3HA-
CAT plasmid (Huynh and Carruthers, 2009) and the construct was linearized with the enzyme

NheI for transfection into the *iΔSERCA* mutant. Clonal cell lines were generated after selection with chloramphenicol and subcloning and termed *iΔSERCA-3HA*.

Expression and purification of TgSERCA recombinant protein—The phosphorylation(P) and nucleotide binding(N) domains of *Tg*SERCA (TGGT1_230420) (nucleotides 1123 to 2415, amino acid residues 375 to 805) were cloned into XmaI and HindIII sites of pQE-80L with primers 13 and 14 to create recombinant protein with a N-terminal 6xHis tag. The resulting plasmid was transformed into *Escherichia coli* BL21-CodonPlus competent cells and expression was induced by addition of 0.4 mM isopropyl β-D-1-thiogalactopyranoside (IPTG) for 4 hours at 37°C. Cells were pelleted and resuspended in equilibration/binding buffer (50 mM Na_3PO_4, 300 mM NaCl, 10 mM Imidazole, 8 M Urea, and protease inhibitor cocktail P-8849). The cells were then sonicated for 80 seconds total and centrifuged at 12,000 rpm for 20 mins at 4°C. The supernatant was filtered through a 0.45 μm membrane and the protein was purified using HisPur Ni-NTA Chromatography Cartridge (Thermo Scientific) following instructions from the manufacturer. Proteins that were unbound were washed off with 12 ml of wash buffer (50 mM Na_3PO_4, 300 mM NaCl, 40 mM imidizole, and 8 M urea), and the recombinant protein was eluted with 5 ml elution buffer (50 mM Na_3PO_4, 300 mM NaCl, 250 mM iImidizole, and 8 M urea). Eluted protein fractions were concentrated and desalted with Amicon Ultra-0.5 mL centrifugal filter (Millipore Sigma).

Anti-TgSERCA antibody generation in Guinea pig—Two Guinea pigs were each immunized with 0.2 mg of purified recombinant protein mixed with equal volume of Freund's Complete Adjuvant (Sigma F5581), followed by two boosts of 0.1 mg antigen mixed with equal volume of Freund's Incomplete Adjuvant (Sigma F5506) for Guinea pig 1 and three boosts for Guinea pig 2. The resulting antibodies were tested at 1:1000 in western blot against RH lysates and were developed with Alexa Fluor 488 goat anti-Guinea pig (1:1000). The antibodies were compared with Dr. Sibley's mouse anti-SERCA antibody (Nagamune *et al.*, 2007a) to confirm size and purity. The anti-*Tg*SERCA antibodies were then affinity purified.

Cytosolic calcium measurements with Fura-2—*T. gondii* tachyzoites were loaded with Fura-2 AM as previously described (Vella *et al.*, 2020; Stasic *et al.*, 2021). Freshly released tachyzoites were washed twice with buffer A plus glucose (BAG; 116 mM NaCl, 5.4 mM KCl, 0.8 mM $MgSO_4$, 50 mM Hepes, pH 7.3, and 5.5 mM glucose), by centrifugation (706 x *g* for 10 min) and re-suspended to a final density of 1×10^9 parasites/ml in loading buffer (BAG plus 1.5% sucrose, and 5 μM Fura2 AM). The suspension was incubated for 26 min at 26 °C with mild agitation. Subsequently, the parasites were washed twice (2000 x *g* for 2 min) with BAG to remove extracellular dye, re-suspended to a final density of 1×10^9 parasites per ml in BAG and kept on ice. For fluorescence measurements, 2×10^7 parasites/mL were placed in a cuvette with 2.5 mL of Ringer's buffer without calcium (155 mM NaCl, 3 mM KCl, 1 mM $MgCl_2$, 3 mM NaH_2PO_4, and 10 mM Hepes, and 10 mM glucose). Fluorescence measurements were done in a Hitachi F-7000 fluorescence spectrofluorometer using the Fura-2 conditions for excitation (340 and 380 nm) and emission (510 nm). The Fura-2 fluorescence response to Ca^{2+} was calibrated from the ratio of 340/380 nm fluorescence values after subtraction of the background fluorescence of the cells at

465 340 and 380 nm as previously described (Grynkiewicz *et al.*, 1985). The Ca^{2+} release rate was
466 defined as the change in Ca^{2+} concentration during the initial 20 s after reagent addition. Delta F
467 was calculated by the difference between the higher calcium peak and basal Ca^{2+}, and recovery
468 was defined as the change of Ca^{2+} concentration after the calcium peak was reached for the
469 subsequent indicated times.
470

Endoplasmic reticulum Ca^{2+} Measurements in permeabilized T. gondii tachyzoites—
Tachyzoites freshly egressed and washed as described above were resuspend to a final density of
$1x10^9$ cells/ml in HBS buffer (135 mM NaCl, 5.9 mM KCl, 1.2 mM $MgCl_2$, 11.6 mM HEPES pH
7.3, 1.5 mM $CaCl_2$, 11.5 mM glucose) containing 1 mg/ml BSA, 0.2 mg/ml of pluronic F127 and
20 μM MagFluo4/AM. The suspension was incubated in a 20°C water bath with mild shaking for
1 h, in the dark. Subsequently, parasites were washed 3 times at 5,000 rpm for 2 min to remove
extracellular dye. The pellet was resuspended in 1.8 ml of CLM buffer (20 mM NaCl, 140 mM
KCl, 20 mM PIPES, pH 7.0) containing 1 mM EGTA at $1x10^9$ cells/ml. Cells were permeabilized
with 44.4 μg/ml digitonin for 10 min. Cells were washed 3 times with CLM containing 1 mM
EGTA at 5,000 rpm for 2 min to remove digitonin and resuspended to a final density of $1x10^9$
tachyzoites/ml and kept in ice. For each test, 50 μl ($5x10^7$) of parasite suspension was added to
1.95 ml of CLM containing 1 mM EGTA and 0.375 mM $CaCl_2$ which results in 220 nM free Ca^{2+}
as calculated with MaxChelator. Fluorescence was measured with an Hitachi F-7000 fluorescence
spectrometer (Excitation at 485 nm and emission at 520 nm). Ratio ($\Delta F/F_0$) was evaluated by
measuring the rate of change in fluorescence over 20 s after reagent addition (linear rate).
486

Strain construction and maintenance—The organelle targeting of GCaMP6f was made by
overlapping PCR. The N-terminal mitochondrial targeting sequence of the *T. gondii* SOD2 gene
(Pino *et al.*, 2007) was used to target GCaMP6f to the mitochondrion. The Gcamp6f for this
construct was amplified by primers 9 and 10 (Table S1). After gel purification of the GCaMP6f
and SOD2 sequences, the mitochondria targeting construct was built by overlapping PCR with the
purified PCR products as template. This construct was then cloned into the Topo-blunt vector.
After the sequence was verified by sequencing, the SOD2-Gcamp6f fragment was removed by
BglII and AvrII digestion and cloned into the same restriction sites of the pDT7S4H3 (Sheiner *et
al.*, 2011) and pCTH3 (Vella *et al.*, 2020) vectors. The pDT7S4H3-SOD2-GCaMP6f construct
was introduced into RH parasites by electroporation. After selection with pyrimethamine, the
parasites were sorted by FACS and then subcloned. Clones were selected based on the dynamic
range of the response to ionomycin. The pCTH3-SOD2-GCaMP6f was introduced into the
iΔTgSERCA mutant by electroporation. After selection with chloramphenicol, the parasites were
sorted by FACS and then subcloned. The expression of GCaMP6f was verified by live cell
imaging, IFA, and western blots. The clone with the largest dynamic range, as evaluated with
Ionomycin, was selected to further use.
503

GCaMP6f fluorescence Measurements—Permeabilized tachyzoites: *T. gondii* tachyzoites
expressing SOD2-GCaMP6 were collected and washed 3 times at 5,000 rpm for 2 min. The
parasite pellet was resuspended in 1.8 ml of BAG buffer containing 0.1 mM EGTA at $1x10^9$

cells/ml. Permeabilization with 44.4μg/ml digitonin for 10 min was done by following the
fluorescence of GCaMP6. Cells were washed 3 times with the same buffer at 5,000 rpm for 2 min
to remove digitonin, resuspended to a final density of $1x10^9$ parasites/ml and kept in ice. 50 μl
($5x10^7$) of the parasite suspension was mixed with 1.95 ml reaction buffer containing 0.1 mM
EGTA.

For measurements with intact parasites, they were collected, washed, and resuspended in BAG
at $1x10^9$ cells/ml for testing. Measurements were done in a Hitachi 7000 fluorescence spectrometer
set at excitation, 485 nm and emission, 509 nm. The ratio ($\Delta F/F_0$) was evaluated by measuring the
rate of change in fluorescence over 20 s after reagent addition (linear rate).
516

Growth, Invasion and Egress assays—Red-green invasion assays were performed as originally
described (Kafsack *et al.*, 2004), modified (Chasen *et al.*, 2017) and adapted to use td-RFP
expressing parasites. The number of tachyzoites used was 2×10^7, and invasion was for 5 min.
Plaque assays were performed as described in (Liu *et al.*, 2014) with modifications. 125
tachyzoites were used for infection of confluent six well plates with hTert fibroblasts followed by
an incubation time of 10 days prior to fixing and staining with crystal violet.

For egress assays, HeLa cells were infected with either parental or *iΔTgSERCA-HA*
tachyzoites. *iΔTgSERCA-HA* tachyzoites were grown in HeLa cells in the absence or presence of
anhydrous tetracycline for the duration of the infection. Forty-eight hours after infection,
parasitophorous vacuoles containing 4-8 parasites were monitored by microscopy after washing
with Ringer buffer. $CaCl_2$ was omitted for experiments in the absence of extracellular calcium,
and the media was supplemented with 100 μM or 1 mM EGTA to chelate contaminating calcium.
For experiments containing $CaCl_2$, 2 mM $CaCl_2$ was added to the calcium-free Ringer buffer right
before the addition of the specific triggers. Drugs were added in calcium free Ringer buffer at the
concentrations indicated: ionomycin (1 μM), thapsigargin (1 μM), zaprinast (100 μM), saponin
(0.01%). Ringer buffer was used as extracellular buffer (EB).

For natural egress, the *iΔTgSERCA* mutant expressing td-tomato RFP was used to infect
confluent host cell monolayers 36 hours before adding ATc and 1 μM compound 1 (pyrrole 4-[2-
(4-fluorophenyl)-5-(1-methylpiperidine-4-yl)-1H-pyrrol-3-yl]pyridine) (Donald *et al.*,
2002) dissolved in ethanol and the culture continued for 24 hours. After treating with cpd1 for 24
hours, cultures showed full vacuoles which differed from the vehicle treated plates (36 hours
cultured plus ATc treatment for 24 hours or 48 hours without ATc), which were fully lysed. The
cpd1 media was removed and the synchronized vacuoles were washed twice, and the media
replaced with media without cpd1. The plates were then moved to the Prewarmed DeltaVision
microscope for observation. After 10 minutes at 37C, egress of the full vacuoles was enumerated.
We counted each plate for 1 minute and we counted at least 100 vacuoles per experiment. 3
independent biological experiments were conducted and summarized.

For replication assays, the hTERT cells were grown on 35 mm Mattek dishes. Each dish was
infected with 50,000 tdTomato-expressing parasites. 24 h after the infection the counting of
parasites per PV was done in a fluorescence microscope. For each experiment, at least 100 PVs
were counted. Results were the average of 3 independents experiments (Li *et al.*, 2021).

Microscopy and western blot analyses—Tachyzoites were grown on hTERT cells on cover slips for ~24 hr, washed twice with buffer A with glucose (BAG, 116 mM NaCl, 5.4 mM KCl, 0.8 mM $MgSO_4$, 50 mM HEPES, pH 7.2, and 5.5 mM glucose) and fixed with 4% formaldehyde for 1 h, followed by permeabilization with 0.3% Triton X-100 for 20 min, and blocking with 3% bovine serum albumin. IFAs were performed as previously described (Miranda *et al.*, 2010). Fluorescence images were collected with an Olympus IX-71 inverted fluorescence microscope with a Photometrix CoolSnapHQ CCD camera driven by DeltaVision software (Applied Precision, Seattle, WA). Superresolution microscopy was performed using a Zeiss ELYRA S1 (SR-SIM) microscope with a high-resolution Axio observer Z1 inverted microscope stand with transmitted (HAL), UV (HBO,) and high-power solid-state laser illumination sources (405/488/561 nm), a 100x oil immersion objective, and an Andor iXon EM-CCD camera and acquired using the ZEN software (Zeiss) with a SIM analysis module. Rat anti-HA antibody from Roche was used at a dilution of 1:25. Mouse anti-HA antibody from Covance was used at a dilution of 1:200.

Western blot analysis was performed as previously described (Liu *et al.*, 2014). Rat anti-HA antibody from Roche was used at a dilution of 1:200. Mouse anti-HA antibody from Covance was used at a dilution of 1:1,000. Secondary goat anti-rat or mouse antibody conjugated with HRP was used at 1:5,000. Mouse anti-α-tubulin at a dilution of 1:5,000 was used for loading control.

Transmission Electron Microscopy—For ultrastructural observations of intracellular *T. gondii* by thin-section transmission electron microscopy (EM), infected human foreskin fibroblast cells were fixed in 2.5% glutaraldehyde in 0.1 mM sodium cacodylate (EMS) and processed as described (Coppens and Joiner, 2003). Ultrathin sections of infected host cells were stained before examination with a Hitachi 7600 EM under 80 kV. For quantitative measurement of distance between organelles, the closest point between *T. gondii's* organelles and ER membrane was measured using ImageJ and was performed on 47 representative electron micrographs at the same magnification for accurate comparison between organelles

Statistical Analysis—Statistical analyses were performed by Student's t-test using GraphPad PRISM version 9. Error bars shown represent mean ± SD (standard deviation) of at least three independent biological replicates. Unpaired two tailed t test performed in all comparisons.

Acknowledgments

The authors thank Dr. Muthugapatti Kandasamy and the Biomedical Microscopy Core of the University of Georgia for the use of the microscopes. The CTEGD Cytometry Shared Resource Laboratory provided access and training to state-of-the-art flow cytometry analyzers. We would like to thank David Sibley for the generous gift of the mouse SERCA antibody and the plasmid for SERCA expression. This work was supported by the U.S. National Institutes of Health grants AI128356, AI154931 and AI174600 to SNJM and R01AI166921 to IC.

References

Arrizabalaga, G., and Boothroyd, J.C. (2004). Role of calcium during Toxoplasma gondii invasion and egress. International journal for parasitology *34*, 361-368.

Arrizabalaga, G., Ruiz, F., Moreno, S., and Boothroyd, J.C. (2004). Ionophore-resistant mutant of Toxoplasma gondii reveals involvement of a sodium/hydrogen exchanger in calcium regulation. J Cell Biol *165*, 653-662.

Bers, D.M., Patton, C.W., and Nuccitelli, R. (1994). A practical guide to the preparation of Ca2+ buffers. Methods Cell Biol *40*, 3-29.

Bisio, H., and Soldati-Favre, D. (2019). Signaling Cascades Governing Entry into and Exit from Host Cells by Toxoplasma gondii. Annu Rev Microbiol *73*, 579-599.

Black, M.W., and Boothroyd, J.C. (2000). Lytic cycle of Toxoplasma gondii. Microbiol Mol Biol Rev *64*, 607-623.

Blader, I.J., Coleman, B.I., Chen, C.T., and Gubbels, M.J. (2015). Lytic Cycle of Toxoplasma gondii: 15 Years Later. Annu Rev Microbiol *69*, 463-485.

Bootman, M.D., and Bultynck, G. (2020). Fundamentals of Cellular Calcium Signaling: A Primer. Cold Spring Harb Perspect Biol *12*.

Borges-Pereira, L., Budu, A., McKnight, C.A., Moore, C.A., Vella, S.A., Hortua Triana, M.A., Liu, J., Garcia, C.R., Pace, D.A., and Moreno, S.N. (2015). Calcium Signaling throughout the Toxoplasma gondii Lytic Cycle: A STUDY USING GENETICALLY ENCODED CALCIUM INDICATORS. J Biol Chem *290*, 26914-26926.

Burgoyne, T., Patel, S., and Eden, E.R. (2015). Calcium signaling at ER membrane contact sites. Biochim Biophys Acta *1853*, 2012-2017.

Camello, C., Lomax, R., Petersen, O.H., and Tepikin, A.V. (2002). Calcium leak from intracellular stores--the enigma of calcium signalling. Cell Calcium *32*, 355-361.

Carruthers, V.B., and Sibley, L.D. (1999). Mobilization of intracellular calcium stimulates microneme discharge in Toxoplasma gondii. Mol Microbiol *31*, 421-428.

Chasen, N.M., Asady, B., Lemgruber, L., Vommaro, R.C., Kissinger, J.C., Coppens, I., and Moreno, S.N.J. (2017). A Glycosylphosphatidylinositol-Anchored Carbonic Anhydrase-Related Protein of Toxoplasma gondii Is Important for Rhoptry Biogenesis and Virulence. mSphere *2*.

Chen, T.W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A., Schreiter, E.R., Kerr, R.A., Orger, M.B., Jayaraman, V., Looger, L.L., Svoboda, K., and Kim, D.S. (2013). Ultrasensitive fluorescent proteins for imaging neuronal activity. Nature *499*, 295-300.

Clapham, D.E. (2007). Calcium signaling. Cell *131*, 1047-1058.

Coppens, I., and Joiner, K.A. (2003). Host but not parasite cholesterol controls Toxoplasma cell entry by modulating organelle discharge. Mol Biol Cell *14*, 3804-3820.

De Stefani, D., Raffaello, A., Teardo, E., Szabo, I., and Rizzuto, R. (2011). A forty-kilodalton protein of the inner membrane is the mitochondrial calcium uniporter. Nature *476*, 336-340.

Del Carmen, M.G., Mondragon, M., Gonzalez, S., and Mondragon, R. (2009). Induction and regulation of conoid extrusion in Toxoplasma gondii. Cell Microbiol *11*, 967-982.

Donald, R.G., Allocco, J., Singh, S.B., Nare, B., Salowe, S.P., Wiltsie, J., and Liberator, P.A. (2002). Toxoplasma gondii cyclic GMP-dependent kinase: chemotherapeutic targeting of an essential parasite protein kinase. Eukaryot Cell *1*, 317-328.

Endo, T., Sethi, K.K., and Piekarski, G. (1982). Toxoplasma gondii: calcium ionophore A23187-mediated exit of trophozoites from infected murine macrophages. Exp Parasitol *53*, 179-188.

Farwell, D.G., Shera, K.A., Koop, J.I., Bonnet, G.A., Matthews, C.P., Reuther, G.W., Coltrera, M.D., McDougall, J.K., and Klingelhutz, A.J. (2000). Genetic and epigenetic changes in human epithelial cells immortalized by telomerase. Am J Pathol *156*, 1537-1547.

Gincel, D., Zaid, H., and Shoshan-Barmatz, V. (2001). Calcium binding and translocation by the voltage-dependent anion channel: a possible regulatory mechanism in mitochondrial function. The Biochemical journal *358*, 147-155.

Grynkiewicz, G., Poenie, M., and Tsien, R.Y. (1985). A new generation of Ca2+ indicators with greatly improved fluorescence properties. J Biol Chem *260*, 3440-3450.

Hager, K.M., Striepen, B., Tilney, L.G., and Roos, D.S. (1999). The nuclear envelope serves as an intermediary between the ER and Golgi complex in the intracellular parasite Toxoplasma gondii. J Cell Sci *112 (Pt 16)*, 2631-2638.

Haller, T., Dietl, P., Deetjen, P., and Volkl, H. (1996). The lysosomal compartment as intracellular calcium store in MDCK cells: a possible involvement in InsP3-mediated Ca2+ release. Cell Calcium *19*, 157-165.

Hortua Triana, M.A., Marquez-Nogueras, K.M., Fazli, M.S., Quinn, S., and Moreno, S.N.J. (2024). Regulation of calcium entry by cyclic GMP signaling in Toxoplasma gondii. J Biol Chem *300*, 105771.

Hortua Triana, M.A., Marquez-Nogueras, K.M., Vella, S.A., and Moreno, S.N.J. (2018). Calcium signaling and the lytic cycle of the Apicomplexan parasite Toxoplasma gondii. Biochim Biophys Acta Mol Cell Res *1865*, 1846-1856.

Huet, D., and Moreno, S.N.J. (2023). Interorganellar Communication Through Membrane Contact Sites in Toxoplasma Gondii. Contact (Thousand Oaks) *6*, 25152564231189064.

Huynh, M.H., and Carruthers, V.B. (2009). Tagging of endogenous genes in a Toxoplasma gondii strain lacking Ku80. Eukaryot Cell *8*, 530-539.

Kafsack, B.F., Beckers, C., and Carruthers, V.B. (2004). Synchronous invasion of host cells by Toxoplasma gondii. Mol Biochem Parasitol *136*, 309-311.

Kuhlbrandt, W. (2004). Biology, structure and mechanism of P-type ATPases. Nat Rev Mol Cell Biol *5*, 282-295.

Li, Z.H., King, T.P., Ayong, L., Asady, B., Cai, X., Rahman, T., Vella, S.A., Coppens, I., Patel, S., and Moreno, S.N.J. (2021). A plastid two-pore channel essential for inter-organelle communication and growth of Toxoplasma gondii. Nat Commun *12*, 5802.

Liu, J., Pace, D., Dou, Z., King, T.P., Guidot, D., Li, Z.H., Carruthers, V.B., and Moreno, S.N. (2014). A vacuolar-H(+) -pyrophosphatase (TgVP1) is required for microneme secretion, host cell invasion, and extracellular survival of Toxoplasma gondii. Mol Microbiol *93*, 698-712.

Lloyd-Evans, E., Morgan, A.J., He, X., Smith, D.A., Elliot-Smith, E., Sillence, D.J., Churchill, G.C., Schuchman, E.H., Galione, A., and Platt, F.M. (2008). Niemann-Pick disease type C1 is a sphingosine storage disease that causes deregulation of lysosomal calcium. Nat Med *14*, 1247-1255.

Long, S., Brown, K.M., Drewry, L.L., Anthony, B., Phan, I.Q.H., and Sibley, L.D. (2017). Calmodulin-like proteins localized to the conoid regulate motility and cell invasion by Toxoplasma gondii. PLoS Pathog *13*, e1006379.

Lourido, S., and Moreno, S.N. (2015). The calcium signaling toolkit of the Apicomplexan parasites Toxoplasma gondii and Plasmodium spp. Cell Calcium *57*, 186-193.

Lovett, J.L., and Sibley, L.D. (2003). Intracellular calcium stores in Toxoplasma gondii govern invasion of host cells. J Cell Sci *116*, 3009-3016.

Luo, S., Ruiz, F.A., and Moreno, S.N. (2005). The acidocalcisome Ca2+-ATPase (TgA1) of Toxoplasma gondii is required for polyphosphate storage, intracellular calcium homeostasis and virulence. Mol Microbiol *55*, 1034-1045.

Luo, S., Vieira, M., Graves, J., Zhong, L., and Moreno, S.N. (2001). A plasma membrane-type Ca(2+)-ATPase co-localizes with a vacuolar H(+)-pyrophosphatase to acidocalcisomes of Toxoplasma gondii. EMBO J *20*, 55-64.

Mallo, N., Ovciarikova, J., Martins-Duarte, E.S., Baehr, S.C., Biddau, M., Wilde, M.L., Uboldi, A.D., Lemgruber, L., Tonkin, C.J., Wideman, J.G., Harding, C.R., and Sheiner, L. (2021). Depletion of a Toxoplasma porin leads to defects in mitochondrial morphology and contacts with the endoplasmic reticulum. J Cell Sci *134*.

Marquez-Nogueras, K.M., Hortua Triana, M.A., Chasen, N.M., Kuo, I.Y., and Moreno, S.N. (2021). Calcium signaling through a transient receptor channel is important for Toxoplasma gondii growth. Elife *10*.

Miranda, K., Pace, D.A., Cintron, R., Rodrigues, J.C., Fang, J., Smith, A., Rohloff, P., Coelho, E., de Haas, F., de Souza, W., Coppens, I., Sibley, L.D., and Moreno, S.N. (2010). Characterization of a novel organelle in Toxoplasma gondii with similar composition and function to the plant vacuole. Mol Microbiol *76*, 1358-1375.

Mogami, H., Nakano, K., Tepikin, A.V., and Petersen, O.H. (1997). Ca2+ flow via tunnels in polarized cells: recharging of apical Ca2+ stores by focal Ca2+ entry through basal membrane patch. Cell *88*, 49-55.

Moreno, S.N., and Zhong, L. (1996). Acidocalcisomes in Toxoplasma gondii tachyzoites. The Biochemical journal *313 (Pt 2)*, 655-659.

Nagamune, K., Beatty, W.L., and Sibley, L.D. (2007a). Artemisinin induces calcium-dependent protein secretion in the protozoan parasite Toxoplasma gondii. Eukaryot Cell *6*, 2147-2156.

Nagamune, K., Moreno, S.N., and Sibley, L.D. (2007b). Artemisinin-resistant mutants of Toxoplasma gondii have altered calcium homeostasis. Antimicrob Agents Chemother *51*, 3816-3823.

Nagamune, K., and Sibley, L.D. (2006). Comparative genomic and phylogenetic analyses of calcium ATPases and calcium-regulated proteins in the apicomplexa. Mol Biol Evol *23*, 1613-1627.

Oliveira Souza, R.O., Jacobs, K.N., Back, P.S., Bradley, P.J., and Arrizabalaga, G. (2022). IMC10 and LMF1 mediate mitochondrial morphology through mitochondrion-pellicle contact sites in Toxoplasma gondii. J Cell Sci *135*.

Ovciarikova, J., Lemgruber, L., Stilger, K.L., Sullivan, W.J., and Sheiner, L. (2017). Mitochondrial behaviour throughout the lytic cycle of Toxoplasma gondii. Sci Rep *7*, 42746.

Ovciarikova, J., Shikha, S., and Sheiner, L. (2022). Nuclear Interactions: A Spotlight on Nuclear Mitochondrial Membrane Contact Sites. Contact (Thousand Oaks) *5*, 25152564221096217.

Pace, D.A., McKnight, C.A., Liu, J., Jimenez, V., and Moreno, S.N. (2014). Calcium entry in Toxoplasma gondii and its enhancing effect of invasion-linked traits. J Biol Chem *289*, 19637-19647.

Papp, S., Dziak, E., Michalak, M., and Opas, M. (2003). Is all of the endoplasmic reticulum created equal? The effects of the heterogeneous distribution of endoplasmic reticulum Ca2+-handling proteins. J Cell Biol *160*, 475-479.

Park, M.K., Petersen, O.H., and Tepikin, A.V. (2000). The endoplasmic reticulum as one continuous Ca(2+) pool: visualization of rapid Ca(2+) movements and equilibration. EMBO J *19*, 5729-5739.

Periasamy, M., and Kalyanasundaram, A. (2007). SERCA pump isoforms: their role in calcium transport and disease. Muscle Nerve *35*, 430-442.

Petersen, O.H., Courjaret, R., and Machaca, K. (2017). Ca(2+) tunnelling through the ER lumen as a mechanism for delivering Ca(2+) entering via store-operated Ca(2+) channels to specific target sites. J Physiol *595*, 2999-3014.

Phillips, M.J., and Voeltz, G.K. (2016). Structure and function of ER membrane contact sites with other organelles. Nat Rev Mol Cell Biol *17*, 69-82.

Pino, P., Foth, B.J., Kwok, L.Y., Sheiner, L., Schepers, R., Soldati, T., and Soldati-Favre, D. (2007). Dual targeting of antioxidant and metabolic enzymes to the mitochondrion and the apicoplast of Toxoplasma gondii. PLoS Pathog *3*, e115.

Rapizzi, E., Pinton, P., Szabadkai, G., Wieckowski, M.R., Vandecasteele, G., Baird, G., Tuft, R.A., Fogarty, K.E., and Rizzuto, R. (2002). Recombinant expression of the voltage-dependent anion channel enhances the transfer of Ca2+ microdomains to mitochondria. J Cell Biol *159*, 613-624.

Rossi, A.M., and Taylor, C.W. (2020). Reliable measurement of free Ca(2+) concentrations in the ER lumen using Mag-Fluo-4. Cell Calcium *87*, 102188.

Sagara, Y., and Inesi, G. (1991). Inhibition of the sarcoplasmic reticulum Ca2+ transport ATPase by thapsigargin at subnanomolar concentrations. J Biol Chem *266*, 13503-13506.

Sheiner, L., Demerly, J.L., Poulsen, N., Beatty, W.L., Lucas, O., Behnke, M.S., White, M.W., and Striepen, B. (2011). A systematic screen to discover and analyze apicoplast proteins identifies a conserved and essential protein import factor. PLoS Pathog *7*, e1002392.

Smith, J.B., Zheng, T., and Lyu, R.M. (1989). Ionomycin releases calcium from the sarcoplasmic reticulum and activates Na+/Ca2+ exchange in vascular smooth muscle cells. Cell Calcium *10*, 125-134.

Spang, A. (2018). The endoplasmic reticulum-the caring mother of the cell. Curr Opin Cell Biol *53*, 92-96.

Stasic, A.J., Dykes, E.J., Cordeiro, C.D., Vella, S.A., Fazli, M.S., Quinn, S., Docampo, R., and Moreno, S.N.J. (2021). Ca(2+) entry at the plasma membrane and uptake by acidic stores is regulated by the activity of the V-H(+) -ATPase in Toxoplasma gondii. Mol Microbiol *115*, 1054-1068.

Thastrup, O., Cullen, P.J., Drobak, B.K., Hanley, M.R., and Dawson, A.P. (1990). Thapsigargin, a tumor promoter, discharges intracellular Ca2+ stores by specific inhibition of the endoplasmic reticulum Ca2(+)-ATPase. Proc Natl Acad Sci U S A *87*, 2466-2470.

Tomavo, S., Slomianny, C., Meissner, M., and Carruthers, V.B. (2013). Protein trafficking through the endosomal system prepares intracellular parasites for a home invasion. PLoS Pathog *9*, e1003629.

Tomova, C., Humbel, B.M., Geerts, W.J., Entzeroth, R., Holthuis, J.C., and Verkleij, A.J. (2009). Membrane contact sites between apicoplast and ER in Toxoplasma gondii revealed by electron tomography. Traffic *10*, 1471-1480.

Vella, S.A., Calixto, A., Asady, B., Li, Z.H., and Moreno, S.N.J. (2020). Genetic Indicators for Calcium Signaling Studies in Toxoplasma gondii. Methods Mol Biol *2071*, 187-207.

Vella, S.A., Moore, C.A., Li, Z.H., Hortua Triana, M.A., Potapenko, E., and Moreno, S.N.J. (2021). The role of potassium and host calcium signaling in Toxoplasma gondii egress. Cell Calcium *94*, 102337.

Vieira, M.C., and Moreno, S.N. (2000). Mobilization of intracellular calcium upon attachment of Toxoplasma gondii tachyzoites to human fibroblasts is required for invasion. Mol Biochem Parasitol *106*, 157-162.

Weiss, L.M., and Dubey, J.P. (2009). Toxoplasmosis: A history of clinical observations. Int J Parasitol *39*, 895-901.

Wenzel, E.M., Elfmark, L.A., Stenmark, H., and Raiborg, C. (2022). ER as master regulator of membrane trafficking and organelle function. J Cell Biol *221*.

Wuytack, F., Raeymaekers, L., and Missiaen, L. (2002). Molecular physiology of the SERCA and SPCA pumps. Cell Calcium *32*, 279-305.

Yuan, Y., Kilpatrick, B.S., Gerndt, S., Bracher, F., Grimm, C., Schapira, A.H., and Patel, S. (2021). The lysosomotrope GPN mobilises Ca(2+) from acidic organelles. J Cell Sci *134*.

FIGURES AND FIGURE LEGENDS:

Figure 1: The role of extracellular calcium in the filling of intracellular stores. A, scheme depicting the Fura-2-AM loading and the experimental set-up. **B**, *T. gondii* tachyzoites loaded with Fura-2 were in suspension in Ringer buffer with 100 μM EGTA. Thapsigargin (TG) was added at 100 seconds, at two different concentrations (1 and 2 μM). **C**, Similar experimental set-up to the one shown in B. 1.8 mM CaCl₂ was added to the suspension at 100 sec, followed by 2 μM TG at 300 sec (*dark blue trace*). The light blue trace shows the same experiment without the addition of CaCl₂. **D**, similar to C but using 40 μM GPN instead of TG. **E**, same experimental set-up to the one shown in C but adding the potassium ionophore nigericin, 10 μM. **F**, same experimental set-up to the one

shown in C but using the mitochondrial uncoupler CCCP. **G**, quantification of the % increase of cytoplasmic calcium by adding extracellular calcium prior to each indicated reagent (TG, thapsigargin; CP, CCCP; NIG, nigericin). **H**, cytosolic calcium measurements following the addition of 1.8 mM extracellular calcium, 1 µM TG prior to the addition of GPN, 40 µM. **I**, cytosolic Calcium measurements following the addition of 1.8 mM extracellular Calcium, 1 µM TG prior to the addition of 10 µM NIG. The quantification shows the comparison of the cytosolic increase with and without the addition of TG. Data are presented as mean $\pm$ SD for G, H and I. *p value*: unpaired two tailed t test performed in all comparisons.

Figure 2: Ca²⁺ uptake by intracellular stores. A, Scheme showing the loading with MagFluo4-AM followed by permeabilization with digitonin of a *T. gondii* tachyzoite (RH parental strain) suspension (IS, intracellular store). **B**, Fluorescence measurements (see Materials and Methods for specifics) of the suspension of parasites loaded with MagFluo4. MgATP was added at 500 µM at 50 seconds. The bar graph shows the quantification of the slope of the increase in fluorescence after adding the SERCA substrate MgATP. The concentration of free calcium was varied, and it is indicated. The calculation of free calcium was done with MaxChelator. **C**, similar experimental set-up to the one shown in B but the concentration of MgATP was varied at the concentrations indicated in the bar graph. Quantification of the slope of fluorescence increase after adding MgATP is shown in the bar graph. **D**, this experiment was done with 500 µM MgATP and 220 nM CaCl₂. Thapsigargin (TG) was added to inhibit SERCA causing calcium to be released from the store. The concentrations used are indicated. The bar graph shows the negative slope after the addition of TG. **E**, similar to D, but adding ionomycin (IO). The concentrations used are indicated and the slopes were measured after the addition of IO. Data are presented as mean ± SD for B-D. *p value*: unpaired two tailed t test performed in all comparisons.

Figure 3: The sarcoplasmic-endoplasmic reticulum calcium ATPase (SERCA) is essential for *T. gondii* replication. A, Scheme showing the strategy used for the down regulation of the expression of TgSERCA by promoter insertion and regulation by 0.5 µg/ml Anhydrotetracyclin (ATc). The mutants generated were termed *iΔTgSERCA* or *iΔTgSERCA-HA* for the C-terminal tagged one. DHFR, Dihydrofolate reductase gene for selection with pyrimethamine; CAT, chloramphenicol acetyl transferase gene for selection with chloramphenicol. **B**, Western blots of *iΔTgSERCA-HA* lysates grown with and without ATc. The expression of TgSERCA is shown by the signal obtained with the anti-HA antibody. **C**, plaque assays comparing the size of plaques formed by tachyzoites (150/well) of the *iΔTgSERCA* mutant grown with and without 0.5 µg/ml ATc for 8 days. Plaques formed by the parental strain *TatiΔku80* are shown for comparison. **D**, quantification of the size of the plaques presented in C. **E**, replication assays using the *iΔTgSERCA_RFP* mutant. Number of parasites per parasitophorous vacuole (PV) obtained 24 h after infection of fibroblast cells is reported and compared between the mutant grown with and without 0.5 µg/ml ATc. Data represented as mean ± SD. *p value*: unpaired two tailed t test performed in all comparisons. **F**, Average number of parasites

per PV counted at 24 h after the initial infection. The graph shown to the right shows the number of parasites per PV of the *iΔTgSERCA* (+ATc) for 24 or 48 hours after the initial infection. **G**, Invasion assays with the *iΔTgSERCA* mutant treated with ATc for 24 h were done using the red-green assay. **H**, Egress assays with the *iΔTgSERCA* mutant previously grown with or without ATc. Egress was triggered with IO (100 or 50 nM) or saponin (0.01%). The protocol for natural egress uses compound 1 and it is detailed in the methods section.

Figure 4: Organellar calcium pools in the *iΔTgserca* mutant. **A**, cytosolic calcium response to the addition of 1 μM TG. The *iΔTgserca* grown with and without ATc was previously loaded with Fura-2 and the cytosolic Ca²⁺ responses were measured. 1 μM TG was added at 200 sec. The purple trace shows the response of the parental cell line grown without ATc and the pink one shows the response of the same mutant grown with ATc for 24 h. The bar graph shows the analysis of the Δ[Ca²⁺] from three independent experiments. **B**, same experimental set-up as the one in A but adding 1.8 mM extracellular Ca²⁺ at 200 sec. **C**, SERCA activity measured with MagFluo4-loaded tachyzoites of

i∆Tgserca mutant previously grown with or without ATc. Parasites were collected and loaded with MagFluo4-AM as described in the Methods section. Digitonin permeabilized parasites were used in the experiment. The concentration of free calcium in the buffer is 220 nM and MgATP (0.125 mM) was added at 100 sec. The response of the control mutant grown without ATc is shown in purple. The other traces show the response of the mutant grown for either 24 or 48 h with ATc. TG was added at 1 µM TG. The bar graph shows the quantification of the slope after adding MgATP. **D**. Ca^{2+} entry measured in Fura-2 loaded *i∆TgSERCA* (±ATc). 1.8 mM extracellular Ca^{2+} was added at 200 sec. The inset shows the ΔF measurement from three independent experiments and showed no significant difference. **E**, Similar conditions to the ones used in A but adding 100 µM Zaprinast. The bar graph shows the analysis of the $\Delta[Ca^{2+}]$ from three independent experiments. **F**, Similar conditions to the ones used in B but adding 1.8 mM extracellular calcium at 200 sec and 100 µM Zaprinast at 400 sec. The bar graph shows the analysis of the $\Delta[Ca^{2+}]$ from 3 independent experiments. **G**, Same as A but adding 40 µM GPN. The bar graph shows the analysis of the $\Delta[Ca^{2+}]$ from three independent experiments. **H**, Same set-up as in F but adding 1.8 mM Ca^{2+} at 200 sec followed by 40 µM GPN. The bar graph shows the analysis of the $\Delta[Ca^{2+}]$ from three independent experiments. Data are presented as mean ± SD. *p* value: unpaired two tailed t test performed in all comparisons.

Figure 5: Mitochondrial Calcium uptake. **A**, Fluorescence image of *T. gondii* tachyzoites expressing *SOD2-GCaMP6f* (pDT7S4H3-SOD2-GCaMP6f). The generation of this cell line is described in material and methods. **B**, *T. gondii* tachyzoites expressing SOD2-GCaMP6f (5×10^7) permeabilized with digitonin as described in the method's section were tested in a buffer with 100 μM EGTA. Ca^{2+} was added at 100 sec at the final concentrations 1, 10, 20, 50, 100 and 200 μM. Maxchelator was used to calculate the amount of Calcium to be added to attain the indicated concentrations. **C**, ΔF was measured as the change in fluorescence between the baseline and the maximum obtained 20 sec after the addition of Ca^{2+} from three independent biological experiments.

D, Fura-2 loaded tachyzoites of the SOD2-GCaMP6f expressing mutant in suspension. The experimental set-up was identical to the one described in Fig. 1A-B. 1.8 mM $CaCl_2$ was added at 400 sec. Fluorescence measurements were done using the conditions for Fura-2. **E**, GCaMP6 fluorescence measurements of parasites expression SOD2-GCaMP6f in their mitochondrion. The fluorometer conditions were set to measure GCaMP6. **F**, 1 µM TG was added at 100 sec followed by 1.8 mM $CaCl_2$ at 400 sec. Fura2 loaded parasites and Fura2 conditions were used. **G**, Same additions as in F but measuring fluorescence of GCaMP6. **H**, Response to the addition of 1 µM TG of the *i∆Tgserca-sod2-gcamp6f* mutant (transfected with pCTH3-SOD2-GCaMP6f plasmid) previously grown in the presence (*pink trace*) or absence (*blue trace*) of ATc. The conditions for the fluorescence measurements are the same used in part G with intact parasites. The bar graph shows the ∆F measurements from three independent experiments. **I**, same as H but using 40 µM GPN. **J**, Same as H but using 100 µM Zaprinast. **K**, Same as H but using 1 µM Ionomycin. Data are presented as mean ± SD from 3 independent biological experiments. *p value*: unpaired two tailed t test performed in all comparisons.

Figure 6: ER membrane contacts with the mitochondrion and the plant like vacuolar compartment (PLVAC). **A**, Super-resolution IFAs of intracellular parasites with the mitochondrion labeled with the αTom40 (green, 1:20,000) antibody and the ER labeled with the αTgcalumenin antibody (an ER calcium binding protein) (red, 1:1000) or the TgSERCA (red 1:1000). **B**, Immunofluorescence (IFA) of extracellular tachyzoites with the same antibodies used for part A. The mitochondrion and ER membranes interact in several regions. **C**, The PLVAC was localized with the αVP1 antibody (green, 1:200) or the αTgCPL antibody (green, 1:500). The ER membranes were labeled with the αTgERC antibody (red). The spots of contact between the ER and the PLVAC are yellow. Scale bars in A-C represent 5 μm. **D**, Transmission Electron Microscopy imaging of the contact sites formed between ER and PLVAC, ER and Apicoplast, ER and mitochondrial. Size bars are 100 nm.

Figure 7: Model showing calcium entry through two different types of Ca^{2+} channels, uptake to the ER by SERCA and distribution to the other organelles via transfer from the ER to the mitochondria, PLVAC and apicoplast. In intracellular parasites, the mitochondria is in close contact with the ER which is constitutively leaking Ca^{2+} into the cytosol. The mitochondria take up Calcium from the ER through an unknown mechanism. VDAC could be involved in the transfer through the outer mitochondrial membrane (Mallo *et al.*, 2021). The PLVAC interacts with the ER and may also interact with the mitochondrion and the apicoplast. TgA1, a calcium ATPase previously characterized may be the pump involved in uptake (Luo *et al.*, 2001; Luo *et al.*, 2005). The mechanism of release is unknown. The TPC was shown to be involved in the formation of contacts between the ER and the apicoplast (Li *et al.*, 2021). Ca^{2+} could leak from the ER through the TgTRPPL-2 channel previously described (Marquez-Nogueras *et al.*, 2021).