The evolution of signalling and eavesdropping in plant-fungal networks

Thomas W. Scott[1,2,3], E. Toby Kiers[4,5] and Stuart A. West[1]

[1]Department of Biology, University of Oxford, 11a Mansfield Road, Oxford, United Kingdom

[2]School of Biology, University of St Andrews, Dyers Brae, St Andrews, United Kingdom

[4]A-life, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands

[5]Society for the Protection of Underground Networks, SPUN, Dover, DE 19901, USA

[3]Author for correspondence: thomas.scott@biology.ox.ac.uk

Abstract

Experiments have shown that when one plant is attacked by a pathogen or herbivore, this can lead to other plants connected to the same mycorrhizal network upregulating their defence mechanisms. It has been hypothesised that this represents signalling, with attacked plants producing a signal to warn other plants of impending harm. We examined the evolutionary plausibility of this and other hypotheses theoretically. We found that the evolution of plant signalling about an attack requires restrictive conditions, and so will rarely be evolutionarily stable. The problem is that signalling about an attack provides a benefit to competing neighbours, even if they are kin, and so reduces the relative fitness of signalling plants. Indeed, selection is often more likely to push plant behaviour in the opposite direction – with plants signalling dishonestly about an attack that has not occurred, or by suppressing a cue that they have been attacked. Instead, we show that there are two viable alternatives that could explain the empirical data: (1) the process of being attacked leads to a cue (information about the attack) which is too costly for the attacked plant to fully suppress; (2) mycorrhizal fungi eavesdrop on their host plants, detect when they are attacked, and then the fungi signal this information to warn other plants in their network. Our results suggest the empirical work that would be required to distinguish between these possibilities.

Significance statement

Experiments have shown that when one plant is attacked by a herbivore, this can lead to other plants connected to the same mycorrhizal network upregulating their defence mechanisms. It has been hypothesised that this represents signalling, with attacked plants producing a signal to warn other plants of impending harm. We found theoretically that plant warning signals are

rarely evolutionarily stable. Instead, we identify two viable alternatives that could explain the empirical data: (1) being attacked leads to a cue (information about the attack) which is too costly for the attacked plant to suppress; (2) mycorrhizal fungi eavesdrop on their host plants, detect when they are attacked, and then the fungi signal this information to warn other plants in their network.

Introduction

Mycorrhizal fungi form symbiotic associations with plant roots, trading nutrients such as phosphorous and nitrogen for plant-derived carbon. These fungi form physical networks of mycelium that can connect roots of different plants and act as potential routes for signalling between those plants (1–5). Several laboratory experiments have provided clear evidence that when one plant in a mycorrhizal network is attacked that this leads to other plants in the network upregulating their defence mechanisms. For example, when a tomato plant is infested with a leaf chewing caterpillar, tomato plants connected to the same network will increase their production of defence enzymes (6). It has been hypothesised that this pattern represents a 'warning signal' in which the attacked plant actively signals to other plants using chemicals transported via the mycorrhizal network. This work has even fuelled narratives in the media that forest trees use mycorrhizal networks to warn other trees of impending danger (7–9).

However, the evolutionary plausibility of this signalling hypothesis remains unclear (10–28). A signal is defined as '*any act or structure that alters the behaviour of other organisms, which evolved owing to that effect, and which is effective because the receiver's response has also evolved*' (29, 30). Consequently, signalling is a form of cooperation, that is only favoured when it provides a benefit to both the sender and the receiver (29, 31). Plants compete with neighbours for resources such as sunlight and nutrients, and so helping a neighbour could be costly to a potential signaller (12). Neighbouring plants could be relatives, which could provide a kin-selected benefit of signalling, but this may require restrictive conditions to also avoid competition between relatives, which can reduce or even negate any benefit of helping relatives (32–36). For instance, a low migration rate could cause relatives to signal to each other, leading to a kin-selected benefit, but it could also cause relatives to compete with each other, negating this benefit. Consequently, it is not clear if signalling about attack would be evolutionarily stable between neighbouring plants who are both relatives and competitors.

Another potential problem for the plant signalling hypothesis is that there are at least two alternative explanations that could possibly explain the experimental data (Fig. 1) (12, 37). One possibility is that the neighbouring plants could be detecting a cue that another plant is being attacked (12). A cue is defined by when a receiver uses some feature of the sender to guide their own behaviour, but this feature has not evolved for that purpose (30). An example of a cue is when a mosquito searching for a mammal to bite will fly up wind if it detects carbon dioxide (30). Mosquitos use carbon as a cue of the presence of a source of blood, but mammals do not produce carbon dioxide to signal their presence to mosquitoes (getting bitten is costly!). In this case, all that is required is that the damage caused by attack causes a plant to produce some cue, such as a released chemical (smell).

A second possible alternative explanation of the experimental data is that the fungus can detect when a plant is being attacked, and that they then signal this information to the other plants to which they are connected (15). This would therefore involve fungi 'eavesdropping' on a cue of predator attack, and it would be the fungus not the attacked plant that is signalling. Fungi could be selected to eavesdrop and signal to plants in their network because they actively trade resources with those individuals, and so could benefit from helping keep them in better condition, to make better trading partners (12, 38–41).

Figure 1. Hypotheses for information transfer in plant-fungal networks. Two plants are connected *via* a mycorrhizal network. One plant is under attack, and this information may be transferred *via* the mycorrhizal network to the other plant, allowing it to upregulate its defence mechanisms. There are three hypotheses regarding how this information is transferred: (*blue*) signalling by the predated plant; (*cyan*) cues, which are potentially vulnerable to suppression by the predated plant; (*orange*) eavesdropping and signalling by the mycorrhizal network.

The difference between these hypotheses matters because they represent different evolutionary outcomes, are favoured for different reasons, and require different empirical tests (30, 42, 43). With plant signalling, the problem is to determine how both the sender and receiver benefit. What conditions would be required for honest signalling to be evolutionary stable, and how could this seemingly altruistic act of warning neighbours be explained in the context of plant-plant competition? Could plants even be favoured to signal dishonestly to harm competitors? In contrast, with a cue, we would need to ask what stimulates the cue to be produced? If the cue provides a benefit to neighbours, then does this cue impose a cost to the producer, by aiding

their competitors? Why don't plants suppress a cue? Finally, in the case of mycorrhizal eavesdropping, we would need to ask why the fungus would be selected to eavesdrop and then produce a signal in response? And what are they eavesdropping on?

We investigated these different hypotheses theoretically. Our aim was to determine their evolutionary plausibility, as well as define the empirical work required to test among them. We first consider selection on plants, examining whether: (a) an attacked plant would be selected to signal that it had been attacked; (b) plants can be selected to signal dishonestly about attack, to harm competitors; (c) plants can be selected to suppress a cue that they are being attacked. We then examine selection on mycorrhizal fungi and ask whether they can be selected to eavesdrop on plants, to determine whether they are being attacked, and then signal that information to neighbouring plants.

Results

We constructed a series of theoretical models to examine selection on signals and cues, from the perspective of both plants and mycorrhizal fungi. We constructed deliberately simple models which are easy to interpret and can be applied across diverse species (44–46).

Plant signalling model. We first examined whether an attacked plant can be selected to produce a warning signal. We assume an infinite population of individual plants, split into patches (demes) of size N (infinite island model)(47). Each generation, with probability p, the population is attacked, for instance by a herbivore or pathogen (with probability $1-p$, the population is not attacked). In generations where the population is attacked, a random individual on each patch, i, is initially attacked, and suffers a fecundity cost of d. This individual then invests x_i into the production of a signal, resulting in an additional fecundity cost of cx_i, where c is the marginal cost of signal production.

The signal produced by the initially-attacked individual is transferred to the other $N-1$ individuals on the patch (receivers). We make no assumptions about how the signal is transferred, so it may be transferred through a common mycorrhizal network, the air, or any other mechanism (19). The signal warns the receivers that an attack is imminent. Consequently, the receivers can prepare for being attacked, and defend themselves, meaning they suffer a

reduced fecundity cost of being attacked. The extent to which receivers, j, respond to the signal, preparing for predator-attack, rather than ignoring the signal, is given by y_j. Preparation for being attacked (defence) incurs a fecundity cost of sx_iy_j, where s is the marginal cost of defence, but it reduces the cost of being attacked, which is now given by $d(1-x_iy_j)$. We assume that the cost of defence is less than being attacked ($s<d$), which ensures that individuals are favoured to defend themselves against attack (rather than let themselves be damaged).

We then allow individuals to produce offspring (juvenile haploid clones) in proportion to their fecundity. A random sample of N juvenile individuals are chosen, for each patch, to survive and form the next adult population (local population regulation). After population regulation, a proportion of the (new) adult population, m, migrate to different and random patches. The remaining proportion, $1-m$, remain on their local patch. This lifecycle then iterates over many generations until an evolutionary end point is reached. In Appendix J, we determined the equilibrium (ESS) levels of signalling investment (x^*) and signal-response (y^*).

Signalling is not evolutionarily stable. We found that individual plants were not favoured to produce warning signals ($x^*=0$) (Fig. 2). Any tendency to honestly signal an impending attack will ultimately be removed by natural selection, to avoid providing a benefit to competitors. Honest signalling is a helping behaviour, that increases the fitness of social partners by warning them about an impending attack. Helping can be potentially favoured if it is directed towards relatives that share genes for helping, termed kin selection (32, 48, 49). When the migration rate (m) is lower, individuals on a patch will be more closely related. However, a lower migration rate also means that the individuals receiving help are in stronger local competition with the helper, which negates the benefit of helping relatives (33). Put simply, there is no benefit in helping one's relative if this comes at an equal cost to another relative (34, 50, 51). We showed in Appendix K that this result also holds if population regulation occurs after migration, rather than before, and if plants signal even if they are not the first plant on the patch to be attacked (obligate versus facultative signalling).

Figure 2. Plant signalling is not evolutionarily stable. Signalling is favoured if plant fitness (w) increases with signalling investment (x), in other words if the selection differential $\frac{dw}{dx}$ is positive. The solid black line is a reference line that plots $\frac{dw}{dx}$ for arbitrary parameter values. Coloured lines show the effect of increasing parameter values. The key thing to note about this figure is that, irrespective of the parameter values (which are varied on the x axis and on the coloured lines), the selection differential is never positive, meaning signalling is never favoured. We assumed: (*reference values*) N=3 (group size), c=0.1 (cost of signalling), s=0 (defence cost), m=0.3 (migration rate); (*alternative values*) N=7, g=1, c=0.6, s=0.05, m=1; x=0, y=1, z=0 (signals initially rare, honest and responded-to).

Alternative population structures. Our model assumed a simple life history where offspring (seeds) can disperse after reproduction. This seems reasonable for plants, where the lack of movement after they have started growing will mean that any neighbours potentially helped will also be competitors. We also assumed that the parameters of the life cycle such as the migration rate determined both the relatedness (genetic similarity) between different individuals on a patch and who competition occurs between. As an alternative approach, we can construct an 'open' model that detaches these model parameters and keeps them as free variables (52). While this can be artificial, it can suggest the kind of conditions that would be required for honest signalling to be favoured. We showed in Appendix K that honest signalling can be favoured when individuals on a patch are highly related, but local competition does not occur between relatives. Although theoretically this is possible, it is hard to imagine a scenario

in which low dispersal could lead to high relatedness among interacting plants, but not local competition. Other hypotheses that could be tested empirically include kin discrimination, if there was a way to only signal to related plants through the fungal network; certain forms of generation overlap; or a private (direct) benefit to helping reduce the local population of insect pests (32, 52–54).

Can dishonest signalling be favoured? Our model also allowed us to investigate the opposite of honest signalling – whether plants could be favoured to produce dishonest signals, where they signalled an attack when this had not happened. This could potentially benefit the dishonestly-signalling plant if it sufficiently reduced the fecundity of their local competitors. In generations where the population is not attacked, a random individual on each deme, i, is given the opportunity to signal an attack even though no attack had occurred (dishonest signalling). This individual invests cx_iz_i into signal production, where z_i denotes dishonesty. The receivers, j, consequently suffer a defence cost of $sx_iy_jz_i$. In Appendix J, we determined the equilibrium (ESS) level of signal dishonesty (z^*) and how it influences signal-response (y^*).

We found that individuals could be favoured to produce dishonest signals, where they signalled an attack even when none had occurred. We found that plants could gain a benefit from dishonest signalling, because it harms their local competitors. However, this leads to selection on the receiver plants to ignore signals, and so dishonest signalling would only be transient, not evolutionarily stable. Dishonest signalling has not been empirically observed, but it has also not been tested for. More generally, dishonest signals can in principle be stable within signalling systems when they are at a low frequency, and so do not completely remove the benefit of responding to signals. For example, fork-tailed drongos produce alarm calls to signal to meerkats when a predator is approaching; but also occasionally produce false alarm calls, in the absence of a predator, to steal food left by the fleeing meerkats (55, 56).

Cue-suppression model. We then examined the case where an attacked plant produces a cue when it is being attacked. This would include chemicals that are produced or released by herbivory or pathogenicity, and which are transmitted by any route, including the air or fungal network. The production of a cue could provide another explanation for the experimental data showing upregulation of defences by the neighbours of attacked plants. Can the plant be selected to suppress this cue? We test this by modifying the assumptions of the previous model so that now, after an individual is attacked, it releases a *cue* rather than a *signal*. The cue

indicates to the other N-1 individuals on the patch (receivers) that an attack is imminent. However, we assume that the attacked plant may pay a cost to suppress the cue, to stop the receivers from learning that an attack is imminent. So now, x_i denotes investment into cue-suppression, rather than signalling. We assume that plants respond to cues by upregulating their defences. In Appendix E, we determined the equilibrium (ESS) level of cue-suppression (x^*).

Cue-suppression is evolutionarily stable. We found that individuals could be favoured to suppress cues (information) of an attack (Fig. 3a). Plants can gain a benefit from cue-suppression because it harms their local competitors (by avoiding helping them). Cue-suppression is favoured when the benefit incurred due to competitors being less able to defend themselves against the attack is smaller than the cost of suppressing the cue (Fig. 3a). More generally, we showed in Appendix H that cue-suppression is favoured with low relatedness between individuals on a patch and local competition. However, since there are many situations in which cue suppression is not favoured, the detection of a cue that another plant is being attacked remains an evolutionarily viable explanation for the experimental data (12).

Fungal eavesdropping model. Finally, we examine selection on mycorrhizal fungi, and ask whether they can be selected to eavesdrop on plants, to determine whether they are being attacked, and then signal that information to the other plants they are connected to (15). To examine this, we modify the assumptions of the first 'plant signalling' model, so that now, as well as the (infinite) population of plants, we additionally model an infinite population of fungi.

The population is split into patches comprising N plants connected by one fungus (which means that each plant is associated with just one fungus). When a plant is attacked, the fungus may 'eavesdrop' on warning signals produced by the initially-attacked plant and signal this information to the other plants it is connected to (recipients). As before, signals may be dishonest and ignored. The difference is that, now, signal investment and signal dishonesty are fungal rather than plant traits (signal response is still a plant trait). We assume that fungi gain a benefit by being connected to plant partners in better condition ('fitter'), because such partners will be better trade partners (38–40, 57). Specifically, a fungus gains a benefit of $bf_{partner}$, where $f_{partner}$ gives the average condition (fecundity) of the plants it is connected to. Fungi reproduce by producing offspring (juveniles) in proportion to their fecundity; each juvenile migrates to a random deme, and a random juvenile is chosen for each deme to survive

and form the next adult fungal population. In Appendix L, we determined the equilibrium (ESS) levels of fungal signalling investment ($\tilde{x}^*$) and plant signal-response (y^*).

Fungal eavesdropping & signalling is evolutionarily stable. We found that mycorrhizal fungi could be selected to eavesdrop on plants, to determine whether they are being attacked, and then signal that information to (warn) the other plants they are connected to. We found that fungi could gain a benefit from this, because it helps their trade partners defend themselves against attacks, allowing them to better transfer essential resources like nitrogen and phosphorus to the fungus. Mathematically, fungal eavesdropping and signalling is favoured when the costs to the fungus of eavesdropping and signalling are less than the benefit to the fungus of having higher quality trade partners (Fig. 3b). In contrast, we showed in Appendix L that fungi are never favoured to signal dishonestly, by tricking plants into upregulating their defences when no attack is imminent, because this would reduce the quality of their trade partners (58).

One possibility we have not considered is if multiple fungal individuals associate with each other or the same plant (59, 60). Previous theory has shown that when more fungi and plants interact within a network this favours more efficient resource trading, and hence helps make trading evolutionarily stable (57). More complex network interactions could similarly influence the evolution of eavesdropping and signalling by fungi (61). For example, could larger networks lead to fungi eavesdropping on other fungi, and hence increased selection for fungal signalling? This would be especially the case if it helped facilitate trading across the network. Or could larger networks select for fungal cheating, where some fungi avoid any cost of signalling? Answering such questions would require an alternative modelling approach, that examined more complex networks. Nonetheless, more complex situations do not change that fungi can gain a benefit from eavesdropping on plants and then passing that information to other plants that they trade with.

Figure 3. The evolution of cue-suppression and eavesdropping. Traits are favoured above the lines, and disfavoured below them (*i.e.* the lines mark the boundary where the trait changes from being favoured / disfavoured, for a given set of parameter values). Solid black lines are reference lines generated with arbitrary parameter values. Coloured lines show the effect of increasing parameter values. (**A**) Suppression of cues of predator-presence is favoured when the net benefit of predator-defence (*d-s*) is greater than the trait cost (*c*). (**B**) Mycorrhizal eavesdropping and signalling is more likely to be favoured when: the benefit of high-quality trade-partners (*b*), *N* (group size) & *d* (predation cost) are high; *s* (defence cost) & *c* are low. We assumed: (A) any value for *N*; (B, *reference values*) *N*=3, *b*=1; (B, *alternative values*) *N*=5, *b*=1.5.

Discussion

We have shown that plants: (1) are unlikely to be selected to signal to their neighbours about the presence of predators (Fig. 2); (2) can be favoured to produce dishonest signals, where they signalled an attack when none had occurred, but that this will select for other plants to ignore signals; (3) can be favoured to pay resources to suppress any information (cue) to neighbours about attack (Fig. 3a). In contrast, mycorrhizal fungi can be selected to eavesdrop on their host plants, detect when they are attacked, and then signal (warn) other plants in their network (Fig. 3b).

Our results do not support the hypothesis of warning signals by plants, passed *via* any route, including common mycorrhizal networks or the air (10–28). This is because these warnings would benefit neighbouring competitors, to the cost of the signalling individual. Furthermore,

we found that not only are plants not expected to signal, but that they can be selected to signal dishonestly or to actively suppress any cues of being predated on. Dishonest signalling or suppression of cues is favoured to harm or avoid helping neighbouring competitors.

For a helping behaviour to be favoured, such as signalling a warning of herbivore or pathogen attack, it would require that helping and competition occur at different scales (economic neighbourhoods) or some method of kin recognition / discrimination (34, 35, 50, 51, 62–64). This seems relatively unlikely for warning signalling in plants, where local interactions can involve both cooperation and competition for resources. The same problem of local competition has been demonstrated empirically in other organisms, such as when fig wasps compete for mates in the closed environment of a fig fruit, or when bacteria compete locally for resources (65, 66). In animals, warnings about the presence of predators have also been argued to be favoured because they also reduce predation on the individual making the warning call – this is different from plants, where the warning arises after attack (67–69). In contrast, local competition favours harming behaviours, because harming neighbours can decrease competition for resources (35, 70–72). Alternative modelling approaches could include explicit spatial structures, such as on a graph or lattice, but these have been shown to lead to analogous results (62, 73–75).

In contrast to the situation for plants, we found that mycorrhizal fungi can be favoured to eavesdrop on their host plants, detect a cue of when they are attacked, and then signal this to (warn) other plants in their network (15). Fungi are selected to eavesdrop and signal because defended plants will maintain better condition and hence become better trade partners. Previous theory has shown that selection for fungi and plants to trade resources with each other is increased when multiple plants and fungi interact in the same network, because this stabilises efficient trading (57).

To conclude, we examined hypotheses for the empirical result that when one plant in a mycorrhizal network is attacked that this leads to other plants in the network upregulating their defence mechanisms (1–6). Our modelling suggests that this is more likely to represent either: a cue produced by plants that is too costly to suppress, or fungi eavesdropping on plants, and then signalling to other plants. Further experiments could test between these possibilities, by examining the underlying mechanism in networks or experimental multiple root systems. How is information conveyed? Where does that information arise from? What are the fitness

consequences for all the individuals involved? More work is needed to understand if and how these ideas can be used in an applied context.

Materials and methods

In the Supplementary Information, comprising Appendices A–L, we analyse and interpret a series of models. The models differ from each other in the assumptions they make about: lifecycle; how demography affects relatedness; what traits can evolve. Appendices J & K present our most general plant signalling models, where signals can evolve to be dishonest and / or ignored. Appendices A–D present special cases of these models, where signals are forced to be honest and responded-to. Appendices E–H present cue-suppression models. Appendix L presents fungal eavesdropping models. Appendix I interprets the models presented in Appendices A–H, setting them in the context of the wider literature on the evolution of helping and harming.

Acknowledgements

We thank Anna Dewar, Ashleigh Griffin, Vasilis Kokkoris, Asher Leeks and George Shillcock for comments on the manuscript. We thank the ERC (834164 & 771387), NERC (NE/V011537/1), HFSP Program Grant (RGP 0029), NWO-VICI (202.012), NWO-Microp program, and Ammodo Foundation for funding.

References

1. B. R. Vanishri, M. Khandappagol, Belowground Plant-Plant Interaction by Common Mycorrhizal Network (CMN). *Int J Curr Microbiol Appl Sci* **7**, 1691–1705 (2018).

2. S. W. Simard, "Mycorrhizal Networks Facilitate Tree Communication, Learning, and Memory" in (2018), pp. 191–213.

3. D. Johnson, L. Gilbert, Interplant signalling through hyphal networks. *New Phytologist* **205**, 1448–1453 (2015).

4. M. A. Gorzelak, A. K. Asay, B. J. Pickles, S. W. Simard, Inter-plant communication through mycorrhizal networks mediates complex adaptive behaviour in plant communities. *AoB Plants* **7**, plv050 (2015).

5. E. K. Barto, J. D. Weidenhamer, D. Cipollini, M. C. Rillig, Fungal superhighways: do common mycorrhizal networks enhance below ground communication? *Trends Plant Sci* **17**, 633–637 (2012).

6. Y. Y. Song, *et al.*, Hijacking common mycorrhizal networks for herbivore-induced defence signal transfer between tomato plants. *Sci Rep* **4**, 3915 (2014).

7. D. Immerwahr, Mother trees and socialist forests: is the 'wood-wide web' a fantasy? *The Guardian* (2024).

8. A. Irwin, The 'Mother Tree' idea is everywhere — but how much of it is real? *Nature* **627**, 718–721 (2024).

9. K. Flinn, The Idea That Trees Talk to Cooperate Is Misleading. *Sci Am* (2021).

10. J. Son, S. Jang, N. Mathevon, C. Ryu, Is plant acoustic communication fact or fiction? *New Phytologist* (2024). https://doi.org/10.1111/nph.19648.

11. M. C. Rillig, A. Lehmann, L. Lanfranco, T. Caruso, D. Johnson, Clarifying the definition of common mycorrhizal networks. *Funct Ecol* (2024). https://doi.org/10.1111/1365-2435.14545.

12. A. van't Padje, M. D. Whiteside, E. T. Kiers, Signals and cues in the evolution of plant–microbe communication. *Curr Opin Plant Biol* **32**, 47–52 (2016).

13. M. A. Gorzelak, "Kin-selected signal transfer through mycorrhizal networks in Douglas-fir.," PhD thesis, University of British Columbia. (2017).

14. Y. Y. Song, S. W. Simard, A. Carroll, W. W. Mohn, R. Sen Zeng, Defoliation of interior Douglas-fir elicits carbon transfer and stress signalling to ponderosa pine neighbors through ectomycorrhizal networks. *Sci Rep* **5**, 8495 (2015).

15. Z. Babikova, D. Johnson, T. Bruce, J. Pickett, L. Gilbert, Underground allies: How and why do mycelial networks help plants defend themselves? *BioEssays* **36**, 21–26 (2014).

16. M. G. A. Van Der Heijden, T. R. Horton, Socialism in soil? The importance of mycorrhizal fungal networks for facilitation in natural ecosystems. *Journal of Ecology* **97**, 1139–1150 (2009).

17. Z. Babikova, *et al.*, Underground signals carried through common mycelial networks warn neighbouring plants of aphid attack. *Ecol Lett* **16**, 835–843 (2013).

18. G. Boyno, S. Demir, Plant-mycorrhiza communication and mycorrhizae in inter-plant communication. *Symbiosis* **86**, 155–168 (2022).

19. M. U. Rasheed, A. Brosset, J. D. Blande, Tree Communication: the Effects of "Wired" and "Wireless" Channels on Interactions with Herbivores. *Current Forestry Reports* **9**, 33–47 (2022).

20. S. W. Simard, "Mycorrhizal Networks Facilitate Tree Communication, Learning, and Memory" in (2018), pp. 191–213.

21. J. Leake, *et al.*, Networks of power and influence: the role of mycorrhizal mycelium in controlling plant communities and agroecosystem functioning. *Canadian Journal of Botany* **82**, 1016–1045 (2004).

22. A. F. Figueiredo, J. Boy, G. Guggenberger, Common Mycorrhizae Network: A Review of the Theories and Mechanisms Behind Underground Interactions. *Frontiers in Fungal Biology* **2** (2021).

23. Y. Y. Song, *et al.*, Interplant Communication of Tomato Plants through Underground Common Mycorrhizal Networks. *PLoS One* **5**, e13324 (2010).

24. Y. Song, M. Wang, R. Zeng, K. Groten, I. T. Baldwin, Priming and filtering of antiherbivore defences among Nicotiana attenuata plants connected by mycorrhizal networks. *Plant Cell Environ* **42**, 2945–2961 (2019).

25. P.-L. Alaux, F. Naveau, S. Declerck, S. Cranenbrouck, Common Mycorrhizal Network Induced JA/ET Genes Expression in Healthy Potato Plants Connected to Potato Plants Infected by Phytophthora infestans. *Front Plant Sci* **11** (2020).

26. L. Gilbert, D. Johnson, "Plant–Plant Communication Through Common Mycorrhizal Networks" in (2017), pp. 83–97.

27. L. Gilbert, D. Johnson, Plant-mediated 'apparent effects' between mycorrhiza and insect herbivores. *Curr Opin Plant Biol* **26**, 100–105 (2015).

28. Z. Babikova, D. Johnson, T. Bruce, J. Pickett, L. Gilbert, How rapid is aphid-induced signal transfer between plants via common mycelial networks? *Commun Integr Biol* **6**, e25904 (2013).

29. A. Grafen, Biological signals as handicaps. *J Theor Biol* **144**, 517–546 (1990).

30. J. Maynard Smith, D. Harper, *Animal Signals* (Oxford University Press, 2003).

31. M. Kirkpatrick, M. J. Ryan, The evolution of mating preferences and the paradox of the lek. *Nature* **350**, 33–38 (1991).

32. W. D. Hamilton, The genetical evolution of social behaviour. I & II. *J Theor Biol* **7**, 1–52 (1964).

33. P. D. Taylor, Altruism in viscous populations-an inclusive fitness model. *Evol Ecol* **6**, 352–356 (1992).

34. S. A. West, I. Pen, A. S. Griffin, Cooperation and Competition Between Relatives. *Science (1979)* **296**, 72–75 (2002).

35.   L. Lehmann, F. Rousset, How life history and demography promote or inhibit the evolution of helping behaviours. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 2599–2617 (2010).

36.   A. F. G. Bourke, *Principles of Social Evolution* (Oxford University Press, 2011).

37.   J. Karst, M. D. Jones, J. D. Hoeksema, Positive citation bias and overinterpreted results lead to misinformation on common mycorrhizal networks in forests. *Nat Ecol Evol* **7**, 501–511 (2023).

38.   E. T. Kiers, *et al.*, Reciprocal Rewards Stabilize Cooperation in the Mycorrhizal Symbiosis. *Science (1979)* **333**, 880–882 (2011).

39.   M. W. Schwartz, J. D. Hoeksema, Specialization and resource trade: biological markets as a model of mutualisms. *Ecology* **79**, 1029–1038 (1998).

40.   J. L. Bronstein, Conditional outcomes in mutualistic interactions. *Trends Ecol Evol* **9**, 214–217 (1994).

41.   J. L. Bronstein, Our Current Understanding of Mutualism. *Q Rev Biol* **69**, 31–51 (1994).

42.   N. Davies, J. Krebs, S. West, *An Introduction to Behavioural Ecology* (John Wiley & Sons, 2012).

43.   S. P. Diggle, A. Gardner, S. A. West, A. S. Griffin, Evolutionary theory of bacterial quorum sensing: when is a signal not a signal? *Philosophical Transactions of the Royal Society B: Biological Sciences* **362**, 1241–1249 (2007).

44.   A. Grafen, "Modelling in behavioural ecology" in *Behavioural Ecology: An Evolutionary Approach*, (1991), pp. 5–31.

45.   G. A. Parker, J. M. Smith, Optimality theory in evolutionary biology. *Nature* **348**, 27–33 (1990).

46.   R. M. May, Uses and Abuses of Mathematics in Biology. *Science (1979)* **303**, 790–793 (2004).

47.   F. Rousset, *Genetic Structure and Selection in Subdivided Populations (MPB-40)* (Princeton University Press, 2004).

48.   A. F. G. Bourke, Hamilton's rule and the causes of social evolution. *Philosophical Transactions of the Royal Society B: Biological Sciences* **369**, 20130362 (2014).

49.   A. F. G. Bourke, The validity and value of inclusive fitness theory. *Proceedings of the Royal Society B: Biological Sciences* **278**, 3313–3320 (2011).

50.   S. A. Frank, *Foundations of Social Evolution* (Princeton University Press, 1998).

51.   D. C. Queller, Genetic relatedness in viscous populations. *Evol Ecol* **8**, 70–73 (1994).

52. G. A. Cooper, S. R. Levin, G. Wild, S. A. West, Modeling relatedness and demography in social evolution. *Evol Lett* **2**, 260–271 (2018).

53. S. Alizon, P. Taylor, Empty sites can promote altruistic behavior. *Evolution (N Y)* **62**, 1335–1344 (2008).

54. A. L. File, J. Klironomos, H. Maherali, S. A. Dudley, Plant Kin Recognition Enhances Abundance of Symbiotic Microbial Partner. *PLoS One* **7**, e45648 (2012).

55. T. P. Flower, M. Gribble, A. R. Ridley, Deception by Flexible Alarm Mimicry in an African Bird. *Science (1979)* **344**, 513–516 (2014).

56. T. Flower, Fork-tailed drongos use deceptive mimicked alarm calls to steal food. *Proceedings of the Royal Society B: Biological Sciences* **278**, 1548–1555 (2011).

57. G. A. K. Wyatt, E. T. Kiers, A. Gardner, S. A. West, A biological market analysis of the plant-mycorrhizal symbiosis. *Evolution (N Y)* **68**, 2603–2618 (2014).

58. R. Ferriere, J. L. Bronstein, S. Rinaldi, R. Law, M. Gauduchon, Cheating and the evolutionary stability of mutualisms. *Proc R Soc Lond B Biol Sci* **269**, 773–780 (2002).

59. M. M. Hart, R. J. Reader, J. N. Klironomos, Plant coexistence mediated by arbuscular mycorrhizal fungi. *Trends Ecol Evol* **18**, 418–423 (2003).

60. M. M. Hart, R. J. Reader, J. N. Klironomos, Life-history strategies of arbuscular mycorrhizal fungi in relation to their successional dynamics. *Mycologia* **93**, 1186–1194 (2001).

61. P. J. CaraDonna, *et al.*, Interaction rewiring and the rapid turnover of plant–pollinator networks. *Ecol Lett* **20**, 385–394 (2017).

62. A. Grafen, M. Archetti, Natural selection of altruism in inelastic viscous homogeneous populations. *J Theor Biol* **252**, 694–710 (2008).

63. T. W. Scott, G. Wild, How to make an inclusive-fitness model. *Proceedings of the Royal Society B: Biological Sciences* **290** (2023).

64. P. D. Taylor, Inclusive fitness in a homogeneous environment. *Proc R Soc Lond B Biol Sci* **249**, 299–302 (1992).

65. A. S. Griffin, S. A. West, A. Buckling, Cooperation and competition in pathogenic bacteria. *Nature* **430**, 1024–1027 (2004).

66. S. A. West, M. G. Murray, C. A. Machado, A. S. Griffin, E. A. Herre, Testing Hamilton's rule with competition between relatives. *Nature* **409**, 510–513 (2001).

67. T. H. Clutton-Brock, *et al.*, Selfish Sentinels in Cooperative Mammals. *Science (1979)* **284**, 1640–1644 (1999).

68. K. Zuberbühler, D. Jenny, R. Bshary, The Predator Deterrence Function of Primate Alarm Calls. *Ethology* **105**, 477–490 (1999).

69. P. A. Bednekoff, Mutualism Among Safe, Selfish Sentinels: A Dynamic Game. *Am Nat* **150**, 373–392 (1997).

70. S. A. West, A. Gardner, Altruism, Spite, and Greenbeards. *Science (1979)* **327**, 1341–1344 (2010).

71. A. Gardner, S. A. West, Spite and the scale of competition. *J Evol Biol* **17**, 1195–1203 (2004).

72. M. Patel, S. A. West, J. M. Biernaskie, Kin discrimination, negative relatedness, and how to distinguish between selfishness and spite. *Evol Lett* **4**, 65–72 (2020).

73. P. D. Taylor, T. Day, G. Wild, Evolution of cooperation in a finite homogeneous graph. *Nature* **447**, 469–472 (2007).

74. A. Grafen, An inclusive fitness analysis of altruism on a cyclical network. *J Evol Biol* **20**, 2278–2283 (2007).

75. P. Taylor, Inclusive fitness in finite populations-effects of heterogeneity and synergy. *Evolution (N Y)* **71**, 508–525 (2017).

Supplementary Information for 'The evolution of signalling and eavesdropping in plant-fungal networks'

Thomas W. Scott[1,2,3], E. Toby Kiers[4,5] and Stuart A. West[1]

[1]Department of Biology, University of Oxford, 11a Mansfield Road, Oxford, United Kingdom

[2]School of Biology, University of St Andrews, Dyers Brae, St Andrews, United Kingdom

[4]A-life, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands

[5]Society for the Protection of Underground Networks, SPUN, Dover, DE 19901, USA

[3]Author for correspondence: thomas.scott@biology.ox.ac.uk

In this Supplementary Information, we analyse and interpret a series of models. The models presented in Appendices A–H (Models A–H) differ from each other in three respects: (1) some are models of the evolution of signalling predator-presence, and others are models of the evolution of suppressing a cue of predator-presence (Maynard Smith & Harper, 2003; Diggle *et al.*, 2007; Padje *et al.*, 2016); (2) some assume (accurately) that relatedness emerges from model parameters (closed models), and others artificially 'detach' relatedness from lifecycle assumptions so that it can be treated as a parameter (open models) (Frank, 1998; Cooper *et al.*, 2018); (3) some assume a lifecycle where population regulation occurs before migration, and others assume that population regulation occurs after migration (Lehmann & Rousset, 2010). The models are formulated and analysed in the following Appendices:

- **Appendix A.** Signalling; closed; population regulation before migration.
- **Appendix B.** Signalling; open; population regulation before migration.
- **Appendix C.** Signalling; closed; population regulation after migration.
- **Appendix D.** Signalling; open; population regulation after migration.
- **Appendix E.** Cue-suppression; closed; population regulation before migration.
- **Appendix F.** Cue-suppression; open; population regulation before migration.
- **Appendix G.** Cue-suppression; closed; population regulation after migration.
- **Appendix H.** Cue-suppression; open; population regulation after migration.

The reason why we analysed (technically inaccurate) open models as well as (accurate) closed models is that, in closed models, 'who you socially interact with' and 'who you compete with' are both determined by model parameters, meaning they are correlated. This makes the effects of kin selection (favouring helping behaviours) and kin competition (disfavouring helping behaviours) hard to disentangle (Taylor, 1992a, 1992b; Queller, 1994; Frank, 1998; Grafen & Archetti, 2008; Lehmann & Rousset, 2010; Scott & Wild, 2023). Open models detach relatedness from model parameters, so that the causal impact of kin selection can be more easily seen, free from associated competitive effects. Examining open models alongside closed ones has a rich tradition in social evolution research (Frank, 1998; Cooper *et al.*, 2018).

The reason why we analysed 'population regulation before migration' and 'population regulation after migration' versions of the models is that, in the former case, individuals only compete with individuals on their native patch, but in the latter case, individuals compete to some extent with individuals on other patches. This is known to have potentially dramatic consequences on the type of social behaviour that evolves (Lehmann & Rousset, 2010). It is therefore important to examine how sensitive conclusions are to lifecycle assumptions.

We do not *interpret* Models A–H in Appendices A–H themselves. We save this for **Appendix I**, where we re-cap and tie together the results of Models A–H, explaining why these results arise and why different models led to different outcomes. We also set these results in the context of the wider (more general) literature on the evolution of helping and harming behaviours in closed and open models of social evolution.

The models presented in Appendix J & H (Models J & H) are generalisations of Models A–D. Specifically, Models A–D are signalling models where the signal is forced to be honest, and individuals are forced to respond to the signal. Models J & H generalise this scenario so that signal honesty is allowed to evolve, as well as the extent to which signals are responded to (rather than ignored). The generalised models are formulated, analysed and interpreted in the following Appendices:

- **Appendix J**. Signalling; open & closed; population regulation before migration; individuals may signal dishonestly and not respond to signals.
- **Appendix K**. Signalling; open & closed; population regulation after migration; individuals may signal dishonestly and not respond to signals.

Finally, in **Appendix L** we formulate, analyse and interpret a model of mycorrhizal eavesdropping and signalling towards plants. The mycorrhizal signal may evolve to be dishonest, and plants may evolve to ignore the signal. This model differs from previous models in that it is a coevolutionary model (*i.e.*, it tracks evolution in two rather than one species), and that the signaller is the fungus rather than the plant.

A *Mathematica* file comprising all calculations in this Supplementary Information, and a *Matlab* file implementing the numerical procedure used to solve Model K, can be accessed at https://github.com/ThomasWilliamScott/Plant_Signalling. The results figures in the main text plot results for Model K (signalling; Fig. 2), Model E (cue-suppression; Fig. 3a), and Model L (eavesdropping; Fig. 3b).

Appendix A
Model A: signalling; closed; population regulation before migration.

We assume an infinite population of individuals (*e.g.*, plants). We assume for simplicity that individuals are haploid and asexual. The population is split into demes of size N (infinite island model) (Rousset, 2004). We focus our analysis on a focal individual, drawn at random from the population.

Each generation, a random individual on each deme is initially attacked by a predator, and suffers a fecundity cost of d. Given that there are N individuals on each deme, each individual has a $1/N$ chance of being initially attacked by the predator. The initially-predated individual may then invest into the production of a signal. If the focal individual is the initially-predated individual on its deme, it invests an amount x_{focal} into signal production. Signal investment incurs a fecundity cost that scales linearly with c. Therefore, the focal individual pays a signalling cost of cx_{focal} when it is initially-predated.

The signal produced by the initially-predated individual is transferred to (and recognised by) the other N-1 individuals on the deme. We make no assumptions about how the signal is transferred, but it may be being transferred through a common mycorrhizal network, or through the air, *etc*. The signal warns the other plants that a predator is in the vicinity. Consequently, the other plants can prepare for being attacked, and defend themselves, meaning they suffer a reduced fecundity cost of predation. Preparation for predator attack (defence) incurs a fecundity cost that scales linearly with s. We denote the average signalling investment amongst the N-1 other plants on the focal individual's deme by x_{others}. This means that, when the focal individual is not the initially-predated individual, it will ultimately suffer a cost of predation that is given by $d(1-x_{others})$, and a defence cost that is given by sx_{others}. We assume that preparing for predator attack is less costly than the fecundity cost of being predated ($d>s$), which ensures that individuals are favoured to defend themselves against predators.

We emphasise that, in our model, only the initially-predated individual has a chance to warn the other individuals about the presence of a predator. Subsequently-predated individuals may still produce the signal, but our assumption is that by then, the predator has already arrived, rendering the signal useless for warning others about predator presence. We allow for the possibility that subsequently-predated (*i.e.*, not initially-predated) individuals can facultatively turn off signal production, so that they do not pay the costs of producing a useless signal. We assume that the focal individual pays a signalling cost of $cx_{focal}(1-g)$ when subsequently-predated, where g denotes the extent to which signal investment is 'facultative' as opposed to

'obligate' (analogously, a non-focal individual pays a signalling cost of $cx_{others}(1-g)$ when subsequently-predated). A fully obligate signal ($g=0$) means that it is produced irrespective of whether the individual is initially-predated (rendering the signal is useful) or subsequently-predated (rendering the signal useless).

After the fecundity costs of predation and signal investment have been incurred by all individuals, each individual produces a large number of offspring (juvenile haploid clones) in proportion to their fecundity. A random sample of N juvenile individuals are chosen, for each deme, to survive and form the next adult population (local population regulation). After population regulation, a proportion of the (new) adult population, m, migrate to a different deme, randomly chosen for each individual. The remaining proportion, $1-m$, remain on their local deme. This lifecycle then iterates over many generations until an evolutionary end point is reached.

We denote the population average investment into signalling by x_{pop}. We denote the average signalling investment on the focal individual's deme (incorporating its own signalling investment) by x_{deme}; this can be written in terms of x_{focal} and x_{others} as

$$x_{deme} = \frac{(N-1)x_{others}}{N} + \frac{x_{focal}}{N}. \tag{A1}$$

We assume that the baseline fecundity is 1, that $0 \leq x_{focal}, x_{others}, x_{whole}, x_{pop} \leq 1$, and that $c + d + s \leq 1$; these assumptions ensure that fecundity never falls below zero. The number of juvenile offspring produced by the focal individual is then proportional to

$$f_{focal} = 1 - \frac{1}{N}(d + cx_{focal})$$
$$- \left(\frac{N-1}{N}\right)\left(d(1 - x_{others}) + cx_{focal}(1 - g) + sx_{others}\right). \tag{A2}$$

We can make sense of the right-hand side of Equation A2 as follows. Baseline fecundity is one (left-hand term). With probability $1/N$, the focal individual is predated first, leading to additively applied costs of predation (d) and signal production (cx_{focal}), leading to the second term. With probability $\frac{N-1}{N}$, the focal individual is not predated first, and can receive a signal of predator presence from another individual on the deme, leading to additively applied costs of predation ($d(1-x_{others})$) and defence (sx_{others}), plus the costs of signal production that are

incurred even when the focal individual is not predated first, owing to obligatory (unconditional) expression ($cx_{focal}(1\text{-}g)$). This leads to the third term.

The number of juvenile offspring produced by an individual drawn randomly from the focal individual's deme (including the focal individual) is proportional to

$$f_{deme} = 1 - \frac{1}{N}(d + cx_{deme})$$
$$- \left(\frac{N-1}{N}\right)(d(1 - x_{deme}) + cx_{deme}(1-g) + sx_{deme}), \qquad (A3)$$

which we can rewrite using Equation A1 (getting rid of the superfluous x_{deme} parameter) as

$$f_{deme} = 1 - \frac{1}{N}\left(d + c\left(\frac{(N-1)x_{others}}{N} + \frac{x_{focal}}{N}\right)\right)$$
$$- \left(\frac{N-1}{N}\right)\left(d\left(1 - \left(\frac{(N-1)x_{others}}{N} + \frac{x_{focal}}{N}\right)\right)\right.$$
$$+ c\left(\frac{(N-1)x_{others}}{N} + \frac{x_{focal}}{N}\right)(1-g)$$
$$\left. + s\left(\frac{(N-1)x_{others}}{N} + \frac{x_{focal}}{N}\right)\right). \qquad (A4)$$

The number of juvenile offspring produced by an individual drawn randomly from the population is proportional to

$$f_{pop} = 1 - \frac{1}{N}(d + cx_{pop}) - \left(\frac{N-1}{N}\right)(d(1 - x_{pop}) + cx_{pop}(1-g) + sx_{pop}). \qquad (A5)$$

The fitness of the focal individual is then given by:

$$w = \frac{f_{focal}}{f_{deme}}, \qquad (A6)$$

where explicit expressions for the fecundity (f_{focal}, f_{deme}) functions are provided in Equations A2 & A4 (Rousset, 2004; Roze & Rousset, 2008; Lehmann & Rousset, 2010). Note that the

f_pop function derived above (Equation A5) does not feature in this fitness function, but it will feature in the fitness function of some subsequent models. 'Fitness' refers to the number of surviving offspring produced by an individual after one full iteration of the lifecycle (Rousset, 2004). Under our lifecycle assumptions, the size of the population remains constant after each full iteration of the lifecycle, and this is reflected by the fact that the average fitness taken over the population is equal to 1.

As a technical aside, we note that, by 'fitness', we strictly mean 'neighbour-modulated' fitness, which counts up fitness effects on a focal individual. This can be conceptually distinguished from 'inclusive' fitness, which counts up fitness effects arising from the actions of a focal individual, though the two fitness measures are formally equivalent under a standard set of mathematical assumptions (Rousset, 2004; Taylor *et al.*, 2007; Scott & Wild, 2023).

Equation A6 reflects the fact that, under our assumption that (local) population regulation occurs before migration, individuals only compete with competitors on their native deme, rather than with individuals on other demes. This is why the fecundity of the focal individual (f_{focal}) is weighted only by the average fecundity of an individual from the native deme (f_{deme}), rather than by the average fecundity of an individual drawn from the population at large (f_{pop}).

Our aim is to examine what level of signal investment evolves at evolutionary equilibrium. To do this, we assume a monomorphic population, where all individuals invest the same amount into signal production, given by x^* (the lack of subscript implies that the value is the same across the population). We then take a random (focal) individual and mutate its (and its identical-by-descent relatives') level of signal investment to a deviant value x, and ask how the fitness of the focal individual changes in response, in effect calculating $\frac{dw}{dx}$. Evaluating $\frac{dw}{dx}$ at equilibrium (x^*), the equilibrium (ESS) level of signal production can then be obtained as: (*i*) $x^*=1$ if $\frac{dw}{dx}|_{x=x^*} > 0$ (maximal signal production); (*ii*) $x^*=0$ if $\frac{dw}{dx}|_{x=x^*} < 0$ (no signal production); (*iii*) $0 \leq x^* \leq 1$ if $\frac{dw}{dx}|_{x=x^*} = 0$, where x^* is specifically obtained as the value for which $\frac{dw}{dx}|_{x=x^*} = 0$ (Taylor & Frank, 1996; Frank, 1998).

To calculate $\frac{dw}{dx}$, we first expand it using the chain rule

$$\frac{dw}{dx} = \frac{\partial w}{\partial x_{focal}} \frac{dx_{focal}}{dx} + \frac{\partial w}{\partial x_{others}} \frac{dx_{others}}{dx}, \qquad (A7)$$

and then proceed to evaluate the derivatives on the right-hand side (Taylor & Frank, 1996; Frank, 1998). Because the population is monomorphic, and because we are interested in the long term evolutionary state of the population, we evaluate these derivatives at the point where $x_{focal}, x_{others}, x_{pop} = x^*$. We trivially obtain

$$\frac{dx_{focal}}{dx}\big|_{x_{focal}, x_{others}, x_{pop}=x^*} = 1. \tag{A8}$$

We obtain the equilibrium-evaluated partial derivatives $\frac{\partial w}{\partial x_{focal}}\big|_{x_{focal}, x_{others}, x_{pop}=x^*}$ and $\frac{\partial w}{\partial x_{others}}\big|_{x_{focal}, x_{others}, x_{pop}=x^*}$ respectively as

$$\frac{\partial w}{\partial x_{focal}}\big|_{x_{focal}, x_{others}, x_{pop}=x^*}$$
$$= -\frac{(N-1)(c(g(-N)+g+N)+d-s)}{N(x^*(N(cg-s)-c(g+N)+d(N-1)+s)-dN+N)} \tag{A9}$$

and

$$\frac{\partial w}{\partial x_{others}}\big|_{x_{focal}, x_{others}, x_{pop}=x^*}$$
$$= \frac{(N-1)(c(g(-N)+g+N)+d-s)}{N(x^*(N(cg-s)-c(g+N)+d(N-1)+s)-dN+N)}. \tag{A10}$$

We obtain $\frac{dx_{others}}{dx}\big|_{x_{focal}, x_{others}, x_{pop}=x^*}$ by first noting that $\frac{dx_{others}}{dx}$ is interpretable as a coefficient of relatedness, since it denotes the marginal (correlated) change in signal production by a social partner as a consequence of a change in signal production by an actor (Taylor & Frank, 1996; Frank, 1998). We can therefore write

$$\frac{dx_{others}}{dx} = r, \tag{A11}$$

where r denotes the relatedness between social partners. We note that r is an 'others-only' measure of relatedness, as opposed to a 'whole-group' measure of relatedness, since it measures the relatedness between an individual and its social partners (who are the other members of the

deme), rather than the relatedness between an individual and a random member of its deme (including itself) (Pepper, 2000).

We cannot evaluate r in general, because it is changing over time and will depend on initial genotype frequencies, but fortunately, we can evaluate it at equilibrium, which is the point that we are interested in anyway, to find

$$\frac{dx_{others}}{dx}\Big|_{x_{focal},x_{others},x_{pop}=x^*} = r^*. \tag{A12}$$

To get an expression for the equilibrium relatedness, r^*, it is useful to first calculate the equilibrium 'whole-group' relatedness for this model, r^*_{whole}, then convert back to find the equilibrium 'others-only' relatedness, r^*. The equilibrium whole-group relatedness can be found by first writing a recursion for how whole group relatedness, r_{whole}, changes across a generation (r'_{whole} denotes whole-group relatedness in the next generation):

$$r'_{whole} = \frac{1}{N} + r_{whole}\left(\frac{N-1}{N}\right)(1-m)^2 \tag{A13}$$

(Taylor, 1992a; Rousset, 2004; Lehmann & Rousset, 2010). We obtained Equation A13 by calculating the average relatedness between a focal individual and a randomly drawn member of the focal individual's deme after one iteration of the lifecycle. With probability $1/N$, the randomly drawn individual is the focal individual itself, leading to a relatedness of 1, hence the first term of the right-hand side of Equation A13. With probability $\frac{N-1}{N}$, the randomly drawn individual is not the focal individual. With probability 1-m, the randomly drawn individual is a non-dispersing individual. With probability 1-m, the focal individual is a non-dispersing individual. This leads to a relatedness of r_{whole}, and combining these factors leads to the second term of the right-hand side of Equation A13. With the remining probability $\frac{N-1}{N}(1-(1-m)^2)$, the randomly drawn individual is related to the focal individual by zero, because one of the two individuals have dispersed – the multiplication by zero is why this term does not feature in Equation A13.

Equilibrium whole-group relatedness is then obtained by setting $r'_{whole} = r_{whole} = r^*_{whole}$ to obtain

$$r^*_{whole} = \frac{1}{N - (N-1)(1-m)^2} \tag{A14}$$

(Taylor, 1992a). Then, we can write down how whole-group relatedness relates to others-only relatedness:

$$r_{whole} = \frac{1}{N} + \left(\frac{N-1}{N}\right)r \tag{A15}$$

(Pepper, 2000). Using Equations A14 & A15, and setting $r_{whole} = r^*_{whole}$ & $r = r^*$, we obtain the equilibrium (others-only) relatedness as

$$r^* = \frac{1}{1 - \frac{(m-2)mN}{(m-1)^2}}, \tag{A16}$$

which we can substitute into Equation A12 to find the equilibrium (long-term) value of $\frac{dx_{others}}{dx_{focal}}$:

$$\frac{dx_{others}}{dx}\Big|_{x_{focal},x_{others},x_{pop}=x^*} = \frac{1}{1 - \frac{(m-2)mN}{(m-1)^2}}. \tag{A17}$$

We can now substitute Equations A8, A9, A10 & A17 into Equation A7 to obtain an explicit expression for $\frac{dw}{dx}\Big|_{x_{focal},x_{others},x_{pop}=x^*}$:

$$\frac{dw}{dx}\Big|_{x_{focal},x_{others},x_{pop}=x^*}$$
$$= \Big((m-2)m(N-1)(c(g-1)N - cg - d + s)\Big)$$
$$/ \Big(((m-2)m(N-1) - 1)(x^*(N(cg - s) - c(g + N) + d(N-1) + s)$$
$$- dN + N)\Big). \tag{A18}$$

As long as there is some migration ($m>0$), the right-hand side of Equation A18 is strictly negative. If the reader would like to check this for themselves by inspection, remember that

most variables are bound between 0 and 1, except for N which is greater than or equal to 2, and also remember that $c + d + s \leq 1$ and $d > s$.

Therefore, in this model, the long-term (ESS) level of signal production is $x^* = 0$. At equilibrium, individuals evolve to *not* signal the presence of predators to their social partners. Any tendency to signal the presence of predators will ultimately be removed by natural selection.

Appendix B

Model B: signalling; open; population regulation before migration.

We now 'open' Model 1, detaching relatedness from demographic parameters (*m* and *N*), so that it is now a parameter (constant) in its own right (Frank, 1998; Cooper *et al.*, 2018). Inspecting Equation A16 reveals that this assumption, of independence of relatedness from demography, is obviously false. But we proceed anyway with this noted.

To analyse this open model, where relatedness varies independently of model parameters, we calculate we calculate $\frac{dw}{dx}|_{x=x^*}$ as before, by evaluating the right-hand derivatives of Equation A7 at the point where $x_{focal}, x_{others}, x_{pop} = x^*$. The only difference is that, in the previous model, we evaluated the $\frac{dx_{others}}{dx}|_{x_{focal}, x_{others}, x_{pop}=x^*}$ derivative, which is interpretable as an equilibrium value of relatedness, in terms of model parameters *m* and *N*, to obtain Equation A17. However, now, we simply evaluate $\frac{dx_{others}}{dx}|_{x_{focal}, x_{others}, x_{pop}=x^*}$ as a constant, *r**. In other words, in the previous model, we substituted Equation A17 (alongside Equations A8, A9 & A10) into Equation A7 to obtain an explicit expression for $\frac{dw}{dx}|_{x_{focal}, x_{others}, x_{pop}=x^*}$, but now, we substitute in Equation A12 instead (still alongside Equations A8, A9 & A10). We obtain

$$\frac{dw}{dx}|_{x_{focal}, x_{others}, x_{pop}=x^*}$$
$$= \frac{(N-1)(r^*-1)(c(g(-N)+g+N)+d-s)}{N(x^*(N(cg-s)-c(g+N)+d(N-1)+s)-dN+N)}. \qquad (B1)$$

As long as relatedness is not maximal (*r**<1), the right-hand side of Equation B1 is strictly negative. Therefore, in this model, the long-term (ESS) level of signal production is $x^* = 0$. This is the same result as the equivalent 'closed' model (Model A). We have shown that 'opening' Model A does not facilitate the evolution of signalling. In the next section, we therefore move onto a slightly different lifecycle, where population regulation occurs after migration, to see if it is more favourable for the evolution of signalling predator-presence.

Appendix C

Model C: signalling; closed; population regulation after migration.

In this Appendix, we formulate and analyse a slightly different version of Model A, where population regulation now occurs after migration, rather than before. Specifically, after each individual has produced a large number of offspring (juvenile haploid clones) in proportion to their fecundity, a proportion of these offspring, *m,* migrate to a different deme, randomly chosen for each juvenile individual. A proportion, *1-m*, remain on the local deme. A random sample of *N* juvenile individuals are then chosen, for each deme, to survive and form the adult population for the next generation (local population regulation).

Under these revised lifecycle assumptions, the fitness of the focal individual is given by

$$w = (1 - m)\frac{f_{focal}}{(1 - m)f_{deme} + mf_{pop}} + m\frac{f_{focal}}{f_{pop}}, \qquad (C1)$$

where explicit expressions for the fecundity (*f$_i$*) functions are provided in Equations A2, A4 & A5 (Rousset, 2004; Roze & Rousset, 2008; Lehmann & Rousset, 2010).

We can make sense of the first term of the right-hand side of Equation C1 as follows. Of the f_{focal} juvenile offspring produced by the focal individual, $(1 - m)f_{focal}$ of these remain on the native deme, leading to the numerator. These juvenile offspring compete with the $(1 - m)f_{deme}$ non-dispersing juvenile offspring produced by an average member of the native deme, as well as the mf_{pop} dispersing juvenile offspring produced by an average member of another deme, leading to the denominator. The second term of the right-hand side of Equation C1 can be understood analogously.

We then analyse the model, as before, by evaluating the right-hand derivatives of the $\frac{dw}{dx}|_{x=x^*}$ expression (Equation A7) at equilibrium. For this version of the model, these right-hand derivates are calculated as:

$$\frac{\partial w}{\partial x_{focal}}\Big|_{x_{focal},x_{others},x_{pop}=x^*}$$

$$= -\frac{c\big((g - 1)N - g\big)((m - 1)^2 - N) + d(m - 1)^2(N - 1) - (m - 1)^2(N - 1)s}{N(x^*(N(cg - s) - c(g + N) + d(N - 1) + s) - dN + N)}, \qquad (C2)$$

$$\frac{\partial w}{\partial x_{others}}\Big|_{x_{focal},x_{others},x_{pop}=x^*}$$

$$= -\Big((N-1)\big(c(m-1)^2\big((g-1)N-g\big)+d(m-2)mN-d(m-1)^2$$

$$-(m-2)m(N-1)s+s\big)\Big)$$

$$/\Big(N(x^*(N(cg-s)-c(g+N)+d(N-1)+s)-dN$$

$$+N)\Big). \tag{C3}$$

The $\frac{dx_{focal}}{dx}\Big|_{x_{focal},x_{others},x_{pop}=x^*}$ and $\frac{dx_{others}}{dx}\Big|_{x_{focal},x_{others},x_{pop}=x^*}$ derivatives are still given by Equations A8 and A17 respectively.

By substituting Equations A8, C2, C3 & A17 into Equation A7, we obtain

$$\frac{dw}{dx}\Big|_{x_{focal},x_{others},x_{pop}=x^*}$$

$$= (c(m-2)mN(g(N-1)-N))$$

$$/\Big(((m-2)m(N-1)-1)(x^*(N(cg-s)-c(g+N)+d(N-1)+s)$$

$$-dN+N)\Big). \tag{C4}$$

As long as there is some migration ($m>0$) and signalling cost ($c>0$), the right-hand side of Equation C4 is strictly negative. This means that the long-term (ESS) level of signal production is $x^* = 0$. This is the same result as Models A & B. Therefore, the lifecycle change (moving population regulation to go after migration) did not facilitate the evolution of signalling. Next, we 'open' this model, to see if the lifecycle change is more favourable to the evolution of signalling in an open-model setting.

Appendix D

Model D: signalling; open; population regulation after migration.

In this Appendix, we formulate and analyse the 'open' version of Model C, where relatedness is now treated (artificially) as a constant (Frank, 1998; Cooper *et al.*, 2018).

As usual, we calculate $\frac{dw}{dx}|_{x=x^*}$ by evaluating its right-hand derivatives (Equation A7). Substituting Equations A8, C2, C3 & A12 into Equation A7, we obtain

$$\frac{dw}{dx}|_{x_{focal},x_{others},x_{pop}=x^*}$$

$$= -((N-1)r^*(c(m-1)^2((g-1)N-g)+d(m-2)mN-d(m-1)^2$$
$$- (m-2)m(N-1)s+s)+c((g-1)N-g)((m-1)^2-N)$$
$$+ d(m-1)^2(N-1)-(m-1)^2(N-1)s)$$
$$/(N(x^*(N(cg-s)-c(g+N)+d(N-1)+s)-dN$$
$$+ N)). \tag{D1}$$

This expression is complex, so we focus on two special cases. First, we consider the case where the migration rate is zero. Substituting *m*=0 into Equation D1, we obtain

$$\frac{dw}{dx}|_{x_{focal},x_{others},x_{pop}=x^*,m=0}$$

$$= \frac{(N-1)(r^*-1)(c(g(-N)+g+N)+d-s)}{N(x^*(N(cg-s)-c(g+N)+d(N-1)+s)-dN+N)}. \tag{D2}$$

As long as relatedness is not maximal ($r^* < 1$), the right-hand side of Equation D2 is strictly negative. This means that, when there is no migration, the long-term (ESS) level of signal production is $x^* = 0$ (no signal production).

Second, we consider the case where the migration rate is one. Substituting *m*=1 into Equation D1, we obtain

$$\frac{dw}{dx}|_{x_{focal},x_{others},x_{pop}=x^*,m=1}$$

$$= \frac{c(g-1)N-cg+(N-1)r^*(d-s)}{x^*(N(cg-s)-c(g+N)+d(N-1)+s)-dN+N}. \tag{D3}$$

We find that this condition is positive, meaning signalling is favoured (leading to $x^*=1$), whenever

$$r^* > \frac{c\left(1 - \frac{N-1}{N}g\right)}{(d-s)\left(\frac{N-1}{N}\right)} \qquad (D4)$$

is satisfied. Conversely, signalling is disfavoured, leading to $x^*=0$, whenever $r^* < \frac{c\left(1-\frac{N-1}{N}g\right)}{(d-s)\left(\frac{N-1}{N}\right)}$. Therefore, signal production is sometimes favoured if there is sufficient migration (m).

Equation D4 is a form of Hamilton's rule, as can be seen by setting $C = c\left(1 - \frac{N-1}{N}g\right)$ and $B = (d-s)\left(\frac{N-1}{N}\right)$ to obtain $r^* > \frac{C}{B}$. The capital 'C' gives the overall cost of the social act (signalling). It is the unit cost of signal production (c), weighted by the proportion of incidences where this cost is actually paid $\left(1 - \frac{N-1}{N}g\right)$. The capital 'B' gives the overall benefit of the social act (signalling). It is the unit benefit of signal production, as manifested as a reduced fecundity cost of being predated (d) minus the cost of predator defence (s), weighted by the proportion of incidences where this benefit is actually received $\left(\frac{N-1}{N}\right)$.

Therefore, unlike in Models A-C, signalling predator-presence can be favoured in this model (if there is sufficient migration and Hamilton's Rule is satisfied). The combination of 'population regulation after migration' and 'relatedness as a parameter' (open model) can facilitate the evolution of signalling predator-presence.

To give even more intuition for this result, we plot it below in Fig. S1. Signalling is favoured above the lines (technically, this corresponds to when $\frac{dw}{dx}|_{x_{focal},x_{others},x_{pop}=x^*}$, given in Equation D1, is greater than zero).

Figure S1. The evolution of signalling in an open model (Model D). Signalling is favoured above the lines, and disfavoured below them. Technically, the lines correspond to when $\frac{dw}{dx}\big|_{x_{focal},x_{others},x_{pop}=x^*}$ (given in Equation D1) is equal to zero. The solid black line is a reference line generated with arbitrary parameter values. Coloured lines show the effect of increasing parameter values. Signalling is more likely to be favoured when: relatedness (r^*), scale of competition (m), group size (N), predation cost (d) and conditionality (g) are high; trait cost (c) and defence cost (s) are low. We assumed: (*reference values*) d=0.1, c=0.001, N=3, s=0.08, g=0; (*alternative values*) d=0.9, c=0.003, N=10, s=0.09, g=1.

Appendix E

Model E: Cue-suppression; closed; population regulation before migration.

We now slightly alter assumptions of Model A, to study a different sort of social behaviour. As before, we assume an infinite population of haploid and asexual individuals, split into demes of size N, and focus on a randomly drawn focal individual. As before, each generation, a random individual on each deme is initially attacked by a predator and suffers a fecundity cost d.

However, in the present model, after the initially-predated individual is attacked by the predator, it releases a *cue* rather than a *signal*. The cue indicates to the other N-1 individuals on the deme that a predator is in the vicinity. However, we assume that the initially-predated individual may suppress the cue, to stop the other N-1 individuals from learning that a predator is present. If the focal individual is the initially-predated individual on its deme, it invests an amount x_{focal} into cue suppression. Investment into cue-suppression incurs a fecundity cost that scales linearly with c. Therefore, the focal individual pays a cue-suppression cost of cx_{focal} when it is initially-predated.

We denote the average cue-suppression investment amongst the N-1 other plants on the focal individual's deme by x_{others}. This means that, when the focal individual is not the initially-predated individual, it suffers a cost of predation of dx_{others}, and a defence cost that is given by $s(1-x_{others})$. We denote the population average investment into cue-suppression by x_{pop}, and the average cue-suppression investment on the focal individual's deme by x_{deme}.

As before, we assume that only the initially-predated individual can convey useful information about predator presence. Specifically, only cues transmitted by the initially-predated individual can warn the other individuals about predator-presence; cues transmitted by subsequently-predated individuals are useless, because the predator has already arrived. We allow for the facultative 'turning off' of cue-suppression when the individual is subsequently-predated. Therefore, the focal individual pays a cue-suppression cost of $cx_{focal}(1-g)$ when subsequently-predated (non-focal individuals pay a signalling cost of $cx_{others}(1-g)$ when subsequently-predated), where g denotes how facultative cue-suppression is.

After social interactions and predation have taken place, the rest of the lifecycle plays out as it does in the Model A: production of a large number of juvenile offspring in proportion to fecundity; random sampling of N juveniles on each deme to establish the next adult

population; migration of (surviving) adults to a random patch with probability *m*. The lifecycle is iterated until an evolutionary end-point is reached.

We emphasise that the only mathematical difference between this model and Model A is that, in Model A, secondarily-predated individuals suffer predation and defence costs of $d(1-x_i)$ & sx_i respectively, but in the present model, secondarily-predated individuals suffer predation and defence costs of dx_i & $s(1-x_i)$ respectively (where *i=others* for the focal individual; *i=deme* for non-focal individuals on the focal individual's deme; *i=pop* for individuals on other demes). The other mathematical details are the same. This means the fecundity functions (Equations A2–A5) are changed to:

$$f_{focal} = 1 - \frac{1}{N}\left(d + cx_{focal}\right)$$
$$- \left(\frac{N-1}{N}\right)\left(dx_{others} + cx_{focal}(1-g) + s(1-x_{others})\right), \qquad (E1)$$

$$f_{deme} = 1 - \frac{1}{N}\left(d + c\left(\frac{(N-1)x_{others}}{N} + \frac{x_{focal}}{N}\right)\right)$$
$$- \left(\frac{N-1}{N}\right)\left(d\left(\frac{(N-1)x_{others}}{N} + \frac{x_{focal}}{N}\right)\right.$$
$$+ c\left(\frac{(N-1)x_{others}}{N} + \frac{x_{focal}}{N}\right)(1-g)$$
$$\left.+ s\left(1 - \left(\frac{(N-1)x_{others}}{N} + \frac{x_{focal}}{N}\right)\right)\right), \qquad (E2)$$

$$f_{pop} = 1 - \frac{1}{N}\left(d + cx_{pop}\right) - \left(\frac{N-1}{N}\right)\left(dx_{pop} + cx_{pop}(1-g) + s(1-x_{pop})\right). \qquad (E3)$$

The fitness of the focal individual is obtained by substituting the fecundity functions (Equations E1 – E3) into Equation A6. With fitness obtained, we proceed, as usual, by evaluating the derivatives on the right-hand side of Equation A7, which we now obtain as

$$\frac{\partial w}{\partial x_{focal}}\Big|_{x_{focal},x_{others},x_{pop}=x^*}$$

$$= -\frac{(N-1)(c(g-1)N - cg + d - s)}{N(x^*(-N(cg+s) + c(g+N) + d(N-1) + s) + d + (N-1)s - N)}, \qquad (E4)$$

$$\frac{\partial w}{\partial x_{others}}\Big|_{x_{focal},x_{others},x_{pop}=x^*}$$

$$= \frac{(N-1)(c(g-1)N - cg + d - s)}{N(x^*(-N(cg+s) + c(g+N) + d(N-1) + s) + d + (N-1)s - N)}. \qquad (E5)$$

We obtain the same expressions for $\frac{dx_{focal}}{dx}\Big|_{x_{focal},x_{others},x_{pop}=x^*}$ and

$\frac{dx_{others}}{dx}\Big|_{x_{focal},x_{others},x_{pop}=x^*}$ as we did for Model A (provided in Equations A8 and A17 respectively). By substituting Equations A8, A17, E4 & E5 into Equation A7, we obtain

$$\frac{dw}{dx}\Big|_{x_{focal},x_{others},x_{pop}=x^*}$$

$$= -\big((m-2)m(N-1)(c(g-1)N - cg + d - s)\big)$$

$$/ \Big(\big((m-2)m(N-1) - 1\big)(x^*(-N(cg+s) + c(g+N) + d(N-1) + s)$$

$$+ d + (N-1)s - N)\Big). \qquad (E6)$$

We can see by setting this to be greater than zero that cue-suppression evolves when

$$\frac{d-s}{N} > c\left(1 - \frac{N-1}{N}g\right). \qquad (E7)$$

Equation E7 can be understood as cue-suppression (harming) being favoured when the cost of the social act to the actor, $c\left(1 - \frac{N-1}{N}g\right)$, is less than the cost of the social act to the competitor, $\frac{d-s}{N}$. The actor cost term $c\left(1 - \frac{N-1}{N}g\right)$ is the unit cost of cue-suppression (c), weighted by the proportion of incidences where this cost is actually paid $\left(1 - \frac{N-1}{N}g\right)$. The competitor cost term $\frac{d-s}{N}$ is the fecundity cost of being predated (d), minus the saved cost of predator defence (s), weighted by the proportion of incidences where these costs can actually be applied by the actor ($1/N$).

Therefore, for this lifecycle (population regulation before migration), suppressing a cue of predator-presence can evolve, but signalling predator-presence cannot evolve (Model A).

Appendix F

Model F: Cue-suppression; open; population regulation before migration.

In this Appendix, we 'open' Model E, by substituting Equation A12 rather than Equation A17 into Equation A7 to obtain

$$\frac{dw}{dx}\Big|_{x_{focal}, x_{others}, x_{pop} = x^*}$$
$$= \big((N-1)(r^* - 1)(c(g-1)N - cg + d - s)\big)$$
$$/\big(N(x^*(-N(cg+s) + c(g+N) + d(N-1) + s) + d + (N-1)s$$
$$- N)\big). \tag{F1}$$

We can see by setting this to be greater than zero that cue-suppression evolves when Equation E7 is satisfied $\left(\frac{d-s}{N} > c\left(1 - \frac{N-1}{N}g\right)\right)$. The conditions are the same for the open and closed versions of the model, which shows that allowing relatedness to vary independently of kin competition does not affect evolutionary outcomes (relatedness does not influence the outcome of social evolution in this lifecycle).

Appendix G

Model G: Cue-suppression; closed; population regulation after migration.

We now amend Model E so that population occurs after migration, rather than before. This means that fitness is obtained by substituting the fecundity functions Equations E1 – E3 into Equation C1 rather than A6, leading to the following derivatives:

$$\frac{\partial w}{\partial x_{focal}}\Big|_{x_{focal},x_{others},x_{pop}=x^*}$$

$$= -\frac{-c((g-1)N-g)((m-1)^2-N)+d(m-1)^2(N-1)-(m-1)^2(N-1)s}{N(x^*(-N(cg+s)+c(g+N)+d(N-1)+s)+d+(N-1)s-N)}, \quad (G1)$$

$$\frac{\partial w}{\partial x_{others}}\Big|_{x_{focal},x_{others},x_{pop}=x^*}$$

$$= -\Big((N-1)\big(-c(m-1)^2((g-1)N-g)+d(m-2)mN-d(m-1)^2$$

$$- (m-2)m(N-1)s+s\big)\Big)$$

$$/\Big(N(x^*(-N(cg+s)+c(g+N)+d(N-1)+s)+d+(N-1)s$$

$$- N)\Big). \quad (G2)$$

We obtain the same expressions for $\frac{dx_{focal}}{dx}\Big|_{x_{focal},x_{others},x_{pop}=x^*}$ and $\frac{dx_{others}}{dx}\Big|_{x_{focal},x_{others},x_{pop}=x^*}$ as we did for Model A (provided in Equations A8 and A17 respectively).

By substituting Equations A8, A17, G1 & G2 into Equation C1, we obtain

$$\frac{dw}{dx}\Big|_{x_{focal},x_{others},x_{pop}=x^*}$$

$$= -\big(c(m-2)mN(g(N-1)-N)\big)$$

$$/\Big(((m-2)m(N-1)-1)(x^*(-N(cg+s)+c(g+N)+d(N-1)+s)$$

$$+ d+(N-1)s-N)\Big). \quad (G3)$$

As long as there is some migration ($m>0$) and cue-suppression cost ($c>0$), the right-hand side of Equation G3 is strictly negative. This means that the long-term (ESS) level of cue-

suppression is $x^* = 0$. Therefore, at equilibrium, individuals evolve to *not* suppress the cue of predator-presence. Any tendency to suppress the cue of predator-presence will ultimately be removed by natural selection. Therefore, cue-suppression could evolve previously (Models E & F), but the change in lifecycle assumption, so that population regulation occurs after migration rather than before, prevents cue-suppression from evolving.

Appendix H

Model H: Cue-suppression; open; population regulation after migration.

We now 'open' Model G, by substituting Equation A12 rather than Equation A17 into Equation C1, to obtain

$$\frac{dw}{dx}\Big|_{x_{focal},x_{others},x_{pop}=x^*}$$
$$= (-(N-1)r^*(-c(m-1)^2((g-1)N-g)+d(m-2)mN-d(m-1)^2$$
$$-(m-2)m(N-1)s+s)+c((g-1)N-g)((m-1)^2-N)$$
$$-d(m-1)^2(N-1)+(m-1)^2(N-1)s)$$
$$/(N(x^*(-N(cg+s)+c(g+N)+d(N-1)+s)+d+(N-1)s$$
$$-N)). \tag{H1}$$

This expression is complex, so we focus on two special cases. First, we consider the case where the migration rate is one. Substituting $m=1$ into Equation 29, we obtain

$$\frac{dw}{dx}\Big|_{x_{focal},x_{others},x_{pop}=x^*,m=1}$$
$$= \big(c(g-1)N-cg-(N-1)r^*(d-s)\big)$$
$$/(-x^*(-N(cg+s)+c(g+N)+d(N-1)+s)-d-Ns+N+s). \tag{H2}$$

As long as there is some relatedness ($r^* > 0$) or cost of cue-suppression ($c>0$), the right-hand side of Equation H2 is strictly negative. This means that, when there is maximal migration, the long-term (ESS) level of signal production is $x^* = 0$ (no cue suppression).

Second, we consider the case where the migration rate is zero. Substituting $m=0$ into Equation H1, we obtain Equation F1, which is positive (meaning cue-suppression is favoured, leading to $x^*=1$) whenever Equation E7 is satisfied $\left(\frac{d-s}{N} > c\left(1 - \frac{N-1}{N}g\right)\right)$. Conversely, cue-suppression is disfavoured, leading to $x^*=0$, whenever $\frac{d-s}{N} < c\left(1 - \frac{N-1}{N}g\right)$. Therefore, cue-suppression is sometimes favoured if there is sufficiently low migration (m).

Therefore, whilst cue-suppression could evolve when population regulation preceded migration (Models E & F), but not in a closed model where migration preceded population

regulation (Model G), opening this latter model (G) allows the results of the previous models (E & F) to be recovered for the zero-migration case.

Appendix I

Supplementary Discussion

<u>Summary of results</u>

To recap, we found that signalling the presence of a predator cannot evolve unless: migration precedes population regulation, relatedness is detached from model parameters (open model) and sufficiently high that Hamilton's Rule is satisfied (Equation D4), migration is sufficiently high. Only in Model D are these conditions satisfied.

We found that suppressing a cue of predator-presence is favoured when cue-suppression causes more harm to others than yourself (Equation E7), and either: population regulation precedes migration (Models E & F); or, migration precedes population regulation, relatedness is detached from model parameters and sufficiently low, migration is sufficiently low (Model H).

To explain these results, we first note that: signalling the presence of a predator is an example of a *helping* behaviour because it increases the fitness (offspring after one lifecycle iteration) of neighbours; suppressing a cue of predator-presence is an example of a *harming* behaviour because it decreases the fitness of neighbours (Rousset, 2004; Lehmann & Keller, 2006; Lehmann & Rousset, 2010). Most of our results are recovered as special cases of previously published results on the evolution of helping and harming behaviours generally (though this doesn't render our theoretical analysis redundant or trivial – our models had novel ecological aspects like predation, which could in principle have caused our results to diverge from previous models).

<u>Models C & G</u>

As first shown formally by Taylor (1992a), in closed island lifecycles where population regulation goes after migration, social behaviours (i.e., helping and harming behaviours) cannot evolve. (Technically, behaviours with social characteristics can evolve, but only if they provide a 'direct' fitness benefit to the actor – in other words, the 'indirect' or 'social' characteristic of the behaviour does not have any bearing on whether it is favoured; Lehmann and Rousset, 2010.) This is why neither signalling predator-presence nor suppressing a cue of predator-presence could evolve in our Models C & G. The reason for this is that, in closed models of this type, social behaviours either serve to help competitors, or harm non-competitors, which precludes the evolution of social behaviours.

To tease this out further, consider a helping behaviour. For helping to evolve (by kin selection), help needs to be directed towards individuals that are: (*i*) sufficiently related to the actor, and (*ii*) in sufficiently weak competition with the actor (Taylor, 1992b, 1992a; Queller, 1994; Grafen & Archetti, 2008; Lehmann & Rousset, 2010; Scott & Wild, 2023). But the closed lifecycle means that, whenever help is directed towards relatives (this occurs when the migration rate is low), the individuals receiving help are also in strong competition with the helper (because the low migration rate leads to intense local competition). There are no combinations of demographic parameters (*N,m*) that simultaneously lead to high relatedness and low competitiveness amongst social partners, which precludes the evolution of helping.

Now consider a harming behaviour. For harming to evolve, harm needs to be directed towards individuals that: (*i*) have sufficiently low relatedness with the actor; (*ii*) are in sufficiently strong competition with the actor (Queller, 1994; Lehmann *et al.*, 2009; Lehmann & Rousset, 2010; West & Gardner, 2010). But the closed lifecycle means that, whenever harm is directed towards non-kin (this occurs when the migration rate is high), the individuals that are harmed are not in strong competition with the harmer (because the high migration rate causes them to compete appreciably on other demes). There are no combinations of demographic parameters (*N,m*) that simultaneously lead to high competitiveness and low relatedness amongst social partners, which precludes the evolution of harming.

Models D & H

Taylor's (1992a) scenario can be 'opened' so that relatedness can vary independently of demographic parameters (*N,m*). Consequently, relatedness can be made to increase, without also increasing the relatedness of competitors, allowing helping behaviours to evolve. Specifically, this requires that, as well as a high value for the relatedness parameter, the migration rate is kept high, leading to low competition between the actor and its social partners. Frank (1998, p114-115) showed that increasing a relatedness parameter, as well as a migration parameter (or in his terminology, a 'spatial scale of density dependent competition' parameter), can allow helping to evolve. This explains why signalling predator-presence (an example of a helping behaviour) could evolve in our Model D.

Analogously, 'opening' the Taylor (1992a) scenario can allow harming behaviours to evolve, by simultaneously decreasing a relatedness parameter, as well as a migration parameter (increasing competitiveness with social partners). This explains why cue-suppression (an example of a harming behaviour) could evolve in our Model H.

Models A & E

Taylor's closed (1992a) scenario can be amended slightly, so that population regulation precedes migration. Lehmann and Rousset (2010) showed that helping behaviours cannot evolve in this setting, but harming behaviours can. This is why signalling predator-presence (a helping behaviour) could not evolve in our Model A, but cue-suppression (a harming behaviour) could evolve in our Model E.

The reason why helping cannot evolve in this lifecycle is that, when population regulation precedes migration, competition occurs completely within demes (rather than between individuals from different demes), leading to maximally local competition. The evolution of helping requires that competitiveness with social partners is low, but this is impossible in this lifecycle (i.e., it cannot be achieved by any combination of demographic parameters, N & m), precluding the evolution of helping.

The reason why harming can evolve in this lifecycle is that, for harming to evolve, there needs to be sufficiently high competition and low relatedness with social partners. In this lifecycle, competition with social partners is maximal (maximally local competition). It turns out that, given the extreme strength of local competition, harming can evolve for any value of relatedness (*i.e.*, it doesn't need to be especially low, because local competition is so intense) (Lehmann & Rousset, 2010).

Models B & F

Lehmann and Rousset's (2010) closed scenario, where population regulation precedes migration, can be 'opened' so that relatedness can vary independently of demographic parameters (*N,m*). However, opening this scenario (*e.g.*, Models B & F) does not alter any of the equivalent closed model results (*e.g.*, Models A & E). Specifically, helping can still never evolve, and harming can (for any relatedness). The reason for this is that, in the open model setting, relatedness to social partners can be varied, but this is inconsequential, because it is the intense (maximal) local competition that is driving results for this lifecycle. Specifically, it precludes helping behaviours, and facilitates harming behaviours, irrespective of relatedness to social partners.

Take-home message

Closed models are more realistic than open ones, because they give a full account of how relatedness emerges from demography. Open models can be useful, particularly as a way of interpreting their counterpart closed models, by detaching relatedness from model parameters,

so that the causal effect of kin selection on model outcomes can be isolated and examined (Frank, 1998; Cooper *et al.*, 2018).

In the closed models considered, suppressing a cue of predator-presence could evolve (if population regulation precedes migration), but signalling the presence of predator presence could not. The take-home message is therefore that suppressing a cue of predator-presence is likely to evolve more permissively than signalling predator-presence.

Appendix J

Model J: signalling; open & closed; population regulation before migration; individuals may signal dishonestly and not respond to signals.

In this appendix, we generalise Model A, so that individuals can choose to signal dishonestly (*i.e.*, warn others of predator-presence when the predator isn't there), and choose not to respond to the signal (*i.e.*, not prepare for predator-attack when warned by another individual). We have written the verbal model description for this appendix entry in full, without referring to previous appendices, so that it can be read in isolation. In this appendix, we examine both 'closed' (emergent relatedness) and 'open' (parameterised relatedness) versions of the model, starting with the closed version.

We assume an infinite population of individuals (*e.g.*, plants). We assume for simplicity that individuals are haploid and asexual. The population is split into demes of size N (infinite island model) (Rousset, 2004). We focus our analysis on a focal individual, drawn at random from the population.

Each generation, with probability p, the population is attacked by predators (with probability 1-p, the population is not attacked). In generations where the population is attacked, a random individual on each deme is initially attacked by a predator, and suffers a fecundity cost of d. Given that there are N individuals on each deme, each individual has a $1/N$ chance of being initially attacked by the predator. The initially-predated individual may then invest into the production of a signal. If the focal individual is the initially-predated individual on its deme, it invests an amount x_{focal} into signal production. Signal investment incurs a fecundity cost that scales linearly with c. Therefore, the focal individual pays a signalling cost of cx_{focal} when it is initially-predated.

The signal produced by the initially-predated individual is transferred to (and recognised by) the other N-1 individuals on the deme. We make no assumptions about how the signal is transferred, but it may be being transferred through a common mycorrhizal network, or through the air, *etc*. The signal warns the other plants that a predator is in the vicinity. Consequently, the other plants can prepare for being attacked, and defend themselves, meaning they can reduce their fecundity cost of predation. Plants may choose the extent to which they actually respond to the signal, preparing for predator-attack, rather than ignoring the signal. Preparation for predator attack (defence) incurs a fecundity cost that scales linearly with s. We denote the average signalling investment amongst the N-1 other plants on the focal individual's

deme by x_{others}. We denote the extent of signal-response by the focal individual by y_{focal}. This means that, when the focal individual is not the initially-predated individual, it will ultimately suffer a cost of predation that is given by $d(1-x_{others}y_{focal})$, and a defence cost that is given by $sx_{others}y_{focal}$. We assume that preparing for predator attack is less costly than the fecundity cost of being predated ($d>s$), which ensures that individuals are favoured to defend themselves against predators.

We emphasise that, in our model, only the initially-predated individual has a chance to warn the other individuals about the presence of a predator. Subsequently-predated individuals may still produce the signal, but our assumption is that by then, the predator has already arrived, rendering the signal useless for warning others about predator presence. We allow for the possibility that subsequently-predated (*i.e.*, not initially-predated) individuals can facultatively turn off signal production, so that they do not pay the costs of producing a useless signal. We assume that the focal individual pays a signalling cost of $cx_{focal}(1-g)$ when subsequently-predated, where g denotes the extent to which signal investment is 'facultative' as opposed to 'obligate' (analogously, a non-focal individual pays a signalling cost of $cx_{others}(1-g)$ when subsequently-predated). A fully obligate signal ($g=0$) means that it is produced irrespective of whether the individual is initially-predated (rendering the signal is useful) or subsequently-predated (rendering the signal useless).

In generations where the population is not attacked by predators (such generations occur with probability $1-p$), a random individual on each deme is given the opportunity to feign being an initially-predated individual, producing the signal of predator-presence even though no predator is present. In such cases, any signal produced is 'dishonest'. If the focal individual is given the opportunity to be the dishonest signaller, it invests $cx_{focal}z_{focal}$ into signal production, where z_{focal} denotes dishonesty, with $z_{focal}=0$ corresponding to no dishonest signal production, and $z_{focal}=1$ corresponding to the same investment in signal production irrespective of whether the predator is present or not. If the focal individual is not given the opportunity to be the dishonest signaller, it will suffer a cost of preparation for predator attack that is given by $sx_{others}y_{focal}z_{others}$, and a cost of obligate (redundant) dishonest signal production of $cx_{focal}(1-g)z_{focal}$.

After the fecundity costs of predation and signal investment have been incurred by all individuals, each individual produces a large number of offspring (juvenile haploid clones) in proportion to their fecundity. A random sample of N juvenile individuals are chosen, for each deme, to survive and form the next adult population (local population regulation). After population regulation, a proportion of the (new) adult population, m, migrate to a different

deme, randomly chosen for each individual. The remaining proportion, *1-m*, remain on their local deme. This lifecycle then iterates over many generations until an evolutionary end point is reached.

We denote the population average investment into signalling by x_{pop}. We denote the average signalling investment on the focal individual's deme (incorporating its own signalling investment) by x_{deme}; this can be written in terms of x_{focal} and x_{others} as $x_{deme} = \frac{(N-1)x_{others}}{N} + \frac{x_{focal}}{N}$ (Equation A1). Similarly, we denote the population average investment into signal response and dishonesty, respectively, by y_{pop} & z_{pop}, and the focal-deme-average investment into signal response and dishonesty, respectively, by y_{deme} & z_{deme}, where

$$y_{deme} = \frac{(N-1)y_{others}}{N} + \frac{y_{focal}}{N}, \qquad (J1)$$

$$z_{deme} = \frac{(N-1)z_{others}}{N} + \frac{z_{focal}}{N}. \qquad (J2)$$

We assume that the baseline fecundity is 1, that $0 \leq x_{focal}, x_{others}, x_{deme}, x_{pop}, y_{focal}, y_{others}, y_{deme}, y_{pop}, z_{focal}, z_{others}, z_{deme}, z_{pop} \leq 1$, and that $c + d + s \leq 1$; these assumptions ensure that fecundity never falls below zero. The number of juvenile offspring produced by the focal individual is then proportional to

$$
\begin{aligned}
f_{focal} = 1 - p\Bigg(&\frac{1}{N}\big(d + cx_{focal}\big) \\
&+ \frac{N-1}{N}\big(d\big(1 - x_{others}y_{focal}\big) + cx_{focal}(1 - g) + sx_{others}y_{focal}\big)\Bigg) \\
&- (1 - p)\Bigg(\frac{1}{N}\big(cx_{focal}z_{focal}\big) + \frac{N-1}{N}\big(cx_{focal}(1 - g)z_{focal}\big) \\
&+ sx_{others}y_{focal}z_{others}\Bigg).
\end{aligned}
\qquad (J3)
$$

We can make sense of the right-hand side of Equation J3 as follows. Baseline fecundity is one (left-hand term). With probability p, a predator attacks. Given predator-attack, with probability $1/N$, the focal individual is predated first, leading to additively applied costs of predation (*d*) and signal production (*cx_{focal}*). Given predator-attack, with probability $\frac{N-1}{N}$, the focal individual is not predated first, and can receive a signal of predator presence from another

individual on the deme, leading to additively applied costs of predation ($d(1-x_{others})$), defence ($sx_{others}y_{focal}$), and the costs of signal production that are incurred even when the focal individual is not predated first, owing to obligatory (unconditional) expression ($cx_{focal}(1-g)$). With probability $1-p$, a predator does not attack. Given predator-absence, with probability $1/N$, the focal individual is given the opportunity to feign predator-presence, leading to an additively applied cost of dishonest signal production ($cx_{focal}z_{focal}$). Given predator-absence, with probability $\frac{N-1}{N}$, the focal individual is not given the opportunity to feign predator-presence, and can receive a dishonest signal of predator presence from another individual on the deme, leading to additively applied costs of defence ($sx_{others}y_{focal}z_{others}$), and the costs of dishonest signal production that are incurred even when the focal individual is not given the opportunity to feign predator-presence, owing to obligatory (unconditional) expression ($cx_{focal}(1-g)z_{focal}$).

The number of juvenile offspring produced by an individual drawn randomly from the focal individual's deme (including the focal individual) is proportional to

$$
\begin{aligned}
f_{deme} = 1 - p & \left(\frac{1}{N}(d + cx_{deme}) \right. \\
& \left. + \frac{N-1}{N}(d(1 - x_{deme}y_{deme}) + cx_{deme}(1-g) + sx_{deme}y_{deme}) \right) \\
& - (1-p)\left(\frac{1}{N}(cx_{deme}z_{deme}) + \frac{N-1}{N}(cx_{deme}(1-g)z_{deme}) \right. \\
& \left. + sx_{deme}y_{deme}z_{deme} \right),
\end{aligned}
\tag{J4}
$$

which we can rewrite using Equations A1, J1 & J2 (getting rid of the superfluous x_{deme}, y_{deme} & z_{deme} parameters) as

$$f_{deme} = 1 - p\left(\frac{1}{N}\left(d + c\left(\frac{(N-1)x_{others}}{N} + \frac{x_{focal}}{N}\right)\right)\right.$$

$$+ \frac{N-1}{N}\left(d\left(1 - \left(\frac{(N-1)x_{others}}{N} + \frac{x_{focal}}{N}\right)\left(\frac{(N-1)y_{others}}{N} + \frac{y_{focal}}{N}\right)\right)\right.$$

$$+ c\left(\frac{(N-1)x_{others}}{N} + \frac{x_{focal}}{N}\right)(1-g)$$

$$\left.\left.+ s\left(\frac{(N-1)x_{others}}{N} + \frac{x_{focal}}{N}\right)\left(\frac{(N-1)y_{others}}{N} + \frac{y_{focal}}{N}\right)\right)\right)$$

$$- (1-p)\left(\frac{1}{N}\left(c\left(\frac{(N-1)x_{others}}{N} + \frac{x_{focal}}{N}\right)\left(\frac{(N-1)z_{others}}{N} + \frac{z_{focal}}{N}\right)\right)\right.$$

$$+ \frac{N-1}{N}\left(\left(\frac{(N-1)x_{others}}{N} + \frac{x_{focal}}{N}\right)(1-g)\left(\frac{(N-1)z_{others}}{N} + \frac{z_{focal}}{N}\right)\right)$$

$$+ s\left(\frac{(N-1)x_{others}}{N} + \frac{x_{focal}}{N}\right)\left(\frac{(N-1)y_{others}}{N} + \frac{y_{focal}}{N}\right)\left(\frac{(N-1)z_{others}}{N}\right.$$

$$\left.\left.\left.+ \frac{z_{focal}}{N}\right)\right)\right). \tag{J5}$$

The number of juvenile offspring produced by an individual drawn randomly from the population is proportional to

$$f_{pop} = 1 - p\left(\frac{1}{N}(d + cx_{pop}) + \frac{N-1}{N}\left(d(1 - x_{pop}y_{pop}) + cx_{pop}(1-g) + sx_{pop}y_{pop}\right)\right)$$

$$- (1-p)\left(\frac{1}{N}(cx_{pop}z_{pop}) + \frac{N-1}{N}(cx_{pop}(1-g)z_{pop})\right.$$

$$\left.+ sx_{pop}y_{pop}z_{pop}\right). \tag{J6}$$

The fitness of the focal individual is then given by Equation A6 $\left(w = \frac{f_{focal}}{f_{deme}}\right)$, where explicit expressions for the fecundity (f_{focal}, f_{deme}) functions are provided in Equations J3 & J5 (Rousset, 2004; Roze & Rousset, 2008; Lehmann & Rousset, 2010). Note that the f_{pop} function derived

above (Equation J6) does not feature in this fitness function, but it will feature in the fitness function of subsequent models. Equation A6 reflects the fact that, under our assumption that (local) population regulation occurs before migration, individuals only compete with competitors on their native deme, rather than with individuals on other demes (see Appendix A for more discussion).

Our aim is to examine what level of signalling, signal-response and signal-dishonesty evolve at evolutionary equilibrium. To do this, we assume a monomorphic population, where all individuals have the same level of signalling as each other, given by x^*, as well as the same level of signal-response as each other, given by y^*, and the same level of signal-dishonesty as each other, given by z^* (the lack of subscripts imply that the values are the same across the population).

We can then take a random (focal) individual and mutate its (and its identical-by-descent relatives') level of signalling to a deviant value x, and ask how the fitness of the focal individual changes in response, in effect calculating $\frac{dw}{dx}$. Analogously, we can mutate a random individual's signal-response to y and calculate $\frac{dw}{dy}$. We can mutate a random individual's signal-dishonesty to z and calculate $\frac{dw}{dz}$. Evaluating these derivatives at equilibrium $(x_{focal}, x_{others}, x_{pop} = x^*, y_{focal}, y_{others}, y_{pop} = y^*, z_{focal}, z_{others}, z_{pop} = z^*)$, the equilibrium (ESS) levels of signalling (x^*), signal-response (y^*) and signal-dishonesty (z^*) can then be obtained, under the assumption of no genetic association (linkage disequilibrium) between traits, as the values for which $\frac{dw}{dx}|_{x=x^*,y=y^*,z=z^*}, \frac{dw}{dy}|_{x=x^*,y=y^*,z=z^*}, \frac{dw}{dz}|_{x=x^*,y=y^*,z=z^*} = 0$ is satisfied (Brown & Taylor, 2010).

We start by calculating $\frac{dw}{dz}|_{x=x^*,y=y^*,z=z^*}$ (mutating signal-dishonesty). We do so by expanding it using the chain rule

$$\frac{dw}{dz} = \frac{\partial w}{\partial z_{focal}} \frac{dz_{focal}}{dz} + \frac{\partial w}{\partial z_{others}} \frac{dz_{others}}{dz}, \qquad (J7)$$

and then evaluating the derivatives on the right-hand side (Taylor & Frank, 1996; Frank, 1998; Brown & Taylor, 2010). Because the population is monomorphic, and because we are interested in the long-term evolutionary state of the population, we evaluate these derivatives at the point

where $x_{focal}, x_{others}, x_{pop} = x^*, y_{focal}, y_{others}, y_{pop} = y^*, z_{focal}, z_{others}, z_{pop} = z^*$. We trivially obtain

$$\frac{dz_{focal}}{dz}\bigg|_{x=x^*, y=y^*, z=z^*} = 1. \tag{J8}$$

We obtain the equilibrium-evaluated partial derivatives $\frac{\partial w}{\partial z_{focal}}\big|_{x=x^*, y=y^*, z=z^*}$ and $\frac{\partial w}{\partial z_{others}}\big|_{x=x^*, y=y^*, z=z^*}$ respectively as

$$\frac{\partial w}{\partial z_{focal}}\bigg|_{x=x^*, y=y^*, z=z^*}$$

$$= \Big((N-1)(p-1)x^*(c(g-1)N - cg + sy^*)\Big)$$

$$/ \Bigg(N\Big(x^*\Big(c((g-1)N - g)((p-1)z^* - p)$$

$$- (N-1)y^*(p(d-s) + (p-1)sz^*)\Big) + N(dp-1)\Big)\Bigg) \tag{J9}$$

and

$$\frac{\partial w}{\partial z_{others}}\bigg|_{x=x^*, y=y^*, z=z^*}$$

$$= -\Big((N-1)(p-1)x^*(c(g-1)N - cg + sy^*)\Big)$$

$$/ \Bigg(N\Big(x^*\Big(c((g-1)N - g)((p-1)z^* - p)$$

$$- (N-1)y^*(p(d-s) + (p-1)sz^*)\Big) + N(dp-1)\Big)\Bigg). \tag{J10}$$

We obtain $\frac{dz_{others}}{dx}\big|_{x=x^*, y=y^*, z=z^*}$ by noting that it is interpretable as a coefficient of relatedness, since it denotes the marginal (correlated) change in signal production by a social partner as a consequence of a change in signal production by an actor (Taylor & Frank, 1996; Frank, 1998). We can therefore write

$$\frac{dz_{others}}{dx}\bigg|_{x=x^*, y=y^*, z=z^*} = r^*, \tag{J11}$$

where r^* is the long-term (equilibrium) value of relatedness. We have already calculated r^* for our demographic assumptions (Appendix A; Equation A16), and we can substitute this into Equation J11 to obtain

$$\frac{dz_{others}}{dx}\Big|_{x=x^*,y=y^*,z=z^*} = \frac{1}{1 - \frac{(m-2)mN}{(m-1)^2}}. \tag{J12}$$

We can now substitute Equations J8, J9, J10 & J12 into Equation J7 to obtain an explicit expression for $\frac{dw}{dz}\big|_{x=x^*,y=y^*,z=z^*}$. Setting $\frac{dw}{dz}\big|_{x=x^*,y=y^*,z=z^*} = 0$ and simplifying, we obtain

$$y^* = \frac{Nc\left(1 - \frac{N-1}{N}g\right)}{s}. \tag{J13}$$

The equilibrium level of signal-dishonesty (z^*) corresponds to the value of z^* for which Equation J13 is satisfied (note that we haven't obtained a specific value for z^* yet). To interpret Equation J13, note that $c\left(1 - \frac{N-1}{N}g\right)$ gives the cost of (dishonest) signalling to the actor. $\frac{y^*s}{N}$ gives the cost of responding to a dishonest signal, borne by a competitor. Signal dishonesty is favoured to increase when it results in a greater cost to competitors than to self, meaning that, at an internal equilibrium, these costs to competitors and self are equalised, leading to $\frac{y^*s}{N} = c\left(1 - \frac{N-1}{N}g\right)$, which with some slight rearrangement gives Equation J13.

Next, we calculate $\frac{dw}{dy}\big|_{x=x^*,y=y^*,z=z^*}$ (mutating signal-response). We do so by expanding it using the chain rule

$$\frac{dw}{dy} = \frac{\partial w}{\partial y_{focal}}\frac{dy_{focal}}{dy} + \frac{\partial w}{\partial y_{others}}\frac{dy_{others}}{dy}, \tag{J14}$$

and then evaluating the derivatives on the right-hand side at equilibrium (Taylor & Frank, 1996; Frank, 1998; Brown & Taylor, 2010). We trivially obtain

$$\frac{dy_{focal}}{dy}\Big|_{x=x^*,y=y^*,z=z^*} = 1. \tag{J15}$$

We obtain the equilibrium-evaluated partial derivatives $\frac{\partial w}{\partial y_{focal}}|_{x=x^*,y=y^*,z=z^*}$ and

$\frac{\partial w}{\partial y_{others}}|_{x=x^*,y=y^*,z=z^*}$ respectively as

$$\frac{\partial w}{\partial y_{focal}}|_{x=x^*,y=y^*,z=z^*}$$
$$= \left((N-1)^2 x^* (p(d-s) + (p-1)sz^*)\right)$$
$$/ \left(N \left(x^* \left((N-1)y^* (p(d-s) + (p-1)sz^*) \right. \right. \right.$$
$$\left. \left. \left. - c((g-1)N-g)((p-1)z^*-p) \right) - dNp + N \right) \right) \tag{J16}$$

and

$$\frac{\partial w}{\partial y_{others}}|_{x=x^*,y=y^*,z=z^*}$$
$$= -\left((N-1)^2 x^* (p(d-s) + (p-1)sz^*)\right)$$
$$/ \left(N \left(x^* \left((N-1)y^* (p(d-s) + (p-1)sz^*) \right. \right. \right.$$
$$\left. \left. \left. - c((g-1)N-g)((p-1)z^*-p) \right) - dNp + N \right) \right). \tag{J17}$$

We obtain $\frac{dy_{others}}{dx}|_{x=x^*,y=y^*,z=z^*}$ by noting that it is interpretable as a coefficient of relatedness, meaning we can write

$$\frac{dy_{others}}{dx}|_{x=x^*,y=y^*,z=z^*} = r^*, \tag{J18}$$

where r^* is the long-term (equilibrium) value of relatedness. Substituting in our explicit expression for r^* (Equation A16), we obtain

$$\frac{dy_{others}}{dx}|_{x=x^*,y=y^*,z=z^*} = \frac{1}{1 - \frac{(m-2)mN}{(m-1)^2}}. \tag{J19}$$

We can now substitute Equations J15, J16, J17 & J19 into Equation J14 to obtain an explicit expression for $\frac{dw}{dy}|_{x=x^*,y=y^*,z=z^*}$. Setting $\frac{dw}{dy}|_{x=x^*,y=y^*,z=z^*} = 0$ and simplifying, we obtain

$$z^* = \frac{p(d-s)}{(1-p)s}. \tag{J20}$$

The equilibrium level of signal-response (y^*) corresponds to the value of y^* for which Equation J20 is satisfied (note that the specific value for y^* is provided in Equation J13). To interpret Equation J20, note that $\frac{N-1}{N}s\big((1-p)z^* + p\big)$ gives the cost of responding to a signal – specifically, it is the defence cost (s) weighted by the proportion of generations where this defence cost is paid $\frac{N-1}{N}\big((1-p)z^* + p\big)$. $\frac{N-1}{N}pd$ gives the benefit of responding to a signal – specifically, it is the benefit of a reduced predation cost (d), weighted by the proportion of generations where benefit can be accrued $\left(\frac{N-1}{N}p\right)$. Signal response is favoured to increase when it results in a greater benefit than cost to the responder, meaning that, at an internal equilibrium, this benefit and cost are equalised, leading to $pd = s\big((1-p)z + p\big)$, which with some rearrangement gives Equation J20.

Finally, we calculate $\frac{dw}{dx}|_{x=x^*,y=y^*,z=z^*}$ (mutating signal investment). We do so by expanding it using the chain rule $\frac{dw}{dx} = \frac{\partial w}{\partial x_{focal}}\frac{dx_{focal}}{dx} + \frac{\partial w}{\partial x_{others}}\frac{dx_{others}}{dx}$ (Equation A7), and evaluating the derivatives on the right-hand side at equilibrium (Taylor & Frank, 1996; Frank, 1998; Brown & Taylor, 2010). We trivially obtain

$$\frac{dx_{focal}}{dx}|_{x=x^*,y=y^*,z=z^*} = 1. \tag{J21}$$

We obtain the equilibrium-evaluated partial derivatives $\frac{\partial w}{\partial x_{focal}}|_{x=x^*,y=y^*,z=z^*}$ and $\frac{\partial w}{\partial x_{others}}|_{x=x^*,y=y^*,z=z^*}$ respectively as

$$\frac{\partial w}{\partial x_{focal}}\Big|_{x=x^*,y=y^*,z=z^*}$$

$$= -\Big((N-1)\Big(c(g(N-1)-N)((p-1)z^*-p)$$

$$+ y^*(p(d-s)+(p-1)sz^*)\Big)\Big)$$

$$/ \Big(N\Big(x^*\Big((N-1)y^*(p(d-s)+(p-1)sz^*)$$

$$- c((g-1)N-g)((p-1)z^*-p)\Big)-dNp+N\Big)\Big) \tag{J22}$$

and

$$\frac{\partial w}{\partial x_{others}}\Big|_{x=x^*,y=y^*,z=z^*}$$

$$= \Big((N-1)\Big(c(g(N-1)-N)((p-1)z^*-p)$$

$$+ y^*(p(d-s)+(p-1)sz^*)\Big)\Big)$$

$$/ \Big(N\Big(x^*\Big((N-1)y^*(p(d-s)+(p-1)sz^*)$$

$$- c((g-1)N-g)((p-1)z^*-p)\Big)-dNp+N\Big)\Big). \tag{J23}$$

We obtain $\frac{dx_{others}}{dx}\Big|_{x=x^*,y=y^*,z=z^*}$ by noting that it is interpretable as a coefficient of relatedness, meaning we can write

$$\frac{dx_{others}}{dx}\Big|_{x=x^*,y=y^*,z=z^*} = r^*, \tag{J24}$$

where r^* is the long-term (equilibrium) value of relatedness. Substituting in our explicit expression for r^* (Equation A16), we obtain

$$\frac{dx_{others}}{dx}\Big|_{x=x^*,y=y^*,z=z^*} = \frac{1}{1 - \frac{(m-2)mN}{(m-1)^2}}. \tag{J25}$$

We can now substitute Equations J21, J22, J23 & J25 into Equation A7 to obtain an explicit expression for $\frac{dw}{dx}|_{x=x^*,y=y^*,z=z^*}$. Instead of setting $\frac{dw}{dx}|_{x=x^*,y=y^*,z=z^*} = 0$ like we did for $\frac{dw}{dy}|_{x=x^*,y=y^*,z=z^*}$ and $\frac{dw}{dz}|_{x=x^*,y=y^*,z=z^*}$, we set $\frac{dw}{dx}|_{x=x^*,y=y^*,z=z^*} > 0$ to see when signalling is favoured by natural selection (rather than when it is at an evolutionary equilibrium). We obtain

$$c(p + (1-p)z^*)\left(1 - \frac{N-1}{N}g\right) < \frac{y^*}{N}(s(p + (1-p)z^*) - dp). \qquad (J26)$$

To interpret Equation J26, note that the left-hand side gives the overall cost of signalling to the actor, and the right-hand side gives the overall cost of signalling incurred by receivers. Signalling is then favoured in this lifecycle if hurts competitors more than oneself. Specifically, the cost of signalling to the actor (left-hand side) is obtained as the cost of signal production $\left(c\left(1 - \frac{N-1}{N}g\right)\right)$ weighted by the proportion of generations where a signal is produced $\left((1-p)z^* + p\right)$. The cost of signalling to receivers (right-hand side) is obtained as the cost of defence (s/N), where this is weighted by the proportion of generations where a signal is produced $\left((1-p)z^* + p\right)$ as well as by the response to the signal (y^*), minus the saved cost of predation (d/N), where this is weighted by the proportion of generations where a predator is present (p) as well as by the response to the signal (y^*).

We can obtain explicit values for y^* and z^*. These explicit values will often be the values of y^* and z^* specified by Equations J13 & J20. However, y^* and z^* must lie in the range 0 to 1, but the value of y^* or z^* specified by Equations J13 & J20 may sometimes go above 1. In such cases, the true (obtained) value for y^* or z^* will be 1. Similarly, the value of y^* or z^* specified by Equations J13 & J20 may sometimes fall below zero. In such cases, the true (obtained) value for y^* or z^* will be 0. Substituting these explicit values for y^* and z^* into Equation J26, we find that Equation J26 is never satisfied, meaning signalling is never favoured ($x^*=0$) (see supplementary *Matlab* file for a numerical demonstration of this).

This recovers the result of Model A. Therefore, the result of Model A, that signalling does not evolve, still holds, after generalising the model to allow dishonest signalling and submaximal signal-response. The reason why this previous result holds is that, in this generalised model, signals either evolve to be relatively more dishonest or honest. Dishonest signals harm competitors by inducing them to defend themselves against predators that aren't

even there, giving the signaller a transient advantage, but inducing competitors to ignore the signal. Ultimately, the signal is ignored, meaning costly signal production is disfavoured. Conversely, honest signals help competitors by warning them about predators, giving the signaller a competitive disadvantage, disfavouring signal production. Therefore, irrespective of how honest or dishonest the signal evolves to be, signal production is ultimately disfavoured in this lifecycle.

We note that 'opening' this model, detaching relatedness from demographic parameters (m and N) so that it is a parameter in its own right, does not affect model outcomes. To appreciate this, note that calculating $\frac{dw}{dz}|_{x=x^*,y=y^*,z=z^*}$ with Equation J11 rather than Equation J12 (i.e., treating r* as a parameter rather than a function of demographic parameters) does not lead to a different expression for $\frac{dw}{dz}|_{x=x^*,y=y^*,z=z^*}$. Similarly, calculating $\frac{dw}{dy}|_{x=x^*,y=y^*,z=z^*}$ with Equation J18 rather than Equation J19 does not lead to a different expression for $\frac{dw}{dy}|_{x=x^*,y=y^*,z=z^*}$. Calculating $\frac{dw}{dx}|_{x=x^*,y=y^*,z=z^*}$ with Equation J24 rather than J25 does not lead to a different expression $\frac{dw}{dx}|_{x=x^*,y=y^*,z=z^*}$. This means that the results of the model (Equations J13, J20 & J26) are unchanged – relatedness does not affect model outcomes in this lifecycle.

Appendix K

Model K: signalling; open & closed; population regulation after migration; individuals may signal dishonestly and not respond to signals.

In this appendix, we alter Model J so that population regulation occurs after migration rather than before. We examine both 'closed' (emergent relatedness) and 'open' (parameterised relatedness) versions of the model, starting with the open version. We found in Appendix D that signalling can evolve in the open version of this lifecycle (with population regulation occurring after migration). However, this result was obtained on the assumptions that signals are honest and individuals respond maximally to signals. In this appendix, we see if this result (that signalling can evolve) still holds when signal-dishonesty and signal-response can evolve.

Our assumption that population regulation occurs after migration means that the fitness of the focal individual is given by Equation C1 $\left(w = (1-m)\frac{f_{focal}}{(1-m)f_{deme}+mf_{pop}} + m\frac{f_{focal}}{f_{pop}} \right)$, where explicit expressions for the fecundity (f_{focal}, f_{deme}, f_{pop}) functions are provided in Equations J3, J5 & J6 (Rousset, 2004; Roze & Rousset, 2008; Lehmann & Rousset, 2010).

Our aim is to examine what level of signalling, signal-response and signal-dishonesty evolve at evolutionary equilibrium, as we did in Appendix J. We do so by first calculating $\frac{dw}{dz}|_{x=x^*,y=y^*,z=z^*}$, $\frac{dw}{dz}|_{x=x^*,y=y^*,z=z^*}$ & $\frac{dw}{dz}|_{x=x^*,y=y^*,z=z^*}$, by expanding them using the chain rule (Equations J7, J14 & A7), and evaluating the associated partial derivatives (Taylor & Frank, 1996; Frank, 1998; Brown & Taylor, 2010). We obtain the following partial derivatives:

$$\frac{\partial w}{\partial x_{focal}}|_{x=x^*,y=y^*,z=z^*}$$

$$= -\Big((m-1)^2(N-1)y^*(p(d-s) + (p-1)sz^*)$$

$$- c(g(N-1) - N)((m-1)^2 - N)\big((p-1)z^* - p\big)\Big)$$

$$/ \Big(N\Big(x^*\big((N-1)y^*(p(d-s) + (p-1)sz^*)$$

$$- c\big((g-1)N - g\big)\big((p-1)z^* - p\big)\big) - dNp + N\Big)\Big), \qquad (K1)$$

$$\frac{\partial w}{\partial y_{focal}}\Big|_{x=x^*,y=y^*,z=z^*}$$

$$= -\big((N-1)x^*((m-1)^2 - N)(p(d-s) + (p-1)sz^*)\big)$$

$$\Big/ \Big(N \Big(x^* \big((N-1)y^*(p(d-s) + (p-1)sz^*) $$

$$- c\big((g-1)N - g\big)\big((p-1)z^* - p\big) \Big) - dNp + N \Big) \Big), \qquad (K2)$$

$$\frac{\partial w}{\partial z_{focal}}\Big|_{x=x^*,y=y^*,z=z^*}$$

$$= \Big((p-1)x^*\big((m-1)^2(N-1)sy^* - c(g(N-1) - N)((m-1)^2 - N)\big) \Big)$$

$$\Big/ \Big(N \Big(x^* \big(c\big((g-1)N - g\big)\big((p-1)z^* - p\big) $$

$$- (N-1)y^*(p(d-s) + (p-1)sz^*) \big) + N(dp-1) \Big) \Big), \qquad (K3)$$

$$\frac{\partial w}{\partial x_{others}}\Big|_{x=x^*,y=y^*,z=z^*}$$

$$= \Big((N-1)\Big(c(m-1)^2(g(N-1) - N)\big((p-1)z^* - p\big) $$

$$- y^*\big((m-2)m(N-1) - 1\big)(p(d-s) + (p-1)sz^*) \Big) \Big)$$

$$\Big/ \Big(N \Big(x^* \big((N-1)y^*(p(d-s) + (p-1)sz^*) $$

$$- c\big((g-1)N - g\big)\big((p-1)z^* - p\big) \Big) - dNp + N \Big) \Big), \qquad (K4)$$

$$\frac{\partial w}{\partial y_{others}}\Big|_{x=x^*,y=y^*,z=z^*}$$

$$= -\big((m-1)^2(N-1)^2 x^*(p(d-s) + (p-1)sz^*)\big)$$

$$\Big/ \Big(N \Big(x^* \big((N-1)y^*(p(d-s) + (p-1)sz^*) $$

$$- c\big((g-1)N - g\big)\big((p-1)z^* - p\big) \Big) - dNp + N \Big) \Big), \qquad (K5)$$

$$\frac{\partial w}{\partial z_{others}}\Big|_{x=x^*,y=y^*,z=z^*}$$

$$= -\Big((N-1)(p-1)x^*\Big(c(m-1)^2(g(N-1)-N)$$

$$+ sy^*(1-(m-2)m(N-1))\Big)\Big)$$

$$/\Big(N\Big(x^*\Big(c((g-1)N-g)((p-1)z^*-p)$$

$$- (N-1)y^*(p(d-s)+(p-1)sz^*)\Big)+N(dp-1)\Big)\Big). \tag{K6}$$

We can now substitute Equations J21, J24, K1 & K4 into Equation A7 to obtain an explicit expression for $\frac{dw}{dx}\big|_{x=x^*,y=y^*,z=z^*}$. Analogously, we can substitute Equations J15, J18, K2 & K5 into Equation J14 to obtain an explicit expression for $\frac{dw}{dy}\big|_{x=x^*,y=y^*,z=z^*}$. We can substitute Equations J8, J11, K3 & K6 into Equation J7 to obtain an explicit expression for $\frac{dw}{dz}\big|_{x=x^*,y=y^*,z=z^*}$.

We find that the explicit value for $\frac{dw}{dy}\big|_{x=x^*,y=y^*,z=z^*}$ is the same in this model as it was for Model J (Equation J20). Setting $\frac{dw}{dy}\big|_{x=x^*,y=y^*,z=z^*} > 0$ and simplifying, we obtain

$$z^* < \frac{p(d-s)}{(1-p)s}. \tag{K7}$$

Setting $\frac{dw}{dz}\big|_{x=x^*,y=y^*,z=z^*} > 0$ and simplifying, we obtain

$$\Big((p-1)x^*\Big((N-1)r^*\Big(c(m-1)^2((g-1)N-g)+sy^*(1-(m-2)m(N-1))\Big)$$

$$+ c((g-1)N-g)((m-1)^2-N)-((m-1)^2(N-1)sy^*)\Big)\Big)$$

$$/\Big(N\Big(x^*\Big(c((g-1)N-g)((p-1)z^*-p)$$

$$- (N-1)y^*(p(d-s)+(p-1)sz^*)\Big)+N(dp-1)\Big)\Big) > 0. \tag{K8}$$

Setting $\frac{dw}{dx}\big|_{x=x^*,y=y^*,z=z^*} > 0$ and simplifying, we obtain

$$r^* > \Big((m-1)^2(N-1)y^*(p(d-s)+(p-1)sz^*)$$

$$-c(g(N-1)-N)((m-1)^2-N)\big((p-1)z^*-p\big)\Big)$$

$$/\Big((N-1)\Big(c(m-1)^2(g(N-1)-N)\big((p-1)z^*-p\big)$$

$$-y^*\big((m-2)m(N-1)-1\big)(p(d-s)+(p-1)sz^*)\Big)\Big). \qquad (K9)$$

Signal-response is favoured if Equation K7 is satisfied. Conversely, signal-response is disfavoured if $\frac{dw}{dy}\big|_{x=x^*,y=y^*,z=z^*} < 0$, and at an internal ESS if $\frac{dw}{dy}\big|_{x=x^*,y=y^*,z=z^*} = 0$. Signal-dishonesty is favoured if Equation K8 is satisfied. Conversely, signal-dishonesty is disfavoured if $\frac{dw}{dz}\big|_{x=x^*,y=y^*,z=z^*} < 0$, and at an internal ESS if $\frac{dw}{dz}\big|_{x=x^*,y=y^*,z=z^*} = 0$. Signalling is favoured Equation K9 is satisfied. Conversely, signalling is disfavoured if $\frac{dw}{dx}\big|_{x=x^*,y=y^*,z=z^*} < 0$, and at an internal ESS if $\frac{dw}{dx}\big|_{x=x^*,y=y^*,z=z^*} = 0$.

Equations K7, K8 & K9 describe how selection acts on each of the three traits (response, dishonesty & signalling). We cannot solve this model analytically, so we use numerical methods to elucidate what trait values obtain in the evolutionary long term (see supplementary *Matlab* file). Our approach assumes that there is no genetic correlation (linkage disequilibrium) between the traits (Brown & Taylor, 2010). Specifically, we assume some initial values for *x, y* & *z* and use them to calculate the selection differential for each of the three traits $(\frac{dw}{dy}\big|_{x=x^*,y=y^*,z=z^*}, \frac{dw}{dz}\big|_{x=x^*,y=y^*,z=z^*}, \frac{dw}{dx}\big|_{x=x^*,y=y^*,z=z^*})$. If the selection differential is positive, we add on a very small increment to the previous trait value; if the selection differential is negative, we subtract a very small increment; if the selection differential is zero, we do not add or subtract an increment. We then use the new trait values to calculate the selection differentials, and repeat the process until trait values have stopped changing. We find that the resulting trait values are independent of initial trait values and of the size of the increment, as long as the increment is sufficiently small. The resulting trait values correspond to the long-term trait values: *x*, y** & *z**.

We find that, in this lifecycle, signals evolve to be honest and responded-to at equilibrium, and that signalling evolves in the same region of parameter space that it did in Model D, which assumed the same lifecycle but forced signals to be honest and responded-to.

Specifically, we find that, when the right-hand side of Equation D1 (which gives an explicit value for $\frac{dw}{dx}\big|_{x_{focal},x_{others},x_{pop}=x^*}$ for Model D) is greater than zero, signalling is favoured in the present model ($x^*=1$), and the signals are favoured to be honest ($z^*=0$) and responded-to ($y^*=1$). Conversely, when the right-hand side of Equation D1 is less than zero, signalling is not favoured in the present model ($x^*=0$), and any signals that are (transiently) present in the population are favoured to be dishonest ($z^*=1$) and ignored ($y^*=0$). The reader may wish to consult Fig. S1, which can be found in Appendix D – signalling, signal honesty and signal response are favoured above the lines.

As can be seen by inspecting Equation D1, a necessary requirement for the evolution of honest signalling is this is that there is sufficient relatedness and migration. The reason for this is that high relatedness leads to social interactions with kin, and high migration leads to competition with non-kin, and these combined conditions are necessary for the evolution of helping behaviours, such as honestly signalling the presence of a predator.

We can 'close' this model by substituting an explicit expression for relatedness (r^*) into our results conditions. Specifically, we substitute Equation A16 into Equations K8 and K9 and use the same numerical method to find long-term trait values. Doing so, we find that signalling can no longer evolve. This recovers the result of Model C, which made the same demographic assumptions (same lifecycle) but didn't allow signal-dishonesty or signal-response to evolve.

Appendix L

Model L: eavesdropping; open & closed; population regulation before migration; individuals may not respond to signals.

In plants, information about the presence of predators is often transmitted between individuals through common (shared) mycorrhizal networks. The mycorrhizal network is an evolutionary agent (individual) in itself, which raises the possibility that the mycorrhizal network might 'eavesdrop' on a plant, to find out when it is being attacked by a predator, and then signal this information to the other plants on the network. We examine this possibility in this Appendix. We examine both 'closed' (emergent relatedness) and 'open' (parameterised relatedness) versions of the model, starting with the closed version.

We assume an infinite populations of plants, as well as an infinite population of fungi. We assume for simplicity that all individuals (plants and fungi) are haploid and asexual. The plant and fungi populations are split into demes, where each deme has N plants and 1 fungus, where the N plants are connected to the fungus (common mycorrhizal network) and therefore to each other. We focus our analysis on a 'focal plant' and a 'focal fungus', drawn at random from the population.

Each generation, with probability p, the population is attacked by predators (with probability $1-p$, the population is not attacked). In generations where the population is attacked, a random plant on each deme is initially attacked by a predator, and suffers a fecundity cost of d. Given that there are N plants on each deme, each plant has a $1/N$ chance of being initially attacked by the predator. The fungus may then invest into 'eavesdropping & signalling' to learn that the initially-predated individual has been attacked, and warn the other N-1 plants on the deme of this. Specifically, the 'focal fungus' invests an amount $\tilde{x}_{focal}$ into eavesdropping & signalling (the tilde over the x denotes fungal rather than plant behaviour). Eavesdropping & signalling investment incurs a fecundity cost that scales linearly with c. Therefore, the focal fungus pays an eavesdropping & signalling cost of $c\tilde{x}_{focal}$.

The signal produced by the fungus is transferred to (and recognised by) the other N-1 individuals on the deme. The signal warns the other plants that a predator is in the vicinity. Consequently, the other plants can prepare for being attacked, and defend themselves, meaning they can reduce their fecundity cost of predation. Plants may choose the extent to which they actually respond to the signal, preparing for predator-attack, rather than ignoring the signal. Preparation for predator attack (defence) incurs a fecundity cost that scales linearly with s. We

assume that there is no genetic correlation between fungal and plant genotypes (i.e., no partner choice), which means that the focal plant's mycorrhizal network (fungus) invests $\tilde{x}_{pop}$ into eavesdropping & signalling, where the *pop* subscript denotes the (fungal) population average investment. We denote the extent of signal-response by the focal individual by y_{focal}. This means that, when the focal individual is not the initially-predated individual, it will ultimately suffer a cost of predation that is given by $d\left(1 - \tilde{x}_{pop}y_{focal}\right)$, and a defence cost that is given by $s\tilde{x}_{pop}y_{focal}$. We assume that preparing for predator attack is less costly than the fecundity cost of being predated (*d>s*), which ensures that individuals are favoured to defend themselves against predators.

In generations where the population is not attacked by predators (such generations occur with probability 1-*p*), fungal individuals are given the opportunity to produce the signal of predator-presence even though no predator is present. In such cases, any signal produced is 'dishonest'. The focal fungus invests $c\tilde{x}_{focal}\tilde{z}_{focal}$ into signal production, where $\tilde{z}_{focal}$ denotes dishonesty, with $\tilde{z}_{focal} = 0$ corresponding to no dishonest signal production, and $\tilde{z}_{focal} = 1$ corresponding to the same investment in signal production irrespective of whether the predator is present or not. Dishonest signalling means that the focal plant suffers a cost of preparation for predator attack that is given by $s\tilde{x}_{pop}y_{focal}\tilde{z}_{pop}$.

We assume that fungi gain a benefit by being connected to 'fitter' (more fecund) plant partners, because such partners will be better trade partners (*i.e.*, they will be in a position where they can acquire and transfer carbon to their fungal mutualist). Specifically, we assume that the focal fungus gains a benefit of $bf_{partner}$, where $f_{partner}$ gives the average fecundity of the plants connected to the focal fungus.

After fecundity effects have been incurred by all individuals, each individual produces a large number of offspring (juvenile haploid clones) in proportion to their fecundity. Population regulation and migration for the plant population then occurs as follows. A random sample of *N* juvenile plants are chosen, for each deme, to survive and form the next adult population (local population regulation). After population regulation, a proportion of the (new) adult plant population, *m,* migrate to a different deme, randomly chosen for each individual. The remaining proportion, *1-m*, remain on their local deme.

Population regulation and migration for the fungal population occurs as follows. Each member of the juvenile population migrates (with certainty) to a different deme, randomly chosen for each individual. A random juvenile individual is then chosen, for each deme, to survive and be the 'adult' fungus (mycorrhizal network) for that deme in the next generation.

Note that the assumption of certain migration followed by population regulation for the fungal population is taken to ensure that population regulation occurs effectively at the global rather than local level. Local population regulation would prevent evolution from occurring in this case, since there is just one genotype per deme, meaning natural selection would be 'choosing' between individuals with the same genotype. This lifecycle then iterates over many generations until an evolutionary end point is reached.

We denote the population average plant investment into signal response by y_{pop}. We denote the average investment into signal response on the focal plant's deme (incorporating its own signal response) by y_{deme}; this can be written in terms of y_{focal} and y_{others} as $y_{deme} = \frac{(N-1)y_{others}}{N} + \frac{y_{focal}}{N}$ (Equation J1). We assume that the baseline fecundity is 1, that $0 \leq \tilde{x}_{focal}, \tilde{x}_{pop}, y_{focal}, y_{others}, y_{deme}, y_{pop}, \tilde{z}_{focal}, \tilde{z}_{pop} \leq 1$, that $c \leq 1$ and that $d + s \leq 1$; these assumptions ensure that fecundity never falls below zero. The number of juvenile offspring produced by the focal plant (f_{focal}) and focal fungus ($\tilde{f}_{focal}$) are then, respectively, proportional to

$$f_{focal} = 1 - p\left(\frac{d}{N} + \frac{N-1}{N}\left(d\left(1 - \tilde{x}_{pop}y_{focal}\right) + s\tilde{x}_{pop}y_{focal}\right)\right)$$
$$- (1-p)s\tilde{x}_{pop}y_{focal}\tilde{z}_{pop} \tag{L1}$$

and

$$\tilde{f}_{focal} = 1 - pc\tilde{x}_{focal} - (1-p)c\tilde{x}_{focal}\tilde{z}_{focal} + bf_{partner}, \tag{L2}$$

where $f_{partner}$ denotes the average fecundity of plants connected to the focal fungus, and is proportional to

$$f_{partner} = 1 - p\left(\frac{d}{N} + \frac{N-1}{N}\left(d\left(1 - \tilde{x}_{focal}y_{pop}\right) + s\tilde{x}_{focal}y_{pop}\right)\right)$$
$$- (1-p)s\tilde{x}_{focal}y_{pop}\tilde{z}_{focal}. \tag{L3}$$

We can make sense of the right-hand side of Equation L1 as follows. Baseline plant fecundity is one (left-hand term). With probability p, a predator attacks. Given predator-attack,

with probability $1/N$, the focal plant is predated first, leading to an additively applied cost of predation (d). Given predator-attack, with probability $\frac{N-1}{N}$, the focal plant is not predated first, and can receive a signal of predator presence from the fungus it is connected to, leading to additively applied costs of predation $\left(d\left(1 - \tilde{x}_{pop}y_{focal}\right)\right)$ and defence ($s\tilde{x}_{pop}y_{focal}$). With probability 1-p, a predator does not attack. Given predator-absence, the fungus connected to the focal plant is given the opportunity to dishonestly signal predator-presence, meaning the focal plant suffers an additively applied cost of defence $\left(s\tilde{x}_{pop}y_{focal}\tilde{z}_{pop}\right)$. Equation L3 is obtained *mutatis mutandis*.

We can make sense of the right-hand side of Equation L2 as follows. Baseline fungus fecundity is one (left-hand term). With probability p, a predator attacks, meaning the focal fungus incurs an additively applied cost of honest signal production $\left(c\tilde{x}_{focal}\right)$. With probability 1-p, a predator does not attack, meaning the focal fungus incurs an additively applied cost of dishonest signal production $\left(c\tilde{x}_{focal}\tilde{z}_{focal}\right)$. The focal fungus also gains an additively applied benefit of being connected to high-quality (fecund) trade partners $\left(bf_{partner}\right)$.

The number of juvenile offspring produced by a plant drawn randomly from the focal plant's deme (including the focal individual) is proportional to

$$f_{deme} = 1 - p\left(\frac{d}{N} + \frac{N-1}{N}\left(d\left(1 - \tilde{x}_{pop}y_{deme}\right) + s\tilde{x}_{pop}y_{deme}\right)\right)$$
$$- (1-p)s\tilde{x}_{pop}y_{deme}\tilde{z}_{pop}, \tag{L4}$$

which we can rewrite using Equation J1 (getting rid of the superfluous y_{deme} parameter) as

$$f_{deme} = 1 - p \left(\frac{d}{N} \right.$$

$$+ \frac{N-1}{N} \left(d \left(1 - \tilde{x}_{pop} \left(\frac{(N-1)y_{others}}{N} + \frac{y_{focal}}{N} \right) \right) \right.$$

$$\left. + s\tilde{x}_{pop} \left(\frac{(N-1)y_{others}}{N} + \frac{y_{focal}}{N} \right) \right) \right)$$

$$\left. - (1-p)s\tilde{x}_{pop} \left(\frac{(N-1)y_{others}}{N} + \frac{y_{focal}}{N} \right) \tilde{z}_{pop}. \right. \tag{L5}$$

The number of juvenile offspring produced by a plant drawn randomly from the population is proportional to

$$f_{pop} = 1 - p \left(\frac{d}{N} + \frac{N-1}{N} \left(d(1 - \tilde{x}_{pop}y_{pop}) + s\tilde{x}_{pop}y_{pop} \right) \right)$$

$$- (1-p)s\tilde{x}_{pop}y_{pop}\tilde{z}_{pop}. \tag{L6}$$

The number of juvenile offspring produced by a fungus drawn randomly from the population is proportional to

$$\tilde{f}_{pop} = 1 - pc\tilde{x}_{pop} - (1-p)c\tilde{x}_{pop}\tilde{z}_{pop} + bf_{pop}. \tag{L7}$$

The fitness of the focal plant is then given by Equation A6 $\left(w = \frac{f_{focal}}{f_{deme}} \right)$, where explicit expressions for the fecundity (f_{focal}, f_{deme}) functions are provided in Equations L1 & L5 (Rousset, 2004; Roze & Rousset, 2008; Lehmann & Rousset, 2010). The fitness of the focal fungus is given by

$$\tilde{w} = \frac{\tilde{f}_{focal}}{\tilde{f}_{pop}}, \tag{L8}$$

where explicit expressions for the fecundity ($\tilde{f}_{focal}$, $\tilde{f}_{pop}$) functions are provided in Equations L2 & L7 (we also need Equation L6 to evaluate the f_{pop} term in Equation L7).

Our aim is to examine what level of fungal eavesdropping & signalling, plant signal-response, and fungal signal-dishonesty evolve at evolutionary equilibrium. To do this, we assume a monomorphic fungus population, where all fungi have the same level of eavesdropping & signalling as each other, given by $\tilde{x}^*$, as well as the same level of signal-dishonesty as each other, given by $\tilde{z}^*$. We additionally assume a monomorphic plant population, where all plants have the same level of signal-response as each other, given by y^*.

We can then take a random (focal) fungus and mutate its (and its identical-by-descent relatives') level of signalling to a deviant value x, and ask how the fitness of the focal fungus changes in response, in effect calculating $\frac{d\tilde{w}}{dx}$ (to reiterate, the tilde over the w is used to refer to fungal rather than plant fitness). Analogously, we can mutate a random fungus' signal-dishonesty to z and calculate $\frac{d\tilde{w}}{dz}$. Additionally, we can mutate a random plant's signal-response to y and calculate $\frac{dw}{dy}$. Evaluating these derivatives at equilibrium ($\tilde{x}_{focal}, \tilde{x}_{pop} = \tilde{x}^*, y_{focal}, y_{others}, y_{pop} = y^*, \tilde{z}_{focal}, \tilde{z}_{pop} = \tilde{z}^*$), the equilibrium (ESS) levels of signalling ($\tilde{x}^*$), signal-response (y^*) and signal-dishonesty ($\tilde{z}^*$) can then be obtained, under the assumption of no genetic association (linkage disequilibrium) between traits, as the values for which $\frac{d\tilde{w}}{dx}|_{\tilde{x}=\tilde{x}^*,y=y^*,\tilde{z}=\tilde{z}^*}, \frac{dw}{dy}|_{\tilde{x}=\tilde{x}^*,y=y^*,\tilde{z}=\tilde{z}^*}, \frac{d\tilde{w}}{dz}|_{\tilde{x}=\tilde{x}^*,y=y^*,\tilde{z}=\tilde{z}^*} = 0$ is satisfied (Frank, 1994; Foster & Wenseleers, 2006; Brown & Taylor, 2010).

We start by calculating $\frac{d\tilde{w}}{dx}|_{\tilde{x}=\tilde{x}^*,y=y^*,\tilde{z}=\tilde{z}^*}$ (mutating signal-dishonesty). We do so by expanding it using the chain rule

$$\frac{d\tilde{w}}{d\tilde{z}} = \frac{\partial \tilde{w}}{\partial \tilde{z}_{focal}} \frac{d\tilde{z}_{focal}}{d\tilde{z}} + \frac{\partial \tilde{w}}{\partial \tilde{z}_{others}} \frac{d\tilde{z}_{others}}{d\tilde{z}}, \qquad (L9)$$

and then evaluating the derivatives on the right-hand side at equilibrium (Taylor & Frank, 1996; Frank, 1998; Brown & Taylor, 2010). Because the plant and fungus populations are monomorphic, and because we are interested in the long-term evolutionary state of the population, we evaluate these derivatives at the point where $\tilde{x}_{focal}, \tilde{x}_{pop} = \tilde{x}^*, y_{focal}, y_{others}, y_{pop} = y^*, \tilde{z}_{focal}, \tilde{z}_{pop} = \tilde{z}^*$. We trivially obtain

$$\frac{d\tilde{z}_{focal}}{d\tilde{z}}\Big|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*} = 1. \tag{L10}$$

We obtain the equilibrium-evaluated partial derivatives $\frac{\partial \widetilde{w}}{\partial \tilde{z}_{focal}}\Big|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*}$ and

$\frac{\partial \widetilde{w}}{\partial \tilde{z}_{others}}\Big|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*}$ respectively as

$$\frac{\partial \widetilde{w}}{\partial \tilde{z}_{focal}}\Big|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*}$$

$$= \big(N(p-1)x^*(bsy^* + c)\big)$$
$$/\big(x^*(N(p-1)z^*(bsy^* + c) + b(N-1)py^*(d-s) - cNp)$$
$$+ N(b(-d)p + b + 1)\big) \tag{L11}$$

and

$$\frac{\partial \widetilde{w}}{\partial \tilde{z}_{others}}\Big|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*} = 0. \tag{L12}$$

Note that, because $\frac{\partial \widetilde{w}}{\partial \tilde{z}_{others}}\Big|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*} = 0$ (Equation L12), this gets rid of the $\frac{d\tilde{z}_{others}}{d\tilde{z}}\Big|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*}$ partial derivate in Equation L9, which means we don't need to evaluate $\frac{d\tilde{z}_{others}}{d\tilde{z}}\Big|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*}$ in order to evaluate the full derivative $\frac{d\widetilde{w}}{d\tilde{z}}\Big|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*}$.

We can now substitute Equations L10, L11 & L12 into Equation L9 to obtain an explicit expression for $\frac{d\widetilde{w}}{d\tilde{z}}\Big|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*}$. Setting $\frac{d\widetilde{w}}{d\tilde{z}}\Big|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*} < 0$ and simplifying, we obtain

$$y^* + \frac{c}{bs} > 0, \tag{L13}$$

which we can see is satisfied (as long as $y^*>0$ or $c>0$), meaning signal dishonesty is disfavoured, resulting in an equilibrium signal dishonesty of

$$\tilde{z}^* = 0, \tag{L14}$$

corresponding to completely honest signals. The reason for this result is that, in this model, the fungi are the signallers, and they can never gain a fitness advantage from signalling dishonestly. Dishonest signalling incurs an energetic cost to the fungus (proportional to c). Furthermore, it reduces the fecundity of the fungus' plant partners, since they invest in predator defence when no predator is present (proportional to s), which in turn reduces the fitness of the fungus as a consequence of having lower quality trade partners (proportional to b). Signalling dishonestly is therefore always bad for the signaller (fungus), meaning fungi are favoured to increase the honesty of their signals.

Next, we calculate $\frac{dw}{dy}\big|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*}$ (mutating signal-response). We do so by expanding it using the chain rule $\frac{dw}{dy} = \frac{\partial w}{\partial y_{focal}}\frac{dy_{focal}}{dy} + \frac{\partial w}{\partial y_{others}}\frac{dy_{others}}{dy}$ (Equation J14), and then evaluating the derivatives on the right-hand side at equilibrium (Taylor & Frank, 1996; Frank, 1998; Brown & Taylor, 2010). We trivially obtain

$$\frac{dy_{focal}}{dy}\big|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*} = 1. \qquad (L15)$$

We obtain the equilibrium-evaluated partial derivatives $\frac{\partial w}{\partial y_{focal}}\big|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*}$ and $\frac{\partial w}{\partial y_{others}}\big|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*}$ respectively as

$$\frac{\partial w}{\partial y_{focal}}\big|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*} = \frac{(N-1)x^*\big((N-1)p(d-s)+N(p-1)sz^*\big)}{N\big(x^*y^*\big((N-1)p(d-s)+N(p-1)sz^*\big)-dNp+N\big)} \qquad (L16)$$

and

$$\frac{\partial w}{\partial y_{others}}\big|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*} = -\frac{(N-1)x^*\big((N-1)p(d-s)+N(p-1)sz^*\big)}{N\big(x^*y^*\big((N-1)p(d-s)+N(p-1)sz^*\big)-dNp+N\big)}. \qquad (L17)$$

We obtain $\frac{dy_{others}}{dx}\big|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*}$ by noting that it is interpretable as a coefficient of relatedness (between plant individuals), meaning we can write

$$\frac{dy_{others}}{dx}\big|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*} = r^*, \qquad (L18)$$

where r^* is the long-term (equilibrium) value of plant relatedness. Substituting in our explicit expression for r^* (Equation A16), we obtain

$$\frac{dy_{others}}{dx}\Big|_{\tilde{x}=\tilde{x}^*,y=y^*,\tilde{z}=\tilde{z}^*} = \frac{1}{1-\frac{(m-2)mN}{(m-1)^2}}. \qquad (L19)$$

We can now substitute Equations L15, L16, L17 & L19 into Equation J14 to obtain an explicit expression for $\frac{dw}{dy}\Big|_{\tilde{x}=\tilde{x}^*,y=y^*,\tilde{z}=\tilde{z}^*}$. Setting $\frac{dw}{dy}\Big|_{\tilde{x}=\tilde{x}^*,y=y^*,\tilde{z}=\tilde{z}^*} > 0$ and simplifying, we obtain

$$\tilde{z}^* < \frac{(N-1)p(d-s)}{N(1-p)s}. \qquad (L20)$$

Signal-response (y^*) is favoured to increase if Equation L20 is satisfied. To interpret Equation L20, note that $s\left((1-p)\tilde{z}^* + \frac{N-1}{N}p\right)$ gives the cost to a plant of responding to a signal – specifically, it is the defence cost (s) weighted by the proportion of generations where this defence cost is paid $\left((1-p)\tilde{z}^* + \frac{N-1}{N}p\right)$. $\frac{N-1}{N}pd$ gives the benefit to a plant of responding to a signal – specifically, it is the benefit of a reduced predation cost (d), weighted by the proportion of generations where benefit can be accrued $\left(\frac{N-1}{N}p\right)$. Signal response is favoured to increase when it results in a greater benefit than cost to the responder (plant), leading to $\frac{N-1}{N}pd > s\left((1-p)\tilde{z} + \frac{N-1}{N}p\right)$, which with some rearrangement gives Equation L20.

Note that the right-hand side of Equation L20, which gives the threshold signal dishonesty under which signal-response is favoured, is almost the same as the right-hand side of Equation J20. The difference is that the right-hand side of Equation L20 is a factor $\frac{N-1}{N}$ greater than the right-hand side of Equation J20. The reason for this difference is that, in the previous model where plants are signallers (Model J), each plant has a generational probability $1/N$ of being 'chosen' to be a potential signaller rather than a potential signal receiver. In the present model where fungi are signallers, in generations where a predator is absent and the fungus signals dishonestly, all plants on the deme receive this signal (*i.e.*, there is no 'chosen' plant that does not receive the signal). This slightly increased probability of receiving a signal

in the present model, relative to Model J, is the reason why the $\frac{N-1}{N}$ factor features in Equation L20.

By substituting our explicit value for signal dishonesty ($\tilde{z}^* = 0$; Equation L14) into our condition for signal-response being favoured (Equation L20), we find that Equation L20 is satisfied (as long as $p > 0$ & $d > 0$). This means that signal-response is favoured, leading to

$$y^* = 1, \qquad (L21)$$

corresponding to maximal response of plants to fungal signals. The reason why plants are favoured to respond maximally to fungal signals in this model is that, at equilibrium, fungal signals are honest, meaning they reliably indicate that a predator is present (Equation L14). It is always in the plants' interests to respond to such signals and defend themselves against the impending predator attack.

Finally, we calculate $\frac{d\widetilde{w}}{d\tilde{x}}|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*}$ (mutating signal investment). We do so by expanding it using the chain rule

$$\frac{d\widetilde{w}}{d\tilde{x}} = \frac{\partial \widetilde{w}}{\partial \tilde{x}_{focal}} \frac{dx_{focal}}{d\tilde{x}} + \frac{\partial \widetilde{w}}{\partial \tilde{x}_{others}} \frac{d\tilde{x}_{others}}{d\tilde{x}}, \qquad (L22)$$

and evaluating the derivatives on the right-hand side at equilibrium (Taylor & Frank, 1996; Frank, 1998; Brown & Taylor, 2010). We trivially obtain

$$\frac{d\tilde{x}_{focal}}{d\tilde{x}}|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*} = 1. \qquad (L23)$$

We obtain the equilibrium-evaluated partial derivatives $\frac{\partial \widetilde{w}}{\partial \tilde{x}_{focal}}|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*}$ and $\frac{\partial \widetilde{w}}{\partial \tilde{x}_{others}}|_{\tilde{x}=\tilde{x}^*, y=y^*, \tilde{z}=\tilde{z}^*}$ respectively as

$$\frac{\partial \widetilde{w}}{\partial \widetilde{x}_{focal}} \Big|_{\widetilde{x}=\widetilde{x}^*, y=y^*, \widetilde{z}=\widetilde{z}^*}$$

$$= -(-N(p-1)z^*(bsy^* + c) - b(N-1)py^*(d-s) + cNp)$$
$$/\big(x^*(N(p-1)z^*(bsy^* + c) + b(N-1)py^*(d-s) - cNp)$$
$$+ N(b(-d)p + b + 1)\big) \tag{L24}$$

and

$$\frac{\partial \widetilde{w}}{\partial \widetilde{x}_{others}} \Big|_{\widetilde{x}=\widetilde{x}^*, y=y^*, \widetilde{z}=\widetilde{z}^*} = 0. \tag{L25}$$

Note that, because $\frac{\partial w}{\partial \widetilde{x}_{others}}\big|_{\widetilde{x}=\widetilde{x}^*, y=y^*, \widetilde{z}=\widetilde{z}^*} = 0$ (Equation L24), this gets rid of the $\frac{d\widetilde{x}_{others}}{d\widetilde{x}}\big|_{\widetilde{x}=\widetilde{x}^*, y=y^*, \widetilde{z}=\widetilde{z}^*}$ partial derivate in Equation L21, which means we don't need to evaluate $\frac{d\widetilde{x}_{others}}{d\widetilde{x}}\big|_{\widetilde{x}=\widetilde{x}^*, y=y^*, \widetilde{z}=\widetilde{z}^*}$ in order to evaluate the full derivative $\frac{d\widetilde{w}}{d\widetilde{z}}\big|_{\widetilde{x}=\widetilde{x}^*, y=y^*, \widetilde{z}=\widetilde{z}^*}$.

We can now substitute Equations L23, L24 & L25 into Equation L22 to obtain an explicit expression for $\frac{d\widetilde{w}}{d\widetilde{x}}\big|_{\widetilde{x}=\widetilde{x}^*, y=y^*, \widetilde{z}=\widetilde{z}^*}$. Setting $\frac{d\widetilde{w}}{d\widetilde{x}}\big|_{\widetilde{x}=\widetilde{x}^*, y=y^*, \widetilde{z}=\widetilde{z}^*} > 0$, substituting in our explicit values for y^* (Equation L21) and $\widetilde{z}^*$ (Equation L14), and simplifying, we obtain

$$c < \frac{N-1}{N} b(d-s), \tag{L26}$$

which is a condition for when fungi are favoured to signal the presence of a predator to the plants that they are connected to. Signalling is favoured when the benefit of signalling to the signaller (fungus), given by the right-hand side of Equation L26, is greater than the cost of signalling to the signaller (fungus), given by the left-hand side of Equation L26. The benefit of signalling to the signaller (fungus) is obtained as the increase in plant fecundity (fitness) as a consequence of responding to the fungal signal and upregulating predator defence (d–s), weighted by the fraction of the plant population who can respond to the fungal signal and therefore obtain this increase in fecundity $\left(\frac{N-1}{N}\right)$, weighted by the benefit to the fungus of having higher-quality trade partners (b). The cost of signalling to the signaller (fungus) is simply given by the cost of signal production (c).

We note that 'opening' this model, detaching relatedness from demographic parameters (m and N) so that it is a parameter in its own right, does not affect model outcomes. To appreciate this, note that calculating $\frac{dw}{dy}\big|_{\tilde{x}=\tilde{x}^*,y=y^*,\tilde{z}=\tilde{z}^*}$ with Equation L18 rather than Equation L19 (i.e., treating r^* as a parameter rather than a function of demographic parameters) does not lead to a different expression for $\frac{dw}{dy}\big|_{\tilde{x}=\tilde{x}^*,y=y^*,\tilde{z}=\tilde{z}^*}$. Relatedness does not feature in the expressions for $\frac{d\tilde{w}}{d\tilde{x}}\big|_{\tilde{x}=\tilde{x}^*,y=y^*,\tilde{z}=\tilde{z}^*}$ and $\frac{d\tilde{w}}{d\tilde{z}}\big|_{\tilde{x}=\tilde{x}^*,y=y^*,\tilde{z}=\tilde{z}^*}$. This means that the results of the model (Equations L14, L21 & L26) are unchanged – relatedness does not affect model outcomes in this lifecycle.

References

Brown, S.P. & Taylor, P.D. (2010) Joint evolution of multiple social traits: a kin selection analysis. *Proceedings of the Royal Society B: Biological Sciences*, **277**, 415–422.

Cooper, G.A., Levin, S.R., Wild, G. & West, S.A. (2018) Modeling relatedness and demography in social evolution. *Evolution Letters*, **2**, 260–271.

Diggle, S.P., Gardner, A., West, S.A. & Griffin, A.S. (2007) Evolutionary theory of bacterial quorum sensing: when is a signal not a signal? *Philosophical Transactions of the Royal Society B: Biological Sciences*, **362**, 1241–1249.

Foster, K.R. & Wenseleers, T. (2006) A general model for the evolution of mutualisms. *Journal of Evolutionary Biology*, **19**, 1283–1293.

Frank, S.A. (1994) Genetics of Mutualism: The Evolution of Altruism between Species. *Journal of Theoretical Biology*, **170**, 393–400.

Frank, S.A. (1998) *Foundations of Social Evolution*. Princeton University Press.

Grafen, A. & Archetti, M. (2008) Natural selection of altruism in inelastic viscous homogeneous populations. *Journal of Theoretical Biology*, **252**, 694–710.

Lehmann, L., Feldman, M.W. & Rousset, F. (2009) On the evolution of harming and recognition in finite panmictic and infinite structured populations. *Evolution*, **63**, 2896–2913.

Lehmann, L. & Keller, L. (2006) The evolution of cooperation and altruism – a general framework and a classification of models. *Journal of Evolutionary Biology*, **19**, 1365–1376.

Lehmann, L. & Rousset, F. (2010) How life history and demography promote or inhibit the evolution of helping behaviours. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **365**, 2599–2617.

Maynard Smith, J. & Harper, D. (2003) *Animal Signals*. Oxford University Press, Oxford, UK.

Padje, A. van't, Whiteside, M.D. & Kiers, E.T. (2016) Signals and cues in the evolution of plant–microbe communication. *Current Opinion in Plant Biology*, **32**, 47–52.

Pepper, J.W. (2000) Relatedness in Trait Group Models of Social Evolution. *Journal of Theoretical Biology*, **206**, 355–368.

Queller, D.C. (1994) Genetic relatedness in viscous populations. *Evolutionary Ecology*, **8**, 70–73.

Rousset, F. (2004) *Genetic Structure and Selection in Subdivided Populations (MPB-40)*. Princeton University Press.

Roze, D. & Rousset, F. (2008) Multilocus models in the infinite island model of population structure. *Theoretical Population Biology*, **73**, 529–542.

Scott, T.W. & Wild, G. (2023) How to make an inclusive-fitness model. *Proceedings of the Royal Society B: Biological Sciences*, **290**.

Taylor, P.D. (1992a) Altruism in viscous populations-an inclusive fitness model. *Evolutionary Ecology*, **6**, 352–356.

Taylor, P.D. (1992b) Inclusive fitness in a homogeneous environment. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, **249**, 299–302.

Taylor, P.D. & Frank, S.A. (1996) How to Make a Kin Selection Model. *Journal of Theoretical Biology*, **180**, 27–37.

Taylor, P.D., Wild, G. & Gardner, A. (2007) Direct fitness or inclusive fitness: how shall we model kin selection? *Journal of Evolutionary Biology*, **20**, 301–309.

West, S.A. & Gardner, A. (2010) Altruism, Spite, and Greenbeards. *Science*, **327**, 1341–1344.