Title: Shifts in vernalization and phenology at the rear edge hold insight into adaptation of

temperate plants to future milder winters

Authors: Antoine Perrier, Megan C. Turner, Laura F. Galloway

Department of Biology, University of Virginia, PO Box 400328, Charlottesville, VA 22904, USA.

Corresponding author contact details:

Antoine Perrier

Department of Biology,

University of Virginia,

PO Box 400328,

Charlottesville, VA 22904, USA

Phone: +1 434 243 0774

Email: cdt9qe@virginia.edu

Orcid ID: https://orcid.org/0000-0002-6447-5692

Type of article: Full paper

Number of words: Summary: 195; main text: 5852 (Introduction: 834, Material and Methods:

2267, Results: 931, Discussion: 1819)

Number of references: 51

Number of figures: 4 (4 in colour)

Number of tables: 2

Number of supporting information: 22, including 7 supporting figures, 11 supporting tables and

4 supporting methods

Summary

- Temperate plants often regulate reproduction through winter cues, such as vernalization, that may be reduced under climate change. Understanding adaptive potential to milder winters may be achieved by studying rear-edge populations, glacial relicts that persist in environments that have warmed since last glaciation.

- We studied how rear-edge populations have adapted to shorter winters and compared them to the rest of the range in the herb *Campanula americana*. Using citizen science, climate data and experimental climate manipulation, we characterize variation in vernalization requirements and reproductive phenology across the range and their potential climatic drivers.

- Rear-edge populations experienced little to no vernalization in nature. In climate manipulation experiments, these populations also had a reduced vernalization requirement, weaker response to changes in vernalization length, and flowered later compared to the rest of the range.

- Our results suggest shifts in phenology and its underlying regulation at the rear edge to compensate for unreliable vernalization cues. Thus, future milder winters may be less detrimental to these populations than more northern ones. Furthermore, our results showcase strong shifts in adaptation at the rear edges of temperate plants' ranges, highlighting their importance in studies on response and adaptation to predicted future climates.

Keywords: *Campanula americana, climate change, clines, flowering time, phenology, rear edge, reproductive cues, vernalization.*

Introduction

Organisms exposed to cyclical environmental changes often evolve mechanisms to sense these fluctuations and time developmental shifts to occur under favorable conditions (Preston & Sandve, 2013). In temperate plants, vernalization, the prolonged exposure to non-lethal winter cold (Chouard, 1960), serves as an important cue so that key life-cycle transitions will occur after winter, e.g. vegetative growth and reproduction (Amasino, 2005). However, relying on such cues may be detrimental under rapidly changing environments. This is of particular concern in the context of ongoing global warming, as it is expected to affect the strength and timing of seasonal temperatures experienced by natural and agricultural species (Willis *et al.*, 2008; Blackman, 2017). Species requiring vernalization will likely be negatively affected by climate warming as shorter and milder winters become more common (Luedeling *et al.*, 2011; Anderson, 2023). Such conditions are expected to lead to a reduction in reproduction (Padhye & Cameron, 2009; Liu *et al.*, 2012; Satake *et al.*, 2013) or shifts in reproductive phenology outside of optimal time-windows (Fitter & Fitter, 2002; Parmesan & Yohe, 2003; Faidiga *et al.*, 2023; Geissler *et al.*, 2023). It is therefore critical to understand response to seasonal cuing, and variation in this response, to determine possible consequences of expected warmer climates for temperate plant species.

Vernalization requirements and cue responses often vary within and between species in response to differences in winter condition (Andrés & Coupland, 2012; Blackman, 2017; Preston & Fjellheim, 2022). This suggests that while these mechanisms are crucial to complete the life cycle, they vary enough to allow persistence across heterogeneous environments. Studies have explored variation in cueing reproduction across environments by testing for clines in vernalization requirements and phenological response to temperature gradients (Blackman, 2017). This body of work provides evidence of a reduction in vernalization requirements with increases in temperature (Wesselingh *et al.*, 1994; Dijk *et al.*, 1997; Boudry *et al.*, 2002; Stinchcombe *et al.*, 2005; Jokela *et al.*, 2015). Reproductive phenology also varies over temperature gradients though a general pattern is less clear. For example, flowering is later with the later arrival of spring-like conditions at high latitudes than lower ones in *Arabidopsis thaliana* (Stinchcombe *et al.*, 2004; Lempe *et al.*, 2005). In contrast, earlier flowering occurs at higher latitudes in other species (e.g. Paccard et al. 2014; Vest and Sobel 2021). These studies reveal the potential for both the mechanism of environmental cuing as well as the pattern of reproductive phenology to evolve to track warming climates.

Populations at the warmer limits of temperate species ranges are particularly important for

understanding adaptation to shorter winters. This part of the range, colloquially known as the "rear

edge" (Hampe & Petit, 2005), often consists of relict populations that have persisted more or less

in place at least since the last glacial maximum (LGM, c. 23 000–19 000 years ago; Hughes et al.

2013). Long-term persistence at the warmer range limit over multiple glacial cycles and under

continuously warming climates after the LGM has likely resulted in adaptation to warmer climates.

With this history, the rear edge may provide insight into how species requiring vernalization adapt

to mild winters, and which adaptations may be best suited to future warming climates. However,

these populations are typically not sampled in studies investigating clines and therefore our

knowledge of phenology and vernalization requirements in populations from the warmest habitats

is limited (but see Vest and Sobel 2021).

Here we investigate whether rear-edge populations show patterns of differentiation

consistent with adaptation to milder winters compared to populations from elsewhere in the range

in the North American herb *Campanula americana*. This species requires vernalization, but

occupies a wide latitudinal and temperature gradient, with ancestral rear-edge populations

occurring in distinctively warmer climates in the southern parts of the range. To test for

differentiation between the rear edge and the rest of the range, we first characterized differences

in flowering phenology across latitudes using range-wide observations in natural populations

gathered from citizen science data, and then tested whether this variation reflects differences in

winter or growing season climate (Experiment 1). We also assessed genetic differences in

phenology by testing for variation in flowering among populations from across latitudes raised

under common greenhouse conditions (Experiment 2). We then evaluated plasticity in

reproductive phenology in response to a range of natural growing season cues by raising

populations sampled across latitudes in common gardens along a latitudinal gradient (Experiment

3). Finally, we tested differences in vernalization requirement and phenological plasticity in

response to these winter cues by raising populations under experimentally manipulated

vernalization length (Experiment 4). In total, these four experiments allow us to comprehensively

test for differences in phenology and its regulation across the range of this species. Results of these

experiment will allow us to understand not only the mechanisms that underly how rear-edge

populations have adapted to their distinct habitats, but will also shed light on how populations

across the range vary in their response to milder winter, and ultimately the mechanisms that may
evolve to allow persistence under future conditions.

Material and methods

Study system

Campanula americana is a monocarpic herb found in open forests, clearings and forests margins
across the eastern USA (Fig. 1a). Seeds germinate in spring or fall, vernalize over winter as rosettes
and then bolt, elongating into a reproductive stalk. Following several months of growth, flower
buds develop and the plants transition to flowering in mid-summer, with flowers opening
progressively over several weeks. Most populations require at least six weeks at $4° C$ for successful
vernalization that cues reproduction (Baskin & Baskin, 1984; Etterson & Galloway, 2002), but
southernmost populations may not always require vernalization to bolt (Kalisz and Wardle 1994).
This species occupies a large range, spanning ~15 ° latitude over ~1800km with a
difference of ~22 °C in minimum annual temperature (Fig. 1a; bio6, inferred from WorldClim 2.0,
Fick and Hijmans 2017). The species comprises three geographically distinct genetic clades
(Barnard-Kubow *et al.*, 2015). Here we focus on the largest clade, found west of the Appalachian
Mountains, that covers most of the range. We limited sampling to low elevations (<600m) to
reduce confounding climate gradients across elevation and latitude. This "Western" clade is
characterized by a history of persistence in glacial refugia in the southern part of the current range
during LGM, followed by postglacial range expansion (Barnard-Kubow *et al.*, 2015; Koski *et al.*,
2019; Prior *et al.*, 2020). We define the "rear edge" as the lower latitudinal third of the range
(below 35° N, Fig. 1a). This part of the range largely overlaps with putative glacial refugia
(Barnard-Kubow *et al.*, 2015). It also has a climate defined as subtropical, characterized by mild
winters and rare freezing events (American meteorological society,
https://glossary.ametsoc.org/wiki/Subtropics), distinct from the continental temperate climate in
more northern parts of the range.

Experiment 1: Variation in flowering phenology in natural populations
We first tested whether rear-edge populations in *C. americana* differ from the rest of the range
based on flowering phenology observed in natural populations across latitudes, and then explore
associations between phenology and both winter and growing season climates.

Inference of flowering phenology in natural populations

We inferred flowering phenology of natural populations from the publicly available repository iNaturalist (https://www.inaturalist.org. Accessed 23/02/2023). We extracted all 10,250 records of *Campanula americana* observed between 2018 and 2022 (earlier records did not cover the distribution well). This set was reduced to 335 observations following pruning to select for low elevation populations, reduce duplicates and oversampling close to cities, and reduce bias of fewer occurrences at the rear edge (Supporting method S1.1, Fig. S1.1). We inferred the day of first flower (recorded as day of the year) based on the photograph provided with each observation (Table S1.1, Fig. S1.2), resulting in a final dataset of 238 observations (Fig. 1a).

Climatic variables

For each observation, we obtained daily mean, maximum, and minimum temperature and daily precipitation for the year of observation and the year before from the PRISM database (https://prism.oregonstate.edu, accessed 01/09/2023) using the *prism* package (Edmund & Bell, 2015) in R (R Core team, 2024). We then generated 17 variables from this data that capture the climate at key life cycle stages, to test associations of phenology with climate (Table S1.2a).

Statistical analysis

We first explored how the day of first flower varied across latitudes. To do this, we compared three mixed-effect models with latitude as a continuous fixed effect and the year of observation as random effect. The first model tested for a linear relationship between the day of first flower and latitude, the second tested a second-degree quadratic relationship, and the third model tested a piecewise linear relationship with one breakpoint. The latter two allow clines with latitude to vary across the range. All analyses were performed in *R*. Model parametrizations are provided in Supporting method S1.2a. Model fits were compared using the corrected Akaike's information criterion (AICc, Sugiura 1978) using *MuMin* (Bartoń, 2023). From this we identified a breakpoint in the relationship between the day of first flower and latitude at ~35 °N (see Results), that we incorporated into the design of subsequent experiments.

We then assessed whether the observed variation in the day of first flower was associated with environmental variables. We first tested which climatic variable best discriminated between

the part of the range below or above the breakpoint in latitude by performing a discriminant
analysis of principal components (DAPC) using the *dapc* function in *adegent* (Jombart, 2008;
Jombart *et al.*, 2010). DAPC was performed on all 17 climate variables (transformed by PCA, 8
PC retained) with the grouping factor of whether observations were located at latitudes above
(group 1) or below (group 2) the predicted breakpoint. We then evaluated whether identified
climate variables changed across the range in mixed effect models that included the latitude of
each observation as a continuous fixed effect, range position of each observation relative to the
estimated breakpoint as a categorical effect, and their interaction. The year of observation was
included as a random effect (model in Supporting method S1.2b). We also tested the effect of
latitude on the day of first flower using the same model structure.

Experiment 2: Variation in reproductive phenology in a common environment
We tested for genetic differences in flowering phenology between *C. americana*'s rear edge and
the rest of the range by raising populations from across latitudes under a common environment in
a greenhouse experiment.

Experimental design
We raised two cohorts of *C. americana* populations that cover a similar latitudinal breadth
below (~4.4° N) and above (~6.6° N) the breakpoint identified in Exp. 1. For the first cohort, we
collected seeds in 23 natural populations (Fig. 2a, Table S2.1) in late summer of 2020 and 2021
and stored them by maternal plant at 4 °C under dark and dry conditions. For the second cohort,
we used greenhouse-reared seeds from 25 populations (Supporting method S2.1), 20 from the first
cohort and five additional populations to yield better sampling (Fig. 2a). Greenhouse produced
seeds have less variation due to maternal effects and collection date. We raised 544 seedlings in
the first cohort (Fall 2021), and 535 in the second cohort (Fall 2022) in similar conditions
(Supporting method S2.1). Conditions were designed to mimic the natural growth cycle of the
species, starting with germination and vegetative growth in growth chambers simulating fall,
followed by vernalization in a cold room simulating winter, and then bolting and reproduction in
a greenhouse with conditions simulating summer.

Recording of traits and statistical analysis

We recorded the date of bolting (presence of a stem) twice a week for the first cohort and weekly for the second. We recorded the date of first flower three times a week for both cohorts. Most plants that bolted also flowered (>98%).

Based on this data, we generated four traits related to reproduction initiation and phenology: *flowering success* (binary, production of at least 1 flower), *time to flowering* (days from vernalization to flowering) and its components *time to bolting* and *time between bolting and flowering* (Table S2.2a). Each was analyzed at the individual level in mixed effect models in *R*, with population *latitude* as a continuous fixed effect, *cohort* a categorical effect, and their interaction. Population and seed family nested within population were random effects (models in Supporting method S2.2). Similar to Exp. 1, we initially tested whether latitude was best described as a linear relationship or a second-degree quadratic relationship for each trait. This was done by comparing model performance based on AICc. We report results only for the best performing model (all results Table S2.3). For this model, we performed a post-hoc test to estimate the slope of the relationship between latitude and reproductive traits within each cohort, cohort effects and the difference in the slope between cohorts using the package *emmeans* (Lenth, 2019).

Experiment 3: Variation in reproductive phenology in a transplant experiment

We tested for differences in phenology in response to spring and summer conditions between the rear edge and the rest of the range by raising populations from across the range in common gardens established on a latitudinal gradient.

Experimental design

We selected 25 populations with similar distribution as Exp. 2 (Fig 3a, Table S2.1), and approx. 15 seed families per population (Table S2.1), to be raised in five common gardens. We used greenhouse-reared seeds for 20 of the populations and field-collected seeds for five. For each garden, we sowed two seeds in each of two pots per seed family (three seeds for field-collected seeds, one if few seeds), thinning to one plant per pot after germination. This gave 2 individuals/family x 15 families/population x 25 populations = 750 individuals per garden, 3750 total. Seeds were planted in fall 2022, germinated for six weeks in growth chambers and vernalized for seven weeks in a cold room (Supporting method S2.1).

Seedlings were acclimated to outdoor conditions for one week (Charlottesville, VA, USA), and then transplanted to one of five common garden sites (CG, Fig. 3a, Tables S2.4): Lexington KY (CG1, 38.1° N), Blaine TN (CG2, 36.2° N), Clemson SC (CG3, 34.7° N), Akron AL (CG4, 32.9° N), and Tallahassee FL (CG5, 30.5° N). These sites span an 850 km latitudinal gradient in roughly 2° steps and reflect the temperature gradient within the sampled range. Transplant for each site occurred when mean daily temperature was consistently above 10°C in spring, estimated using PRISM daily mean temperature 2018 to 2022. Sowing was timed for each site such that vernalization ended one week before the projected transplant date.

Garden locations were chosen to mimic the natural habitat of the species. In each site, individuals were transplanted in 30 spatial blocks of 25 individuals, with one individual per population per block. Sites were raked to reduce initial competition. Individuals were then transplanted into the exposed ground with the substrate they were raised in and watered once. Each site was surrounded by a 2m high fence to exclude large herbivores. No further intervention was performed. One data logger (iButton®, Maxim Integrated Products, Inc) was placed in each site in the shade, 1.5m above ground, to monitor air temperature every hour for the length of the experiment. This data revealed that plants experienced additional days of vernalization after transplant (Table S2.4a), but less so in the southernmost garden. The Experiment was stopped 153 to 170 days after transplant (Table S2.4b), when most plants had reached fruit maturation but seeds were not yet dispersing. Unfortunately, plants in CG4 experienced very low survival (<10%), so this site was removed from analysis.

Recording of trait and statistical analysis

Data was collected three times, about a month apart, in the three northern sites (GC1-GC3), and only two times for the southern site (GC5). In each round of data collection, we recorded survival, transition to bolting and to flowering (here production of at least one bud) and inferred the flowering date as in Exp. 1 (Table S2.5).

Based on this data, we generated three binary traits at the plant level: *transition to bolting* and *transition from bolting to flowering* estimated at each time point, *flowering success* based on the first time point to provide a "snapshot" of flowering phenology. We estimated *day of first flower* (day of the year), as well as *time to flowering since transplant* (days from transplant) (Table S2.2b). The effects of latitude (of population origin), common garden (categorical, four sites) and

263   their interaction on each of these traits was analyzed in mixed effect models (Supporting method

264   S2.2). Preliminarily analysis tested whether latitude was best described by a linear or a quadratic

265   relationship, and results are only reported for the best model (Table S2.3). If models performed

266   equally well, the quadratic one was retained for ease of comparison among traits. Post-hoc tests

267   were performed for the best models.

269   Experiment 4: Variation in vernalization requirements and response across the range

We tested for differentiation in vernalization requirement and in phenological response to

vernalization length between the rear edge and the rest of the range by subjecting populations from

across latitudes to experimentally manipulated vernalization length.

Experimental design

We selected 12 populations with greenhouse-reared seeds, representing the same latitudinal range

as Exp.2 (Fig. 4a, Table S2.1). We selected approx. 15 families per population (Table S2.1), and

sowed two seeds in each of nine pots per seed family, resulting in 45 individuals per population

after thinning to one seedling per pot. Pots were then randomly assigned to one of three

vernalization treatments (see below), resulting in 3 individuals/family x 15 families/population x

12 populations = 540 individuals per treatment, 1620 total. Seedlings were raised from germination

in fall 2022 to flowering in 2023 (Supporting method S2.1), but with the difference that treatments

differed in vernalization length, lasting either six, four or two weeks. Sowing was timed such that

all three treatments finished vernalization at the same time (Fig. S2.1).

Recording of trait and statistical analysis

We recorded survival, bolting and flowering three times a week during the greenhouse phase.

Bolting was recorded for 52 days after vernalization and flowering for 141. We then estimated

flowering success and its two components *transition to bolting* and *transition from bolting to*

flowering, as well as *time to flowering* and its two components *time to bolting*, and *time between*

bolting and flowering (Table S2.2a). Similar to Exp. 2, the effects of latitude, vernalization

treatment (categorical with three levels) and their interaction on each of these traits was analyzed

in mixed effect models (Supporting method S2.2). For the three phenology traits, the two-week

treatment was excluded as few individuals bolted. Preliminarily analysis tested whether latitude

was best described by a linear or a quadratic relationship (Table S2.3), results are reported for the
best model and post-hoc tests were performed.

Results

Experiment 1: Variation in flowering phenology in natural populations

In natural populations, the relationship between the day of first flower and latitude was best
described as a quadratic or piecewise relationship (Table S1.3, Fig S1.3), with both models
performing equally well (ΔAICc < 2). The earliest flowering was at 36.64° N and 34.98° N
respectively, with flowering at increasingly later dates towards lower and higher latitudes.
Subsequent analyzes were parameterized with the breakpoint from the piecewise model because it
matched best our *a priori* delimitation of the rear edge (~35 °N), though use of the minimum from
the quadratic model yielded similar results (not shown). Climates of populations north and south
of this breakpoint were best discriminated by the length of vernalization as well as the time
between the start of the growing season and flowering (Table S1.2b). The other climate variables
had negligeable contributions.

The day of first flower and the two climate variables (length of vernalization, time between
the start of the growing season and flowering) showed strong clines across latitudes, with the slope
depending on their position relative to the breakpoint (L*BP, Table 1). Flowering was earliest at
the breakpoint, with much later flowering per unit of latitude toward low latitudes (26 days over
~4.5 °) than towards high latitudes (18 days over ~9.9 °, Table 1, Fig. 1b). The length of
vernalization increased with latitude (Table 1, Fig. 1c), but days of vernalization accumulated more
rapidly north of the breakpoint than to the south, where vernalization was limited. The time
between the start of the growing season and flowering was substantially shorter and relatively
insensitive to latitude to the north (65 to 51 days predicted), whereas it increased sharply to the
south of the breakpoint (62 to 130 days predicted; Table 1, Fig. 1d).

Experiment 2: Variation in reproductive phenology in a common environment
Rear-edge populations showed differentiation in reproductive phenology from the rest of the range
when raised in a common controlled environment. Populations from low latitudes had greater
flowering success but flowered later relative to those in the rest of the range (Table 2, Fig. 2, Table
S2.6). Populations from the highest latitudes also flowered later than central populations, but this

was weaker than in low latitude populations. Later flowering was due to a delay in time to bolting in northern populations, but a longer developmental period between bolting and the opening of the first flower in southern populations (Table 2, Table S2.6, Fig. S2.2). Patterns across latitudes were consistent between cohorts, though a greater proportion of plants flowered in the second cohort.

Experiment 3: Variation in reproductive phenology in a transplant experiment

Rear-edge populations showed strong differentiation from the rest of the range in the common gardens, indicated by delayed reproductive phenology relative to those from more central sites. Southern populations experienced smaller flowering success (except CG5), due to a reduction in both transitions to bolting and from bolting to flowering (Table 2, Fig. 3b, Tables S2.7, Fig S2.3a, b). These populations also flowered later, both in day of first flower and time to flowering since transplanting (Table 2, Fig. 3c, Tables S2.7, Fig S2.3c). Populations at high latitudes also had smaller flowering success and a moderate delay in phenology compared to central populations in all gardens, but differences were weaker than in low latitude populations (except CG5).

Patterns of trait variation among southern populations were generally consistent across garden sites despite differences in spring and summer climate (Fig. 3). In contrast, northern populations exhibited plasticity in flowering phenology across gardens. For example, in the most southern garden (CG5), northern populations showed a stronger delay in phenology relative to central populations than in any of the other sites (Table 2, Fig. 3, Tables S2.7, Fig S2.3).

Variation in phenology across gardens was generally linked to transplant time. The phenology "snapshot", flowering success, was significantly smaller in northern sites compared to southern sites (except for CG1 vs CG2; Fig. 3b, Table S2.7b), indicating a delay in flowering in northern sites. In line, the day of first flower generally occurred later in northern sites compared to southern sites (except for CG1 vs CG2; Table 2, Table S2.7b, Fig. S2.3c). However, these differences were almost eliminated when flowering was measured relative to transplant date (Fig. 3c, Table S2.7b).

Experiment 4: Variation in vernalization requirements and response across the range

Vernalization requirements of rear-edge populations were strongly differentiated from the rest of the range, characterized by a reduced need for vernalization and smaller plastic response to variation in vernalization length. In keeping with this pattern, there was a decrease in flowering

success in populations from higher latitudes that was exacerbated when vernalization was reduced (Table 2, Fig. 4b, Tables S2.8). In northern populations, a reduction in vernalization from six weeks to two weeks yielded a large reduction in flowering success. In contrast, southernmost populations had high flowering success regardless of vernalization treatment. Most of this variation was due to variation in transition to bolting among populations (Table 2, Tables S2.8, Fig. S2.4a).

Differentiation in response to vernalization was also observed for phenology, with rear-edge populations again showing smaller plastic response to variation in vernalization. In the six-week treatment, populations from low latitudes flowered later compared to more central populations (Fig. 4c). Reducing vernalization led to a general delay in time to flowering except for in the southernmost populations which changed little (Fig. 4c, Tables S2.8). Differences in time to flowering across treatments were mainly due to changes in the time between bolting and flowering, rather than changes in time to bolting (Table 2, Tables S2.8, Fig. S2.4).

Discussion

Understanding how relict rear-edge populations at the warmer range limits of temperate species differ from younger populations at higher latitudes can help us predict how populations across the range may respond and adapt to future climates. We found that in the North American herb *Campanula americana,* rear-edge populations occurring in the lower latitudinal part of the range have evolved to rely less on winter cues to regulate phenology compared to the rest of the range. This suggests that critical life cycle mechanisms, such as the regulation of phenology, may evolve rapidly to adapt to changing environments. In addition, contrary to classic expectations, we found the warmer range limit may be less affected by climate change's mild winters compared to higher latitude regions of the range. Finally, our results suggest that northern populations will suffer losses in reproduction with a reduction in winter cues although, over time, they may adapt to these changes similarly to the rear edge.

Natural rear-edge populations differ in phenology from the rest of the range

In natural *C. americana* populations, timing of flowering had opposing clines across latitudes (Exp. 1, Fig. 1b). The breakpoint of the clines was ~35 °N, coinciding with our *a priori* definition of the rear-edge (below) relative to the rest of the range (above). The day of first flower was the

earliest at the breakpoint, becoming progressively later towards higher and lower latitudes.
Previous work has found latitudinal clines in flowering are most often continuous in temperate
plants (e.g. Stinchcombe et al. 2004). These clines are thought to reflect continuous differences in
cues such as photoperiod or vernalization used to signal the onset of favorable growing seasons
(Preston & Sandve, 2013; Blackman, 2017; Preston & Fjellheim, 2022). However, rear-edge
populations have typically been neglected in such studies, so we don't know how common the
opposing clines seen here are. Opposing clines suggest a divergence in the response to seasonal
cues, or a change in the cues used to regulate phenology between the rear edge and the rest of the
range.
396
Most of the range shows adaption to winter conditions typical for temperate plants
We found that differentiation in the day of first flower between rear-edge populations and the rest
of the range was explained by differences in the length of vernalization (Exp. 1), suggesting
vernalization may serve as environmental cue for flowering. In nature, populations occurring
above 35 °N experience increasing winter length towards higher latitudes (Fig. 1c). These longer
winters were associated with an increase in vernalization required for reproduction in both
greenhouse experiments (Exp. 2, 4). Fewer plants flowered from high-latitude populations exposed
to short vernalization periods, and this response was strongest in the northernmost populations
which had a complete loss in reproduction under limited vernalization. A longer vernalization
requirement in high latitudes is common for temperate plant taxa (Wesselingh *et al.*, 1994; Boudry
et al., 2002; Jokela *et al.*, 2015), and may reflect an adaptation to prevent premature flowering in
areas with long winters and detrimental early spring frost events (Inouye, 2000).
Differentiation in the day of first flower across latitudes was also explained by the time
between flowering and the start of the growing season. Above 35 °N, flowering in nature occurred
somewhat earlier relative to the onset of the growing season with increasing latitudes (ca. 14 days
earlier across ca. 10° of latitude to the north, Exp. 1, Fig. 1d). This is also common among
temperate plants (e.g. Paccard et al. 2014; Vest and Sobel 2021), and is thought to be associated
with adaptation to complete reproduction under shorter growing seasons (Griffith & Watson,
2005). A similar shift in phenology was reported along an elevational gradient in *C. americana*,
where high elevation populations with shorter growing seasons had accelerated reproduction
compared to low elevation populations (Haggerty & Galloway, 2011).

In the manipulative experiments, populations from locations north of the rear edge
exhibited plasticity in flowering phenology in response to vernalization length. While generally
flowering slightly later than central populations, they flowered even later when vernalization was
short (Exp. 4, Fig. 4c) or when raised in southern climates (Exp. 3, Fig. 3c), with northernmost
populations exhibiting the strongest delay. This plasticity of later flowering under shorter
vernalization may simply be a passive byproduct of inadequate cuing. Specifically, limited
vernalization may partially cue the onset of reproduction, therefore resulting in a longer transition
to reproduction. The larger plasticity in timing of flowering in northern populations may be due to
greater vernalization requirements, increasing the mismatch between required and experimental
vernalization length. Indeed, the shortfall for vernalization for northern populations in the
manipulative experiments may have masked earlier flowering relative to the start of the growing
season as observed in nature. Alternatively, greater plasticity in the northernmost populations
could be that this plasticity has evolved to modulate reproductive phenology in response to thaws
in late winter, extending the period prior to flowering in short winters and thereby reducing
negative effects of these "false springs."

Winter cue requirements and responses are shifted at the rear-edge where winters are milder
Rear-edge populations showed distinct patterns of reproductive phenology compared to the rest of
the range, suggesting differences in adaptation in response to milder, shorter winters. In nature,
rear-edge populations experience less vernalization, with some not experiencing any vernalization.
Southern *C. americana* populations reflect this climate difference by having reduced vernalization
requirements, including southernmost populations which achieve a high percentage of flowering
(>90%) even under very limited vernalization (Exp. 4, Fig. 4b; Kalisz & Wardle, 1994). Reduced
vernalization requirements at the rear edge likely evolved in response to local shorter, warmer
winters where vernalization was an unreliable cue. Such patterns have been found in other plants,
in some cases leading up to a complete loss of a vernalization requirement (Wesselingh *et al.*,
1994; Dijk *et al.*, 1997; Boudry *et al.*, 2002; Stinchcombe *et al.*, 2005; Jokela *et al.*, 2015).
Flowering in natural rear-edge populations was later relative to the onset of the growing
season towards lower latitudes, with an almost ten-fold steeper cline in flowering time relative to
the rest of the range. Flowering was also much later in rear-edge populations under common
greenhouse conditions (Exp. 2, Fig. 2c), as well as in common gardens (Exp. 3, Fig. 3c). Such a

dramatic difference in phenology between populations along temperature gradients has rarely been

documented in temperate plants. The exception, wild beets, similar to *C. americana* has later

flowering in southern populations and variation in vernalization along a latitudinal gradient (Dijk

et al., 1997). While it is possible that other environmental factors such as day length cue

differences in phenology (Blackman, 2017), this seems unlikely for rear-edge populations in our

study. Phenology in southernmost populations was not affected by variation in winter length (Exp.

4, Fig. 3c), growing season climate (Exp. 3, Fig. 3c), or planting date (Exp. 3, Fig. S2.3c),

suggesting that the time to flowering is strongly genetically determined in these populations rather

than relying on an alternative environmental cue to vernalization.

Together, our results suggest a dramatic shift in the regulation of flowering phenology at

the rear edge, with less reliance on vernalization to initiate and time flowering. Instead, phenology

in these populations seems to be regulated by an environmentally insensitive delay in flowering,

contrasting with the plasticity in flowering time observed in northern populations. This evolved

delay may represent an adaptation to compensate for the extremely early start of the growing

season, thus allowing reproduction to track the warmest period of the year. A previous artificial

selection study revealed trade-offs between early flowering and reproductive fitness in *C.*

americana (Burgess *et al.*, 2007), supporting an evolutionary advantage to delay flowering when

the growing season length does not strongly limit reproductive output.

The rear edge will differ in response to climate change compared to the rest of the range

Our study provides insight into how temperate plants may vary in their response to climate change

across their range. Warming climates are often predicted to result in range shifts towards higher

latitudes and altitudes through colonization at the colder edge and extinctions at the warmer edge

(Thomas *et al.*, 2004; Hampe & Petit, 2005; Lenoir & Svenning, 2013). In temperate plants that

rely on vernalization to cue reproduction, one may expect that populations at the warmer range

limits will be especially sensitive to climate change as warming winters may not meet vernalization

requirements, thus preventing the transition to reproduction (*e.g.* Padhye and Cameron 2009;

Satake et al. 2013). However, in contrast to this prediction, we found that rear-edge populations of

C. americana are likely to be the least affected by a reduction in winter length. Rather, populations

at mid- and high latitudes may be at risk because their sensitivity in the initiation and timing of

reproduction in response to winter length may lead to a substantial fitness loss under milder

winters. Reduced sensitivity at the rear edge may be common across temperate plants. Lower latitude populations often have a reduced vernalization requirement (Wesselingh *et al.*, 1994; Dijk *et al.*, 1997; Boudry *et al.*, 2002; Stinchcombe *et al.*, 2005; Jokela *et al.*, 2015) They also show strong local adaptation (Bontrager *et al.*, 2021), likely facilitated by a history of long term exposure to increasingly warmer climates. This result contributes to a growing body of literature questioning the expected population decline under climate change at the rear edge (Vilà-Cabrera *et al.*, 2019).

Postglacial colonization history drives divergence in adaptation to changing climates

Our study highlights the importance of studying the effects of warming climates in a phylogeographic context, as the divergence in adaptation between northern and southern parts of the range may also reflect different histories of exposure to climate change. The ~35°N breakpoint in the day of first flower coincides with the transition between populations persisting in former glacial refugia in southern parts of the range, *i.e.* the rear edge, and more northern populations that have established after last glacial maximum (Barnard-Kubow et al. 2015; Koski et al. 2019; Prior et al. 2020). Post-glacial colonization in temperate plants often results from populations establishing in newly suitable habitats, tracking the gradual retreat of ice sheets as Earths' global temperatures rose during the Holocene (Hewitt, 2000, 2004). In this context, *C. americana* populations at higher latitudes may have retained flowering patterns closer to the ancestral state, while rear-edge populations may have adapted to climates that have warmed over the last millennia such that they are now subtropical. Reduced vernalization requirements and shifts in phenology at the rear edge may thus be a relatively recent, post-glacial adaptation, diverging from an ancestral state of relying on winter cues to regulate phenology. Despite historical and contemporary climate change progressing at very different temporal scales, the evolution of reduced vernalization requirements since the LGM suggests that evolution of lower vernalization requirements in response to warming climate may be possible in populations at mid- and high latitude, and may be facilitated by genetic variation occurring at the rear-edge.

Acknowledgements

This work was supported by the Swiss national Foundation (P2BSP3_195363) and the University of Virginia College of Arts and Sciences. We are grateful to David Westneat (University of Kentucky Ecological Research and Education Center Field Station, Lexington, KY), Gordon Burghardt (Blaine, TN), Trevor Stamey (Clemson University Experimental Forest, Clemson, SC), Jayne Lampley and John Walton (University of Alabama Tanglewood Station, Akron, AL), Theresa Jepsen (Florida state University Mission Road Research Facility, Tallahassee, FL) for logistical support in establishing field sites. For their help in raising of plants, collecting data, performing crosses and setting up the common garden experiments we thank A. Burricks, C. Claussen, S. Cox, L. Elhady, M. Gower-Fici, S. Kelly, O. Keenan, A. López, K. Lamb, H. Makowski, M. Marcich & E. Scott. We are also grateful to D. Brown, J. Collins, A. Diamond, L. Elliott, E. Galloway, F. Griffith, I. Guenther, J. Hansen, H. Horne, J. Kees, M. Kohout, R. Laporte, D. Reed & B. Sutherland for seed collection in natural populations. Collection permits were provided by the Florida Department of Environmental Protection, the Tennessee Department of Environment and Conservation, and Missouri Department of Conservation.

Conflict of Interest Statement

The authors declare that there is no conflict of interest.

Author Contributions

All authors contributed to the study design. AP collected seeds in the field and performed crosses, AP and MT raised plants and analyzed the data, AP wrote the manuscript with input from LG.

Data availability statement

iNaturalist observations, climate data and phenotypic data for Experiments 2, 3 and 4 will be stored on Dryad and made publicly available after initial acceptance.

References

Amasino RM. **2005**. Vernalization and flowering time. *Current Opinion in Biotechnology* **16**: 154–158.

Anderson JT. **2023**. The consequences of winter climate change for plant performance. *American Journal of Botany* **110**: e16252.

Andrés F, Coupland G. **2012**. The genetic basis of flowering responses to seasonal cues. *Nature Reviews. Genetics* **13**: 627–639.

Barnard-Kubow KB, Debban CL, Galloway LF. **2015**. Multiple glacial refugia lead to genetic structuring and the potential for reproductive isolation in a herbaceous plant. *American Journal of Botany* **102**: 1842–1853.

Bartoń K. **2023**. MuMIn: Multi-Model Inference. URL https://CRAN.R-project.org/package=MuMIn.

Baskin JM, Baskin CC. **1984**. The ecological life cycle of *Campanula americana* in Northcentral Kentucky. *Bulletin of the Torrey Botanical Club* **111**: 329–337.

Blackman BK. **2017**. Changing responses to changing seasons: Natural variation in the plasticity of flowering time. *Plant Physiology* **173**: 16–26.

Bontrager M, Usui T, Lee-Yaw JA, Anstett DN, Branch HA, Hargreaves AL, Muir CD, Angert AL. **2021**. Adaptation across geographic ranges is consistent with strong selection in marginal climates and legacies of range expansion. *Evolution* **75**: 1316–1333.

Boudry P, Mccombie H, Van Dijk H. **2002**. Vernalization requirement of wild beet *Beta vulgaris* ssp. *maritima*: among population variation and its adaptive significance. *Journal of Ecology* **90**: 693–703.

Burgess KS, Etterson JR, Galloway LF. **2007**. Artificial selection shifts flowering phenology and other correlated traits in an autotetraploid herb. *Heredity* **99**: 641–648.

Chouard P. **1960**. Vernalization and its relations to dormancy. *Annual Review of Plant Physiology* **11**: 191–238.

Dijk HV, Boudry P, McCombre H, Vernet P. **1997**. Flowering time in wild beet (*Beta vulgaris* ssp. *maritima*) along a latitudinal cline. *Acta Oecologica* **18**: 47–60.

Edmund H, Bell K. **2015**. prism: Download data from the Oregon prism project. URL https://CRAN.R-project.org/package=prism.

Etterson JR, Galloway LF. **2002**. The influence of light on paternal plants in *Campanula americana* (Campanulaceae): pollen characteristics and offspring traits. *American Journal of Botany* **89**: 1899–1906.

Faidiga AS, Oliver MG, Budke JM, Kalisz S. **2023**. Shifts in flowering phenology in response to spring temperatures in eastern Tennessee. *American Journal of Botany* **110**: e16203.

Fick SE, Hijmans RJ. **2017**. WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology* **37**: 4302–4315.

Fitter AH, Fitter RSR. **2002**. Rapid changes in flowering time in British plants. *Science* **296**: 1689–1691.

Geissler C, Davidson A, Niesenbaum RA. **2023**. The influence of climate warming on flowering phenology in relation to historical annual and seasonal temperatures and plant functional traits. *PeerJ* **11**: e15188.

Griffith TM, Watson MA. **2005**. Stress avoidance in a common annual: reproductive timing is important for local adaptation and geographic distribution. *Journal of Evolutionary Biology* **18**: 1601–1612.

Haggerty BP, Galloway LF. **2011**. Response of individual components of reproductive phenology to growing season length in a monocarpic herb. *Journal of Ecology* **99**: 242–253.

Hampe A, Petit RJ. **2005**. Conserving biodiversity under climate change: the rear edge matters. *Ecology Letters* **8**: 461–467.

Hewitt G. **2000**. The genetic legacy of the Quaternary ice ages. *Nature* **405**: 907–913.

Hewitt GM. **2004**. Genetic consequences of climatic oscillations in the Quaternary (KJ Willis, KD Bennett, and D Walker, Eds.). *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* **359**: 183–195.

Hughes PD, Gibbard PL, Ehlers J. **2013**. Timing of glaciation during the last glacial cycle: evaluating the concept of a global 'Last Glacial Maximum' (LGM). *Earth-Science Reviews* **125**: 171–198.

Inouye D w. 2000. The ecological and evolutionary significance of frost in the context of climate change. *Ecology Letters* **3**: 457–463.

Jokela V, Trevaskis B, Seppänen MM. **2015**. Genetic variation in the flowering and yield formation of timothy (*Phleum pratense* L.) accessions after different photoperiod and vernalization treatments. *Frontiers in Plant Science* **6**.

Jombart T. **2008**. adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics* **24**: 1403–1405.

Jombart T, Devillard S, Balloux F. **2010**. Discriminant analysis of principal components: a new method for the analysis of genetically structured populations. *BMC Genetics* **11**: 94.

Kalisz S, Wardle GM. **1994**. Life history variation in *Campanula americana* (Campanulaceae): population differentiation. *American Journal of Botany* **81**: 521–527.

Koski MH, Layman NC, Prior CJ, Busch JW, Galloway LF. **2019**. Selfing ability and drift
load evolve with range expansion. *Evolution Letters* **3**: 500–512.

Lempe J, Balasubramanian S, Sureshkumar S, Singh A, Schmid M, Weigel D. **2005**. Diversity
of flowering responses in wild *Arabidopsis thaliana* strains (J Doebley, Ed.). *PLoS Genetics* **1**: e6.

Lenoir J, Svenning J-C. **2013**. Latitudinal and elevational range shifts under contemporary
climate change. In: Levin SA, ed. Encyclopedia of Biodiversity (Second Edition). Waltham:
Academic Press, 599–611.

Lenth R. **2019**. emmeans: estimated marginal means, aka least-squares means. URL
https://CRAN.R-project.org/package=emmeans.

Liu Y, Mu J, Niklas KJ, Li G, Sun S. **2012**. Global warming reduces plant reproductive output
for temperate multi-inflorescence species on the Tibetan plateau. *New Phytologist* **195**: 427–436.

Luedeling E, Girvetz EH, Semenov MA, Brown PH. **2011**. Climate change affects winter chill
for temperate fruit and nut trees. *PLOS ONE* **6**: e20155.

Paccard A, Fruleux A, Willi Y. **2014**. Latitudinal trait variation and responses to drought in
Arabidopsis lyrata. *Oecologia* **175**: 577–587.

Padhye SR, Cameron AC. **2009**. Vernalization responses of *Campanula* 'Birch Hybrid'. *Journal*
of the American Society for Horticultural Science **134**: 497–504.

Parmesan C, Yohe G. **2003**. A globally coherent fingerprint of climate change impacts across
natural systems. *Nature* **421**: 37–42.

Preston JC, Fjellheim S. **2022**. Flowering time runs hot and cold. *Plant Physiology* **190**: 5–18.

Preston JC, Sandve SR. **2013**. Adaptation to seasonality and the winter freeze. *Frontiers in Plant*
Science **4**.

Prior CJ, Layman NC, Koski MH, Galloway LF, Busch JW. **2020**. Westward range expansion
from middle latitudes explains the Mississippi River discontinuity in a forest herb of eastern North
America. *Molecular Ecology* **29**: 4473–4486.

R Core Team. 2024. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.URL https://www.R-project.org/.

Satake A, Kawagoe T, Saburi Y, Chiba Y, Sakurai G, Kudoh H. **2013**. Forecasting flowering
phenology under climate warming by modelling the regulatory dynamics of flowering-time genes.
Nature Communications **4**: 2303.

Stinchcombe JR, Caicedo AL, Hopkins R, Mays C, Boyd EW, Purugganan MD, Schmitt J.
2005. Vernalization sensitivity in *Arabidopsis thaliana* (Brassicaceae): the effects of latitude and
FLC variation. *American Journal of Botany* **92**: 1701–1707.

Stinchcombe JR, Weinig C, Ungerer M, Olsen KM, Mays C, Halldorsdottir SS, Purugganan MD, Schmitt J. 2004. A latitudinal cline in flowering time in *Arabidopsis thaliana* modulated by the flowering time gene *FRIGIDA*. *Proceedings of the National Academy of Sciences* **101**: 4712–4717.

Sugiura N. **1978**. Further analysts of the data by akaike' s information criterion and the finite corrections: Further analysts of the data by akaike' s. *Communications in Statistics - Theory and Methods* **7**: 13–26.

Thomas CD, Cameron A, Green RE, Bakkenes M, Beaumont LJ, Collingham YC, Erasmus BFN, de Siqueira MF, Grainger A, Hannah L, *et al.* **2004**. Extinction risk from climate change. *Nature* **427**: 145–148.

Vest K, Sobel JM. **2021**. Variation in seasonal timing traits and life history along a latitudinal transect in *Mimulus ringens*. *Journal of Evolutionary Biology* **34**: 1803–1816.

Vilà-Cabrera A, Premoli AC, Jump AS. **2019**. Refining predictions of population decline at species' rear edges. *Global Change Biology* **25**: 1549–1560.

Wesselingh RA, Klinkhamer PGL, Jong TJ, Schlatmann EGM. **1994**. A latitudinal cline in vernalization requirement in *Cirsium vulgare*. *Ecography* **17**: 272–277.

Willis CG, Ruhfel B, Primack RB, Miller-Rushing AJ, Davis CC. **2008**. Phylogenetic patterns of species loss in Thoreau's woods are driven by climate change. *Proceedings of the National Academy of Sciences of the United States of America* **105**: 17029–17033.

Supporting information

Supporting material 1: Flowering phenology in natural populations (Experiment 1)

Fig. S1.1: Pruning of iNaturalist observations used in Experiment 1

Fig. S1.2: Example of flowering stages attributed to each observation

Fig. S1.3: Variation in flowering time across latitudes for the two best models

Table S1.1 Estimation of first flower date in natural populations based on flowering stage

Table S1.2a: Description of the 17 climatic variables used in Experiment 1

Table S1.2b: Contribution to the discriminant function and range of the 17 climatic variables used in Experiment 1

Table S1.3: Comparison of three models describing variation in the day of first flower across latitude

Supporting method S1.1: Selection of observations for estimation of day of first flower

Supporting method S1.2: Parametrization of analyses

Supporting material 2: Flowering phenology under experimentally manipulated conditions (Experiments 2, 3, 4)

Fig. S2.1: Design of Experiment 4

Fig. S2.2: Variation in time to bolting (a) and time between bolting and flowering (b) in Experiment 2

Fig. S2.3: Patterns of transition to bolting (a), transition from bolting to flowering (b) and day of first flower (c) in Experiment 3.

Fig. S2.4: Patterns of transition to bolting (a), transition from bolting to flowering (b), time to bolting (c), and time between bolting and flowering (D) in Experiment 4

Table S2.1: Populations used in Experiments 2, 3 and 4

Table S2.2a: Dependent variables analyzed in Experiments 2 & 4

Table S2.2b: Dependent variables analyzed in Experiment 3

Table S2.3: Comparison of model fit for linear and quadratic relationship with latitude for Experiments 2, 3 and 4

Table S2.4a: Conditions in each common garden

Table S2.4b: Dates of transplanting and phenology estimation in each common garden

Table S2.5: Estimation of first flower date in the common garden experiment based on flowering
stages.
Table S2.6: Estimates of the effect of latitude, cohort and their interaction on traits in Experiment
2
Table S2.7a: Estimates of the effect of latitude on traits in Experiment 3
Table S2.7b: Comparison of traits between common gardens in Experiment 3
Table S2.7c: Comparison of the effect of latitude on traits between common gardens in
Experiment 3
Table S2.8a: Estimates of the effect of latitude across three vernalization treatments on traits in
Experiment 4
Table S2.8b: Comparison of the effect of vernalization treatments on traits in Experiment 4
Table S2.8c: Comparison of the effect of latitude on traits between vernalization treatments in
Experiment 4
Supporting method S2.1: Raising of plants and seed rearing in Experiment 2
Supporting method S2.2: Parametrization of the hierarchical mixed-effect models used in
Experiments 2, 3, 4

Tables

Table 1: Test of variation in the day of first flower, length of vernalization and time between the start of the growing season and flowering across the latitudinal range of *Campanula americana* relative to a predicted breakpoint.

Dependent variable	Latitude (L)				Breakpoint (BP)			L * BP			R^2m	R^2c
	$\beta < BP$	$\beta > BP$	χ^2		β	χ^2		β	χ^2			
Day of first flower	-5.96	1.88	23.59	***	-273.10	39.52	***	7.85	37.84	***	0.18	0.18
Length of vernalization	2.63	10.75	13.57	***	-289.84	130.99	***	8.12	119.59	***	0.93	0.95
Time between flowering and the start of the growing season	-15.58	-1.37	107.87	***	-493.68	85.97	***	14.25	82.81	***	0.46	0.54

All dependent variables were assumed to follow Gaussian distributions. Each model was optimized with the *bobyqa* optimizer to improve convergence. Test statistics include the slope of each effect (β), the χ^2-value, and the marginal and conditional R^2 of the model. For the effect of latitude, β was estimated for the parts of the range below or above the predicted breakpoint (BP) in latitude (34.98 °N). *** $P<0.001$. Results for random effects are not shown.

Dependent variable	N	Latitude (L) X^2	Categorical effect (C) X^2	L * C X^2	R^2m	R^2c
Experiment 2			*Cohort*			
Flowering success †	994	**5.95****	**6.05***	3.62 (*)	0.51	0.52
Time to flowering	881	**9.40****	**50.82*****	**36.63*****	0.27	0.65
Time to bolting	893	**25.70****	0.46	3.02	0.15	0.51
Time between bolting and flowering	877	**40.08*****	**28.19*****	**31.25*****	0.46	0.73
Experiment 3			*Common garden*			
Flowering success †	2231	**31.60*****	**143.91*****	**58.07*****	0.39	0.51
Transition to bolting †	100	2.72 (*)	**3.23***	**3.63****	0.35	-
Transition from bolting to flowering †	2341	**7.09***	**59.31*****	**20.16****	0.44	0.56
Day of first flower	2136	**40.09*****	**1365.27*****	**161.40*****	0.53	0.70
Time to flowering (since transplanting)	2136	**40.09*****	**87.05*****	**161.40*****	0.34	0.58
Experiment 4			*Vernalization Trt*			
Flowering success †	1544	**97.33*****	**143.24*****	**74.50*****	0.69	0.80
Transition to bolting †	1544	**71.72*****	**143.11*****	**42.26*****	0.67	0.82
Transition from bolting to flowering †	979	**20.00*****	**18.26*****	**16.80*****	0.30	0.44
Time to flowering	711	2.92	**127.80*****	**152.03*****	0.20	0.64
Time to bolting	786	**14.49*****	**16.39*****	**13.63****	0.18	0.45
Time between bolting and flowering	711	0.66	**98.51*****	**117.23*****	0.33	0.67

Binary variables (†) were analyzed by models predicting non-zeros on the logit scale. All other dependent variables were assumed to follow Gaussian distributions. Each model was optimized with the *bobyqa* optimizer to improve convergence. Test statistics include the χ^2-value, and the marginal and conditional R^2 of the model. χ^2-value with P-values < 0.05 are in bold; (*) $P<0.1$, * $P<0.05$, ** $P<0.01$, *** $P<0.001$. The slope of each effect is reported in Table S2.6, S2.7 and S2.8. Results for random effects are not shown. Note that *transition to bolting* in Exp. 3 was analyzed at the population level in a simple linear model without random effects, we therefore report the F-value instead of χ^2, and R^2 instead of marginal and conditional R^2.

Figures

Fig. 1: Variation in flowering phenology observed in nature and associated climatic factors
across latitudes (Exp. 1). (**a**) Location of *C. americana* observations (black dots). The range of
the species is outlined with a gray line; red arrows represent the latitudinal delimitation of the rear
edge. Shading indicates the latitudinal temperature gradient based on minimum temperature of the
coldest month (bio6, WorldClim 2.0, (Fick & Hijmans, 2017). (**b**) Day of first flower inferred from
observations in 2018-2022 (**a**). (**c**) The length of vernalization and (**d**) the time between the start
of the growing season and flowering inferred for each observation from PRSIM climate data
(https://prism.oregonstate.edu). These variables best explained the differences in the day of first
flower below (red) or above (black) the identified breakpoint (34.98° N, dashed vertical lines) in
the latitudinal cline in day of first flower. Solid lines represent the significant model predicted
slope of the relationship between each variable and latitude, with the 95% confidence interval
indicated as shading. Test statistics reported in Table 1.

Fig. 2: Variation in reproductive phenology in a common environment (Exp. 2). (**a**) Location of populations (dots). The range of
the species is shaded in gray; red arrows represent the latitudinal delimitation of the rear edge. Flowering success (**b**) and time to
flowering (**c**) were recorded for two cohorts of populations raised in a common controlled environment. Lines represent the significant
model-predicted relationship between traits estimated at the individual level and population latitude, with the 95% confidence interval
indicated as shading (cohort 1) or dotted lines (cohort 2). Test statistics reported in Table 2 and Table S2.6.

Fig. 3: Range-wide variation in reproductive phenology in common gardens (Exp. 3). (**a**) Location of populations and common gardens. The range of the species is shaded in gray; red arrows representing the latitudinal delimitation of the rear edge. Populations are indicated by white dots, and common gardens by colored triangles. CG4 (gray) was not included in the analyses because of low survival. Flowering success in collection round 1 (**b**) and time to flowering since transplanting (**c**) were recorded for populations sampled across the range and raised in five common gardens across a latitudinal gradient. Colors distinguish the different gardens as depicted in (A), and their latitude is indicated with arrows along the x-axis. Lines represent the significant model-predicted relationship for each garden between traits estimated at the individual level and home latitude of the population, with the 95% confidence interval indicated by shading. Test statistics reported in Table 2 and Tables S2.7.

Fig. 4: Effect of vernalization length on range-wide variation in flowering success and phenology (Exp. 4). (**a**) Location of populations (dots). The range of the species is shaded in gray; red arrows represent the latitudinal delimitation of the rear edge. Flowering success (**b**), and time to flowering (**c**) were recorded populations sampled across the range and raised in three vernalization treatments (six, four or two weeks) in controlled conditions. For time to flowering, the 2-week treatment is not shown as too few individuals flowered. Lines represent the significant model-predicted relationship between each trait estimated at the individual level and population latitude in each treatment, with the 95% confidence interval indicated as shading. Test statistics reported in Table 2 and Tables S2.8.