Selective inhibition in CA3: A mechanism for stable pattern completion through heterosynaptic plasticity

Gyeongtae Kim
Department of Mathematical Sciences
Ulsan National Institute of Science and Technology (UNIST)
Ulsan, Republic of Korea
kgt1220@unist.ac.kr

Pilwon Kim*
Department of Mathematical Sciences
Ulsan National Institute of Science and Technology (UNIST)
Ulsan, Republic of Korea
pwkim@unist.ac.kr

ABSTRACT

Neural assemblies representing different engrams compete for successful retrieval in the CA3 region of the hippocampus, but the detailed mechanisms underlying their formation remain elusive. Recent research reveals that hippocampal inhibitory neurons respond selectively to stimuli and exhibit diverse plasticity, suggesting their important role in engram formation. Conventional attractor network models for CA3 typically employ global inhibition, where inhibitory neurons uniformly suppress the activity of excitatory neurons. However, these models may not fully capture the complex competition dynamics arising from sparse distributed coding and may not accurately reflect the specific roles of inhibitory neurons in the competition between neural assemblies during memory retrieval. We propose an engram formation mechanism in CA3 using a spiking neural network model, highlighting the critical role of the association between excitatory and inhibitory neurons through heterosynaptic plasticity. In our model, inhibitory neurons are associated with specific neural assemblies during encoding and selectively inhibit the excitatory neurons involved in competing assemblies during retrieval. To explore the functional advantages and validate the proposed mechanisms, we developed a simplified, feed-forward model of the dentate gyrus (DG). Combined with the activation of inhibitory neurons in CA3, this model led to sparsely distributed engrams in CA3, with configurations that included partially overlapping, competing engrams. The results demonstrated that selective inhibition provides more stable pattern completion and enhances retrieval performance compared to global inhibition alone. Additionally, the observed neural activity in the hippocampal subregions of the model was consistent with experimental findings regarding these regions' roles in pattern separation and pattern completion.

1 Introduction

The hippocampus is a crucial region for processing episodic memory (O'Keefe and Nadel, 1978; Tulving, 1983). In this region, engrams—neural circuits representing the physical traces of memories—are stored and retrieved (Semon et al., 1923; Josselyn and Tonegawa, 2020). Engrams are specifically formed in CA3 in a sparse and distributed manner, facilitated by sparse activations from the DG (Wixted et al., 2014). This process, known as pattern separation, creates distinct and non-overlapping representations of similar events or stimuli. It is one of the key functions of the hippocampus, which reduces interference between memories (Marr, 1971; McClelland et al., 1995; Berron et al., 2016; Rolls, 2016; Knierim and Neunuebel, 2016). Successful pattern separation is achieved when distinct representations in CA3 are formed regardless of similarities in external information (Leutgeb et al., 2007; Treves et al., 2008; Yassa and

Stark, 2011; Santoro, 2013).

While the DG generates sparse and highly distinct representations that substantially differ from the original inputs, the hippocampus can still successfully retrieve specific memories through CA3. This process, the reconstruction of a complete memory from partial or incomplete cues, is called pattern completion (Marr, 1971; McClelland et al., 1995; Norman and O'Reilly, 2003; Lee et al., 2004; Rolls, 2016; Knierim and Neunuebel, 2016; Guzman et al., 2016). It relies on two critical hippocampal pathways: the direct perforant path (PP) from the entorhinal cortex (EC) to CA3, and the recurrent collateral (Rc) connections within CA3. The direct PP encodes features that enable neural assemblies in CA3 to recognize similarities between incoming cues and stored memories. CA3 functions as an attractor network, where Rc connections are crucial for activating the full neural assembly (Hopfield, 1982; Káli and Dayan, 2000; Rolls, 2007; Rennó-Costa et al., 2014). By amplifying the activity of excitatory neurons associated with a specific engram, Rc connections ensure accurate retrieval of the intended memory while suppressing interference from competing memories. Although the processes of pattern separation and pattern completion in the hippocampus are well understood, the underlying mechanisms related to engrams in CA3 remain elusive, particularly concerning the role of inhibitory neurons in these processes. Attractor network models describe pattern completion in CA3 in the context of memory competition (Hopfield, 1982; Káli and Dayan, 2000; Rolls, 2007; Rennó-Costa et al., 2014), but they do not specifically address the mechanisms underlying engram formation. Most of these models assume completely separate neural assemblies that compete through a winner-takes-all process. They commonly employ global inhibition, where inhibitory neurons suppress all excitatory neurons.

Recent research has highlighted the crucial role of inhibitory neurons in the memory process (Raven and Aton, 2021; Cummings et al., 2021). Stimulus selectivity was primarily associated with excitatory neurons (Usrey et al., 2003; Niell and Stryker, 2008). However, recent studies reveal that many GABAergic neurons exhibit greater stimulus selectivity than previously recognized.(Li et al., 2014; Khan et al., 2018). In addition, increasing evidence shows the presence of synaptic plasticity mechanisms in inhibitory neurons. Both long-term potentiation (LTP) and long-term depression (LTD) have been observed at inhibitory neuron synapses, similar to what has been previously observed at excitatory synapses (Perez et al., 2001; Owen et al., 2013; Cohen et al., 2016; Korotkova et al., 2010; He et al., 2021). This plasticity contributes to engram formation and refinement by modulating inhibitory control in response to experience. Extending beyond their contribution to sparse distributed coding (Acsády et al., 1998; Rolls, 2013), these observations suggest that inhibitory neurons play a broader role in memory encoding, storage, and retrieval processes.

Building on the evidence discussed above, our work aims to explore the roles of inhibitory neurons in memory processing using a modeling approach. We propose that inhibitory neurons offer functional advantages by providing selective inhibition in memory competition, beyond the effects of global inhibition, potentially enhancing the stability and efficiency of memory retrieval. We developed a spiking neural network model that preserves the structural and connectional properties of the hippocampus, with a focus on the trisynaptic circuit—the pathway from the EC to CA3 via the DG. Drawing on evidence of heterosynaptic plasticity observed in various connections (Tsukamoto et al., 2003; Hunt et al., 2013; Chater and Goda, 2021; Jenks et al., 2021), we propose a mechanism of engram formation where heterosynaptic plasticity between excitatory and inhibitory neurons in CA3 naturally induces selective inhibition during memory competition.

In the proposed model, selective inhibition is characterized by inhibitory neurons that are associated with a specific neural assembly in CA3: these neurons inhibit excitatory neurons in competing assemblies while sparing those within their own assembly. To investigate the effect of selective inhibition on pattern completion, we simplified the DG into a feed-forward structure. This simplification enables instant sparse distributed coding and, together with the proposed engram formation mechanism, creates various competitive conditions. These conditions result in situations within CA3 where multiple overlapping engrams of varying sizes compete during memory retrieval.

We conducted pattern separation and pattern completion tasks across the entire hippocampal model, made possible by the stable retrieval performance. The resulting activation patterns in each subregion were then compared with experimental data. The analysis shows that selective inhibition by inhibitory neurons enhances the stability and efficiency of competition between neural assemblies. These findings provide novel insights into the role of inhibitory neurons in pattern completion and separation within the hippocampus.

2 Results

2.1 Model architecture and learning mechanisms

We constructed a spiking neural network that preserves hippocampal structure to model the CA3 engram formation process and the competition between engrams during retrieval ('Materials and methods'). The model focuses on investigating the stability of pattern completion under varying conditions. To achieve this, engrams in CA3 should be naturally formed for arbitrary inputs via the DG, resulting in a sparse distributed representation. However, modeling DG pattern separation is challenging due to its intricate circuitry, complex synaptic connectivity, and heterogeneous neuronal types (Jinde et al., 2013; Cayco-Gajic et al., 2017; Hainmueller and Bartos, 2020). Therefore, we simplified

Figure 1: Overview of model architecture and learning. **(A)** The model includes hippocampal subfields: DG, CA3, and CA1, as well as the superficial and deep layers of the EC. The model operates in two distinct phases aligned with theta oscillations, where encoding and retrieval are processed separately. **(A1)** Information flow during the encoding phase. The superficial EC sends inputs to the DG and deep EC, but not directly to CA3. Synaptic plasticity occurs at the silenced connections during encoding, specifically at the PP from the superficial EC to CA3, Rc, and Sc. **(A2)** Information flow during the retrieval phase. The superficial EC sends inputs directly to CA3, which retrieves memories in the deep EC through Rc, Sc, and CA1. **(B)** Symmetric STDP kernel used during the encoding phase. **(C)** Synaptic plasticity across various connections. Abbreviations: ext: external input; mf: mossy fiber; exc: excitatory neuron; inh: inhibitory neuron.

our approach by focusing solely on spatial pattern separation for non-sequential inputs, represented as 16-dimensional binary vectors at a fixed frequency ('Materials and methods'). Under these conditions, we designed a feedforward form of the DG, which we will explain in more detail in the next section, 'Feedforward Form of the Dentate Gyrus and Its Inherent Ability for Spatial Pattern Separation.'

Additionally, we simplified CA1's role to reinstating original information from CA3 engrams ('Materials and Methods'). All neurons in the network were modeled using the Izhichevich model (Izhikevich and Edelman, 2008) distinguishing hippocampal neurons as either excitatory or inhibitory ('Materials and Methods'). The model operates in distinct phases according to theta oscillation theory (Hasselmo and Stern, 2014). Theta oscillations, rhythmic brain waves typically in the 4-8 Hz range, are prominent in the hippocampus and play a crucial role in memory processes. Several theories

have been proposed to explain the specific functions of theta waves (Hasselmo and Stern, 2014; Drieu and Zugaro, 2019; Herweg et al., 2020). One theory suggests that theta oscillations divide information processing into discrete time windows, optimizing memory by coordinating information flow across hippocampal subregions (Bragin et al., 1995; Colgin, 2016; dos Santos et al., 2018; Kragel et al., 2020; Reddy et al., 2021; Ter Wal et al., 2021; Lowet et al., 2023; Gattas et al., 2023).

During the encoding phase, the superficial EC sends inputs to CA3 via the DG and to the deep EC, which then sends inputs to CA1 (Figure 1A1). Four types of connections—direct PP, Rc, Schaffer collateral (Sc), and the connection from CA1 to deep EC—are silenced but still undergo learning through heterosynaptic plasticity, except for the connection to deep EC (Figure 1C). We applied symmetric spike-timing-dependent plasticity (STDP) across the model for simplicity, unifying the various STDP types that exist for different connections based on the symmetric STDP observed in the CA3 recurrent circuit ('Materials and Methods' and Figure 1B) (Wittenberg and Wang, 2006; Mishra et al., 2016; Rebola et al., 2017; Ecker et al., 2022).

During the retrieval phase, all connections that were silenced during encoding are activated, while those active during encoding are silenced. The superficial EC sends inputs to CA3 via the direct PP, where competition among neural assemblies occurs, ultimately selecting the dominant engram for pattern completion. The winning engram in CA3 activates CA1, which in turn triggers the neurons in the deep EC (Figure 1A2). Each phase lasts 120 ms, providing sufficient time for information storage and retrieval. Because there is no interaction between the encoding and retrieval phases in the model, we did not synchronize these phases with oscillatory cycles for most tasks, nor did we base oscillation generation on biological processes such as those in the medial septum (Cheng and Yakel, 2015; Robinson et al., 2016; Unal et al., 2018; Müller and Remy, 2018; Nuñez and Buño, 2021; Gedankien et al., 2023). Our primary focus was on the distinct information flows for each phase to simplify the model.

2.2 Engram formation with selective inhibition through heterosynaptic plasticity

This section details the mechanism of engram formation in the model, with a focus on the novel role of selective inhibition mediated by heterosynaptic plasticity. Mossy fibers from the DG project to both pyramidal neurons and inhibitory neurons in the CA3 region, facilitating sparse distributed coding (Acsády et al., 1998; Mori et al., 2007; Torborg et al., 2010; Treviño et al., 2011; Cosgrove et al., 2010; Prince et al., 2016). These fibers transmit the outputs of pattern separation performed by the DG. We specifically concentrated on mossy fiber inputs to inhibitory neurons in CA3 due to their role in feedforward inhibition, which regulates pyramidal neuron activity (Ruediger et al., 2011; Morrison et al., 2016; Stefanelli et al., 2016; Guo et al., 2018; Viana da Silva et al., 2019; Josselyn and Tonegawa, 2020; Giorgi and Marinelli, 2021). Our hypothesis extends beyond the conventional view of feedforward inhibition as merely a sparsification mechanism (Acsády et al., 1998; Rolls, 2013). We propose that it facilitates selective inhibition, enhancing the competitive dynamics between neural assemblies through heterosynaptic plasticity.

Heterosynaptic plasticity, in which synaptic changes in one set of synapses influence another set, is well-documented (Tsukamoto et al., 2003; Hunt et al., 2013; Chater and Goda, 2021; Jenks et al., 2021). This form of plasticity occurs when the strength of a synapse changes due to activity at a different, nearby synapse. For example, the activation of one synapse can lead to the strengthening or weakening of neighboring synapses, even if those synapses were not directly activated. This mechanism enables more complex forms of learning compared to traditional Hebbian plasticity. Multiple studies have demonstrated that heterosynaptic plasticity occurs in the direct PP, Rc, and Sc pathways (Tsukamoto et al., 2003; Hunt et al., 2013; Chater and Goda, 2021; Jenks et al., 2021). However, most of this evidence pertains to excitatory neurons. In light of this evidence, we hypothesize that heterosynaptic plasticity also occurs from excitatory to inhibitory neurons in CA3. Building on this evidence and our hypothesis, the proposed model incorporates heterosynaptic plasticity at each type of connection (Figure 1C).

During the encoding phase, the inhibitory neurons suppress most of the excitatory neurons. The remaining excitatory neurons, along with the activated inhibitory neurons, form a neural assembly through heterosynaptic plasticity (Figure 2A). In addition, these remaining excitatory neurons form associations with activated neurons in the superficial EC and CA1 by strengthening direct PP and Sc connections, respectively. These strengthened connections contribute to engram formation (Figure 1A2).

The proposed learning mechanism ensures that inhibitory neurons within an engram do not inhibit the excitatory neurons of the same engram (Figure 2C). However, inhibitory neurons are not limited to a single engram; instead, they exhibit stimulus selectivity, aligning with recent findings that they can show selective responses to stimuli (Li et al., 2014; Khan et al., 2018; Cummings et al., 2021). In our model, we refer to this stimulus selectivity as information selectivity, as it reflects the ability of neurons to store specific pieces of information based on the encoded stimuli. We examined the information storage capacity of each CA3 neuron by assessing the amount of information retained after encoding 500 arbitrary inputs. As shown in Figure 2B, most excitatory neurons in CA3 (approximately 85%) store only one or two pieces of information. In contrast, inhibitory neurons have a broader range of storage capacities, with some storing significantly more information. These results suggest that while inhibitory neurons tend to be information-selective, this does not imply that they are restricted to storing only a single piece of information.

Figure 2: Illustrations and properties for selective inhibition. **(A)** The process of engram formation in CA3 provides the formed engram with the capability for selective inhibition. Inhibitory neurons activated by mossy fibers inhibit most of the excitatory neurons. The surviving excitatory neurons, along with the activated inhibitory neurons, form a neural assembly through STDP. **(B)** Distribution of CA3 neurons based on the number of stored memories. **(C)** Characterization of CA3 connections with inhibitory neurons for global versus selective inhibition. In the case of global inhibition, inhibitory neurons suppress excitatory neurons without regard to which engram they represent. In contrast, for selective inhibition, inhibitory neurons associated with a specific engram do not inhibit excitatory neurons within that same engram. **(D)** The inhibitory influence of one engram on others as a function of the degree of overlap between engrams.

We also investigated how the overlap between CA3 engrams, specifically the shared excitatory neurons within CA3, affects selective inhibition. To evaluate this, we measured the ratio of suppressed excitatory neurons in each neural assembly caused by inhibitory neurons associated with another assembly. This approach allowed us to assess the extent to which one engram inhibits others. We generated 500 engrams in CA3 corresponding to 500 inputs. Engrams in each pair were selected based on their overlap ratio ('Materials and Methods'), and we divided the overlap ratio into 25 intervals, each representing a 4% range. The inhibition ratio was then averaged within each interval. As shown in Figure 2D, the inhibition power of an engram over other engrams decreases as the overlap ratio between engrams increases, suggesting that selective inhibition power is reduced with higher overlap. This observation may underscore the importance of sparse distributed coding in CA3 (Marr, 1971; McClelland et al., 1995; Norman and O'Reilly, 2003; Rolls, 2016; Knierim and Neunuebel, 2016; Guzman et al., 2016).

The strengthened connections from excitatory to inhibitory neurons facilitate the earlier activation of inhibitory neurons during retrieval. These activated inhibitory neurons then inhibit most excitatory neurons, except those within the same neural assemblies in CA3, effectively guiding the retrieval cue toward a single memory. Because we initialized the connection weights from excitatory to inhibitory neurons as nonzero ('Materials and Methods' and Figure 5A3), global

Figure 3: The DG in the model has a feedforward structure consisting of the hilus layer and the GCL. Both layers contain excitatory and inhibitory neurons: excitatory neurons are represented in gray, and inhibitory neurons are represented in red. Superficial EC sends inputs to all types of neurons in the DG. Inhibitory neurons in the GCL also receive inputs from excitatory neurons in the hilus layer, which are regulated by inhibitory neurons in the hilus layer. Finally, excitatory neurons in the GCL, regulated by inhibitory neurons within the GCL, send outputs to CA3 via mossy fibers.

inhibition becomes effective once a sufficient number of neurons in one assembly have begun firing.

We propose that selective inhibition provides targeted suppression of competing engrams, while global inhibition helps maintain overall network stability. This inhibition mechanism could lead to more accurate pattern completion and reduced interference between memories with similar information. To demonstrate the stability of pattern completion with selective inhibition under natural conditions, we first need to construct a DG model that generates engrams in CA3 with various characteristics. This model will enable a comprehensive evaluation of our inhibition mechanism across diverse scenarios of engram competition. The following section details the construction of this DG model and assesses its performance in sparse distributed coding through pattern separation.

2.3 Feedforward form of the dentate gyrus and its inherent ability for spatial pattern separation

To equip the DG with spatial pattern separation capabilities, we modeled its structure based on real anatomical features. We excluded the recurrent circuits, creating a feedforward configuration. Numerous modeling studies have demonstrated the pattern separation performance of the DG (Myers and Scharfman, 2009; Faghihi and Moustafa, 2015; Chavlis et al., 2017; Cayco-Gajic et al., 2017; Madar et al., 2019; Modhej et al., 2020; Guzman et al., 2021; Kim and Lim, 2022), and it is well-established that a feedforward DG can achieve this function (Cayco-Gajic et al., 2017). In the proposed model, the DG consists of two distinct layers: the hilus layer and the granule cell layer (GCL) (Figure 3). Although the real DG contains various neuron types in each layer, such as hilar interneurons, mossy cells, basket cells, and granule cells (Freund and Buzsáki, 1996; Amaral et al., 2007; Danielson et al., 2017; Hainmueller and Bartos, 2020), we simplified these into excitatory and inhibitory neurons using the Izhikevich neuron model ('Materials and Methods'). Therefore, each layer includes both excitatory and inhibitory neurons (Figure 3).

During the encoding phase, inhibitory neurons in the GCL receive two types of excitatory inputs: direct excitation from the superficial EC and excitation from the hilus layer. The latter is modulated by inhibitory neurons within the hilus layer, known to play an important role in pattern separation through the disynaptic effect (Kim and Lim, 2022). Both inputs detect features of the given input pattern, leading to the activation of inhibitory neurons in the GCL for most input patterns (Figures 3 and 4A). Excitatory neurons in the GCL become active when they simultaneously escape inhibition from GCL inhibitory neurons and receive excitatory input from the superficial EC. This mechanism transforms low-dimensional information into very sparse, high-dimensional information. The resulting activation patterns of excitatory neurons in the GCL stimulate both excitatory and inhibitory neurons in CA3 via mossy fibers, creating sparse distributed representations. We passed 10 arbitrary inputs through the DG, and Figure 4A illustrates how these inputs are separated within the excitatory neuron population in the GCL.

To evaluate the pattern separation performance of our DG model, we presented 1,000 input patterns and observed the

Figure 4: DG spike raster and pattern separation performance. **(A)** Raster plot showing DG activity during the encoding of 10 different inputs (A to J). The activity is displayed for various regions, including the superficial EC, excitatory and inhibitory neurons in the hilus layer, and excitatory and inhibitory neurons in the GCL. For better visualization, 50 neurons are arbitrarily selected from the 100 excitatory neurons in the hilus layer and the 400 inhibitory neurons in the GCL. Additionally, 16 excitatory neurons in the GCL, involved in encoding the 10 inputs (from a total of 800 neurons), are shown. **(B)** Pattern separation performance of the DG. The graph plots the separation performance against input similarity. **(C)** Distribution of engram pairs based on the similarity between activation patterns in the superficial EC (input) and those of excitatory neurons in the GCL of the DG (output).

activation patterns of excitatory neurons in the GCL. We calculated the overlap degree for each pair of patterns using the measure suggested by Löffler and Gupta (2022), with the discrimination value defined as one minus the overlap value ('Materials and Methods'). The output discrimination values were averaged within each of the 25 windows, with each window spanning 0.04 units (Figure 4B). This figure shows that the output discrimination values rapidly approach 1.0 when input discrimination values exceed approximately 0.1, indicating effective pattern separation. High output discrimination is maintained for input discrimination values above this threshold. Figure 4C illustrates the distribution of input and output pairs based on their similarity, comparing input pairs from the superficial EC with output pairs from the DG. The analysis indicates that a significant majority of input pairs (over 85%) are fully separated, as reflected by the high percentage of output pairs with minimal similarity. Both Figures 4B and 4C underscore the DG's spatial pattern separation ability in the model, demonstrating that a feedforward form is sufficient for this function even without any learning.

2.4 Sparse distributed coding for memory storage and retrieval

The proposed model, equipped with a feedforward structure of the DG, successfully separates patterns and projects inputs into a high-dimensional space. his process contributes to the sparse distributed representation of memories in

Figure 5: Encoding the various inputs. **(A)** Weight matrices learned through STDP during encoding. Weights represent the fixed connection weights multiplied by the learned peak conductances. Although there are 2,400 CA3 excitatory neurons, only those involved in the encoding of 10 specific examples are selected and arranged for visualization. **(A1)** Learned direct PP weight matrix from superficial EC neurons to CA3 excitatory neurons. **(A2)** Learned excitatory Rc weight matrix. **(A3)** Learned Rc weight matrix from CA3 excitatory neurons to CA3 inhibitory neurons. **(A4)** Learned Rc weight matrix from CA3 inhibitory neurons to CA3 excitatory neurons. **(A5)** Learned Sc weight matrix from CA3 excitatory neurons to CA1 neurons. **(B)** Raster plot showing network activity during the encoding of 10 different inputs (A to J). The activity is depicted for different regions, including the superficial EC, excitatory neurons in the GCL, CA3 excitatory and inhibitory neurons, CA1, and deep EC. Other types of neurons in the DG are omitted for clarity. As in **(A)**, only the CA3 excitatory neurons involved in the 10 specific examples are

Figure 5 continued on next page

CA3, allowing for a diverse range of neural assembly sizes and relationships among assemblies. First, we assessed the storage capacity of the CA3 region in the model to ensure it could adequately support our investigation into the stability of pattern completion. We did this by examining the number of distinct engrams formed in CA3 relative to the number

Figure 5 continued

selected for visualization. **(C)** CA3 capacity is depicted as the number of engrams formed versus the number of inputs. **(D)** Correlation between input size and CA3 engram size, with a linear regression.

of inputs (Figure 5C). The graph shows a consistent and steady increase in the number of engrams within CA3, rising by approximately 200 engrams as the number of inputs increases from 800. This capacity is sufficient for investigating the stability of selective inhibition in pattern completion.

Additionally, we analyzed the correlation between the size of the input pattern and the size of the neural assembly in CA3. The size of the patterns in each subregion is defined as the number of firing neurons in the superficial EC and CA3 regions, respectively. Figure 5D shows no correlation between input size and engram size in CA3, indicating that each engram in CA3 typically consists of around 10 excitatory neurons. This suggests that the DG effectively separates input features before they reach CA3.

Figures 5B and 6A illustrate the firing patterns for 10 arbitrary inputs during the encoding and retrieval phases, respectively. Instead of modifying the synaptic weights directly, we updated the peak conductances, which function similarly to conventional synaptic weights ('Materials and Methods'). This approach keeps the synaptic weights for each connection fixed, thereby maintaining the connectivity ratio. For readability, however, we will refer to the learned weights as the product of the peak conductance and synaptic weights for each connection (Figure 5A).

During encoding, inputs separated by the DG result in distinct firing patterns of excitatory neurons in the GCL, which subsequently activate distinct patterns of neurons in CA3, forming sparsely distributed patterns (Figure 5A2). The learned weights for Rc connections, which are connections from excitatory neurons to other excitatory neurons, show some weights deviating from the diagonal axis, indicating sparsely distributed storage. In contrast, no diagonal patterns were observed for other types of connections, particularly those involving inhibitory neurons in CA3 (Figures 5A3 and 5A4). This suggests that while inhibitory neurons are selective to inputs (Figure 2B), they are not specifically constrained to particular engrams.

To observe engram competition during the retrieval phase, we presented an input with 0.5 similarity to both engrams E and G 10 times during retrieval. This input, sent to the superficial EC, flows to CA3 via the direct PP, acting as a retrieval cue. Noise was introduced to CA3 excitatory neurons at a frequency of 3.5 Hz to simulate various factors present in a real brain, which are essential for the attractor network (Webb et al., 2011). Under these conditions, the model successfully retrieved either input G or E in 80% of the trials. Failures were defined as instances where no clear winner emerged from the competition, resulting in interference or no activation in the deep EC (Figure 6A). An analysis of the average firing rate over a 30 ms window for each CA3 neural assembly revealed that only one assembly displayed a significantly high firing rate of approximately 80 to 120 Hz during successful pattern completion, while others exhibited rates of less than about 10 Hz (Figures 6B1 and 6B2). This suggests that the dominant assembly effectively suppressed the activity of competing assemblies. In cases of failed pattern completion, no dominant assembly was observed (Figure 6B3).

These results demonstrate the model's ability to retrieve memories based on their similarity to retrieval cues. While the DG processes input features to create distinct representations before they reach CA3, the direct PP preserves the original features, allowing for more direct associations with similar memories. To evaluate the function of the direct PP, we deactivated it during retrieval and used mossy fibers to transmit information to CA3. We tested varying percentages of partial cues for each memory to assess the role of the direct PP in capturing input completeness (Figure 7A). Figure 7C illustrates retrieval performance, averaged over 5 samples for each percentage of partial cue. The results show significantly higher success rates when the direct PP is intact, compared to when it is silenced. This difference occurs because the DG perceives the cue as distinct information, activating entirely different neurons in the CA3 region through pattern separation. In contrast, the direct PP stimulates neurons associated with memories similar to the cue (Figures 7B1 and 7B2).

2.5 Stability of pattern completion with selective inhibition under various conditions.

We demonstrated that the model can encode inputs in a sparse and distributed form through pattern separation by the DG and retrieve these inputs via the direct PP through pattern completion. Additionally, the model is well-suited for investigating the stability of pattern completion. The sparsely distributed nature of engrams in CA3 exhibits various properties that may influence competition during pattern completion. We identified four key conditions affecting this competition: (1) disparity in input strength to CA3 engrams, (2) overlap between CA3 engrams, (3) the number of competing engrams for a single cue, and (4) size differences between CA3 engrams, defined by the number of excitatory neurons involved in each engram. We examined the impact of these conditions on engram competition and compared the model's retrieval performance under selective inhibition with that under global inhibition. To simulate the effects of global inhibition, we maintained constant initial peak conductances from excitatory to inhibitory neurons in CA3. By

Figure 6: Retrieval for learned inputs. **(A)** Raster plot showing network activity in response to a cue similar to inputs E and G among 10 different inputs (A to J). The depicted layers and neurons correspond to those in Figure 3B. The retrieval cue is presented 10 times, resulting in either successful retrieval (E, G) or retrieval failure, as observed in the activity patterns of the deep EC. **(B)** Firing rate over time for neural assemblies in CA3 during retrieval, illustrating different scenarios of memory retrieval. **(B1)** Example of successful retrieval of input G, showing a dominant firing rate for the neural assembly representing G. **(B2)** Example of successful retrieval of input E, showing a dominant firing rate for the neural assembly representing E. **(B3)** Example of retrieval failure, where no single neural assembly dominates the activity.

not applying plasticity to the connections from excitatory to inhibitory neurons in CA3, we removed the capability for selective inhibition.

2.5.1   Robust pattern completion against input strength disparity

According to the encoding specificity principle, retrieval is most effective when the retrieval cue closely matches the features present during the initial encoding of the memory (Tulving and Thomson, 1973; Ji, 2023). However, retrieval can become challenging when competing memories share significant similarities with the retrieval cue, leading to interference. While such interference can naturally result in the generalization of competing memories, it becomes problematic when distinguishing between these memories is necessary. One approach to address this challenge is

Figure 7: The effect of the direct perforant path for pattern completion during retrieval. **(A)** Illustration of different percentages of partial cues for a learned input A. **(B)** Firing rate heatmap for CA3 neurons in response to a cue during retrieval (120 ms). Since the model may retrieve different outputs in each trial for a given cue, the heatmaps display the firing pattern with the highest likelihood of retrieval across 20 repetitions. **(B1)** Example heatmap for a cue in the intact direct PP condition, showing similar firing patterns regardless of the cue variation. **(B2)** Example heatmap for a cue when the direct PP is silenced, displaying different firing patterns for each cue variation. **(C)** Retrieval performance as a function of the percentage of partial cues under two conditions: with direct PP (blue) and without direct PP (red). Performance is measured as the mean retrieval success rate, averaged over 5 samples for each cue percentage, except for the 100% cue case, which is based on a single sample. The success rate for each sample is averaged over 20 repetitions.

by providing a biased input to one memory, thereby reducing the similarity of another memory to the cue (Deco and Rolls, 2006). However, when two memories contain highly similar information, introducing a biased input becomes challenging and may lead to poor retrieval performance. This, in turn, reduces the effectiveness of the dentate gyrus in distinguishing highly similar information.

We explored the role of selective inhibition in stabilizing pattern completion, even when the cue was uniformly similar to the memories encoded by the two competing CA3 engrams. We modulated input bias by adjusting the similarity of cues to one memory relative to another. Specifically, we varied the number of common neurons shared between the cue and the encoded input (Figure 8A). To specifically assess the impact of input bias, we selected 15 pairs of inputs, each stored in CA3 engrams of the same size, with no overlap between engrams (Figure 8B). Each input pair was tested with five different input bias levels, and retrieval was tested 50 times for each level.

Our results showed that when the cue was uniformly similar to the memories encoded by two CA3 engrams, retrieval performance under global inhibition was approximately 60%. This suggests that non-biased inputs can hinder engram competition, increasing the likelihood of interference between memories, even when they are fully separated (Figure 8D). In contrast, under selective inhibition, retrieval performance remained consistently high, around 90%, regardless of disparities in input strength This consistency is likely because selective inhibition enhances the contrast between competing memories, effectively amplifying small differences in input strength. These findings indicate that selective inhibition can improve the robustness of memory retrieval in the hippocampus by maintaining pattern completion performance even when cues are not biased toward a particular memory.

2.5.2 Pattern completion with overlapping engrams

Next, we examined the impact of overlap between engrams (Figure 8E). In CA3, engrams are encoded using sparse distributed representations, which can result in varying degrees of overlap. Although there may be mechanisms that could mitigate interference from overlapping engrams, such overlaps still pose a significant risk of interference, potentially reducing retrieval performance. To investigate this effect, we prepared 60 input pairs, each stored in CA3

Figure 8: The effect of input strength disparity and overlap ratio between CA3 engrams on retrieval. **(A)** Schematic of the retrieval cue with different input strengths for each engram. **(B)** Example of the learned Rc weight matrix in the input strength disparity task. **(C)** Firing rate over time for neural assemblies in CA3 during retrieval in the input strength disparity task. **(C1)** and **(C2)** show results for the same cue with the same learned samples. **(C1)** Example where engram A is retrieved. **(C2)** Example when B is retrieved. **(D)** Retrieval performance at different levels of input strength disparity, comparing selective inhibition (blue) and global inhibition (red). Performance is measured as the mean retrieval success rate, averaged over 15 samples for each input bias level. Each sample's success rate is averaged over 50 repetitions. **(E)** Schematic of the retrieval cue with overlapping engrams. **(F)** Example of the learned Rc weight matrix in the overlap task. **(G)** Firing rate over time for neural assemblies in CA3 during retrieval in the overlap task. **(G1)** and **(G2)** show results for the same cue with the same learned samples.
Figure 8 continued on next page

engrams of the same size, while varying the overlap ratio from 0 to 0.3 (Figure 8F). Each pair of memories received uniformly similar partial cues, and retrieval was tested 50 times.

Figure 8 continued

(G1) Example when A is retrieved. **(G2)** Example when B is retrieved. **(H)** Retrieval performance at different overlap ratios between neural assemblies, comparing selective inhibition (blue) and global inhibition (red). Performance is measured as the mean retrieval success rate, averaged over 15 samples for each overlap ratio. Each sample's success rate is averaged over 50 repetitions.

Figure 9: The effect of the number of engrams in CA3 on retrieval. **(A)** Schematic of the retrieval cue with multiple engrams competing for retrieval. **(B)** Example of the learned Rc weight matrix in the task where multiple engrams compete. **(C)** Firing rate over time for neural assemblies in CA3 during retrieval. **(C1)** to **(C6)** show results for the same cue with the same learned samples. **(C1)** Example when A is retrieved. **(C2)** Example when B is retrieved. **(C3)** Example when C is retrieved. **(C4)** Example when D is retrieved. **(C5)** Example when E is retrieved. **(C6)** Example when F is retrieved. **(D)** Retrieval performance across different numbers of engrams, comparing selective inhibition (blue) and global inhibition (red). Performance is measured as the mean retrieval success rate, averaged over 15 samples for each condition involving different numbers of engrams. Each sample's success rate is averaged over 50 repetitions.

The results showed that retrieval performance generally declined as the overlap ratio between engrams increased. However, selective inhibition consistently outperformed global inhibition at all overlap ratios (Figure 8H). Specifically, selective inhibition maintained about about 70% performance until the overlap ratio reached 0.1, whereas global inhibition drastically decreased to about 30% at the same overlap ratio. Although selective inhibition also experienced a sharp decline, dropping to about 35% as the overlap ratio increased to 0.3, it still performed better than global inhibition. These findings suggest that while the effectiveness of selective inhibition diminishes with higher overlap, it still mitigates interference between overlapping engrams more effectively than global inhibition. This indicates that selective inhibition helps maintain memory integrity, even in the presence of potential interference.

2.5.3 Pattern completion performance relative to the number of competing engrams

We also considered scenarios where multiple memories compete for retrieval from a single cue (Figure 9A). As the number of memories associated with a cue increases, it becomes more challenging for the cue to retrieve specific memories effectively (Watkins and Watkins, 2013; Parkin, 2007), leading to a natural decline in retrieval performance. Nonetheless, studying the effects of selective inhibition in this context is valuable, as it helps us understand its role in managing situations with multiple competing memories. To explore this, we prepared 75 sets of inputs, each stored in CA3 engrams of the same size and without overlap, while varying the number of competing engrams from 2 to 6. (Figure 9B). As in other tasks, each pair of memories received uniformly similar partial cues, and retrieval was tested 50 times.

Figure 9D shows that retrieval performance generally decreases as the number of competing engrams increases. For selective inhibition, retrieval performance gradually decreased from about 80% to about 60% as the number of competing engrams increased. In contrast, under global inhibition, performance remained steady at about 60% when the number of competing engrams was 3, but then drastically dropped to about 30% as the number of competing engrams increased to 6. These data suggest that selective inhibition can mitigate some of the negative effects more effectively than global inhibition, even as the number of competing memories increases.

2.5.4 Reduced sensitivity to size variations of the engram in CA3 under selective inhibition

In the model, the size of the engram in CA3 shows no correlation with input information (Figure 5D), suggesting that engram size should not inherently affect pattern completion. Ideally, input similarity should have a greater influence than engram size to prevent scenarios where a less similar memory is easier to retrieve because it just has a larger engram. We investigated the impact of size differences to determine whether a larger engram has an advantage in retrieval and how selective inhibition manages this size disparity compared to global inhibition (Figure 10A). We prepared 75 sets of input pairs, each stored in CA3 engrams with varying sizes and no overlap between engrams (Figure 10B). Each pair of memories received uniformly similar partial cues, and retrieval was tested 50 times.

The results show that selective inhibition consistently maintained about 80% retrieval performance, outperforming global inhibition, which achieved about 60% (Figure 10D). This result is consistent with the findings in Figure 8D. However, the size of the engram influenced which memory was retrieved. We conducted 50 trials to investigate which memories were retrieved and calculated the retrieval ratio by comparing the difference in retrieval counts for each memory type to the total number of successful retrievals. Memories stored in larger engrams were more likely to dominate the retrieval process. This effect was particularly pronounced under global inhibition, with larger engrams accounting for about 30% of retrievals, compared to about 20% under selective inhibition (Figure 10E). This competition revealed a tendency for memories stored in larger engrams to be retrieved more frequently, highlighting the potential limitations of global inhibition in managing engram competition.

Our findings demonstrate that selective inhibition provides a robust mechanism for pattern completion across various conditions that affect engram competition. It enhances retrieval performance regardless of input strength disparity, mitigates the negative effects of overlap between engrams, maintains higher performance levels as the number of competing memories increases, and effectively manages size differences between engrams. These results underscore the functional advantages of selective inhibition over global inhibition in reliable memory retrieval. They support our hypothesis that connections from excitatory to inhibitory neurons can be learned during encoding via heterosynaptic plasticity and suggest a potential role for this mechanism in managing complex memory dynamics within the hippocampus.

2.6 Neuronal representations across hippocampal subregions during pattern separation and pattern completion task.

We expanded our investigation to focus on pattern separation across the entire hippocampal model, building on our observations of stable pattern completion under various conditions facilitated by selective inhibition. Previous studies have shown that distinct hippocampal subregions exhibit different patterns of neuronal activation (Yassa and Stark, 2011; Knierim and Neunuebel, 2016). To explore this, we conducted 105 trials, varying the similarity between inputs A and B (Figures 11A and 11B). Input A was presented during a single encoding phase lasting 120 ms, while input B was presented over 7.2 seconds across 30 phases, encompassing both encoding and retrieval (Figure 11A).

To quantify the discrimination between neuronal patterns A and B, we averaged the discrimination values across 30 phases. Our analysis revealed distinct computational characteristics for each hippocampal subregion. The DG exhibited robust pattern separation, effectively distinguishing input patterns, particularly when input dissimilarity (Δ input) exceeded about 0.2. In contrast, CA1 showed a linear relationship between input and output discrimination, indicating a more gradual differentiation process (Figure 11C). This finding aligns with evidence suggesting that the DG has a unique capacity to discriminate between similar inputs (Neunuebel and Knierim, 2014; Berron et al., 2016). In the CA3 region, Δ output tended to be lower than in CA1 when Δ input was below about 0.2, indicating reduced efficiency in pattern separation for highly similar inputs. However, as Δ input increased beyond 0.2, CA3's Δ output surpassed that

Figure 10: The effect of the size of engrams in CA3 on retrieval. **(A)** Schematic of the retrieval cue with engrams of different sizes. **(B)** Example of the learned Rc weight matrix in the task where engrams have varying sizes. **(C)** Firing rate over time for neural assemblies in CA3 during retrieval. **(C1)** and **(C2)** show results for the same cue with the same learned samples. **(C1)** Example when A is retrieved. **(C2)** Example when B is retrieved. **(D)** RRetrieval performance as a function of engram size differences, comparing selective inhibition (blue) and global inhibition (red). Performance is measured as the mean retrieval success rate, averaged over 15 samples for each size difference. Each sample's success rate is averaged over 50 repetitions. **(E)** Retrieval dominance of memories stored in larger engrams across different size differences, comparing selective inhibition (blue) and global inhibition (red). Retrieval dominance indicates the percentage of times the memory stored in the larger engram is retrieved.

of CA1, suggesting enhanced separation capabilities for more distinct inputs (Figure 11C). These results are consistent with previous studies (Lee et al., 2004; Leutgeb et al., 2004; Guzowski et al., 2004; Yassa and Stark, 2011; Knierim and Neunuebel, 2016).

We investigated these Δ output patterns by separating the encoding and retrieval phases to understand the reasons for these behaviors. During encoding, CA3 closely mirrored the pattern observed in the DG, while CA1 displayed a more linear response (Figure 11D). This similarity arises because CA3 receives input from the DG during the encoding phase in the model. At Δ input levels below approximately 0.2, the reduced output discrimination in CA3 is attributed to a failure of pattern separation in the DG, where the model incorrectly perceived two distinct inputs as identical. This finding explains the lower output discrimination in CA3 compared to CA1, as shown in Figure 11C.

During retrieval, the pattern in CA3 more closely resembled that of CA1 than it did during encoding, consistent with the model's design, where CA3 transmits information to CA1 (Figure 11E). Notably, the Δ output of CA3 during retrieval was lower than during encoding, despite input B being provided as an intact cue. This resulted in CA3 appearing to have lower pattern separation performance than the DG in Figure 11C, even though CA3 retained the outcomes of the DG's pattern separation during encoding (Figure 11D). This effect was observed even at high Δ outputs (around

Figure 11: **(A)** Schematic of the pattern separation task. Input A is presented to the model during a single encoding phase (120 ms), while Input B is presented over 7.2 seconds (30 phases), encompassing both encoding and retrieval phases. Input B's similarity to Input A is varied across 105 trials to evaluate the model's pattern separation ability. **(B)** Examples of learned Rc weight matrices for **(B1)** highly similar (Δ Input = 0.92) and **(B2)** less similar (Δ Input = 0.17) input patterns. **(C)** Neural representation changes across subregions in the model as a function of input discrimination (Δ Input). The Δ outputs are averaged over 30 phases. **(D)** Neural representation changes across subregions in the model as a function of input discrimination (Δ Input) during the encoding phase. **(E)** Neural representation changes across subregions in the model as a function of input discrimination (Δ Input) during the retrieval phase.

0.8), where input B did not consistently converge to its own representation due to the competing influence of pattern completion involving input A (Figure 11E).

To gain deeper insights into the dynamics of pattern completion, we designed a specific task to assess this process. We investigated how varying degrees of similarity between retrieval cues and stored memories impact retrieval accuracy. Specifically, we used an input pair with zero similarity (A and B) as the stored memories. During retrieval, we introduced cues with graded similarities to inputs A and B, maintaining a linear proportional relationship between each cue's similarity to A and B (Figure 12A). For example, if a cue was 70% similar to A, it would be 30% similar to B. This linear progression enables a more precise analysis of the transition between retrieving different memories compared to arbitrary cue selection.

The results demonstrated a strong linear relationship between cue similarity and retrieval accuracy. Given that inputs A and B were entirely dissimilar, both CA3 and CA1 exhibited highly similar linear response patterns. Notably, the cues did not converge on a dominant memory, even when there was a small degree of similarity to one of the inputs. This finding suggests that the hippocampal model's pattern completion mechanism is linearly influenced by the degree of similarity between the retrieval cue and the stored memory (Figures 12C and 12D). The linear relationship indicates that even minimally similar cues can lead to accurate memory retrieval, highlighting the model's sensitivity and precision in pattern completion. These insights into the specific mechanisms of pattern completion within our hippocampal model deepen our understanding of memory processes.

Figure 12: **(A)** Schematic of the pattern completion task. During the encoding phase (120 ms), inputs A and B are presented to the model with a similarity of 0. In the retrieval phase, 100 cues are presented, each having a linearly proportional relationship in their similarity to A and B, ensuring that the sum of these similarities equals 1. Each cue is presented for 7.2 seconds, and 30 retrieval phases are observed without further learning. **(B)** The learned Rc weight matrix after training with inputs A and B. **(C)** Retrieval performance as a function of the similarity between input A and the cue (Δ A$-$Cue). Performance is measured by the probability of retrieving Input A (blue) or Input B (purple) in response to the presented cue across 30 retrieval phases. **(D)** Neural representation changes relative to the discrimination between input A and the cues (Δ A$-$Cue) during retrieval. The Δ outputs are averaged over 30 phases.

3 Discussion

We developed a spiking neural network model based on theta oscillation theory that captures the structural and connectivity properties of the hippocampus. In our exploration of memory processing with this model, we focused on the contributions of inhibitory neurons in CA3. Our findings suggest that the activation of these neurons, along with heterosynaptic plasticity during encoding, can lead to selective inhibition during retrieval. Thus, the role of inhibitory neurons in CA3 includes both selective inhibition and sparse distributed coding. In the proposed model, inhibitory neurons associated with specific neural assemblies in CA3 selectively suppress excitatory neurons involved in competing assemblies.

Interestingly, inhibitory neurons do not suppress excitatory neurons within the same engram. This occurs because only excitatory neurons that are not suppressed by inhibitory neurons survive to form neural assemblies. These roles of inhibitory neurons in memory processing are supported by experimental evidence, including (1) feedforward inhibition of the dentate gyrus via mossy fibers, (2) plasticity within inhibitory neurons, (3) heterosynaptic plasticity across the hippocampus, and (4) the segregation of encoding and retrieval phases by theta oscillation.

3.1 Model validation for memory formation

During encoding in the model, inhibitory neurons in CA3 receive inputs from the DG via mossy fibers. These inhibitory neurons then suppress most excitatory neurons in CA3, determining which neurons remain active. Some observations from the literature, although not all fully specific to the hippocampus, align with the proposed model: Granule cells in

the DG are known to recruit feedforward inhibition via mossy fibers (Ruediger et al., 2011; Guo et al., 2018; Viana da Silva et al., 2019). Parvalbumin-positive (PV+) inhibitory neurons in the hippocampus are crucial for memory storage (Korotkova et al., 2010; He et al., 2021; Raven and Aton, 2021). Inhibitory neurons influence the size of the engram (Morrison et al., 2016; Stefanelli et al., 2016; Josselyn and Tonegawa, 2020; Giorgi and Marinelli, 2021). These experimental findings underscore the importance of inhibitory neurons in memory formation, indicating that the model effectively captures their functional and structural roles.

In the model, inhibitory neurons have stimulus selectivity because the DG processes and separates external inputs before transmitting them to CA3. This feature parallels the stimulus selectivity observed primarily in inhibitory interneurons within the visual and auditory cortices (Li et al., 2014; Khan et al., 2018; Cummings et al., 2021), implying that similar mechanisms may exist in the hippocampus. In this context, inhibitory neurons in sensory areas play a more active role in information processing than previously thought. In line with these findings, the model supports the view that inhibitory neurons in the hippocampus actively participate in both encoding and retrieval processes while exhibiting stimulus selectivity.

Observations of plasticity in inhibitory neurons indicate their synaptic interactions with excitatory neurons (Pelletier and Lacaille, 2008; Korotkova et al., 2010; Perez et al., 2001; Owen et al., 2013; Cohen et al., 2016; He et al., 2021). In the model, this plasticity enables inhibitory neurons to modulate their synaptic strengths with excitatory neurons, thereby facilitating selective inhibition during engram competition. Selective inhibition positions inhibitory neurons as active participants in retrieval processes, underscoring their potential contribution to pattern completion.

To effectively harness the properties of inhibitory neurons mentioned above, the model employs two key mechanisms: theta oscillation theory (Hasselmo and Stern, 2014; Kragel et al., 2020; Reddy et al., 2021; Ter Wal et al., 2021; Lowet et al., 2023; Gattas et al., 2023) and heterosynaptic plasticity (Tsukamoto et al., 2003; Hunt et al., 2013; Chater and Goda, 2021; Jenks et al., 2021). Theta oscillation theory posits that distinct phases of the theta rhythm modulate different information flows in the hippocampus, with encoding occurring at the trough and retrieval at the peak. The hippocampus maintains these distinct phases through phasic changes in membrane potential (Hasselmo and Stern, 2014; Malezieux et al., 2020).

In the model, reflecting these phenomena, synaptic activity related to retrieval is suppressed during encoding. During engram formation in CA3, heterosynaptic plasticity strengthens connections between inhibitory and excitatory neurons, even in the absence of direct stimulation. These mechanisms enable inhibitory neurons to play an active role in both sparse distributed coding for pattern separation and selective inhibition for pattern completion.

Proposed model illustrates the rapid process of hippocampal memory formation, completed in just 120 milliseconds. This rapid storage capability reinforces the model's functional plausibility by effectively replicating the rapid memory formation observed in the hippocampus (Moser and Moser, 2003; Lee et al., 2015; Liao and Losonczy, 2024). Moreover, the model demonstrated stable pattern completion across various realistic retrieval scenarios in the hippocampus, including those with evenly distributed input cues, overlapping engrams, multiple competing engrams, and engrams of varying sizes. This stability suggests that the model effectively captures key aspects of how the hippocampus retrieves memories under different engram configurations.

3.2 Alignment with cognitive evidence

The enhanced stability of the model, driven by selective inhibition, allowed us to investigate firing patterns across hippocampal subregions during pattern separation and completion by quantitatively controlling input similarity. The distinct functions of hippocampal subregions during pattern separation and completion are well-documented in the literature (Yassa and Stark, 2011; Knierim and Neunuebel, 2016). The DG is particularly robust in supporting pattern separation (Leutgeb et al., 2007; Santoro, 2013; Neunuebel and Knierim, 2014; Berron et al., 2016). In CA1, output discrimination is thought to be linearly related to input discrimination, meaning that output differences increase at a constant rate as input differences increase (Guzowski et al., 2004; Yassa and Stark, 2011; Knierim and Neunuebel, 2016). CA3, however, exhibits more complex dynamics because it participates in both pattern separation and pattern completion. Some studies suggest that CA3 is more involved in pattern separation, while others emphasize its greater role in pattern completion. (Leutgeb et al., 2004; Lee et al., 2004; Guzowski et al., 2004).

The model successfully replicates the distinct roles of hippocampal subregions, demonstrating strong pattern separation in the DG and a consistent linear response in CA1. In CA3, the model aligns with the evidence discussed earlier, displaying either pattern separation or pattern completion depending on the degree of input similarity. We were able to effectively investigate CA3's behavior by clearly separating the encoding and retrieval phases. When the difference between inputs is small, the difference between output pairs in CA3 is reduced compared to CA1. This occurs because pattern separation in the DG does not reach the discrimination threshold during encoding. As a result, the model interprets the two similar inputs as identical and performs pattern completion.

When the difference between inputs is large, CA3 shows less pattern separation than the DG, but still more than CA1. This reduced separation is likely due to mixed outcomes, where two completely different memories are retrieved. In other words, a cue might still trigger pattern completion for less similar memories, even when it closely resembles

a specific memory. This is supported by the observation that retrieval probability depends linearly on the similarity between the cue and the memory. Such linearity prevents the dominance of biased memories during retrieval, partially capturing the complexity of real-world information. Because the proposed model aligns with established experimental data, it may serve as a valuable tool for understanding hippocampal functions. By examining phase-specific dynamics in the model, we gain new insights into how the hippocampus handles conflicting information.

3.3 Limitations and future works

The proposed model has several limitations that should be addressed in future research. One key aspect is the assumption that heterosynaptic plasticity in the connections from excitatory to inhibitory neurons in CA3 drives selective inhibition. Our model suggests that such plasticity could be pivotal for selective inhibition. However, conclusive evidence for heterosynaptic plasticity at these specific connections is still lacking, underscoring the need for further experimental investigations to validate this mechanism.

By simplifying the DG to focus solely on spatial pattern separation with a fixed input activation frequency, we overlook its established roles in temporal and rate pattern separation (Pernía-Andrade and Jonas, 2014; Madar et al., 2019; Löffler and Gupta, 2022). As a result of this simplification, certain details, such as the number of neurons in each subregion, connection probabilities between subregions, and synaptic weights, do not fully align with biological specifics. Since these differences may impact the memory retrieval process, the observed linear relationship between similarity and retrieval probability in the pattern completion task should be interpreted with caution, warranting further experimental validation and larger-scale modeling.

We limited CA1's role to reinstatement of original information from CA3 engrams, which overlooks the neuronal diversity and complex functions of CA1 (Masurkar, 2018; Jeong and Singer, 2022). The model disregards cellular-level processes, feedback connections, and molecular signaling pathways, such as those mediated by cAMP response element-binding protein (CREB) (Pelletier and Lacaille, 2008; Josselyn and Tonegawa, 2020; Bickle, 2021). Since these elements are key components of memory processes, their omission may limit the quantitative accuracy of the model's outcomes.

In future work, we will extend the model to incorporate sequential learning, exploring how inhibitory neurons contribute to engram formation and competition across sequences of information. This extension could deepen our understanding of hippocampal function in processing sequences, offering new insights into cognitive processes such as replay, sequence compression, and predictive coding.

3.4 Conclusion

Our research highlights the active and dynamic role of inhibitory neurons in CA3 engram formation and stable pattern completion. This work provides a fresh perspective on established hippocampal functions and opens new avenues for experimental investigation. As our understanding of hippocampal circuitry and function continues to evolve, models like ours will be essential in bridging the gap between cellular mechanisms and cognitive phenomena, advancing research in both neuroscience and artificial intelligence.

4 Methods and Materials

4.1 Model structure

We constructed a spiking neural network model of the hippocampal trisynaptic circuit (superficial EC to CA1 via DG and CA3) to investigate memory formation and competition dynamics. The model preserves key pathways, including the mossy fiber projections from DG and the direct PP from superficial EC. We focused solely on spatial pattern processing, excluding temporal sequences for simplicity. The model's input layer consists of 16 neurons, each firing at a fixed rate of 50 Hz. This configuration allows for 2^{16} unique spatial patterns, providing sufficient variability to assess the model's performance while maintaining computational feasibility.

Table 1: Parameters of Izhikevich model for excitatory neurons and inhibitory neurons.

	C	k	V_{rest}	V_{th}	V_{peak}	a	d	c	d
Exc	80	3	-60	-50	50	0.01	5	-60	10
Inh	20	1	-55	-40	25	0.15	8	-55	200

Given our focus on spatial patterns, we considered only spatial pattern separation in DG, omitting temporal and rate separation mechanisms. This simplification necessarily leads to deviations from biological details, such as the number of neurons in each subregion, connection probabilities between subregions, and synaptic weights (Tables 2 and 3). The parameters were derived from the study by Chen (2017) and were optimized with significant modifications through iterative manual adjustments to ensure the model's proper functioning. We initialized and fixed connection weights with specific values for each connection during learning to maintain connection probabilities. Rather than adjusting these connection weights, we modified the peak conductance, which is analogous to conventional connection weights. These were initialized to 0, except for connections from inhibitory neurons to excitatory neurons in CA3, to establish global inhibition as the default state (detailed explanation in the next section).

The CA1 region in the model is simplified, consisting only of excitatory neurons. While this simplification omits some of the intricate microcircuitry found in biological CA1 (Masurkar, 2018; Jeong and Singer, 2022), it preserves the crucial function of interfacing between CA3 and EC, facilitating the reinstatement of original information by CA3 engrams. We designed the CA1-EC connections so that each CA1 neuron maintains consistent bidirectional connections with specific EC neurons. Although this arrangement is not strictly biological, it enables CA3 engrams to effectively encode and decode information between hippocampal and cortical representations.

We implemented the model using Python with NumPy. The following sections detail the specific neuron models, learning rules, and analysis methods used to investigate memory dynamics within this structure.

4.2 Single neuron models

Each neuron was described using the Izhikevich model (Izhikevich and Edelman, 2008) following as

$$C\dot{v} = k(v - v_r)(v - v_t) - u - I_{syn} \tag{1}$$

$$\dot{u} = a\{b(v - v_r) - u\} \text{ if } v \geq 30mV, \text{ then } v = c, u = u + d \tag{2}$$

where C is the membrane capacitance, v is the membrane potential, v_r is the resting potential, v_t is the threshold potential, and u is the recovery variable (the difference of all inward and outward voltage-gated currents) (Izhikevich and Edelman, 2008). I_{syn} is the summation of synaptic currents from other neurons, detailed in the next section. Each parameter, including the dimensionless values k, a, b, c, and d, follows the values specified by Izhikevich and Edelman (2008) for different neuron types. Based on this paper, the excitatory neurons use the parameters for pyramidal cells, and the inhibitory neurons use those for basket interneurons (Table 1).

4.3 Synapse models

The total synaptic current model also follows (Izhikevich and Edelman, 2008; Chen, 2017), which consists of four basic conductances from AMPA, NMDA, GABA$_A$, and GABA$_B$ channels with the simplified voltage-gating of the NMDA channel ($[[(v + 80)/60]^2/[1 + ((v + 80)/60)^2]$).

Table 2: Parameters for neurons organizing each layer.

	N	τ_{AMPA}	τ_{NMDA}	τ_{GABA_A}	τ_{GABA_B}	W_{Be}	W_{Bi}
superficial EC	16	5	-	-	-	5	-
deep EC	16	5	-	-	-	5	-
Hilus(Exc)	100	5	-	15	-	5	10
Hilus(Inh)	16	5	-	-	-	5	-
GCL(Exc)	800	5	-	15	-	5	10
GCL(Inh)	400	5	-	-	-	7	7
CA3(Exc)	2400	5	30	8	30	4, 0.75, 20	10
CA3(Inh)	120	5	30	8	30	5	10
CA1	200	5	30	-	-	4	-

$$I_{syn} = g_{AMPA}(v - 0) + g_{NMDA}\frac{\left[(v+80)/60\right]^2}{1 + \left[(v+80)/60\right]^2}(v - 0) + g_{GABA_A}(v + 70) + g_{GABA_B}(v + 90) \quad (3)$$

$$\begin{cases} g = g + q & \text{if } v \geq 30mV \\ \dfrac{dg(t)}{dt} = -\dfrac{1}{\tau}g(t) & \text{otherwise} \end{cases} \quad (4)$$

where the reversal potentials of AMPA and NMDA channels are 0 mV and reversal potentials for GABA$_A$ and GABA$_B$ channels are -70 mV and -90 mV respectively. The ratio of NMDA to AMPA receptors was set to 5:5, and the ratio of GABA$_A$ to GABA$_B$ receptors was set to 9:1 for all of the neurons in CA3 (Chen et al., 2013). For simplification, the synapse models of neurons in the DG included only AMPA and GABA$_A$ receptors because learning does not occur in this region of the model (Table 2).

Synaptic conductance decreases exponentially with different time constants and increases by the peak conductance q upon the arrival of an action potential at the synapse (Chen and Wang, 2007). The total excitatory and inhibitory synaptic currents were limited to maximum values of W_{Be} and W_{Bi} respectively (Table 2). For excitatory neurons in CA3, the total synaptic current was differentiated based on the region of origin of the sending neuron—superficial EC, DG, or CA3—to preserve the distinct influences of each region. Additionally, some connections featured delayed transmission times to the postsynaptic neuron to ensure stable learning (Table 3).

4.4 STDP learning

Learning was implemented using symmetric STDP (Mishra et al., 2016; Ecker et al., 2022). Instead of directly adjusting the synaptic weights W, we updated the peak conductance q to maintain the connectivity ratio within the network. Synaptic weights, which are typically involved in scaling the current, were initialized and held constant throughout the learning process to preserve the network's initial structure and stability (Chen and Wang, 2007). This approach

Table 3: Parameters for connections between layers.

	p_{conn}	W	t_d
Superficial EC→Hilus (Exc)	0.3125	3	-
Superficial EC→Hilus (Inh)	One-to-One	3	-
Superficial EC→GCL (Exc)	0.0625	2	10
Superficial EC→GCL (Inh)	0.5	4	-
Superficial EC→CA3 (Exc)	0.25	2	-
Superficial EC→Deep EC	One-to-One	3	-
Deep EC→CA1	0.125	0.75	-
Hilus (Inh)→Hilus (Exc)	0.0625	4	-
Hilus (Exc)→GCL (Inh)	0.5	4	-
GCL (Inh)→GCL (Exc)	0.0013	3	-
GCL (Exc)→CA3 (Exc)	0.0125	2	3
GCL (Exc)→CA3 (Inh)	0.025	2	-
CA3 (Exc)→CA3 (Exc)	0.25	1	5
CA3 (Exc)→CA3 (Inh)	0.25	1	-
CA3 (Exc)→CA1	0.5	0.45	-
CA3 (Inh)→CA3 (Exc)	0.25	5	-
CA1→Deep EC	0.125	2	15

ensures that the modifications introduced by STDP do not disrupt the overall connectivity but instead refine the existing connections. (Table 3).

$$q = q + F(\Delta t), \ F(\Delta t) = \begin{cases} A_+ \exp\left(-\dfrac{\Delta t}{\tau_+}\right) & \text{at } t_{post}, \text{ if } t_{pre} < t_{post} \\ A_- \exp\left(\dfrac{\Delta t}{\tau_-}\right) & \text{at } t_{pre}, \text{ if } t_{post} < t_{pre} \end{cases} \tag{5}$$

where $\Delta t = t_{post} - t_{pre}$ represents the time difference between action potentials, and $A_\pm$ determines the maximum amount of synaptic modification, which decays exponentially with time constants $\tau_\pm$. The peak conductance is constrained within the range $[0, q_{max}]$, where $q_{max} = 3$ nS. Given the symmetric nature of the STDP rule adopted, $A_\pm$ was set to 0.2 nS, and $\tau_\pm$ was set to 62.5 ms.

4.5 Similarity analysis

Our model focuses exclusively on spatial input patterns and spatial pattern separation. Therefore, the similarity between firing patterns was assessed by calculating the spatial overlap, as suggested by Löffler and Gupta (2022).

$$O_{ij} = \frac{n_i}{n_i + n_j} \tag{6}$$

where n_i represents the number of activated neurons in pattern i, and n_{ij} denotes the number of neurons activated in both patterns i and j. The discrimination value Δ between patterns i and j is calculated as $1 - O_{ij}$. The firing pattern is determined based on neuron activation, except for the case of CA3. For CA3 patterns during retrieval, only neurons firing at rates exceeding 25 Hz are considered, ensuring that only significant activity is captured and noise is effectively filtered out. This threshold helps to isolate meaningful firing patterns that contribute to memory retrieval, thereby enhancing the accuracy of the pattern analysis.

References

Acsády, L., Kamondi, A., Sík, A., Freund, T., and Buzsáki, G. (1998). Gabaergic cells are the major postsynaptic targets of mossy fibers in the rat hippocampus. *Journal of Neuroscience*, 18(9):3386–3403.

Amaral, D. G., Scharfman, H. E., and Lavenex, P. (2007). The dentate gyrus: fundamental neuroanatomical organization (dentate gyrus for dummies). *Progress in Brain Research*, 163:3–22.

Berron, D., Schütze, H., Maass, A., Cardenas-Blanco, A., Kuijf, H. J., Kumaran, D., and Düzel, E. (2016). Strong evidence for pattern separation in human dentate gyrus. *Journal of Neuroscience*, 36(29):7569–7579.

Bickle, J. (2021). The first two decades of creb-memory research: data for philosophy of neuroscience. *AIMS Neuroscience*, 8(3):322–339.

Bragin, A., Jando, G., Nadasdy, Z., Hetke, J., Wise, K., and Buzsaki, G. (1995). Gamma (40-100 hz) oscillation in the hippocampus of the behaving rat. *Journal of Neuroscience*, 15(1):47–60.

Cayco-Gajic, N. A., Clopath, C., and Silver, R. A. (2017). Sparse synaptic connectivity is required for decorrelation and pattern separation in feedforward networks. *Nature Communications*, 8:1116.

Chater, T. E. and Goda, Y. (2021). My neighbour hetero—deconstructing the mechanisms underlying heterosynaptic plasticity. *Current Opinion in Neurobiology*, 67:106–114.

Chavlis, S., Petrantonakis, P. C., and Poirazi, P. (2017). Dendrites of dentate gyrus granule cells contribute to pattern separation by controlling sparsity. *Hippocampus*, 27(1):89–110.

Chen, K. and Wang, L. (2007). *Trends in Neural Computation*. Studies in Computational Intelligence. Springer Nature.

Chen, Y. (2017). Mechanisms of winner-take-all and group selection in neuronal spiking networks. *Frontiers in Computational Neuroscience*, 11.

Chen, Y., Mckinstry, J. L., and Edelman, G. M. (2013). Versatile networks of simulated spiking neurons displaying winner-take-all behavior. *Frontiers in Computational Neuroscience*, 7.

Cheng, Q. and Yakel, J. L. (2015). The effect of 7 nicotinic receptor activation on glutamatergic transmission in the hippocampus. *Biochemical Pharmacology*, 97(4):439–444.

Cohen, S. M., Ma, H., Kuchibhotla, K. V., Watson, B. O., Buzsáki, G., Froemke, R. C., and Tsien, R. W. (2016). Excitation-transcription coupling in parvalbumin-positive interneurons employs a novel cam kinase-dependent pathway distinct from excitatory neurons. *Neuron*, 90(2):292–307.

Colgin, L. L. (2016). Rhythms of the hippocampal network. *Nature Reviews Neuroscience*, 17(4):239–249.

Cosgrove, K. E., Galván, E. J., Meriney, S. D., and Barrionuevo, G. (2010). Area ca3 interneurons receive two spatially segregated mossy fiber inputs. *Hippocampus*, 20(9):1003–1009.

Cummings, K. A., Lacagnina, A. F., and Clem, R. L. (2021). Gabaergic microcircuitry of fear memory encoding. *Neurobiology of Learning and Memory*, 184.

Danielson, N. B., Turi, G. F., Ladow, M., Chavlis, S., Petrantonakis, P. C., Poirazi, P., and Losonczy, A. (2017). In vivo imaging of dentate gyrus mossy cells in behaving mice. *Neuron*, 93(3):552–559.e4.

Deco, G. and Rolls, E. T. (2006). Decision-making and weber's law: a neurophysiological model. *European Journal of Neuroscience*, 24(3):901–916.

dos Santos, V. L., van de Ven, G. M., Morley, A., Trouche, S., Campo-Urriza, N., and Dupret, D. (2018). Parsing hippocampal theta oscillations by nested spectral components during spatial exploration and memory-guided behavior. *Neuron*, 100(4):940–952.e7.

Drieu, C. and Zugaro, M. (2019). Hippocampal sequences during exploration: Mechanisms and functions. *Frontiers in Cellular Neuroscience*, 13.

Ecker, A., Bagi, B., Vértes, E., Steinbach-Németh, O., Karlócai, M. R., Papp, O. I., Miklós, I., Hájos, N., Freund, T. F., Gulyás, A. I., and Káli, S. (2022). Hippocampal sharp wave-ripples and the associated sequence replay emerge from structured synaptic interactions in a network model of area ca3. *eLife*, 11:e71850.

Faghihi, F. and Moustafa, A. A. (2015). A computational model of pattern separation efficiency in the dentate gyrus with implications in schizophrenia. *Frontiers in Systems Neuroscience*, 9:42.

Freund, T. F. and Buzsáki, G. (1996). Interneurons of the hippocampus. *Hippocampus*, 6(4):347–470.

Gattas, S., Larson, M. S., Mnatsakanyan, L., Sen-Gupta, I., Vadera, S., Swindlehurst, A. L., Rapp, P. E., Lin, J. J., and Yassa, M. A. (2023). Theta mediated dynamics of human hippocampal-neocortical learning systems in memory formation and retrieval. *Nature Communications*, 14(1):8505.

Gedankien, T., Tan, R. J., Qasim, S. E., Moore, H., McDonagh, D., Jacobs, J., and Lega, B. (2023). Acetylcholine modulates the temporal dynamics of human theta oscillations during memory. *Nature Communications*, 14(1):5283.

Giorgi, C. and Marinelli, S. (2021). Roles and transcriptional responses of inhibitory neurons in learning and memory. *Frontiers in Molecular Neuroscience*, 14.

Guo, N., Soden, M. E., Herber, C., Kim, M. T., Besnard, A., Lin, P., Ma, X., Cepko, C. L., Zweifel, L. S., and Sahay, A. (2018). Dentate granule cell recruitment of feedforward inhibition governs engram maintenance and remote memory generalization. *Nature Medicine*, 24(4):438–449.

Guzman, S. J., Schlögl, A., Espinoza, C., Zhang, X., Suter, B. A., and Jonas, P. (2021). How connectivity rules and synaptic properties shape the efficacy of pattern separation in the entorhinal cortex–dentate gyrus–ca3 network. *Nature Computational Science*, 1(12):830–842.

Guzman, S. J., Schlögl, A., Frotscher, M., and Jonas, P. (2016). Synaptic mechanisms of pattern completion in the hippocampal ca3 network. *Science*, 353(6304):1117–1123.

Guzowski, J. F., Knierim, J. J., and Moser, E. I. (2004). Ensemble dynamics of hippocampal regions ca3 and ca1. *Neuron*, 44(4):581–584.

Hainmueller, T. and Bartos, M. (2020). Dentate gyrus circuits for encoding, retrieval and discrimination of episodic memories. *Nature Reviews Neuroscience*, 21:153–168.

Hasselmo, M. E. and Stern, C. E. (2014). Theta rhythm and the encoding and retrieval of space and time. *NeuroImage*, 85(Pt 2):656–666.

He, X., Li, J., Zhou, G., Hu, H., Duan, S., and Ma, H. (2021). Gating of hippocampal rhythms and memory by synaptic plasticity in inhibitory interneurons. *Neuron*, 109(6):1013–1028.e9.

Herweg, N. A., Solomon, E. A., and Kahana, M. J. (2020). Theta oscillations in human memory. *Trends in Cognitive Sciences*, 24(3):208–227.

Hopfield, J. J. (1982). Neural networks and physical systems with emergent collective computational abilities. *Proceedings of the National Academy of Sciences of the United States of America*, 79(8):2554–2558.

Hunt, D. L., Puente, N., Grandes, P., and Castillo, P. E. (2013). Bidirectional nmda receptor plasticity controls ca3 output and heterosynaptic metaplasticity. *Nature Neuroscience*, 16:1049–1059.

Izhikevich, E. M. and Edelman, G. M. (2008). Large-scale model of mammalian thalamocortical systems. *Proceedings of the National Academy of Sciences of the United States of America*, 105(9):3593–3598.

Jenks, K. R., Tsimring, K., Ip, J. P. K., Zepeda, J. C., and Sur, M. (2021). Heterosynaptic plasticity and the experience-dependent refinement of developing neuronal circuits. *Frontiers in Neural Circuits*, 15.

Jeong, N. and Singer, A. C. (2022). Learning from inhibition: Functional roles of hippocampal ca1 inhibition in spatial learning and memory. *Current Opinion in Neurobiology*, 76:102604.

Ji, L. (2023). The validity of the specific coding principle: The explanations from cognitive psychology and psycho-analysis. *Journal of Education, Humanities and Social Sciences*, 8:1259–1263.

Jinde, S., Zsiros, V., and Nakazawa, K. (2013). Hilar mossy cell circuitry controlling dentate granule cell excitability. *Frontiers in Neural Circuits*, 7.

Josselyn, S. A. and Tonegawa, S. (2020). Memory engrams: Recalling the past and imagining the future. *Science*, 367(6473):eaaw4325.

Káli, S. and Dayan, P. (2000). The involvement of recurrent connections in area ca3 in establishing the properties of place fields: a model. *Journal of Neuroscience*, 20(19):7463–7477.

Khan, A. G., Poort, J., Chadwick, A., Blot, A., Sahani, M., Mrsic-Flogel, T. D., and Hofer, S. B. (2018). Distinct learning-induced changes in stimulus selectivity and interactions of gabaergic interneuron classes in visual cortex. *Nature Neuroscience*, 21(6):851–859.

Kim, S.-Y. and Lim, W. (2022). Disynaptic effect of hilar cells on pattern separation in a spiking neural network of hippocampal dentate gyrus. *Cognitive Neurodynamics*, 16(6):1427–1447.

Knierim, J. J. and Neunuebel, J. P. (2016). Tracking the flow of hippocampal computation: Pattern separation, pattern completion, and attractor dynamics. *Neurobiology of Learning and Memory*, 129:38–49.

Korotkova, T., Fuchs, E. C., Ponomarenko, A., von Engelhardt, J., and Monyer, H. (2010). Nmda receptor ablation on parvalbumin-positive interneurons impairs hippocampal synchrony, spatial representations, and working memory. *Neuron*, 68(3):557–569.

Kragel, J. E., VanHaerents, S., Templer, J. W., Schuele, S., Rosenow, J. M., Nilakantan, A. S., and Bridge, D. J. (2020). Hippocampal theta coordinates memory processing during visual exploration. *eLife*, 9:e52108.

Lee, I., Yoganarasimha, D., Rao, G., and Knierim, J. J. (2004). Comparison of population coherence of place cells in hippocampal subfields ca1 and ca3. *Nature*, 430(6998):456–459.

Lee, S. W., O'Doherty, J. P., and Shimojo, S. (2015). Neural computations mediating one-shot learning in the human brain. *PLOS Biology*, 13(4):1–36.

Leutgeb, J. K., Leutgeb, S., Moser, M.-B., and Moser, E. I. (2007). Pattern separation in the dentate gyrus and ca3 of the hippocampus. *Science*, 315(5814):961–966.

Leutgeb, S., Leutgeb, J. K., Treves, A., Moser, M.-B., and Moser, E. I. (2004). Distinct ensemble codes in hippocampal areas ca3 and ca1. *Science*, 305(5688):1295–1298.

Li, L.-y., Xiong, X. R., Ibrahim, L. A., Yuan, W., Tao, H. W., and Zhang, L. I. (2014). Differential receptive field properties of parvalbumin and somatostatin inhibitory neurons in mouse auditory cortex. *Cerebral Cortex*, 25(7):1782–1791.

Liao, Z. and Losonczy, A. (2024). Learning, fast and slow: Single- and many-shot learning in the hippocampus. *Annual Review of Neuroscience*, 47(2):187–209.

Lowet, E., Sheehan, D. J., Chialva, U., De Oliveira Pena, R., Mount, R. A., Xiao, S., Zhou, S. L., an Tseng, H., Gritton, H., Shroff, S., Kondabolu, K., Cheung, C., Wang, Y., Piatkevich, K. D., Boyden, E. S., Mertz, J., Hasselmo, M. E., Rotstein, H. G., and Han, X. (2023). Theta and gamma rhythmic coding through two spike output modes in the hippocampus during spatial navigation. *Cell Reports*, 42(8):112906.

Löffler, H. and Gupta, D. S. (2022). A model of pattern separation by single neurons. *Frontiers in Computational Neuroscience*, 16.

Madar, A. D., Ewell, L. A., and Jones, M. V. (2019). Pattern separation of spiketrains in hippocampal neurons. *Scientific Reports*, 9:5282.

Malezieux, M., Kees, A. L., and Mulle, C. (2020). Theta oscillations coincide with sustained hyperpolarization in ca3 pyramidal cells, underlying decreased firing. *Cell Reports*, 32(1):107868.

Marr, D. (1971). Simple memory: A theory for archicortex. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences*, 262:23–81.

Masurkar, A. V. (2018). Towards a circuit-level understanding of hippocampal ca1 dysfunction in alzheimer's disease across anatomical axes. *Journal of Alzheimer's Disease & Parkinsonism*, 8(1):412.

McClelland, J. L., McNaughton, B. L., and O'Reilly, R. C. (1995). Why there are complementary learning systems in the hippocampus and neocortex: insights from the successes and failures of connectionist models of learning and memory. *Psychological Review*, 102(3):419–457.

Mishra, R. K., Kim, S., Guzman, S. J., and Jonas, P. (2016). Symmetric spike timing-dependent plasticity at ca3–ca3 synapses optimizes storage and recall in autoassociative networks. *Nature Communications*, 7(1):11552.

Modhej, N., Bastanfard, A., Teshnehlab, M., and Raiesdana, S. (2020). Pattern separation network based on the hippocampus activity for handwritten recognition. *IEEE Access*, 8:212803–212817.

Mori, M., Gähwiler, B. H., and Gerber, U. (2007). Recruitment of an inhibitory hippocampal network after bursting in a single granule cell. *Proceedings of the National Academy of Sciences*, 104(18):7640–7645.

Morrison, D. J., Rashid, A. J., Yiu, A. P., Yan, C., Frankland, P. W., and Josselyn, S. A. (2016). Parvalbumin interneurons constrain the size of the lateral amygdala engram. *Neurobiology of Learning and Memory*, 135:91–99.

Moser, E. I. and Moser, M.-B. (2003). One-shot memory in hippocampal ca3 networks. *Neuron*, 38(2):147–148.

Müller, C. and Remy, S. (2018). Septo-hippocampal interaction. *Cell and Tissue Research*, 373(3):565–575.

Myers, C. E. and Scharfman, H. E. (2009). A role for hilar cells in pattern separation in the dentate gyrus: a computational approach. *Hippocampus*, 19(4):321–337.

Neunuebel, J. P. and Knierim, J. J. (2014). Ca3 retrieves coherent representations from degraded input: direct evidence for ca3 pattern completion and dentate gyrus pattern separation. *Neuron*, 81(2):416–427.

Niell, C. M. and Stryker, M. P. (2008). Highly selective receptive fields in mouse visual cortex. *Journal of Neuroscience*, 28(30):7520–7536.

Norman, K. A. and O'Reilly, R. C. (2003). Modeling hippocampal and neocortical contributions to recognition memory: A complementary-learning-systems approach. *Psychological Review*, 110(4):611–646.

Nuñez, A. and Buño, W. (2021). The theta rhythm of the hippocampus: From neuronal and circuit mechanisms to behavior. *Frontiers in Cellular Neuroscience*, 15.

O'Keefe, J. and Nadel, L. (1978). *The Hippocampus as A Cognitive Map*. Clarendon Press.

Owen, S. F., Tuncdemir, S. N., Bader, P. L., Tirko, N. N., Fishell, G., and Tsien, R. W. (2013). Oxytocin enhances hippocampal spike transmission by modulating fast-spiking interneurons. *Nature*, 500(7463):458–462.

Parkin, A. (2007). Levels of processing and the cue overload principle. *Quarterly Journal of Experimental Psychology - QUART J EXP PSYCHOL*, 32:427–434.

Pelletier, J. G. and Lacaille, J.-C. (2008). Long-term synaptic plasticity in hippocampal feedback inhibitory networks. *Progress in Brain Research*, 169:241–250.

Perez, Y., Morin, F., and Lacaille, J.-C. (2001). A hebbian form of long-term potentiation dependent on mglur1a in hippocampal inhibitory interneurons. *Proceedings of the National Academy of Sciences*, 98(16):9401–9406.

Pernía-Andrade, A. and Jonas, P. (2014). Theta-gamma-modulated synaptic currents in hippocampal granule cells in vivo define a mechanism for network oscillations. *Neuron*, 81(1):140–152.

Prince, L. Y., Bacon, T. J., Tigaret, C. M., and Mellor, J. R. (2016). Neuromodulation of the feedforward dentate gyrus-ca3 microcircuit. *Frontiers in Synaptic Neuroscience*, 8:32.

Raven, F. and Aton, S. J. (2021). The engram's dark horse: How interneurons regulate state-dependent memory processing and plasticity. *Frontiers in Neural Circuits*, 15.

Rebola, N., Carta, M., and Mulle, C. (2017). Operation and plasticity of hippocampal ca3 circuits: implications for memory encoding. *Nature Reviews Neuroscience*, 18(4):208–220.

Reddy, L., Self, M. W., Zoefel, B., Poncet, M., Possel, J. K., Peters, J. C., Baayen, J. C., Idema, S., VanRullen, R., and Roelfsema, P. R. (2021). Theta-phase dependent neuronal coding during sequence learning in human single neurons. *Nature Communications*, 12(1):4839.

Rennó-Costa, C., Lisman, J. E., and Verschure, P. F. M. J. (2014). A signature of attractor dynamics in the ca3 region of the hippocampus. *PLOS Computational Biology*, 10(5):1–15.

Robinson, J., Manseau, F., Ducharme, G., Amilhon, B., Vigneault, E., El Mestikawy, S., and Williams, S. (2016). Optogenetic activation of septal glutamatergic neurons drive hippocampal theta rhythms. *Journal of Neuroscience*, 36(10):3016–3023.

Rolls, E. T. (2007). *Memory, attention, and decision-making: A unifying computational neuroscience approach*. Oxford University Press.

Rolls, E. T. (2013). A quantitative theory of the functions of the hippocampal ca3 network in memory. *Frontiers in Cellular Neuroscience*, 7.

Rolls, E. T. (2016). Pattern separation, completion, and categorisation in the hippocampus and neocortex. *Neurobiology of Learning and Memory*, 129:4–28.

Ruediger, S., Vittori, C., Bednarek, E., and et al. (2011). Learning-related feedforward inhibitory connectivity growth required for memory precision. *Nature*, 473:514–518.

Santoro, A. (2013). Reassessing pattern separation in the dentate gyrus. *Frontiers in Behavioral Neuroscience*, 7.

Semon, R. W., Duffy, B., and Lee, V. (1923). *Revival: Mnemic Psychology*. Routledge.

Stefanelli, T., Bertollini, C., Lüscher, C., Muller, D., and Mendez, P. (2016). Hippocampal somatostatin interneurons control the size of neuronal memory ensembles. *Neuron*, 89(5):1074–1085.

Ter Wal, M., Linde-Domingo, J., Lifanov, J., Roux, F., Kolibius, L. D., Gollwitzer, S., Lang, J., Hamer, H., Rollings, D., Sawlani, V., Chelvarajah, R., Staresina, B., Hanslmayr, S., and Wimber, M. (2021). Theta rhythmicity governs human behavior and hippocampal signals during memory-dependent tasks. *Nature Communications*, 12(1):7048.

Torborg, C. L., Nakashiba, T., Tonegawa, S., and McBain, C. J. (2010). Control of ca3 output by feedforward inhibition despite developmental changes in the excitation-inhibition balance. *Journal of Neuroscience*, 30(46):15628–15637.

Treves, A., Tashiro, A., Witter, M., and Moser, E. (2008). What is the mammalian dentate gyrus good for? *Neuroscience*, 154(4):1155–1172.

Treviño, M., Vivar, C., and Gutiérrez, R. (2011). Excitation-inhibition balance in the ca3 network–neuronal specificity and activity-dependent plasticity. *European Journal of Neuroscience*, 33(10):1771–1785.

Tsukamoto, M., Yasui, T., Yamada, M. K., Nishiyama, N., Matsuki, N., and Ikegaya, Y. (2003). Mossy fibre synaptic nmda receptors trigger non-hebbian long-term potentiation at entorhino-ca3 synapses in the rat. *The Journal of Physiology*, 546:665–675.

Tulving, E. (1983). *Elements of episodic memory*. Oxford University Press.

Tulving, E. and Thomson, D. M. (1973). Encoding specificity and retrieval processes in episodic memory. *Psychological Review*, 80(5):352–373.

Unal, G., Crump, M. G., Viney, T. J., Éltes, T., Katona, L., Klausberger, T., and Somogyi, P. (2018). Spatio-temporal specialization of gabaergic septo-hippocampal neurons for rhythmic network activity. *Brain Structure and Function*, 223(5):2409–2432.

Usrey, W. M., Sceniak, M. P., and Chapman, B. (2003). Receptive fields and response properties of neurons in layer 4 of ferret visual cortex. *Journal of Neurophysiology*, 89(2):1003–1015.

Viana da Silva, S., Zhang, P., Haberl, M. G., Labrousse, V., Grosjean, N., Blanchet, C., Frick, A., and Mulle, C. (2019). Hippocampal mossy fibers synapses in ca3 pyramidal cells are altered at an early stage in a mouse model of alzheimer's disease. *Journal of Neuroscience*, 39(21):4193–4205.

Watkins, M. and Watkins, O. (2013). Cue-overload theory and the method of interpolated attributes. *Bulletin of the Psychonomic Society*, 7:289–291.

Webb, T. J., Rolls, E. T., Deco, G., and Feng, J. (2011). Noise in attractor networks in the brain produced by graded firing rate representations. *PLOS ONE*, 6(9):1–15.

Wittenberg, G. M. and Wang, S. S.-H. (2006). Malleability of spike-timing-dependent plasticity at the ca3–ca1 synapse. *Journal of Neuroscience*, 26(24):6610–6617.

Wixted, J. T., Squire, L. R., Jang, Y., Papesh, M. H., Goldinger, S. D., Kuhn, J. R., Smith, K. A., Treiman, D. M., and Steinmetz, P. N. (2014). Sparse and distributed coding of episodic memory in neurons of the human hippocampus. *Proceedings of the National Academy of Sciences*, 111(26):9621–9626.

Yassa, M. A. and Stark, C. E. L. (2011). Pattern separation in the hippocampus. *Trends in Neurosciences*, 34:515–525.