Identifying unmeasured heterogeneity in microbiome data via quantile thresholding (QuanT)

Jiuyao Lu[1], Glen A. Satten[2], Katie A. Meyer[3], Lenore J. Launer[4], Wodan Ling[5], Ni Zhao[6,*]

Corresponding Authors: Ni Zhao, nzhao10@jhu.edu

[1]Department of Statistics and Data Science, The Wharton School, University of Pennsylvania, 265 South 37th Street, 19104 PA, USA. jiuyaolu@wharton.upenn.edu

[2]Department of Gynecology and Obstetrics, Emory University School of Medicine, 101 Woodruff Circle, 30322 Atlanta, USA. gsatten@emory.edu

[3]Nutrition Research Institute and Department of Nutrition, University of North Carolina, 500 Laureate Way, 28081 Kannapolis, USA. ktmeyer@email.unc.edu

[4]Laboratory of Epidemiology and Population Science, NIA, NIH, 7201 Wisconsin Ave, 20814 Bethesda, USA. lenore_launer@nih.gov

[5]Division of Biostatistics, Department of Population Health Sciences, Weill Cornell Medicine, 402 East 67th Street, 10065 New York, USA. wol4002@med.cornell.edu

[6]Department of Biostatistics, Johns Hopkins Bloomberg School of Public Health, 615 N Wolfe St, 21205 Baltimore, USA. nzhao10@jhu.edu

Abstract

Background: Microbiome data, like other high-throughput data, suffer from technical heterogeneity stemming from differential experimental designs and processing. In addition to measured artifacts such as batch effects, there is heterogeneity due to unknown or unmeasured

factors, which lead to spurious conclusions if unaccounted for. With the advent of large-scale

multi-center microbiome studies and the increasing availability of public datasets, this issue

becomes more pronounced. Current approaches for addressing unmeasured heterogeneity in

high-throughput data were developed for microarray and/or RNA sequencing data. They

cannot accommodate the unique characteristics of microbiome data such as sparsity and over-

dispersion.

Results: Here, we introduce Quantile Thresholding (QuanT), a novel non-parametric approach

for identifying unmeasured heterogeneity tailored to microbiome data. QuanT applies quantile

regression across multiple quantile levels to threshold the microbiome abundance data and

uncovers latent heterogeneity using thresholded binary residual matrices. We validated QuanT

using both synthetic and real microbiome datasets, demonstrating its superiority in capturing

and mitigating heterogeneity and improving the accuracy of downstream analyses, such as

prediction analysis, differential abundance tests, and community-level diversity evaluations.

Conclusions: We present QuanT, a novel tool for comprehensive identification of unmeasured

heterogeneity in microbiome data. QuanT's distinct non-parametric method markedly enhances

downstream analyses, serving as a valuable tool for data integration and comprehensive

analysis in microbiome research.

Keywords: microbiome data, unmeasured heterogeneity, batch effects, zero inflation,

conditional quantile regression

Background

Next-generation sequencing technology has facilitated large-scale microbiome profiling to

explore associations with various disease and health outcomes. Given the large scale, samples

must be collected and profiled in different runs or even from various studies. Furthermore,

there has been a significant increase in the availability of publicly accessible microbiome data[1-

3], and integrating these diverse datasets can be invaluable in advancing scientific discoveries[4,

5]. When integrating these diverse datasets, it is crucial for researchers to recognize potential

variations within these datasets, which might not be immediately apparent to external

researchers. Much of this unwanted variation is attributable to hidden technical or biomedical

heterogeneity, such as instrumental drift, undocumented differences in study design, or even

lab labeling errors. Failing to account for these hidden heterogeneities can obscure the true

biological signals and lead to spurious findings. The situation can be even worse when

including publicly available data[1-3], because sources of variability such as batch labels may not

be included in the distribution data.

In genomic studies, numerous statistical methods have been developed to correct for

batch effects when batch indicators are available, with ComBat being among the most widely

adopted. In the microbiome field, several methods have also been proposed to address batch

effects when indicators are known, including approaches such as MMUPHin[6] and DEBIAS-M[7]

and ConQuR[8], among others. These approaches rely on the availability of batch indicators. In

contrast, no methods are designed to address *unmeasured/unknown* heterogeneity in

microbiome studies, although techniques developed for genomic data, such as surrogate

variable analysis (SVA)[9] and remove unwanted variation (RUV)[10], may be applicable. SVA is

a two-step procedure for identifying surrogate variables (detection of hidden factors and then

construction of surrogate variables), a general framework that was adapted by subsequent

methods, including ours here. RUV employs factor analysis on control genes (via spike-in

experiment or housekeeping gene estimation) to identify hidden heterogeneity. However, SVA

and RUV are not ideally suited for microbiome data. First, they do not address the large number

of zeros and the over-dispersion in microbiome data. Second, they only detect the underlying

structure in the (possibly log-transformed) mean expression while ignoring other possible

changes in distribution. Third, RUV requires users to specify the number of hidden directions:

incorrect specification of these parameters may result in false positive or negative discoveries.

Here, we present the Quantile Thresholding (QuanT) approach, the first tool for

comprehensive identification of unmeasured heterogeneity in microbiome data. QuanT applies

quantile regression[11] at multiple quantile levels to taxa relative abundances, and then forms

dichotomized residuals. QuanT distills information from these dichotomized residuals to

output a set of unit vectors that represent the hidden heterogeneity. As a non-parametric

approach, QuanT is robust to the complex distributional attributes of microbiome data, handles

sparseness in a natural way, and can detect heterogeneity beyond simple mean shifts. The

quantile surrogate variables (QSVs) output by QuanT can be easily incorporated into

downstream analysis. QSVs can be treated as known variables and embedded into any existing

analytic workflows (e.g., regression models) that allow for covariate adjustment. Alternatively,

QuanT can be combined with the recently developed microbiome batch effect removal method,

ConQuR[8], to produce batch-free microbial profiles for downstream analysis.

89 **Methods**

90 **Overview of QuanT**

We first give an overview of the method before delving into a detailed explanation of each step.

We propose a two-stage algorithm, depicted in Fig. 1, to identify unmeasured heterogeneity in

microbiome relative abundance. Similar to existing methods such as SVA, our algorithm,

QuanT, first employs matrix decomposition to detect hidden directions pertaining to

unmeasured factors in the data, followed by a second stage to refine these detected directions.

In the first stage, we fit quantile regression models at multiple quantile levels to the relative

abundance of each taxon with the primary variable of interest and important covariates, and

then generate dichotomized, centered quantile residuals. These columns of residuals are

concatenated into a single matrix of discretized residuals R. We then conduct a singular value

decomposition (SVD) of R to uncover the systematic structure of the residuals and select

singular vectors based on their directional information and variability. These selected vectors,

termed stage-1 singular vectors, capture systematic variation independent of the known

variables.

The stage-1 singular vectors are unrelated to the known variables as these known

variables were included as covariates in the quantile regression models. As a result, adjusting

for stage-1 singular vectors in downstream analysis would not correct for the confounding

effects of the unmeasured heterogeneity (although efficiency may be improved). Therefore, in

the second stage, we perform a "correlation calibration" step, in which we weight the "features"

(i.e., combinations of quantile level and taxon). Specifically, we upweight features that are

strongly marginally associated with the stage-1 singular vectors but weakly associated with the

covariates, and downweight those with the opposite characteristics. Our second stage is an

iterative procedure such that in each iteration, we fit quantile regression models and decompose

a matrix of discretized residuals, similar to the first stage, but using the weighted features for

model fitting. Each iteration will yield new "iterative" singular vectors (iSVs) by decomposing

the residual matrix. The iSVs at convergence, i.e., when the resulting iSVs are close enough to

the iSVs in the previous iteration, are called Quantile Surrogate Variables (QSVs), which are

the output of our algorithm. This kind of correlation calibration step has been shown to be

necessary in similar tasks when analyzing gene expression data[9, 12-15].

Fig. 1 | **Flowchart of QuanT.** V.O.I: variable of interest; QSV: quantile surrogate variable; $\tau_1 \ldots \tau_k$: quantile levels. Stage 1 conducts quantile regression for each taxon at multiple quantile levels using V.O.I (Z) and covariates (X). Residuals are thresholded to create a binary matrix R, which is analyzed through singular value decomposition to identify key directions. Stage 2 implements iterative feature weighting and correlation calibration. Stage 2 starts by fitting quantile regression and thresholding without covariates at multiple quantile levels, followed by the weighting of taxon and quantile levels so that the singular vectors of the weighted residual matrix $R^{(u)\prime}$ are highly associated with the prior stage singular vectors but minimally associated with X and Z. The singular vectors at convergence of Stage 2 are QSVs, the output of QuanT.

Comparing to the existing approaches such as SVA and RUV, QuanT offers several novel modifications/improvements to make it suitable for microbiome data, which is typically sparser and over-dispersed. First, by using quantile regressions instead of least squares or generalized

linear models, QuanT can gather information from the whole distribution of the taxa rather

than just the mean, which is critical for microbiome data. Secondly, SVA and RUV search for

significant directions by identifying singular vectors that represent a disproportionately high

amount of variation. QuanT, on the other hand, relies on both the direction of a singular vector

and the variance it explains, and thus allows us to detect significant directions that may harbor

important underlying structure but explain only a modest proportion of variation.

We now describe our notation and proceed to a more in-depth explanation of the method's

components.

Notation

Assume the relative abundance data form a $n \times p$ matrix Y, in which Y_{ij} is the relative

abundance of the j-th taxon in the i-th sample. We denote by $Y_{.j}$ the j-th column of Y

comprising data on the j-th taxon, and treat it as the outcome in the proposed regression

framework. Suppose that in addition each sample i has a primary variable of interest (denoted

by Z_i) and a set of L covariates (denoted by X_i) to adjust for. We use quantile regression at

each of K quantile levels in our regression framework; these levels are represented by the

vector, $\vec{\tau}$. By default, we set the quantile levels as an equally spaced sequence from 0.1 to 0.9

with a step size of 0.005 corresponding to K=161.

Note that some taxon-quantile combinations may not convey useful information. For example,

if 70% of the observations for a specific taxon j are zeros, any quantile levels $\vec{\tau}$ less than 70%

will be irrelevant and are thus disregarded for this taxon. Thus, the effective length of $\vec{\tau}$ varies

by taxon. This approach makes $\vec{\tau}$ data adaptive, utilizing more quantile levels for taxa with

more informative distributions.

Stage 1: Obtaining singular vector directions that capture the underlying heterogeneity

structure.

Quantile thresholding

For each feasible pair of taxon j and quantile level τ_k, we fit the quantile regression

model $Q_{\tau_k}(Y_{\cdot j}|X, Z) = \alpha(\tau_k) + \sum_{l=1}^{L} X_{il}\beta_l(\tau_k) + Z_i\gamma(\tau_k)$ to the data $Y_{\cdot j}$ in which

$Q_{\tau_k}(Y|X, Z) = \inf\{y: F_{Y|X,Z}(y) > \tau_k)\}$ is the τ_k -th quantile function. Parameters

$(\hat{\alpha}_j(\tau_k), \hat{\beta}_{lj}(\tau_k), \hat{\gamma}_j(\tau_k))$ are obtained by solving the minimization problem

$$(\hat{\alpha}_j(\tau_k), \hat{\beta}_{lj}(\tau_k), \hat{\gamma}_j(\tau_k)) = \arg\min_{\alpha,\beta,\gamma} \sum_{i=1}^{n} \rho_{\tau_k}\left(Y_{ij} - \alpha(\tau_k) - \sum_{l=1}^{L} X_{il}\beta_l(\tau_k) - Z_i\gamma(\tau_k)\right),$$

where ρ_{τ_k} is the check function[16] $\rho_{\tau_k(x)} = \tau_k x - I\{x < 0\}x$.

After fitting the quantile regression model, we threshold the microbial abundances against the

estimated quantiles and center them as $I\{Y_{ij} > \hat{\alpha}_j(\tau_k) + \sum_{l=1}^{L} X_{il}\hat{\beta}_{lj}(\tau_k) + Z_i\hat{\gamma}_j(\tau_k)\} -$

$(1 - \tau_k)$. For each pair of taxon j and quantile level τ_k, this yields a n-dimensional vector of

discretized (binary) centered residuals. We gather the results from different pairs into a matrix

R with n rows (samples) and no more than pK columns (processed features) as we omit the

uninformative pairs of taxon and quantile level. R captures the heterogeneity structure such as

the residual matrix constructed by SVA but is more robust and distribution-free and handles

zero cells in a natural way.

Singular vector selection

We conduct SVD of the binarized and centered residual matrix R to obtain

$$177 \qquad\qquad\qquad\qquad R \; = \; U\Sigma V.$$

We aim at selecting a subset of the singular vectors, i.e., columns of U, to represent the

important structure in the data. We use an integrated approach to select the singular vectors.

Specifically, we select singular vectors 1) whose distributions are statistically different from

the marginal distribution of the singular vectors if there is no underlying structure in the binary

centered residual matrix (a.k.a., complete null, details are discussed in the next paragraph), and

2) which explain a large proportion of variability of the binary centered residual matrix. We

use the SelectiveSeqStep procedure in the knockoff paper[17] to achieve this goal based on the

two criteria simultaneously. This dual goal selection is a novel feature of QuanT. The detailed

rationale and procedure are as follows.

Under the complete null that there is no underlying structure in the binary centered

residual matrix, each singular vector direction should point to a uniformly distributed point on

the n-dimensional sphere; this implies that the components of each singular vector should

follow a Gaussian distribution[18]. Thus, we can test whether a singular vector is an important

direction by testing the normality of each singular vector direction using the Lilliefors test[19] or

the Anderson-Darling test[20] or the Shapiro-Francia test[21] and obtain its corresponding p-value.

We also order the singular vectors based on the size of their corresponding singular values. We

aim to select from the top m singular vectors which have p-values smaller than a cutoff c (e.g.,

0.05). The choice of m is determined by controlling the false discovery rate (denoted by q) of

the selected singular vectors. Under the null hypothesis, p-value is uniformly distributed; thus

the ratio between the number of singular vectors with p-values greater than c and those with p-

values less than c should be $\frac{1-c}{c}$. Roughly speaking, the singular vectors with p-values $> c$

can be considered generally null, and therefore, we can estimate the number of falsely

discovered singular vectors based on them, which is $\#\{j \leq m: p_j > c\}/\frac{1-c}{c}$, where #

represents the size of a set. This is similar to the idea of q-value[22-24], a commonly used approach

for multiple testing problems. The proportion of falsely discovered singular vectors, if we

select the top m singular vectors, are estimated by $\frac{\#\{j \leq m: p_j > c\}}{\#\{j \leq m: p_j \leq c\} \vee 1} / \frac{1-c}{c}$. We pick the m such that

this proportion is controlled below a false discovery rate threshold q if the top m singular

vectors are selected and would exceed q if the top $m + 1$ singular vectors were selected:

206
$$\hat{m} = \min\left\{m: \frac{\#\{j \leq m: p_j > c\}}{\#\{j \leq m: p_j \leq c\} \vee 1} / \frac{1-c}{c} > q\right\} - 1.$$

After determining $\hat{m}$, we select the singular vectors from the top $\hat{m}$ singular vectors with p-

values smaller than c. Note that our approach allows for a non-monotonic selection of singular

vectors, yet primarily focus on those that account for a substantial portion of variability in the

data.

We define Φ to be the matrix with the s-th column, denoted as Φ_s, equal to the s-th stage-1

singular vector.

Stage 2: Iterative feature weighting and correlation calibration

Similar to SVA[9, 12-15], we conduct a second stage to refine the stage-1 singular vectors that we

call correlation calibration. The reason is that, by regressing out X and Z to construct the

binarized centered residuals in the quantile regression stage, the selected singular vectors are,

loosely speaking, uncorrelated with the primary variable Z and covariates X (if we had used

linear regression, this statement would be exact). Thus, they are not able to capture unmeasured

confounding effects, as confounders must be correlated with Z as well. Nonetheless, it is the

unmeasured confounders that pose the greatest threat to the integrity of a study.

Because the stage-1 singular vectors represent the important features of the full set of

residuals R, we seek to reweight the columns of R that will allow us to identify directions that

not only reflect the underlying data structure but also maintain potential correlations with X

and Z, thereby enhancing the robustness of downstream analysis against hidden confounding

factors.

To accomplish this, we first conduct a similar thresholding procedure as in stage 1, but

on a reduced quantile regression model $Q_{\tau_k}(Y_{\cdot j}) = \alpha_j(\tau_k)$, i.e., without regressing X and Z out,

and construct the binarized and centered residuals (denoted by R'). We then weight columns

(combinations of taxon and quantile levels) of R' to prioritize those that represent underlying

data structure while being minimally associated with X and Z. Specifically, we use an iterative

procedure that balances the two goals—capturing the underlying heterogeneity and

maintaining appropriate correlation with X and Z.

In the first iteration, for each feasible pair of taxon j and quantile level τ_k, we fit two

pairs of quantile regression models to the columns of R'. The first pair compares the reduced

quantile regression model $Q_{\tau_k}(Y) = \alpha(\tau_k)$ to the model that includes the stage-1 singular

vectors $Q_{\tau_k}(Y|\Phi) = \alpha(\tau_k) + \sum_{s=1}^{\hat{m}} \Phi_s \theta_s(\tau_k)$. The second pair compares the model with the

stage-1 singular vectors $Q_{\tau_k}(Y|\Phi) = \alpha(\tau_k) + \sum_{s=1}^{\hat{m}} \Phi_s \theta_s(\tau_k)$ to the full model

$Q_{\tau_k}(Y|X, Z, \Phi) = \alpha(\tau_k) + \sum_{l=1}^{L} X_{il} \beta_l(\tau_k) + Z_i \gamma(\tau_k) + \sum_{s=1}^{\hat{m}} \Phi_s \theta_s(\tau_k)$. Note that the two

models in the first pair can assess whether the taxon-quantile combination can capture

information in the stage-1 singular vectors, and the second pair of models allows us to assess

whether the taxon-quantile combination convey information about the known covariates,

conditional on the stage-1 singular vectors. Thus, these two pairs of models facilitate the

identification of directions that relate to the stage-1 singular vectors without being overly

influenced by X and Z.

For each pair of hypotheses at each taxon j and quantile level τ_k, we calculate a rank-

based pseudo-F statistic[25] to generate a p-value, comparing whether the larger model fits the

data better than the smaller model. With this, we obtain two sets of p-values, with $p_{jk}^{(1)}$ $(p_{jk}^{(2)})$

corresponding to the p-value comparing the first (second) pair of models for taxon j and

quantile level τ_k. Intuitively, a small $p_{jk}^{(1)}$ implies that the stage-1 singular vectors

corresponding to taxon j and quantile level τ_k convey important information on the data

structure; yet a large $p_{jk}^{(2)}$ suggests that taxon j and quantile level τ_k is less impacted by the

known covariates given the corresponding stage-1 singular vectors. Thus, we will weight each

column of R' so that columns with small $p_{jk}^{(1)}$ values and large $p_{jk}^{(2)}$ values receive a higher

weight.

To be specific, we apply the local FDR method[26] to the two sets of p-values

separately to obtain two sets of local FDR estimates. This transforms each set of p-values to a

set of local FDR estimates. Each taxon-quantile pair corresponds to two p-values before the

transformation, and hence corresponds to two local FDR estimates after the transformation.

These two local FDR estimates answer the following questions: given the observed data for

this specific taxon-quantile pair, what is the posterior probability that it is impacted by the

unmeasured heterogeneity and what is the posterior probability that it is not impacted by the

known covariates. We then use the product of the two local FDR estimates as the weight for

the taxon-quantile pair. We multiply each column of R' by its weight and obtain a reweighted

matrix $R^{(u)'}$. We then conduct SVD to obtain a new set of singular vectors, which we denote

as $\Psi^{(1)}$, in which the superscripts indicate the iteration cycle, and we are now at iteration 1.

Note that R' is obtained without regressing out X and Z, and thus, $\Psi^{(1)}$ can be correlated with

X and Z. Note also that the number of the new singular vectors $\Psi^{(1)}$ can be larger than the

stage-1 singular vectors Φ. We seek to only maintain the directions that are most correlated

with Φ. To do this, for each column, Φ_s, we search for the columns of $\Psi^{(1)}$ that have the

highest absolute correlation with Φ_s. In other words, for each column Φ_s, we choose the d_s-

th column of $\Psi^{(1)}$ in which $d_s = \arg\max_d |\text{corr}(\Phi_s, \Psi_d^{(1)})|$. The final selected singular

vectors are called the iSVs at the first iteration.

For later iterations, we replace the stage-1 singular vectors in the two pairs of regression

models with the iSV of the previous iteration and follow the same procedure. The iteration

stops when iSVs from two adjacent iterations are the close enough, or when we reach a

sufficient number (T_{max}, with default = 10) of iterations. In the second case, we calculate the

final singular vector as a weighted average of the last $T_{max}/2$ iterations. The final singular

vectors are the QSV output of our algorithm. Please refer to the Algorithm 1 for individual

steps.

Algorithm 1: QuanT algorithm

Input: Y, X, Z

Output: Ψ (quantile surrogate variables)

---

1. Stage 1 residualization:

   For each feasible pair of taxon j and quantile level τ_k, fit quantile regression

   $Q_{\tau_k}(Y_{.j}|X, Z) = \alpha_j(\tau_k) + \sum_{l=1}^{L} X_{il} \beta_{lj}(\tau_k) + Z_i \gamma_j(\tau_k);$

   Compute $r_{jk} = I\{Y_{ij} > \hat{\alpha}_j(\tau_k) + \sum_{l=1}^{L} X_{il} \hat{\beta}_{lj}(\tau_k) + Z_i \hat{\gamma}_j(\tau_k)\} - (1 - \tau_k);$

Stack r_{jk} across j and k to form matrix R.

2. Conduct SVD on $R = U\Sigma V$.

3. Apply normality test to each singular vector to get p-values. From the top m singular vectors, select singular vectors whose p-value is less than a cutoff c using the SelectiveSeqStep procedure. Here, m is selected so that the FDR is smaller than a prespecified value of q. These form the matrix consisting of stage-1 singular vectors Φ.

4. For each feasible pair of taxon j and quantile level τ_k, fit the reduced quantile regression model $Q_{\tau_k}(Y._j) = \alpha_j(\tau_k)$;

   Compute $r'_{jk} = I\{Y_{ij} > \tilde{\alpha}_j(\tau_k)\} - (1 - \tau_k)$, where $\tilde{\alpha}_j(\tau_k)$ is the estimate under this reduced model.

   Stack r'_{jk} across j and k to form matrix R'.

5. Iterative procedure for correlation calibration:

   a. Initialize $\Psi^{(0)} \leftarrow \Phi$, set $u \leftarrow 0$

   b. Repeat until convergence or until $u = T_{max}$ (maximum number of iteration)

      i. For each taxon j and quantile τ_k, compare pairs of models to obtain $p_{jk}^{(1)}$ and $p_{jk}^{(2)}$:

   Pair 1: $Q_{\tau_k}(Y) = \alpha(\tau_k)$,

   vs. $Q_{\tau_k}(Y|\Psi^{(u)}) = \alpha(\tau_k) + \sum_{s=1}^{\hat{m}} \Psi_s^{(u)} \theta_s(\tau_k)$;

Pair 2: $Q_{\tau_k}\big(Y|\Psi^{(u)}\big) = \alpha(\tau_k) + \sum_{s=1}^{\widehat{m}} \Psi_s^{(u)} \theta_s(\tau_k),$

vs. $Q_{\tau_k}\big(Y|X, Z, \Psi^{(u)}\big) = \alpha(\tau_k) + \sum_{l=1}^{L} X_{il}\, \beta_l(\tau_k) + Z_i \gamma(\tau_k) +$

$\sum_{s=1}^{\widehat{m}} \Psi_s^{(u)} \theta_s(\tau_k)$

    ii.   Compute Weights:

Apply the local FDR method to $p_{jk}^{(1)}$ to obtain $q_{jk}^{(1)}$;

Apply the local FDR method to $p_{jk}^{(2)}$ to obtain $q_{jk}^{(2)}$;

Calculate the weight as $w_{jk} = \left(1 - q_{jk}^{(1)}\right) q_{jk}^{(2)}$;

Form $R^{(u)\prime}$ by multiplying the corresponding column of R' by w_{jk} for each combination of j and τ_k

    iii. Conduct SVD on $R^{(u)\prime}$.

    iv. Select the set of singular vectors of $R^{(u)\prime}$ that shares the strongest absolute correlation with the stage-1 singular vectors Φ. Denote these as $\Psi^{(u+1)}$.

    v.   Convergence check: If $\left\|\Psi^{(u+1)} - \Psi^{(u)}\right\|_F \leq \epsilon$, then break and set $\Psi = \Psi^{(u+1)}$; else $u \leftarrow u + 1$.

c.   If convergence is not reached after T_{max} iterations:

    i.   Calculate the weight $\bar{w}_{jk}$ as the average of w_{jk} in the last $\frac{T_{max}}{2}$ iterations;

    ii. Form $R^{(u)\prime}$ by multiplying the corresponding column of R' by $\bar{w}_{jk}$ for each combination of j and τ_k.

iii. Conduct SVD on $R^{(u)'}$.

iv. Select the set of singular vectors of $R^{(u)'}$ that shares the strongest absolute correlation with the stage-1 singular vectors Φ. Denote these as Ψ.

6. Return Ψ.

282
283

Singular vectors selection when no stage-1 singular vectors are found

In rare cases, we may encounter the situation where no singular vector is selected in the first

stage. This usually happens when the hidden heterogeneity variable is normally distributed or

when the sample size is not large enough. In this case, we provide an alternative approach that

selects important singular vectors solely based on the proportion of variabilities explained. This

approach is very similar to the principal component/eigengene selection methods in SVA,

except that the least squares regression models are replaced by appropriate quantile regressions.

The detailed algorithm can be found in the Algorithm 2. Note that this variant in our algorithm

only applies to calculating the stage-1 singular vectors; we still need the same iterative

correlation calibration procedure as in the standard QuanT algorithm (Algorithm 1).

Algorithm 2: alternative stage-1 algorithm when no singular vector is selected through the normality test

1-2 Step 1 – 2 is the same as the Algorithm 1.

3. Let d_k be the k-th singular value. For the k-th singular vector, construct the test statistic as $T_k = \frac{d_k^2}{\sum d_l^2}$

4. Form a matrix R^0 by permuting each row of R independently to remove any structure in the matrix. Denote the (i, j) entry of R^0 by r_{ij}^0.

5. Fit the models $Q_{\tau_k}(r_{\cdot j}^0 | X, Z) = \alpha(\tau_k) + \sum_{l=1}^{L} X_{il} \beta_l(\tau_k) + Z_i \gamma(\tau_k)$, and obtain the binarized and centered residual matrix, denote by R^{0*}

6. Conduct SVD of R^{0*} and calculate statistic $T_k^0 = \frac{d_{0k}^2}{\sum d_{0l}^2}$

7. Repeat steps 5-7 for a total of B times to obtain null statistics $T_K^{0b}, b = 1, \ldots B$. We set B as 10.

8. Compute the p-value for singular vector k as

$$p_k = \frac{\#\{T_K^{0b} \geq T_k; \ b = 1, \ldots B\}}{B}.$$

We conservatively force monotonicity among the p-values. Thus, set $p_k = max\{p_{k-1}, p_k\}$.

9. Select the singular vectors with $p_k \leq 0.1$.

10. Carry out correlation calibration (steps 6-7 in Algorithm 1)

Simulation settings

We conducted simulations to evaluate QuanT's performance when the true signal is known.

We simulated data that mimicked two datasets from the integrative Human Microbiome

Project[27]: a gut microbiome dataset from the Inflammatory Bowel Disease Multi-omics

Database (IBDMDB) project[28], and a vaginal microbiome dataset from the Multi-Omic

Microbiome Study: Pregnancy Initiative (MOMS-PI) project[29]. These two datasets represent

microbiome data from two body sites that are of great research interest. They also have

contrasting attributes in terms of sparsity and over-dispersion level, which makes them useful

for evaluating our proposed method QuanT. We used two methods to simulate data that

mimicked the template data: the parametric Dirichlet-Multinomial (DM) model and the non-

parametric approach in MIDASim[30]. We additionally simulated a third scenario where

microbiome absolute abundances were generated using a log-linear model, in order to evaluate

QuanT's performance in compositional settings.

For each sample i, we simulated a clinical variable of interest (denoted by Z_i) and a hidden

batch variable (denoted as B_i) through a bivariate Bernoulli distribution, with both variables

having a marginal mean of 0.5. A correlation of 0.3 was set between them. Note that because

the "batch" variable is simulated to be correlated with both the microbiome data and Z, thus, it

is a confounder. Here, we use the "batch" variable to represent all the hidden heterogeneities.

For the parametric simulations, we adopted the simulation setup from a previous study[31]

to simulate taxa abundances using the Dirichlet-Multinomial (DM) distribution. In our

simulations, we designated the batch variable to impact most taxa (referred to as Batch Taxa),

while the variable of interest, Z, influenced only a small subset (referred to as Clinical Taxa).

Specifically, for the IBDMDB dataset, we randomly selected 250 of the 301 taxa to be Batch

Taxa, among which 40 were also Clinical Taxa. Similarly, for the MOMS-PI dataset, we

selected 200 of the 233 taxa as Batch Taxa, with 50 of these being Clinical Taxa. We first fitted

the DM model to the template datasets (IBDMDB and MOMS-PI) and obtained the estimated

mean vectors π_0. Subsequently, we formed two additional vectors π_B and π_Z to incorporate

batch and clinical effects. Both vectors were initially set to be the same as π_0. We then

randomly permuted the frequencies within π_B and π_Z corresponding to the Batch Taxa and

Clinical Taxa. Finally, a sample-specific frequency was defined as $\pi_i = (1 - w_B B_i -$

$w_z Z_i)\pi_0 + w_B B_i \pi_B + w_z Z_i \pi_z$, where (w_B, w_z) determine the underlying batch and clinical

effects. We considered three combinations for these parameters: $(w_B, w_z) \in$

$\{(0,0.5), (0.5,0.5), (0.8,0.2)\}$. When $w_B = 0$, there is no underlying hidden

heterogeneity/batch effect. The count data for each sample were generated using the DM

distribution with π_i and an overdispersion parameter of 0.02.

For the non-parametric simulations, we used the Microbiome Data Simulator

(MIDASim)[30], which is designed to generate microbiome data that capture the underlying

distributional and correlation structure of a template dataset. It offers a nonparametric quantile

matching approach, and a parametric approach based on log-linear models. Here, we utilized

the non-parametric version because it captures the underlying data structure more accurately.

Via MIDASim, for each template data set, we first generated vectors of relative abundances

and zero-proportions. After choosing the Batch Taxa and the Clinical Taxa, we generated

sample-specific relative abundance and zero-proportion vectors in the same way as in the DM

simulation and also used the same (w_B, w_z) combinations. Count data for each sample were

generated by inputting the sample-specific vectors into MIDASim and choosing to use its non-

parametric approach.

In the third simulation scenario, we again simulated microbiome read counts using the

Dirichlet–Multinomial model, but introduced batch and clinical effects differently. Instead of

modifying the mean frequency vectors π_B and π_Z as in the first simulation, we first generated

read counts from the fitted DM model using π_0, and then directly adjusted the counts.

Specifically, for samples with $B_i = 1$, read counts of Batch Taxa were multiplied by w_B; for

samples with $Z_i = 1$, read counts of Clinical Taxa were multiplied by w_z . We considered three

combinations for these parameters: $(w_B, w_z) \in \{(1,4), (4,4), (5,3)\}$.

Results

To rigorously assess QuanT, we conducted extensive simulation studies based on real-world gut and virginal microbiome datasets from the HMP2 projects[27]. We also applied QuanT and competing methods to three curated microbiome datasets, each of which has distinct sources of technical heterogeneity: a gut microbiome study on cardiovascular diseases, which displays a moderate technical variation arising from sequencing in multiple runs; an integrated dataset compiled from various HIV gut microbiome studies; and a consolidated dataset encompassing multiple colon cancer studies. Through both visual and quantitative comparisons, we demonstrate that QuanT outperforms existing methods by robustly and comprehensively uncovering underlying microbiome data heterogeneity, which is beyond mere shifts in means. When incorporated into the analytical workflow, QuanT effectively reduces the influence of unwanted heterogeneity, enhancing false discovery rate control, maintaining sensitivity of discovery, as well as improving prediction accuracy.

Application to the CARDIA study, a single large-scale epidemiological investigation

We assessed QuanT using multiple datasets. We first applied it to a study containing traditional batch variation: samples are collected under one protocol but handled in different batches. The Coronary Artery Risk Development in Young Adults (CARDIA) Study[32], initiated in 1985-1986, recruited young adults aged between 18 and 30 with the goal of uncovering determinants of cardiovascular disease (CVD) throughout adult life. Subject enrollment was balanced according to race, gender, education, and age. Extensive data on risk factors related to CVD have been collected, including health behaviors, medical history (including medication use), and clinical risk factors, such as blood pressure. Stool samples were also collected for some visits. Here, we used the microbiome data generated from the

stool samples collected during the Year 30 follow-up (2015-2016). The samples were

processed through four DNA extraction/library preparation batches, and subsequently

sequenced through seven runs using the Illumina MiSeq 2x300 platform. Specifically, each

DNA extraction batch was allocated to two sequencing runs, except for the final batch, which

was sequenced by itself during the seventh run. Consequently, the data were processed in

seven distinct batches. Following sequencing, forward reads were processed through the

DADA2[33] pipeline for quality control and derivation of amplicon sequence variants (ASVs),

and taxonomy was assigned using the Silva reference database[34]. The data were agglomerated

to the genus level, and taxa that did not yield any reads in all samples were omitted from the

dataset. After preprocessing, the data contains 633 samples and 145 taxa. Analyses were

restricted to participants for whom all covariate data were available.

In this study, we considered high blood pressure (HBP) (non-HBP = 0, HBP = 1) as

the variable of interest, and gender (Male = 0, Female = 1) and race (White = 0, Black = 1) as

adjustment covariates. To assess QuanT and the competing methods, we treated the batch

indicators as unknown and used these methods to identify the now-hidden

heterogeneities/batches. The performance of these methods was assessed based on three

criteria: 1) consistency and sensitivity in downstream differential abundance analysis for

individual taxa; 2) enhancement in prediction accuracy for the variable of interest; and 3) the

efficacy of batch effect removal while preserving the effect from the variable of interest at the

entire community level. For SVA, we used the svaseq function, which added a pseudo count

of 1 to all counts and then applied log transformation as described in Leek (2014)[15], and used

the iterative reweighted algorithm[12]. The default parameter setting was used. For RUV, we

used the RUVg approach in Risso *et al.* (2014)[10] with an estimated group of null taxa as

controls, rather than RUVr and RUVs as these approaches can only identify hidden directions

that are uncorrelated with the variables of interest. For hyperparameters, we selected control

taxa using the Benjamini–Hochberg procedure on edgeR p-values at a 0.05 threshold, and set

the number of latent factors following the guidance in Risso *et al.* (2014)[10].

Fig. 2a and 2b compare the consistency of differential abundance analysis results

across various models. Here, differential abundance analysis was conducted using linear

regression on relative abundances of each taxon, controlling for age, gender, and identified

hidden variables. For benchmarking purposes, analyses were conducted adjusting for true

batch/group indicators (TG), which was regarded as the gold standard, and with no correction

(NC), which served as the baseline. Fig. 2a shows the number of differentially abundant taxa

identified from the CARDIA dataset by Benjamini-Hochberg adjusted p-value < 0.05. TG

leads to 4 discoveries. Considering these as true signals, NC identifies the 4 true discoveries

and has 1 additional false discovery; QuanT identifies 3 true discoveries and has 1 additional

false discovery. SVA and RUV lead to no discoveries. Therefore, QuanT demonstrates

superior performance compared to SVA and RUV when applied to the CARDIA dataset,

identifying more potential true discoveries while maintaining a reasonable false discovery

proportion. Fig. 2b shows the top K detected differentially abundant taxa. Treating TG as the

gold standard and calculating the proportion of overlapping discoveries, QuanT has

comparable performance with NC and SVA, and is substantially better than RUV.

Fig. 2c shows the prediction accuracy using the original and the ConQuR corrected

microbiome data, where the corrected data were adjusted for TG, or surrogate variables

estimated by SVA, RUV or QuanT. Random forest models were fit with five-fold cross-

validation to predict HBP status, and the area under the receiver operating characteristic curve

(ROC-AUC) was used to evaluate the accuracy. We note that this is not a genuine predictive

analysis, but rather a metric to quantify how well the corrected microbiome data can explain

the variable of interest. As shown in Fig. 2c, RUV and QuanT corrected data predicts HBP

more accurately than using TG or SVA, and prediction using the unadjusted data (NC) leads

to the worst accuracy. Fig. 2d shows the principal coordinate analysis (PCoA) plots using the

Bray-Curtis distance, comparing with the uncorrected and ConQuR-corrected datasets.

Assessed by the PERMANOVA R^2 (which evaluates the multivariate association between the

microbial community and a certain variable, here quantifies the proportion of microbiome data

variability explained by the variable), correction with TG leads to the most significant

reduction in batch effects, which was expected. Visually, it not only equalizes the mean and

dispersion (centroid and size of the ellipses) across groups but also removes higher-order

batch/group effects (shown by angles of the ellipses). Correction for QuanT, SVA, and RUV's

estimated surrogate variables leads to comparable efficacy in reducing batch effects and

maintaining the effects of high blood pressure (HBP).

It is worth noting that the CARDIA study has well-balanced batch designs; thus the

analysis with NC leads to reasonable and comparable results as the analyses with adjustments.

Fig. 2 | Evaluation on the CARDIA data. NC: no batch correction; TG: correction using true group indicators. **a** Heatmap representing the number of differentially abundant taxa identified in the CARDIA dataset by Benjamini-Hochberg adjusted p-value < 0.05; diagonal elements are numbers detected, and off-diagonal elements are numbers detected in common. **b** Proportions of overlapping taxa in the top K detected differentially abundant taxa between each method and TG in the CARDIA dataset. **c** Average cross-validated area under the receiver operating characteristic curves (ROC-AUC) of predicting HBP from the ConQuR-corrected microbiome data via random forest. Larger values indicate stronger predictive capabilities. **d** PCoA plots of the ConQuR-corrected data based on Bray-Curtis distance, with PERMANOVA R^2 of batch ID and HBP.

Application to HIVRC, an integrated dataset with multiple HIV studies

Our second example used data from the HIV re-analysis consortium (HIVRC)[5]. In this consortium study, raw 16s rRNA gene sequencing data were solicited from 22 studies that investigated the relationship between the gut microbiome and HIV infection. Data from 17 of these studies were subsequently processed through a common pipeline with taxa assigned using

Resphera Insight[35]. Note that the studies in the final-processed data have distinct experimental

designs and sequencing protocols (Supp. Tab. 2 of Tuddenham, et al. 2020[17]). Thus, the data

have more pronounced variability than that observed within a single study, such as the

CARDIA study.

We selected two studies with similar library sizes (5000-5500) but very different study

designs (an observational study in the US[36] and a clinical trial in Europe[37]), which include 37

and 24 samples respectively. After preprocessing, the data contains 126 taxa. We applied

QuanT, SVA, and RUV to the data aggregated at the genus level. HIV status (Negative = 0,

Positive = 1) was regarded as the variable of interest, while age, gender and BMI were

considered as covariates. We retained complete cases only and conducted the same set of

analyses as on the CARDIA data.

Fig. 3a and 3b compare the consistency of differential abundance analysis results

obtained analyzing these data using various methods. Fig. 3a shows the number of

differentially abundant taxa identified in the HIVRC dataset by Benjamini-Hochberg adjusted

p-value < 0.05. We identified 19 discoveries adjusting for true group indicators (TG), which

were considered as the gold standard. QuanT leads to 17 true discoveries and 1 false discovery.

NC leads to 11 true discoveries and 16 false discoveries. SVA leads to 11 true discoveries and

21 false discoveries. RUV leads to 8 true discoveries and 5 false discoveries. Therefore, QuanT

again performs better than the other methods when applied to the HIVRC dataset and has the

lowest FDR compared to competitors. In Fig. 3b, we consider the top K detected differentially

abundant taxa. QuanT is the most consistent with the gold standard, TG, followed by NC, SVA,

and RUV.

Fig. 3c shows the average cross-validated ROC-AUC of predicting HIV from the

ConQuR-corrected data via random forest. TG, SVA, RUV, and QuanT all perform perfectly

and are better than no correction. Fig. 3d shows the PCoA plots using Bray-Curtis distance,

comparing with the uncorrected and ConQuR-corrected datasets. Visually, TG, RUV and

QuanT lead to better harmonization of the microbiome data from the two distinct HIV studies.

As quantified by PERMANOVA R^2, QuanT leads to the greatest reduction of the effects of the

unwanted heterogeneity, followed by RUV, SVA and TG. On the other hand, SVA maintains

the HIV effects the most, followed by TG, QuanT and RUV (figure 3d).

Fig. 3 | Evaluation on the HIVRC data. NC: no batch correction; TG: correction using true group indicators. **a**

Heatmap representing the number of differentially abundant taxa identified in the HIVRC dataset by Benjamini-

Hochberg adjusted p-value < 0.05; diagonal elements are numbers detected, off-diagonal elements are numbers

detected in common. **b** Proportions of overlapping taxa in the top K detected differentially abundant taxa between

each method and TG in the HIVRC dataset. **c** Average cross-validated area under the receiver operating

characteristic curves (ROC-AUC) of predicting HIV from the ConQuR-corrected data via random forest. Larger

values indicate stronger predictive capabilities. **d** PCoA plots of the ConQuR-corrected data based on the Bray-

Curtis dissimilarity, with PERMANOVA R^2 of study ID and HIV.

Application to an integrated microbiome dataset in colon cancers

Our third example came from two colorectal cancer (CRC) studies[38, 39] from the

curatedMetagenomicData package[40]. The CuratedMetagenomicData data package offers

standardized, curated, and analysis-ready metagenomic datasets derived from human

microbiome studies, facilitating comprehensive meta-analyses and cross-study comparisons

by ensuring consistent data processing and annotation. The bacterial, fungal, and archaeal

taxonomic abundances for each sample were calculated with MetaPhlAn3[41], and metabolic

functional potential was calculated with HUMAnN3[41]. We selected two CRC studies[38, 39]

from this package, which include 154 and 80 stool samples, respectively. After filtering, 166

genera were included in the data. These studies collectively cover samples from three distinct

health conditions: colorectal cancer (CRC), adenoma, and disease-free controls, with CRC

subdivided into adenocarcinoma and carcinoma, and adenoma into simple adenoma and

advanced adenoma.

We applied QuanT, SVA, and RUV to this CRC dataset to find the hidden heterogeneity

directions, treating the true study indicators as unknown. We considered the five phenotype

categories (adenocarcinoma, carcinoma, adenoma, advanced adenoma, control) as the primary

variable of interest, age (Adult = 0, Senior = 1) and gender (Male = 0, Female = 1) as the

covariates to adjust for.

Fig. 4a and 4b compare the consistency of differential abundance analysis results across

various methods. Fig. 4a shows the number of differentially abundant taxa identified in the

CRC dataset by Benjamini-Hochberg adjusted p-value < 0.05. TG (gold standard) leads to 8

discoveries. QuanT leads to 5 true discoveries with 1 false discovery. NC leads to 6 true

discoveries and 24 false discoveries. SVA leads to 6 true discoveries and 23 false discoveries.

RUV leads to 1 true discovery with no false discovery. Therefore, NC and SVA lead to inflated

false discovery proportion when applied to the CRC dataset. RUV and QuanT have reasonable

false discovery proportion control, while QuanT is notably better at identifying true signals

than RUV. In Fig. 3b, we consider the top K detected differentially abundant taxa. QuanT is

the most consistent with the gold standard, TG. NC, SVA, and RUV perform similarly are all

outperformed by QuanT.

Fig. 4c presents the prediction accuracy, measured by negative cross entropy, for the

five phenotype groups using the original and ConQuR-corrected CRC microbiome data.

Here, QuanT performs the best, followed by SVA, RUV, TG, and NC. Fig. 4d displays the

PCoA plots using Bray-Curtis distance. Visually, QuanT leads to similar data harmonization

as using the true study IDs, and outperforms SVA, RUV and NC. This is numerically

confirmed by the PERMANOVA R^2 of the study ID and the phenotypes of interest.

Fig. 4 | Evaluation on the CRC data. NC: no batch correction; TG: correction using true group indicators. **a**

Heatmap representing the number of differentially abundant taxa identified in the CRC dataset by Benjamini-

Hochberg adjusted p-value < 0.05; diagonal elements are numbers detected, off-diagonal elements are numbers

detected in common. **b** Proportions of overlapping taxa in the top K detected differentially abundant taxa between

each method and TG in the CRC dataset. **c** Average cross-validated negative cross entropy of predicting HIV

from the ConQuR-corrected data via random forest. Larger values indicate stronger predictive capabilities. **d.**

PCoA plots of the ConQuR-corrected data based on the Bray-Curtis dissimilarity, with PERMANOVA R^2 of

study ID and Phenotype.

Evaluation using simulated data

We applied SVA, RUV and QuanT to the simulated datasets to identify the underlying hidden

heterogeneity, treating the batch variables as unknown. Like in the CARDIA, HIVRC, and

CRC analyses, we used linear regression models to conduct differential abundance analysis,

adjusting for hidden variables estimated by SVA, RUV, and QuanT. When data are simulated

from MIDASim, as shown in Fig. 5a, QuanT leads to better FDR control compared to SVA

and RUV. Fig. 5b shows that the discoveries by QuanT align more closely with the gold

standard than SVA, RUV, and the NC methods. Here, the gold standard varies: in the absence

of batch effects, NC is the gold standard, while when batch effects are present, the model

correcting for true batch ID is the gold standard. More simulation results based on MIDASim

(Supp. Fig. 2,4,6) and results based on parametric simulations (Supp. Fig. 1,3,5) lead to similar

conclusions.

We also applied ConQuR to remove the true batch, SVs and QSV effects and generate

a "batch-free" dataset. Integrating QuanT with ConQuR in simulations (Supp. Fig. 7-10)

yielded microbial profiles where confounding batch effects were eliminated, while the effects

of the variable of interest were preserved. Finally, QuanT demonstrates scalable computational

performance for large-scale microbiome studies, as evidenced in Tab. 1 and Tab. 2.

For our simulation scenario 3, we applied LinDA[42] and LDM-CLR[43], two modern tools

that are designed for compositional hypothesis in microbiome data. These results (Supp. Fig.

11-16) align with the conclusions from our other simulation studies.

In short, QuanT demonstrates improved performance compared to existing methods,

on simulated datasets and biological datasets, regardless of the effect size and the type of source

of unmeasured heterogeneity.

Fig. 5 | Evaluation on the simulated data. The data are simulated by MIDASim using the IBDMDB and MOMS-PI datasets as templates. **a** FDR of the differential abundance analysis via adjusted linear regression with BH-adjusted p-value < 0.05. **b** Proportions of overlapping taxa in the top K detected differentially abundant taxa between each method and the gold standard in simulated data. Simulation setups vary across different datasets (rows), and different combinations of batch and key variable effects (columns).

Computation of QuanT

We have streamlined the process of quantile regression and matrix decomposition within

QuanT by employing strategies for enhanced efficiency. The algorithm executes quantile

regression across various taxon and quantile levels independently, which lends itself well to

parallel processing. We leverage the doParallel package in R to expedite this computation. In

the second stage of QuanT, where SVD is performed, we focus on a few leading singular

vectors because it is generally improbable for a singular vector that accounts for minimal

variance to represent a critical direction established in the first stage. Consequently, it is

sufficient to compute only the leading singular vectors of the binarized residual matrix during

the correlation calibration phase. For this reason, QuanT only calculates the top 15 principal

components using the RSpectra R package.

Tables 1 and 2 present the computational time of QuanT for datasets with different

numbers of samples and taxa. Table 1 illustrates the scalability of QuanT's computational time

with respect to various numbers of samples (n) and taxa (p), benchmarked against SVA and

RUV. For each configuration of n and p, we generated ten datasets using the DM simulation

with parameters $(w_B, w_z) = (0.5, 0.25)$ and the IBDMDB dataset as the template. We also

compared the computational time of QuanT, SVA, and RUV for analyzing the three real-world

datasets (CARDIA, HIVRC, and CRC) (Tab. 2). Overall, QuanT exhibits slower computation

than SVA and RUV due to its multiple quantile regression procedures. Nevertheless, the run

time remains practical and the method is applicable to reasonably large microbiome datasets.

Table 1 | The computation time (in seconds) of QuanT with different numbers of samples

and taxa, compared with SVA and RUV. The data are simulated from the Dirichlet-

Multinomial distribution using the IBDMDB dataset as the template.

	p=100			p=200			p=300		
	SVA	RUV	QuanT	SVA	RUV	QuanT	SVA	RUV	QuanT
n=250	0.4	0.2	37.9	0.9	0.4	83.3	1.5	0.6	138.0
n=500	1.5	0.5	56.5	2.7	0.7	113.9	4.2	1.0	193.2
n=750	4.0	0.6	78.8	6.0	1.3	157.5	8.5	1.6	256.4
n=1000	8.5	0.8	103.3	11.7	1.4	208.0	15.0	1.9	334.7

599 **Table 2 | The computation time (in seconds) of QuanT for the three datasets analyzed,**

600 **compared with SVA and RUV.**

	n	p	SVA	RUV	QuanT
CARDIA	633	145	4.1	2.0	107.0
HIVRC	61	126	0.1	0.6	52.8
CRC	234	166	0.9	5.9	79.1

Discussion

Identifying unmeasured heterogeneity is a challenging yet important task in microbiome data

analysis. Failing to identify and adjust for the unmeasured heterogeneity can drastically

increase the false discovery rate and hinder the detection of true signals. Due to lack of methods

specifically tailored to microbiome data, researchers often use methods developed for genomic

studies. However, these methods do not account for the challenging characteristics of

microbiome data, such as sparsity and overdispersion. In addition, they focus solely on the

information underlying the mean expression and ignore other aspects of the data distribution.

As shown the "Results" section, these methods lead to increased false discoveries and/or

reduced power when analyzing microbiome studies.

We present QuanT, the first comprehensive tool to identify unmeasured heterogeneity

in microbiome data. QuanT-uses quantile regression models and so can thoroughly investigate

the underlying data structure at multiple quantiles. QuanT outputs a set of vectors as surrogates

(QSV) for the unmeasured heterogeneity, which offers users the flexibility to incorporate them

in different types of downstream analysis. These vectors can be treated as known covariates

and adjusted for in regression models. They can also be used in conjunction with a batch effect

removal method, such as ConQuR, to generate microbial profiles without the effects of hidden

heterogeneity.

We applied QuanT to three microbiome datasets and a wide range of simulated datasets

to assess its performance, in comparison with competing methods. The microbiome datasets

include large-scale epidemiological studies with samples processed in different sequencing

runs, and curated microbiome datasets integrated from various publicly available studies. Our

simulations are comprehensive, generating microbiome profiles that mimic real-world

microbiome studies in gut and vaginal environments and using two distinct simulation

mechanisms. Visually and numerically, we showed that QuanT improved subsequent

differential abundance analysis, offering better FDR control and improved statistical power,

and providing results that align better with the gold standard in most setups. Frequently, the

results of differential abundance analysis following QuanT are comparable to those when the

true labels of batches or studies are known. It also offers comparable or improved performance

in predicting the variable of interest, as well as reducing the impact of hidden heterogeneity

while preserving key signals at the community level.

Despite its advantages, QuanT has several limitations and potentials that could be

investigated in future research. First, nowadays, it is common that microbiome studies are

conducted in a longitudinal manner to investigate the dynamics of the microbiome. Such

studies typically produce abundance data collected across multiple time points from the same

individuals. For longitudinal data, one tentative approach is to incorporate time as an

additional adjustment variable into the QuanT framework. However, this approach ignores

the correlations between repeated samples from the same person and may lead to spurious

conclusions. In contrast, in a well-designed study, multiple samples from the same individual

are usually sequenced on the same plate. In such cases, baseline microbiome data can be used

for the identification of unmeasured heterogeneity. Sensitivity analyses may then be

conducted by leveraging microbiome samples that share similar batch structures—for

example, by performing analyses separately on baseline samples and on samples collected at

later time points, thereby assessing the consistency of results across these subsets. Secondly,

QuanT leads to well-controlled false discovery rate in our real and extensive simulation

experiments, although it lacks theoretical guarantee. However, theoretical guarantees might

be derived given strict assumptions about the data generation mechanism, which, in practice,

is usually less important. Finally, a potential extension of QuanT is to identify unmeasured

heterogeneity in gene expression data, especially in single cell and spatial transcriptomics

data which have similar sparsity and over-dispersion as do microbiome data. It would be

worth investigating how QuanT performs when applied to these data types, which might

require some adaptations. However, these aspects are beyond the scope of this paper, and we

leave them to future studies.

Conclusions

In summary, we propose a versatile approach, QuanT, to uncover concealed heterogeneities

within microbiome data. QuanT stands out from existing approaches by detecting hidden data

structures beyond the mean and by easing the constraints of distributional assumptions. When

integrated into the analysis pipeline, QuanT effectively mitigates the confounding influences

of undesired latent variables, enhances false discovery rate control, and preserves statistical

power.

List of abbreviations

QuanT: quantile thresholding

SVA: surrogate variable analysis

RUV: remove unwanted variation

SVD: singular value decomposition

iSV: iterative singular vectors

QSV: quantile surrogate variables

VOI: variable of interest

CARDIA: Coronary Artery Risk Development in Young Adults

CVD: cardiovascular disease

HBP: high blood pressure

TG: true group

NC: no correction

ROC-AUC: area under the receiver operating characteristic curve

PCoA: principal coordinate analysis

HIVRC: HIV re-analysis consortium

CRC: colorectal cancer

IBDMDB: Inflammatory Bowel Disease Multi-omics Database

MOMS-PI: Multi-Omic Microbiome Study: Pregnancy Initiative

DM: Dirichlet-Multinomial

MIDASim: Microbiome Data Simulator

Declarations

688
Ethics approval and consent to participate

Not applicable. Because this study only involved secondary analysis of existing, de-identified

datasets, it is not considered human subject research.

Consent for publication

Not applicable.

Availability of data and material

The CARDIA data can be shared upon request by submitting a CARDIA Data Set Request-

Intent to Analyze Form to the CARDIA Coordinating Center, University of Alabama at

Birmingham, at https://www.cardia.dopm.uab.edu/publications-2/publications-documents.

The HIVRC data are available in Synapse under accession code syn18406854. The CRC data

are from the curatedMetagenomicData package at

https://waldronlab.io/curatedMetagenomicData/. The Inflammatory Bowel Disease Multi-

omics Database (IBDMDB) project[28] and the vaginal microbiome dataset from Multi-Omic

Microbiome Study[29]: Pregnancy Initiative (MOMS-PI) project, which were used as templates

for simulation studies, are provided in the Bioconductor R package, HMP2Data[44].

The R package QuanT is available at https://github.com/jiuyaolu/QuanT in formats

appropriate for Macintosh, Windows, or Linux systems. A vignette demonstrating use of the

package is included (can be accessed at https://jiuyaolu.github.io/QuanT/QuanT.html).

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported in part by R01GM147162 (Lu, Satten, and Zhao), R01HL155417

(Ling), R01GM151301 (Ling) and R01GM155734 (Ling). The Coronary Artery Risk

Development in Young Adults Study (CARDIA) is supported by contracts 75N92023D00002,

75N92023D00003, 75N92023D00004, 75N92023D00005, and 75N92023D00006 from the

National Heart, Lung, and Blood Institute (NHLBI). The HIVRC data used in this study are

from work that was supported by the HIV Microbiome Re-analysis Consortium.

Authors' contributions

JL, GS, NZ and WL developed the method, analyzed the data, and wrote the manuscript. JL

prepared the software package and vignette. LL and KM contributed to the dataset generation

and manuscript development with technical input. All authors read and approved the final

manuscript.

Acknowledgements

The authors thank Drs. Susan A. Tuddenham, James White, Wei Li A Koay, Cynthia L. Sears,

and all members on the HIV Microbiome Re-analysis Consortium for collecting and processing

the HIVRC dataset.

References

1.   Oliveira, F.S. et al. MicrobiomeDB: a systems biology platform for integrating, mining
and analyzing microbiome experiments. *Nucleic acids research* **46**, D684-D691
(2018).
2.   Dhariwal, A., Chong, J., Habib, S., King, I.L., Agellon, L.B. & Xia, J. MicrobiomeAnalyst:
a web-based tool for comprehensive statistical, visual and meta-analysis of
microbiome data. *Nucleic acids research* **45**, W180-W188 (2017).
3.   Stewart, C.J. et al. Temporal development of the gut microbiome in early childhood
from the TEDDY study. *Nature* **562**, 583-588 (2018).
4.   Duvallet, C., Gibbons, S.M., Gurry, T., Irizarry, R.A. & Alm, E.J. Meta-analysis of gut
microbiome studies identifies disease-specific and shared responses. *Nature
communications* **8**, 1784 (2017).
5.   Tuddenham, S.A., Koay, W.L.A., Zhao, N., White, J.R., Ghanem, K.G. & Sears, C.L. The
impact of human immunodeficiency virus infection on gut microbiota α-diversity: an
individual-level meta-analysis. *Clinical Infectious Diseases* **70**, 615-627 (2020).
6.   Ma, S. et al. Population structure discovery in meta-analyzed microbial communities
and inflammatory bowel disease using MMUPHin. *Genome Biology* **23**, 208 (2022).
7.   Austin, G.I. et al. Processing-bias correction with DEBIAS-M improves cross-study
generalization of microbiome-based prediction models. *Nature Microbiology*, 1-15
(2025).
8.   Ling, W. et al. Batch effects removal for microbiome data via conditional quantile
regression. *Nature Communications* **13**, 1-14 (2022).
9.   Leek, J.T. & Storey, J.D. Capturing heterogeneity in gene expression studies by
surrogate variable analysis. *PLoS genetics* **3**, e161 (2007).
10. Risso, D., Ngai, J., Speed, T.P. & Dudoit, S. Normalization of RNA-seq data using factor
analysis of control genes or samples. *Nature biotechnology* **32**, 896-902 (2014).
11. Koenker, R. & Bassett Jr, G. Regression quantiles. *Econometrica: journal of the
Econometric Society*, 33-50 (1978).

12.   Leek, J.T. & Storey, J.D. A general framework for multiple testing dependence.
761   *Proceedings of the National Academy of Sciences* **105**, 18718-18723 (2008).

762   13.   Leek, J.T. et al. Tackling the widespread and critical impact of batch effects in high-
763   throughput data. *Nature Reviews Genetics* **11**, 733-739 (2010).

764   14.   Leek, J.T., Johnson, W.E., Parker, H.S., Jaffe, A.E. & Storey, J.D. The sva package for
765   removing batch effects and other unwanted variation in high-throughput
766   experiments. *Bioinformatics* **28**, 882-883 (2012).

767   15.   Leek, J.T. Svaseq: removing batch effects and other unwanted noise from sequencing
768   data. *Nucleic acids research* **42**, e161-e161 (2014).

769   16.   Koenker, R. & Hallock, K.F. Quantile regression. *Journal of economic perspectives* **15**,
770   143-156 (2001).

771   17.   Barber, R.F. & Candès, E.J. Controlling the false discovery rate via knockoffs. (2015).

772   18.   Goldstein, S., Lebowitz, J.L., Tumulka, R. & Zanghi, N. Any orthonormal basis in high
773   dimension is uniformly distributed over the sphere. (2017).

774   19.   Lilliefors, H.W. On the Kolmogorov-Smirnov test for normality with mean and
775   variance unknown. *Journal of the American statistical Association* **62**, 399-402
776   (1967).

777   20.   Anderson, T.W. & Darling, D.A. Asymptotic theory of certain" goodness of fit" criteria
778   based on stochastic processes. *The annals of mathematical statistics*, 193-212
779   (1952).

780   21.   Royston, P. A pocket–calculator algorithm for the shapiro–francia test for non–
781   normality: An application to medicine. *Statistics in medicine* **12**, 181-184 (1993).

782   22.   Storey, J.D. A direct approach to false discovery rates. *Journal of the Royal Statistical
783   Society Series B: Statistical Methodology* **64**, 479-498 (2002).

784   23.   Storey, J.D. The positive false discovery rate: a Bayesian interpretation and the q-
785   value. *The annals of statistics* **31**, 2013-2035 (2003).

786   24.   Storey, J.D. & Tibshirani, R. Statistical significance for genomewide studies.
787   *Proceedings of the National Academy of Sciences* **100**, 9440-9445 (2003).

788   25.   Gutenbrunner, C., Jurečková, J., Koenker, R. & Portnoy, S. Tests of linear hypotheses
789   based on regression rank scores. *Journaltitle of Nonparametric Statistics* **2**, 307-331
790   (1993).

791   26.   Efron, B., Tibshirani, R., Storey, J.D. & Tusher, V. Empirical Bayes analysis of a
792   microarray experiment. *Journal of the American statistical association* **96**, 1151-1160
793   (2001).

794   27.   Integrative, H. The Integrative Human Microbiome Project: dynamic analysis of
795   microbiome-host omics profiles during periods of human health and disease. *Cell
796   host & microbe* **16**, 276-289 (2014).

797   28.   Lloyd-Price, J. et al. Multi-omics of the gut microbial ecosystem in inflammatory
798   bowel diseases. *Nature* **569**, 655-662 (2019).

799   29.   Fettweis, J.M. et al. The vaginal microbiome and preterm birth. *Nature medicine* **25**,
800   1012-1021 (2019).

801   30.   He, M., Zhao, N. & Satten, G.A. MIDASim: a fast and simple simulator for realistic
802   microbiome data. *Microbiome* **12**, 135 (2024).

31. Hu, Y.-J. & Satten, G.A. Testing hypotheses about the microbiome using the linear
decomposition model (LDM). *Bioinformatics* **36**, 4106-4115 (2020).
32. Friedman, G.D. et al. CARDIA: study design, recruitment, and some characteristics of
the examined subjects. *Journal of clinical epidemiology* **41**, 1105-1116 (1988).
33. Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A. & Holmes, S.P.
DADA2: High-resolution sample inference from Illumina amplicon data. *Nature*
methods **13**, 581-583 (2016).
34. Callahan, B. Silva taxonomic training data formatted for DADA2 (Silva version 132).
Zenodo (2018).
35. Daquigan, N., Seekatz, A.M., Greathouse, K.L., Young, V.B. & White, J.R. High-
resolution profiling of the gut microbiome reveals the extent of Clostridium difficile
burden. *NPJ biofilms and microbiomes* **3**, 35 (2017).
36. Lozupone, C.A. et al. Alterations in the gut microbiota associated with HIV-1
infection. *Cell host & microbe* **14**, 329-339 (2013).
37. Villar-García, J. et al. Impact of probiotic Saccharomyces boulardii on the gut
microbiome composition in HIV-treated patients: a double-blind, randomised,
placebo-controlled trial. *PloS one* **12**, e0173802 (2017).
38. Feng, Q. et al. Gut microbiome development along the colorectal adenoma–
carcinoma sequence. *Nature communications* **6**, 6528 (2015).
39. Thomas, A.M. et al. Metagenomic analysis of colorectal cancer datasets identifies
cross-cohort microbial diagnostic signatures and a link with choline degradation.
Nature medicine **25**, 667-678 (2019).
40. Pasolli, E. et al. Accessible, curated metagenomic data through ExperimentHub.
Nature methods **14**, 1023-1024 (2017).
41. Beghini, F. et al. Integrating taxonomic, functional, and strain-level profiling of diverse
microbial communities with bioBakery 3. *elife* **10**, e65088 (2021).
42. Zhou, H., He, K., Chen, J. & Zhang, X. LinDA: linear models for differential abundance
analysis of microbiome compositional data. *Genome biology* **23**, 95 (2022).
43. Hu, Y.-J. & Satten, G.A. Compositional analysis of microbiome data using the linear
decomposition model (LDM). *Bioinformatics* **39**, btad668 (2023).
44. Stansfield, J., Smirnova, E., Zhao, N., Fettweis, J., Waldron, L. & Dozmorov, M.
HMP2Data: 16s rRNA sequencing data from the Human Microbiome Project 2. *R*
package version **1** (2020).
836