Intercellular compartmentation of trehalose 6-phosphate metabolism

in *Setaria viridis* leaves

Tomás Tonetti[1,#], Bruno E. Rojas[1,#], Leandro E. Lucero[1], Regina Feil[2], Paula Calace[3], Mariana Saigo[3], Stéphanie Arrivault[2], Mark Stitt[2], John E. Lunn[2], Carlos M. Figueroa[1,*]

[1]Instituto de Agrobiotecnología del Litoral, UNL, CONICET, FBCB, Santa Fe, Argentina

[2]Max Planck Institute of Molecular Plant Physiology, Potsdam-Golm, Germany

[3]Centro de Estudios Fotosintéticos y Bioquímicos, UNR, CONICET, FCBF, Rosario, Argentina

[#]Equal contribution

[*]Correspondence: carfigue@fbcb.unl.edu.ar

ORCID

TT, 0000-0002-3820-9411; BER, 0000-0002-3126-8363; LEL, 0000-0003-4453-8964; RF, 0000-0002-9936-1337; MSaigo, 0000-0002-2312-8782; SA, 0000-0003-0516-6950; MStitt, 0000-0002-4900-1763; JEL, 0000-0001-8533-3004; CMF, 0000-0003-4047-0480.

Running title

Compartmentation of Tre6P metabolism in *Setaria viridis*

Date of submission: 16[th] August 2024

Number of figures: 6

Supplementary data: 9 tables and 9 figures

Word count: 7458

Highlight

Trehalose 6-phosphate, a sugar signalling metabolite, is present mainly in the bundle sheath cells of *Setaria viridis* leaves, implicating it in the regulation of phloem loading.

Abstract

Trehalose 6-phosphate (Tre6P) is a signal metabolite that links carbon metabolism with plant development. Our current understanding of Tre6P metabolism and signalling is predominantly based on studies in *Arabidopsis thaliana*. Some features could be adapted to the specific physiology, anatomy, and life cycle of this C_3 eudicot model species, and thus might not be representative of other angiosperms. To better understand the regulation of carbon metabolism by Tre6P in monocot C_4 species we used *Setaria viridis*, which has been widely adopted as a model for the major C_4 NADP-malic enzyme subtype crop species, such as maize (*Zea mays*), sorghum (*Sorghum bicolor*) and sugarcane (*Saccharum officinarum*). In this work, we analysed the levels of transcripts encoding Tre6P-related enzymes in different tissues and cell types from *S. viridis*. The *TREHALOSE-6-PHOSPHATE SYNTHASE1* transcript, its encoded protein (TPS1, the enzyme responsible for Tre6P synthesis) and Tre6P were mainly located in bundle sheath cells of *S. viridis*. Our results show that Tre6P is predominately synthesized and located in bundle sheath and associated cells, where it could play a fundamental role in the regulation of sucrose levels by modulating phloem loading.

Keywords

bundle sheath cells, C$_4$ photosynthesis, mesophyll cells, phloem loading, *Setaria viridis*, sucrose, trehalose 6-phosphate

Abbreviations

BSC, bundle sheath cells; MC, mesophyll cells; NADP-MDH, NADP-dependent malate dehydrogenase; NADP-ME, NADP-dependent malic enzyme; PEPC, phospho*enol*pyruvate carboxylase; PRK, phosphoribulokinase; SWEET, SUGARS WILL EVENTUALLY BE EXPORTED TRANSPORTERS; Tre6P, trehalose 6-phosphate; TPS, Tre6P synthase; TPP, Tre6P phosphatase; TRE, trehalase.

Introduction

Trehalose is a non-reducing disaccharide widely distributed in nature (Figueroa and Lunn, 2016). In plants, the pathway of trehalose synthesis occurs through the phosphorylated intermediate trehalose 6-phosphate (Tre6P), which is synthesized from UDP-glucose (UDPGlc) and glucose 6-phosphate (Glc6P) by Tre6P synthase (TPS; EC 2.4.1.15) and dephosphorylated to trehalose by Tre6P phosphatase (TPP; EC 3.1.3.12; Cabib and Leloir, 1958; Avonce *et al.*, 2006). Trehalose is then hydrolysed to glucose by trehalase (TRE; EC 3.2.1.28). Gene families coding for TPS and TPP are present in species from all of the main taxonomic groups of green plants (Avonce *et al.*, 2006; Lunn, 2007; Lunn *et al.*, 2014; Figueroa and Lunn, 2016). Plant TPSs are divided into two distinct clades: class I and class II. Class I TPSs are responsible for Tre6P synthesis, while class II TPSs (which lack some of the glycosyltransferase active site residues involved in substrate binding and catalysis) are putatively involved in Tre6P perception and/or signalling (Avonce *et al.*, 2006; Lunn, 2007; Vandesteene *et al.*, 2010; Lunn *et al.*, 2014; Delorge *et al.*, 2015; Figueroa and Lunn, 2016; Van Leene *et al.,* 2022). In the last two decades, studies performed with mutant and transgenic lines demonstrated the importance of Tre6P for all stages of plant development (Eastmond *et al.*, 2002; van Dijken *et al.*, 2004; Gomez *et al.*, 2006; Gomez *et al.*, 2010; Wahl *et al.*, 2013; Fichtner *et al.*, 2020; Ponnu *et al.*, 2020; reviewed in Fichtner and Lunn, 2021).

Schluepmann *et al.* (2003) demonstrated the critical role played by Tre6P in plants by over-expressing the genes coding for the *Escherichia coli* TPS (otsA) and TPP (otsB) in *Arabidopsis thaliana*. Later, various genetic and biochemical approaches showed that Tre6P is a sucrose-signalling metabolite that functions as an indicator of the metabolic status of the plant (Lunn *et al*., 2006; Carillo *et al.*, 2013; Martins *et al.*, 2013; Nunes *et al*., 2013; Yadav *et al.*, 2014). The Tre6P-sucrose nexus model postulates that Tre6P is both a signal and a

negative regulator of sucrose levels in plants, thus keeping its concentration within an optimal

range (Lunn *et al.*, 2014; Yadav *et al.*, 2014; Figueroa *et al.* 2016; Figueroa and Lunn, 2016).

In *A. thaliana* leaves, Tre6P is synthesized primarily in phloem parenchyma (bundle

sheath) cells and in the companion cell-sieve element complex of the phloem (Fichtner *et al.*,

2020). The phloem parenchyma cells are symplastically connected with mesophyll cells

(MC) via plasmodesmata, allowing Tre6P to diffuse into the MC to regulate sucrose

production (Fichtner and Lunn, 2021). Tre6P regulates the synthesis and transport of sucrose

by influencing photoassimilate partitioning between sucrose or organic and amino acids

during the day (Figueroa *et al.*, 2016), transitory starch degradation during the night and

towards the end of the light period (Martins *et al.*, 2013; dos Anjos *et al.*, 2018; Ishihara *et*

al., 2022), and expression of *SWEET* (*SUGARS WILL EVENTUALLY BE EXPORTED*

TRANSPORTER) genes, which encode transporters involved in sucrose phloem loading

(Oszvald *et al.*, 2018; Fichtner *et al.*, 2021). Therefore, Tre6P coordinates the synthesis and

transport of sucrose and amino acids from source leaves with the demand from growing and

sink tissues (Martins *et al.*, 2013; Figueroa *et al.*, 2016; dos Anjos *et al.*, 2018). In addition,

Tre6P exerts widespread transcriptional repression of photosynthesis, gluconeogenesis and

carbon starvation responses, induces amino acid synthesis, nucleotide synthesis and ribosome

formation and interacts with light signaling and many phytohormone signaling pathways

(Avidan *et al.*, 2024).

Our current understanding of Tre6P metabolism and signalling in plants is mainly

based on studies in *A. thaliana* (Fichtner and Lunn, 2021; Rojas *et al.* 2023). This model

eudicot is a rosette-forming, annual weed species that performs C_3 photosynthesis and grows

predominantly in temperate zones. Some features of Tre6P metabolism and signalling in *A.*

thaliana might be adapted to its specific physiology, anatomy and life cycle, and so might

differ from other species. Tre6P metabolism and signalling have also been investigated in

maize (*Zea mays*), a domesticated monocot species that performs C_4 photosynthesis and originated in subtropical regions. These studies have revealed the importance of Tre6P metabolism and signalling in the coordination of metabolism in source and sink zones, inflorescence and shoot branching, floral identity, tillering, seed development and tolerance to water deficit (Satoh-Nagasawa *et al.*, 2006; Henry *et al.*, 2014, 2015; Nuccio *et al.*, 2015; Czedik-Eysenberg *et al.*, 2016; Bledsoe *et al.*, 2017; Oszvald *et al.*, 2018; Claeys *et al.*, 2019; Dong *et al.*, 2019; Klein *et al.*, 2022). However, how far Tre6P metabolism and signalling are adapted to the specialized C_4 photosynthetic metabolism and leaf anatomy of this species is poorly understood.

The C_4 cycle is a carbon-concentrating mechanism that increases the intracellular concentration of CO_2 around ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco), thereby suppressing the oxygenation reaction of this enzyme and the need for photorespiration. The C_4 pathway increases photosynthetic efficiency under warm and high-light conditions, and C_4 plants generally have higher water- and nitrogen-use efficiencies than C_3 plants (Trípodi *et al.* 2021; Rojas *et al.* 2023). C_4 photosynthesis has evolved independently multiple times in different plant taxonomic groups and this involved a series of anatomical and biochemical adaptations (Sage, 2004). In the MC of C_4 plants, phospho*enol*pyruvate carboxylase (PEPC) performs a primary fixation of CO_2 into oxaloacetate, a C_4 acid, which is then converted into malate or aspartate by the action of NADP-malate dehydrogenase (NADP-MDH) or aspartate aminotransferase, respectively. These C_4 acids diffuse to the adjacent bundle sheath cells (BSC), where decarboxylation of C_4 acids is followed by a secondary fixation via Rubisco and the Calvin-Benson cycle (Maier *et al.*, 2011; Wang *et al.*, 2014). According to the principal enzyme used in the decarboxylation step, C_4 plants are broadly classified into NADP-malic enzyme (NADP-ME), NAD-malic enzyme (NAD-ME), and phospho*enol*pyruvate carboxykinase (PEPCK)

subtypes (Hatch *et al.*, 1975). Nevertheless, C_4 plants can use various carbon fixation

pathways at the same time, and these can vary depending on the physiological and

environmental conditions (Chapman and Hatch, 1981; Wang *et al.*, 2014; Sales *et al.*, 2018;

Cacefo *et al.*, 2019).

Tre6P regulates photoassimilate partitioning between sucrose and organic acids and

amino acids in *A. thaliana* via coordinated changes in the activities of nitrate reductase, PEPC

and, by inference, the mitochondrial pyruvate dehydrogenase (Figueroa *et al*., 2016).

Artificially increased levels of Tre6P reduced the mono-ubiquitination and increased the

phosphorylation status of PEPC, thus activating the enzyme. These changes increased carbon

flux into organic acids and the TCA cycle to provide carbon skeletons for amino acid

synthesis. Given that PEPC is responsible for primary carbon fixation in the C_4 pathway,

Tre6P might therefore play a direct role in regulating photosynthetic carbon fixation in C_4

plants. This raises the question of whether Tre6P is also involved in the regulation of

photoassimilate partitioning to amino acids in C_4 plants and, if so, how flux into metabolites

in the C_4 cycle and flux into the TCA cycle and amino acid synthesis are reconciled. Tre6P

also regulates the mobilization of transitory starch reserves in *A. thaliana* leaves and

provision of sucrose at night (Martins *et al*., 2013; dos Anjos *et al*., 2018; Ishihara *et al*.,

2022). In contrast to C_3 plants, where sucrose and starch synthesis both occur within MC, in

C_4 plants starch synthesis is restricted to the BSC, while sucrose is made predominantly in

MC (Lunn and Furbank, 1997; 1999). If Tre6P also regulates photoassimilate partitioning and

transitory starch turnover in C_4 plants, this cellular compartmentation of sucrose and starch

metabolism would add a further layer of complexity. It has been proposed that expression of

the genes encoding the SWEET and SUT/SUC (sucrose-H^+ symporters) transporters had to

be adapted to the specialized Kranz anatomy in C_4 species, and was potentially a critical

element in the evolution of C_4 plants (Furbank and Kelly, 2021). Mutation of the three maize

ZmSWEET13 genes produced plants with impaired sugar phloem loading and drastic effects

on growth (Bezrutczyk *et al.*, 2018). As Tre6P has been implicated in the transcriptional

regulation of *SWEET* genes (Oszvald *et al*., 2018; Fichtner *et al*., 2021), it is possible that

Tre6P also plays a key role in modulating phloem loading during the establishment of Kranz

anatomy in developing leaves of C_4 species.

164   Currently, we have little knowledge about Tre6P signalling in the leaves of C_4 plants

(Rojas *et al.*, 2023). By extrapolation from its known functions in leaves of C_3 plants, we

hypothesize that Tre6P plays a role in the regulation of carbon fixation, photoassimilate

partitioning and phloem loading in the leaves of C_4 plants. If so, Tre6P signalling in these

species may have unique features, reflecting their specialized metabolism and anatomy. To

better understand how Tre6P signalling operates in C_4 crops, we investigated the

compartmentation of Tre6P in leaves of *Setaria viridis* (wild foxtail millet), a model species

for translational research in C_4 plants (Brutnell *et al.*, 2010; Doust *et al.*, 2019). Our results

show that Tre6P metabolism takes place mainly in the BSC, where it could modulate sucrose

levels by controlling phloem loading.

Materials and methods

Plant growth and harvest conditions

Seeds of *Setaria viridis* (L.) P. Beauv. accession A10.1 were obtained from the Germplasm

Resources Information Network (GRIN; http://www.ars-grin.gov/), United States Department

of Agriculture (PI 669942). To increase the germination rate, the seed coat was abraded with

sandpaper (Van Eck and Swartwood, 2015). Seeds were surface sterilized with 0.55 g l^{-1}

sodium hypochlorite containing 0.1% (v/v) Tween-20 for 5 minutes, followed by three

washes with sterile deionized water. The seeds were then sown on plates with half-strength

Murashige and Skoog medium (PhytoTech Labs; https://phytotechlab.com/) supplemented

with 0.8% (w/v) agar. Plates were placed in a growth chamber at 28°C with an irradiance of

120 µmol m^{-2} s^{-1}, provided by white fluorescent tubes, under a long-day photoperiod regime

(16 h light/8 h dark). After five days, seedlings were transferred to individual 8-cm diameter

pots containing Klasmann TS1 substrate and irrigated with 1X Hoagland solution. Plants

were grown in a Conviron Adaptis A1000 controlled environment chamber

(https://www.conviron.com/) with an irradiance of 350 µmol m^{-2} s^{-1}, provided by white

fluorescent tubes, with a long-day photoperiod regime, at 28/22°C (day/night), and 50%

relative humidity for 2 weeks. For metabolite fractionation experiments, plants were grown in

a walk-in phytotron with an irradiance of 550 µmol m^{-2} s^{-1}, provided by LEDs, at 28/22°C

(day/night) and 65% relative humidity. Harvested plants had eight fully developed leaves and

inflorescences were not macroscopically visible. The blades from the 5th and 6th true leaves

(counting from the bottom of the plant) were harvested in the middle of the photoperiod. The

leaf material used for isolation of MC and BSC was immediately processed and then stored at

-80°C. For enzymatic and metabolite analyses, leaf samples were rapidly quenched in liquid

nitrogen and stored at -80°C until use. Samples for the analysis of transcripts and proteins

(seedlings, inflorescences, the fourth internode and the flag leaf) were frozen in liquid

nitrogen and stored at -80°C until use.

Phylogenetic analysis

TPS, TPP and TRE protein sequences from *S. viridis* (genome version 2.1) were identified

using the keyword "trehalose" at the Phytozome 13 server (https://phytozome.jgi.doe.gov/).

We followed the same approach to retrieve TPS and TPP protein sequences from *Ananas*

comosus (genome version 3) and *Musa acuminata* (genome version 1). Sequences were

manually curated to remove incomplete and duplicated entries. TPS and TPP protein

sequences from *A. thaliana*, *Brachypodium distachyon*, *Oryza sativa*, *Phaseolus vulgaris*,

Populus trichocarpa and *Z. mays* were obtained from previous publications (Yang *et al.*,

2012; Henry *et al.*, 2014; Barraza *et al.*, 2016; Wang *et al.*, 2019). Sequences were aligned

with the CLUSTALW tool available at the GenomeNet server (https://www.genome.jp/tools-

bin/clustalw). Distance-based phylogenetic trees were reconstructed with the SeaView

software version 4.3.0 (Gouy *et al.*, 2010) using the BioNJ algorithm (Gascuel, 1997) and

1000 replicates. Figures were prepared with the FigTree program version 1.3.1

(http://tree.bio.ed.ac.uk/software/figtree/). Sequences used to reconstruct the trees were

labelled as shown in Supplementary Table S1.

Separation of mesophyll and bundle sheath cells

Fractions highly enriched in MC or BSC strands were obtained from fresh leaves following

the methods described by John *et al.* (2014), with the modifications introduced by Calace *et*

al. (2021). In both cases, each replicate consisted of the material obtained from 20 plants.

These samples were used for the analysis of transcripts and proteins.

To determine the distribution of metabolites, frozen leaf tissue was finely ground in

liquid nitrogen and the frozen tissue powder was passed through a series of nylon meshes of

different pore sizes (200, 80, and 41 µm). The material that was retained on each filter and

the one that passed through the 41 µm filter were collected to obtain the samples named

F200, F80, F40, and FT, respectively. The F200 fraction, depending on the grinding, often

has larger whole leaf bits but tends to be enriched in BSC; the F80 fraction is especially

enriched in BSC; the F40 fraction is usually a mix of BSC and MC; and FT fraction is

especially enriched in MC (Stitt and Heldt, 1985). This method is suitable for the

quantification of metabolites with short half-lives (Arrivault *et al.*, 2009, 2017). Metabolites

and enzymes, including cell-specific marker enzymes, were measured in each fraction, and

their distribution between MC and BSC were calculated by linear regression, as described by

Stitt and Heldt (1985). Two C_4 photosynthetic enzymes were used as markers for each cell-

type: PEPC and NADP-MDH for MC and NADP-ME and phosphoribulokinase (PRK) for

BSC. Each biological replicate consisted of the material obtained from 20 plants.

Total RNA extraction and cDNA synthesis

Total RNA was extracted from 50 mg fresh weight (FW) of finely ground tissue with the

TransZol reagent (TransGen Biotech; https://www.transgenbiotech.com/), following the

manufacturer's instructions. After RNA extraction, samples were treated with DNase

(Promega; https://www.promega.com) and RNA concentration was quantified using a

NanoDrop 2000 (Thermo Scientific; https://www.thermofisher.com). cDNA was synthesized

with the MultiScribe Reverse Transcriptase (Applied Biosystems;

https://www.thermofisher.com), according to the manufacturer's instructions.

Determination of transcript levels

Real-time reverse transcriptase-qPCR (RT-qPCR) reactions were performed in a final volume

of 8 µl in 96-well microplates (Applied Biosystems) with the iQ SYBR Green Supermix

(Bio-Rad; https://www.bio-rad.com), 0.2 µM of each primer, and an appropriate dilution of

the sample. Transcript abundance was calculated with the comparative C_t method

(Schmittgen and Livak, 2008) and normalization was performed with the *SvKIN* transcript

(Sevir.4G252700; Martins *et al.*, 2016), which showed similar C_t values across all the

analysed samples (Supplementary Fig. S1). All the primers used in this work are listed in

Supplementary Table S2.

 Production of a polyclonal antiserum against SvTPS1

The protein coding region of the class I *TPS* gene (SvTPS1; Uniprot ID A0A4U6VAG9) was

codon-optimized and *de novo* synthesized (BioBasic; https://www.biobasic.com/) for

expression in *E. coli* and cloned into the pET28b expression vector (Novagen;

https://www.merckmillipore.com) between the NdeI and SacI restriction sites, in frame with a

His$_6$-tag at the N-terminus. The resulting vector was used to transform *E. coli* BL21 (DE3)

cells (Invitrogen; https://www.thermofisher.com). Transformed cells were grown in

Lysogeny Broth medium, supplemented with 50 µg ml^{-1} kanamycin at 37°C, with shaking at

200 rpm until the optical density at 600 nm reached approximately 0.6. Synthesis of the

recombinant protein was induced by addition of isopropyl-β-D-thiogalactopiranoside to a

final concentration of 1 mM and cells were incubated for 16 h at 25°C and 200 rpm after

induction. The recombinant SvTPS1 protein was mainly present in the insoluble fraction;

thus, it was purified from inclusion bodies (Jenö and Horst, 2002). This protein was used to

raise a polyclonal antiserum in rabbits at ICIVET Litoral (UNL-CONICET, Argentina).

Antibodies were affinity-purified using a nitrocellulose membrane strip containing the

recombinant SvTPS1, as described by Fang (2012).

Protein extraction

Plant material was homogenized in a pre-cooled mortar with liquid nitrogen. For denaturing

protein extraction, 20 mg FW of frozen tissue were extracted with 200 µl of 125 mM Tris-

HCl pH 6.8, 2% (w/v) sodium dodecyl sulphate (SDS), 20% (v/v) glycerol, 1.4 M 2-

mercaptoethanol, and 0.05% (w/v) Bromophenol Blue. After adding the buffer, samples were

vortex-mixed and heated for 5 min at 95°C with agitation. Samples were then cooled to room

temperature and centrifuged at 20,000 x *g* for 10 min to separate the protein extract from

insoluble tissue debris. Alternatively, 50 mg FW of frozen tissue were resuspended in 500 µl

of 10 mM Tris-HCl pH 8.0, 10% (v/v) trichloroacetic acid (TCA), 25 mM ammonium

acetate, and 1 mM EDTA (Rojas *et al.*, 2021). Samples were incubated at 4°C for 30 min and

centrifuged at 20,000 x *g* for 15 min. The supernatant was discarded and the pellet was

washed with 500 µl of 5% (w/v) TCA and then with 500 µl of 90% (v/v) acetone, then

allowed to dry before the protein pellet was dissolved in 50 µL of Laemmli buffer for

analysis by SDS-PAGE.

Protein extraction for enzymatic assays was performed as described by Gibon *et al.*

(2004), with minor modifications. Samples (20 mg FW of frozen tissue) were supplemented

with 10 mg of polyvinylpolypyrrolidone and then resuspended in 500 µl of 50 mM HEPES-

KOH pH 7.5, 10 mM $MgCl_2$, 1 mM EDTA, 1 mM EGTA, 1 mM benzamidine, 1 mM ε-

aminocaproic acid, 0.02 mM leupeptin, 1 mM DTT, 1 mM phenylmethylsulfonyl fluoride,

0.1% (v/v) Triton X-100, and 20% (v/v) glycerol. Samples were incubated on ice for 10 min

and then centrifuged at 20,000 x *g* and 4°C for 10 min to separate soluble proteins from tissue

debris.

Protein concentration was determined with the Bradford reagent (Bradford, 1976),

using a standard curve constructed with bovine serum albumin.

Immunodetection of SvTPS1

Protein extracts were subjected to a 10% SDS-PAGE and electrotransferred to a 0.45 µm

nitrocellulose membrane (Amersham; https://www.cytivalifesciences.com/). To check lane

loadings, membranes were stained with 0.01% (w/v) Ponceau S in 1% (v/v) acetic acid. After

washing with TBST solution [50 mM Tris-HCl pH 8.0; 150 mM NaCl; 0.2% (v/v) Tween-

20], membranes were blocked for 1 h with TBST supplemented with 0.2% (w/v) fat-free

dried milk powder. Membranes were incubated overnight at 4°C with purified anti-SvTPS1

in TBST supplemented with 0.2% (w/v) fat-free dried milk powder. After washing, the

membranes were incubated for 30 min at room temperature with goat anti-rabbit IgG (H&L)

conjugated to horseradish peroxidase (Agrisera AS09 602; https://www.agrisera.com/).

Proteins were detected using the ECL Bright reagent (Agrisera AS16 ECL-N) and exposed to

X-ray films (AGFA; https://www.agfa.com/corporate/) for 1 min. The sizes of the

immunoreactive bands were estimated from the relative mobility (Rf) *versus* logarithm of

molecular weight (logMM) plot constructed with protein standards, using data from three

independent experiments.

Metabolite extraction and measurement

Phosphorylated intermediates and organic acids were extracted with a mixture of

chloroform:methanol (3:7) and then measured by anion-exchange high-performance liquid

chromatography coupled to tandem mass spectrometry (LC-MS/MS), as described by Lunn *et*

al. (2006), with the modifications introduced by Figueroa *et al.* (2016). Calvin-Benson cycle

intermediates and aspartate were measured in chloroform-methanol extracts by ion-pair

reverse phase LC-MS/MS, as described by Arrivault *et al.* (2009). Pyruvate,

phospho*enol*pyruvate (PEP), dihydroxyacetone phosphate (DHAP) and 3-phosphoglycerate

(3PGA) were enzymatically measured in TCA extracts (Arrivault *et al.*, 2017). The content of

sucrose, glucose, and fructose in ethanolic extracts was determined following the method of

Stitt *et al.* (1989), while the amount of starch was measured using the insoluble residue from

the ethanolic extraction, as described by Hendriks *et al.* (2003). The datasets containing

metabolite levels and the associated calculations are available in Supplementary Tables S3-

S5.

Enzyme activity assays

PRK activity was measured spectrophotometrically using a continuous assay (Leegood,

1990), with minor modifications. The assay mixture contained: 50 mM Tricine-KOH pH 8.0,

1 mM ATP, 50 mM KCl, 0.3 mM NADH, 10 mM $MgCl_2$, 2 mM PEP, 2 U pyruvate kinase,

and 2 U lactate dehydrogenase and the reaction was started by adding 0.5 mM ribulose 5-

phosphate. PEPC, NADP-MDH, and NADP-ME activities were measured as described by

Ashton *et al.* (1990), with minor modifications. The PEPC assay mixture contained: 50 mM

Tricine-KOH pH 8.0, 5 mM $MgCl_2$, 2 mM DTT, 1 mM $KHCO_3$, 5 mM Glc6P, 0.3 mM

NADH, and 0.4 U malate dehydrogenase and the reaction was started with 2 mM PEP. The

NADP-MDH assay mixture contained: 50 mM Tricine-KOH pH 8.0, 70 mM KCl, 1 mM

EDTA, 1 mM DTT, and 0.3 mM NADPH and the reaction was started with 1 mM

oxaloacetate (freshly prepared). The NADP-ME assay mixture contained: 50 mM Tricine-

KOH pH 8.0, 0.5 mM $NADP^+$, 0.1 mM EDTA, and 2 mM $MgCl_2$ and the reaction was

started with 5 mM malate. All reactions were performed in a final volume of 200 µl at 25°C

in 96-well microplates. The reduction/oxidation of $NAD(P)^+$/NAD(P)H was followed by

measuring the absorbance at 340 nm in a Biotek ELx808 microplate reader. The amount of

crude extract added to the reaction was chosen to ensure a linear response of the signal. One

unit (U) of enzyme activity is defined as the amount of extract producing 1 µmol of product

in 1 min under the specified assay conditions.

Results

Phylogenetic analysis of Tre6P-related enzymes

To identify the proteins involved in Tre6P metabolism in *S. viridis*, we performed keyword

searches against its deduced proteome. We found 21 putative Tre6P-related enzymes: 10

TPSs, 10 TPPs, and 1 TRE (Supplementary Table S1), as previously observed for *Setaria*

italica (Henry et al., 2020), a domesticated relative of *S. viridis*. After removing duplicated

sequences or splicing variants (for SvTPP1, SvTPP8, and SvTPP10), phylogenetic trees were

reconstructed for TPS (Fig. 1) and TPP (Fig. 2) proteins.

S. viridis TPS sequences were split into two major clades containing class I and class

II TPSs, as previously described for TPSs from *A. thaliana*, poplar, rice, pea and maize

(Leyman *et al.*, 2001; Avonce *et al.*, 2006; Lunn, 2007; Yang *et al.*, 2012; Henry *et al.*, 2014;

Barraza *et al.*, 2016). *S. viridis* has only one class I TPS (SvTPS1), which diverges from the

other nine class II TPS proteins (SvTPS2-10; Fig. 1). All *S. viridis* TPS sequences displayed

the glycosyl-transferase family 20 domain (InterPro ID IPR001830) and the TPP domain

(InterPro ID IPR003337) that are typical of plant TPS proteins. Inspection of the primary

sequence of *S. viridis* TPSs in detail showed that only SvTPS1 had the characteristic residues

involved in UDPGlc and Glc6P binding (Supplementary Fig. S2). The class II TPS sequences

from *S. viridis* (SvTPS2-10) were distributed into several branches, most of which contained

proteins from eudicot and monocot species. The exceptions were the monocot-specific sub-

clade containing SvTPS2/3/9 and that containing *M. acuminata* TPS8 and TPS16, which

branched at the base of the class II TPS tree (Fig. 1), an observation initially made by Lunn

(2007). A detailed inspection of *S. viridis* class II TPS sequences showed substitutions in the

substrate-binding residues of the glucosyltransferase domain that might explain their apparent

lack of catalytic activity (Supplementary Fig. S2; Gibson *et al.*, 2002; Vandesteene *et al.*,

2010). The phylogenetic and sequence analyses indicate that SvTPS1 is the enzyme solely

responsible for Tre6P synthesis in *S. viridis*.

The *S. viridis* TPP family was separated into two major clades: three in Clade A and

the other seven in Clade B (Fig. 2). All TPP sequences from *S. viridis* had the TPP domain

(InterPro ID IPR003337) and the three conserved active site motifs previously described for

these proteins (Avonce *et al.*, 2006; Lunn, 2007; Vandesteene *et al.*, 2012; Henry *et al.*, 2014;

Kerbler *et al.*, 2023); thus, all 10 *S. viridis* TPP proteins are likely to be catalytically active

(Supplementary Fig. S3). Among these are representatives of the monocot-specific

RAMOSA3-type sub-clade (Fig. 2), which modify inflorescence branching, sex

determination and tillering in maize (Satoh-Nagasawa et al., 2006; Claeys et al. 2019; Klein

et al. 2022), independently of their catalytic activity. Particularly, SvTPP10 and SvTPP2, the

orthologues of RAMOSA3 (ZmTPP10) and ZmTPP4, respectively (Fig. 2).

Differential expression and compartmentation of Tre6P-related transcripts

To analyse the expression pattern of the identified *S. viridis TPS*, *TPP* and *TRE* genes, we

performed real-time RT-qPCR analysis of mRNA extracted from the aerial part of five-days

old seedlings, whole leaves (pool of leaves 5 and 6), the flag leaf, whole inflorescences, and

the fourth internode of the stem (all samples harvested in the middle of the day). In addition,

we measured the levels of transcripts encoding enzymes involved in the C_4 cycle with

differential expression in MC (*SvNADP-MDH* and *SvPEPC*) and BSC (*SvNADP-ME* and

SvRbcS, which codes for the small subunit of Rubisco) from *S. viridis* and maize leaves

(Chang *et al.*, 2012; John *et al.*, 2014; Li *et al.*, 2010). We also analysed the transcripts

encoding the sucrose transporters *SvSWEET13a* and *SvSWEET13b*, which have been detected

in BSC from *S. viridis* leaves (Chen *et al.*, 2022) and are putatively regulated by Tre6P

(Oszvald *et al.*, 2018; Fichtner *et al.*, 2021). These transcripts were detected in all samples,

with the highest expression in fully developed leaves and seedlings (Supplementary Fig. S4).

SvTPS1 was mainly detected in whole leaves and the internode and, to a lesser extent,

in seedlings, the flag leaf and the inflorescence. Class II *TPS* transcripts were differentially

expressed among the analysed tissues; most of these transcripts were highly expressed in

whole leaves and seedlings but only marginally in the internode and the inflorescence. The

SvTPS3 transcript showed the highest level in seedlings and whole leaves; the levels of

SvTPS6 and *SvTPS9* were similar to those observed for *SvTPS1* in whole leaves; the *SvTPS7*

transcript displayed the highest level in the flag leaf; whereas the levels of *SvTPS7* and *SvTPS6* were comparable but lower than for *SvTPS1* in the internode (Supplementary Fig. S4).

SvTPP* transcripts also showed a differential spatial distribution, being higher in whole leaves and internode samples. Among clade A *TPP* transcripts, *SvTPP1* showed the highest levels in whole leaves, the inflorescence and the internode, with a maximum in the latter, while *SvTPP8* displayed the highest levels in seedlings and the flag leaf. Regarding clade B *TPP* transcripts, *SvTPP2* was consistently the one with the highest level in all the samples, showing a maximum in the fourth internode. The highest level of the *SvTPP10* transcript (the orthologue of *RAMOSA3*) was observed in the fourth internode. *SvTRE* was detected in all the samples and its abundance was maximal in the internode (Supplementary Fig. S4).

To determine the compartmentation of these transcripts in mature leaves, we applied fractionation protocols (John *et al.*, 2014; Calace *et al.*, 2021) to enrich samples in BSC and MC, followed by RNA extraction and analysis by RT-qPCR (Fig. 3). The marker transcripts *SvNADP-MDH* and *SvPEPC* were highly enriched in the MC fraction, while *SvNADP-ME* and *SvRbcS* were mainly detected in the BSC fraction (Fig. 3A). The *SvTPS1* transcript was almost exclusively found in the BSC fraction (Fig. 3B). It should be noted that BSC strands, while enriched in BSC, also contain phloem (companion cells and sieve elements) and xylem tissue. In *A. thaliana* leaves*,* the AtTPS1 protein is localized primarily in phloem parenchyma (bundle sheath) cells and in the companion cell-sieve element complex of the phloem (Fichtner *et al*., 2020). Thus, some of the *SvTPS1* transcript we attribute to the BSC might be in the phloem complex. The *SvTPS3* and *SvTPS10* transcripts were also preferentially located in the BSC fraction, while *SvTPS6* and *SvTPS8* also showed higher levels in BSC than MC, although with a lower enrichment (Fig. 3B). Conversely, *SvTPS4* and *SvTPS7* transcripts were more abundant in the MC fraction; indeed, *SvTPS7* was not even detected in BSC (Fig.

3B). Among the *TPP* transcripts, *SvTPP1* displayed similar levels in both fractions; *SvTPP2*,

SvTPP3, *SvTPP7* and *SvTPP9* showed a significant enrichment in the BSC fraction; and

SvTPP4 was only detected in the BSC fraction (Fig. 3C). The *SvTRE* transcript was equally

distributed in both cell types, whereas the *SvSWEET13a* and *SvSWEET13b* transcripts were

highly enriched in BSC (Fig. 3D). It is important to note that Fig. 3 does not include

transcripts that were neither detected in BSC nor MC fractions.

Tissue- and cell-specific distribution of the SvTPS1 protein

We recently performed a proteomic analysis with samples enriched in MC and BSC from *S.*

viridis leaves (Calace *et al.*, 2021). In our dataset (Supplementary Table S1 in Calace *et al.*,

2021), which contains 1334 proteins from MC and 1268 proteins from BSC, we observed that

SvTPS1 was detected in two out of four biological replicates from the BSC fraction and was

absent from the MC fraction (Fig. 4A). To further investigate the distribution of the SvTPS1

protein in different cells and tissues, we raised an antiserum against recombinant SvTPS1. To

test the specificity of the antiserum, we extracted proteins from MC and BSC enriched

samples using TCA and Laemmli buffers. The antiserum recognized two protein bands in the

BSC fraction of approximately 112 and 118 kDa (Fig. 4B-C), in agreement with the predicted

molecular mass of SvTPS1 (108.4 kDa). Extraction with TCA is expected to rapidly denature

any proteases in the samples (Rojas *et al.*, 2021), indicating that the double band does not

result from proteolytic cleavage of the SvTPS1 protein during the extraction procedure. We

then analysed the distribution of SvTPS1 in different tissues. The protein was detected in all

the analysed samples, with maximal signals in whole leaves and the internode (Fig. 4D),

which resembles the distribution observed for the *SvTPS1* transcript (Supplementary Fig. S4).

Intercellular distribution of Tre6P

To determine the intercellular distribution of Tre6P, we prepared fractions enriched in MC or

BSC from *S. viridis* leaves by sequential filtering of frozen leaf tissue powder through a

series of nylon meshes with different pore sizes according to Stitt and Heldt (1985). We

performed two independent experiments (Experiments 1 and 2, Supplementary Tables S3 and

S4, respectively), each one consisting of four biological replicates. Supplementary Fig. S5

shows the results obtained for marker enzymes from Experiment 1. The recovery of marker

enzymes in Experiments 1 and 2 varied between 57 and 87%, depending on the enzyme and

the experiment (Supplementary Table S6). The relative distribution of these enzymatic

activities indicated that the fractionation method was successfully adapted to *S. viridis* leaves,

as the results resemble those obtained for maize leaves (Stitt and Heldt, 1985; Arrivault *et al.*,

2017; Denton *et al.*, 2017). We then measured ~30 metabolites (including Tre6P) by LC-

MS/MS, whereas pyruvate, PEP, 3PGA, DHAP, soluble sugars and starch were determined

by enzymatic assays. With these data, we calculated the proportion of each metabolite in BSC

and MC (Supplementary Tables S3-S5).

For Experiment 2, we estimated that ~83% of Tre6P was in the BSC fraction (Fig. 5

and Supplementary Table S4). As already noted, some of this Tre6P might be in the phloem

complex itself, rather than BSC. Other metabolites that were predominantly in the BSC

fraction included pyruvate, 3PGA, sucrose and starch; conversely, DHAP was mainly found

in the MC fraction, while PEP was almost equally distributed between BSC and MC (Fig. 5

and Supplementary Table S4). It should be noted that we estimated similar proportions for

Tre6P, DHAP and starch in Experiment 1 (Supplementary Tables S3 and S5).

The levels of Tre6P and sucrose are tightly correlated across many treatments and

conditions in *A. thaliana* leaves (Yadav *et al.*, 2014). To test whether soluble sugars were

correlated with Tre6P in *S. viridis* leaves, we measured these metabolites in whole leaves

harvested every four hours during the light period, from Zeitgeber time (ZT) 0 to 16 h

(Supplementary Fig. S6 and Supplementary Table S7). At the whole leaf level, Tre6P was not

significantly correlated with glucose, fructose or sucrose (Fig. 6). As previously mentioned,

the leaves of C_4 plants have an additional complexity compared to those from C_3 plants, due

to the arrangement of MC and BSC in the characteristic Kranz anatomy. Thus, we performed

the same analysis using data from the second fractionation experiment (Supplementary Fig.

S6 and Supplementary Table S4). As shown in Fig. 6, across cell-type-enriched fractions,

Tre6P was significantly correlated with sucrose and fructose and, to a lesser extent, with

glucose.

Discussion

Tre6P metabolism is critical for plant growth and development (Fichtner and Lunn, 2021)

and recent reports indicate that it might be an important metabolic target to engineer crops

with improved agronomic traits (Kretzschmar *et al.*, 2015; Nuccio *et al.*, 2015; Griffiths *et*

al., 2016; Gabriel *et al.*, 2021). However, most of our knowledge on this signal molecule is

based on studies performed with the model plant *A. thaliana*, a eudicot that performs C_3

photosynthesis (Rojas *et al.*, 2023). Here, we studied the compartmentation of Tre6P

metabolism in *S. viridis*, a model system for studying C_4 photosynthesis and associated traits

(Brutnell *et al.*, 2010; Doust *et al.*, 2019).

To gain an overview of Tre6P and trehalose metabolism in *S. viridis*, we searched the

genome sequence to identify all recognisable *TPS*, *TPP* and *TRE* genes, and then performed a

phylogenetic analysis of the corresponding protein sequences. The number of genes found in

this species (10 *TPS*, 10 *TPP* and 1 *TRE*) is the same as in its domesticated relative *S. italica*

(Henry *et al.*, 2020). The topologies of the reconstructed phylogenetic trees in this work (Figs

1-2) were in good agreement with those published by Henry *et al.* (2014) for Tre6P-related

enzymes from maize, reinforcing the suitability of *S. viridis* species as a model system for

photosynthesis research in crops. The *S. viridis* genome sequence has only one annotated

gene model for *TPS1* (*Sevir.3G180900.1*), whereas *S. italica* has two gene models for *TPS1*

(*Seita.3G176900.1* and *Seita.3G176900.2*), in which the alternative combination of the first

two exons could produce proteins with different N-terminus, with predicted molecular masses

of 108.4 and 94.9 kDa, respectively (Supplementary Fig. S7). The *Seita.3G176900.1* and

Seita.3G176900.2 transcripts displayed a differential response to light (Henry *et al.*, 2020),

suggesting their encoded polypeptides could have particular biological functions. A detailed

comparison of the *Sevir.3G180900* and *Seita.3G176900* DNA sequences showed eight

differences, but only from the fifth intron towards the 3´-end (Supplementary Fig. S7).

Therefore, an alternative transcript could also be produced from the *Sevir.3G180900* gene.

The different transcriptional variants of these closely related species (wild *vs* domesticated) is

of interest for further research.

The expression of transcripts encoding enzymes involved in Tre6P metabolism in *S.*

viridis showed a differential pattern among the tissues analysed in this work. Analysis of

Tre6P-related transcripts in different tissues of *S. viridis* was not previously reported,

although other works showed their response in leaves subjected to different treatments,

including light (Henry *et al.*, 2014, 2020), salt (Henry *et al.*, 2015), high light and

temperature (Anderson *et al.*, 2021). The *SvTPS1* transcript and SvTPS1 protein abundance

was highest in internode samples (Supplementary Fig. S4 and Fig. 4D), which are rich in

vascular tissue. This is in good agreement with data from *A. thaliana*, where the TPS1 protein

was mainly detected in the vasculature of leaves, roots and reproductive tissues (Fichtner *et*

al., 2020). We could not find Tre6P-related transcripts in the RNAseq analysis performed by

Martin *et al.* (2016) using developing internodes from *S. viridis*, probably due to differences

in the sampling procedures and analysis between both works. Other authors have reported

that *A. thaliana TPS1* transcripts are relatively abundant in the protovasculature of

developing leaves and the shoot apical meristem (Wahl *et al.*, 2013), and in differentiated

phloem cells from the inflorescence stem (Shi *et al.*, 2021).

The transcripts encoding class II TPS proteins in *S. viridis* showed different levels according to the analysed tissue, which resembles data reported for class II TPS transcripts from *A. thaliana* (Ramon *et al.*, 2009). As previously mentioned, our analyses were performed with samples harvested in the middle of the light period, and thus did not capture the diurnal fluctuations previously observed in *A. thaliana* (Usadel *et al.*, 2008) and maize (Henry *et al.*, 2014; Borba *et al.*, 2023). The transcript *SvTPS3* displayed high levels in seedlings and whole leaves (Supplementary Fig. S4). The protein encoded by this transcript was located in a sub-clade exclusively containing class II TPS proteins from monocots, well separated from all other class II TPS sequences from eudicots (Fig. 1). This particular branch contains sequences from PACMAD (*S. viridis* and maize) and BOP/BEP (rice and *B. distachyon*) species, as well as from monocots outside the Poaceae that were included in our analysis (*A. comosus* and *M. acuminata*). This suggests that the monocot-specific class II TPS sub-clade arose before the divergence of the orders Poales (including the Poaceae and Bromeliaceae) and Zingeberales (including the Musaceae). The *SvTPS6* transcript was mainly detected in whole leaves and the fourth internode, while abundance of the *SvTPS7* transcript was highest in the flag leaf and the fourth internode (Supplementary Fig. S4). The proteins encoded by these transcripts were located in the same sub-clade as *A. thaliana* TPS7 (Fig. 1), whose expression was mainly detected in leaf primordia, cotyledons, roots (concentrated in the vasculature of the mature root and evenly distributed in the meristematic zone) and flowers (in ovules and pollen; Ramon *et al.*, 2009). Recently, Van Leene *et al.* (2022) showed that all class II TPS proteins from *A. thaliana*, including TPS7, interact with and potentially act as regulators of the SUCROSE-NON-FERMENTING1-RELATED

PROTEIN KINASE1 (SnRK1). Whether these proteins play the same role in *S. viridis* and

other C$_4$ grasses has to be further investigated.

Transcripts encoding TPP proteins from *S. viridis* also showed a particular distribution

pattern, according to the tissue and cell type under analysis, as previously reported for *A.*

thaliana (Vandesteene *et al.*, 2012). The *SvTPP2* transcript was mainly detected in the fourth

internode, with lower but significant levels in seedlings, whole leaves and the inflorescence

(Supplementary Fig. S4). The protein encoded by this transcript is the orthologue of ZmTPP4

and was located in the monocot-specific sub-clade that contains RAMOSA3 (ZmTPP10; Fig.

2). The levels of *SvTPP1* were higher in whole leaves, the inflorescence, and the fourth

internode, compared to other tissues, while levels of *SvTPP8* were higher in seedlings, the

flag leaf and the fourth internode (Supplementary Fig. S4). The SvTPP1 and SvTPP8 protein

sequences were located in clade A, which contains *A. thaliana* TPPA, TPPF and TPPG (Fig.

2). There is evidence of recent duplication of genes encoding *A. thaliana* clade A TPPs,

associated with whole-genome duplications within the Brassicaceae, with a high degree of

paralogue retention and evidence of neofunctionalization (Vandesteene *et al.*, 2012). Thus, it

seems feasible that the duplication and diversification of clade A *S. viridis* TPPs observed in

Fig. 2 occurred independently of those in *A. thaliana* (Kerbler *et al.*, 2023).

We found that several Tre6P-related transcripts were preferentially located in BSC of

S. viridis leaves (Fig. 3). To compare our results with previously published datasets, we

retrieved RNAseq data from studies performed with enriched MC and BSC fractions from *S.*

viridis and maize (John *et al.*, 2014; Denton *et al.*, 2017; Supplementary Fig. S8 and

Supplementary Table S8). Our analysis showed that the *TPS1* transcript was enriched in BSC

from *S. viridis* and maize leaves (Fig. 3B and Supplementary Fig. S8), as previously shown

for the *A. thaliana TPS1* transcript (Aubry *et al.*, 2014). It is important to recall that our

experimental data originate from samples harvested in the middle of the day (ZT 8 h), as *TPS*

and *TPP* transcripts fluctuate during the diel cycle (Usadel *et al.*, 2008, Borba *et al.*, 2023);

particularly, the maize *TPS1* transcript displayed the highest level at ZT 8 h (Borba *et al.*,

2023), indicating that this time of day is optimal for analysing the cellular distribution of

TPS1 transcripts. Regarding the class II *TPS* transcripts, the levels of *SvTPS3* and *SvTPS10*

were significantly higher in BSC than in MC when analysed by RT-qPCR (Fig. 3B);

however, these patterns were not the same in RNAseq data from *S. viridis* and maize. As

shown in Supplementary Fig. S8, *SvTPS9* and *SvTPS10* displayed the highest enrichment in

S. viridis BSC, whereas *ZmTPS11* (which clustered together with *SvTPS7*; Fig. 1) showed

high enrichment in BSC from almost all leaf segments of maize leaves. Abundance of

SvTPP1 (the orthologue of *ZmTPP6*) was similar in MC and BSC (Fig. 3C), which agrees

with RNAseq data from *S. viridis* and maize (Supplementary Fig. S8). Our RT-qPCR results

showed greater abundance of *SvTPP2, SvTPP3, SvTPP7* and *SvTPP9* in BSC than in MC

(Fig. 3C), and the same trend was observed in the RNAseq data from *S. viridis*

(Supplementary Fig. S8). Similarly, *ZmTPP1*, *ZmTPP2*, *ZmTPP7* and *ZmTPP8* (the

orthologues of *SvTPP7*, *SvTPP3*, *SvTPP2* and *SvTPP3*, respectively; Fig. 2) were enriched in

BSC across all fractions of the maize leaf, while *ZmTPP4* and *ZmTPP9* (the orthologues of

SvTPP2 and *SvTPP5*, respectively; Fig. 2) were enriched in BSC from the middle and the tip

of the maize leaf (fractions 1 to 3, Supplementary Fig. S8). The levels of the *SvTRE* transcript

were similar in MC and BSC, as reported in RNAseq studies of *S. viridis* and maize leaves

(Fig. 3D and Supplementary Fig. S8).

Cell-specific expression of transcripts in MC and BSC is regulated by transcriptional

and post-transcriptional mechanisms (Reeves *et al.*, 2017). Furthermore, due to translational

and post-translational regulation, transcript abundance is not necessarily a reliable indicator

of protein abundance (Gibon *et al.*, 2004; Wang *et al.*, 2011; Reeves *et al.*, 2017). Thus, we

investigated the distribution of the SvTPS1 polypeptide in different tissues and cell types.

Immunodetection assays detected the SvTPS1 protein in whole leaves and the fourth

internode (Fig. 4D), as previously observed in the transcript abundance analysis

(Supplementary Fig. S4). The localization of SvTPS1 in the BSC fraction (Fig. 4) is

consistent with the proteomic data from Calace *et al*. (2021). The antiserum raised against

SvTPS1 consistently recognized two immunoreactive protein bands, particularly in BSC (Fig.

4B-C). Based on our results, we hypothesized that these proteoforms of SvTPS1 may result

from two splicing variants or a post-translational modification. The current gene model for

SvTPS1 does not predict the existence of a splicing variant; however, *S. italica* has two

splicing variants for *SiTPS1* (Henry *et al.*, 2020). As previously mentioned, the sequences

encoding *SvTPS1* and *SiTPS1* are identical up to the fifth intron (Supplementary Fig. S7),

which opens the possibility of an alternative splicing variant for *SvTPS1*. In that case, the

alternative polypeptide should have a 135-residues shorter N-terminus (i.e., a difference of

13.5 kDa). Regarding the existence of a putative post-translational modification, it is

important to note that *A. thaliana* TPS1 is phosphorylated and potentially sumoylated

(Fichtner *et al.*, 2020; Fichtner and Lunn, 2021). The phosphorylation sites found on the *A.*

thaliana TPS1 are not conserved in SvTPS1 but the putative sumoylation site (Lys^{902} in *A.*

thaliana TPS1, Lys^{919} in SvTPS1) is well-conserved (Supplementary Fig. S2; Fichtner *et al.*,

2020; Fichtner and Lunn, 2021). Considering the theoretical molecular mass of SvTPS1

(108.4 kDa) and the estimated sizes of the immunoreactive bands (112 and 118 kDa, Fig. 4B-

C), the occurrence of a post-translational modification seems more plausible, as the two

putative splicing variants of SvTPS1 should have 108.4 and 94.9 kDa, respectively

(Supplementary Fig. S7). Overall, our results show that the only class I TPS from *S. viridis* is

predominantly located in BSC, suggesting this is the major location for Tre6P synthesis in *S.*

viridis leaves.

We adapted a method, originally developed for maize, that allows metabolite analysis

in MC and BSC enriched fractions (Stitt and Heldt, 1985). We found that the proportion of

Tre6P was higher in BSC than in MC. Similar results were observed for pyruvate, 3PGA and

starch, as expected for an NADP-ME C_4 species (Fig. 5 and Supplementary Tables S3-S5).

Similarly, ribulose 1,5-bisphosphate and other intermediates of the Calvin-Benson cycle were

almost exclusively located in BSC (Supplementary Table S5). The amount of malate was also

higher in BSC than MC. This result is counterintuitive, as the concentration gradient of

malate is expected to be the opposite; i.e., a higher concentration in MC than BSC, to drive

diffusion of malate from the MC to the BSC. However, it is important to consider that maize

leaves have a large pool of malate (up to 60% of the total malate) that is not actively involved

in the C_4 cycle, including a potentially large pool in the vacuole (Arrivault *et al.*, 2017). Thus,

there still might be a concentration gradient which could move malate from the cytosol of

MC to cytosol of BSC in *S. viridis* leaves, although this has to be proven, for instance, by a

combination of $^{13}CO_2$ labelling and leaf fractionation, as previously done with maize leaves

(Arrivault *et al.*, 2017). Two intermediates of the Calvin-Benson cycle showed contrasting

distributions: 3PGA was predominantly located in the BSC, while DHAP was mainly in the

MC (Fig. 5). This is similar to the intercellular distributions of 3PGA and DHAP reported for

maize leaves (Leegood, 1985; Stitt and Heldt, 1985; Lunn and Furbank, 1997; Arrivault *et*

al., 2017), where there is intercellular shuttling of these metabolites. Maize, in common with

other NADP-ME-type C_4 species, has mostly agranal chloroplasts in the BSCs that lack

photosystem II, restricting the photosynthetic electron transport chain to cyclic electron flow

and production of ATP, but little or no NADPH. It has been proposed that a low

$NADPH:NADP^+$ ratio in the BSC chloroplasts is needed to maintain flux through the NADP-

ME enzyme in the direction of malate decarboxylation (Wang *et al.*, 2014). This reaction

provides about half of the NADPH needed for reduction of 1,3-bis-phosphogylcerate to

glyceraldehyde 3-phosphate in the Calvin-Benson cycle. To supplement this, 3PGA is

exported from the BSC chloroplasts and moves to the MCs, where it enters the chloroplasts

for conversion to triose-phosphates, mainly DHAP, which then moves back to the BSCs and

re-enters the Calvin-Benson cycle in the BSC chloroplasts. High concentrations of 3PGA and

DHAP in the BSC and MC, respectively, drive diffusion of these metabolites in opposite

directions between the two cell types. The asymmetric distribution of 3PGA and DHAP in *S.*

viridis suggests that intercellular shuttling of 3PGA and DHAP is also a feature of C_4

photosynthesis in this NADP-ME-type species.

Pyruvate was also predominantly located in the BSC, consistent with its production

from malate decarboxylation, via NADP-ME, in the BSCs, and diffusion into the MCs for

conversion to PEP via PPDK. PEP was evenly distributed between MC and BSC (Fig. 5 and

Supplementary Table S5). These distributions differ from those reported for maize, where

PEP showed a preferential increase in the BSC and pyruvate was located mainly in the MC

(Arrivault *et al.*, 2017). Such differences could be explained by i) the existence of metabolic

gradients along the leaf, or ii) the presence of metabolite pools that are not actively involved

in C_4 photosynthesis, for example, if located in the vacuole, which could mask metabolite

gradients between the cytosol of MC and that of BSC (Szecowka *et al.*, 2013; Arrivault *et al.*,

2017; Denton *et al.*, 2017). Concerning the first hypothesis, proteomic, transcriptomic and

metabolomic studies along maize leaf sections have shown a gradient that goes from a

heterotrophic (in the base) to a photosynthetic (in the tip) metabolism (Majeran *et al.*, 2010;

Pick *et al.*, 2011). In fact, Denton *et al.* (2017) found significant metabolic differences across

the longitudinal gradient of maize leaves, while changes were rather small between MC and

BSC. Regarding the second hypothesis, labelling studies showed that, for some metabolites,

pools that are not involved in photosynthesis form a considerable part of the total pool both in

C_3 and C_4 plants (Szecowka *et al.*, 2013; Arrivault *et al.*, 2017). Particularly, the relative

abundance of the unlabelled pyruvate isotopomer after 60 min of $^{13}CO_2$ labelling in maize

leaves was 17% (Arrivault *et al.*, 2017), which points to the existence of a considerable

inactive pool of pyruvate.

A limitation of our study is that samples for metabolite analysis were collected from

the middle section of the leaf blade at one defined time-point in the middle of the light period,

so we could not capture metabolic dynamics. It would be interesting to perform labelling

studies followed by cellular enrichment to obtain a detailed scenario of the metabolic

fluctuations that occur in *S. viridis* leaves. Furthermore, overall gradients between the MC

and BSC might not reflect the gradients between the cytosol of these two cell types, if there

were active transport of metabolites across the plastid envelope or the pH gradient across the

envelope membrane influences their distribution between the chloroplast stroma and the

cytosol, as is known to be the case for 3PGA (Fliege *et al.*, 1978; Stitt *et al.*, 1980). The

relation between the overall intercellular metabolite gradient and the gradient between the

cytosols of the BSC and MC will also be affected by the relative contribution of chloroplasts

and cytosol to cytoplasmic volume. Overall, our study indicates that the metabolite gradients

reported in maize may not be generalizable and underlines the need for studies of metabolite

gradients in other C_4 species. Just as different biochemical solutions have arisen in different

species to concentrate CO_2 into the BSC (Furbank and Kelly, 2021), different strategies may

also have been followed to generate the intercellular gradients in the cytosol that are needed

to drive metabolite movement between the BSC and MC. Overall, our fractionation data

provide the first insights into the distributions of key photosynthetic intermediates in *S.*

viridis. These are consistent with the photosynthetic fluxes and intercellular metabolite

movements that we would expect to see in an NADP-ME-type C_4 species, demonstrating the

reliability of this experimental approach.

The C_4 cycle involves metabolic fluxes within and between MC and BSC, whereas

other metabolic pathways can be restricted to one particular cell type. For example, sucrose

synthesis occurs predominantly in MC, while starch metabolism takes place in BSC

(Leegood, 1985; Stitt and Heldt, 1985; Lunn and Furbank, 1997; Furbank and Kelly, 2021).

To better understand the compartmentation of metabolic pathways in *S. viridis* leaves, we

combined transcriptomic and proteomic data previously published by John *et al.* (2014) and

Calace *et al.* (2021), respectively (Supplementary Table S9). Supplementary Fig. S9 shows

that transcripts and proteins related to sucrose and starch synthesis are preferentially located

in MC and BSC, respectively, whereas the components of the C_4 cycle are distributed

between MC and BSC, as expected for an NADP-ME subtype species.

The main aim of the cell fractionation experiments in our study was to determine the

cellular localization of Tre6P. Our transcript and protein analyses showed that the SvTPS1

enzyme is predominantly located in the BSC and associated vascular tissues, and essentially

absent from MC (Figs 3-4), indicating that Tre6P is synthesized in BSC but not in MC. We

found that approximately 83% of Tre6P was present in the BSC fraction and, considering the

previously mentioned limitations, this value could be even higher. It is worth noting that

some of the Tre6P in the BSC fraction may be in the phloem complex because, although the

cell separation method allows the sample to be enriched with BSC strands, this enrichment is

partial and may bring with it remains from other cells types. This would make sense

considering that, in *A. thaliana*, Tre6P metabolism has been reported to take place in both

phloem parenchyma cell (equivalent to the BSC) and in the companion cell-sieve element

complex of the phloem (Fichtner *et al.*, 2020). As previously mentioned, sucrose synthesis

seems to occur in MC of *S. viridis* leaves (Supplementary Fig. S9); however, the amount of

sucrose was higher in the BSC fraction than in the MC fraction (Supplementary Fig. S6).

This result is puzzling, as sucrose levels are expected to be higher in the MC than the BSC, to

drive its diffusion to the BSC. We thus speculate that i) sucrose is sequestered either in some

compartment other than the cytosol in the BSC or other cell types that are enriched in the

BSC fractions; or ii) that a large part of the sucrose in the leaf is in the phloem.

Lunn *et al.* (2006) showed there is a significant positive correlation between Tre6P

and sucrose levels in samples from *A. thaliana* seedlings and rosettes. The correlation

between Tre6P and sucrose persisted even in transgenic lines expressing the *E. coli* otsA and

otsB proteins under the control of the *35S* promoter, although sucrose levels were markedly

lower or higher, respectively, which led to the development of the Tre6P-sucrose nexus

model (Yadav *et al.*, 2014). Tre6P and sucrose levels fluctuated during the light period in *S.*

viridis leaves (Supplementary Fig. S6) but, unlike *A. thaliana*, we did not observe a

significant correlation at the whole-leaf level (Fig. 6). Conversely, we found a significant

correlation between Tre6P and sucrose across leaf fractions containing different proportions

of MC and BSC (Fig. 6). The Tre6P-sucrose nexus model postulates that Tre6P is a signal

and a negative regulator of sucrose levels (Yadav *et al.*, 2014). According to this model, if

sucrose levels rise too high, Tre6P rises in parallel and triggers mechanisms to lower the level

of sucrose, either by reducing sucrose synthesis or increasing its degradation or transport.

Regarding the first alternative, research performed with ethanol-inducible *A. thaliana*

transgenic lines showed that artificially increased Tre6P levels decreased the amount of

sucrose by diverting newly fixed carbon to organic and amino acids during the day, through

the post-translational activation of PEPC and nitrate reductase (Figueroa *et al.*, 2016). As

previously mentioned, the initial CO_2 fixation step in C_4 plants is catalysed by PEPC, which

is located in MC. Our results strongly suggest that Tre6P is mainly produced and

accumulated in BSC of *S. viridis* leaves. Thus, if the Tre6P-PEPC connection is operational

in C_4 species, this would imply that Tre6P has to diffuse from BSC to MC; indeed, a small

amount of Tre6P was actually found in MC (Fig. 5). The asymmetrical distribution of Tre6P

suggests that it could be sensing sucrose levels in BSC and then act as a signal that moves

into the MC to potentially modulate PEPC activity and/or sucrose synthesis, which is also

located in MC (Supplementary Fig. S9).

Another possibility is that Tre6P regulate sucrose levels by controlling phloem

loading. TPS1 has been found in the vasculature of *A. thaliana* leaves; particularly, in the

phloem parenchyma and the companion cell/sieve element complex (Fichtner *et al.*, 2020).

Tre6P has been shown to modulate the levels of *ZmSWEET13a,c* and *AtSWEET11-14*

transcripts in maize and *A. thaliana*, respectively, which encode sucrose transporters involved

in phloem loading (Oszvald *et al.*, 2018; Fichtner *et al.*, 2021). Single-cell transcriptomics in

A. thaliana, rice and maize showed that *SWEET* transcripts were preferentially detected in

BSC (Bezrutczyk *et al.*, 2021; Shi *et al.,* 2021; Hua *et al.,* 2021; Tao *et al.*, 2022). Moreover,

the *ZmSWEET13a-c* transcripts and the SWEET13a protein were specifically detected in the

abaxial BSC of major and intermediate veins from maize leaves (Bezrutczyk *et al.*, 2021),

while both SWEET13a and SWEET13b proteins were detected in the abaxial BSC of

intermediate veins from *S. viridis* leaves (Chen *et al.*, 2022). Based on these data, it has been

proposed that sucrose moves from adaxial to abaxial cells via plasmodesmata and is then

exported by SWEET13 to the apoplast, from where it is taken up by transporters from the

SUT/SUC (sucrose-H$^+$ symporter) family located in the companion cells (Bezrutczyk *et al.*,

2021; Chen *et al.*, 2022). Transcripts encoding Tre6P-related enzymes (specifically,

ZmTPS10, *ZmTPS12*, *ZmTPP1* and *ZmTPP2*; Supplementary Table S8) have been detected in

abaxial BSC from maize leaves (Bezrutczyk *et al.*, 2021), suggesting Tre6P metabolism

could also be specifically located in these cells. The maize *zmsweet13a-c* triple mutant is

impaired in sugar phloem loading, thus producing drastic effects on growth (Bezrutczyk *et*

al., 2018). In *S. viridis*, such mutants have not yet been identified, but there is compelling

evidence that it could be an apoplastic phloem-loading species, and thus SvSWEET13a-b

might function in phloem loading (Chen *et al.*, 2022). Based on this evidence, and the fact

that Tre6P metabolism is mostly located in BSC, we speculate that Tre6P modulates sucrose

export in leaves of C_4 plants by controlling the expression, and thereby maximal activity, of

the SWEET13 transporters, although this hypothesis remains to be tested. If confirmed, this

would implicate Tre6P in the reconfiguring of phloem loading that is a common feature of C_4

monocots, and potentially an essential element in the evolution of Kranz anatomy and C_4

photosynthesis (Furbank and Kelly, 2021; Rojas *et al*., 2023). To conclude, our results

indicate that Tre6P is synthesized and mainly located in the BSC of *S. viridis*, which play a

central role in phloem loading in C_4 species and represent a strategic site at the interface

between source and sink tissues. This localization allows Tre6P to monitor the levels of

sucrose in BSC and, via diffusion into the MC, potentially regulate CO_2 fixation and sucrose

biosynthesis as well as sucrose export from the leaves (Rojas *et al*., 2023). Our results point

to Tre6P signalling being adapted to the specialized metabolism and anatomy of C_4 plants,

and raise the possibility that such adaptations were a prerequisite for the evolution of C_4

photosynthesis from C_3 ancestors.

Supplementary Data

Supplementary Figure S1. C_t values for the reference transcript *SvKIN* in different tissues

and cell types from *S. viridis*.

Supplementary Figure S2. Multiple sequence alignment of *S. viridis* TPS proteins.

Supplementary Figure S3. Multiple sequence alignment of *S. viridis* TPP proteins.

Supplementary Figure S4. Relative abundance of transcripts encoding Tre6P-related

enzymes in different *S. viridis* tissues.

Supplementary Figure S5. Activities of marker enzymes in different fractions from *S.*

viridis leaves.

Supplementary Figure S6. Analysis of metabolites from *S. viridis* leaves.

Supplementary Figure S7. Gene models for the class I TPS from *S. viridis* and *S. italica*.

Supplementary Figure S8. Analysis of transcripts encoding Tre6P-related enzymes in BSC

and MC from *S. viridis* and maize leaves.

Supplementary Figure S9. Analysis of selected metabolic pathways located in MC and BSC

of *S. viridis* leaves.

Supplementary Table S1. TPS and TPP protein sequences used for reconstruction of

phylogenetic trees.

Supplementary Table S2. Sequences of the primers used to measure transcript levels by

qPCR.

Supplementary Table S3. Metabolite and enzyme data and calculations (Experiment 1).

Supplementary Table S4. Metabolite and enzyme data and calculations (Experiment 2).

Supplementary Table S5. Calculation of metabolite fractions in MC and BSC.

Supplementary Table S6. Recovery of marker enzymes in the fractionation experiments.

Supplementary Table S7. Metabolite data from *S. viridis* leaves during a day time-course.

Supplementary Table S8. Relative transcript abundance in MC and BSC from *S. viridis* and

maize.

Supplementary Table S9. Analysis of selected metabolic pathways using transcriptomic and

proteomic data from *S. viridis* leaves.

Acknowledgements

TT, BER and PC are fellows from CONICET. LEL, MSaigo and CMF are researchers from

CONICET. CMF thanks NME, CF and MF for their joy and support.

Author contributions

Conceptualization: MStitt, JEL and CMF; Investigation: TT, BER, LEL, PC, RF, SA and

CMF; Formal analysis: all authors; Writing - Original Draft: TT, BER and CMF; Writing -

Review & Editing: all authors.

Conflict of interest

No conflict of interest declared.

Funding

This work was supported by Agencia Nacional de Promoción de la Investigación, el

Desarrollo Tecnológico y la Innovación [PICT-2018-00865 and PICT-2020-00260];

Universidad Nacional del Litoral [CAI+D 2020]; Consejo Nacional de Investigaciones

Científicas y Técnicas [PIP 2021-2023]; Agencia Santafesina de Ciencia, Tecnología e

Innovación [PEIC I+D 2021-021]; and the Max Planck Society [Partner Group for Plant

Biochemistry].

Data availability

All data supporting the findings of this study are available within the paper and within its

supplementary data published online.

References

Anderson CM, Mattoon EM, Zhang N, *et al. 2021. High light and temperature reduce photosynthetic efficiency through different mechanisms in the C$_4$ model *Setaria viridis*. Communications Biology **4**, 1092.

Arrivault S, Guenther M, Ivakov A, Feil R, Vosloh D, van Dongen JT, Sulpice R, Stitt M. 2009. Use of reverse-phase liquid chromatography, linked to tandem mass spectrometry, to profile the Calvin cycle and other metabolic intermediates in Arabidopsis rosettes at different carbon dioxide concentrations. Plant Journal **59**, 826–839.

Arrivault S, Obata T, Szecówka M, Mengin V, Guenther M, Hoehne, M, Fernie AR, Stitt M. 2017. Metabolite pools and carbon flow during C$_4$ photosynthesis in maize: $^{13}CO_2$ labeling kinetics and cell type fractionation. Journal of Experimental Botany **68**, 283–298.

Ashton AR, Burnell JN, Furbank RT, Jenkins CLD, Hatch MD. 1990. Enzymes of C$_4$ photosynthesis. In: Lea PJ, ed. Methods in plant biochemistry, Volume 3, Enzymes of primary metabolism. London, 39-72.

Aubry S, Smith-Unna RD, Boursnell CM, Kopriva S, Hibberd JM. 2014. Transcript residency on ribosomes reveals a key role for the *Arabidopsis thaliana* bundle sheath in sulfur and glucosinolate metabolism. Plant Journal **78**, 659–673.

Avidan O, Martins MCM, Feil R, Lohse M, Giorgi FM, Schlereth A, Lunn JE, Stitt M. 2024. Direct and indirect responses of the Arabidopsis transcriptome to an induced increase in trehalose 6-phosphate. Plant Physiology, in press. https://doi.org/10.1093/plphys/kiae196

Avonce N, Mendoza-Vargas A, Morett E, Iturriaga G. 2006. Insights on the evolution of trehalose biosynthesis. BMC Evolutionary Biology **6**, 1-15.

Barraza A, Contreras-Cubas C, Estrada-Navarrete G, Reyes JL, Juárez-Verdayes MA, Avonce N, Quinto C, Díaz-Camino C, Sanchez F. 2016. The class II trehalose 6-phosphate synthase gene *PvTPS9* modulates trehalose metabolism in *Phaseolus vulgaris* nodules.

Frontiers in Plant Science **7**, 1589.

Bezrutczyk M, Hartwig T, Horschman M, Char SN, Yang J, Yang B, Frommer WB, Sosso D. 2018. Impaired phloem loading in *zmsweet13a,b,c* sucrose transporter triple knock-out mutants in *Zea mays*. New Phytologist **218**, 594-603.

Bezrutczyk M, Zöllner NR, Kruse CPS, Hartwig T, Lautwein T, Köhrer K, Frommer WB, Kim JY. 2021. Evidence for phloem loading via the abaxial bundle sheath cells in maize leaves. Plant Cell **33**, 531–547.

Bledsoe SW, Henry C, Griffiths CA, Paul MJ, Feil R, Lunn JE, Stitt M, Lagrimini LM. 2017. The role of Tre6P and SnRK1 in maize early kernel development and events leading to stress-induced kernel abortion. BMC Plant Biology **17**, 1-17.

Borba AR, Reyna-Llorens I, Dickinson PJ, *et al*. 2023. Compartmentation of photosynthesis gene expression in C_4 maize depends on time of day. Plant Physiology **193,** 2306–2320.

Bradford MM. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Analytical Biochemistry **72**, 248–254.

Brutnell TP, Wang L, Swartwood K, Goldschmidt A, Jackson D, Zhu XG, Kellogg E, Van Eck J. 2010. *Setaria viridis*: a model for C_4 photosynthesis. The Plant Cell **22**, 2537–2544.

Cabib E, Leloir LF. 1958. The biosynthesis of trehalose phosphate. Journal of Biological Chemistry **231**, 259–275.

Cacefo V, Ribas AF, Zilliani RR, Neris DM, Domingues DS, Moro AL, Vieira LGE. 2019. Decarboxylation mechanisms of C_4 photosynthesis in Saccharum spp.: Increased PEPCK activity under water-limiting conditions. BMC Plant Biology **19**, 1–14.

Calace P, Tonetti T, Margarit E, Figueroa CM, Lobertti C, Andreo CS, Gerrard

Wheeler MC, Saigo M. 2021. The C_4 cycle and beyond: diverse metabolic adaptations accompany dual-cell photosynthetic functions in Setaria. Journal of Experimental Botany **72**, 7876–7890.

Carillo P, Feil R, Gibon Y, Satoh-Nagasawa N, Jackson D, Bläsing OE, Stitt M, Lunn JE. 2013. A fluorometric assay for trehalose in the picomole range. Plant Methods **9**, 1-15.

Chang YM, Wen-Yu L, Chun-Chieh AS, *et al.* 2012. Characterizing regulatory and functional differentiation between maize mesophyll and bundle sheath cells by transcriptomic analysis. Plant Physiology **160**, 165–177.

Chapman KSR, Hatch MD. 1981. Aspartate decarboxylation in bundle sheath cells of *Zea mays* and its possible contribution to C_3 photosynthesis. Functional Plant Biology **8**, 237–248.

Chen L, Ganguly DR, Shafik SH, Ermakova M, Pogson BJ, Grof CP, Sharwood RE, Furbank RT. 2022. Elucidating the role of SWEET13 in phloem loading of the C_4 grass *Setaria viridis*. Plant Journal **109**, 615-632.

Claeys H, Vi SL, Xu X, *et al.* 2019. Control of meristem determinacy by trehalose 6-phosphate phosphatases is uncoupled from enzymatic activity. Nature Plants **5**, 352–357.

Czedik-Eysenberg A, Arrivault S, Lohse M, *et al.* 2016. The interplay between carbon availability and growth in different zones of the growing maize leaf. Plant Physiology **172**, 943-967.

Delorge I, Figueroa CM, Feil R, Lunn JE, Van Dijck P. 2015. Trehalose-6-phosphate synthase 1 is not the only active TPS in *Arabidopsis thaliana*. Biochemical Journal **466**, 283–290.

Denton AK, Maß J, Külahoglu C, Lercher MJ, Bräutigam A, Weber APM. 2017. Freeze-quenched maize mesophyll and bundle sheath separation uncovers bias in previous tissue-specific RNA-Seq data. Journal of Experimental Botany **68**, 147–160.

Devi K, Prathima PT, Gomathi R, Manimekalai R, Lakshmi K, Selvi A. 2019. Gene expression profiling in sugarcane genotypes during drought stress and rehydration. Sugar Tech **21**, 717–733.

Dong Z, Xiao Y, Govindarajulu R, Feil R, Siddoway ML, Nielsen T, Lunn JE, Hawkins J, Whipple C, Chuck G. 2019. The regulatory landscape of a core maize domestication module controlling bud dormancy and growth repression. Nature Communications **10**, e3810.

dos Anjos L, Pandey PK, Moraes TA, Feil R, Lunn JE, Stitt M. 2018. Feedback regulation by trehalose 6-phosphate slows down starch mobilization below the rate that would exhaust starch reserves at dawn in Arabidopsis leaves. Plant Direct **2**, e00078.

Doust AN, Brutnell TP, Upadhyaya HD, Van Eck J. 2019. Editorial: setaria as a model genetic system to accelerate yield increases in cereals, forage crops, and bioenergy grasses. Frontiers in Plant Science **10**, 1211.

Eastmond PJ, van Dijken AJ, Spielman M, Kerr A, Tissier AF, Dickinson HG, Jones JDG, Smeekens SC, Graham IA. 2002. Trehalose-6-phosphate synthase 1, which catalyses the first step in trehalose synthesis, is essential for arabidopsis embryo maturation. Plant Journal **29**, 225–235.

Fang L. 2012. Antibody purification from western blotting. Bio-protocol **2**, e133.

Fichtner F, Barbier FF, Annunziata MG, Feil R, Olas JJ, Mueller-Roeber B, Stitt M, Beveridge CA, Lunn JE. 2021. Regulation of shoot branching in Arabidopsis by trehalose 6-phosphate. New Phytologist **229**, 2135–2151.

Fichtner F, Lunn JE. 2021. The role of trehalose-6-phosphate (Tre6P) in plant metabolism and development. Annual Review of Plant Biology **72**, 737–760.

Fichtner F, Olas JJ, Feil R, Watanabe M, Krause U, Hoefgen R, Stitt M, Lunn JE. 2020. Functional features of TREHALOSE-6-PHOSPHATE SYNTHASE1, an essential enzyme in arabidopsis. Plant Cell **32**, 1949–1972.

Figueroa CM, Feil R, Ishihara H, *et al.* 2016. Trehalose 6-phosphate coordinates organic and amino acid metabolism with carbon availability. Plant Journal **85**, 410–423.

Figueroa CM, Lunn JE. 2016. A tale of two sugars: trehalose-6-phosphate and sucrose. Plant Physiology **172**, 7–27.

Fliege R, Flügge UI, Werdan K, Heldt HW. 1978. Specific transport of inorganic phosphate, 3-phosphoglycerate and triosephosphates across the inner membrane of the envelope in spinach chloroplasts. Biochimica et Biophysica Acta **502**, 232-247.

Furbank RT and Kelly S. 2021. Finding the C_4 sweet spot: cellular compartmentation of carbohydrate metabolism in C_4 photosynthesis. Journal of Experimental Botany **72**, 6018-6026.

Gabriel C, Fernhout JJ, Fichtner F, Feil R, Lunn JE, Kossmann J, Lloyd JR, van der Vyver C. 2021. Genetic manipulation of trehalose-6-phosphate synthase results in changes in the soluble sugar profile in transgenic sugarcane stems. Plant Direct **5**, 1–18.

Gascuel O. 1997. BIONJ: An improved version of the NJ algorithm based on a simple model of sequence data. Molecular Biology and Evolution **14**, 685–695.

Gibon Y, Blaesing OE, Hannemann J, Carillo P, Höhne M, Hendriks JHM, Palacios N, Cross J, Selbig J, Stitt M. 2004. A robot-based platform to measure multiple enzyme activities in Arabidopsis using a set of cycling assays: comparison of changes of enzyme activities and transcript levels during diurnal cycles and in prolonged darkness. The Plant Cell **16**, 3304–3325.

Gibson RP, Turkenburg JP, Charnock SJ, Lloyd R, Davies GJ. 2002. Insights into trehalose synthesis provided by the structure of the retaining glucosyltransferase OtsA. Chemistry and Biology **9**, 1337–1346.

Gomez LD, Baud S, Gilday A, Li Y, Graham IA. 2006. Delayed embryo development in the Arabidopsis trehalose-6-phosphate synthase 1 mutant is associated with altered cell wall

structure, decreased cell division and starch accumulation. Plant Journal **46**, 69–84.

Gomez LD, Gilday A, Feil R, Lunn JE, Graham IA. 2010. AtTPS1-mediated trehalose 6-phosphate synthesis is essential for embryogenic and vegetative growth and responsiveness to ABA in germinating seeds and stomatal guard cells. Plant Journal **64**, 1–13.

Gouy M, Guindon S, Gascuel O. 2010. SeaView version 4: a multiplatform graphical user interface for sequence alignment and phylogenetic tree building. Molecular Biology and Evolution **27**, 221-224.

Griffiths CA, Sagar R, Geng Y, *et al.* 2016. Chemical intervention in plant sugar signalling increases yield and resilience. Nature **540**, 574–578.

Hatch MD, Kagawa T, Craig S. 1975. Subdivision of C_4-pathway species based on differing C_4 acid decarboxylating systems and ultrastructural features. Functional Plant Biology **2**, 111–128.

Hendriks JH, Kolbe A, Gibon Y, Stitt M, Geigenberger P. 2003. ADP-glucose pyrophosphorylase is activated by posttranslational redox-modification in response to light and to sugars in leaves of arabidopsis and other plant species. Plant Physiology **133**, 838–849.

Henry C, Bledsoe SW, Griffiths CA, Kollman A, Paul MJ, Sakr S, Lagrimini LM. 2015. Differential role for trehalose metabolism in salt-stressed maize. Plant Physiology **169**, 1072–1089.

Henry C, Bledsoe SW, Siekman A, Kollman A, Waters BM, Feil R, Stitt M, Lagrimini LM. 2014. The trehalose pathway in maize: conservation and gene regulation in response to the diurnal cycle and extended darkness. Journal of Experimental Botany **65**, 5959–5973.

Henry C, Watson-Lazowski A, Oszvald M, Griffiths C, Paul MJ, Furbank RT, Ghannoum O. 2020. Sugar sensing responses to low and high light in leaves of the C_4 model grass *Setaria viridis*. Journal of Experimental Botany **71**, 1039–1052.

Hua L, Stevenson SR, Reyna-Llorens I, Xiong H, Kopriva S, Hibberd JM. 2021. The bundle sheath of rice is conditioned to play an active role in water transport as well as sulfur assimilation and jasmonic acid synthesis. Plant Journal **107**, 268-286.

Ishihara H, Alseekh S, Feil R, *et al*. 2022. Rising rates of starch degradation during daytime and trehalose 6-phosphate optimize carbon availability. Plant Physiology **189**, 1976-2000.

Jenö P, Horst M. 2002. Electroelution of proteins from polyacrylamide gels. The Protein Protocols Handbook. Springer, 207–214.

John CR, Smith-Unna RD, Woodfield H, Covshoff S, Hibberd JM. 2014. Evolutionary convergence of cell-specific gene expression in independent lineages of C_4 grasses. Plant Physiology **165**, 62–75.

Kerbler SM, Armijos-Jaramillo V, Lunn JE, Vicente R. 2023. The trehalose 6-phosphate phosphatase family in plants. Physiologia Plantarum **175**, e14096.

Klein H, Gallagher J, Demesa-Arevalo E, *et al*. 2022. Recruitment of an ancient branching program to suppress carpel development in maize flowers. Proceedings of the National Academy of Sciences USA **119**, e2115871119.

Kretzschmar T, Pelayo MAF, Trijatmiko KR, *et al.* 2015. A trehalose-6-phosphate phosphatase enhances anaerobic germination tolerance in rice. Nature Plants **1**, 1-5.

Leegood RC. 1985. The intercellular compartmentation of metabolites in leaves of *Zea mays* L. Planta **164**,163 171.

Leegood RC. 1990. Enzymes of the Calvin cycle. In: Lea PJ, ed. Methods in plant biochemistry, Volume 3, Enzymes of primary metabolism. London: Academic Press, 15–37.

Leyman B, Van Dijck P, Thevelein JM. 2001. An unexpected plethora of trehalose biosynthesis genes in *Arabidopsis thaliana*. Trends in Plant Science **6**, 510-513.

Li P, Ponnala L, Gandotra N, *et al*. 2010. The developmental dynamics of the maize leaf transcriptome. Nature Genetics **42**, 1060-1067.

Lunn JE and Furbank RT. 1997. Localisation of sucrose-phosphate synthase and starch in leaves of C_4 plants. Planta **202**, 106–111.

Lunn JE and Furbank RT. 1999. Sucrose biosynthesis in C_4 plants. New Phytologist **143**, 221–237.

Lunn JE, Delorge I, Figueroa CM, Van Dijck P, Stitt M. 2014. Trehalose metabolism in plants. Plant Journal **79**, 544–567.

Lunn JE, Feil R, Hendriks JH, Gibon Y, Morcuende R, Osuna D, Scheible WR, Carillo P, Hajirezaei MR, Stitt M. 2006. Sugar-induced increases in trehalose 6-phosphate are correlated with redox activation of ADPglucose pyrophosphorylase and higher rates of starch synthesis in *Arabidopsis thaliana*. Biochemical Journal **397**, 139–148.

Lunn JE. 2007. Gene families and evolution of trehalose metabolism in plants. Functional Plant Biology **34**, 550–563.

Maier A, Zell MB, Maurino VG. 2011. Malate decarboxylases: evolution and roles of NAD(P)-ME isoforms in species performing C_4 and C_3 photosynthesis. Journal of Experimental Botany **62**, 3061–3069.

Majeran W, Friso G, Ponnal L, *et al.* 2010. Structural and metabolic transitions of C_4 leaf development and differentiation defined by microscopy and quantitative proteomics in maize. Plant Cell **22, 3509–3542.

Martin AP, Palmer WM, Brown C, Abel C, Lunn JE, Furbank RT, Grof CP. 2016. A developing *Setaria viridis* internode: an experimental system for the study of biomass generation in a C_4 model species. Biotechnology for Biofuels **9**, 1–12.

Martins MC, Hejazi M, Fettke J, *et al.* 2013. Feedback inhibition of starch degradation in Arabidopsis leaves mediated by trehalose 6-phosphate. Plant Physiology **163, 1142–1163.

Martins PK, Mafra V, de Souza WR, Ribeiro AP, Vinecky F, Basso MF, da Cunha BADB, Kobayashi AK, Molinari HBC. 2016. Selection of reliable reference genes for RT-

qPCR analysis during developmental stages and abiotic stress in *Setaria viridis*. Scientific Reports **6**, 1-10.

Nuccio ML, Wu J, Mowers R, *et al.* 2015. Expression of trehalose-6-phosphate phosphatase in maize ears improves yield in well-watered and drought conditions. Nature Biotechnology **33**, 862–869.

Nunes C, O'Hara LE, Primavesi LF, Delatte TL, Schluepmann H, Somsen GW, Silva AB, Fevereiro PS, Wingler A, Paul MJ. 2013. The trehalose 6-phosphate/SnRK1 signaling pathway primes growth recovery following relief of sink limitation. Plant Physiology **162**,1720-1732.

Oszvald M, Primavesi LF, Griffiths CA, Cohn J, Basu SS, Nuccio ML, Paul MJ. 2018. Trehalose 6-phosphate regulates photosynthesis and assimilate partitioning in reproductive tissue. Plant Physiology **176**, 2623–2638.

Pick TR, Brautigam A, Schluter U, *et al.* 2011. Systems analysis of a maize leaf developmental gradient redefines the current C_4 model and provides candidates for regulation. Plant Cell **23**, 4208–4220.

Ponnu J, Schlereth A, Zacharaki V, Dzialo MA, Abel C, Feil R, Schmid M, Wahl V. 2020. The trehalose 6-phosphate pathway impacts vegetative phase change in *Arabidopsis thaliana*. Plant Journal **104**, 768-780.

Ramon M, De Smet I, Vandesteene L, Naudts M, Leyman B, Van Dijck P, Rolland F, Beeckman T, Thevelein JM. 2009. Extensive expression regulation and lack of heterologous enzymatic activity of the Class II trehalose metabolism proteins from *Arabidopsis thaliana*. Plant, Cell and Environment **32**, 1015-1032.

Reeves G, Grangé-Guermente MJ, Hibberd JM. 2017. Regulatory gateways for cell-specific gene expression in C_4 leaves with Kranz anatomy. Journal of Experimental Botany **68**, 107–116.

Rojas BE, Hartman MD, Figueroa CM, Iglesias AA. 2021. Proteolytic cleavage of *Arabidopsis thaliana* phospho*enol*pyruvate carboxykinase-1 modifies its allosteric regulation. Journal of Experimental Botany **72**, 2514-2524.

Rojas BE, Tonetti T, Figueroa CM. 2023. Trehalose 6-phosphate metabolism in C_4 species. Current Opinion in Plant Biology **72**, 102347.

Sage RF. 2004. The evolution of C_4 photosynthesis. New Phytologist **161**, 341–370.

Sales CRG, Ribeiro R V., Hayashi AH, Marchiori PER, Silva KI, Martins MO, Silveira JAG, Silveira NM, Machado EC. 2018. Flexibility of C_4 decarboxylation and photosynthetic plasticity in sugarcane plants under shading. Environmental and Experimental Botany **149**, 34–42.

Satoh-Nagasawa N, Nagasawa N, Malcomber S, Sakai H, Jackson D. 2006. A trehalose metabolic enzyme controls inflorescence architecture in maize. Nature **441**, 227–230.

Schluepmann H, Pellny T, van Dijken A, Smeekens S, Paul MJ. 2003. Trehalose 6-phosphate is indispensable for carbohydrate utilization and growth in *Arabidopsis thaliana*. Proceedings of the National Academy of Sciences USA **100**, 6849–6854.

Schmittgen TD, Livak KJ. 2008. Analyzing real-time PCR data by the comparative CT method. Nature Protocols **3**, 1101–1108.

Shi D, Jouannet V, Agustí J, Kaul V, Levitsky V, Sanchez P, Mironova VV, Greb T. 2021. Tissue-specific transcriptome profiling of the Arabidopsis inflorescence stem reveals local cellular signatures. Plant Cell **33**, 200-223.

Stitt M, Heldt HW. 1985. Control of photosynthetic sucrose synthesis by fructose-2,6-bisphosphate : Intercellular metabolite distribution and properties of the cytosolic fructosebisphosphatase in leaves of *Zea mays* L. Planta **164**, 179–188.

Stitt M, Lilley RM, Gerhardt R, Heldt HW. 1989. Metabolite levels in specific cells and subcellular compartments of plant leaves. Methods in Enzymology **174**, 518–552.

Stitt M, Wirtz W, and Heldt HW. 1980. Metabolite levels in the chloroplast and extrachloroplast compartments of spinach protoplasts. Biochimica et Biophysica Acta **593**, 85-102.

Szecowka M, Heise R, Tohge T, *et al.* 2013. Metabolic fluxes in an illuminated Arabidopsis rosette. The Plant Cell **25**, 694–714.

Tao S, Liu P, Shi Y, *et al*. 2022. Single-cell transcriptome and network analyses unveil key transcription factors regulating mesophyll cell development in maize. Genes **13**, 374.

Trípodi KEJ, Rojas BE, Iglesias AA, Podestá FE. 2021. CAM plants as crops: metabolically flexible, hardy plants for a changing world. In: Handbook of plant and crop physiology, 4th Edition, CRC press, Mohammad Pessarakli (Ed), 1083-1098.

Usadel B, Bläsing OE, Gibon Y, Retzlaff K, Höhne M, Günther M, Stitt M. 2008. Global transcript levels respond to small changes of the carbon status during progressive exhaustion of carbohydrates in Arabidopsis rosettes. Plant Physiology **146**, 1834-1861.

van Dijken AJ, Schluepmann H, Smeekens SC. 2004. Arabidopsis trehalose-6-phosphate synthase 1 is essential for normal vegetative growth and transition to flowering. Plant Physiology **135**, 969–977.

Van Eck J, Swartwood K. 2015. *Setaria viridis*. In: Wang K, ed. Methods in Molecular Biology. Humana Press Inc., 57–67.

Van Leene J, Eeckhout D, Gadeyne A, *et al*. 2022. Mapping of the plant SnRK1 kinase signalling network reveals a key regulatory role for the class II T6P synthase-like proteins. Nature Plants **8**, 1245–1261.

Vandesteene L, Lopez-Galvis L, Vanneste K, *et al.* 2012. Expansive evolution of the trehalose-6-phosphate phosphatase gene family in Arabidopsis. Plant Physiology **160**, 884–896.

Vandesteene L, Ramon M, Le Roy K, Van Dijck P, Rolland F. 2010. A single active

trehalose-6-P synthase (TPS) and a family of putative regulatory TPS-like proteins in Arabidopsis. Molecular Plant **3**, 406–419.

Wahl V, Ponnu J, Schlereth A, Arrivault S, Langenecker T, Franke A, Feil R, Lunn JE, Stitt M, Schmid M. 2013. Regulation of flowering by trehalose-6-phosphate signaling in *Arabidopsis thaliana*. Science **339**, 704–707.

Wang L, Peterson R, Brutnell T. 2011. Regulatory mechanisms underlying C_4 photosynthesis. New Phytologist **190**, 9–20.

Wang S, Ouyang K, Wang K. 2019. Genome-wide identification, evolution, and expression analysis of *TPS* and *TPP* gene families in *Brachypodium distachyon*. Plants **8**, 362.

Wang Y, Bräutigam A, Weber APM, Zhu XG. 2014. Three distinct biochemical subtypes of C_4 photosynthesis? A modelling analysis. Journal of Experimental Botany **65**, 3567–3578.

Yadav UP, Ivakov A, Feil R, *et al.* 2014. The sucrose-trehalose 6-phosphate (Tre6P) nexus: specificity and mechanisms of sucrose signalling by Tre6P. Journal of Experimental Botany **65**, 1051–1068.

Yang HL, Liu YJ, Wang CL, Zeng QY. 2012. Molecular evolution of trehalose-6-phosphate synthase (TPS) gene family in populus, arabidopsis and rice. PLoS ONE **7**, e42438.

Figure legends

Figure 1. Phylogenetic relationships of *Setaria viridis* TPS proteins. TPS protein sequences were collated from *Setaria viridis* (Sv) and other selected species: Ac, *Ananas comosus*; At, *Arabidopsis thaliana*; Bd, *Brachypodium dystachion*; Mc, *Musa acuminata*; Os, *Oryza sativa*; Pt, *Populus trichocarpa*; Pv, *Phaseolus vulgaris*; Zm, *Zea mays* (see Supplementary Table S1 for gene/protein identifiers). A distance-based tree was reconstructed using the BioNJ algorithm. Numbers in nodes represent the bootstrap obtained for 1000 replicates. The tree is rooted on the *Escherichia coli* TPS (otsA). Sub-clades in the tree are coloured according to the position of *A. thaliana* TPS sequences: AtTPS1/2/3 (red), AtTPS5 (orange), AtTPS6 (yellow), AtTPS7 (green), AtTPS8/9/10 (blue) and AtTPS11 (violet), with the exception of the monocot-specific sub-clade (brown). *S. viridis* sequences are highlighted in grey.

Figure 2. Phylogenetic relationships of *Setaria viridis* TPP proteins. TPP protein sequences were collated from *Setaria viridis* (Sv) and other selected species: Ac, *Ananas comosus*; At, *Arabidopsis thaliana*; Bd, *Brachypodium dystachion*; Mc, *Musa acuminata*; Os, *Oryza sativa*; Pt, *Populus trichocarpa*; Pv, *Phaseolus vulgaris*; Zm, *Zea mays* (see Supplementary Table S1 for gene/protein identifiers). A distance-based tree was reconstructed using the BioNJ algorithm. Number in nodes represent the bootstrap obtained for 1000 replicates. The trees is rooted on the *Escherichia coli* TPP (otsB). Sub-clades in the tree are coloured according to the position of *A. thaliana* TPP sequences: AtTPPA/F/G (red) and AtTPPB/C/D/E/H/I/J (blue), with the exception of the monocot-specific sub-clade containing RAMOSA3 (ZmTPP10; brown). *S. viridis* sequences are highlighted in grey.

Figure 3. Relative expression of transcripts encoding Tre6P-related enzymes in MC and BSC from *S. viridis* leaves. Relative expression data were calculated with the $2^{-\Delta Ct}$ method, using the *SvKIN* transcript as a reference. Data are the mean ± standard error of 4 biological replicates. Statistical analysis was performed using a two-tailed t-test (unequal variance). Significant differences are indicated with asterisks: *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$. nd, not detected.

Figure 4. Immunodetection of SvTPS1 in different tissues and cell types of *S. viridis*. (A) Protein abundance in MC and BSC from *S. viridis* leaves. Data were retrieved from Calace *et. al*. (2021). MC, grey bars; BSC, black bars; AU, arbitrary units; nd, not detected. SvPEPC, Sevir.4G143500; SvNADP-MDH, Sevir.6G256400; SvRbcS, Sevir.3G276100; SvNADP-ME; Sevir.5G132500; SvTPS1, Sevir.3G180900. Data are the mean ± standard error of 4 biological replicates (n=4), except for SvTPS1, which was only detected in two BSC samples (n=2). Statistical analysis was performed using a two-tailed t-test (unequal variance). Significant differences are indicated with asterisks: **, $p<0.01$; ***, $p<0.001$. (B) Immunodetection of SvTPS1 in MC and BSC from *S. viridis* leaves extracted with TCA. Each lane was loaded with extract from 2 mg fresh weight of tissue. (C) Immunodetection of SvTPS1 in MC and BSC from *S. viridis* leaves extracted with Laemmli buffer. Each lane was loaded with extract from 2 mg fresh weight of tissue. (D) Immunodetection of SvTPS1 in different tissues and cell types from *S. viridis*. Each lane was loaded with 25 µg of protein. IF; inflorescence, FL; flag leaf, IN; internode, SL; seedlings, MC; mesophyll cells, BSC; bundle sheath cells, WL; whole leaf.

Figure 5. Distribution of selected metabolites in BSC and MC from *S. viridis* leaves. The abundances of each metabolite in BSC and MC were calculated as described in the methods

section from metabolite/PRK *vs* PEPC/PRK plots (as shown in Supplementary Figure S5). Data are the mean of 4 biological replicates (except for starch, n=3) from one representative experiment (Experiment 2, Supplementary Table S4).

Figure 6. Correlation analysis of metabolites in fractions of *S. viridis* leaves. Data presented in panels A-C correspond to whole-leaf samples from the time-course experiment (Supplementary Table S7), while data presented in panels D-F correspond to one representative fractionation experiment (Experiment 2, Supplementary Table S4). Pearson correlation coefficients (r) between Tre6P and glucose (A and D), fructose (B and E) or sucrose (C and F) were calculated with a 95% confidence interval.

Figure 1. Phylogenetic relationships of *Setaria viridis* TPS proteins. TPS protein sequences were collated from *Setaria viridis* (Sv) and other selected species: Ac, *Ananas comosus*; At, *Arabidopsis thaliana*; Bd, *Brachypodium dystachion*; Mc, *Musa acuminata*; Os, *Oryza sativa*; Pt, *Populus trichocarpa*; Pv, *Phaseolus vulgaris*; Zm, *Zea mays* (see Supplementary Table S1 for gene/protein identifiers). A distance-based tree was reconstructed using the BioNJ algorithm. Numbers in nodes represent the bootstrap obtained for 1000 replicates. The tree is rooted on the *Escherichia coli* TPS (otsA). Sub-clades in the tree are coloured according to the position of *A. thaliana* TPS sequences: AtTPS1/2/3 (red), AtTPS5 (orange), AtTPS6 (yellow), AtTPS7 (green), AtTPS8/9/10 (blue) and AtTPS11 (violet), with the exception of the monocot-specific sub-clade (brown). *S. viridis* sequences are highlighted in grey.

Figure 2. Phylogenetic relationships of *Setaria viridis* TPP proteins. TPP protein sequences were collated from *Setaria viridis* (Sv) and other selected species: Ac, *Ananas comosus*; At, *Arabidopsis thaliana*; Bd, *Brachypodium dystachion*; Mc, *Musa acuminata*; Os, *Oryza sativa*; Pt, *Populus trichocarpa*; Pv, *Phaseolus vulgaris*; Zm, *Zea mays* (see Supplementary Table S1 for gene/protein identifiers). A distance-based tree was reconstructed using the BioNJ algorithm. Number in nodes represent the bootstrap obtained for 1000 replicates. The trees is rooted on the *Escherichia coli* TPP (otsB). Sub-clades in the tree are coloured according to the position of *A. thaliana* TPP sequences: AtTPPA/F/G (red) and AtTPPB/C/D/E/H/I/J (blue), with the exception of the monocot-specific sub-clade containing RAMOSA3 (ZmTPP10; brown). *S. viridis* sequences are highlighted in grey.

Figure 3. Relative expression of transcripts encoding Tre6P-related enzymes in MC and BSC from *S. viridis* leaves. Relative expression data were calculated with the $2^{-\Delta Ct}$ method, using the *SvKIN* transcript as a reference. Data are the mean ± standard error of 4 biological replicates. Statistical analysis was performed using a two-tailed t-test (unequal variance). Significant differences are indicated with asterisks: *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$. nd, not detected.

Figure 4. Immunodetection of SvTPS1 in different tissues and cell types of *S. viridis*. (A) Protein abundance in MC and BSC from *S. viridis* leaves. Data were retrieved from Calace *et. al*. (2021). MC, grey bars; BSC, black bars; AU, arbitrary units; nd, not detected. SvPEPC, Sevir.4G143500; SvNADP-MDH, Sevir.6G256400; SvRbcS, Sevir.3G276100; SvNADP-ME; Sevir.5G132500; SvTPS1, Sevir.3G180900. Data are the mean ± standard error of 4 biological replicates (n=4), except for SvTPS1, which was only detected in two BSC samples (n=2). Statistical analysis was performed using a two-tailed t-test (unequal variance). Significant differences are indicated with asterisks: **, p<0.01; ***, p<0.001. (B) Immunodetection of SvTPS1 in MC and BSC from *S. viridis* leaves extracted with TCA. Each lane was loaded with extract from 2 mg fresh weight of tissue. (C) Immunodetection of SvTPS1 in MC and BSC from *S. viridis* leaves extracted with Laemmli buffer. Each lane was loaded with extract from 2 mg fresh weight of tissue. (D) Immunodetection of SvTPS1 in different tissues and cell types from *S. viridis*. Each lane was loaded with 25 µg of protein. IF; inflorescence, FL; flag leaf, IN; internode, SL; seedlings, MC; mesophyll cells, BSC; bundle sheath cells, WL; whole leaf.

Figure 5. Distribution of selected metabolites in BSC and MC from *S. viridis* leaves. The abundances of each metabolite in BSC and MC were calculated as described in the methods section from metabolite/PRK *vs* PEPC/PRK plots (as shown in Supplementary Figure S5). Data are the mean of 4 biological replicates (except for starch, n=3) from one representative experiment (Experiment 2, Supplementary Table S4).

Figure 6. Correlation analysis of metabolites in fractions of *S. viridis* leaves. Data presented in panels A-C correspond to whole-leaf samples from the time-course experiment (Supplementary Table S7), while data presented in panels D-F correspond to one representative fractionation experiment (Experiment 2, Supplementary Table S4). Pearson correlation coefficients (r) between Tre6P and glucose (A and D), fructose (B and E) or sucrose (C and F) were calculated with a 95% confidence interval.