Intercellular compartmentation of trehalose 6-phosphate metabolism

in *Setaria viridis* leaves

Tomás Tonetti[1,#], Bruno E. Rojas[1,#], Regina Feil[2], Camila Seimandi[1], Leandro E. Lucero[1],

Julieta V. Cabello[1], Paula Calace[3], Mariana Saigo[3], Stéphanie Arrivault[2], Mark Stitt[2], John E.

Lunn[2], Carlos M. Figueroa[1,*]

[1]Instituto de Agrobiotecnología del Litoral, UNL, CONICET, FBCB, Santa Fe, Argentina

[2]Max Planck Institute of Molecular Plant Physiology, Potsdam-Golm, Germany

[3]Centro de Estudios Fotosintéticos y Bioquímicos, UNR, CONICET, FCBF, Rosario,

Argentina

[#]Equal contribution

[*]Correspondence: carfigue@fbcb.unl.edu.ar

ORCID

TT, 0000-0002-3820-9411; BER, 0000-0002-3126-8363; RF, 0000-0002-9936-1337; CS,

0009-0007-2180-1666; LEL, 0000-0003-4453-8964; MSaigo, 0000-0002-2312-8782; SA,

0000-0003-0516-6950; MStitt, 0000-0002-4900-1763; JEL, 0000-0001-8533-3004; CMF,

0000-0003-4047-0480.

Running title

Compartmentation of Tre6P metabolism in *Setaria viridis*

Date of submission: 16[th] May 2025

Number of figures: 6

Supplementary data: 9 tables and 13 figures

Word count: 6990

Highlight

Trehalose 6-phosphate, a sugar signalling metabolite, is mainly present in the bundle sheath

cells of *Setaria viridis* leaves.

Abstract

Trehalose 6-phosphate (Tre6P) is a signal metabolite that links carbon metabolism with plant

development. Our current understanding of Tre6P metabolism and signalling is

predominantly based on studies in *Arabidopsis thaliana*. Some features could be adapted to

the specific physiology, anatomy, and life cycle of this C_3 eudicot model species, and thus

might not be representative of other angiosperms. To better understand Tre6P metabolism in

monocot C_4 species we used *Setaria viridis*, which has been widely adopted as a model for

the major C_4 NADP-malic enzyme subtype crop species, such as maize (*Zea mays*), sorghum

(*Sorghum bicolor*) and sugarcane (*Saccharum officinarum*). In this work, we analysed the

levels of transcripts encoding Tre6P-related enzymes in different tissues and cell types from

S. viridis. The *TREHALOSE-6-PHOSPHATE SYNTHASE1* transcript, its encoded protein

(TPS1, the enzyme responsible for Tre6P synthesis) and Tre6P were mainly located in bundle

sheath cells of *S. viridis*. Our results show that Tre6P is predominately synthesized and

located in bundle sheath and associated cells.

Keywords

bundle sheath cells, C$_4$ photosynthesis, mesophyll cells, *Setaria viridis*, sucrose, trehalose 6-phosphate

Abbreviations

BSC, bundle sheath cells; MC, mesophyll cells; NADP-MDH, NADP-dependent malate dehydrogenase; NADP-ME, NADP-dependent malic enzyme; PEPC, phospho*enol*pyruvate carboxylase; PRK, phosphoribulokinase; SWEET, SUGARS WILL EVENTUALLY BE EXPORTED TRANSPORTERS; Tre6P, trehalose 6-phosphate; TPS, Tre6P synthase; TPP, Tre6P phosphatase; TRE, trehalase.

Introduction

Trehalose is a non-reducing disaccharide widely distributed in nature (Figueroa and Lunn, 2016). In plants, the pathway of trehalose synthesis occurs through the phosphorylated intermediate trehalose 6-phosphate (Tre6P), which is synthesized from UDP-glucose (UDPGlc) and glucose 6-phosphate (Glc6P) by Tre6P synthase (TPS; EC 2.4.1.15) and dephosphorylated to trehalose by Tre6P phosphatase (TPP; EC 3.1.3.12; Cabib and Leloir, 1958; Avonce *et al.*, 2006). Trehalose is then hydrolysed to glucose by trehalase (TRE; EC 3.2.1.28). Gene families coding for TPS and TPP are present in species from all of the main taxonomic groups of green plants (Avonce *et al.*, 2006; Lunn, 2007; Lunn *et al.*, 2014; Figueroa and Lunn, 2016). Plant TPSs are divided into two distinct clades: class I and class II. Class I TPSs are responsible for Tre6P synthesis, while class II TPSs (which lack some of the glycosyltransferase active site residues involved in substrate binding and catalysis) are putatively involved in Tre6P perception and/or signalling (Avonce *et al.*, 2006; Lunn, 2007; Vandesteene *et al.*, 2010; Lunn *et al.*, 2014; Delorge *et al.*, 2015; Figueroa and Lunn, 2016; Van Leene *et al.,* 2022). In the last two decades, studies performed with mutant and transgenic lines demonstrated the importance of Tre6P for all stages of plant development (Eastmond *et al.*, 2002; van Dijken *et al.*, 2004; Gomez *et al.*, 2006; Gomez *et al.*, 2010; Wahl *et al.*, 2013; Fichtner *et al.*, 2020; Ponnu *et al.*, 2020; reviewed in Fichtner and Lunn, 2021).

In *Arabidopsis thaliana* leaves, Tre6P is synthesized primarily in phloem parenchyma (bundle sheath) cells and in the companion cell-sieve element complex of the phloem (Fichtner *et al.*, 2020). The phloem parenchyma cells are symplastically connected with mesophyll cells (MC) via plasmodesmata, allowing Tre6P to diffuse into the MC to regulate sucrose production (Fichtner and Lunn, 2021). Tre6P regulates the synthesis and transport of sucrose by influencing photoassimilate partitioning between sucrose or organic and amino

acids during the day (Figueroa *et al.*, 2016), transitory starch degradation during the night and

towards the end of the light period (Martins *et al.*, 2013; dos Anjos *et al.*, 2018; Ishihara *et*

al., 2022), and expression of *SWEET* (*SUGARS WILL EVENTUALLY BE EXPORTED*

TRANSPORTER) genes, which encode transporters involved in sucrose phloem loading

(Oszvald *et al.*, 2018; Fichtner *et al.*, 2021). Therefore, Tre6P coordinates the synthesis and

transport of sucrose and amino acids from source leaves with the demand from growing and

sink tissues (Martins *et al.*, 2013; Figueroa *et al.*, 2016; dos Anjos *et al.*, 2018). In addition,

Tre6P exerts widespread transcriptional repression of photosynthesis, gluconeogenesis and

carbon starvation responses, induces amino acid synthesis, nucleotide synthesis and ribosome

formation and interacts with light signaling and many phytohormone signaling pathways

(Avidan *et al.*, 2024).

Our current understanding of Tre6P metabolism and signalling in plants is mainly

based on studies in *A. thaliana* (Fichtner and Lunn, 2021; Rojas *et al.* 2023). This model

eudicot is a rosette-forming, annual weed species that performs C_3 photosynthesis and grows

predominantly in temperate zones. Some features of Tre6P metabolism and signalling in *A.*

thaliana might be adapted to its specific physiology, anatomy and life cycle, and so might

differ from other species. Tre6P metabolism and signalling have also been investigated in

maize (*Zea mays*), a domesticated monocot species that performs C_4 photosynthesis and

originated in subtropical regions. These studies have revealed the importance of Tre6P

metabolism and signalling in the coordination of metabolism in source and sink zones,

inflorescence and shoot branching, floral identity, tillering, seed development and tolerance

to water deficit (Satoh-Nagasawa *et al.*, 2006; Henry *et al.*, 2014, 2015; Nuccio *et al.*, 2015;

Czedik-Eysenberg *et al.*, 2016; Bledsoe *et al.*, 2017; Oszvald *et al.*, 2018; Claeys *et al.*, 2019;

Dong *et al.*, 2019; Klein *et al.*, 2022). However, how far Tre6P metabolism and signalling are

adapted to the specialized C_4 photosynthetic metabolism and leaf anatomy of this species is

poorly understood.

111   Tre6P regulates photoassimilate partitioning between sucrose and organic acids and

amino acids in *A. thaliana* via coordinated changes in the activities of nitrate reductase,

phospho*enol*pyruvate carboxylase (PEPC) and, by inference, the mitochondrial pyruvate

dehydrogenase (Figueroa *et al.*, 2016). Artificially increased levels of Tre6P reduced the

mono-ubiquitination and increased the phosphorylation status of PEPC, thus activating the

enzyme. These changes increased carbon flux into organic acids and the TCA cycle to

provide carbon skeletons for amino acid synthesis. Given that PEPC is responsible for

primary carbon fixation in the C_4 pathway, Tre6P might therefore play a direct role in

regulating photosynthetic carbon fixation in C_4 plants. This raises the question of whether

Tre6P is also involved in the regulation of photoassimilate partitioning to amino acids in C_4

plants and, if so, how flux into metabolites in the C_4 cycle and flux into the TCA cycle and

amino acid synthesis are reconciled. Tre6P also regulates the mobilization of transitory starch

reserves in *A. thaliana* leaves and provision of sucrose at night (Martins *et al.*, 2013; dos

Anjos *et al.*, 2018; Ishihara *et al.*, 2022). In contrast to C_3 plants, where sucrose and starch

synthesis both occur within MC, in C_4 plants starch synthesis is restricted to the bundle

sheath cells (BSC), while sucrose is made predominantly in MC (Lunn and Furbank, 1997;

1999). If Tre6P also regulates photoassimilate partitioning and transitory starch turnover in

C_4 plants, this cellular compartmentation of sucrose and starch metabolism would add a

further layer of complexity. It has been proposed that expression of the genes encoding the

SWEET and SUT/SUC (sucrose-H^+ symporters) transporters had to be adapted to the

specialized Kranz anatomy in C_4 species, and was potentially a critical element in the

evolution of C_4 plants (Furbank and Kelly, 2021). Mutation of the three maize *ZmSWEET13*

genes produced plants with impaired sugar phloem loading and drastic effects on growth

(Bezrutczyk *et al.*, 2018). As Tre6P has been implicated in the transcriptional regulation of

SWEET genes (Oszvald *et al.*, 2018; Fichtner *et al.*, 2021), it is possible that Tre6P also plays

a key role in modulating phloem loading during the establishment of Kranz anatomy in

developing leaves of C_4 species.

Currently, we have little knowledge about Tre6P signalling in the leaves of C_4 plants

(Rojas *et al.*, 2023). By extrapolation from its known functions in leaves of C_3 plants, we

hypothesize that Tre6P plays a role in the regulation of carbon fixation, photoassimilate

partitioning and phloem loading in the leaves of C_4 plants. If so, Tre6P signalling in these

species may have unique features, reflecting their specialized metabolism and anatomy. For

instance, the *TREHALOSE-6-PHOSPHATE SYNTHASE1* transcript was mainly found in

BSC of several C_4 species (Benning *et al.*, 2023). To better understand how Tre6P signalling

operates in C_4 crops, we investigated the compartmentation of Tre6P in leaves of *Setaria*

viridis (wild foxtail millet), a model species for translational research in plants performing

NADP-dependent malic enzyme (NADP-ME) C_4 photosynthesis (Brutnell *et al.*, 2010; Doust

et al., 2019). Our results show that the *TPS1* transcript, its encoded protein (TPS1) and Tre6P

are mainly located in BSC.

Materials and methods

Plant growth and harvest conditions

Seeds of *Setaria viridis* (L.) P. Beauv. accession A10.1 were obtained from the Germplasm

Resources Information Network (GRIN; http://www.ars-grin.gov/), United States Department

of Agriculture (PI 669942). To increase the germination rate, the seed coat was abraded with

sandpaper (Van Eck and Swartwood, 2015). Seeds were surface sterilized with 0.55 g l^{-1}

sodium hypochlorite containing 0.1% (v/v) Tween-20 for 5 minutes, followed by three

washes with sterile deionized water. The seeds were then sown on plates with half-strength

Murashige and Skoog medium (PhytoTech Labs; https://phytotechlab.com/) supplemented

with 0.8% (w/v) agar. Plates were placed in a growth chamber at 28°C with an irradiance of

120 µmol m^{-2} s^{-1}, provided by white fluorescent tubes, under a long-day photoperiod regime

(16 h light/8 h dark). After five days, seedlings were transferred to individual 8-cm diameter

pots containing Klasmann TS1 substrate and irrigated with 1X Hoagland solution. For the

analysis of transcripts and proteins, plants were grown in a Conviron Adaptis A1000

controlled environment chamber (https://www.conviron.com/) with an irradiance of 350

µmol m^{-2} s^{-1}, provided by white fluorescent tubes, with a long-day photoperiod regime, at

28/22°C (day/night), and 50% relative humidity for 2 weeks. For metabolite fractionation

experiments, plants were grown in a walk-in phytotron with an irradiance of 550 µmol m^{-2} s^{-}

^{1}, provided by LEDs, at 28/22°C (day/night) and 65% relative humidity. Harvested plants had

eight fully developed leaves and inflorescences were not macroscopically visible. The blades

from the 5th and 6th true leaves (counting from the bottom of the plant) were harvested in the

middle of the photoperiod. The leaf material used for isolation of MC and BSC was

immediately processed and then stored at -80°C. For enzymatic and metabolite analyses, leaf

samples were rapidly quenched in liquid nitrogen and stored at -80°C until use. Samples for

the analysis of transcripts and proteins (seedlings, inflorescences, the fourth internode and the

flag leaf) were frozen in liquid nitrogen and stored at -80°C until use.

Phylogenetic analysis

TPS, TPP and TRE protein sequences from *S. viridis* (genome version 2.1) were identified

using the keyword "trehalose" at the Phytozome 13 server (https://phytozome.jgi.doe.gov/).

We followed the same approach to retrieve TPS and TPP protein sequences from *Ananas*

comosus (genome version 3) and *Musa acuminata* (genome version 1). Sequences were

manually curated to remove incomplete and duplicated entries. TPS and TPP protein

sequences from *A. thaliana*, *Brachypodium distachyon*, *Oryza sativa*, *Phaseolus vulgaris*,

Populus trichocarpa, *Triticum aestivum* and *Z. mays* were obtained from previous

publications (Yang *et al.*, 2012; Henry *et al.*, 2014; Barraza *et al.*, 2016; de Haro *et al.*, 2019;

Wang *et al.*, 2019; Du *et al.*, 2022; Griffiths *et al.*, 2025). Sequences were aligned with the

CLUSTALW tool available at the GenomeNet server (https://www.genome.jp/tools-

bin/clustalw). Distance-based phylogenetic trees were reconstructed with the SeaView

software version 4.3.0 (Gouy *et al.*, 2010) using the BioNJ algorithm (Gascuel, 1997) and

1000 replicates. Figures were prepared with the FigTree program version 1.3.1

(http://tree.bio.ed.ac.uk/software/figtree/). Sequences used to reconstruct the trees were

labelled as shown in Supplementary Table S1.

Separation of mesophyll and bundle sheath cells

Fractions highly enriched in MC or BSC strands were obtained from fresh leaves following

the methods described by John *et al.* (2014), with the modifications introduced by Calace *et*

al. (2021). In both cases, each replicate consisted of the material obtained from 20 plants.

These samples were used for the analysis of transcripts and proteins. To assess the suitability

of the fractionation procedure, we performed western blots using commercial antibodies

raised against the large subunit of Rubisco (RbcL; Agrisera AS03 037;

https://www.agrisera.com/) and PEPC (Agrisera AS09 458), as described below for the

immunoblotting of SvTPS1. Supplementary Fig. S1 shows that Rubisco and PEPC were

exclusively detected in BSC and MC fractions, respectively.

To determine the distribution of metabolites, frozen leaf tissue was finely ground in

liquid nitrogen and the frozen tissue powder was passed through a series of nylon meshes of

different pore sizes (200, 80, and 41 µm). The material that was retained on each filter and

the one that passed through the 41 µm filter were collected to obtain the samples named

F200, F80, F40, and FT, respectively. The F200 fraction, depending on the grinding, often

has larger whole leaf bits but tends to be enriched in BSC; the F80 fraction is especially

enriched in BSC; the F40 fraction is usually a mix of BSC and MC; and FT fraction is

especially enriched in MC (Stitt and Heldt, 1985). This method is suitable for the

quantification of metabolites with short half-lives (Arrivault *et al.*, 2009, 2017). Metabolites

and enzymes, including cell-specific marker enzymes, were measured in each fraction, and

their distribution between MC and BSC were calculated by linear regression, as described by

Stitt and Heldt (1985). Two C_4 photosynthetic enzymes were used as markers for each cell-

type: PEPC and NADP-dependent malate dehydrogenase (NADP-MDH) for MC and NADP-

ME and phosphoribulokinase (PRK) for BSC. Each biological replicate consisted of the

material obtained from 20 plants.

Total RNA extraction and cDNA synthesis

Total RNA was extracted from 50 mg fresh weight (FW) of finely ground tissue with the

TransZol reagent (TransGen Biotech; https://www.transgenbiotech.com/), following the

manufacturer's instructions. After RNA extraction, samples were treated with DNase

(Promega; https://www.promega.com) and RNA concentration was quantified using a

NanoDrop 2000 (Thermo Scientific; https://www.thermofisher.com). cDNA was synthesized

with the MultiScribe Reverse Transcriptase (Applied Biosystems;

https://www.thermofisher.com), according to the manufacturer's instructions.

Determination of transcript levels

Real-time reverse transcriptase-qPCR (RT-qPCR) reactions were performed in a final volume

of 8 µl in 96-well microplates (Applied Biosystems) with the iQ SYBR Green Supermix

(Bio-Rad; https://www.bio-rad.com), 0.2 µM of each primer, and an appropriate dilution of

the sample. Transcript abundance was calculated with the comparative C_t method

(Schmittgen and Livak, 2008) and normalization was performed with the *SvKIN* transcript

(Sevir.4G252700; Martins *et al.*, 2016), which showed similar C_t values across all the

analysed samples (Supplementary Fig. S2). All the primers used in this work are listed in

Supplementary Table S2.

Production of a polyclonal antiserum against SvTPS1

The protein coding region of the class I *TPS* gene (SvTPS1; Uniprot ID A0A4U6VAG9) was

codon-optimised and *de novo* synthesized (BioBasic; https://www.biobasic.com/) for

expression in *E. coli* (Supplementary File S1) and cloned into the pET28b expression vector

(Novagen; https://www.merckmillipore.com) between the NdeI and SacI restriction sites, in

frame with a His$_6$-tag at the N-terminus. The resulting vector was used to transform *E. coli*

BL21 (DE3) cells (Invitrogen; https://www.thermofisher.com). Transformed cells were

grown in Lysogeny Broth medium, supplemented with 50 µg ml^{-1} kanamycin at 37°C, with

shaking at 200 rpm until the optical density at 600 nm reached approximately 0.6. Synthesis

of the recombinant protein was induced by addition of isopropyl-β-D-thiogalactopiranoside to

a final concentration of 1 mM and cells were incubated for 16 h at 25°C and 200 rpm after

induction. The recombinant SvTPS1 protein was mainly present in the insoluble fraction;

thus, it was purified from inclusion bodies (Jenö and Horst, 2002). This protein was used to

raise a polyclonal antiserum in rabbits at ICIVET Litoral (UNL-CONICET, Argentina).

Antibodies were affinity-purified using a nitrocellulose membrane strip containing the

recombinant SvTPS1, as described by Fang (2012).

Protein extraction

Plant material was homogenized in a pre-cooled mortar with liquid nitrogen. For denaturing

protein extraction, 20 mg FW of frozen tissue were extracted with 200 µl of 125 mM Tris-

HCl pH 6.8, 2% (w/v) sodium dodecyl sulphate (SDS), 20% (v/v) glycerol, 1.4 M 2-

mercaptoethanol, and 0.05% (w/v) Bromophenol Blue. After adding the buffer, samples were

vortex-mixed and heated for 5 min at 95°C with agitation. Samples were then cooled to room

temperature and centrifuged at 20,000 x g for 10 min to separate the protein extract from

insoluble tissue debris. Alternatively, 50 mg FW of frozen tissue were resuspended in 500 µl

of 10 mM Tris-HCl pH 8.0, 10% (v/v) trichloroacetic acid (TCA), 25 mM ammonium

acetate, and 1 mM EDTA (Rojas *et al.*, 2021). Samples were incubated at 4°C for 30 min and

centrifuged at 20,000 x g for 15 min. The supernatant was discarded and the pellet was

washed with 500 µl of 5% (w/v) TCA and then with 500 µl of 90% (v/v) acetone, then

allowed to dry before the protein pellet was dissolved in 50 µL of Laemmli buffer for

analysis by SDS-PAGE.

Protein extraction for enzymatic assays was performed as described by Gibon *et al.*

(2004), with minor modifications. Samples (20 mg FW of frozen tissue) were supplemented

with 10 mg of polyvinylpolypyrrolidone and then resuspended in 500 µl of 50 mM HEPES-

KOH pH 7.5, 10 mM $MgCl_2$, 1 mM EDTA, 1 mM EGTA, 1 mM benzamidine, 1 mM ε-

aminocaproic acid, 0.02 mM leupeptin, 1 mM DTT, 1 mM phenylmethylsulfonyl fluoride,

0.1% (v/v) Triton X-100, and 20% (v/v) glycerol. Samples were incubated on ice for 10 min

and then centrifuged at 20,000 x g and 4°C for 10 min to separate soluble proteins from tissue

debris.

Protein concentration was determined with the Bradford reagent (Bradford, 1976),

using a standard curve constructed with bovine serum albumin.

Immunoblotting

Protein extracts were subjected to a 10% SDS-PAGE and electrotransferred to a 0.45 μm

nitrocellulose membrane (Amersham; https://www.cytivalifesciences.com/). To check lane

loadings, membranes were stained with 0.01% (w/v) Ponceau S in 1% (v/v) acetic acid. After

washing with TBST1 solution [50 mM Tris-HCl pH 8.0; 150 mM NaCl; 0.2% (v/v) Tween-

20], membranes were blocked for 1 h with TBST1 supplemented with 0.2% (w/v) fat-free

dried milk powder. Membranes were incubated overnight at 4°C with purified anti-SvTPS1

in TBST1 supplemented with 0.2% (w/v) fat-free dried milk powder. After washing, the

membranes were incubated for 30 min at room temperature with goat anti-rabbit IgG (H&L)

conjugated to horseradish peroxidase (Agrisera AS09 602). Proteins were detected using the

ECL Bright reagent (Agrisera AS16 ECL-N) and exposed to X-ray films (AGFA;

https://www.agfa.com/corporate/) for 1 min. The sizes of the immunoreactive bands were

estimated from the relative mobility (Rf) *versus* logarithm of molecular weight (logMM) plot

constructed with protein standards, using data from three independent experiments.

Immunoprecipitation of SvTPS1

Proteins were extracted from 100 mg FW of BSC with 400 μl of a buffer containing 100 mM

Bicine pH 9.0, 10% (v/v) glycerol, 0.1% (v/v) Triton X-100, 5 mM 2-mercaptoethanol, 1 mM

EDTA, 1 mM EGTA, 2 mM phenylmethylsulfonyl fluoride and 1 X Protease Inhibitor

Cocktail   III   (Calbiochem;   https://www.sigmaaldrich.com/).   Immunoprecipitation   was

performed   as   described   by   Hartman   *et al.*   (2023),   with   minor   modifications.   Briefly,

antibody-bead conjugates were prepared by adding 5 μl of polyclonal anti-SvTPS1 antibody

to a microcentrifuge tube containing 200 μl of TBST2 [50 mM Tris-HCl pH 8.0; 150 mM

NaCl; 0.05% (v/v) Tween-20] and 75 μl of protein A-Sepharose beads (Sigma). Conjugates

were incubated 90 min at room temperature with gently mixing and then washed with TBST2

to remove unbound antibodies. The protein extract (150 μl) was incubated with the antibody-

bead conjugate for 16 h at 4°C with gently mixing. After washing with TBST2, the pellet was

resuspended with Laemmli buffer and heated at 95°C for 10 min. The sample was then

centrifuged at room temperature and 10,000 x g for 5 min and the resulting supernatant was

separated by SDS-PAGE. The gel was stained using colloidal Coomassie Blue G-250.

Controls were processed in parallel either without antibodies or the protein extract.

Mass spectrometry analysis

Tryptic peptides from the gel fragment were obtained using the protocol of Link and LaBaer

(2009) and analysed by mass spectrometry (MS) at the Mass Spectrometry Unit of the

Institute of Molecular and Cellular Biology of Rosario (UNR-CONICET), Argentina. The

sample (3 µl) was injected on an nanoHPLC Ultimate3000 (Thermo Scientific) and peptides

were separated using a nano column EASY-Spray ES901 (15 cm × 50 µm ID, PepMap RSLC

C18). The mobile phase flow rate was 300 nl/min using 0.1% formic acid in water (solvent

A) and 0.1% formic acid and 100% acetonitrile (solvent B). The gradient profile was set as

follows: 4-30% solvent B for 10 min, 30%-80% solvent B for 2 min and 90% solvent B for 1

min. MS analysis was performed using a Q-Exactive HF mass spectrometer (Thermo

Scientific). For ionization, 1,9 kV of liquid junction voltage and 250°C of capillary

temperature were used. The full scan method employed a m/z 375–2000 mass selection, an

Orbitrap resolution of 120,000 (at m/z 200), a target automatic gain control (AGC) value of

1e6 and a maximum injection time of 100 ms. After the survey scan, the 5 most intense

precursor ions were selected for MS/MS fragmentation.

Fragmentation was performed with a normalized collision energy of 27 eV and

MS/MS scans were acquired with a dynamic first mass, AGC target was 5e5, resolution of

30000 (at m/z 200), isolation window of 1.4 m/z units and maximum IT was 55 ms. Charge

state screening was enabled to reject unassigned, singly charged, and equal or more than six

protonated ions. A dynamic exclusion time of 10 s was used to discriminate against

previously selected ions.

MS data were analysed with Proteome Discoverer 2.4.1.15, using standardized

workflows. The mass spectra raw file was searched against the *Setaria viridis* deduced

proteome (UniProt ID UP000298652). Precursor and fragment mass tolerance were set to 10

ppm and 0.02 Da, respectively, allowing 2 missed cleavages, with Cys carbamidomethylation

(+57.021 Da) as fixed modification, and the following dynamic modifications: Met oxidation

(+15.995 Da), N-term acetylation (+42.011 Da), Met loss (-131.040 Da) and Met loss plus

acetylation (-89.030 Da).

The raw proteomics data were deposited to the PRIDE repository

(https://www.ebi.ac.uk/pride/) with the dataset identifier PXD057139.

Immunohistochemistry

Tissue fixation, dehydration, assembly of the paraffin block and microtome cutting were

performed as described by Cabello and Chan (2019), with minor modifications. Sections of

0.5-1.0 cm in length from the sixth leaf of three-week-old plants (without the midrib) were

fixed at 25°C for 1 h in a solution containing 3.7% (v/v) formaldehyde, 5% (v/v) acetic acid

and 47.5% (v/v) ethanol. To further fix and discolour the tissue, sections were incubated in

70% (v/v) ethanol for two weeks. Sections were dehydrated through a series of ethanol

solutions (80, 90, 96 and 100%; 30 min each), followed by incubation in ethanol-xylene

solutions (3:1; 1:1 and 1:3; 1 h each), and finally stored overnight in 100% xylene at room

temperature. To obtain the paraffin blocks, sections were incubated through a series of

xylene-paraffin solutions (3:1, 1:1 and 1:3; 1 h each). The samples were placed into plastic

casts and finally embedded in Histoplast (Biopack; https://www.biopack.com.ar/). Each block

was incubated overnight at room temperature to ensure solidification. Cross-sections (10 μm

thick) were obtained using a Microtome RM2125 (Leica Microsystems; https://www.leica-

microsystems.com/) and mounted on slides coated with 50 mg/ml poly-D-Lys (Sigma) in

10 mM Tris-HCl pH 8.0 and dried for 16 h at 37°C. Paraffin was removed with 100% xylene

for 15 min at room temperature and sections were rehydrated using a series of ethanol (100,

96, 90, 80, 70 and 50%; 1 min each) to finish in distilled water.

Immunolocalization of proteins in *S. viridis* leaves was performed as described by

Sauer and Friml (2010), with minor modifications. Mounted cross-sections were rinsed three

times in PBS (10 mM sodium phosphate, 130 mM NaCl, pH 7.4) and blocked with PBS with

the addition of 1% (w/v) BSA and 0.1% (v/v) Tween-20 (blocking solution) at room

temperature for 1 h in a humid chamber. The blocking solution was removed and sections

were incubated for 2 h at room temperature with the anti-RbcL (Agrisera AS03 037; dilution

1:1000) or the purified anti-SvTPS1 (see above; dilution 1:100) in blocking solution. Tissue

sections were washed three times with blocking solution and then incubated with Cy2-

conjugated   goat   anti-rabbit   IgG   (Jackson   ImmunoResearch   111-225-144;

https://www.jacksonimmuno.com/; dilution 1:200). Samples were incubated at room

temperature for 1 h, washed three times with blocking solution and air-dried. Sections

incubated only with the secondary antibody were used as a negative control. Finally, samples

were mounted with 85% (v/v) glycerol and a coverslip. Micrographs were taken with a SP8

LIGHTNING confocal microscope (Leica Microsystems) using the 20X objective and the

Cy2 bis NHS ester fluorophore channel (excitation 492 nm, emission 510 nm). Images were

processed using the LAS X Office 1.4.7.28982 software (Leica Microsystems) and figures

were prepared with Inkscape 0.92.2 (https://inkscape.org/).

Metabolite extraction and measurement

Phosphorylated intermediates and organic acids were extracted with a mixture of

chloroform:methanol (3:7) and then measured by anion-exchange high-performance liquid

chromatography coupled to tandem mass spectrometry (LC-MS/MS), as described by Lunn *et*

al. (2006), with the modifications introduced by Figueroa *et al.* (2016). Calvin-Benson cycle

intermediates and aspartate were measured in chloroform-methanol extracts by ion-pair

reverse   phase   LC-MS/MS,   as   described   by   Arrivault   *et al.*   (2009).   Pyruvate,

phospho*enol*pyruvate (PEP), dihydroxyacetone phosphate (DHAP) and 3-phosphoglycerate

(3PGA) were enzymatically measured in TCA extracts (Arrivault *et al.*, 2017). The content of

sucrose, glucose, and fructose in ethanolic extracts was determined following the method of

Stitt *et al.* (1989), while the amount of starch was measured using the insoluble residue from

the ethanolic extraction, as described by Hendriks *et al.* (2003). The datasets containing

metabolite levels and the associated calculations are available in Supplementary Tables S3-

S5.

Enzyme activity assays

PRK activity was measured spectrophotometrically using a continuous assay (Leegood,

1990), with minor modifications. The assay mixture contained: 50 mM Tricine-KOH pH 8.0,

1 mM ATP, 50 mM KCl, 0.3 mM NADH, 10 mM $MgCl_2$, 2 mM PEP, 2 U pyruvate kinase,

and 2 U lactate dehydrogenase and the reaction was started by adding 0.5 mM ribulose 5-

phosphate. PEPC, NADP-MDH, and NADP-ME activities were measured as described by

Ashton *et al.* (1990), with minor modifications. The PEPC assay mixture contained: 50 mM

Tricine-KOH pH 8.0, 5 mM $MgCl_2$, 2 mM DTT, 1 mM $KHCO_3$, 5 mM Glc6P, 0.3 mM

NADH, and 0.4 U malate dehydrogenase and the reaction was started with 2 mM PEP. The

NADP-MDH assay mixture contained: 50 mM Tricine-KOH pH 8.0, 70 mM KCl, 1 mM

EDTA, 1 mM DTT, and 0.3 mM NADPH and the reaction was started with 1 mM

oxaloacetate (freshly prepared). The NADP-ME assay mixture contained: 50 mM Tricine-

KOH pH 8.0, 0.5 mM $NADP^+$, 0.1 mM EDTA, and 2 mM $MgCl_2$ and the reaction was

started with 5 mM malate. All reactions were performed in a final volume of 200 µl at 25°C

in 96-well microplates. The reduction/oxidation of $NAD(P)^+$/NAD(P)H was followed by

measuring the absorbance at 340 nm in a Biotek ELx808 microplate reader. The amount of

crude extract added to the reaction was chosen to ensure a linear response of the signal. One

unit (U) of enzyme activity is defined as the amount of extract producing 1 µmol of product

in 1 min under the specified assay conditions.

Results

Phylogenetic analysis of Tre6P-related enzymes

To identify the proteins involved in Tre6P metabolism in *S. viridis*, we performed keyword

searches against its deduced proteome. We found 21 putative Tre6P-related enzymes: 10

TPSs, 10 TPPs, and 1 TRE (Supplementary Table S1), as previously observed for *Setaria*

italica (Henry et al., 2020), a domesticated relative of *S. viridis*. After removing duplicated

sequences or splicing variants (for SvTPP1, SvTPP8, and SvTPP10), phylogenetic trees were

reconstructed for TPS and TPP proteins (Fig. 1 and Supplementary Fig. S3, respectively).

S. viridis TPS sequences were split into two major clades containing class I and class

II TPSs, as previously described for TPSs from *A. thaliana*, poplar, rice, pea, wheat and

maize (Leyman *et al*., 2001; Avonce *et al.*, 2006; Lunn, 2007; Yang *et al.*, 2012; Henry *et al.*,

2014; Barraza *et al.*, 2016; Paul *et al.*, 2018; de Haro *et al.*, 2019). *S. viridis* has only one

class I TPS (SvTPS1), which diverges from the other nine class II TPS proteins (SvTPS2-10;

Fig. 1). All *S. viridis* TPS sequences displayed the glycosyl-transferase family 20 domain

(InterPro ID IPR001830) and the TPP domain (InterPro ID IPR003337) that are typical of

plant TPS proteins. Inspection of the primary sequence of *S. viridis* TPSs in detail showed

that only SvTPS1 had the characteristic residues involved in UDPGlc and Glc6P binding

(Supplementary Fig. S4). The class II TPS sequences from *S. viridis* (SvTPS2-10) were

distributed into several branches, most of which contained proteins from eudicot and

monocot species. The exceptions were the monocot-specific sub-clade containing

SvTPS2/3/9 and that containing *M. acuminata* TPS8 and TPS16, which branched at the base

of the class II TPS tree (Fig. 1), an observation initially made by Lunn (2007). A detailed

inspection of *S. viridis* class II TPS sequences showed substitutions in the substrate-binding

residues of the glucosyltransferase domain that might explain their apparent lack of catalytic

activity (Supplementary Fig. S4; Gibson *et al.*, 2002; Vandesteene *et al.*, 2010). The

phylogenetic and sequence analyses indicate that SvTPS1 is the enzyme solely responsible

for Tre6P synthesis in *S. viridis*.

The *S. viridis* TPP family was separated into two major clades: three members in

clade A and the other seven in clade B (Supplementary Fig. S3). All TPP sequences from *S.*

viridis had the TPP domain (InterPro ID IPR003337) and the three conserved active site

motifs previously described for these proteins (Avonce *et al.*, 2006; Lunn, 2007; Vandesteene

et al., 2012; Henry *et al.*, 2014; Kerbler *et al.*, 2023); thus, all 10 *S. viridis* TPP proteins are

likely to be catalytically active (Supplementary Fig. S5). Among these, we found SvTPP10

and SvTPP2, the orthologues of RAMOSA3 (ZmTPP10) and ZmTPP4, respectively

(Supplementary Fig. S3), which modify inflorescence branching, sex determination and

tillering in maize (Satoh-Nagasawa et al., 2006; Claeys et al. 2019; Klein et al. 2022),

independently of their catalytic activity.

Differential expression and compartmentation of Tre6P-related transcripts

To analyse the expression pattern of the identified *S. viridis TPS*, *TPP* and *TRE* genes, we

performed real-time RT-qPCR analysis of mRNA extracted from the aerial part of five-days

old seedlings, whole leaves (pool of leaves 5 and 6), the flag leaf, whole inflorescences, and

the fourth internode of the stem (all samples harvested in the middle of the day). In addition,

we measured the levels of transcripts encoding enzymes involved in the C_4 cycle with

differential expression in MC (*SvNADP-MDH* and *SvPEPC*) and BSC (*SvNADP-ME* and

SvRbcS, which codes for the small subunit of Rubisco) from *S. viridis* and maize leaves

(Chang *et al.*, 2012; John *et al.*, 2014; Li *et al.*, 2010). We also analysed the transcripts

encoding the sucrose transporters *SvSWEET13a* and *SvSWEET13b*, which have been detected

in BSC from *S. viridis* leaves (Chen *et al.*, 2022) and are putatively regulated by Tre6P

(Oszvald *et al.*, 2018; Fichtner *et al.*, 2021). These transcripts were detected in all samples,

with the highest expression in fully developed leaves and seedlings (Supplementary Fig. S6).

SvTPS1 was mainly detected in whole leaves and the internode and, to a lesser extent,

in seedlings, the flag leaf and the inflorescence. Class II *TPS* transcripts were differentially

expressed among the analysed tissues; most of these transcripts were highly expressed in

whole leaves and seedlings but only marginally in the internode and the inflorescence. The

SvTPS3 transcript showed the highest level in seedlings and whole leaves; the levels of

SvTPS6 and *SvTPS9* were similar to those observed for *SvTPS1* in whole leaves; the *SvTPS7*

transcript displayed the highest level in the flag leaf; whereas the levels of *SvTPS7* and

SvTPS6 were comparable but lower than for *SvTPS1* in the internode (Supplementary Fig.

S6).

SvTPP transcripts also showed a differential spatial distribution, being higher in whole

leaves and internode samples. Among clade A *TPP* transcripts, *SvTPP1* showed the highest

levels in whole leaves, the inflorescence and the internode, with a maximum in the latter,

while *SvTPP8* displayed the highest levels in seedlings and the flag leaf. Regarding clade B

TPP transcripts, *SvTPP2* was consistently the one with the highest level in all the samples,

showing a maximum in the fourth internode. The highest level of the *SvTPP10* transcript (the

orthologue of *RAMOSA3*) was observed in the fourth internode. *SvTRE* was detected in all

the samples and its abundance was maximal in the internode (Supplementary Fig. S6).

To determine the compartmentation of these transcripts in mature leaves, we applied

fractionation protocols (John *et al.*, 2014; Calace *et al.*, 2021) to enrich samples in BSC and

MC, followed by RNA extraction and analysis by RT-qPCR (Fig. 2). The marker transcripts

SvNADP-MDH and *SvPEPC* were highly enriched in the MC fraction, while *SvNADP-ME*

and *SvRbcS* were mainly detected in the BSC fraction (Fig. 2A). The *SvTPS1* transcript was

almost exclusively found in the BSC fraction (Fig. 2B). It should be noted that BSC strands,

while enriched in BSC, also contain phloem (companion cells and sieve elements) and xylem

tissue. In *A. thaliana* leaves*,* the AtTPS1 protein is localized primarily in phloem parenchyma

(bundle sheath) cells and in the companion cell-sieve element complex of the phloem

(Fichtner *et al*., 2020). Thus, some of the *SvTPS1* transcript we attribute to the BSC might be

in the phloem complex. The *SvTPS3* and *SvTPS10* transcripts were also preferentially located

in the BSC fraction, while *SvTPS6* and *SvTPS8* also showed higher levels in BSC than MC,

although with a lower enrichment (Fig. 2B). Conversely, *SvTPS4* and *SvTPS7* transcripts

were more abundant in the MC fraction; indeed, *SvTPS7* was not even detected in BSC (Fig.

2B). Among the *TPP* transcripts, *SvTPP1* displayed similar levels in both fractions; *SvTPP2*,

SvTPP3, *SvTPP7* and *SvTPP9* showed a significant enrichment in the BSC fraction; and

SvTPP4 was only detected in the BSC fraction (Fig. 2C). The *SvTRE* transcript was equally

distributed in both cell types, whereas the *SvSWEET13a* and *SvSWEET13b* transcripts were

highly enriched in BSC (Fig. 2D). It is important to note that Fig. 2 does not include

transcripts that were neither detected in BSC nor MC fractions.

Tissue- and cell-specific distribution of the SvTPS1 protein

We recently performed a proteomic analysis with samples enriched in MC and BSC from *S.*

viridis leaves (Calace *et al.*, 2021). In our dataset (Supplementary Table S1 in Calace *et al.*,

2021), which contains 1334 proteins from MC and 1268 proteins from BSC, we observed that

SvTPS1 was detected in two out of four biological replicates from the BSC fraction and was

absent from the MC fraction (Fig. 3A). To further investigate the distribution of the SvTPS1

protein in different cells, we used an antiserum raised against recombinant SvTPS1 and

proteins extracted from MC and BSC enriched samples with TCA and Laemmli buffers. The

antiserum recognized two protein bands in the BSC fraction of approximately 112 and 118

kDa (Fig. 3B-C), in agreement with the predicted molecular mass of SvTPS1 (108.4 kDa).

Extraction with TCA is expected to rapidly denature any proteases in the samples (Rojas *et*

al., 2021), indicating that the double band does not result from proteolytic cleavage of the

SvTPS1 protein during the extraction procedure. We then analysed the distribution of

SvTPS1 in different tissues. The protein was detected in all the analysed samples, with

maximal signals in whole leaves and the internode (Fig. 3D), which resembles the

distribution observed for the *SvTPS1* transcript (Supplementary Fig. S6). The loading

controls and full-size images that support Figs 3B-D are presented in Supplementary Fig. S7,

which also shows that the purified anti-TPS1 antiserum cross-reacted with a protein of ~50

kDa in samples extracted with Laemmli buffer.

To confirm the identity of the protein recognized by the polyclonal antibody, we

immunoprecipitated SvTPS1 from samples enriched in BSC. A protein band corresponding to

~115 kDa was sliced from the gel and analysed by LC-MS (Supplementary Fig. S8).

Supplementary Table S6 shows that the most abundant protein from *S. viridis* present in the

sample was TPS1 (UniProt ID A0A4U6VAG9), with a coverage of 36%, 28 unique peptides

and 51 spectra. We analysed in detail the MS data for immunoprecipitated SvTPS1 but we

did not detect any post-translationally modified peptide that could explain the existence of

two immunoreactive bands, probably due to the extraction and/or immunoprecipitation

procedures employed in this work.

To confirm the location of SvTPS1 on *S. viridis* leaves we performed an

immunohistochemical assay. Cross-sections of *S. viridis* leaves were incubated with purified

anti-SvTPS1, anti-RbcL (positive control for BSC) or without primary antibody (negative

control), and then with goat anti-rabbit IgG conjugated with Cy2. As shown in Fig. 4 and

Supplementary Fig. S9, SvTPS1 was preferentially detected in BSC, although we also

observed fluorescence in some MC. This result was consistent with the signal observed for

the large subunit of Rubisco, which was mainly detected in BSC and, to a lesser extent, in

MC (Fig. 4 and Supplementary Fig. S9). The negative control did not show any differential

signal (Fig. 4 and Supplementary Fig. S9). Overall, these results indicated that the TPS1

protein is mainly located in BSC of *S. viridis* leaves.

Intercellular distribution of Tre6P

To determine the intercellular distribution of Tre6P, we prepared fractions enriched in MC or

BSC from *S. viridis* leaves by sequential filtering of frozen leaf tissue powder through a

series of nylon meshes with different pore sizes according to Stitt and Heldt (1985). We

performed two independent experiments (Experiments 1 and 2, Supplementary Tables S3 and

S4, respectively), each one consisting of four biological replicates. Supplementary Fig. S10

shows the results obtained for marker enzymes from Experiment 1. The recovery of marker

enzymes in Experiments 1 and 2 varied between 57 and 87%, depending on the enzyme and

the experiment (Supplementary Table S7). The relative distribution of these enzymatic

activities indicated that the fractionation method was successfully adapted to *S. viridis* leaves,

as the results resemble those obtained for maize leaves (Stitt and Heldt, 1985; Arrivault *et al.*,

2017; Denton *et al.*, 2017). We then measured ~30 metabolites (including Tre6P) by LC-

MS/MS, whereas pyruvate, PEP, 3PGA, DHAP, soluble sugars and starch were determined

by enzymatic assays. With these data, we calculated the proportion of each metabolite in BSC

and MC (Supplementary Tables S3-S5).

For Experiment 2, we estimated that ~83% of Tre6P was in the BSC fraction (Fig. 5

and Supplementary Table S4). As already noted, some of this Tre6P might be in the phloem

complex itself, rather than BSC. Other metabolites that were predominantly in the BSC

fraction included pyruvate, 3PGA, sucrose and starch; conversely, DHAP was mainly found

in the MC fraction, while PEP was almost equally distributed between BSC and MC (Fig. 5 and Supplementary Table S4). It should be noted that we estimated similar proportions for Tre6P, DHAP and starch in Experiment 1 (Supplementary Tables S3 and S5).

The levels of Tre6P and sucrose are tightly correlated across many treatments and conditions in *A. thaliana* leaves (Yadav *et al.*, 2014). To test whether soluble sugars were correlated with Tre6P in *S. viridis* leaves, we measured these metabolites in whole leaves harvested every four hours during the light period, from Zeitgeber time (ZT) 0 to 16 h (Supplementary Fig. S11 and Supplementary Table S8). At the whole leaf level, Tre6P was not significantly correlated with glucose, fructose or sucrose (Fig. 6). As previously mentioned, the leaves of C_4 plants have an additional complexity compared to those from C_3 plants, due to the arrangement of MC and BSC in the characteristic Kranz anatomy. Thus, we performed the same analysis using data from the second fractionation experiment (Supplementary Fig. S11 and Supplementary Table S4). As shown in Fig. 6, across cell-type-enriched fractions, Tre6P was significantly correlated with sucrose and fructose and, to a lesser extent, with glucose.

Discussion

Tre6P metabolism is critical for plant growth and development (Fichtner and Lunn, 2021) and recent reports indicate that it might be an important metabolic target to engineer crops with improved agronomic traits (Kretzschmar *et al.*, 2015; Nuccio *et al.*, 2015; Griffiths *et al.*, 2016; Gabriel *et al.*, 2021). However, most of our knowledge on this signal molecule is based on studies performed with the model plant *A. thaliana*, a eudicot that performs C_3 photosynthesis (Rojas *et al.*, 2023). Here, we studied the compartmentation of Tre6P metabolism in *S. viridis*, a model system for studying C_4 photosynthesis and associated traits (Brutnell *et al.*, 2010; Doust *et al.*, 2019).

To gain an overview of Tre6P and trehalose metabolism in *S. viridis*, we searched the

genome sequence to identify all recognisable *TPS*, *TPP* and *TRE* genes, and then performed a

phylogenetic analysis of the corresponding protein sequences. The number of genes found in

this species (10 *TPS*, 10 *TPP* and 1 *TRE*) is the same as in its domesticated relative *S. italica*

(Henry *et al.*, 2020). The topologies of the reconstructed phylogenetic trees in this work (Fig.

1 and Supplementary Fig. S3) were in good agreement with those published by Henry *et al.*

(2014) for Tre6P-related enzymes from maize, reinforcing the suitability of *S. viridis* species

as a model system for photosynthesis research in crops.

The expression of transcripts encoding enzymes involved in Tre6P metabolism in *S.*

viridis showed a differential pattern among the tissues analysed in this work. Analysis of

Tre6P-related transcripts in different tissues of *S. viridis* was not previously reported,

although other works showed their response in leaves subjected to different treatments,

including light (Henry *et al.*, 2014, 2020), salt (Henry *et al.*, 2015), high light and

temperature (Anderson *et al.*, 2021). The *SvTPS1* transcript and SvTPS1 protein abundance

was highest in internode samples (Supplementary Fig. S6 and Fig. 3D), which are rich in

vascular tissue. This is in good agreement with data from *A. thaliana*, where the TPS1 protein

was mainly detected in the vasculature of leaves, roots and reproductive tissues (Fichtner *et*

al., 2020). We could not find Tre6P-related transcripts in the RNAseq analysis performed by

Martin *et al.* (2016) using developing internodes from *S. viridis*, probably due to differences

in the sampling procedures and analysis between both works. Other authors have reported

that *A. thaliana TPS1* transcripts are relatively abundant in the protovasculature of

developing leaves and the shoot apical meristem (Wahl *et al.*, 2013), and in differentiated

phloem cells from the inflorescence stem (Shi *et al.*, 2021).

The transcripts encoding class II TPS proteins in *S. viridis* showed different levels

according to the analysed tissue, which resembles data reported for class II TPS transcripts

from *A. thaliana* (Ramon *et al.*, 2009). As previously mentioned, our analyses were

performed with samples harvested in the middle of the light period, and thus did not capture

the diurnal fluctuations previously observed in *A. thaliana* (Usadel *et al.*, 2008) and maize

(Henry *et al.*, 2014; Borba *et al.*, 2023). The *SvTPS3* transcript displayed high levels in

seedlings and whole leaves (Supplementary Fig. S6). The protein encoded by this transcript

was located in a sub-clade exclusively containing class II TPS proteins from monocots, well

separated from all other class II TPS sequences from eudicots (Fig. 1). This particular branch

contains sequences from PACMAD (*S. viridis* and maize) and BOP/BEP (wheat, rice and *B.*

distachyon) species, as well as from monocots outside the Poaceae that were included in our

analysis (*A. comosus* and *M. acuminata*). This suggests that the monocot-specific class II TPS

sub-clade arose before the divergence of the orders Poales (including the Poaceae and

Bromeliaceae) and Zingeberales (including the Musaceae). The *SvTPS6* transcript was

mainly detected in whole leaves and the fourth internode, while abundance of the *SvTPS7*

transcript was highest in the flag leaf and the fourth internode (Supplementary Fig. S6). The

proteins encoded by these transcripts were located in the same sub-clade as *A. thaliana* TPS7

(Fig. 1), whose expression was mainly detected in leaf primordia, cotyledons, roots

(concentrated in the vasculature of the mature root and evenly distributed in the meristematic

zone) and flowers (in ovules and pollen; Ramon *et al.*, 2009). Recently, Van Leene *et al.*

(2022) showed that all class II TPS proteins from *A. thaliana*, including TPS7, interact with

and potentially act as regulators of the SUCROSE-NON-FERMENTING1-RELATED

PROTEIN KINASE1 (SnRK1). Whether these proteins play the same role in *S. viridis* and

other C_4 grasses has to be further investigated.

Transcripts encoding TPP proteins from *S. viridis* also showed a particular distribution

pattern, according to the tissue and cell type under analysis, as previously reported for *A.*

thaliana (Vandesteene *et al.*, 2012). The *SvTPP2* transcript was mainly detected in the fourth

internode, with lower but significant levels in seedlings, whole leaves and the inflorescence

(Supplementary Fig. S6). The protein encoded by this transcript is the orthologue of

ZmTPP4; both were located in clade B, which also contains RAMOSA3 (ZmTPP10;

Supplementary Fig. S3). The levels of *SvTPP1* were higher in whole leaves, the

inflorescence, and the fourth internode, compared to other tissues, while levels of *SvTPP8*

were higher in seedlings, the flag leaf and the fourth internode (Supplementary Fig. S6). The

SvTPP1 and SvTPP8 protein sequences were located in clade A, which contains *A. thaliana*

TPPA, TPPF and TPPG (Supplementary Fig. S3). There is evidence of recent duplication of

genes encoding *A. thaliana* clade A TPPs, associated with whole-genome duplications within

the Brassicaceae, with a high degree of paralogue retention and evidence of

neofunctionalization (Vandesteene *et al.*, 2012). Thus, it seems feasible that the duplication

and diversification of clade A *S. viridis* TPPs observed in Supplementary Fig. S3 occurred

independently of those in *A. thaliana* (Kerbler *et al.*, 2023).

We found that several Tre6P-related transcripts were preferentially located in BSC of

S. viridis leaves (Fig. 2). To compare our results with previously published datasets, we

retrieved RNAseq data from studies performed with enriched MC and BSC fractions from *S.*

viridis and maize (John *et al.*, 2014; Denton *et al.*, 2017; Supplementary Fig. S12 and

Supplementary Table S9). Our analysis showed that the *TPS1* transcript was enriched in BSC

from *S. viridis* and maize leaves (Fig. 2B and Supplementary Fig. S12), as previously shown

for the *A. thaliana TPS1* transcript (Aubry *et al.*, 2014). A similar analysis was performed by

Benning *et al.* (2023), who used publicly available RNA deep-sequencing data to map the

distribution of sugar sensing genes in leaves of C_3 and C_4 species; among these, they found

that *TPS1* transcripts from C_4 species were mainly located in BSC. It is important to recall

that our experimental data originate from samples harvested in the middle of the day (ZT 8

h), as *TPS* and *TPP* transcripts fluctuate during the diel cycle (Usadel *et al.*, 2008, Borba *et*

al., 2023); particularly, the maize *TPS1* transcript displayed the highest level at ZT 8 h

(Borba *et al.*, 2023), indicating that this time of day is optimal for analysing the cellular

distribution of *TPS1* transcripts. Regarding the class II *TPS* transcripts, the levels of *SvTPS3*

and *SvTPS10* were significantly higher in BSC than in MC when analysed by RT-qPCR (Fig.

2B); however, these patterns were not the same in RNAseq data from *S. viridis* and maize. As

shown in Supplementary Fig. S12, *SvTPS9* and *SvTPS10* displayed the highest enrichment in

S. viridis BSC, whereas *ZmTPS11* (which clustered together with *SvTPS7*; Fig. 1) showed

high enrichment in BSC from almost all leaf segments of maize leaves. Abundance of

SvTPP1 (the orthologue of *ZmTPP6*) was similar in MC and BSC (Fig. 2C), which agrees

with RNAseq data from *S. viridis* and maize (Supplementary Fig. S12). Our RT-qPCR results

showed greater abundance of *SvTPP2, SvTPP3, SvTPP7* and *SvTPP9* in BSC than in MC

(Fig. 2C), and the same trend was observed in the RNAseq data from *S. viridis*

(Supplementary Fig. S12). Similarly, *ZmTPP1, ZmTPP2, ZmTPP7* and *ZmTPP8* (the

orthologues of *SvTPP7, SvTPP3, SvTPP2* and *SvTPP3*, respectively; Supplementary Fig. S3)

were enriched in BSC across all fractions of the maize leaf, while *ZmTPP4* and *ZmTPP9* (the

orthologues of *SvTPP2* and *SvTPP5*, respectively; Supplementary Fig. S3) were enriched in

BSC from the middle and the tip of the maize leaf (fractions 1 to 3, Supplementary Fig. S12).

The levels of the *SvTRE* transcript were similar in MC and BSC, as reported in RNAseq

studies of *S. viridis* and maize leaves (Fig. 2D and Supplementary Fig. S12).

Cell-specific expression of transcripts in MC and BSC is regulated by transcriptional

and post-transcriptional mechanisms (Reeves *et al.*, 2017). Furthermore, due to translational

and post-translational regulation, transcript abundance is not necessarily a reliable indicator

of protein abundance (Gibon *et al.*, 2004; Wang *et al.*, 2011; Reeves *et al.*, 2017). Thus, we

investigated the distribution of the SvTPS1 polypeptide in different tissues and cell types.

Immunoblotting assays detected the SvTPS1 protein in whole leaves and the fourth internode

(Fig. 3D), as previously observed in the transcript abundance analysis (Supplementary Fig.

S6). The localization of SvTPS1 in the BSC fraction of *S. viridis* leaves by immunoblotting

(Fig. 3B-C) is consistent with the proteomic data from Calace *et al.* (2021), presented in Fig.

3A.

The antiserum raised against SvTPS1 consistently recognized two immunoreactive

protein bands, particularly in BSC (Fig. 3B-C). The identity of the protein detected by the

anti-SvTPS1 was confirmed by immunoprecipitation and mass spectrometry (Supplementary

Fig. S8). Based on our results, we hypothesized that these proteoforms of SvTPS1 may result

from two splicing variants or a post-translational modification. The *S. viridis* genome

sequence has only one annotated gene model for *TPS1* (*Sevir.3G180900.1*), whereas *S. italica*

has two gene models for *TPS1* (*Seita.3G176900.1* and *Seita.3G176900.2*), in which the

alternative combination of the first two exons could produce proteins with different N-

terminus, with predicted molecular masses of 108.4 and 94.9 kDa, respectively

(Supplementary Fig. S13). The *Seita.3G176900.1* and *Seita.3G176900.2* transcripts displayed

a differential response to light (Henry *et al.*, 2020), suggesting their encoded polypeptides

could have particular biological functions. A detailed comparison of the *Sevir.3G180900* and

Seita.3G176900 DNA sequences showed eight differences, but only from the fifth intron

towards the 3´-end (Supplementary Fig. S13). Therefore, an alternative transcript could also

be produced from the *Sevir.3G180900* gene. Regarding the existence of a putative post-

translational modification, it is important to note that *A. thaliana* TPS1 is phosphorylated and

potentially sumoylated (Fichtner *et al.*, 2020; Fichtner and Lunn, 2021). The phosphorylation

sites found on the *A. thaliana* TPS1 are not conserved in SvTPS1 but the putative

sumoylation site (Lys^{902} in *A. thaliana* TPS1, Lys^{919} in SvTPS1) is well-conserved

(Supplementary Fig. S4; Fichtner *et al.*, 2020; Fichtner and Lunn, 2021). Considering the

theoretical molecular mass of SvTPS1 (108.4 kDa) and the estimated sizes of the

immunoreactive bands (112 and 118 kDa, Fig. 3B-C), the occurrence of a post-translational

modification seems more plausible, as the two putative splicing variants of SvTPS1 should

have 108.4 and 94.9 kDa, respectively (Supplementary Fig. S13).

To confirm the immunoblotting results obtained with enriched fractions, we

performed immunohistochemistry on cross-sections of *S. viridis* leaves. SvTPS1 was

preferentially detected in BSC (Fig. 4 and Supplementary Fig. S9). We did not detect any

significant signal in the vascular bundle, suggesting SvTPS1 is not present in this

compartment. This finding represents a major difference with the observation made for *A.*

thaliana TPS1, which was primarily detected in the companion cell-sieve element complex

and, to a lesser extent, in the phloem parenchyma (BSC) and xylem parenchyma cells

(Fichtner *et al.*, 2020). Overall, our results show that the only class I TPS from *S. viridis* is

predominantly located in BSC, suggesting this is the major location for Tre6P synthesis in *S.*

viridis leaves.

We adapted a method, originally developed for maize, that allows metabolite analysis

in MC and BSC enriched fractions (Stitt and Heldt, 1985). We found that the proportion of

Tre6P was higher in BSC than in MC. Similar results were observed for pyruvate, 3PGA and

starch, as expected for an NADP-ME C_4 species (Fig. 5 and Supplementary Tables S3-S5).

Similarly, ribulose 1,5-bisphosphate and other intermediates of the Calvin-Benson cycle were

almost exclusively located in BSC (Supplementary Table S5). The amount of malate was also

higher in BSC than MC. This result is counterintuitive, as the concentration gradient of

malate is expected to be the opposite; i.e., a higher concentration in MC than BSC, to drive

diffusion of malate from the MC to the BSC. However, it is important to consider that maize

leaves have a large pool of malate (up to 60% of the total malate) that is not actively involved

in the C_4 cycle, including a potentially large pool in the vacuole (Arrivault *et al*., 2017). Thus,

there still might be a concentration gradient which could move malate from the cytosol of

MC to cytosol of BSC in *S. viridis* leaves, although this has to be proven, for instance, by a

combination of $^{13}CO_2$ labelling and leaf fractionation, as previously done with maize leaves

(Arrivault *et al.*, 2017). Two intermediates of the Calvin-Benson cycle showed contrasting

distributions: 3PGA was predominantly located in the BSC, while DHAP was mainly in the

MC (Fig. 5). This is similar to the intercellular distributions of 3PGA and DHAP reported for

maize leaves (Leegood, 1985; Stitt and Heldt, 1985; Lunn and Furbank, 1997; Arrivault *et*

al., 2017), where there is intercellular shuttling of these metabolites. Pyruvate was also

predominantly located in the BSC, consistent with its production from malate

decarboxylation, via NADP-ME, in the BSCs, and diffusion into the MCs for conversion to

PEP via PPDK. PEP was evenly distributed between MC and BSC (Fig. 5 and Supplementary

Table S5). These distributions differ from those reported for maize, where PEP showed a

preferential increase in the BSC and pyruvate was located mainly in the MC (Arrivault *et al.*,

2017). Such differences could be explained by i) the existence of metabolic gradients along

the leaf, or ii) the presence of metabolite pools that are not actively involved in C_4

photosynthesis, for example, if located in the vacuole, which could mask metabolite gradients

between the cytosol of MC and that of BSC (Szecowka *et al.*, 2013; Arrivault *et al.*, 2017;

Denton *et al.*, 2017).

A limitation of our study is that samples for metabolite analysis were collected from

the middle section of the leaf blade at one defined time-point in the middle of the light period,

so we could not capture metabolic dynamics. It would be interesting to perform labelling

studies followed by cellular enrichment to obtain a detailed scenario of the metabolic

fluctuations that occur in *S. viridis* leaves. Furthermore, overall gradients between the MC

and BSC might not reflect the gradients between the cytosol of these two cell types, if there

were active transport of metabolites across the plastid envelope or the pH gradient across the

envelope membrane influences their distribution between the chloroplast stroma and the

cytosol, as is known to be the case for 3PGA (Fliege *et al*., 1978; Stitt *et al.*, 1980). The

relation between the overall intercellular metabolite gradient and the gradient between the

cytosols of the BSC and MC will also be affected by the relative contribution of chloroplasts

and cytosol to cytoplasmic volume. Overall, our study indicates that the metabolite gradients

reported in maize may not be generalizable and underlines the need for studies of metabolite

gradients in other C_4 species. Just as different biochemical solutions have arisen in different

species to concentrate CO_2 into the BSC (Furbank and Kelly, 2021), different strategies may

also have been followed to generate the intercellular gradients in the cytosol that are needed

to drive metabolite movement between the BSC and MC. Overall, our fractionation data

provide the first insights into the distributions of key photosynthetic intermediates in *S.*

viridis. These are consistent with the photosynthetic fluxes and intercellular metabolite

movements that we would expect to see in an NADP-ME-type C_4 species, demonstrating the

reliability of this experimental approach.

776   The main aim of the cell fractionation experiments in our study was to determine the

cellular localization of Tre6P. Our transcript and protein analyses showed that the SvTPS1

enzyme is predominantly located in the BSC and associated vascular tissues, and essentially

absent from MC (Figs 2-4), indicating that Tre6P is synthesized in BSC but not in MC. We

found that approximately 83% of Tre6P was present in the BSC fraction (Fig. 5) and,

considering the previously mentioned limitations, this value could be even higher. It is worth

noting that some of the Tre6P in the BSC fraction may be in the phloem complex because,

although the cell separation method allows the sample to be enriched with BSC strands, this

enrichment is partial and may bring with it remains from other cells types. This would make

sense considering that, in *A. thaliana*, Tre6P metabolism has been reported to take place in

both phloem parenchyma (equivalent to the BSC) and in the companion cell-sieve element

complex of the phloem (Fichtner *et al.*, 2020). As previously mentioned, sucrose synthesis

seems to occur in MC of *S. viridis* leaves; however, the amount of sucrose was higher in the

BSC fraction than in the MC fraction (Supplementary Table S5). This result is puzzling, as

sucrose levels are expected to be higher in the MC than the BSC, to drive its diffusion to the

BSC. We thus speculate that i) sucrose is sequestered either in some compartment other than

the cytosol in the BSC or other cell types that are enriched in the BSC fractions; or ii) that a

large part of the sucrose in the leaf is in the phloem.

Lunn *et al.* (2006) showed there is a significant positive correlation between Tre6P

and sucrose levels in samples from *A. thaliana* seedlings and rosettes. The correlation

between Tre6P and sucrose persisted even in transgenic lines expressing the *E. coli* TPS

(otsA) and TPP (otsB) proteins under the control of the *35S* promoter, although sucrose levels

were markedly lower or higher, respectively, which led to the development of the Tre6P-

sucrose nexus model (Yadav *et al.*, 2014). Tre6P and sucrose levels fluctuated during the

light period in *S. viridis* leaves (Supplementary Fig. S11) but, unlike *A. thaliana*, we did not

observe a significant correlation at the whole-leaf level (Fig. 6). Conversely, we found a

significant correlation between Tre6P and sucrose across leaf fractions containing different

proportions of MC and BSC (Fig. 6). The Tre6P-sucrose nexus model postulates that Tre6P is

a signal and a negative regulator of sucrose levels (Yadav *et al.*, 2014). According to this

model, if sucrose levels rise too high, Tre6P rises in parallel and triggers mechanisms to

lower the level of sucrose, either by reducing sucrose synthesis or increasing its degradation

or transport. Regarding the first alternative, research performed with ethanol-inducible *A.*

thaliana transgenic lines showed that artificially increased Tre6P levels decreased the amount

of sucrose by diverting newly fixed carbon to organic and amino acids during the day,

through the post-translational activation of PEPC and nitrate reductase (Figueroa *et al.*,

2016). As previously mentioned, the initial CO_2 fixation step in C_4 plants is catalysed by

PEPC, which is located in MC. Our results strongly suggest that Tre6P is mainly produced

and accumulated in BSC of *S. viridis* leaves. Thus, if the Tre6P-PEPC connection is

operational in C_4 species, this would imply that Tre6P has to diffuse from BSC to MC;

indeed, a small amount of Tre6P was actually found in MC (Fig. 5). The asymmetrical

distribution of Tre6P suggests that it could be sensing sucrose levels in BSC and then act as a

signal that moves into the MC to potentially modulate PEPC activity and/or sucrose

synthesis, which is also located in MC. The molecular mechanism underlying the putative

transport of Tre6P from BSC to MC deserves to be further studied but is beyond the scope of

this work.

Another possibility is that Tre6P regulate sucrose levels by controlling phloem

loading. TPS1 has been found in the vasculature of *A. thaliana* leaves; particularly, in the

phloem parenchyma and the companion cell/sieve element complex (Fichtner *et al.*, 2020).

Tre6P has been shown to modulate the levels of *ZmSWEET13a,c* and *AtSWEET11-14*

transcripts in maize and *A. thaliana*, respectively, which encode sucrose transporters involved

in phloem loading (Oszvald *et al.*, 2018; Fichtner *et al.*, 2021). Single-cell transcriptomics in

A. thaliana, rice and maize showed that *SWEET* transcripts were preferentially detected in

BSC (Bezrutczyk *et al.*, 2021; Shi *et al.,* 2021; Hua *et al.,* 2021; Tao *et al.*, 2022). Moreover,

the *ZmSWEET13a-c* transcripts and the SWEET13a protein were specifically detected in the

abaxial BSC of major and intermediate veins from maize leaves (Bezrutczyk *et al.*, 2021),

while both SWEET13a and SWEET13b proteins were detected in the abaxial BSC of

intermediate veins from *S. viridis* leaves (Chen *et al.*, 2022). Based on these data, it has been

proposed that sucrose moves from adaxial to abaxial cells via plasmodesmata and is then

exported by SWEET13 to the apoplast, from where it is taken up by transporters from the

SUT/SUC (sucrose-H^+ symporter) family located in the companion cells (Bezrutczyk *et al.*,

2021; Chen *et al.*, 2022). The maize *zmsweet13a-c* triple mutant is impaired in sugar phloem

loading, thus producing drastic effects on growth (Bezrutczyk *et al*., 2018). In *S. viridis*, such

mutants have not yet been identified, but there is compelling evidence that it could be an

apoplastic phloem-loading species, and thus SvSWEET13a-b might function in phloem

loading (Chen *et al.*, 2022). Based on this evidence, and the fact that Tre6P metabolism is

mostly located in BSC, we speculate that Tre6P modulates sucrose export in leaves of C_4

plants by controlling the expression, and thereby maximal activity, of the SWEET13

transporters, although this hypothesis remains to be tested. If confirmed, this would implicate

Tre6P in the reconfiguring of phloem loading that is a common feature of C_4 monocots, and

potentially an essential element in the evolution of Kranz anatomy and C_4 photosynthesis

(Furbank and Kelly, 2021; Rojas *et al.*, 2023).

To conclude, our results indicate that Tre6P is synthesized and mainly located in the

BSC of *S. viridis*, which play a central role in phloem loading in C_4 species and represent a

strategic site at the interface between source and sink tissues. This localization would allow

Tre6P to monitor the levels of sucrose in BSC and, via diffusion into the MC, potentially

regulate CO_2 fixation and sucrose biosynthesis as well as sucrose export from the leaves

(Rojas *et al.*, 2023).

Supplementary Data

Supplementary File S1. Optimised DNA sequence for the recombinant expression of

SvTPS1 in *E. coli* cells.

Supplementary Figure S1. Analysis of MC and BSC fractions for transcript and protein

determinations.

Supplementary Figure S2. C_t values for the reference transcript *SvKIN* in different tissues

and cell types from *S. viridis*.

Supplementary Figure S3. Phylogenetic relationships of *Setaria viridis* TPP proteins.

Supplementary Figure S4. Multiple sequence alignment of *S. viridis* TPS proteins.

Supplementary Figure S5. Multiple sequence alignment of *S. viridis* TPP proteins.

Supplementary Figure S6. Relative abundance of transcripts encoding Tre6P-related

enzymes in different *S. viridis* tissues.

Supplementary Figure S7. Immunoblotting of SvTPS1 in different tissues and cell types of

S. viridis.

Supplementary Figure S8. Analysis by mass spectrometry of immunoprecipitated SvTPS1.

Supplementary Figure S9. Immunolocalization of SvTPS1 in cross-sections of *S. viridis*

leaves.

Supplementary Figure S10. Activities of marker enzymes in different fractions from *S.*

viridis leaves.

Supplementary Figure S11. Analysis of metabolites from *S. viridis* leaves.

Supplementary Figure S12. Analysis of transcripts encoding Tre6P-related enzymes in BSC

and MC from *S. viridis* and maize leaves.

Supplementary Figure S13. Gene models for the class I TPS from *S. viridis* and *S. italica*.

Supplementary Table S1. TPS and TPP protein sequences used for reconstruction of

phylogenetic trees.

Supplementary Table S2. Sequences of the primers used to measure transcript levels by

qPCR.

Supplementary Table S3. Metabolite and enzyme data and calculations (Experiment 1).

Supplementary Table S4. Metabolite and enzyme data and calculations (Experiment 2).

Supplementary Table S5. Calculation of metabolite fractions in MC and BSC.

Supplementary Table S6. Analysis of immunoprecipitated SvTPS1 by mass spectrometry.

Supplementary Table S7. Recovery of marker enzymes in the fractionation experiments.

Supplementary Table S8. Metabolite data from *S. viridis* leaves during a day time-course.

Supplementary Table S9. Relative transcript abundance in MC and BSC from *S. viridis* and

maize.

Acknowledgements

TT, BER and PC are fellows from CONICET. CS is a fellow from Agencia I+D+i. LEL,

MSaigo and CMF are researchers from CONICET. CMF thanks NME, CF and MF for their

everyday support. We thank to the Mass Spectrometry Unit from IBR (UNR-CONICET) for

the analysis of immunoprecipitated SvTPS1; Agustín Arce for help with statistical analysis;

Andrés Dekanty for providing the Cy2-conjugated goat anti-rabbit IgG; and Sergio Gonzalez

for technical assistance with the confocal microscope.

Author contributions

Conceptualization: MStitt, JEL and CMF; Investigation: TT, BER, RF, CS, LEL, JVC, PC,

SA and CMF; Formal analysis: all authors; Writing - Original Draft: TT, BER and CMF;

Writing - Review & Editing: all authors.

Conflict of interest

No conflict of interest declared.

Funding

This work was supported by Agencia Nacional de Promoción de la Investigación, el

Desarrollo Tecnológico y la Innovación [PICT-2018-00865 and PICT-2020-00260];

Universidad Nacional del Litoral [CAI+D 2020]; Consejo Nacional de Investigaciones

Científicas y Técnicas [PIP 2021-2023]; Agencia Santafesina de Ciencia, Tecnología e

Innovación [PEIC I+D 2021-021]; and the Max Planck Society [Partner Group for Plant

Biochemistry].

Data availability

915 All data supporting the findings of this study are available within the paper and within its

916 supplementary data published online. The raw MS data resulting from the analysis of the

917 immunoprecipitated TPS1 were deposited to the PRIDE repository with the dataset identifier

918 PXD057139.

References

Anderson CM, Mattoon EM, Zhang N, *et al.* 2021. High light and temperature reduce photosynthetic efficiency through different mechanisms in the C_4 model *Setaria viridis*. Communications Biology **4**, 1092.

Arrivault S, Guenther M, Ivakov A, Feil R, Vosloh D, van Dongen JT, Sulpice R, Stitt M. 2009. Use of reverse-phase liquid chromatography, linked to tandem mass spectrometry, to profile the Calvin cycle and other metabolic intermediates in Arabidopsis rosettes at different carbon dioxide concentrations. Plant Journal **59**, 826–839.

Arrivault S, Obata T, Szecówka M, Mengin V, Guenther M, Hoehne, M, Fernie AR, Stitt M. 2017. Metabolite pools and carbon flow during C_4 photosynthesis in maize: $^{13}CO_2$ labeling kinetics and cell type fractionation. Journal of Experimental Botany **68**, 283–298.

Ashton AR, Burnell JN, Furbank RT, Jenkins CLD, Hatch MD. 1990. Enzymes of C_4 photosynthesis. In: Lea PJ, ed. Enzymes of primary metabolism. Methods in plant biochemistry, Volume 3. Academic Press, London, 39-72.

Aubry S, Smith-Unna RD, Boursnell CM, Kopriva S, Hibberd JM. 2014. Transcript residency on ribosomes reveals a key role for the *Arabidopsis thaliana* bundle sheath in sulfur and glucosinolate metabolism. Plant Journal **78**, 659–673.

Avidan O, Martins MCM, Feil R, Lohse M, Giorgi FM, Schlereth A, Lunn JE, Stitt M. 2024. Direct and indirect responses of the Arabidopsis transcriptome to an induced increase in trehalose 6-phosphate. Plant Physiology, in press. https://doi.org/10.1093/plphys/kiae196

Avonce N, Mendoza-Vargas A, Morett E, Iturriaga G. 2006. Insights on the evolution of trehalose biosynthesis. BMC Evolutionary Biology **6**, 1-15.

Barraza A, Contreras-Cubas C, Estrada-Navarrete G, Reyes JL, Juárez-Verdayes MA, Avonce N, Quinto C, Díaz-Camino C, Sanchez F. 2016. The class II trehalose 6-phosphate synthase gene *PvTPS9* modulates trehalose metabolism in *Phaseolus vulgaris* nodules.

Frontiers in Plant Science **7**, 1589.

Benning UF, Chen L, Watson-Lazowski A, Henry C, Furbank RT, Ghannoum O. 2023. Spatial expression patterns of genes encoding sugar sensors in leaves of C_4 and C_3 grasses. Annals of Botany **131**, 985-1000.

Bezrutczyk M, Hartwig T, Horschman M, Char SN, Yang J, Yang B, Frommer WB, Sosso D. 2018. Impaired phloem loading in *zmsweet13a,b,c* sucrose transporter triple knock-out mutants in *Zea mays*. New Phytologist **218**, 594-603.

Bezrutczyk M, Zöllner NR, Kruse CPS, Hartwig T, Lautwein T, Köhrer K, Frommer WB, Kim JY. 2021. Evidence for phloem loading via the abaxial bundle sheath cells in maize leaves. Plant Cell **33**, 531–547.

Bledsoe SW, Henry C, Griffiths CA, Paul MJ, Feil R, Lunn JE, Stitt M, Lagrimini LM. 2017. The role of Tre6P and SnRK1 in maize early kernel development and events leading to stress-induced kernel abortion. BMC Plant Biology **17**, 1-17.

Borba AR, Reyna-Llorens I, Dickinson PJ, *et al*. 2023. Compartmentation of photosynthesis gene expression in C_4 maize depends on time of day. Plant Physiology **193**, 2306–2320.

Bradford MM. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Analytical Biochemistry **72**, 248–254.

Brutnell TP, Wang L, Swartwood K, Goldschmidt A, Jackson D, Zhu XG, Kellogg E, Van Eck J. 2010. *Setaria viridis*: a model for C_4 photosynthesis. The Plant Cell **22**, 2537–2544.

Cabello JV, Chan RL. 2019. Arabidopsis and sunflower plants with increased xylem area show enhanced seed yield. Plant Journal **99**, 717-732.

Cabib E, Leloir LF. 1958. The biosynthesis of trehalose phosphate. Journal of Biological

Chemistry **231**, 259–275.

Calace P, Tonetti T, Margarit E, Figueroa CM, Lobertti C, Andreo CS, Gerrard Wheeler MC, Saigo M. 2021. The C_4 cycle and beyond: diverse metabolic adaptations accompany dual-cell photosynthetic functions in Setaria. Journal of Experimental Botany **72**, 7876–7890.

Chang YM, Wen-Yu L, Chun-Chieh AS, *et al.* 2012. Characterizing regulatory and functional differentiation between maize mesophyll and bundle sheath cells by transcriptomic analysis. Plant Physiology **160**, 165–177.

Chen L, Ganguly DR, Shafik SH, Ermakova M, Pogson BJ, Grof CP, Sharwood RE, Furbank RT. 2022. Elucidating the role of SWEET13 in phloem loading of the C_4 grass *Setaria viridis*. Plant Journal **109**, 615-632.

Claeys H, Vi SL, Xu X, *et al.* 2019. Control of meristem determinacy by trehalose 6-phosphate phosphatases is uncoupled from enzymatic activity. Nature Plants **5**, 352–357.

Czedik-Eysenberg A, Arrivault S, Lohse M, *et al.* 2016. The interplay between carbon availability and growth in different zones of the growing maize leaf. Plant Physiology **172**, 943-967.

de Haro LA, Arellano SM, Novák O, Feil R, Dumón AD, Mattio MF, Tarkowská D, Llauger G, Strnad M, Lunn JE, Pearce S, Figueroa CM, Del Vas M. 2019. Mal de Río Cuarto virus infection causes hormone imbalance and sugar accumulation in wheat leaves. BMC Plant Biology **19**, 112.

Delorge I, Figueroa CM, Feil R, Lunn JE, Van Dijck P. 2015. Trehalose-6-phosphate synthase 1 is not the only active TPS in *Arabidopsis thaliana*. Biochemical Journal **466**, 283–290.

Denton AK, Maß J, Külahoglu C, Lercher MJ, Bräutigam A, Weber APM. 2017. Freeze-quenched maize mesophyll and bundle sheath separation uncovers bias in previous

tissue-specific RNA-Seq data. Journal of Experimental Botany **68**, 147–160.

Devi K, Prathima PT, Gomathi R, Manimekalai R, Lakshmi K, Selvi A. 2019. Gene expression profiling in sugarcane genotypes during drought stress and rehydration. Sugar Tech **21**, 717–733.

Dong Z, Xiao Y, Govindarajulu R, Feil R, Siddoway ML, Nielsen T, Lunn JE, Hawkins J, Whipple C, Chuck G. 2019. The regulatory landscape of a core maize domestication module controlling bud dormancy and growth repression. Nature Communications **10**, e3810.

dos Anjos L, Pandey PK, Moraes TA, Feil R, Lunn JE, Stitt M. 2018. Feedback regulation by trehalose 6-phosphate slows down starch mobilization below the rate that would exhaust starch reserves at dawn in Arabidopsis leaves. Plant Direct **2**, e00078.

Doust AN, Brutnell TP, Upadhyaya HD, Van Eck J. 2019. Editorial: setaria as a model genetic system to accelerate yield increases in cereals, forage crops, and bioenergy grasses. Frontiers in Plant Science **10**, 1211.

Du L, Li S, Ding L, Cheng X, Kang Z, Mao H. 2022. Genome-wide analysis of trehalose-6-phosphate phosphatases (TPP) gene family in wheat indicates their roles in plant development and stress response. BMC Plant Biology **22**, 120.

Eastmond PJ, van Dijken AJ, Spielman M, Kerr A, Tissier AF, Dickinson HG, Jones JDG, Smeekens SC, Graham IA. 2002. Trehalose-6-phosphate synthase 1, which catalyses the first step in trehalose synthesis, is essential for arabidopsis embryo maturation. Plant Journal **29**, 225–235.

Fang L. 2012. Antibody purification from western blotting. Bio-protocol **2**, e133.

Fichtner F, Barbier FF, Annunziata MG, Feil R, Olas JJ, Mueller-Roeber B, Stitt M, Beveridge CA, Lunn JE. 2021. Regulation of shoot branching in Arabidopsis by trehalose 6-phosphate. New Phytologist **229**, 2135–2151.

Fichtner F, Lunn JE. 2021. The role of trehalose-6-phosphate (Tre6P) in plant metabolism

and development. Annual Review of Plant Biology **72**, 737–760.

Fichtner F, Olas JJ, Feil R, Watanabe M, Krause U, Hoefgen R, Stitt M, Lunn JE. 2020. Functional features of TREHALOSE-6-PHOSPHATE SYNTHASE1, an essential enzyme in arabidopsis. Plant Cell **32**, 1949–1972.

Figueroa CM, Feil R, Ishihara H, *et al.* 2016. Trehalose 6-phosphate coordinates organic and amino acid metabolism with carbon availability. Plant Journal **85**, 410–423.

Figueroa CM, Lunn JE. 2016. A tale of two sugars: trehalose-6-phosphate and sucrose. Plant Physiology **172**, 7–27.

Fliege R, Flügge UI, Werdan K, Heldt HW. 1978. Specific transport of inorganic phosphate, 3-phosphoglycerate and triosephosphates across the inner membrane of the envelope in spinach chloroplasts. Biochimica et Biophysica Acta **502**, 232-247.

Furbank RT and Kelly S. 2021. Finding the C_4 sweet spot: cellular compartmentation of carbohydrate metabolism in C_4 photosynthesis. Journal of Experimental Botany **72**, 6018-6026.

Gabriel C, Fernhout JJ, Fichtner F, Feil R, Lunn JE, Kossmann J, Lloyd JR, van der Vyver C. 2021. Genetic manipulation of trehalose-6-phosphate synthase results in changes in the soluble sugar profile in transgenic sugarcane stems. Plant Direct **5**, 1–18.

Gascuel O. 1997. BIONJ: An improved version of the NJ algorithm based on a simple model of sequence data. Molecular Biology and Evolution **14**, 685–695.

Gibon Y, Blaesing OE, Hannemann J, Carillo P, Höhne M, Hendriks JHM, Palacios N, Cross J, Selbig J, Stitt M. 2004. A robot-based platform to measure multiple enzyme activities in Arabidopsis using a set of cycling assays: comparison of changes of enzyme activities and transcript levels during diurnal cycles and in prolonged darkness. The Plant Cell **16**, 3304–3325.

Gibson RP, Turkenburg JP, Charnock SJ, Lloyd R, Davies GJ. 2002. Insights into

trehalose synthesis provided by the structure of the retaining glucosyltransferase OtsA. Chemistry and Biology **9**, 1337–1346.

Gomez LD, Baud S, Gilday A, Li Y, Graham IA. 2006. Delayed embryo development in the Arabidopsis trehalose-6-phosphate synthase 1 mutant is associated with altered cell wall structure, decreased cell division and starch accumulation. Plant Journal **46**, 69–84.

Gomez LD, Gilday A, Feil R, Lunn JE, Graham IA. 2010. AtTPS1-mediated trehalose 6-phosphate synthesis is essential for embryogenic and vegetative growth and responsiveness to ABA in germinating seeds and stomatal guard cells. Plant Journal **64**, 1–13.

Gouy M, Guindon S, Gascuel O. 2010. SeaView version 4: a multiplatform graphical user interface for sequence alignment and phylogenetic tree building. Molecular Biology and Evolution **27**, 221-224.

Griffiths CA, Sagar R, Geng Y, *et al.* 2016. Chemical intervention in plant sugar signalling increases yield and resilience. Nature **540**, 574–578.

Griffiths CA, Xue X, Miret JA, Salvagiotti F, Acevedo-Siaca LG, Gimeno J, Reynolds MP, Hassall KL, Halsey K, Puranik S, Oszvald M, Kurup S, Davis BG, Paul MJ. 2025. Membrane-permeable trehalose 6-phosphate precursor spray increases wheat yields in field trials. Nature Biotechnology, in press. doi: 10.1038/s41587-025-02611-1.

Hartman MD, Rojas BE, Ferrero DML, Leyva A, Durán R, Iglesias AA, Figueroa CM. 2023. Phosphorylation of aldose-6-phosphate reductase from *Prunus persica* leaves. Plant Physiology and Biochemistry **194**, 461-469.

Hendriks JH, Kolbe A, Gibon Y, Stitt M, Geigenberger P. 2003. ADP-glucose pyrophosphorylase is activated by posttranslational redox-modification in response to light and to sugars in leaves of arabidopsis and other plant species. Plant Physiology **133**, 838–849.

Henry C, Bledsoe SW, Griffiths CA, Kollman A, Paul MJ, Sakr S, Lagrimini LM. 2015.

Differential role for trehalose metabolism in salt-stressed maize. Plant Physiology **169**, 1072–1089.

Henry C, Bledsoe SW, Siekman A, Kollman A, Waters BM, Feil R, Stitt M, Lagrimini LM. 2014. The trehalose pathway in maize: conservation and gene regulation in response to the diurnal cycle and extended darkness. Journal of Experimental Botany **65**, 5959–5973.

Henry C, Watson-Lazowski A, Oszvald M, Griffiths C, Paul MJ, Furbank RT, Ghannoum O. 2020. Sugar sensing responses to low and high light in leaves of the C_4 model grass *Setaria viridis*. Journal of Experimental Botany **71**, 1039–1052.

Hua L, Stevenson SR, Reyna-Llorens I, Xiong H, Kopriva S, Hibberd JM. 2021. The bundle sheath of rice is conditioned to play an active role in water transport as well as sulfur assimilation and jasmonic acid synthesis. Plant Journal **107**, 268-286.

Ishihara H, Alseekh S, Feil R, *et al*. 2022. Rising rates of starch degradation during daytime and trehalose 6-phosphate optimize carbon availability. Plant Physiology **189**, 1976-2000.

Jenö P, Horst M. 2002. Electroelution of proteins from polyacrylamide gels. The Protein Protocols Handbook. Springer, 207–214.

John CR, Smith-Unna RD, Woodfield H, Covshoff S, Hibberd JM. 2014. Evolutionary convergence of cell-specific gene expression in independent lineages of C_4 grasses. Plant Physiology **165**, 62–75.

Kerbler SM, Armijos-Jaramillo V, Lunn JE, Vicente R. 2023. The trehalose 6-phosphate phosphatase family in plants. Physiologia Plantarum **175**, e14096.

Klein H, Gallagher J, Demesa-Arevalo E, *et al*. 2022. Recruitment of an ancient branching program to suppress carpel development in maize flowers. Proceedings of the National Academy of Sciences USA **119**, e2115871119.

Kretzschmar T, Pelayo MAF, Trijatmiko KR, *et al.* 2015. A trehalose-6-phosphate phosphatase enhances anaerobic germination tolerance in rice. Nature Plants **1**, 1-5.

Leegood RC. 1985. The intercellular compartmentation of metabolites in leaves of *Zea mays* L. Planta **164**,163 171.

Leegood RC. 1990. Enzymes of the Calvin cycle. In: Lea PJ, ed. Enzymes of primary metabolism. Methods in plant biochemistry, Volume 3. Academic Press, London, 15–37.

Leyman B, Van Dijck P, Thevelein JM. 2001. An unexpected plethora of trehalose biosynthesis genes in *Arabidopsis thaliana*. Trends in Plant Science **6**, 510-513.

Li P, Ponnala L, Gandotra N, *et al*. 2010. The developmental dynamics of the maize leaf transcriptome. Nature Genetics **42**, 1060-1067.

Link AJ, Labaer J. 2009. In-gel trypsin digest of gel-fractionated proteins. Cold Spring Harbor Protocols **2009**, pdb.prot5110.

Lunn JE and Furbank RT. 1997. Localisation of sucrose-phosphate synthase and starch in leaves of C_4 plants. Planta **202**, 106–111.

Lunn JE and Furbank RT. 1999. Sucrose biosynthesis in C_4 plants. New Phytologist **143**, 221–237.

Lunn JE, Delorge I, Figueroa CM, Van Dijck P, Stitt M. 2014. Trehalose metabolism in plants. Plant Journal **79**, 544–567.

Lunn JE, Feil R, Hendriks JH, Gibon Y, Morcuende R, Osuna D, Scheible WR, Carillo P, Hajirezaei MR, Stitt M. 2006. Sugar-induced increases in trehalose 6-phosphate are correlated with redox activation of ADPglucose pyrophosphorylase and higher rates of starch synthesis in *Arabidopsis thaliana*. Biochemical Journal **397**, 139–148.

Lunn JE. 2007. Gene families and evolution of trehalose metabolism in plants. Functional Plant Biology **34**, 550–563.

Martin AP, Palmer WM, Brown C, Abel C, Lunn JE, Furbank RT, Grof CP. 2016. A developing *Setaria viridis* internode: an experimental system for the study of biomass generation in a C_4 model species. Biotechnology for Biofuels **9**, 1–12.

Martins MC, Hejazi M, Fettke J, *et al.* 2013. Feedback inhibition of starch degradation in Arabidopsis leaves mediated by trehalose 6-phosphate. Plant Physiology **163**, 1142–1163.

Martins PK, Mafra V, de Souza WR, Ribeiro AP, Vinecky F, Basso MF, da Cunha BADB, Kobayashi AK, Molinari HBC. 2016. Selection of reliable reference genes for RT-qPCR analysis during developmental stages and abiotic stress in *Setaria viridis*. Scientific Reports **6**, 1-10.

Nuccio ML, Wu J, Mowers R, *et al.* 2015. Expression of trehalose-6-phosphate phosphatase in maize ears improves yield in well-watered and drought conditions. Nature Biotechnology **33**, 862–869.

Oszvald M, Primavesi LF, Griffiths CA, Cohn J, Basu SS, Nuccio ML, Paul MJ. 2018. Trehalose 6-phosphate regulates photosynthesis and assimilate partitioning in reproductive tissue. Plant Physiology **176**, 2623–2638.

Paul MJ, Gonzalez-Uriarte A, Griffiths CA, Hassani-Pak K. 2018. The role of trehalose 6-phosphate in crop yield and resilience. Plant Physiology **177**, 12-23.

Ponnu J, Schlereth A, Zacharaki V, Dzialo MA, Abel C, Feil R, Schmid M, Wahl V. 2020. The trehalose 6-phosphate pathway impacts vegetative phase change in *Arabidopsis thaliana*. Plant Journal **104**, 768-780.

Ramon M, De Smet I, Vandesteene L, Naudts M, Leyman B, Van Dijck P, Rolland F, Beeckman T, Thevelein JM. 2009. Extensive expression regulation and lack of heterologous enzymatic activity of the Class II trehalose metabolism proteins from *Arabidopsis thaliana*. Plant, Cell and Environment **32**, 1015-1032.

Reeves G, Grangé-Guermente MJ, Hibberd JM. 2017. Regulatory gateways for cell-specific gene expression in C_4 leaves with Kranz anatomy. Journal of Experimental Botany **68**, 107–116.

Rojas BE, Hartman MD, Figueroa CM, Iglesias AA. 2021. Proteolytic cleavage of

Arabidopsis thaliana phospho*enol*pyruvate carboxykinase-1 modifies its allosteric regulation. Journal of Experimental Botany **72**, 2514-2524.

Rojas BE, Tonetti T, Figueroa CM. 2023. Trehalose 6-phosphate metabolism in C$_4$ species. Current Opinion in Plant Biology **72**, 102347.

Satoh-Nagasawa N, Nagasawa N, Malcomber S, Sakai H, Jackson D. 2006. A trehalose metabolic enzyme controls inflorescence architecture in maize. Nature **441**, 227–230.

Sauer M, Friml J. 2010. Immunolocalization of proteins in plants. In: Hennig L, Köhler C, eds. Plant Developmental Biology. Methods in Molecular Biology, Volume 655. Humana Press, Totowa, 253-263.

Schmittgen TD, Livak KJ. 2008. Analyzing real-time PCR data by the comparative CT method. Nature Protocols **3**, 1101–1108.

Shi D, Jouannet V, Agustí J, Kaul V, Levitsky V, Sanchez P, Mironova VV, Greb T. 2021. Tissue-specific transcriptome profiling of the Arabidopsis inflorescence stem reveals local cellular signatures. Plant Cell **33**, 200-223.

Stitt M, Heldt HW. 1985. Control of photosynthetic sucrose synthesis by fructose-2,6-bisphosphate□: Intercellular metabolite distribution and properties of the cytosolic fructosebisphosphatase in leaves of *Zea mays* L. Planta **164**, 179–188.

Stitt M, Lilley RM, Gerhardt R, Heldt HW. 1989. Metabolite levels in specific cells and subcellular compartments of plant leaves. Methods in Enzymology **174**, 518–552.

Stitt M, Wirtz W, and Heldt HW. 1980. Metabolite levels in the chloroplast and extrachloroplast compartments of spinach protoplasts. Biochimica et Biophysica Acta **593**, 85-102.

Szecowka M, Heise R, Tohge T, *et al.* 2013. Metabolic fluxes in an illuminated Arabidopsis rosette. The Plant Cell **25**, 694–714.

Tao S, Liu P, Shi Y, *et al*. 2022. Single-cell transcriptome and network analyses unveil key

transcription factors regulating mesophyll cell development in maize. Genes **13**, 374.

Usadel B, Bläsing OE, Gibon Y, Retzlaff K, Höhne M, Günther M, Stitt M. 2008. Global transcript levels respond to small changes of the carbon status during progressive exhaustion of carbohydrates in Arabidopsis rosettes. Plant Physiology **146**, 1834-1861.

van Dijken AJ, Schluepmann H, Smeekens SC. 2004. Arabidopsis trehalose-6-phosphate synthase 1 is essential for normal vegetative growth and transition to flowering. Plant Physiology **135**, 969–977.

Van Eck J, Swartwood K. 2015. *Setaria viridis*. In: Wang K, ed. Agrobacterium Protocols. Methods in Molecular Biology, Volume 1223. Springer, New York, 57–67.

Van Leene J, Eeckhout D, Gadeyne A, *et al*. 2022. Mapping of the plant SnRK1 kinase signalling network reveals a key regulatory role for the class II T6P synthase-like proteins. Nature Plants **8**, 1245–1261.

Vandesteene L, Lopez-Galvis L, Vanneste K, *et al.* 2012. Expansive evolution of the trehalose-6-phosphate phosphatase gene family in Arabidopsis. Plant Physiology **160**, 884–896.

Vandesteene L, Ramon M, Le Roy K, Van Dijck P, Rolland F. 2010. A single active trehalose-6-P synthase (TPS) and a family of putative regulatory TPS-like proteins in Arabidopsis. Molecular Plant **3**, 406–419.

Wahl V, Ponnu J, Schlereth A, Arrivault S, Langenecker T, Franke A, Feil R, Lunn JE, Stitt M, Schmid M. 2013. Regulation of flowering by trehalose-6-phosphate signaling in *Arabidopsis thaliana*. Science **339**, 704–707.

Wang L, Peterson R, Brutnell T. 2011. Regulatory mechanisms underlying C_4 photosynthesis. New Phytologist **190**, 9–20.

Wang S, Ouyang K, Wang K. 2019. Genome-wide identification, evolution, and expression analysis of *TPS* and *TPP* gene families in *Brachypodium distachyon*. Plants **8**, 362.

Yadav UP, Ivakov A, Feil R, *et al.* 2014. The sucrose-trehalose 6-phosphate (Tre6P) nexus: specificity and mechanisms of sucrose signalling by Tre6P. Journal of Experimental Botany **65**, 1051–1068.

Yang HL, Liu YJ, Wang CL, Zeng QY. 2012. Molecular evolution of trehalose-6-phosphate synthase (TPS) gene family in populus, arabidopsis and rice. PLoS ONE **7**, e42438.

Figure legends

Figure 1. Phylogenetic relationships of *Setaria viridis* TPS proteins. TPS protein sequences were collated from *Setaria viridis* (Sv) and other selected species: Ac, *Ananas comosus*; At, *Arabidopsis thaliana*; Bd, *Brachypodium dystachion*; Mc, *Musa acuminata*; Os, *Oryza sativa*; Pt, *Populus trichocarpa*; Pv, *Phaseolus vulgaris*; Ta, *Triticum aestivum*; Zm, *Zea mays* (see Supplementary Table S1 for gene/protein identifiers). A distance-based tree was reconstructed using the BioNJ algorithm. Numbers in nodes represent the bootstrap obtained for 1000 replicates. The tree is rooted on the *Escherichia coli* TPS (otsA). Sub-clades in the tree are coloured according to the position of *A. thaliana* TPS sequences: AtTPS1/2/4 (red), AtTPS5 (orange), AtTPS6 (yellow), AtTPS7 (green), AtTPS8/9/10 (blue) and AtTPS11 (violet), with the exception of the monocot-specific sub-clade (brown). *S. viridis* sequences are highlighted in grey.

Figure 2. Relative expression of transcripts encoding Tre6P-related enzymes in MC and BSC from *S. viridis* leaves. Relative expression data were calculated with the $2^{-\Delta Ct}$ method, using the *SvKIN* transcript as a reference. Data are the mean ± standard error of 4 biological replicates. Statistical analysis was performed using a two-tailed t-test (unequal variance). Significant differences are indicated with asterisks: *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$. nd, not detected.

Figure 3. Immunoblotting of SvTPS1 in different tissues and cell types of *S. viridis*. (A) Protein abundance in MC and BSC from *S. viridis* leaves. Data were retrieved from Calace *et. al*. (2021). MC, grey bars; BSC, black bars; AU, arbitrary units; nd, not detected. SvPEPC, Sevir.4G143500; SvNADP-MDH, Sevir.6G256400; SvRbcS, Sevir.3G276100; SvNADP-ME; Sevir.5G132500; SvTPS1, Sevir.3G180900. Data are the mean ± standard error of 4

biological replicates (n=4), except for SvTPS1, which was only detected in two BSC samples (n=2). Statistical analysis was performed using a two-tailed t-test (unequal variance). Significant differences are indicated with asterisks: **, $p<0.01$; ***, $p<0.001$. (B) Immunoblotting of SvTPS1 in MC and BSC from *S. viridis* leaves extracted with TCA. Each lane was loaded with extract from 2 mg fresh weight of tissue. (C) Immunoblotting of SvTPS1 in MC and BSC from *S. viridis* leaves extracted with Laemmli buffer. Each lane was loaded with extract from 2 mg fresh weight of tissue. (D) Immunoblotting of SvTPS1 in different tissues and cell types from *S. viridis*. Each lane was loaded with 25 µg of protein. IF; inflorescence, FL; flag leaf, IN; internode, SL; seedlings, MC; mesophyll cells, BSC; bundle sheath cells, WL; whole leaf. The loading controls and full-size images are presented in Supplementary Fig. S7.

Figure 4. Immunolocalization of SvTPS1 in cross-sections of *S. viridis* leaves. Top row, anti-SvTPS1 (1/100); middle row, anti-RbcL (1/1000); bottom row, negative control (without primary antibody). The secondary antibody (anti-rabbit IgG, 1/200) was conjugated with Cy2. The green channel shows the Cy2 fluorescence and the merged images show the Cy2 signal superimposed on the bright-field images. The panel was prepared with digitally-maximized images (2.5X). Bars=10 µm. Non-maximized images are presented in Supplementary Figure S9. The panel shows representative images of three independent experiments (n=3). bs, bundle sheath; vb, vascular bundle; m, mesophyll; ue, upper epidermis; le, lower epidermis.

Figure 5. Distribution of selected metabolites in BSC and MC from *S. viridis* leaves. The abundances of each metabolite in BSC and MC were calculated as described in the methods section from metabolite/PRK *vs* PEPC/PRK plots (as shown in Supplementary Figure S6).

Data are the mean of 4 biological replicates (except for starch, n=3) from one representative experiment (Experiment 2, Supplementary Table S4).

Figure 6. Correlation analysis of metabolites in fractions of *S. viridis* leaves. Data presented in panels A-C correspond to whole-leaf samples from the time-course experiment (Supplementary Table S8), while data presented in panels D-F correspond to one representative fractionation experiment (Experiment 2, Supplementary Table S4). Pearson correlation coefficients (r) between Tre6P and glucose (A and D), fructose (B and E) or sucrose (C and F) were calculated with a 95% confidence interval.

Figure 1. Phylogenetic relationships of *Setaria viridis* TPS proteins. TPS protein sequences were collated from *Setaria viridis* (Sv) and other selected species: Ac, *Ananas comosus*; At, *Arabidopsis thaliana*; Bd, *Brachypodium dystachion*; Mc, *Musa acuminata*; Os, *Oryza sativa*; Pt, *Populus trichocarpa*; Pv, *Phaseolus vulgaris*; Ta, *Triticum aestivum*; Zm, *Zea mays* (see Supplementary Table S1 for gene/protein identifiers). A distance-based tree was reconstructed using the BioNJ algorithm. Numbers in nodes represent the bootstrap obtained for 1000 replicates. The tree is rooted on the *Escherichia coli* TPS (otsA). Sub-clades in the tree are coloured according to the position of *A. thaliana* TPS sequences: AtTPS1/2/4 (red), AtTPS5 (orange), AtTPS6 (yellow), AtTPS7 (green), AtTPS8/9/10 (blue) and AtTPS11 (violet), with the exception of the monocot-specific sub-clade (brown). *S. viridis* sequences are highlighted in grey.

Figure 2. Relative expression of transcripts encoding Tre6P-related enzymes in MC and BSC from *S. viridis* leaves. Relative expression data were calculated with the $2^{-\Delta Ct}$ method, using the *SvKIN* transcript as a reference. Data are the mean ± standard error of 4 biological replicates. Statistical analysis was performed using a two-tailed t-test (unequal variance). Significant differences are indicated with asterisks: *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$. nd, not detected.

Figure 3. Immunoblotting of SvTPS1 in different tissues and cell types of *S. viridis*. (A) Protein abundance in MC and BSC from *S. viridis* leaves. Data were retrieved from Calace *et. al*. (2021). MC, grey bars; BSC, black bars; AU, arbitrary units; nd, not detected. SvPEPC, Sevir.4G143500; SvNADP-MDH, Sevir.6G256400; SvRbcS, Sevir.3G276100; SvNADP-ME; Sevir.5G132500; SvTPS1, Sevir.3G180900. Data are the mean ± standard error of 4 biological replicates (n=4), except for SvTPS1, which was only detected in two BSC samples (n=2). Statistical analysis was performed using a two-tailed t-test (unequal variance). Significant differences are indicated with asterisks: **, p<0.01; ***, p<0.001. (B) Immunoblotting of SvTPS1 in MC and BSC from *S. viridis* leaves extracted with TCA. Each lane was loaded with extract from 2 mg fresh weight of tissue. (C) Immunoblotting of SvTPS1 in MC and BSC from *S. viridis* leaves extracted with Laemmli buffer. Each lane was loaded with extract from 2 mg fresh weight of tissue. (D) Immunoblotting of SvTPS1 in different tissues and cell types from *S. viridis*. Each lane was loaded with 25 µg of protein. IF; inflorescence, FL; flag leaf, IN; internode, SL; seedlings, MC; mesophyll cells, BSC; bundle sheath cells, WL; whole leaf. The loading controls and full-size images are presented in Supplementary Fig. S7.

Figure 4. Immunolocalization of SvTPS1 in cross-sections of *S. viridis* leaves. Top row, anti-SvTPS1 (1/100); middle row, anti-RbcL (1/1000); bottom row, negative control (without primary antibody). The secondary antibody (anti-rabbit IgG, 1/200) was conjugated with Cy2. The green channel shows the Cy2 fluorescence and the merged images show the Cy2 signal superimposed on the bright-field images. The panel was prepared with digitally-maximized images (2.5X). Bars=10 µm. Non-maximized images are presented in Supplementary Figure S9. The panel shows representative images of three independent experiments (n=3). bs, bundle sheath; vb, vascular bundle; m, mesophyll; ue, upper epidermis; le, lower epidermis.

Figure 5. Distribution of selected metabolites in BSC and MC from *S. viridis* leaves. The abundances of each metabolite in BSC and MC were calculated as described in the methods section from metabolite/PRK *vs* PEPC/PRK plots (as shown in Supplementary Figure S6). Data are the mean of 4 biological replicates (except for starch, n=3) from one representative experiment (Experiment 2, Supplementary Table S4).

Figure 6. Correlation analysis of metabolites in fractions of *S. viridis* leaves. Data presented in panels A-C correspond to whole-leaf samples from the time-course experiment (Supplementary Table S8), while data presented in panels D-F correspond to one representative fractionation experiment (Experiment 2, Supplementary Table S4). Pearson correlation coefficients (r) between Tre6P and glucose (A and D), fructose (B and E) or sucrose (C and F) were calculated with a 95% confidence interval.