Crosslinked agarose-gelatine beads as a substrate for investigating biofilms of bacterial pathogens

Dan Roizman[1§], Maren Herzog[1§], Arpita Nath[1], Nivetha Pachaimuthu[2], Ahmad Hujeirat[1], Benno

Kuropka[3], Jens Rolff[1*], Alexandro Rodríguez-Rojas[1,2*]

[1]Evolutionary Biology, Institut für Biologie, Freie Universität Berlin, Berlin, Germany.

[2]Small Animal Internal Medicine, Clinic for Small Animals, University of Veterinary Medicine, Vienna,

Austria.

[3]Institute of Chemistry and Biochemistry, Freie Universität Berlin, Berlin, Germany.

* Jens.Rolff@fu-berlin.de, Alexandro.Rojas@vetmeduni.ac.at

Abstract

Treating chronic bacterial infections is challenging due to the formation of biofilms, making bacteria

less susceptible to antimicrobials. *In vitro* models have limitations in replicating biofilm physiology.

To address this problem, we have created a hydrogel substrate that combines crosslinked agarose

and gelatine presented as beads, providing stability and resistance to autoclaving. Bacterial

pathogens rapidly colonise these biogel beads when submerged in liquid culture. The substrate was

tested with *Escherichia coli*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus*, showing more

robust biofilm growth than its glass bead counterpart. Additionally, this led to increased virulence

factor production and served as a reservoir for biofilm quorum sensing molecules. These features

closely resemble clinical situations, suggesting a more accurate representation of biofilm-associated

infections than current approaches. This new substrate offers a practical and convenient model for

studying biofilms of bacterial pathogens, providing an efficient solution to the research community

and holding promise for future breakthroughs.

Introduction

Biofilms are microbial communities embedded in a self-produced extracellular gelatinous matrix. According to fossil records, they date back 3.5 billion years and facilitate cell adhesion and surface attachment [1,2]. Biofilms are ubiquitous and can be found in diverse habitats, ranging from the mammalian gut to deep subsurface rocks [3]. It is a predominant lifestyle on Earth, and estimations say 40-80% of bacterial and archaeal cells exist in biofilm form [4].

Bacterial biofilms are particularly associated with chronic infections and can withstand high antibiotic concentrations [5]. Biofilms are implicated in infections related to medical devices and other persistent conditions [6]. During infection, bacterial biofilms are often associated with treatment failure [7]. The lack of consensus among the diverse biofilm cultivation methods and analysis techniques further complicates the issue [8].

The most widely used method for biofilm formation involves microtiter plates; initially developed by Fletcher in 1977 for studying bacterial attachment, this technique has since been refined and used for quantifying biofilm biomass [9]. Several methodologies exist to study biofilm formation under simulated natural conditions. One example is the modified Robbins device [10], which utilises a flow system to circulate liquid-containing microorganisms [11]. Advances in fluorescence-based microscopy, such as confocal microscopy, have significantly contributed to our understanding of biofilm mechanisms when combined with later open [12] and closed flow chamber systems [11]. However, a significant limitation of contemporary systems is that they are conducted on artificial surfaces, primarily glass or plastic substrates [13]. Other materials such as silicone, Teflon, PTFE, polypropylene, titanium, or other metals such as stainless steel or copper are also commonly used. Data extrapolated from research on biofilm bactericidal compounds conducted on abiotic surfaces, such as polycarbonate filters, may not be entirely applicable to biofilms growing on biotic surfaces, such as skin or mucoid tissues, due to ultrastructural and physiological variations. These limitations underscore the need for new approaches to studying biofilms [14].

In 1881, Robert Koch used gelatine to develop the first solid media. However, its practicality was compromised due to its susceptibility to bacterial digestion and melting temperature below 37°C. These limitations prompted a search for alternative agents. A suitable substitute was found when Angelina Hesse, the wife and assistant of Walther Hesse, an associate of Robert Koch, proposed using agar to prepare solid cultures in 1887 [15]. Agar offered several advantages over gelatine,

including resistance to bacterial degradation and a higher melting temperature, making it more
practical for microbial growth [16]. Apart from its stability, agar alone is not an accurate model for
recapturing conditions closer to *in vivo* infections [17]. Nonetheless, procedures based on fast
planktonic growth and single colony isolation at 37°C rendered gelatine obsolete, although it
represented the closest proxy to a host environment [16]. Notably, bacterial pathogens discovered
and described in that era using Koch's postulates still comprise most of the priority list 150 years
later [18,19]. The increased challenges of handling those infections necessitate a paradigm shift. Its
rationale is twofold: balancing the focus between biofilm and planktonic lifestyles and having better
substrates that mimic organic surfaces, making them more comparable to within-host scenarios.

Agarose, a purified derivative of agar, is used to manufacture chromatography media for
biomolecule purification through crosslinking its polysaccharide molecules [20,21]. The same
crosslinking method serves as the basis for the work presented here. Our goal was to enhance the
biocompatibility of the substrate by incorporating gelatine, inspired by an emulsification method to
create bacteria-embedded agarose beads [22–24]. Additionally, we found advantages in the
substrate shape of the *in vitro* biofilm models that employ glass beads as a substrate [25–27]. The
ability of bacteria to attach to glass beads and grow, forming aggregates or biofilms, was well
documented as early as the 1940s by researchers like Heukelekian and Heller in 1940 [28] and Zobell
in 1943 [29], even before the term "biofilm" was coined. Building on previous research on biofilms,
we develop a biological hydrogel-based substrate to enhance existing *in vitro* biofilm models for
pathogenic bacteria. Our method combines and stabilises the advantages of agar and gelatine, the
two classic substrates for bacterial growth in semisolid media. We also took advantage of the
protocols established for glass bead biofilm [25] and accordingly shaped our substrate as a bead.

We chose three well-established bacterial models to study the biofilms of bacterial pathogens on the
surface of crosslinked agarose gelatine. Specifically, we examined reference strains from three
bacterial families of the ESKAPE group [30], consistently listed as major concern pathogens [18,31].
Among these are two notable pathogenic bacteria, one gram-negative and one gram-positive:
Pseudomonas aeruginosa and *Staphylococcus aureus*. We also employed the non-pathogenic
Escherichia coli K-12 model, the best-established bacterial model system since its initial isolation a
hundred years ago [32]. Although *E. coli* is generally considered a moderate biofilm former, its
extensive study history allows for in-depth analysis due to all genetic toolboxes and the accuracy of
modern genome databases such as EcoCyc [33].

Results

Generation of agarose-gelatine beads and properties

First, we have focused on the two traditional substrates for semisolid media in microbiology: agarose and gelatine matrices. Due to their biological compatibility and low reactivity, we prepared a crosslinked polymer based on these materials to serve as a substrate. We explored various concentrations of reagents and the crosslinking conditions to create a set of conditions that could be used as a starting point. The method we adopted, based on the catalytic crosslinking activity of divinyl sulphone (DVS), which was first demonstrated by Porath, Låås & Janson [21], which, immediately following adaptation, showed great flexibility for diverse experiments in biofilm analysis. We attempted to develop optimal properties that would enable the fabrication of spherical beads. Combining 2% agarose and 2% gelatine produced the most favourable results for generating biogel spherical structures. Lower concentrations (ranging from 0.5 to 1%) led to easily deformable beads, while higher concentrations (ranging between 3 and 8% each) solidified rapidly, posing handling challenges. These structured, spherical beads were formed by depositing desired volumes of the hot mixture of agarose-gelatine in ice-cold mineral oil, as illustrated in Fig. 1A-E. Some preliminary experiments also confirmed that agarose-alone beads showed a lower biofilm growth than the combined beads. Gelatine-alone beads showed lower stiffness and deficient crosslinking compared to agarose or agarose-gelatine combination.

Initially, we used a multi-stepper pipette to dispense the hot mixture (2% agarose-gelatine each, at 60°C) into the ice-cold mineral oil container (Supplementary Video S2). The drops of agarose-gelatine assumed a spherical shape and gradually sank to the bottom of the mineral oil while solidifying due to the low temperature. This manual method is effective if a small number of beads are required. We also developed a high-throughput system to improve efficiency and streamline the process. This involved using a magnetic stirrer with heating to maintain the solution in a homogeneous liquid state and employing a peristaltic pump to achieve a continuous flow of drops, enabling us to create thousands of beads with consistent sizes in a short period. We estimate that the above system can generate between 60 to 75 beads per minute. By standardising the drop volume to approximately 40 µl by modulating the pump flow rate, we achieved an average diameter of approximately 4 mm, ensuring a highly reproducible procedure (see Fig. 1, Fig. S1, Supplementary Video S1 and Table S1 for further details). The coefficient of variation remained below 2% across five independent batches (Figure S3, Table S1). This approach enabled the bulk creation of agarose-gelatine beads (AGBs), a typical size of glass beads used for biofilm cultivation [34]. The crosslink of

the beads using DVS is a method akin to fabricating Sepharose, a chromatographic matrix commonly

used for biomolecule purification [20]. Our protocol also demonstrated high reproducibility, as

shown in Fig. S1, and the biogel showed significant stiffness and thermostability properties. Unlike

its non-crosslinked counterpart, this biogel can withstand autoclaving without structural

degradation. We also found that the beads need to remain in liquid throughout the production

process and later for long-term storage since desiccation and later reconstitution are disruptive for

the structure in contrast to chromatographic media. When possible, we tried to keep beads at the

relevant medium for a specific experiment, starting from the autoclave step in that medium.

145
146
Next, we characterised the stability of the beads, which remained stable for at least five months

after crosslinking and autoclaving. During the storage period in the culture medium, the AGBs did

not change the aspect or diameter (Fig. 1F-G). The crosslinking not only provided the chemical

stability to allow the beads to be autoclaved, but we also determined that the crosslinking

significantly increased the beads' strength, manifested by a significant increase in stiffness as

measured by Young's modulus of an atomic force microscope (Fig. 1H).

***E. coli*, *P. aeruginosa* and *S. aureus* form robust biofilms on the agarose-gelatine beads**

E. coli, *P. aeruginosa,* and *S. aureus* all formed biofilms on the surface of the AGBs, which were

compared to glass beads. Quantification of bacterial biofilm cells by counting colony-forming units

per millilitre (CFU/ml) revealed significantly more biomass for bacterial cells grown on AGBs than on

glass beads for all strains (Fig. 2). We measured these values during the maturation process of

biofilm at 24 and 48 hours. The number of CFUs for all three bacterial species was significantly

higher for AGBs compared to glass beads at 24 and 48 hours. In preliminary experiments, we

observed a higher variation of the bacterial count coming from the AGBs than from the glass beads.

We suspected the heterogeneity was due to protocol limitations, so we modified the bacterial

recovery protocol by incorporating mechanical and chemical methods. The method included using

some enzymes to digest the extracellular matrix (see M&M and SM). We fundamentally used DNase

I and papain, which helped in better dispersion and a good level of single cells in suspension without

affecting bacterial viability, which is crucial for a satisfactory enumeration.

167
168
Agarose-gelatine beads are compatible with scanning electron microscopy and live fluorescence

microscopy

The transparency of agarose is particularly advantageous for live fluorescence microscopy, enabling

clear visualisation of bacterial cells in real-time without interference. Furthermore, the

biocompatibility of agarose-gelatine ensures that the bacterial cells remain viable and maintain their

physiological states, leading to more reliable and relevant results in biofilm research. As a proof of

principle, we constructed fluorescent strains of *E. coli*, *P. aeruginosa* and *S. aureus* on the surface of

the AGBs using fluorescence confocal microscopy (Fig. 3A). We designed a 3D adapter (M&M, Fig.

S4) for a more simplified and stable microscopy observation to study biofilms on 4mm AGBs (Fig.

3B).

Another relevant technique to study bacterial biofilms is scanning electron microscopy (SEM). Due to

their hydrogel nature and spherical shape, the AGBs required special treatment (see M&M section)

to fix, dehydrate, and keep the integrity of the structure of the beads and preserve the bacterial

biofilm. The best results were obtained by drying all the beads with the critical point dryer. Then, we

could perform scanning electron microscopy of *P. aeruginosa* and *S. aureus* biofilms (Fig. 3C). The

gelatine component offers a suitable texture that facilitates sample preparation and enhances

electron imaging, capturing fine details of biofilm architecture.

Quantitative proteomics (liquid chromatography mass spectrometry, LC-MS) shows that biofilms

on Agarose-Gelatine beads and glass beads have different physiology

Our investigation into the differences between bacterial biofilms growing on AGBs and glass beads,

in terms of gene expression and bacterial physiological changes, was conducted with the rigorous

method of label-free quantitative proteomics. This approach allowed us to identify distinct protein

expression profiles in biofilms grown on agarose-gelatine beads (AGBs) compared to glass beads for

three bacterial species - *E. coli*, *P. aeruginosa*, and *S. aureus*. It's worth noting that we employed a

modified protocol ad doc for this study with variations from previous works [35,36]. We identified

and quantified 1113 proteins for *E. coli*, 1500 for *P. aeruginosa,* and 1251 for *S. aureus*

(Supplementary Table S1, S2, and S3, respectively). Proteins showing at least a 2-fold change in their

relative intensity between glass beads vs. agarose-gelatine beads, with an FDR-adjusted *p*-value <

0.05, were considered significantly changed. Significantly, we observed notable abundance changes

in important virulence factors for *P. aeruginosa* and *S. aureus*, potentially indicating differences in

surface sensing due to chemical composition (Fig. 4).

In the biofilms of *E. coli*, the 50S ribosomal subunit protein L35 (RpmI) had a noticeable increase in

abundance when grown on AGBs compared to glass beads (log2 fold change (log2FC) = 2.5).

Similarly, the small 30S ribosomal subunit S16 (RpsP) and ribosomal silencing factor (RsfS) were

upregulated, but to a lesser extent. RsfS is known to be expressed in slowly growing cells and inhibits

translation, indicating that the population of AGBs is in the mature phase [37]. On glass beads, there

is a distinct upregulation of 50S ribosomal subunits RpmH and RplU, or subunits L34 and L21; at

log2FC = -5.2, RpmH was the most differentially abundant protein. Apart from this striking difference in ribosomal structural content, several other ribosome-associated proteins were upregulated on glass: RsmH, Sra, YgaM and Rmf. RsmH is 16S-rRNA-m4C1402-methyltransferase, ubiquitous in bacteria and eukaryotic organelles; it facilitates a modification of the ribosomal P translation initiation site, allowing different codon usage for more efficient ribosome biogenesis [38]. The physiological roles of Sra and YgaM are less known but considered regulatory [37]. The former tends to bind active 70S Ribosomes during stationary phase and maturation, and the latter provides ribosomes with an anchoring point to the membrane. Rmf, or Ribosome modulation factor, is essential for the reversible formation of 70S-90S-100S Ribosome Dimers in gammaproteobacteria [37]. There is a higher expression of NusA (log2FC = 2.9) on glass beads, which acts as a transcription terminator factor and chaperone of rRNA, increasing their numbers; NusA is expressed following stress such as cold shock [39]. Overall, we see different ways of maturing and diversifying the ribosomes in size, structure, activity, and biogenesis. This trend is evident in all three strains, with potentially more regulatory shifts and stress response adaptations in biofilms on glass beads. The data also shows a notable change in cell division via Z-Ring formation through FtsZ polymerisation, an essential process for cell elongation and division. Notably, ZipA, which anchors FtsZ to the membrane, was significantly more abundant (log2FC = 2.2) on AGBs, contrasting with the upregulation of ZapA (log2FC = -2.1) on glass beads, influencing cell division dynamics [40].

In *P. aeruginosa*, important virulence factors were found to be more abundant in AGBs than in glass beads. Elastase (LasB) showed the most significant change in relative abundance (log2FC = 5.1). Elastase is regulated by the LasR/LasI quorum-sensing system, indicating increased virulence factor activity in biofilms grown on AGBs [41,42]. PrpL (log2FC = 3.2), also known as Protease IV or lysyl-endopeptidase [41,42], which can cleave fibrinogen and disrupt the activity of the immune system, was also elevated. Several proteins involved in pyoverdine biosynthesis, including PvdA, PvdJ, PvdD, and PvdL, also showed an increased abundance in AGB-biofilms. Pyoverdine's primary function is iron chelation from host cells, which is involved in *in vivo* infections [43]. Another significantly upregulated protein in AGBs is the aminopeptidase Lap, also known as PaAP, which cleaves extracellular proteins [42] and PhzM, involved in pyocyanin biosynthesis [44]. The only virulence-related upregulated protein on glass beads was AmrZ, also known as AlgZ (log2FC = 3.0). This important transcription factor regulates Alginate production and motility [45,46]. Conversely, among the downregulated proteins on AGBs were components of the Sec translocon system, such as YajC (log2FC = -4.4) and SecF (log2FC = -2.5), indicating reduced protein export and altered membrane dynamics [47]. The second and third most abundant proteins on glass beads are uncharacterised and

coded by the genes PA0049 and PA5305 (with log2FC values of -4.2 and -3.7, respectively). PA0049
expression is under the stringent redundant response regulator DksA in *P. aeruginosa* [48]. A large
proportion of proteins are probable, putative, uncharacterised, or hypothetical for *P. aeruginosa* (18
of the 55 abundant proteins on AGBs and 113 of the 240 on glass beads, between 30-50%), detected
to be above our specified threshold for significant change. *S. aureus* showed a similar trend, making
comparative analysis more intricate for both organisms relative to *E. coli*.

We observed a similar trend in *S. aureus* biofilms as in *P. aeruginosa*, where we found higher
virulence factor expression in AGBs than in glass beads. However, *S. aureus* showed a more complex
response with more differentially expressed proteins for both substrates (247 proteins on AGBs
versus 297 on glass beads). *S. aureus* had about twice the number of proteins with altered levels
compared to *E. coli* and *P. aeruginosa*. These results indicate that different species have differential
responses in the class of proteins and the number of genes that respond to the surface where the
biofilms develop.

DeoC was the highest expressed protein on AGBs (log2FC = 5.0), and it is usually part of the inducible
deo operon in bacteria, enabling the metabolism of exogenous nucleosides [49]. DeoC, or
deoxyribose phosphate aldolase, is a duplicated gene in *S. aureus*, and both copies are highly
upregulated on AGBs (log2FC = 2.5), along with other genes in this operon, DeoB and DeoD [50]. The
second most abundant protein on AGBs was MsrB (4.9 log2FC), a methionine sulfoxide reductase
that protects against oxidative stress and is a marker for mature biofilms [51]. DltC, critical in
teichoic acid biosynthesis, cell envelope overall charge, and virulence, was next on the list [52]. The
most abundant protein on glass beads was SAOUHSC_02161 (8.0 log2FC), an MHC class II analogue
protein of unknown function, under the SaeRS two-component system regulon and considered an
attachment protein [53].

Several transcription factors and additional regulators were significantly increased on glass: SarR,
FapR, Rex, RecX, MsrR, and Spx; while on AGBs, seven other transcription factors were upregulated:
ArcR, NusB, NusG, CodY, NusA, SrrA, RsbW and VraR. SarR downregulates the expression of SarA,
which usually activates the *agr* operon quorum sensing system, leading to increased production of
exotoxins required for virulence [54]. FapR is a regulator of membrane lipid biosynthesis and lipid
homeostasis [55]. Rex is a redox-responsive repressor activated by the ratios of NADH:NAD in cells;
its regulon enzymes are involved in many metabolic pathways [56]. RecX is an anti-recombinase as it
inhibits RecA under accumulated DNA damage [57]. MsrR regulates the cell envelope homeostasis

and virulence factor production by structural means, abundant in antibiotic-tolerant cells [58]. Spx is
a global effector that accumulates cells during various stresses and regulates many pathways, most
notably thioredoxin, by a unique mechanism [59]. In biofilms on AGBs, the most abundant regulator
is ArcR, which is critical for anaerobic metabolism, mainly allowing the metabolism of arginine over
sugars [60]. Nus is an accessory protein forming a complex with RNA polymerase, enhancing
bacterial rRNA transcription and synthesis [61]. CodY regulates the *agr* virulence system depending
on population density and nutrient scarcity [62]. SrrA is part of the two-component system SrrAB
that increases virulence factor production through *agr* at low oxygen levels [54]; RsbW is a serine-
protein kinase acting as an anti-sigma factor to SigB, inhibiting transcription after binding [63]; and
VraR is part of the two-component system VraSR, which is upregulated in response to cell-wall
damage affecting peptidoglycan homeostasis [64].
287
Regarding other pathways leading to virulence factor abundance, we observed a differential
upregulation on glass beads, notably the biosynthesis pathway of staphyloxanthin: AldH1, CrtM,
CrtP, and CrtN. Additional virulence factors that were more abundant on glass beads comprise two
fibrinogen-binding proteins (SAOUHSC_01114, SAOUHSC_01110), immunoglobulin binding Sbi and
two uncharacterised leukocidin-like proteins (SAOUHSC_02241, SAOUHSC_02243). Several subunits
of Phenol-Soluble Modulin are upregulated (psmA1/A2 and psmA4). ClfA, or clumping factor,
facilitates cells' clumping, thus defending them from phagocytosis. There is a notable abundance of
extracellular active enzymes such as serine protease SplE and Coagulase (SAOUHSC_00192) and a
plethora of known staphylococcal antigens (IsaB, SsA2, SsL10, SsaA). On AGBs, we can detect the
upregulation of virulence-promoting factors, such as TraP and LuxS. TraP is involved in the
upregulation of the *agr* operon in a cell density-dependent manner [65], LuxS is the enzyme which
produces autoinducer-2 molecule, which in turn results in enhanced biofilm phenotypes [66]. Other
notable enzymes involved in pathogenesis were also more abundant on AGBs, including lipase-1
LipA, catalase katA, and components of the type VII secretion system components, EsxA and EsaG,
important for intracellular infection of the host [67].
303

The AGBs are suitable for the detection of biofilm-deficient strains

Our model system presents a valuable tool for detecting or screening genes and conditions
influencing biofilm formation. To validate this concept and assess the capability of AGBs to detect
variations in biofilm formation, we examined two bacterial strains known for their deficiencies in
biofilm formation. Previous biofilm research has shown that type I fimbriae, produced by the *fim*
operon, are crucial for forming submerged biofilms. Additionally, type I pili, including the mannose-

specific adhesin FimH, are for initial surface attachment [68]. Earlier studies reported that disrupting

the *pfpI* gene in *P. aeruginosa* decreased the bacterium's ability to form biofilms [69]. When we

cultivated both mutant strains in AGBs and compared them with their respective parental wild-type

strains, we observed a significant decrease in biofilm formation (Fig. 5A). This finding indicates that

our AGB system is suitable for detecting differences in biofilm formation between strains.

Chemical modification of agarose-gelatine beads is a valuable strategy for investigating anti-

biofilm compounds

One of the significant advantages of crosslinked agarose is its potential for conjugating diverse

chemical compounds to its surface. This feature benefits our biofilm model and has practical

implications for investigating antibiofilm compounds to protect surfaces, a crucial aspect of biofilm

research. To demonstrate this concept, we chose melittin, an antimicrobial peptide known for its

anti-biofilm properties [70]. Following the protocol described by March *et al.*, we used the cyanogen

bromide method to couple melittin to agarose-gelatine beads [71]. Cultivating biofilms of our three

bacterial models on melittin-decorated AGBs revealed a dramatic decrease in bacterial attachment

to the bead surfaces compared to untreated and BSA-coupled beads. For all three bacterial

species—*E. coli*, *P. aeruginosa*, and *S. aureus*—the number of biofilm-forming bacteria was reduced

by 4 to 5 orders of magnitude compared to the controls (Fig. 5B). Biofilms grown on BSA-coupled

beads exhibited the highest biofilm formation, indicating that the chemicals used for protein

coupling to the AGBs were not responsible for the observed biofilm inhibition. Notably, final counts

of planktonic bacteria did not differ in wells where bacterial species were incubated, indicating that

the observed effects were specific to biofilm formation.

Agarose-gelatine bead biofilms can be quantified with a crystal violet assay

A popular and convenient method for quantifying bacterial biofilms is to measure the amount of

crystal violet stain retained by the biofilms [72]. We tested this method with our AGBs and

compared the performance of the crystal violet staining with biofilms grown on glass beads. Initially,

when using the standard protocol, we observed that the beads accumulated a significant amount of

background crystal violet compared to the glass beads. To address this issue, we made some

modifications to the protocol. By adjusting the crystal violet concentration, incubation temperature,

and washing times, we optimised the protocol for the AGBs. As a result, we confirmed through the

crystal violet assay (see Methods and Materials section) that bacteria growing on the surface of

AGBs form more biofilm than their glass bead counterparts (Fig. 5C).

Agarose-gelatine beads can serve as small molecule reservoirs

In contrast to other abiotic biofilm substrates, the permeability of crosslinked agarose could serve as

a reservoir of small molecules such as nutrients, quorum-sensing molecules, drugs, etc. This property

can be advantageous in certain experiments as it may occur in *in vivo* biofilms where bacteria are

attached to host surfaces, such as mucosal layers or necrotic tissue debris. Our experiments involve

two bacterial models that produce detectable substances that could accumulate within the agarose

beads. We observed that the agarose beads consistently appeared yellower for *S. aureus* or greener

for *P. aeruginosa*, indicating the diffusion of pigments into the inner part of the beads (Fig. 6A-B). For

P. aeruginosa, upregulation of the biosynthetic pathway of Pyoverdine was evident in AGBs; in

contrast, upregulation of the Staphyloxanthin was apparent on glass beads. To try to quantify

changes in the concentration of those pigments in bead-containing samples, we adapted some

chemical and physical protocols (Materials and Methods) to extract them into the solution that

could be measured with a plate reader relative to CFU/mL numbers (Fig. 6A-C); A similar approach to

this protocol has been taken in a recent publication [73]. We also propose that this hydrogel could

facilitate the diffusion of soluble molecules, including nutrients and bacterial quorum-sensing

molecules while retaining some molecules within its structure as a reservoir, as observed in many *in*

vivo systems. We estimated the quantity of N-acyl-homoserine-lactones (AHL) in the *P. aeruginosa*

PAO1 bead-containing samples using a previously described colourimetric method [74] with

adaptations. The assay indicates a significantly higher lactone content in AGB samples than glass

(Fig. 6D).

Discussion

Our biofilm model has significant potential for a wide range of research applications (see Fig. 7).

AGBs can be used to test anti-biofilm molecules, evaluate surface protection compounds, study

quorum sensing mechanisms, investigate biofilms on foreign bodies, and conduct microscopy and

'omics profiling studies of biofilms. Additionally, our system is compatible with traditional

experimental approaches, including crystal violet quantification and live and scanning electron

microscopy.

One attractive application of our system is the ease of conjugating molecules to the surface of the

beads. The chemical modification of the AGBs is a great advantage, as demonstrated in this study by

conjugation with melittin. The attachment of this antimicrobial and antibiofilm peptide significantly

reduced biofilm formation by *E. coli*, *P. aeruginosa*, and *S. aureus*. Melittin, a small cationic peptide

composed of 26 amino acid residues [75], is a principal constituent of bee venom and is used as a

tool of social immunity to protect their colony from pathogen invasion [76]. This property of easy

compound conjugation can also be used to investigate what elements stimulate biofilm attachment
and avoid certain materials or use them in beneficial biofilms [77].

Agarose-gelatine beads exhibit a unique capability to serve as reservoirs for small molecules. They
allow for the gradual release of nutrients, quorum-sensing molecules, drugs, and other minor
compounds, thus providing a more accurate simulation of *in vivo* biofilm conditions. This feature is
advantageous for experimental setups replicating complex environments where bacteria adhere to
host surfaces or necrotic tissue debris [78]. The diffusion of virulence factors such as
staphyloxanthin, pyoverdine, pyocyanin, and quorum-sensing molecules exemplified this capability.
Staphyloxanthin is an interesting example because the proteomic analysis shows downregulation of
its biosynthesis. It is present in cell membranes, protecting lipids, proteins, and DNA from ROS [79].
The main enzymes in ROS defence are catalases and superoxide dismutase, while staphyloxanthin
provides additional protection [80]. Clinical samples from diabetic patients' foot ulcers have shown
that staphyloxanthin impairs wound healing and enhances *S. aureus* survival in presence of ROS [81].
It is known that a substantial reduction in staphyloxanthin production occurs in correlation with
biofilm maturation [82]. Additionally, staphyloxanthin accumulation within the AGBs, as we have
shown, could be responsible for inhibiting pigment production.

The biofilm system based on AGBs proves effective for detecting genetic factors influencing biofilm
formation, as demonstrated by studying mutant strains deficient in biofilm formation, explicitly
focusing on the *fimH* of *E. coli* [83] and the *pfpI* gene in *P. aeruginosa* [69]. Comparing biofilm
formation in AGBs between mutant and wild-type strains revealed a significant reduction in the
mutants' biofilm formation, validating the AGB system's capability to detect variations in biofilm
formation under controlled conditions.

The quantitative proteomic profiling performed in this study revealed distinct protein expression
profiles for *E. coli*, *P. aeruginosa*, and *S. aureus* biofilms grown on AGBs compared to glass beads.
Some studies suggest that bacterial growth on abiotic surfaces significantly differs from the bacteria
from *in vivo* biofilm-related infections [84]. Many more differentially expressed proteins are crucial
virulence factors for *P. aeruginosa* and *S. aureus*, likely reflecting differences in surface sensing. For
E. coli, there were significant differences in ribosomal protein expression in biofilms on AGBs and
glass; A similar trend was observed for *P. aeruginosa* and *S. aureus*. In *P. aeruginosa*, essential
virulence factors such as elastase (LasB) and the pyoverdine synthesis pathway were more abundant
in biofilms on AGBs. For *S. aureus*, biofilms on AGBs and glass beads exhibited differences in protein

expression and gene regulation; it exhibits differential expression of factors for host cell recognition
and utilisation of host resources. The study identified several regulatory proteins that mediate the
expression and effects of the quorum sensing system. In conclusion, biofilms from AGBs and glass
beads represent different phenotypes, with the former possibly resembling a mature and
established biofilm and the latter resembling a phenotype closer to a bloodstream infection.
It does not escape our attention that this new biofilm model offers several applications. Firstly, it
may enable *in vivo* experiments that investigate biofilms in the context of infections associated with
foreign objects, providing crucial insights into the dynamics and treatment of such infections.
Secondly, the hydrogel material could be potentially used in bioengineering and 3D printing as a
biopolymer for fabrication, opening avenues for advanced tissue engineering and regenerative
medicine. Lastly, AGBs can serve as an effective model for understanding the role of biofilms in
bacterial resistance to antibiotics, allowing the scientific community to conduct experimental
evolution studies that could lead to novel strategies for combating antibiotic resistance and biofilm-
associated infections. Besides, because of the hydrogel nature of the AGBs, there is also a potential
utilisation to study liquid air interface biofilm.
428
Conclusions
In this study, we have introduced a new substrate based on crosslinked agarose-gelatine hydrogel in
the form of beads as a novel strategy to investigate biofilms of bacterial pathogens. We have shown
that this method is compatible with various biofilm structural studies and quantitative techniques,
including live microscopy techniques such as fluorescence and confocal microscopy, as well as
scanning electron microscopy. Our model is also compatible with other biofilm quantitative
techniques, including crystal violet staining, and we have developed a protocol for the enumeration
of viable cells, which maximises recovery and dispersion using enzymatic and mechanical disruption
of the biofilm. The agarose-gelatine bead model provides a new opportunity to study biofilm
physiology, as the permeability of the hydrogel beads allows the diffusion of small molecules,
including drugs, nutrients, quorum sensing signalling molecules, and other substances such as
siderophores and other secretion products. We have demonstrated that the beads maintain their
properties for an extended period, at least several months.

Materials and Methods

Generation of agarose-gelatine beads. To produce the bead substrate, 4 g of each agarose and

gelatine were added to 200 ml (2% each w/v) of distilled water in a 400 ml bottle and autoclaved for

20 minutes at 121°C and 1 atmosphere. After sterilisation and throughout the production process,

the liquid mixture was kept liquid by placing the flask on a hot plate of the stirrer at 65°C with

moderate mixing. In a separate container, 500 ml of mineral oil (Merck, Germany) was placed in an

ice water bath. The oil was previously degasified to remove air bubbles using a vacuum pump. We

used two procedures to produce the agarose-gelatine beads. First, we used a multistep pipette

(Eppendorf, Germany) to spot around 40 µl of substrate in the cold mineral oil, ensuring that the tip

was between 1 to 2 cm from the surface of the oil to avoid deformation of the beads while

solidifying or partitioning of the drops (See Fig. S1). A second and higher throughput method

incorporated a peristaltic pump P-1 (Pharmacia Biotech, Sweden) that allows a continuous and

semiautomatic dropping of the beads into the mineral oil (Fig. 1). At the end of the silicon tubing, a 2

mm internal diameter stainless steel capillary was incorporated for a more precise holding and

stable drops, two silicone tubes with a diameter of 3 mm and a length of 0.5 m were extended with

one metal tube of the same diameter and a length of 20 cm. The flexible ends were connected to the

peristaltic pump, while the ends with the metal tubes opened into the bottle with the substrate or

above the collection beaker, respectively. The movement of the peristaltic pump resulted in a stable

output of uniform drops of the agarose-gelatine mix, which solidified upon entering the inert, cooled

mineral oil, forming ~4 mm Agarose-gelatine beads (AGBs). After processing, the beads were sieved

and washed with distilled water with at least 50 volumes until the oil was removed, allowing them to

rest for five minutes between washes to allow the oil separation from the aqueous phase.

Crosslinking of the agarose-gelatine beads. The washed and clean agarose beads were crosslinked

using divinyl sulfone (DVS) using a modified method from the original described elsewhere [21]. The

beads were weighed to calculate the required amount of DVS (v/w) (Merck, Germany).

Approximately 500 ml of 0.5 M sodium carbonate buffer, pH 11, was used per 100 grams of gel

beads (five volumes). The beads were gently transferred with the help of a funnel to an 800 ml flask,

capped, and the crosslinking reaction was initiated by adding DVS, a final concentration of 0.5%. The

reaction was allowed to proceed on a shaker at room temperature overnight, covering the beaker

with parafilm to prevent evaporation. The next day, the beads were thoroughly washed with distilled

water by decanting and sieving until a neutral pH was achieved. To remove residual amounts of

unreacted vinyl groups, we placed the beads in five volumes of 0.5 M sodium carbonate buffer, pH 9,

and added beta-mercaptoethanol for a final concentration of 0.5%. The flask was incubated with

shaking overnight in the dark at room temperature. Afterwards, 500 ml of distilled water was used

to wash the beads five times. To remove the residual unreacted vinyl group, the beads were covered

with five volumes of 1 M sodium carbonate, pH 10, and glycine was added for a final concentration

of 10% (w/v). The reaction was also conducted overnight at room temperature with shaking. The

next day, the beads were washed three times with five volumes of 0.1 M carbonate-bicarbonate

buffer at pH 10 and a 0.1 M citrate buffer at pH 4.0. The beads were rewashed with five volumes of

distilled water three times and added directly to a flask containing five volumes of bacterial culture

media lysogeny broth (LB) or Mueller-Hinton broth (MHB). The bottles were autoclaved and stored

at +4°C until use. For a step-by-step protocol, see Protocol I of supplementary materials.

Microscopy of Bead Surfaces. To visualise the intact surfaces of agarose-gelatine glass and beads,

we utilised a VHX-X1 microscope (Keyence, Austria). Glass beads were directly placed into a 45 mm

petri dish for observation. AGBs were first removed from the preservation medium (LB). Then, the

excess liquid was carefully pipetted off before placing the beads into a separate 45 mm petri dish for

direct examination at different magnifications.

Bacterial strains and cultivation conditions. *Escherichia coli* K-12 MG1655 [85], *Escherichia coli*

BW25113, and its derivative mutant *E. coli* BW25113 *fimH::*Kn, both from the Keio collection [86], *P.*

aeruginosa PAO1 [87], *P. aeruginosa PA14* [88] *and its derivative mutant P. aeruginosa PA14*

*pfpI::*MAR2xT7[88], and *Staphylococcus aureus* SH1000 [89] were the strains used for different

experiments across this study. Generally, bacteria were cultivated on Petri dishes from frozen

glycerol stocks on lysogeny broth (LB) agar and incubated overnight at 37°C. The next day, a single

colony was inoculated into 10 ml of liquid LB in 50 ml Falcon tubes and shaken at 200 r.p.m.

overnight at 37°C. The total cell density was determined by measuring the optical density (OD) in a

spectrophotometer at 600 nm. The inoculums were diluted to approximately 10^6 CFU/ml per culture

for biofilm inoculation.

Biofilm cultivation. For biofilm culturing, two beads were aseptically transferred to each well of a

non-binding polypropylene 24-well microplate (Carl Roth, Germany). Overnight cultures of *P.*

aeruginosa, S. aureus SH1000, and *E. coli* K-12 MG1655 in LB medium were diluted to an optical

density of approximately 1x10^5 CFU/ml and added to the microplate containing the beads (1 ml per

well). The 24-well microplate was placed in a multiwell plate shaker with moderate shaking (20-30

r.p.m.) at 37°C for 24 and 48 hours. After 24 hours of incubation, the beads were gently washed with

sterile 0.9% NaCl saline solution by pipetting, and a fresh medium was added.

Biofilm viable bacterial enumeration. After biofilm cultivation, each bead was transferred with

sterilised tweezers into a 1.5 ml microcentrifuge tube (Eppendorf, Germany) containing 1 ml saline.

The samples were sonicated in an ultrasonic bath to detach the biofilm from the bead surface and

separate the bacteria at 40 kHz for 3 min. Before and after sonicating, each microcentrifuge tube

contained a bead and a vortex vigorously mixed with the saline solution. The tubes were centrifuged

at 10,000 x g for 3 minutes, and the pellets resuspended in 200 µl of digestion buffer (150 mM NaCl,

50 mM Tris-buffer, 10 mM $MgCl_2$, 5 mM L-cysteine, 5 mg/ml DNAse I, 2 mg/ml RNAse A, pH 7.5) and

incubated at 37°C for 15 minutes. Afterwards, 20 µl of papain stock solution (10 mg/ml) was added

to each tube and incubated for another 15 minutes at 37°C in moderate shaking. The tubes were

centrifuged at 10,000 x g for 3 minutes and resuspended in 1 ml of 0.9% NaCl. A volume of 20 µl of

each sample was transferred to the wells of the first row of a 96-well microplate. Using a

multichannel pipette, all samples were diluted in 1:10 steps (20 µl sample; 180 µl saline) from the

first row to the last. Droplets of 10 µl for each dilution were spotted on pre-dried LB agar plates (for

45 minutes before use in a laminar flow cabinet). The wells containing the bacterial dilution were

carefully mixed by pipetting before spotting the drops. After drying the droplets (approximately 5–

10 minutes), plates were incubated for 16 hours at 30°C. The next day, the microcolonies were

observed and, if necessary, allowed to grow for another two hours at 37°C. Then, the colonies were

counted for the first dilution, which resulted in a countable range of 5–35 colonies. The bacterial

number was determined, considering the dilution factor, and expressed as CFU/ml.

GFP-tagging. To perform live fluorescent microscopy of bacterial biofilms on the surface of agarose-

gelatin beads, a constitutive green fluorescent protein (GFP) gene was inserted into the

chromosomes of *E. coli* MG1655, *P. aeruginosa* PAO1 and *S. aureus* SH1000 using ADDGENE

Plasmids (pGRG36-Kn-PA1-GFP, pMF230 and pTH100 accordingly) and following the relevant

methodologies published elsewhere [90–92]. Briefly, *E. coli* DH5alpha were used as plasmid donors

carrying either the pGRG36-Kn-PA1-GFP or the pMF230, and *E. coli* DC10b was used for plasmid

pTH100 for integrating GFP expression cassette into the chromosome of the strains used in this

research accordingly. Donor strains were grown under specified conditions using LB broth and

relevant antibiotics. Plasmids were purified using the Pure Yield Plasmid Miniprep System (Promega,

USA) following the manufacturer's instructions and stored at -20°C until use. Mid-exponential

cultures (100 ml) of *E. coli* MG1655, *P. aeruginosa* PAO1, and *S. aureus* SH1000 were grown until an

OD_{600} between 0.4 and 0.6 was reached. *E. coli* and *S. aureus* acceptors were made

electrocompetent using previously described protocols [93,94] transformed with the relevant

plasmids using electroporation (Gene Pulser Xcell™, Biorad, USA) using the manufacturer's preset

protocols for *E. coli* and *S. aureus*. *P. aeruginosa* was transformed by triparental conjugation,

essentially as described [91] using the helper strain *E. coli* S17-1-*lambda-pir* carrying pRK2013 [95].

Growth on LB Agar and a fluorescence-adapted stereomicroscope checked the phenotypes of

marker antibiotic resistance and green fluorescence, respectively. Positive clones were selected,

grown, and kept as glycerol stocks at -80°C until use.

Crystal violet staining. Qualitative quantitation of all biofilm components was performed using

crystal violet staining [96]. For the substrate comparison, AGBs (4 mm) and glass beads (4 mm,

Merck, Germany) were inoculated with *P. aeruginosa* PAO1 and *S. aureus* SH1000. After the

maturation of the biofilm in 24-well polypropylene plates (Carl Roth, Germany) and a final wash, the

beads in each well were covered with 500 µl of NaCl 0.9% saline solution and another 500 µl of a

0.1% crystal violet (Merck, Germany) in saline for 1 min. Using a single-channel pipette, the staining

solution was aspirated and discarded. The beads were carefully rinsed by adding 1 ml of 45°C

prewarmed distilled water several times, with 5 min incubation time between washes. The washing

process was repeated until the water was clear or the wash had an OD_{592} below 0.01. To solubilise

crystal violet retained by the biofilm, 1 ml of a mixed solution of 95% Ethanol [96] and 0.05% of

Triton X-100 [97] were added to each well. The beads were incubated with gentle shaking for 30

minutes at 45°C. 200 µL of each sample were then transferred to the wells of a 96-well plate. The

amount of crystal violet was estimated by absorbance at 592 nm.

Scanning electron microscopy. To visualise biofilm formation on the surface of the bead substrate,

images of colonised Agarose-gelatine beads were taken using a scanning electron microscope (SEM)

(EVO MA 10, Carl Zeiss, Germany). Before removal from the 24-well plates where the biofilm

developed, the samples were washed with saline, as described above. The following fixation steps

were carried out under the safety cabinet due to the high toxicity of the chemicals used. Six samples

of mature biofilm produced by *E. coli* K12, *P. aeruginosa* PAO1 and *S. aureus* SH1000, grown on

AGBs, were collected in a 10 ml glass jar using sterile forceps. Fixation was performed by covering

the samples with 5 ml of 2.5% glutaraldehyde (glutaraldehyde 25%, Carl Roth, Germany) in 0.05 M

cacodylate buffer for 1 hour. The fixing solution was stepwise substituted by 0.05 M cacodylate

buffer for further preparation, ensuring the beads never dried out completely. The samples were

then rinsed three times with 0.05 M cacodylate buffer, each lasting 10 minutes. Osmium tetroxide,

whose primary purpose is contrasting, was used as a secondary fixation agent [98]. Two millilitres of

2% osmium tetroxide (ReagentPlus®, 99.8%, Merck, Germany) in distilled water were added to the

samples immediately after removal of the buffer and incubated for an additional 30 minutes.

Another washing with cacodylate buffer followed as previously described. Dehydration of the fixed

colonised AGBs was performed using alcohol at growing concentrations. This was done as a

preparation for the critical point drying with carbon dioxide (CO_2). The samples were transferred

directly into the critical point dryer's magazine (Foissner vessel, CPD 030, BAL-TEC-Leica, Germany)

for better handling. The alcohol series consisted of 30%, 50%, 70%, 90%, and 2x 100% alcohol with a

molecular sieve. Starting with the lowest concentration, the Foissner vessel was placed in a beaker

filled with alcohol and incubated for 10 minutes. The dehydration process was continued

sequentially, using the highest alcohol concentration, including a molecular sieve. The Foissner

vessel with the samples was now placed in the cabin of the critical point dryer, and the remaining

water and alcohol were entirely replaced by liquid CO_2. Liquid CO_2 changes into the gaseous phase at

31°C and 74 bar. In this way, the surface structure of a sample can be preserved, which would be

damaged in the presence of water during the transition from the liquid to the gaseous state. After

drying, the samples were mounted on 12 mm aluminium stubs and coated with gold using a

sputtering system (SCD 040, BALZERS, Liechtenstein). For an even gold layer on the preparations, the

air in the coater was replaced by inert Argon. The samples were then placed in the SEM and

recorded at working distances from 6.5 mm to 12 mm and magnifications between 50 and 27 k-

times.

Confocal fluorescence microscopy. Initially, the AGBs were cultivated with the *E. coli* GFP-expressing

strain, as described before. After maturation and a last washing step with saline, the samples were

transferred initially into a chambered coverslip with 8 wells and a glass bottom for high-end

microscopy (μ-Slide 8 Well, Ibidi, Germany). The beads in the chambers were covered with saline,

the coverslip was closed with a lid, and the images were acquired with an inverted confocal laser

scanning microscope (SP8-1, Leica, Germany) equipped with a 40x objective. GFP was excited from

488 nm and detected at 510 nm. Both single images and Z-stacks were created from the bead

surface or cross sections and later edited in ImageJ.

For live fluorescence microscopy, we follow the same procedure as described before for E. *coli* for

confocal microscopy with the three species of GFP-expressing strains with some variations. We have

developed a 3D adapter to stabilise the beads on a normal microscope coverslip (60x 25 mm) for

microscope observation of biofilms on 4mm AGBs. The adapter was designed in AutoCAD and

printed using a Form 4B 3D printer (Formlabs, USA). It has precise external dimensions of 25mm x 25

mm and a central 5 mm hole to hold 4mm beads securely (Fig. S4). The 3D_bead_adapter.stl file is

also provided as a supplementary file. This adapter provided a stable and optimal imaging surface

for live microscopy of biofilm of *E. coli*, *P. aeruginosa* and *S. aureus*, which was performed employing

a DMi 8 inverted microscope (Leica, Germany) equipped with a 100X oil immersion objective (Fig.

3B).

Coupling of melittin to the agarose-gelatine beads. Using the same procedure that is employed for affinity chromatography of crosslinked agarose media, we modified the method described by March et al [99] to couple melittin with the agarose beads. The washed agarose beads, as described in a previous section of this manuscript, were first activated with cyanogen bromide. A volume of 500 µl of coupling buffer (0.1 M Sodium bicarbonate, pH 9.5 in distilled water) along with the amino ligand to be coupled, in our case melittin (5 mg/ml, Genescript, USA) and Bovine serum albumin (BSA) (1 mg/ml, Merck) was used per bead. The coupling reaction was carried out for 20 hours at 4°C. Thereafter, the beads were washed with 20 volumes of buffer containing 0.1 M sodium acetate, pH 4, 2 M urea and 0.1 M sodium bicarbonate, pH 10. The beads were then dialysed against 100 volumes of sterile NaCl 9% solution and stored at 4°C. All operations were made under sterile conditions under a laminar flow with filtered sterilised buffer to prevent bead contamination.

Staphyloxanthin accumulation. *S. aureus* SH1000 biofilms growing in AGBs or glass beads were grown in 2 ml of M9 medium with moderate shaking at 37°C for 48 hours with a medium change every 24 hours. The M9 medium (Merck, Germany) used for this experiment was supplemented with 0.4% casamino acids (Carl Roth, Germany), 5 mM MgSO4 (Carl Roth, Germany), and 0.5% glucose as carbon sources (final concentrations). Glucose and magnesium salt were added after autoclaving. Cultures were centrifuged for 1 minute at 10,000 x g, and the supernatant was removed. Each bead was washed twice with 1 ml of NaCl 0.9% solution and added 500 µL of methanol and incubated at 55°C for 30 minutes. Tubes were centrifuged for 2 minutes at 10,000 x g. 300 µL from each tube was added to each well of a polystyrene 96-well plate and then quantified at OD_{450} on a GloMax Explorer plate reader (Promega, USA). Five biological replicates were performed for each group. For normalisation, all OD_{450} values were divided by the CFU count of each bead. Normalised production of staphyloxanthin was expressed as a ratio of OD_{450}/CFU, and AGBs and glass beads were compared.

Quantification of the virulence factors pyocyanin and pyoverdine. To investigate how the virulence of *P. aeruginosa* PAO1 is affected by different substrates, we collected the bacterial supernatant from six biofilm-containing samples on glass or AGBs from all wells of a 24-well plate. The collected supernatant was transferred into 15 ml Falcon tubes and then centrifuged at 4,000 x g for 15 minutes at 20°C. After centrifugation, the resulting supernatant was separated from the bacterial pellets and used for further purification and quantification of pyocyanin concentration. To extract Pyocyanin, we followed a method with some modifications. We incubated 5 ml of the supernatant

from each sample with 1 ml chloroform for 10 minutes, with vortexing in between. This process

formed an organic phase (blue colour) at the bottom of the tubes. We then pipetted 900 μl from the

organic phase and transferred it into 1.5 ml microcentrifuge tubes. After centrifugation for 1 minute

at 10,000 x g to remove impurities, 700 μL of the organic phase was transferred to a new

microcentrifuge tube and further extracted with 700 μl 0.1N HCl solution. From each sample, 7x100

μl of the aqueous phase formed at the top (red colour) was transferred to a polystyrene 96-well

plate. Absorbance was measured at OD_{520}, with the 0.1N HCl solution used as a blank. We

performed five experimental replicates for each group. To normalise the results, we divided all

measured values by each experiment's average group CFU counts. The normalised production of

pyocyanin was expressed as a ratio of the measured values to CFU and then compared. For

pyoverdine, we used an identical setup as for pyocyanin, with identical culture conditions and

number of replications. The 24-hour cultures' supernatants were centrifuged at 10,000 x g for 10

minutes, and the pigment level was determined by measuring its fluorescence, with excitation at

405 nm and emission at 450 nm.

AHL level determination. The AHL activities of the *P. aeruginosa* PAO1 were assessed using a

previously described colourimetric method [74]. Briefly, the bacterial isolates were cultured

overnight in 5 mL of sterile Muller-Hinton broth (Merck, Germany) at 37 ºC. Then, 1.5 ml of the

overnight cultures were aseptically transferred into sterile centrifuge tubes (Eppendorf, Germany)

and centrifuged at 10,000 x g for 15 min. The cell pellets were discarded, and this process was

repeated three times. The resulting supernatants were filtered through 0.2 μm membrane filters

(Sartorius, Germany) to remove any cell debris. The filtrates were mixed with ethyl acetate and

shaken for 10 min. Afterwards, the mixture was left to stand for five minutes in a separating funnel,

resulting in the formation of an upper organic layer and a lower aqueous layer. The upper layer was

collected in sterile tubes, while the lower portion was subjected to two additional extractions

following the same procedure. The collected upper portions from each sample were combined and

dried in an oven at 40°C. For liquid-liquid extraction (LLE) [100], 40 μL of each dried extract was

prepared and inoculated into 96-well polystyrene flat-bottom tissue culture microplate wells. Then,

50 μL of a 1:1 mixture of hydroxylamine (2 M) and NaOH (3.5 M) was added to each well and mixed

with the sample. Subsequently, an equal amount of a 1:1 mixture of ferric chloride (10% in 4 M HCl)

and 95% ethanol was added. The OD was measured at 520 nm. The AHL extracts were concentrated

by drying in an oven at 40°C overnight and stored at -20°C for further analysis. The presence of

lactone compounds was indicated by a dark brown colour in all samples, although in some cases,

repetitive pipetting of the mixtures caused a colour change to yellow, depending on the lactone

concentrations. To ensure that an acidic pH was maintained, we monitored the pH throughout the

experiment, as under alkaline conditions, AHLs undergo rapid inactivation through pH-dependent

lactonolysis. This reaction hydrolyses the homoserine lactone ring, resulting in the open-ring form

corresponding to N-acyl-homoserine. It's worth noting that this lactonolysis reaction can be

reversed by acidification.

Agarose-gelatine stiffness measurement. The stiffness of agarose-gelatine beads, both with and

without crosslinking, was measured using a static compression test described elsewhere [101]. The

beads (4 mm) were directly used for the measurement. Then, beads were placed at room

temperature overnight for temperature equilibration before the mechanical testing. Young's

modulus was determined by fitting a line to the linear region of the stress-strain curve. Five samples

were used for each measurement.

Sample preparation for proteomics of bacterial biofilms. Each experimental condition consisted of

six replicates. A volume of 500 µl of denaturation buffer (6M urea/2 M thiourea in 10 mM HEPES pH

8.0) was added into each sample consisting of three beads of glass, agarose-gelatine and polystyrene

substrate from multiwell plates. The samples went for five freeze-thaw cycles to cause lysis and

protein release. The resulting lysate was used for in-solution protein digestion as described

previously [102]. Briefly, proteins, re-suspended in denaturation buffer, were reduced by the

addition of 1 µl of 10 mM DTT dissolved in 50 mM ammonium bicarbonate (ABC) and incubated for

30 minutes, followed by a 20-minute alkylation reaction with 1 µl of 55 mM iodoacetamide. As a first

digestion step, Lysyl endopeptidase (LysC, Wako, Japan) resuspended in 50 mM ABC was added to

each tube in a ratio of 1 µg per 50 µg of total proteins and incubated for 3 hours. After pre-digestion

with LysC, protein samples were diluted four times with 50 mM ABC and subjected to overnight

trypsin digestion using 1 µg/reaction of sequencing grade modified trypsin (Promega, USA), also

diluted before use in 50 mM ABC. All in-solution protein digestion steps were performed at room

temperature. After adding iodoacetamide, the samples were protected from the light until the

digestion was stopped by acidification, adding 5% acetonitrile and 0.3% trifluoroacetic acid (final

concentrations). The samples were micro-purified and concentrated using the Stage-tip protocol

[102], and the eluates were vacuum-dried. For a step-by-step protocol, see Protocol IV of

supplementary materials.

Nano liquid chromatography-mass spectrometry (LC-MS) and data analysis. Peptides were

reconstituted in 30 µl of 0.05% trifluoroacetic acid, 4% acetonitrile in water, and 2 µl were analysed

by an Ultimate 3000 reversed-phase capillary nano liquid chromatography system connected to a Q

Exactive HF mass spectrometer (Thermo Fisher Scientific). Samples were injected and concentrated
on a trap column (PepMap100 C18, 3 µm, 100 Å, 75 µm i.d. x 2 cm, Thermo Fisher Scientific)
equilibrated with 0.05% TFA in water. After switching the trap column inline, LC separations were
performed on a capillary column (Acclaim PepMap100 C18, 2 µm, 100 Å, 75 µm i.d. x 25 cm, Thermo
Fisher Scientific) at an eluent flow rate of 300 nl/min. Mobile phase A contained 0.1% formic acid in
water, and mobile phase B contained 0.1% formic acid in 80% acetonitrile/20% water. The column
was pre-equilibrated with 5% mobile phase B followed by an increase of 5 - 44% mobile phase B in
35 min. Mass spectra were acquired in a data-dependent mode utilising a single MS survey scan (m/z
300–1650) with a resolution of 60,000 and MS/MS scans of the 15 most intense precursor ions with
a resolution of 15,000. The dynamic exclusion time was set to 20 seconds, and automatic gain
control was set to $3x10^6$ and $1x10^5$ for MS and MS/MS scans, respectively.

MS and MS/MS raw data were analysed using the MaxQuant software package (version 2.0.3.0) with
the implemented Andromeda peptide search engine [103]. Data were searched against the
reference protein sequence databases downloaded from UniProt: *E. coli* (4448 proteins, Proteome
ID UP000000625), *P. aeruginosa* (5564 proteins, Proteome ID UP000002438), and *S. aureus* (2889
proteins, Proteome ID UP000008816). Filtering and statistical analysis were performed using the
software Perseus version 1.6.14 [104]. Missing values are imputed using the default settings. Mean
log2 fold protein LFQ intensity differences between experimental groups (glass beads vs. AGBs) were
calculated in Perseus using student's t-tests, and *p*-values were adjusted via false discovery rate
(FDR) of 0.05 as a cut-off.

Statistical analysis. The student's t-test was used for pair comparisons if sample data followed a
normal distribution and had homogenous variance. Otherwise, the Mann-Whitney U-test was used
instead. For multiple comparisons, we used one-way ANOVA with the Holm-Šídák test for parametric
data or the Kruskal-Wallis test for non-parametric ones. All statistical tests were done in GraphPad
(Prism, version 9) or R version 4.3.3 [105].

Acknowledgements: We want to thank the assistance of the Core Facility BioSupraMol (Freie
Universität Berlin), which was supported by the Deutsche Forschungsgemeinschaft (DFG).

Author Contribution: The authors confirm their contribution to the paper as follows: study
conception and design: DR and ARR. Data collection and analysis: MH, DR, AN, BK, NP, ARR. Analysis
and interpretation of results: MH, DR, AN, BK, NP, JR and ARR. Draft manuscript preparation: DR, MH,
AN, ARR. Manuscript editing and correction: all authors. Supervision: JR and ARR. Funding and

resources: JR and ARR. All authors reviewed the results and approved the final version of the

manuscript.

Data Availability Statement: The data, including the raw data files are available within the

manuscript and the supplementary information files.

Funding: This research was part of a multi-institutional project funded by the Volkswagen Stiftung for

DR. We also thank the OneHealth PhD program funded by Vetmeduni, WWTF Vienna to ARR, and the

program Erasmus-Mundus-Master to NP and ARR.

Competing interests: The authors have declared that no competing interests exist.

Abbreviations: DVS, divinyl sulphone; AGBs, agarose-gelatine beads; CFU, colony-forming units; SEM,

scanning electron microscopy; LC-MS, liquid chromatography mass spectrometry; OD, optical

density; GFP, green fluorescent protein.

References
769
1.   Stoodley P, Sauer K, Davies DG, Costerton JW. Biofilms as complex differentiated
communities.   Annu   Rev   Microbiol.   2002;56:   187–209.
doi:10.1146/ANNUREV.MICRO.56.012302.160705
2.   Römling U. Is biofilm formation intrinsic to the origin of life? Environ Microbiol. 2023;25: 26–
39. doi:10.1111/1462-2920.16179
3.   Bar-On YM, Milo R. Towards a quantitative view of the global ubiquity of biofilms. Nature
Reviews Microbiology 2019 17:4. 2019;17: 199–200. doi:10.1038/s41579-019-0162-0
4.   Flemming H-C, Wuertz S. Bacteria and archaea on Earth and their abundance in biofilms. Nat
Rev Microbiol. 2019;17: 247–260. doi:10.1038/s41579-019-0158-9
5.   Bjarnsholt T. The role of bacterial biofilms in chronic infections. APMIS Suppl. 2013; 1–51.
doi:10.1111/apm.12099
6.   Del Pozo JL. Biofilm-related disease. Expert Rev Anti Infect Ther. 2018;16: 51–65.
doi:10.1080/14787210.2018.1417036
7.   Høiby N, Bjarnsholt T, Moser C, Bassi GL, Coenye T, Donelli G, et al. ESCMID guideline for the
diagnosis and treatment of biofilm infections 2014. Clin Microbiol Infect. 2015;21 Suppl 1: S1-
25. doi:10.1016/j.cmi.2014.10.024
8.   Azeredo J, Azevedo NF, Briandet R, Cerca N, Coenye T, Costa AR, et al. Critical review on
biofilm   methods.   https://doi.org/101080/1040841X20161208146.   2016;43:   313–351.
doi:10.1080/1040841X.2016.1208146
9.   Fletcher M. The effects of culture concentration and age, time, and temperature on bacterial
attachment   to   polystyrene.   https://doi.org/101139/m77-001.   2011;23:   1–6.
doi:10.1139/M77-001
10.   Kharazmi A, Giwercman B, Høiby N. Robbins device in biofilm research. Methods Enzymol.
1999;310: 207–15. doi:10.1016/s0076-6879(99)10018-1
11.   Azeredo J, Azevedo NF, Briandet R, Cerca N, Coenye T, Costa AR, et al. Critical review on
biofilm   methods.   https://doi.org/101080/1040841X20161208146.   2016;43:   313–351.
doi:10.1080/1040841X.2016.1208146

| 797 | 12. | Tolker-Nielsen T, Sternberg C. Methods for studying biofilm formation: flow cells and confocal laser scanning microscopy. Methods Mol Biol. 2014;1149: 615–29. doi:10.1007/978-1-4939-0473-0_47 |

12. Tolker-Nielsen T, Sternberg C. Methods for studying biofilm formation: flow cells and confocal
laser scanning microscopy. Methods Mol Biol. 2014;1149: 615–29. doi:10.1007/978-1-4939-
0473-0_47

13. Gomes IB, Meireles A, Gonçalves AL, Goeres DM, Sjollema J, Simões LC, et al. Standardized
reactors for the study of medical biofilms: a review of the principles and latest modifications.
https://doi.org/101080/073885512017380601.   2017;38:   657–670.
doi:10.1080/07388551.2017.1380601

14. Vyas HKN, Xia B, Mai-Prochnow A. Clinically relevant in vitro biofilm models: A need to mimic
and   recapitulate   the   host   environment.   Biofilm.   2022;4:   100069.
doi:10.1016/J.BIOFLM.2022.100069

15. Austin B. The value of cultures to modern microbiology. Antonie Van Leeuwenhoek.
2017;110: 1247–1256. doi:10.1007/s10482-017-0840-8

16. Das N, Triparthi N, Basu S, Bose C, Maitra S, Khurana S. Progress in the development of gelling
agents for improved culturability of microorganisms. Front Microbiol. 2015;6: 698.
doi:10.3389/FMICB.2015.00698

17. Koch R. Über den augenblicklichen Stand der bacteriologischen Choleradiagnose. Deutsche
Medizinische Wochenschrift. 1893;19: 523–525. doi:10.1055/s-0028-1143698

18. World Health Organization. WHO Bacterial Priority Pathogens List, 2024: bacterial pathogens
of public health importance to guide research, development and strategies to prevent and
control   antimicrobial   resistance.   Geneva;   2024   May.   Available:
https://www.who.int/publications/i/item/9789240093461

19. EUCAST. The European Committee on Antimicrobial Susceptibility Testing. In: Breakpoint
tables for interpretation of MICs and zone diameters, Version 14.0 [Internet]. 1 Jan 2024.
Available: https://www.eucast.org/clinical_breakpoints

20. Hjertén S. The preparation of agarose spheres for chromatography of molecules and
particles. Biochimica et Biophysica Acta (BBA) - Specialized Section on Biophysical Subjects.
1964;79: 393–398. doi:10.1016/0926-6577(64)90020-8

21. Porath J, Låås T, Janson JC. Agar derivatives for chromatography, electrophoresis and gel-
bound enzymes. III. Rigid agarose gels cross-linked with divinyl sulphone (dvs). J Chromatogr.
1975;103: 49–62. doi:10.1016/s0021-9673(00)83800-5

22. Cash HA, Woods DE, McCullough B, Johanson WG, Bass JA. A rat model of chronic respiratory
infection with Pseudomonas aeruginosa. Am Rev Respir Dis. 1979;119: 453–459.
doi:10.1164/ARRD.1979.119.3.453

23. Facchini M, De Fino I, Riva C, Bragonzi A. Long term chronic Pseudomonas aeruginosa airway
infection in mice. J Vis Exp. 2014. doi:10.3791/51019

24. Rodríguez-Rojas A, Maciá MDMD, Couce A, Gómez C, Castañeda-García A, Oliver A, et al.
Assessing the emergence of resistance: the absence of biological cost in vivo may
compromise fosfomycin treatments for P. aeruginosa infections. PLoS One. 2010/04/27.
2010;5: e10193. doi:10.1371/journal.pone.0010193

25. Konrat K, Schwebke I, Laue M, Dittmann C, Levin K, Andrich R, et al. The Bead Assay for
Biofilms: A Quick, Easy and Robust Method for Testing Disinfectants. PLoS One. 2016;11.
doi:10.1371/JOURNAL.PONE.0157663

26. Roizman D, Vidaillac C, Givskov M, Yang L. In Vitro Evaluation of Biofilm Dispersal as a
Therapeutic Strategy To Restore Antimicrobial Efficacy. Antimicrob Agents Chemother.
2017;61. doi:10.1128/AAC.01088-17

27. Er B, Demirhan B, Onurdaꞷ FK, Özgacar SÖ, Öktem AB. Antimicrobial and antibiofilm effects
of selected food preservatives against Salmonella spp. isolated from chicken samples. Poult
Sci. 2014;93: 695–701. doi:10.3382/PS.2013-03404

28. Heukelekian H, Heller A. Relation between Food Concentration and Surface for Bacterial
Growth. J Bacteriol. 1940;40: 547. doi:10.1128/JB.40.4.547-558.1940

29.   Zobell CE. The Effect of Solid Surfaces upon Bacterial Activity. J Bacteriol. 1943;46: 39.
doi:10.1128/JB.46.1.39-56.1943

30.   Miller WR, Arias CA. ESKAPE pathogens: antimicrobial resistance, epidemiology, clinical
impact and therapeutics. Nat Rev Microbiol. 2024. doi:10.1038/s41579-024-01054-w

31.   Murray CJ, Shunji Ikuta K, Sharara F, Swetschinski L, Robles Aguilar G, Gray A, et al. Global
burden of bacterial antimicrobial resistance in 2019: a systematic analysis. The Lancet.
2022;0. doi:10.1016/S0140-6736(21)02724-0

32.   Browning DF, Hobman JL, Busby SJW. Laboratory strains of Escherichia coli K-12: things are
seldom what they seem. Microb Genom. 2023;9. doi:10.1099/mgen.0.000922

33.   Mackie A, Paley S, Keseler IM, Shearer A, Paulsen IT, Karp PD. Addition of Escherichia coli K-12
growth observation and gene essentiality data to the EcoCyc database. J Bacteriol. 2014;196:
982–8. doi:10.1128/JB.01209-13

34.   Richter AM, Konrat K, Osland AM, Brook E, Oastler C, Vestby LK, et al. Evaluation of Biofilm
Cultivation Models for Efficacy Testing of Disinfectants against Salmonella Typhimurium
Biofilms. Microorganisms. 2023;11. doi:10.3390/microorganisms11030761

35.   Rodríguez-Rojas A, Nath A, El Shazely B, Santi G, Kim JJ, Weise C, et al. Antimicrobial Peptide
Induced-Stress Renders Staphylococcus aureus Susceptible to Toxic Nucleoside Analogs.
Front Immunol. 2020;11: 1686. doi:10.3389/fimmu.2020.01686

36.   Rodríguez-Rojas A, Kim JJ, Johnston PR, Makarova O, Eravci M, Weise C, et al. Non-lethal
exposure to H2O2 boosts bacterial survival and evolvability against oxidative stress. PLoS
Genet. 2020;16: e1008649. doi:10.1371/journal.pgen.1008649

37.   Prossliner T, Skovbo Winther K, Sørensen MA, Gerdes K. Ribosome Hibernation. Annu Rev
Genet. 2018;52: 321–348. doi:10.1146/annurev-genet-120215-035130

38.   Kimura S, Suzuki T. Fine-tuning of the ribosomal decoding center by conserved methyl-
modifications in the Escherichia coli 16S rRNA. Nucleic Acids Res. 2010;38: 1341–52.
doi:10.1093/nar/gkp1073

39.   Brandi A, Giangrossi M, Paoloni S, Spurio R, Giuliodori AM, Pon CL, et al. Transcriptional and
post-transcriptional events trigger de novo infB expression in cold stressed Escherichia coli.
Nucleic Acids Res. 2019;47: 4638–4651. doi:10.1093/nar/gkz187

40.   Krupka M, Sobrinos-Sanguino M, Jiménez M, Rivas G, Margolin W. Escherichia coli ZipA
Organizes FtsZ Polymers into Dynamic Ring-Like Protofilament Structures. mBio. 2018;9.
doi:10.1128/mBio.01008-18

41.   Rust L, Pesci EC, Iglewski BH. Analysis of the Pseudomonas aeruginosa elastase (lasB)
regulatory region. J Bacteriol. 1996;178: 1134–1140. doi:10.1128/JB.178.4.1134-1140.1996

42.   Axelrad I, Safrin M, Cahan R, Suh S-J, Ohman DE, Kessler E. Extracellular proteolytic activation
of Pseudomonas aeruginosa aminopeptidase (PaAP) and insight into the role of its non-
catalytic   N-terminal   domain.   PLoS   One.   2021;16:   e0252970.
doi:10.1371/journal.pone.0252970

43.   Dell'Anno F, Vitale GA, Buonocore C, Vitale L, Palma Esposito F, Coppola D, et al. Novel
Insights on Pyoverdine: From Biosynthesis to Biotechnological Application. Int J Mol Sci.
2022;23. doi:10.3390/IJMS231911507

44.   Parsons JF, Greenhagen BT, Shi K, Calabrese K, Robinson H, Ladner JE. Structural and
functional analysis of the pyocyanin biosynthetic protein PhzM from Pseudomonas
aeruginosa. Biochemistry. 2007;46: 1821–8. doi:10.1021/bi6024403

45.   Ramsey DM, Baynham PJ, Wozniak DJ. Binding of Pseudomonas aeruginosa AlgZ to sites
upstream of the algZ promoter leads to repression of transcription. J Bacteriol. 2005;187:
4430–4443. doi:10.1128/JB.187.13.4430-4443.2005

46.   Kalia M, Resch MD, Cherny KE, Sauer K. The Alginate and Motility Regulator AmrZ is Essential
for the Regulation of the Dispersion Response by Pseudomonas aeruginosa Biofilms.
mSphere. 2022;7: e0050522. doi:10.1128/msphere.00505-22

47.   L T, I C. Pushing the Envelope: The Mysterious Journey Through the Bacterial Secretory
898   Machinery, and Beyond. Front Microbiol. 2021;12. doi:10.3389/FMICB.2021.782900

899   48.   Fortuna A, Bähre H, Visca P, Rampioni G, Leoni L. The two Pseudomonas aeruginosa DksA
900   stringent response proteins are largely interchangeable at the whole transcriptome level and
901   in the control of virulence-related traits. Environ Microbiol. 2021;23: 5487–5504.
902   doi:10.1111/1462-2920.15693

903   49.   Lomax MS, Greenberg GR. Characteristics of the deo operon: role in thymine utilization and
904   sensitivity to deoxyribonucleosides. J Bacteriol. 1968;96: 501–14. doi:10.1128/jb.96.2.501-
905   514.1968

906   50.   Sanchez-Herrero JF, Bernabeu M, Prieto A, Hüttener M, Juárez A. Gene Duplications in the
907   Genomes of Staphylococci and Enterococci. Front Mol Biosci. 2020;7: 160.
908   doi:10.3389/fmolb.2020.00160

909   51.   Baum KR, Ahmad Z, Singh VK. Regulation of Expression of Oxacillin-Inducible Methionine
910   Sulfoxide Reductases in Staphylococcus aureus. Int J Microbiol. 2015;2015.
911   doi:10.1155/2015/617925

912   52.   Koprivnjak T, Mlakar V, Swanson L, Fournier B, Peschel A, Weiss JP. Cation-induced
913   transcriptional regulation of the dlt operon of Staphylococcus aureus. J Bacteriol. 2006;188:
914   3622–30. doi:10.1128/JB.188.10.3622-3630.2006

915   53.   Nagel A, Michalik S, Debarbouille M, Hertlein T, Gesell Salazar M, Rath H, et al. Inhibition of
916   Rho Activity Increases Expression of SaeRS-Dependent Virulence Factor Genes in
917   Staphylococcus aureus, Showing a Link between Transcription Termination, Antibiotic Action,
918   and Virulence. mBio. 2018;9. doi:10.1128/mBio.01332-18

919   54.   Jenul C, Horswill AR. Regulation of Staphylococcus aureus Virulence. Microbiol Spectr.
920   2019;7. doi:10.1128/microbiolspec.GPP3-0031-2018

921   55.   Albanesi D, de Mendoza D. FapR: From Control of Membrane Lipid Homeostasis to a
922   Biotechnological Tool. Front Mol Biosci. 2016;3: 64. doi:10.3389/fmolb.2016.00064

923   56.   Ravcheev DA, Li X, Latif H, Zengler K, Leyn SA, Korostelev YD, et al. Transcriptional regulation
924   of central carbon and energy metabolism in bacteria by redox-responsive repressor Rex. J
925   Bacteriol. 2012;194: 1145–57. doi:10.1128/JB.06412-11

926   57.   Kiran K, Patil KN. Characterization of Staphylococcus aureus RecX protein: Molecular insights
927   into negative regulation of RecA protein and implications in HR processes. J Biochem.
928   2023;174: 227–237. doi:10.1093/jb/mvad039

929   58.   Hübscher J, McCallum N, Sifri CD, Majcherczyk PA, Entenza JM, Heusser R, et al. MsrR
930   contributes to cell surface characteristics and virulence in Staphylococcus aureus. FEMS
931   Microbiol Lett. 2009;295: 251–60. doi:10.1111/j.1574-6968.2009.01603.x

932   59.   Pamp SJ, Frees D, Engelmann S, Hecker M, Ingmer H. Spx is a global effector impacting stress
933   tolerance and biofilm formation in Staphylococcus aureus. J Bacteriol. 2006;188: 4861–70.
934   doi:10.1128/JB.00194-06

935   60.   Makhlin J, Kofman T, Borovok I, Kohler C, Engelmann S, Cohen G, et al. Staphylococcus aureus
936   ArcR controls expression of the arginine deiminase operon. J Bacteriol. 2007;189: 5976–86.
937   doi:10.1128/JB.00592-07

938   61.   Yang X, Luo MJ, Yeung ACM, Lewis PJ, Chan PKS, Ip M, et al. First-In-Class Inhibitor of
939   Ribosomal RNA Synthesis with Antimicrobial Activity against Staphylococcus aureus.
940   Biochemistry. 2017;56: 5049–5052. doi:10.1021/acs.biochem.7b00349

941   62.   Roux A, Todd DA, Velázquez J V, Cech NB, Sonenshein AL. CodY-mediated regulation of the
942   Staphylococcus aureus Agr system integrates nutritional and population density signals. J
943   Bacteriol. 2014;196: 1184–96. doi:10.1128/JB.00128-13

944   63.   Miyazaki E, Chen JM, Ko C, Bishai WR. The Staphylococcus aureus rsbW (orf159) gene
945   encodes an anti-sigma factor of SigB. J Bacteriol. 1999;181: 2846–51.
946   doi:10.1128/JB.181.9.2846-2851.1999

64. Tajbakhsh G, Golemi-Kotra D. The dimerization interface in VraR is essential for induction of
the cell wall stress response in Staphylococcus aureus: a potential druggable target. BMC
Microbiol. 2019;19: 153. doi:10.1186/s12866-019-1529-0
65. Haag AF, Bagnoli F. The Role of Two-Component Signal Transduction Systems in
Staphylococcus aureus Virulence Regulation. Curr Top Microbiol Immunol. 2017;409: 145–
198. doi:10.1007/82_2015_5019
66. Hardie KR, Heurlier K. Establishing bacterial communities by "word of mouth": LuxS and
autoinducer 2 in biofilm development. Nat Rev Microbiol. 2008;6: 635–43.
doi:10.1038/nrmicro1916
67. Burts ML, Williams WA, DeBord K, Missiakas DM. EsxA and EsxB are secreted by an ESAT-6-
like system that is required for the pathogenesis of Staphylococcus aureus infections. Proc
Natl Acad Sci U S A. 2005;102: 1169–74. doi:10.1073/pnas.0405620102
68. Pratt LA, Kolter R. Genetic analysis of Escherichia coli biofilm formation: roles of flagella,
motility, chemotaxis and type I pili. Mol Microbiol. 1998;30: 285–293. doi:10.1046/J.1365-
2958.1998.01061.X
69. Rodriguez-Rojas A, Blazquez J, Rodríguez-Rojas A, Blázquez J. The Pseudomonas aeruginosa
pfpI gene plays an antimutator role and provides general stress protection. J Bacteriol.
2008/11/26. 2009;191: 844–850. doi:JB.01081-08 [pii] 10.1128/JB.01081-08
70. Baracchi D, Francese S, Turillazzi S. Beyond the antipredatory defence: honey bee venom
function as a component of social immunity. Toxicon. 2011;58: 550–557.
doi:10.1016/J.TOXICON.2011.08.017
71. March SC, Parikh I, Cuatrecasas P. A simplified method for cyanogen bromide activation of
agarose for affinity chromatography. Anal Biochem. 1974;60: 149–152. doi:10.1016/0003-
2697(74)90139-0
72. O'Toole GA, Kolter R. Initiation of biofilm formation in Pseudomonas fluorescens WCS365
proceeds via multiple, convergent signalling pathways: a genetic analysis. Mol Microbiol.
1998/06/19. 1998;28: 449–461.
73. Fekete-Kertész I, Berkl Z, Buda K, Fenyvesi É, Szente L, Molnár M. Quorum quenching effect of
cyclodextrins on the pyocyanin and pyoverdine production of Pseudomonas aeruginosa. Appl
Microbiol Biotechnol. 2024;108: 271. doi:10.1007/s00253-024-13104-7
74. Yang YH, Lee TH, Kim JH, Kim EJ, Joo HS, Lee CS, et al. High-throughput detection method of
quorum-sensing molecules by colorimetry and its applications. Anal Biochem. 2006;356: 297–
299. doi:10.1016/J.AB.2006.05.030
75. Terwilligert TC, Eisenbergg D. The structure of melittin. II. Interpretation of the structure.
Journal of Biological Chemistry. 1982;257: 6016–6022. doi:10.1016/S0021-9258(20)65098-0
76. Ownby CL, Powell JR, Jiang MS, Fletcher JE. Melittin and phospholipase A2 from bee (Apis
mellifera) venom cause necrosis of murine skeletal muscle in vivo. Toxicon. 1997;35: 67–80.
doi:10.1016/S0041-0101(96)00078-5
77. Mukhi M, Vishwanathan AS. Beneficial Biofilms: a Minireview of Strategies To Enhance
Biofilm Formation for Biotechnological Applications. Appl Environ Microbiol. 2022;88.
doi:10.1128/AEM.01994-21
78. Kreve S, Reis ACD. Bacterial adhesion to biomaterials: What regulates this attachment? A
review. Jpn Dent Sci Rev. 2021;57: 85–96. doi:10.1016/J.JDSR.2021.05.003
79. Lang S, Livesley MA, Lambert PA, Littler WA, Elliott TSJ. Identification of a novel antigen from
Staphylococcus epidermidis. FEMS Immunol Med Microbiol. 2000;29: 213–220.
doi:10.1111/J.1574-695X.2000.TB01525.X
80. Clauditz A, Resch A, Wieland KP, Peschel A, Götz F. Staphyloxanthin plays a role in the fitness
of Staphylococcus aureus and its ability to cope with oxidative stress. Infect Immun. 2006;74:
4950–4953.   doi:10.1128/IAI.00204-06/ASSET/0B500F87-F8F6-43EC-8C7E-
02D1C33F109E/ASSETS/GRAPHIC/ZII0080661110005.JPEG

81.   Campbell AE, Mccready-Vangi AR, Uberoi A, Planet PJ, Gardner SE, Grice Correspondence EA.
Variable staphyloxanthin production by Staphylococcus aureus drives strain-dependent
effects on diabetic wound-healing outcomes. Cell Rep. 2023;42: 113281.
doi:10.1016/j.celrep.2023.113281

82.   Perez-Lopez MI, Mendez-Reina R, Trier S, Herrfurth C, Feussner I, Bernal A, et al. Variations in
carotenoid content and acyl chain composition in exponential, stationary and biofilm states
of Staphylococcus aureus, and their influence on membrane biophysical properties. Biochim
Biophys Acta Biomembr. 2019;1861: 978–987. doi:10.1016/J.BBAMEM.2019.02.001

83.   Pratt LA, Kolter R. Genetic analysis of Escherichia coli biofilm formation: roles of flagella,
motility, chemotaxis and type I pili. Mol Microbiol. 1998;30: 285–293. doi:10.1046/J.1365-
2958.1998.01061.X

84.   Belas R. Biofilms, flagella, and mechanosensing of surfaces by bacteria. Trends Microbiol.
2014;22: 517–27. doi:10.1016/j.tim.2014.05.002

85.   Blattner FR, Plunkett G, Bloch CA, Perna NT, Burland V, Riley M, et al. The complete genome
sequence of Escherichia coli K-12. Science. 1997;277: 1453–1462.
doi:10.1126/SCIENCE.277.5331.1453

86.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, et al. Construction of Escherichia
coli K-12 in-frame, single-gene knockout mutants: the Keio collection. Mol Syst Biol.
2006/06/02. 2006;2: 2006 0008. doi:msb4100050 [pii] 10.1038/msb4100050

87.   Stover CK, Pham XQ, Erwin AL, Mizoguchi SD, Warrener P, Hickey MJ, et al. Complete genome
sequence of Pseudomonas aeruginosa PA01, an opportunistic pathogen. Nature. 2000/09/13.
2000;406: 959–964. doi:10.1038/35023079

88.   Liberati NT, Urbach JM, Miyata S, Lee DG, Drenkard E, Wu G, et al. An ordered, nonredundant
library of Pseudomonas aeruginosa strain PA14 transposon insertion mutants. Proc Natl Acad
Sci U S A. 2006/02/16. 2006;103: 2833–2838. doi:0511100103 [pii]
10.1073/pnas.0511100103

89.   Horsburgh MJ, Aish JL, White IJ, Shaw L, Lithgow JK, Foster SJ. sigmaB modulates virulence
determinant expression and stress resistance: characterization of a functional rsbU strain
derived from Staphylococcus aureus 8325-4. J Bacteriol. 2002;184: 5457–67.
doi:10.1128/JB.184.19.5457-5467.2002

90.   Yang DD, Alexander A, Kinnersley M, Cook E, Caudy A, Rosebrock A, et al. Fitness and
Productivity Increase with Ecotypic Diversity among Escherichia coli Strains That Coevolved in
a Simple, Constant Environment. Appl Environ Microbiol. 2020;86. doi:10.1128/AEM.00051-
20

91.   Nivens DE, Ohman DE, Williams J, Franklin MJ. Role of alginate and its O acetylation in
formation of Pseudomonas aeruginosa microcolonies and biofilms. J Bacteriol. 2001;183:
1047–57. doi:10.1128/JB.183.3.1047-1057.2001

92.   de Jong NWM, van der Horst T, van Strijp JAG, Nijland R. Fluorescent reporters for markerless
genomic integration in Staphylococcus aureus. Sci Rep. 2017;7: 43889.
doi:10.1038/srep43889

93.   Dagert M, Ehrlich SD. Prolonged incubation in calcium chloride improves the competence of
Escherichia coli cells. Gene. 1979;6: 23–8. doi:10.1016/0378-1119(79)90082-9

94.   Schenk S, Laddaga RA. Improved method for electroporation of Staphylococcus aureus. FEMS
Microbiol Lett. 1992;73: 133–8. doi:10.1016/0378-1097(92)90596-g

95.   de Lorenzo V, Eltis L, Kessler B, Timmis KN. Analysis of Pseudomonas gene products using
lacIq/Ptrp-lac plasmids and transposons that confer conditional phenotypes. Gene. 1993;123:
17–24. doi:10.1016/0378-1119(93)90533-9

96.   O'Toole GA, Kolter R. Flagellar and twitching motility are necessary for Pseudomonas
aeruginosa biofilm development. Mol Microbiol. 1998/10/29. 1998;30: 295–304. Available:
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&
list_uids=9791175

97. Moskowitz SM, Foster JM, Emerson J, Burns JL. Clinically Feasible Biofilm Susceptibility Assay for Isolates of Pseudomonas aeruginosa from Patients with Cystic Fibrosis. J Clin Microbiol. 2004;42: 1915. doi:10.1128/JCM.42.5.1915-1922.2004

98. Boyles J, Anderson L, Hutcherson P. A new fixative for the preservation of actin filaments: fixation of pure actin filament pellets. J Histochem Cytochem. 1985;33: 1116–1128. doi:10.1177/33.11.3902963

99. March SC, Parikh I, Cuatrecasas P. A simplified method for cyanogen bromide activation of agarose for affinity chromatography. Anal Biochem. 1974;60: 149–152. doi:10.1016/0003-2697(74)90139-0

100. Wang J, Quan C, Wang X, Zhao P, Fan S. Extraction, purification and identification of bacterial signal molecules based on N-acyl homoserine lactones. Microb Biotechnol. 2011;4: 479–90. doi:10.1111/j.1751-7915.2010.00197.x

101. Lu C, Zheng J, Yoshitomi T, Kawazoe N, Yang Y, Chen G. How Hydrogel Stiffness Affects Adipogenic Differentiation of Mesenchymal Stem Cells under Controlled Morphology. ACS Appl Bio Mater. 2023;6: 3441–3450. doi:10.1021/ACSABM.3C00159

102. Rappsilber J, Mann M, Ishihama Y. Protocol for micro-purification, enrichment, pre-fractionation and storage of peptides for proteomics using StageTips. Nat Protoc. 2007;2: 1896–906. doi:10.1038/nprot.2007.261

103. Tyanova S, Temu T, Cox J. The MaxQuant computational platform for mass spectrometry-based shotgun proteomics. Nat Protoc. 2016;11: 2301–2319. doi:10.1038/nprot.2016.136

104. Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, et al. The Perseus computational platform for comprehensive analysis of (prote)omics data. Nat Methods. 2016;13: 731–40. doi:10.1038/nmeth.3901

105. R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2022.

Legends to the figures

Fig. 1. A setup for generating crosslinked agarose-gelatine beads and their properties. In the initial step (A), a mixture of gelatine-agarose (2% each in distilled water) is prepared and placed on a magnetic stirrer with a hot plate to maintain liquidity. A peristaltic pump continuously pumps this solution, forming drops deposited into ice-cold mineral oil to create millimetre-scale beads. Subsequently (B), the mineral oil is removed, and washing steps, including detergent, are performed to eliminate oil residues. Following this (C), the beads undergo crosslinking using divinyl sulphone (DVS), with the crosslinker later removed through additional washing steps. After removal of the crosslinker, the beads are transferred to a bottle and autoclaved at 121°C for 15 minutes at 1 atmosphere (D). Once autoclaved, the beads can be stored for several months until use. In the final step (E), beads are transferred to a 24-well multiwell plate, where the chosen bacteria are inoculated, and the biofilm is established, as previously described. Swelling stability post-crosslinking and autoclaving for five months in two standard culture media: LB (F) and MHB (G). The bars represent the mean of 5 beads (N=5), and the error bars represent the standard deviation. The crosslinking significantly increased the stiffness of the beads as measured via Young's modulus of an atomic force microscope (H). We performed a student's t-test, and the four asterix indicate that the p-values were below 0.0001. Comparative surface view of AGB and glass bead acquired with a Keyence VHX-X1 microscope (I). See the Material and Methods section and supplementary video S1 for additional details.

Fig. 2. Comparative biofilm formation of *E. coli*, *P. aeruginosa*, and *S. aureus* in agarose-gelatine beads and glass after 24 and 48 hours. Boxplots of bacterial count from six biofilm-containing beads for each species grown on the surface of the beads at 24 and 48 hours (N=6). The samples were compared via the Wilcoxon-Mann-Whitney test, and the two asterix indicate that the p-values were below $p<0.01$. The Boxes extend from the 25 to the 75 percentiles, and the whiskers represent minimal and maximal values, the middle line representing the mean.

Fig. 3. Microscopy of AGB biofilm formed by GFP-labelled *E. coli, P. aeruginosa and S. aureus* using
fluorescence, confocal laser scanning microscopy and Scanning electron microscopy. AGBs were allowed to
mature into a biofilm over 48 hours (A). A Z-stack comprising 29 images revealed a surface section of the AGB
densely covered with an intact *E. coli* biofilm, and a single cross-section image extracted from the Z-stack
displayed an AGB uniformly coated with *E. coli* biofilm. Later, the imaging process was made less tedious using
a 3D printed adapter (See Material and Methods and Fig. S4), which also allows imaging with an oil immersion
lens (B). After 48 hours of cultivation for *E. coli, S. aureus* and *P. aeruginosa* at different magnifications using
Scanning Electron Microscopy, biofilms on the surface of an AGB (C).

1110
Fig. 4. Comparison of relative wide proteome expression between agarose-gelatine beads and glass beads
for bacterial biofilms of *E. coli, P. aeruginosa*, and *S. aureus*. Volcano plots show relative protein intensity
changes between AGBs and glass beads measured by LC-MS and label-free quantification. Orange dots denote
proteins significantly more abundant in AGB biofilms, while blue dots indicate proteins more abundant in glass
beads biofilms. Proteins are considered significantly changed if they show at least a 2-fold change in their
relative abundance (log2 fold change >1 or <-1) with an FDR adjusted *p*-value < 0.05, as indicated by the
orange lines.

1118
Fig. 5. Diverse experiments with biofilm formation, including biofilm deficient strains, antibiofilm surface
treatment and crystal violet staining. (A) Two biofilm-deficient strains of *E. coli* (Δ*fimH*) and *P. aeruginosa*
(Δpfpl) on AGBs after 24 and 48 hours. Boxplots of bacterial count from biofilms of *E. coli* and *P. aeruginosa*
mutants compared to their respective parental wild-type strains (N=6). (B) Comparative biofilm formation of *E.*
coli, P. aeruginosa and *S. aureus* on melittin-decorated agarose-gelatine beads after 48 hours of incubation.
Boxplots of bacterial counts from biofilms attached to melittin-coupled, uncoupled, and BSA-coupled after 48
hours of incubation (N=6). (C) Biofilm formation of *P. aeruginosa* and *S. aureus* on AGBs vs. glass beads after 48
hours of incubation using the modified crystal violet staining protocol. Bar charts represent the average
absorption of the dye at 592nm that remained bound to the biofilms, a qualitative measure of biofilm
formation capacity. We measured ten repetitions for each species on each substratum (N=10). The samples
were compared via the Wilcoxon-Mann-Whitney test. One asterisk represents $p<0.05$, two asterisks for
$p<0.01$, three for p<0.001 and four for $p<0.0001$.

1131
Fig. 6. Comparative amounts of different virulence factors from *P. aeruginosa* and *S. aureus* on agarose-
gelatine beads after 48 hours of incubation. Boxplots of (A) pyoverdine, (C) pyocyanin (produced by *P.*
aeruginosa), and (B) staphyloxanthin (produced by *S. aureus*) accumulation from biofilms after 48 hours of
incubation (N=6). (D) Estimating N-Acyl homoserine lactones in *P. aeruginosa* shows the possible effect of
diffused molecules on AGBs on quorum sensing regulated activity. The samples were compared via the
Wilcoxon-Mann-Whitney test. One asterisk represents $p<0.05$, two asterisks for $p<0.01$, three for $p<0.001$ and
four for $p<0.0001$. The Boxes extend from the 25 to the 75 percentiles, and the whiskers represent minimal
and maximal values, with the middle line representing the mean.

1140
Fig. 7. Summary of the potential applications of the new model of crosslinked agarose-gelatine beads. These
beads, with their biocompatibility, ease of preparation, and versatile physical properties, hold significant
potential as substrates for biofilm-related research and applications. Their adaptability to mimic various
environmental conditions, from nutrient-rich to nutrient-limited, provides a versatile platform for studying
biofilms under different scenarios. They can be used to investigate antibiofilm agents, evaluate the efficacy of
various compounds in disrupting or inhibiting biofilm formation, and develop new antibacterial treatments and
coatings. Moreover, these beads can facilitate studying quorum sensing (QS) mechanisms within biofilms.
Furthermore, the beads are compatible with various analytical techniques, including live fluorescence and
scanning electron microscopy.

1150
1151

Agarose-gelatine bead biofilm model applications

Omics of biofilm

LC-MS identification
RNA-seq and Proteomics

Biofilm microscopy

Quorum sensing

Surface modification

In vivo experiments

Experimental evolution

MIC

Anti-biofilm testing

peristaltic pump

cold mineral oil

agarose-gelatine mix

ice water bath

A

B

oil removal and washing steps

crosslinking

C

crosslinker removal and washing steps

D

autoclaving

200
150
100
50

E

biofilm inoculation

F

Swelling over time in LB

Bead diameter [mm]

Months

G

Swelling over time in MHB

Bead diameter [mm]

Months

H

Young's modulus

Stiffness Value (MPa)

**

Cross-linked beads

Non-cross-linked beads

I

AGB

glass bead

2000µm

2000µm

50µm

50µm

50µm

50µm