Recon8D: A metabolic regulome network from oct-omics and machine learning

Ryan Schildcrout[1], Kirk Smith[1, 2, 3], Rupa Bhowmick[1], Yuntao Lu[1], Suraj Menon[1], Minali Kapadia[1], Emily Kurtz[4], Anya Coffeen-Vandeven[5], Srikar Nelakuditi[1], Sriram Chandrasekaran[1, 6, 7]

[1] Department of Biomedical Engineering, University of Michigan, Ann Arbor.

[2] Programs in Metabolism and Medical & Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[3] Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA

[4] Department of Chemical Engineering, University of Michigan, Ann Arbor.

[5] Department of Biological Chemistry, University of Michigan, Ann Arbor

[6] Program in Chemical Biology, University of Michigan, Ann Arbor.

[7] Rogel Cancer Center, University of Michigan Medical School, Ann Arbor, MI, 48109, USA.

* Correspondence: csriram@umich.edu

Abstract

To explore multiomic regulation of the metabolome, we used machine learning to predict metabolomic variation across ~1000 different cancer cell lines with matched omics data from 8 biomolecular classes: genomics (copy-number and mutations), epigenomics (histone post-translational modifications (PTMs) and DNA-methylation), transcriptomics and RNA splice variants, non-coding transcriptomics (miRNA and lncRNA), proteomics, and phosphoproteomics. Overall, the metabolome is tightly associated with the transcriptome, with coding and non-coding RNAs emerging as top predictors. Peripheral metabolites are predictable via levels of corresponding enzymes, while those in central metabolism require combinatorial predictors in signaling and redox pathways, and may not reflect corresponding pathway expression. We reconstruct multiomic interaction subnetworks for highly predictable metabolites, and YAP1 signaling emerged as a top global predictor across 4 omic layers. We prioritize predictive multiomic features for single-cell and spatial metabolomics assays. Top predictors were enriched for synthetic-lethal interactions and synergistic combination therapies that target compensatory metabolic modulators.

Introduction

Variation in metabolite levels between cells may arise due to differences in regulation at the transcriptional, post-transcriptional, translational, and post-translational levels[1,2]. Understanding metabolic regulation in a holistic manner remains challenging due to the complex interconnected nature of metabolic networks, comprising thousands of reactions, that are controlled by an equally complex web of regulatory interactions[3]. In addition to thermodynamics and nutrient availability, intracellular metabolite levels are driven by enzyme activity, which is the basis for the metabolome's relationship with proteogenomic networks. Because of their wide availability, transcriptomics or proteomics data have been used to predict the metabolome with limited success[4–7]. However, these studies focus on a limited set of omics data and do not examine the relative influence and predictive power of each omics class on the metabolome[8–18]. Further, it is unclear if certain metabolites are more predictable than others.

Here, we leverage natural metabolic variation across hundreds of cancer cell lines to infer a multiomic predictor network of the metabolome (Recon8D) using machine learning (ML). Our approach utilizes 10 omics inputs from 8 biomolecular classes from the Cancer Cell Line Encyclopedia (CCLE), allowing for direct comparison of each class on predicting metabolite variation. We hypothesized that the extent of predictability is indicative of the strength of relationship with the metabolome. Similar assumptions have been used in numerous studies to assess relationships between transcription factors and target genes[19–22]. Although many metabolic regulatory relationships have been documented, they represent a small fraction of the number of interactions that likely exist in biological systems. ML algorithms can fill this gap by learning relationships from experimental data. Our approach uncovered numerous mechanistic relationships and identified significant multiomic predictors of metabolomic variation that could serve as markers for either individual metabolites or the overall metabolome for *in vivo*, single-cell or spatial omics studies that require limited measurements[23–26]. Finally, our discovery of synthetic lethal interactions among top metabolite predictors can help design combination therapies by targeting compensatory mechanisms.

Results

Recon8D model construction using machine learning

Analogous to gene regulatory inference with ML, wherein the expression value of each gene is taken as the learning objective, here we use metabolite level variation as the prediction task in a regression problem using multiomic data as input[21,27]. Each omics class is used individually to predict the metabolome, enabling rational comparisons of predictive capabilities across omics. The output is an 'influence' network model that implicitly captures regulatory events at the transcriptomic, proteomic or post-translational levels. Our approach used omics from the CCLE, which include biomolecular assays from up to ~1,000 cancer cell types[28], to predict the corresponding metabolome. We used 8 biomolecular classes: genomics – 1. copy number variation (CNV) and 2. mutations, epigenomics – 3. DNA methylation and 4. histone PTMs, transcriptomics – 5a. coding transcripts and 5b. RNA splicing, non-coding transcriptomics – 6a. miRNA and 6b. lncRNA, 7. proteomics, and 8. phosphoproteomics to train ML models for predicting relative levels of each metabolite across matched cell lines (Figure 1, S. Table 1).

Figure 1: Graphical overview of our approach to predict the metabolome from multiomics and inferring top global and metabolite-specific predictors. Each omics class is utilized in isolation to train a predictive model for each metabolite in a metabolomic panel, and the accuracies and feature importances thereof are used to reconstruct Recon8D metabolic regulatory networks.

Random forest regression ML algorithms were used to generate metabolomic predictions, as they generally outperformed other ML methods based on cross-validation in our benchmarking (S. Figure 1). Random forests have also outperformed other methods in the inference of gene regulatory networks in prior studies[21,27]. We further make use of inherent characteristics of random forests to obtain the importance of each feature in prediction tasks. Higher feature importances indicate a greater possibility that a gene or protein regulates the target metabolite. Separate random forest models were trained iteratively for each of 225 metabolites in the CCLE metabolome using each of the omics features in an input set to create a total of 2,025 models (Figure 2A, 2B). Model performance was evaluated using cross-validation, and significantly predictable metabolites were identified based on a Bonferroni-corrected P value of 0.05 to account for false discovery rate (Figure 2C). We aimed to avoid bias in our models by employing a wide array of omics without incorporating information from previously known molecular interactors, as some omics classes are studied far more in depth than others. This framework enables rational comparisons across omics classes regardless of previous knowledge on their regulatory influence.

100
Figure 2. Random forest model performance indicates transcriptomics, DNA methylation, and non-coding
transcriptomics are accurate predictors of metabolomic variation. A. Representative examples of 1-

methylnicotinamide predictions using transcriptomic, miRNA, and proteomic inputs. Each omics class is used
individually to predict relative 1-methylnicotinamide levels, which yields a predicted metabolite variation array
which can be compared to the true values. Scatter plots of predicted vs true values in cross-validation help visualize
the accuracy of each model. Finally, feature importance analysis identifies the most influential features on the model
output. In this case, the model identified NNMT, the enzyme involved in 1-methylnicotinamide synthesis, as the top
feature predictive of 1-methylnicotinamide. **B.** Number of features in each input dataset. **C.** Number of significantly
predicted (FDR p-value < 0.05) metabolites per input omics class for each of 225 metabolites measured in the CCLE
panel, also represented as a percentage out of 225. **D.** Box plots showing the distribution of Pearson's R values for
predicted vs true values from each omics model type over 225 metabolite models. **E.** Clustered heatmap showing
correlation values for all non-lipid metabolites (columns) with select metabolites labeled.

ML model performance indicates metabolic variation reflects transcriptional differences between cell lines

Across all ML algorithms and validation procedures, transcriptomics consistently generated the
most accurate predictions (Figure 2, S. Figures 1, 3, and 4). While histone PTMs and
phosphoproteomics were not as accurate as other omics classes, they had fewer features by an
order of magnitude, showcasing that small feature sets can still impart meaningful predictions.
The ability of each omics class to make accurate predictions independent of feature set size
instills confidence in the hypothesis that the ML models are picking up mechanistic relationships
and are not overfitting to the data. For example, with the seventh-most number of features out of
9 omics classes, miRNA inputs produced the fourth-most significant predictions. In contrast,
genomics (CNV) had the most features, yet only the sixth-highest predictive performance. We
further validated this assumption with benchmarking models trained on the top 40 features per
dataset identified using holdout validation with a training set to testing set ratio of 75 to 25 (S.
Figure 1). The transcriptomic and miRNA data achieved high accuracy (96% of metabolites
predicted with FDR p-value < 0.05) while only using 40 features. The top 40 features can thus
potentially serve as a reduced set of molecular markers that can be measured for reconstructing
the metabolome (S. Table 2).

Box plots and clustered heatmaps of Pearson's correlations (model predictions versus true
relative metabolite levels) illustrate the differences in predictive power across omics inputs
(Figure 2D, E). While box plots help visualize predictability across the metabolome, the heatmap
of individual models is useful for identifying metabolites that are differentially predictable with
certain omics inputs. While some metabolites are highly predictable across omics layers, such as
1-methylnicotinamide and phosphocreatine, others have stark differences in predictability. For
example, anthranilic acid, 2-deoxyadenosine, and alpha-glycerophosphocholine predictions were
more accurate with transcriptomic and proteomic inputs. This suggests that individual
metabolites can be influenced preferentially at particular omics levels. Conversely, some
metabolites, such as pipecolic acid, do not have high correlations with any model, suggesting
that their mode of regulation might not be sufficiently captured in these data.

The ability to quantify associations with the metabolome across omics classes is vital for
prioritizing the study of novel regulators. Globally, model results indicate that non-coding RNAs
are highly associated with metabolomic variation across cell lines as compared to other omics
classes, highlighting their importance for further study in context with metabolic regulatory
networks.

Control and validation experiments show consistency and robustness in predictive capability

To account for confounding factors that may drive spurious correlations, such as tissue type, culture conditions, and tumor sample origin, several control experiments were conducted. We trained models using omics subsets separated by growth type (adherent and suspension), cancer lineage (lung and hematopoietic/lymphoid tissue, as well as all lineages except lung or hematopoietic/lymphoid tissue), and by tumor sample origin (primary and metastatic) (S. Figure 3). Out of the 8 control experiments conducted, confidence scores were calculated for each feature (0-8) depending on controls where the feature appeared in the top 20 (Methods). The goal of this system is to assess the ability of the models to identify regulator-metabolite relationships rather than assess overall accuracy. Top features discussed throughout are shown with corresponding confidence scores (CS: 0-8), and those discussed in the text have a CS of at least 2, implying that they were observed in the overall model and at least two other control conditions. While the raw accuracies of the models in these experiments are lower due to loss of observations (cell lines) per dataset, thereby limiting training data, similar trends in the predictive power of each omic subset are observed across controls. Models controlled for tumor origin show a high level of concordance in accuracy and feature importance. Overall, transcriptomics had the highest or second-highest predictive accuracy across all omics models in all control experiments.

To assess the robustness of the top features in independent datasets, validation experiments were conducted where top features from CCLE-trained models were compared to corresponding relationships between features and metabolites in the NCI60 cancer cell line panel[29], similar to the control experiments described above. Random forest models were trained on NCI60 genomics (CNV), transcriptomics, miRNA, proteomics, and phosphoproteomics data, and top features from the CCLE models were compared to the top features from the NCI60 models (Methods). CCLE-identified features showed significantly higher feature importance scores in the NCI60 data as compared to the whole feature set (S. Tables 3-4). To further assess the generalizability of this framework to single-cell datasets, wherein the metabolome is challenging to measure, we performed a similar analysis described above with single-cell transcriptomics of the cell cycle and compared with matched metabolome from synchronized cells[30,31]. CCLE-identified features show significantly higher feature importance scores in the single-cell transcriptomics data than expected out of random chance (S. Table 5). These experiments demonstrate generalizability of the feature-metabolite associations from CCLE-trained models across cell line panels.

ML Feature Importance Analysis reveals global and metabolite-specific predictors

For each metabolite, we looked at top predictors from each omics-based model based on feature importance scores (Methods). The transcriptomics model not only had high predictive power, but also recalled several known mechanistic metabolite-gene relationships. For some metabolites, the mRNA transcript of the corresponding enzyme that synthesizes the metabolite was the top predictor. These include choline O-acetyltransferase gene (CS: 4) for the metabolite acetylcholine, choline dehydrogenase (CS: 4) for betaine, cystathionine beta-synthase (CS: 7) for cystathionine, and glycerol 3-phosphate dehydrogenase (CS: 7) for alpha-glycerophosphate. For other metabolites, the consumption or catabolic degradation was more predictive of their levels.

For example, cytidine deaminase (CS: 6) for cytidine, and monoamine oxidase A (CS: 2) for
serotonin. Several metabolites' corresponding transporters also featured as top predictors,
including SLC6A6 (CS: 8) for taurine and SLC6A8 (CS: 8) for phosphocreatine. Nicotinamide
and tryptophan metabolism shared several top predictors, suggesting shared regulation.
Nicotinamide phosphoribosyltransferase (NAMPT) (CS: 4) was a top predictor of NAD.
Indoleamine 2,3-dioxygenase (IDO) (CS: 6) and tryptophan 2,3-dioxygenase (CS: 5) were most
predictive of kynurenine. Anthranilic acid was predicted by kynureninase (CS: 6), and both
methylnicotinamide and niacinamide were predicted by nicotinamide N-methyltransferase
(NNMT) (CS: 8 and 4, respectively).
206
The proteomics model also recapitulates several known relationships between metabolites and
specific enzymes that impact their levels. For example, the top predictor in the proteomics model
for cystathione was cystathionine beta-synthase (CBS) (CS: 5), and the top predictor for
phosphoenol pyruvate was pyruvate kinase (PKM) (CS: 3). Thymine and uracil's top predictor
was dihydropyrimidine dehydrogenase (DPYD) (CS: 2 and 5, respectively) - the initial and rate-
limiting enzyme in uracil and thymidine catabolism. Similarly, malic enzyme (ME2) (CS: 4),
part of the malate-aspartate shuttle, was a top predictor of aspartate levels, while glutaminase
(GLS) (CS: 3) was a top predictor of glutamine.
215
While many metabolites were predicted preferentially by a single top feature, others were
predicted by multiple features, suggesting that they might be involved in multiple pathways and
may be regulated combinatorially. Such metabolites will be harder to predict using
transcriptomic variation of a single gene. For example, citrate is an intermediate in multiple core
metabolic pathways, and its concentration remains low in cancer due to its rapid conversion into
acetyl-CoA and oxaloacetate[32]. As opposed to metabolites such as taurine where there is one
clear regulator, citrate's feature importances show a more balanced regulatory influence among
several top features (Figure 3D). The top predictor for taurine was SLC6A6 (CS: 8), the taurine
transporter, and was shown by our models to have an extremely high influence in predicting
relative taurine levels as compared to even the second most important feature by Shapley
analysis (Figure 3D). However, citrate's feature importance plot does not show one clear top
predictor, but rather an array of features that appear to impact relative citrate levels similarly. In
this way, ML can uncover complex metabolic control by multiple inputs in addition to strong
associations between single features and metabolites. Interestingly, metabolites that had a
distinct single predictor like anthranilic acid, N-carbamoyl-beta-alanine, and adenosine were also
more generalizable to NCI60 (S. Table 3). This may also be related to the limited ability of ML
to tease out complex combinatorial relationships using small data such as the NCI60 panel.
233
Analysis of genomic mutations data revealed several well-known associations between genetic
regulators and metabolite variation in cancers. Due to its binary structure, mutation data was
used to predict metabolite variation using support vector machines (SVMs) (S. Figure 1). While
relatively less accurate than other omics across the metabolome, the mutation ML models
nevertheless recapitulated known associations between important cancer markers and the
metabolome. Out of the 107 metabolites predicted significantly with mutation data, mutual
information calculations between mutation features and metabolites revealed a handful of critical
cancer-related genes consistently appearing in the top 20 features across all metabolites (S. Table
6). For example, TP53 appeared in the top 20 features 20 times, APC 19 times, CDKN2A 17

times, PTEN 16 times, RB1 14 times, and MUC16 12 times. TP53 was a top feature for several nucleotides and APC was associated several amino acids and lipids, consistent with their known effects on cancer metabolism (S. Table 7).

The ML models also have the ability to identify potential novel associations between features and metabolites. While the top feature for 7 out of 10 of the most accurately predicted metabolites from transcriptomics was found to be mechanistically connected to its respective metabolite, the remaining 3 metabolites–oxalate, kynurenic acid, and alpha-hydroxybutyrate–had YAP1 (CS: 3, 3, and 3, respectively), a signaling protein, as the top predictor. Oxalate has been found to be associated with YAP1[33], but the association between kynurenic acid or alpha-hydroxybutyrate and YAP1 has yet to be established.

Comparison of predictive capabilities between omics is particularly useful in the case of lncRNAs, where very few features have known functions, yet show tight associations with the metabolome. Overall, lncRNAs produced highly accurate metabolome predictions, generally second-best behind transcriptomics of protein coding genes. Interesting patterns emerged in the lncRNA data, such as MIR3936HG being the top feature for carnitine, acetyl-, propionyl-, malonyl-, butyryl-, and valeryl-carnitine (CS: 7, 7, 6, 6, 4, and 8, respectively). Top lncRNA features with high confidence scores (>5) prioritized for further study in context with their predicted metabolites are provided in the supplementary materials (S. Table 8).

The ML models also revealed long range metabolite-enzyme interactions. Top features in the proteomics model for glutathione, a key redox metabolite, included SLC7A11 (CS: 3) and ALDH3A1 (CS: 3). SLC7A11 is a cysteine and glutamate transporter (xCT), and its activity has been shown to influence glutathione levels[34]. Aldehyde dehydrogenase protects cells from oxidative stress and consumes redox cofactors. Interestingly, SLC7A11 and ALDH3A1 also form a synthetic lethal pair, suggesting that they may compensate for each other's activity in maintaining redox balance[35].

Interestingly, the top predictors for a majority of the metabolome across various omics data involved either signaling proteins like YAP1 or redox enzymes such as GPX8 (Figure 3B and C). YAP1 was among the top 2 predictors in the transcriptomics model for 45 metabolites including numerous amino acids and nucleotides, while GPX8 was among the top 2 predictors for 23 metabolites including various lipids. This suggests that the combination of growth signaling and redox homeostasis influences a significant part of the metabolome.

Global feature importances from control experiments also identified features integral to specific cellular states. For example, the ML models built using primary or metastatic cell lines revealed distinct miRNA predictors that are known drivers of tumor phenotypes in each condition. miR-142-5p, which has been shown to limit cell invasion and migration[36], was the second most important global feature from the miRNA models and the most important in the primary tumor controls. miR-142 was also a top global feature in the DNA methylation data, 4th in whole-data models and 3rd in primary controls, but is not a top predictor in the metastatic ML models. In the metastatic samples, miR-142 was ranked 233rd out of 734 miRNAs, showing that its relation to metabolite level variation is blunted in the metastatic state. Additionally, miR-181a, which is known to promote proliferation and invasion[37], is the 44th most important feature in the primary

289 tumor control models versus 17th in the metastatic controls, highlighting its association with
290 metabolism in metastatic cells in particular.
291
292 ***Top predictors are close topologically to the corresponding metabolites***
293
To further substantiate the ability of ML to identify metabolic regulatory interactions, we
mapped features from genomic, DNA methylation, transcriptomic, and proteomic models to
reactions present in the Recon3D model (Figure 3A, Methods). Recon3D is a metabolic network
model built on biochemical reactions curated from literature[38]. It is a comprehensive collection
of known metabolic reactions in human cells, making it a prime candidate for systematically
validating ML models and uncovering topological relationships between top features and their
respective predicted metabolites. To ensure overlap with Recon3D, ML models were re-trained
on the Recon3D metabolic gene subset. The resulting metabolic subset ML models showed
similar accuracy to whole-data models (S. Figure 4B and C). Feature importances from
metabolic gene-only models and whole-data models were also conserved significantly (S. Figure
4D and E).
305
306 The top 5 features for each metabolite from all the metabolic ML models displayed a substantial
307 number of overlaps (92-96%) with either the corresponding reactions directly associated with
308 each metabolite or their first-degree neighbors (S. Table 9). This suggests an arms-length or 1-
309 step influence strategy wherein the top predictors for ~95% of the metabolites are within 1 step
310 of the reactions involving those metabolites.
311
While the top features were on average topologically closer to the corresponding metabolite,
consistent with prior studies[6,39], we found distinctions for metabolites in peripheral or central
metabolism (Figure 3C). For example, in the transcriptomics ML models, the top features for
only 50% of the metabolites in central carbon metabolism overlapped with either the
corresponding reaction or first-degree neighboring reactions. In contrast, the top features for 85%
of the metabolites in peripheral metabolism were direct or first-degree neighbors, suggesting that
peripheral metabolites are more influenced by direct enzymatic regulation, as opposed to those in
central carbon metabolism, which are likely influenced more by signaling (Figure 3C).
320
To further explore if signaling is a larger player in central carbon metabolism than peripheral
metabolism, we determined if top features in each category were signaling enzymes for all non-
lipid metabolites. We found that 31% of metabolites in central carbon metabolism had signaling
enzymes as the top feature compared to only 12.5% for peripheral metabolites (Figure 3C).
These findings suggest that central carbon metabolism is influenced to a greater extent by
signaling than peripheral metabolism.

Figure 3. Global insights across all 225 ML models reveal distinct patterns of predictability. A. Workflow for topologically mapping features to Recon3D. Top 1, 2, and 5 features are identified from ML, and are assessed for overlap with genes present in the reactions for its predicted metabolite (both the enzyme target and the 1st degree

neighbor). **B.** Pie chart showing the percentage (of 225) of top features from transcriptomic inputs. YAP1 and GPX8 account for 15.6% and 5.8% of top 1 features, respectively. **C.** Percentage of top 1 features from biologically relevant feature subsets using transcriptomics inputs. 42% (94) of top features (out of 225 metabolite models, metabolic gene-only) are redox enzymes, and 35% (79) of top features (out of 225 metabolite models, whole-data) are signaling enzymes. 17% (16) of top features (out of 95 metabolites that overlapped between CCLE and Recon3D) were involved in the primary reaction for their respective metabolite, and 81% (77) of top features (out of 95 metabolites) were involved within the 1st neighboring reaction for their respective metabolite. 50% (5/10) top features from the central carbon metabolites and 85% (72/85) of peripheral metabolites (from 95 total metabolites that overlapped between CCLE and Recon3D) were involved within the 1st neighboring reaction for their respective metabolites. Lastly, out of 136 non-lipid metabolite models, 31% (5/16) of top features were signaling enzymes for central carbon metabolites, and 13% (15/120) of top features were signaling enzymes for peripheral metabolites. **D.** Contrasting Shapley importance features for citrate and taurine from transcriptomics models. While the taurine transporter SLC6A6 is clearly shown to be the most important feature in predicting relative levels of taurine, citrate does not have one clear top predictor. In this way, feature importance analysis of citrate shows high order, complex influence by multiple features. **E.** Heatmap of log2-scaled feature importance scores for 136 non-lipid metabolites. The top 5 features for each omics class are shown for every metabolite in order of transcriptomics feature importance (descending). This shows a set of metabolites that have 1 strong predictor (top of heatmap) and those metabolites, primarily related to central metabolism, that have multiple predictors (bottom of the heatmap).

YAP1 and GPX8 are globally important features across the metabolome

After identifying the most influential features locally for individual metabolite models, we next pinpointed features that were globally important across the metabolome using the average rank of the features for all 225 metabolite models (Methods, S. Table 10). These results were then scaled by Z-score and top features for each input omics class were identified. Important features of interest were checked for overlap between whole-data models and controls for robustness.

Some top features were several standard deviations above the mean for certain omics classes, such as YAP1, GPX8, and SCD in the transcriptomics models, with Z-scores of 69, 61, and 59, respectively. Well known metabolic regulators such as mTOR also featured in the global top regulators. mTOR phosphorylation was a top 5 feature for several metabolites in central and nucleotide metabolism including oxalate, succinate, xanthine, and phosphogluconate; similarly, phosphorylation of Rictor, which interacts with mTOR was a top feature for amino acids including arginine, consistent with their role in amino acid sensing and regulation[40].

Histone PTMs are increasingly being recognized for their role in sensing and regulating metabolism[41–43]. The CCLE epigenomics data measures total PTM levels and lacks gene-level resolution. Total levels of H3K9 trimethylation and H3K27 di- & trimethylation were the top features in histone PTMs; these are well known repressive marks, and importantly, are also sensitive to the metabolic state of the cell. Histone methylation is sensitive to the level of one-carbon metabolites like SAM and one-carbon sink metabolites such as methyl nicotinamide[42].

Interestingly, top global predictors showed consistency across omics classes. The most intriguing example of this was YAP1, which was found to be the most important global feature for transcriptomics and phosphoproteomics, as well as the seventh for DNA methylation. YAP1 is a coactivator in Hippo signaling, which is essential for growth signaling, oncogenic adaptations, and the epithelial mesenchymal transition (EMT)[44]. YAP1 was the most important feature for 35 metabolite models using transcriptomics, 2 using proteomics, 5 using DNA methylation, and 132 using phosphoproteomics. Other examples of cross-omic consistencies for top features were

GPX8 (transcriptomics and proteomics), miR142 (methylation and miRNA), EPHA2 (transcriptomics and RNA splicing), and ERBB2 (DNA methylation and proteomics). Not only do different omics display conserved global regulators, but globally important features have established relationships. For example, the top global feature from RNA splicing, EPHA2, has been shown to induce accumulation and activation of YAP1, subsequently promoting glutamine metabolism[45].

To validate the association between YAP1 transcripts and the 35 metabolites for which it was the top predictor, we used the DepMap Data Explorer to quantify the relationships between two isoforms of YAP1 protein and each metabolite with linear regression (S. Table 11)[46]. We found that most associations were statistically significant (FDR p-value < 0.05), thus increasing confidence that YAP1 acts as an important predictor for a wide array of metabolites.

Construction of multiomic metabolite regulome networks for top predictive metabolites

The Recon8D network, containing the top 10 predictors identified from random forest feature importance scores from each of the omic layers for all 225 metabolites is available in S. Table 12 and on the [GitHub page](#). To understand how top omics features interact with one another as well as with the metabolite of interest, metabolite regulome networks were constructed using top features identified by Shapley importance analysis (Methods) (Figure 4A). Shapley importance analysis for the top 3 most accurately predicted metabolites for each omics class was conducted to assess the directionality and relative magnitude of feature-metabolite associations (S. Figure 2).

We further focused on 1-methylnicotinamide and phosphocreatine, which were the most highly predictable metabolites (lowest P values and largest correlations) across omics classes (Figure 4A, S. Figure 6). Relationships between miRNA and RNA were identified using TarBase[47], and between DNA methylation, RNA, and proteins using BioGrid[48]. The interconnectedness of these networks highlights the model's ability to identify rational biochemical relationships. For example, nicotinamide N-methyltransferase (NNMT), the enzyme involved in the methylation of nicotinamide, was the top predictor of 1-methylnicotinamide levels from both the transcriptomic (CS: 8) and proteomic (CS: 6) models. NNMT has also been shown to modulate histone methylation[49,50]. Similarly, solute carrier family 6 member 8 (SLC6A8) (CS: 8), a creatine transporter, was the top predictor of phosphocreatine in the transcriptomics model, and creatine kinase (CKMT1A) (CS: 3), the enzyme responsible for phosphorylating creatine to create phosphocreatine, was the third most important phosphocreatine predictor in the proteomics model (S. Figure 6). Shapley analysis also identifies the directionality of those relationships. For example, NNMT is positively correlated with 1-methylnicotinamide concentrations (Figure 4A, right). Similarly, hsa-miR-200c, which was identified as a regulator of NNMT translation via TarBase, is negatively correlated with 1-methylnicotinamide levels. The interconnectedness of the top predictive features across models suggests that the relative levels of these metabolites involve high-order interactions between each of these omics layers.

Physical interactions are frequent among top metabolite predictors

It is difficult to tease out metabolic regulation by individual biomolecules given buffering and
combinatorial regulation. By integrating physical and genetic interactions with multiomic
associations, we hypothesized that we could understand how groups of regulators complement
each other and maintain metabolic homeostasis. We overlaid protein-protein interaction (PPI)
data from the STRING database, miRNA-gene interaction data from the TarBase database and
metabolite-protein interactions from Recon3D with Recon8D interactions (Figure 4B; S. Figure
5; Methods)[51]. We further analyzed the consistency between the top features in Recon8D with
highly connected nodes in the PPI, Recon3D and miRNA networks. We computed a distance
matrix between the nodes in the multiple networks and compared it with ML feature
importances. Interactions that show high consistency between Recon8D and other interaction
networks were visualized (Figure 4B).
436
This cross-network analysis highlighted several shared interactions between omic layers,
especially between various signaling regulators in proteomics and phosphoproteomics, with
transcription factors in transcriptomics and DNA methylation. Chek1 in phosphoproteomics
shared common patterns of interactions with the transcription factor HOXA5 (DNA
methylation); MET and STAT3 (phosphoproteomics) shared similar interactions with each other
and with the transcriptional regulators WIZ (in DNA methylation) and the microRNA miR-1246.
Similarly, enzymes in fatty acid metabolism shared cross-omic similarity with transcriptional
regulators. Lipoic Acid Synthetase (LIAS) enzyme shared similar targets to H3K36
trimethylation, and fatty acid synthase (FASN) with miR-142-3p. Acyl-CoA Synthetase
(ACSF3; DNA methylation) had similar targets to aldo/keto reductase AKR7A2 (splicing) with
ferritin (FTL), which are all related to redox homeostasis. Genomic copy number changes in
Neuroblastoma Breakpoint Family Member 1 (NBPF1) had similar interactions as the GPX8
gene methylation and the histone demethylase KDM3B proteomic levels. (S. Table 13). These
results suggest that Recon8D predicted interactions are consistent with known molecular
interactions, and our cross-omic analysis highlights potential shared regulation between omic
layers.
453

456
457
Figure 4: Integrating protein-protein, miRNA-gene and metabolite-gene interactions with Recon8D
interactions reveals high consistency between feature importance and topological distribution. **A.** Proposed
metabolic regulome network from top 8 features from histone PTM, DNA methylation, miRNA, transcriptomic, and
proteomic-trained models for predicting 1-methylnicotinamide (highest mean Shap value) as determined by Shapley
analysis. **B.** This figure presents a comparative analysis of relationships between top features derived from Recon8D
multi-omics data, including transcriptomics, proteomics, phosphoproteomics, genomics (CNV), methylation,

miRNA, splicing and epigenomics, and their corresponding metabolites. Top global features from each omics
dataset are organized in distinct, color-coded circular layouts surrounding a central circle of metabolites. The size of
each node reflects the number of metabolites associated with that feature. Edges connecting top features to
metabolites illustrate the consistency in feature importance and topological distribution across the datasets. Edges
between top feature factors represent correlations in their metabolite-consistency profiles, with edge width
corresponding to the strength of the correlation. The node selection is based on the top features across the 8 omics
datasets. For example, Yap1 shows high consistency with 11 metabolites based on distance of Yap1 to these
metabolites in the PPI/miRNA/Recon3D networks and their feature importance in Recon8D, and shares similar
metabolite interaction patterns as miR-320b. Met and Stat3 also share similar metabolite interaction patterns with
each other and with WIZ (in DNA methylation).

Synthetic lethal interactions reveal compensatory control mechanisms

Next, inspired by the synthetic lethal (SL) interaction centered around glutathione highlighted
earlier, we systematically assessed for SL interactions across all top metabolite predictors in
Recon8D. Since metabolite levels are controlled by multiple factors, we hypothesized that the SL
interaction analysis can identify compensatory regulators for each metabolite. We compiled
curated experimental human SL interactions from SynLethDB and tested if these interactions are
enriched among the top 10 predictors of each metabolite (S. Table 14)[52]. The transcriptome
model had the most striking enrichment for SL interactions among top predictors for each
metabolite, with 52 interactions involving 45 metabolites (27-fold higher than random chance, p-
value = 1×10^{-34}) (Figure 5A, 5D).

Several SL interactions involve a combination of signaling factors, suggesting that combinations
of signaling pathways influence variation in metabolite levels, and each gene in the SL pair may
compensate for the other's absence. For example, the top predictors of nucleotides and
nucleotide-precursors showed several SL interactions involving growth signaling factors like
AXL, YAP1, CCN1, PTPN14, CSK, and HRAS. Of note, YAP1 and PTPN14 are both top
predictors of hypoxanthine, and they both form an SL pair and also physically interact[53].

SL interactions among top predictors also highlight potential long range regulatory effects.
Several interactions involved both a growth signaling and a redox-related gene and were
connected to metabolites in central metabolism (Figure 5D, 5E). For example, the top predictive
network for lactate contained an SL interaction between the redox enzyme GPX8 and CAV2,
which is involved in cell signaling, growth, and lipid metabolism. Similarly, the top predictors
for glyceraldehyde-3-phosphate and dihydroxyacetone phosphate, intermediates in glycolysis,
also had an SL interaction between GPX8 and the tyrosine kinase AXL.

Synergistic drug combinations converge on nucleotide metabolism pathways

Synthetic lethality is directly related to drug synergy, wherein drugs targeting proteins in
compensatory pathways show enhanced potency when combined. The abundance of SL
interactions among top predictors in Recon8D suggests that drugs that target predictors of
specific metabolites are likely to be synergistic. To test this hypothesis, we utilized curated drug-
drug interaction data in various cancer cell lines from DrugCombDB which includes 448,555
drug combinations covering 2,887 unique drugs and 124 human cancer cell lines[54]. The drugs
were mapped onto known targets using data from DrugBank[55]. We tested if synergistic
interactions were enriched among drugs that target the top 10 predictors of each metabolite. The

512 transcriptome and phosphoproteome models had the strongest enrichment for synergistic
513 interactions among top predictors for each metabolite (Figure 5B). Drug combinations that target
514 top predictors in the transcriptomic model were highly likely to be synergistic with an average
515 bliss interaction score greater than +4, while randomly selected combinations were likely to be
516 antagonistic (bliss score < -5) (Figure 5B).
517
Similar to observations with SL interactions, several synergistic interactions involved a
combination of signaling pathway inhibitors. We next analyzed metabolites that were associated
with synergistic interactions. Notably, nucleotides and nucleotide precursors showed numerous
synergistic interactions across a wide range of drugs targeting diverse signaling and metabolic
pathways. Guanosine, inosine, pantothenate, thymine, xanthine and hypoxanthine were among
the top metabolites associated with most synergistic drug combinations (Figure 5B, S. Table 15).
Out of 104 drug combinations associated with these metabolites, 100 were synergistic (S. Table
15). Drugs that target diverse signaling pathways may lead to similar downstream effects on
nucleotide metabolism. For example, several tyrosine kinase inhibitors including sorafenib and
sunitinib are predicted to be associated with these nucleotides (S. Table 15). These drugs are
known to affect levels of nucleotides including guanine, adenine, adenosine, guanosine,
hypoxanthine, and inosine[56]. To further validate these associations, we used the DepMap Data
Explorer to quantify the relationships between cellular sensitivity to these signaling drugs and
levels of nucleotides, and found several statistically significant associations (S. Table 16)[46].
These results confirm the connection predicted by our analysis. In contrast, antagonistic drug
interactions were surprisingly associated with metabolites in the TCA cycle. Citrate, isocitrate,
and fumarate were the top metabolites associated with most antagonistic drug combinations.
Several drug combinations associated with these TCA cycle metabolites involve drugs such as
Sirolimus, Everolimus and Tetracycline that are known to reduce the activity of the TCA cycle
(S. Table 15)[57,58]. Drugs that perturb TCA cycle activity may lead to antagonism with agents that
require TCA cycle activity. Thus, Recon8D can provide insights into understanding
combinatorial metabolic regulation, synthetic lethality, and drug synergy.
540

Figure 5. Top metabolite predictors are enriched for synthetic lethality (SL) and drug synergy. A. Frequency of SL gene pairs among top 10 influencers using genomic, transcriptomic, proteomic, phosphoproteomic and multiomic (combination of top 10 influencers from genomic, transcriptomic, proteomic, phosphoproteomic ML models). This is compared with frequency of SL pairs obtained from random sampling using the corresponding omics data. SL pairs were obtained from SynLethDB 2.0 which contains 35,943 human SLs. **B**. Average Bliss drug interaction score (positive values implies synergy) for pairs of drugs that target top predictors of a given metabolite derived from various omics. Top metabolites associated with synergy or antagonism are highlighted. Drug interaction data was derived from DrugCombDB. **C**. High confidence genetic and physical interactions among the top 10 global predictors of phosphoproteomics ML model from the STRING database. **D**. High confidence genetic and physical interactions among the top 10 global predictors of transcriptomics ML model from the STRING database. **E**. High-confidence genetic and physical interactions among the top 10 predictors of the metabolite glutathione.

Conclusion

Cellular metabolism is controlled by multiple layers of regulatory networks. Here, we quantified the influence of potential regulatory interactors on metabolomic variation from matched omics across 8 biomolecular classes. The metabolome was surprisingly highly predictable from

transcriptomic data. While the absolute levels of metabolites may depend on thermodynamics,
variation may result from differences in transcriptional regulation. Further, the relationship
between transcripts and metabolites is nonlinear, and a combination of transcripts influences
metabolite levels. In many cases, there is no direct correspondence between metabolites and
enzymes they are involved with. Signaling-related transcripts, proteins, and phosphoproteins
were top predictors across omics layers. A second common global predictor was redox
metabolism, exemplified by GPX enzymes, that influence glutathione levels and overall redox
state of the cell. Levels of histone PTMs, miRNA and phosphoproteins were also highly
predictive of the metabolome despite having the fewest features by an order of magnitude.
570
Analysis of feature importance levels of individual metabolites revealed that metabolites in
peripheral metabolism, like taurine, were most predictable via levels of enzymes related to those
metabolites, while metabolites in central metabolism were more predictable by a combination of
signaling and redox regulators. Because central carbon metabolism has been more extensively
studied than peripheral metabolism, we would expect to find strong clear associations with
central carbon metabolism. However, we find the opposite, that central carbon metabolites often
do not have a clear top predictor, suggesting that these metabolites are more prone to
combinatorial regulation. We found these patterns to hold true in independent datasets from
NCI60 and single-cell omics. Our analysis thus uncovered key systems-level insights on
metabolome variation between cells, the generalizability of ML models, and predictability of
individual metabolites.
582
Although we make use of ML to predict the metabolome, such models are not directly
transferrable in predicting metabolomics from other sources due to several factors, including
differences between *in vitro* and *in vivo* conditions, and between cell line panels, and limited
features overlap within and across omics. Furthermore, one of the key observations from this
study is related to the predictability of specific metabolites, such as those in central metabolism,
which can affect the generalizability of ML models that aim to predict metabolomes. While
many metabolites were predicted in our Recon8D model to be preferentially influenced by a
single top feature, others were predicted by multiple features, suggesting that they may be
regulated combinatorially. Such metabolites will be harder to predict using a single omics
feature. Hence, the focus of Recon8D is not the prediction of absolute metabolite levels, but
rather the associations between features and metabolites, as reflected in our validation
experiments.
595
A limitation of our study is that our analysis was done in cancer cell lines in culture and may not
reflect metabolism *in vivo*; however, patterns observed here related to transcriptional variation,
and topological relatedness of gene-metabolite associations have also been observed *in vivo* in
humans and model systems[39,59,60]. Further, the 'guilt by association' strategy used by network
inference methods may not identify causative mechanisms. We address this through integration
of multiple layers of orthogonal omics information that we found to be consistent with each
other[61]. Further, these interactions are robust across hundreds of cell lines in independent
datasets spanning multiple tissue lineages and growth conditions. Our model also correctly
recalls several known interactions reported in literature. The above suggests that the statistically
inferred associations may represent real interactions.
606

Another limitation is that network associations driven by systems-level data are descriptive, and
findings are not necessarily mechanistic, even among high confidence features. However, along
with extensive documentation of Recon8D associations reflected in known molecular interactors,
these findings suggest strong mathematical relationships between the metabolome and other
omics classes. Experimentally validating proposed interactions from this study, even for a small
percentage of interactions, would require an immense effort. We hope that these prioritized
associations may inform rational exploration of novel metabolic regulatory interactors.
614
Additionally, some high-throughput measurements do not provide a comprehensive panel of
molecules that exist within a cell. For example, the metabolite panel used for this study consisted
of 225 metabolites, a fraction of those that exist within a cell across distinct cellular
compartments. Furthermore, there exist more proteins in human cells than used in this study, and
the 90 phosphoproteins present were measured using a targeted approach, and therefore are not
representative of the entire cellular array of phosphoproteins. To make a strong inference as to
the relative influence of these classes on the metabolome, these analyses would need to be
conducted on more comprehensive, untargeted data. Similarly, differences in sample quantities
can influence accuracies, where omics with too few samples (cell lines) may not perform as well
(S. Figure 1, S. Table 1).
625
Beyond the relative predictability of the metabolome from omics inputs, we find that top
predictors from ML are mechanistically related to the associated metabolites. For example, in the
transcriptomics ML model using genes present in Recon3D, the metabolites' top predictors were
enzymes closely associated, either directly (17%) or first-degree neighbor (81%), with those
metabolites. The top predictors for each metabolite across different omics layers were highly
enriched for genetic and pharmacological interactions. This further supports the ability of our
approach to uncover mechanistic interactions and highlights potential multiomic compensatory
buffering between top metabolomic predictors. Finally, we find that drugs that target top
predictors in nucleotide metabolism are likely to be more synergistic. This builds upon our
observation of increased synthetic lethality among the top predictors of each metabolite. Notably,
a wide range of signaling drugs intersect with nucleotide metabolism and are predominantly
synergistic when used in combination (mean bliss score > 10, S. Table 16, Figure 5B). Our
analysis suggests that these drugs, which target signaling pathways, may also impact nucleotide
homeostasis. This is consistent with a prior study that found combinations of Mek and PI3
Kinase inhibitors show synergy via nucleotide metabolism[62]. In contrast, drug antagonism was
associated with metabolites in the TCA cycle. A similar observation was made in bacteria
wherein antibiotics that have opposing effects on TCA cycle and bacterial respiration are
antagonistic[63]. Several drugs require active growth and metabolism to be potent. Inhibition of
TCA cycle may result in growth arrest which may lead to reduced potency of growth-dependent
drugs. In contrast, cells with nucleotide imbalances continue to grow[64]. This may result in errors
in DNA replication and transcription leading to cell death. Overall, Recon8D may provide a
rational framework for metabolic engineering and designing synergistic combination therapies.
648
649
650 **Methods**
651
652 *Data Acquisition and Preprocessing*

Omics from the Cancer Cell Line Encyclopedia (CCLE) were obtained from the Broad Institute (Cambridge, MA) Dependency Map Portal (with the exception of proteomics and phosphoproteomics, which were obtained from Nusinow et al and Li et al, respectively)[28,65,66]. These include genomics (copy number variation and mutations), epigenomics (bulk histone H3 modifications (MRM) and DNA methylation (RRBS)), transcriptomics (RNAseq), RNA splicing (RNAseq, exon inclusion ratio), miRNA-omics (expression), lncRNA (expression), proteomics (mass spectrometry), phosphoproteomics (RPPA), and metabolomics (LC-MS). The number of cell lines used for training can be found in S. Table 1.

Data preprocessing involved replacing missing values (NaN) with zeros, taking the median value of repeated observations (if required), standardizing cell line or feature names across datasets, and removing features with 0 variance. Only lncRNA, RNA splicing, and mutation data had features with zero variance (38, 1,131 and 146 features, respectively). Once removed, there were 3,767, 2,957 and 17,203 features used for model training, respectively. RNA splicing ensemble IDs were converted to common names using BioTools.fr[67]. Each data type was merged independently with the metabolomics dataset by common cell line, where each matrix was organized by cell lines (rows) and features (columns). Input datasets for the models containing metabolically relevant genes were filtered for genes present in the Recon3D genome-scale metabolic model[68]. The same data preprocessing steps and scaling methods were used for validation experiments with NCI60 data, which were obtained from the National Cancer Institute CellMiner portal[29,69].

Machine Learning Model Training

ML models that predict metabolomic variation were trained on individual omic data from the CCLE. A separate model was trained and tested for each metabolite using 5-fold cross-validation, where data were normalized by Z-score within each CV fold. Statistically significant predictions were determined using a Bonferroni-corrected P value (0.05/225) to account for false discovery rate. Because the number of features was much greater than the number of observations in these data, it was important to select algorithms that penalize large coefficients and limit overfitting. Random forests were primarily used for this study, although several algorithms were tested for benchmarking, including Ridge Regression, Lasso Regression, XGBoost, and SVMs[70–74]. Each of these algorithms is easily interpretable (provides feature importance scores or coefficients) and characterizes different mathematical relationships. Non-linear SVMs are more difficult to interpret, and therefore mutual information calculations were required to provide surrogate feature importance scores. The Python script for these algorithms can be found at the Recon8D GitHub repository. The trained random forest models, along with the CCLE data used for model training, can be found on the Recon8D Synapse page.

Random forests are comprised of many decision trees, and are classified as an ensemble learning method, where they aggregate many iterations of decisions. This algorithm works by randomly selecting a training set from the input data (bootstrapping) and using that training set to dictate a unique decision tree[75]. These predictions are averaged over all the decision trees, which is what allows for more accurate predictions from the overall model than any individual tree could provide[75]. Ultimately, we chose random forests for the bulk of our analyses for their balanced and interpretable framework as well as their overall high accuracy in comparison to other

698 algorithms. More complex models can be challenging to interpret, and simpler models may not
699 identify non-linear or combinatorial relationships.
700
Hyperparameters
Random forests were trained using Sci-kit Learn's Random Forest Regressor algorithm with the
following parameters: (random_state=0, n_estimators=100, n_jobs=-1, max_depth=10). These
parameters were selected after considering computation time while maintaining sufficient depth.
While random forests are robust to changes in hyperparameters, we elected to not tune such
parameters to avoid overfitting. A similar approach was taken for XGBoost and SVMs.
707
708 For ridge and lasso regression, an array of alphas was used in conjunction with cross validation
709 to determine the appropriate regularization term for each model. We used an array containing
710 alpha values of 1e-2, 0.1, 1, 10, 100, and 1000. The best alphas were saved for each cross-
711 validated model and applied to final model training using the whole data.
712
Model Evaluation and Interpretation
Performance of each model was assessed using Pearson's correlation between predicted and true
metabolite concentrations. FDR corrected P values were also calculated for assessing
significance. Cross-validation was used to assess overall accuracy of each data type, and a final
model trained on the entire data set was used to examine feature importances. Feature
importance scores were generated from each random forest model and were analyzed locally (for
each metabolite) and globally (averaged across all metabolite models for a given omics type).
Global importances were sorted by median rank and scaled by Z-score (S. Table 10).
721
722 Mutation feature importances for SVMs were calculated using mutual information—a model-
723 agnostic method for assessing the dependency between a feature and a metabolite. This was done
724 in lieu of directly interpretable model characteristics of non-linear SVMs. Instances of mutations
725 in the top 20 features for each metabolite (similar to control experiment confidence scores) were
726 added to assess the impact of each mutation in predicting the metabolome.
727
728 Feature importances for select predicted metabolites were also quantified using Shapley analysis,
729 a game-theoretic approach that calculates feature importances by adding each feature into the
730 model sequentially, and repeating that process for all orderings of features, thereby assessing the
731 influence of each feature on the model as a whole[76,77]. The three metabolites with the lowest P
732 values were selected for Shapley analysis for each data type (S. Figure 2).
733
Consistencies across control experiments were quantified with a confidence scoring system,
where the top 20 features were extracted from whole-data and control models. We then tallied
the occurrence of each of the top 20 features conserved in the top 20 features of each control
experiment, out of 8 experiments, leading to a confidence scoring system ranging from 0 to 8 for
each feature. The 8 experiments included models generated on the following subsets: lung
cancers, hematopoietic and lymphoid cancers, all cancers except lung, all cancers except
hematopoietic and lymphoid, suspension cell culture, adherent cell culture, primary tumor
samples, and metastatic samples.
742

The top one, two, five and ten features (quantified by random forest feature importance scores)
from genomics (CNV), DNA methylation, transcriptomics and proteomics models (for metabolic
gene-only models) were mapped back to the Recon3D metabolic model reactions. This was
accomplished by matching metabolites from the CCLE panel to metabolites present in the
Recon3D metabolic model. Common metabolic cofactors were removed to avoid
overrepresenting neighboring reactions, including ATP, ADP, water, hydrogen, phosphate, and
diphosphate, which may be found in the recon_mapping section of the [GitHub repository](#).
Functions from the COBRA toolbox allowed for the identification of all reactions associated
with the relevant metabolites (both primary and first-degree neighbors), as well as the genes
associated with each reaction[78]. Once identified, gene names were converted to BiggIDs and
matched to each set of reactions. The number of metabolite models that had at least one top
feature present in the relevant set of reactions (out of the 95 metabolites in both the CCLE panel
and Recon3D) were compiled (S. Table 9). We performed this same analysis on a set of random
transcripts, which show fewer overlaps (S. Table 9). A random set of 5 transcripts was taken for
each metabolite and overlaps were counted. This process was repeated 50 times, and the median
value of overlaps is represented in the table.
759
Model Validation in independent datasets
The top features from CCLE-trained models were isolated and used to analyze relationships
between transcripts and metabolites in the NCI60 data and single-cell data. Random forest
feature importance scores from NCI60-trained and single-cell-trained models were used in
comparison to CCLE outputs, where top features as identified by CCLE models were isolated in
NCI60 data to examine if they were larger than by random chance. Z-score-scaled feature
importance ranks were used for this analysis. Student's t tests and Kolmogrov-Smirnov tests
compared isolated feature importances to the entire feature set. Data were stacked to create one
distribution among all metabolites (pooled). Aggregated importances showed significant
differences between the top features and overall feature set. These analyses were conducted on
sets of 1, 3, 5, 10, 50, and 100 top features from each metabolite model (except for
phosphoproteomics, which only had 90 features, and thus the 100-feature subset was not used).
772
Effect of feature size, missing features and metabolic features
We explored how well metabolic genes alone could predict the metabolome. Genomic, DNA
methylation, transcriptomic, and proteomic datasets were filtered for genes present in the
Recon3D metabolic model[68], and new models were trained on the metabolic gene subset. The
difference in feature size between omics classes can affect their predictive power, and filtering
for genes present in Recon3D standardized these data to provide mechanistic information within
the context of known metabolic networks. Each filtered dataset had a similar number of features,
which provided an avenue through which to assess the impact of feature size on model accuracy.
781
We first filtered the feature sets for metabolically relevant genes from Recon3D (S. Figure 4A).
With fewer than 2,000 genes, transcriptomics inputs predicted over 99% of metabolites
statistically significantly, and other feature sets conferred similar accuracy to whole-data models
in predicting each of the 225 metabolites (S. Figure 4B). The median correlation for each omics
class was slightly lower in the metabolic gene-only models as compared to the whole-data
models, but the distributions were similar overall (S. Figure 4C). Similar accuracies from the
metabolic gene-only models and the whole-data models illustrate the lack of impact of feature

size on model predictions. We further explored this by training models on a random set of 2,000
genes, which also had similar accuracies. While models built on randomly selected features
might not provide mechanistic insights, they show that the metabolome is highly predictable
within a reasonable range of input feature sizes. Feature importances from metabolic gene-only
models and whole-data models were also conserved significantly (S. Figure 4D and E).
794
Both the number of training samples (cell lines) and features can impact accuracy. Proteomics
and phosphoproteomics had fewer cell lines for training than the rest of the omics (S. Table 1),
which could explain lower accuracies comparatively. Additionally, we hypothesize that the
reduced feature size in the proteomic data may account for the reduced accuracy in these models
compared to transcriptomics. We interrogated this by comparing the transcriptomic and
proteomic datasets to see if there were discrepancies in the feature set that might influence
accuracy. Between the 416 genes that were present in the transcriptomics feature set but were
absent from proteomics, 11 of those genes were the top 1 predictor in a metabolite model. These
missing features explain the discrepancy in accuracy (~15%) between transcriptomic and
proteomic inputs, suggesting that access to more comprehensive proteomic data might be more
predictive of relative metabolite levels.
806
Statistical analysis
All P values were corrected using Bonferroni multiple hypothesis correction. Statistical
significance of the overlaps with synthetic lethal interactions from SynLethDB 2.0 and drug
interactions from DrugCombDB were assessed by creating 50 sets of random top features for
each metabolite in all omics models and quantifying the overlap. This was then compared with
the actual overlap using a t-test.
813
Multi-Omics Topological Analysis Based on Metabolic, miRNA and PPI Networks
The metabolic network data was obtained from the Recon3D genome-scale metabolic model.
Protein-protein interaction (PPI) data was sourced from the STRING database (https://string-
db.org/), and miRNA-gene interaction data was sourced from the TarBase database
(https://dianalab.e-ce.uth.gr/tarbasev9). A metabolic network linking metabolites to genes was
constructed based on the metabolite-reaction-gene relationships provided by the Recon3D
genome-scale metabolic model. The PPI network and miRNA-gene network were constructed
directly using the pairwise relationships extracted from the STRING and TarBase databases,
respectively. The Construction of Distance Matrices and Rank Difference Analysis involves two
steps. First, the top features (top 20 for epigenomics, top 40 for phosphoproteomics, top 100 for
other omics) were selected by feature importance ranking, mapped to the PPI network, and used
to compute an n×l distance matrix (n: number of top genes in the PPI network; l: number of
shared genes/proteins between the metabolic and PPI networks). Next, an l×m metabolic distance
matrix was generated from the positions of genes in the Recon3D metabolic network (m:
metabolites common to the metabolic network and CCLE metabolomics data). The final n×m
omics-metabolite distance matrix was obtained by multiplying these matrices. For miRNA, a
three-step process was used: first, a distance matrix based on miRNA-gene interactions was
generated, followed by multiplication with the previous two matrices to yield the final omics-
metabolite distance matrix. To compare the ranking differences of top features between the
distance and feature importance matrices, both were converted into ranks, and the absolute
differences were normalized to a range of 0 to 1. For each metabolite, the top 10% and bottom

10% of feature-metabolite relationships were classified as high and low consistency,
respectively. Cross-omic edges were determined based on similarity of metabolite interaction
patterns based on the Pearson's correlation coefficient, and a threshold of +0.4 was used to
visualize significant cross-omic similarities.
839
840
841

Acknowledgments
This work was supported by faculty start-up funds from the University of Michigan (UM),
Camille and Henry Dreyfus Foundation, the Rogel Cancer Center at UM, R35 GM13779501
from NIH to SC, and the Advanced Proteogenomics of Cancer (T32 CA140044) Program to RS.
We thank Deepak Nagrath, Maciek Antoniewicz, Analisa Difeo and Yatrik Shah for suggestions
and critical feedback on the project.

848
849 Competing interests
850 The authors declare no competing interests.
851
852 Data availability
853 All datasets are provided in figures, the supplementary materials, and in the GitHub repository.
854
855
Author contributions
S.C, R.S designed and performed research, and wrote the manuscript; K.S, R.B, Y.L, S.M,
A.C.V, E.K, M.K, S.N performed research.

References

1. Nielsen, J. Systems Biology of Metabolism. *Annu. Rev. Biochem.* **86**, 245–275 (2017).

2. Chubukov, V., Gerosa, L., Kochanowski, K. & Sauer, U. Coordination of microbial

metabolism. *Nat. Rev. Microbiol.* **12**, 327–340 (2014).

3. Designing and interpreting 'multi-omic' experiments that may change our understanding of

biology - PMC. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7477987/ (2024).

4. Cavicchioli, M. V., Santorsola, M., Balboni, N., Mercatelli, D. & Giorgi, F. M. Prediction of

Metabolic Profiles from Transcriptomics Data in Human Cancer Cell Lines. *Int. J. Mol. Sci.*

23, 3867 (2022).

5. Zelezniak, A. *et al.* Machine Learning Predicts the Yeast Metabolome from the Quantitative

Proteome of Kinase Knockouts. *Cell Syst.* **7**, 269-283.e6 (2018).

6. Ortmayr, K., Dubuis, S. & Zampieri, M. Metabolic profiling of cancer cells reveals genome-

wide crosstalk between transcriptional regulators and metabolism. *Nat. Commun.* **10**, 1841

(2019).

7. Gonçalves, E. *et al.* Systematic Analysis of Transcriptional and Post-transcriptional

Regulation of Metabolism in Yeast. *PLoS Comput. Biol.* **13**, e1005297 (2017).

8. Brunk, E. *et al.* Characterizing posttranslational modifications in prokaryotic metabolism

using a multiscale workflow. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 11096–11101 (2018).

9. Covert, M. W., Knight, E. M., Reed, J. L., Herrgard, M. J. & Palsson, B. O. Integrating high-

throughput and computational data elucidates bacterial networks. *Nature* **429**, 92–96

(2004).

10. Chandrasekaran, S. & Price, N. D. Probabilistic integrative modeling of genome-scale

metabolic and regulatory networks in Escherichia coli and Mycobacterium tuberculosis.

Proc. Natl. Acad. Sci. U. S. A. **107**, 17845–17850 (2010).

11. Daran-Lapujade, P. *et al.* The fluxes through glycolytic enzymes in Saccharomyces

cerevisiae are predominantly regulated at posttranscriptional levels. *Proc. Natl. Acad. Sci.*

U. S. A. **104**, 15753–15758 (2007).

12. Oliveira, A. P. *et al.* Regulation of yeast central metabolism by enzyme phosphorylation.

Mol. Syst. Biol. **8**, 623 (2012).

13. Shen, F., Boccuto, L., Pauly, R., Srikanth, S. & Chandrasekaran, S. Genome-scale network

model of metabolism and histone acetylation reveals metabolic dependencies of histone

deacetylase inhibitors. *Genome Biol.* **20**, 49 (2019).

14. Zaslaver, A. *et al.* Just-in-time transcription program in metabolic pathways. *Nat. Genet.* **36**,

486–491 (2004).

15. Zhao, S. *et al.* Regulation of Cellular Metabolism by Protein Lysine Acetylation. *Science*

327, 1000–1004 (2010).

16. Lempp, M. *et al.* Systematic identification of metabolites controlling gene expression in E.

coli. *Nat. Commun.* **10**, 4463 (2019).

17. Yu, L. *et al.* Modeling the Genetic Regulation of Cancer Metabolism: Interplay between

Glycolysis and Oxidative Phosphorylation. *Cancer Res.* **77**, 1564–1574 (2017).

18. Jia, D. *et al.* Elucidating cancer metabolic plasticity by coupling gene regulation with

metabolic pathways. *Proc. Natl. Acad. Sci.* **116**, 3909–3918 (2019).

19. Badia-i-Mompel, P. *et al.* Gene regulatory network inference in the era of single-cell multi-

omics. *Nat. Rev. Genet.* **24**, 739–754 (2023).

20. Hecker, M., Lambeck, S., Toepfer, S., van Someren, E. & Guthke, R. Gene regulatory

network inference: Data integration in dynamic models—A review. *Biosystems* **96**, 86–103

(2009).

21. Huynh-Thu, V. A., Irrthum, A., Wehenkel, L. & Geurts, P. Inferring Regulatory Networks from

Expression Data Using Tree-Based Methods. *PLOS ONE* **5**, e12776 (2010).

22. Rychel, K. *et al.* iModulonDB: a knowledgebase of microbial transcriptional regulation

derived from machine learning. *Nucleic Acids Res.* **49**, D112–D120 (2021).

23. Hu, T. *et al.* Single-cell spatial metabolomics with cell-type specific protein profiling for tissue

systems biology. *Nat. Commun.* **14**, 8260 (2023).

24. Patti, G. J., Yanes, O. & Siuzdak, G. Metabolomics: the apogee of the omics trilogy. *Nat.*

Rev. Mol. Cell Biol. **13**, 263–269 (2012).

25. Alexandrov, T. Spatial Metabolomics and Imaging Mass Spectrometry in the Age of Artificial

Intelligence. *Annu. Rev. Biomed. Data Sci.* **3**, 61–87 (2020).

26. Liu, Y. *et al.* High-plex protein and whole transcriptome co-mapping at cellular resolution

with spatial CITE-seq. *Nat. Biotechnol.* **41**, 1405–1409 (2023).

27. Zhao, M., He, W., Tang, J., Zou, Q. & Guo, F. A comprehensive overview and critical

evaluation of gene regulatory network inference technologies. *Brief. Bioinform.* **22**, bbab009

(2021).

28. Barretina, J. *et al.* The Cancer Cell Line Encyclopedia enables predictive modelling of

anticancer drug sensitivity. *Nature* **483**, 603–607 (2012).

29. Shoemaker, R. H. The NCI60 human tumour cell line anticancer drug screen. *Nat. Rev.*

Cancer **6**, 813–823 (2006).

30. Ahn, E., Kumar, P., Mukha, D., Tzur, A. & Shlomi, T. Temporal fluxomics reveals oscillations

in TCA cycle flux throughout the mammalian cell cycle. *Mol. Syst. Biol.* **13**, 953 (2017).

31. Schwabe, D., Formichetti, S., Junker, J. P., Falcke, M. & Rajewsky, N. The transcriptome

dynamics of single cells during the cell cycle. *Mol. Syst. Biol.* **16**, e9946 (2020).

32. Icard, P. *et al.* Understanding the Central Role of Citrate in the Metabolism of Cancer Cells

and Tumors: An Update. *Int. J. Mol. Sci.* **22**, 6587 (2021).

33. Li, L. *et al.* YAP/ACSL4 Pathway-Mediated Ferroptosis Promotes Renal Fibrosis in the

Presence of Kidney Stones. *Biomedicines* **11**, 2692 (2023).

34. Jyotsana, N., Ta, K. T. & DelGiorno, K. E. The Role of Cystine/Glutamate Antiporter

SLC7A11/xCT in the Pathophysiology of Cancer. *Front. Oncol.* **12**, (2022).

35. Okazaki, S. *et al.* Synthetic lethality of the ALDH3A1 inhibitor dyclonine and xCT inhibitors in

glutathione deficiency-resistant cancer cells. *Oncotarget* **9**, 33832–33843 (2018).

36. Li, H. *et al.* miR-142-5p Inhibits Cell Invasion and Migration by Targeting DNMT1 in Breast

Cancer. *Oncol. Res.* **28**, 885–897 (2022).

37. Zhai, Z. *et al.* MiR-181a-5p facilitates proliferation, invasion, and glycolysis of breast cancer

through NDRG2-mediated activation of PTEN/AKT pathway. *Bioengineered* **13**, 83–95

(2022).

38. Brunk, E. *et al.* Recon3D enables a three-dimensional view of gene variation in human

metabolism. *Nat. Biotechnol.* **36**, 272–281 (2018).

39. Benedetti, E. *et al.* A multimodal atlas of tumour metabolism reveals the architecture of

gene–metabolite covariation. *Nat. Metab.* **5**, 1029–1044 (2023).

40. Zhang, S. *et al.* Regulation of mTORC1 by amino acids in mammalian cells: A general

picture of recent advances. *Anim. Nutr.* **7**, 1009–1023 (2021).

41. Reid, M. A., Dai, Z. & Locasale, J. W. The impact of cellular metabolism on chromatin

dynamics and epigenetics. *Nat. Cell Biol.* **19**, 1298–1306 (2017).

42. Su, X., Wellen, K. E. & Rabinowitz, J. D. Metabolic control of methylation and acetylation.

Curr. Opin. Chem. Biol. **30**, 52–60 (2016).

43. King, J., Patel, M. & Chandrasekaran, S. Metabolism, HDACs, and HDAC Inhibitors: A

Systems Biology Perspective. *Metabolites* **11**, 792 (2021).

44. Calvo, F. *et al.* Mechanotransduction and YAP-dependent matrix remodelling is required for

the generation and maintenance of cancer-associated fibroblasts. *Nat. Cell Biol.* **15**, 637–

646 (2013).

45. Edwards, D. N. *et al.* Regulation of cancer glutamine metabolism by EphA2 RTK-dependent

activation of transcriptional co-activators YAP and TAZ. *Sci. Signal.* **10**, eaan4667 (2017).

46. DepMap Data Explorer 2.0. https://depmap.org/portal/data_explorer_2/ (2024).

47. Skoufos, G. *et al.* TarBase-v9.0 extends experimentally supported miRNA–gene interactions

to cell-types and virally encoded miRNAs. *Nucleic Acids Res.* **52**, D304–D310 (2024).

48. Oughtred, R. *et al.* The BioGRID database: A comprehensive biomedical resource of

curated protein, genetic, and chemical interactions. *Protein Sci. Publ. Protein Soc.* **30**, 187–

200 (2021).

49. Couto, J. P. *et al.* Nicotinamide N-methyltransferase sustains a core epigenetic program that

promotes metastatic colonization in breast cancer. *EMBO J.* **42**, e112559 (2023).

50. Liang, G. *et al.* Nicotinamide N-methyltransferase and liver diseases. *Genes Dis.* **10**, 1883–

1893 (2022).

51. Szklarczyk, D. *et al.* STRING v11: protein–protein association networks with increased

coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic*

Acids Res. **47**, D607–D613 (2019).

52. Wang, J. *et al.* SynLethDB 2.0: a web-based knowledge graph database on synthetic

lethality for novel anticancer drug discovery. *Database J. Biol. Databases Curation* **2022**,

baac030 (2022).

53. PTPN14 interacts with and negatively regulates the oncogenic function of YAP | Oncogene.

https://www.nature.com/articles/onc2012147 (2024).

54. Liu, H. *et al.* DrugCombDB: a comprehensive database of drug combinations toward the

discovery of combinatorial therapy. *Nucleic Acids Res.* **48**, D871–D881 (2020).

55. Wishart, D. S. *et al.* DrugBank 5.0: a major update to the DrugBank database for 2018.

Nucleic Acids Res. **46**, D1074–D1082 (2018).

56. You, X. *et al.* Metabolic reprogramming and redox adaptation in sorafenib-resistant

leukemia cells: detected by untargeted metabolomics and stable isotope tracing analysis.

Cancer Commun. **39**, 17 (2019).

57. Chatzispyrou, I. A., Held, N. M., Mouchiroud, L., Auwerx, J. & Houtkooper, R. H.

Tetracycline antibiotics impair mitochondrial function and its experimental use confounds

research. *Cancer Res.* **75**, 4446–4449 (2015).

58. Ariaans, G. *et al.* Everolimus decreases [U-13C]glucose utilization by pyruvate carboxylase

in breast cancer cells in vitro and in vivo. *Biomed. Pharmacother. Biomedecine*

Pharmacother. **173**, (2024).

59. Nanda, S. *et al.* Systems-level transcriptional regulation of Caenorhabditis elegans

metabolism. *Mol. Syst. Biol.* **19**, e11443 (2023).

60. Quantifying absolute gene expression profiles reveals distinct regulation of central carbon

metabolism genes in yeast | eLife. https://elifesciences.org/articles/65722 (2024).

61. Yugi, K., Kubota, H., Hatano, A. & Kuroda, S. Trans-Omics: How To Reconstruct

Biochemical Networks Across Multiple 'Omic' Layers. *Trends Biotechnol.* **34**, 276–290

(2016).

62. Shorstova, T. *et al.* Reprogramming of Nucleotide Metabolism Mediates Synergy between

Epigenetic Therapy and MAP Kinase Inhibition. *Mol. Cancer Ther.* **20**, 64–75 (2021).

63. Lobritz, M. A. *et al.* Antibiotic efficacy is linked to bacterial cellular respiration. *Proc. Natl.*

Acad. Sci. **112**, 8173–8180 (2015).

64. Diehl, F. F. *et al.* Nucleotide imbalance decouples cell growth from cell proliferation. *Nat.*

Cell Biol. **24**, 1252–1264 (2022).

65. Nusinow, D. P. *et al.* Quantitative Proteomics of the Cancer Cell Line Encyclopedia. *Cell*

180, 387-402.e16 (2020).

66. Li, H. *et al.* The landscape of cancer cell line metabolism. *Nat. Med.* **25**, 850–860 (2019).

67. Dashboard - Genomics Biotools. https://www.biotools.fr/ (2024).

68. King, Z. A. *et al.* BiGG Models: A platform for integrating, standardizing and sharing

genome-scale models. *Nucleic Acids Res.* **44**, D515–D522 (2016).

69. Shankavaram, U. T. *et al.* CellMiner: a relational database and query tool for the NCI-60

cancer cell lines. *BMC Genomics* **10**, 277 (2009).

70. RandomForestRegressor. *scikit-learn* https://scikit-

learn/stable/modules/generated/sklearn.ensemble.RandomForestRegressor.html.

71. sklearn.linear_model.Ridge. *scikit-learn* https://scikit-

learn/stable/modules/generated/sklearn.linear_model.Ridge.html (2024).

72. Lasso. *scikit-learn* https://scikit-

learn/stable/modules/generated/sklearn.linear_model.Lasso.html.

73. XGBoost Python Package — xgboost 3.0.2 documentation.

https://xgboost.readthedocs.io/en/stable/python/index.html.

74. SVR. *scikit-learn* https://scikit-learn/stable/modules/generated/sklearn.svm.SVR.html (2024).

75. Breiman, L. Random Forests. *Mach. Learn.* **45**, 5–32 (2001).

76. An introduction to explainable AI with Shapley values — SHAP latest documentation.

https://shap.readthedocs.io/en/latest/example_notebooks/overviews/An%20introduction%20

to%20explainable%20AI%20with%20Shapley%20values.html (2024).

77. Lundberg, S. M. *et al.* Explainable machine-learning predictions for the prevention of

hypoxaemia during surgery. *Nat. Biomed. Eng.* **2**, 749–760 (2018).

78. Heirendt, L. *et al.* Creation and analysis of biochemical constraint-based models using the

COBRA Toolbox v.3.0. *Nat. Protoc.* **14**, 639–702 (2019).

1028

1029 **Supplementary Materials**

1030
Supplemental Figure 1: Results from training an array of ML models on each input dataset. Bar charts show the
number of significantly predicted (FDR p-value < 0.05) metabolites (of 225) per input data type, also represented as
a percentage out of 225. These algorithms did not significantly outperform Random forests, and therefore were not
selected for in-depth analysis. **A.** Random Forest results for comparison (also shown in Figure 2) **B.** Results from
Random Forest benchmarking experiments using the top 40 features per input dataset as determined using 75-25%
holdout validation. **C.** XGBoost **D.** Support Vector Machine (SVM) with RBF kernel. Highlighted in green are
results from binary mutation data **E.** Ridge regression **F.** Lasso regression
1038

1039

1040
Supplemental Figure 2: Shapley analysis for top 3 metabolites (lowest P value) for each omics class from random
forest models. Each dot on the Shapley beeswarm plots corresponds to a cell line observation. Red data points
indicate high feature values and blue points indicate low feature values. High SHAP values correspond with higher
metabolic concentration outputs from the model, and low SHAP values correspond with lower outputs. Therefore, a
gene with a high feature value and SHAP value (red and to the right) illustrates a positive correlation, and a high
feature value and a low SHAP value (red and to the left) illustrates a negative correlation. **A.** Genomics (CNV) **B.**
DNA methylation **C.** Histone PTMs **D.** Transcriptomics **E.** RNA Splicing **F.** miRNA **G.** lncRNA **H.** Proteomics **I.**
Phosphoproteomics
1049

A

Lung Cancers

Hematopoietic and Lymphoid Cancers

B

Adherent Cell Culture

Suspension Cell Culture

1052
Supplemental Figure 3: Control experiment results (random forests), showing the number of significant predictions
(FDR p-value < 0.05) out of 225 metabolite models (also represented as a percentage out of 225), the average
correlations across all models, and the number of cell lines in each data set (left to right). **A.** Models trained on
samples from particular cancer lineages. **B.** Models trained on samples grouped by culture conditions. **C.** Models
trained on samples grouped by tumor origin. **D.** Models trained on all cancer lineages excluding lung cancers or
excluding hematopoietic and lymphoid cancers.

1059
Supplemental Figure 4: Results from models trained on genes present in the Recon3D metabolic model. **A.**
Number of features in each omics class before and after filtering for genes present in the Recon3D metabolic model.
B. Number of significant predictions (FDR p-value < 0.05) from metabolic gene-only trained models, also
represented as a percentage out of 225. **C.** Box plots showing distribution of correlations (true vs predicted) from
models built on whole data versus metabolic gene-only models (right). **D.** Shapley analysis for top 3 metabolites
(lowest P value) from metabolic gene-only models identifying the top 8 most important features. Each dot on the
Shapley beeswarm plots corresponds to a cell line observation. Red data points indicate high feature values and blue
points indicate low feature values. High SHAP values correspond with higher metabolic concentration outputs from
the model, and low SHAP values correspond with lower outputs. Therefore, a gene with a high feature value and
SHAP value (red and to the right) illustrates a positive correlation, and a high feature value and a low SHAP value
(red and to the left) illustrates a negative correlation. **E.** Shapley analysis for top 3 metabolites (lowest P value) from
models built on whole data identifying the top 8 most important features. These show that feature importances from
metabolic gene-only models and whole-data models were conserved. For example, NNMT is the top predictor for 1-
methylnicotinamide, SLC6A8 for phosphocreatine, and DPYD for N-carbamoyl-beta-alanine in both the metabolic
gene-only and whole-data models. DPYD is involved in N-carbamoyl-beta-alanine biosynthesis.

Supplemental Figure 5: Integrating protein-protein, miRNA-gene and metabolite-gene interactions with Recon8D interactions. This figure presents a comparative analysis of relationships between top feature factors derived from Recon8D multi-omics data, including transcriptomics, proteomics, phosphoproteomics, genomics, methylation, miRNA, splicing and epigenomics, and their corresponding metabolites. Top features from each omics dataset are organized in distinct, color-coded circular layouts surrounding a central circle of metabolites. The size of each node reflects the number of metabolites associated with that feature. Edges between top features represent correlations in their metabolite-consistency profiles, with edge width corresponding to the strength of the correlation. This figure presents a comparative analysis of low consistency relationships between top feature factors derived from multi-omics data. Edges connecting top feature factors to metabolites illustrate the variability (in contrast to Figure 5, which shows consistency) in feature importance and topological distribution across the datasets. The figure demonstrates how feature importance and topological distribution can diverge (low consistency) across multi-omics data. For example, Yap1 (phosphoproteomics) shows low consistency with 60 metabolites based on distance of Yap1 to these metabolites in the PPI/miRNA/Recon3D networks and their feature importance in Recon8D.

1090
1091 **Supplemental Figure 6:** Proposed metabolic regulome network from top 8 features from miRNA, transcriptomic, and
1092 proteomic-trained models for predicting phosphocreatine (highest mean Shap values) as determined by Shapley analysis.

Supplemental Table 1: Number of cell lines used to train each model

	Genomics	Mutations	DNA methylation	Histone PTMs	Transcript-omics	RNA Splicing	miRNA	lncRNA	Proteomics	Phospho-proteomics
Cell Lines	921	925	828	893	904	900	916	916	373	296

1094
1095
Supplemental Table 2: Top 40 global features as identified from 75-25% holdout validation random forests.

Genomics (CNV)	Histone PTMs	DNA methylation	Transcriptomics	RNA Splicing	miRNA	lncRNA	Proteomics	Phosphoproteomics
LEP1P4	H3K9me3K14ac0	CAPR2	FAM18A	EPHA2	hsa-miR-142-8p	MIR22HG	PLIN2	YAP1
ADAM9	H3K27me0K36me0	CISM	KLF13	CLSTN1	hsa-miR-642-3p	TSPOAP1-AS1	PTPN13	BRAF
FAM20A	H3.3K27me0K36me0	MIR142	MbO	PLEKHB2	hsa-miR-22	SNHG4	CEP290	PIK3CA
TIMP3	H3K4me2	CFTR	SCD	ACOT7	hsa-miR-156	LINC00625	HSPG2	RB1
MEF2C-AS1	H3K27me0K36me0	HDAC7	INSIG1	RPL35	hsa-miR-186	FOXP4-AS1	FUNDC2	ERBB3
FAM45B	H3K27me1K36me3	YAP1	YAP1	L3C	hsa-miR-375	DLGAP1-AS1	TNS21	BRC
MIR4426LS	H3K4me1	TCF3	DNAC1	CLSTN1	hsa-miR-24	LINC00335	PLEK	BIK4EBP1
BTNL3	H3K9ac1K14ac0	SLC2AM	RNLS	DVL1	hsa-miR-31	MIR199-2HG	MARCKS	ADC
VMD1	H3K27ac1K36me0	EPHA2	MT3A	NIPB	hsa-miR-22a	DLGAP1-AS1	DENND4A	BAD
CTSW	H3K18ac0K28ac1	DENND4B	DURPH2	MINCR1	hsa-miR-21	SNHG18	RCBD1	ERBB2
ILRAS	H3K4ac1	UMCLL1-AS1	PLIN2	S2RD1	hsa-miR-130a	HOTAIRM1	RAD17	ERALPHA
LINC0321	H3K27me1K36me2	LANCL2	TRAF3IP2	GNB1	hsa-miR-27b	PAXB-AS1	TREX1	MTOR
SEMA5D	H3K9me2K14ac1	PCF11	SSFN	ICMT	hsa-miR-574-3p	CEMULINC	UTP23	EGR1
UGT3B17	H3K27me0K36me2	VAUP1	EDP	PARK7	hsa-miR-146a	SNHG9	COBLL1	MFK3P65
PTGS2	H3K9me1K14ac0	MIR9797	TNFRSF11A	MIP	hsa-miR-223	DDN-AS1	FLAD1	MAPK
FAM20C	H3K79me1	KIAA1522	CCDC80	NADK	hsa-miR-574-5p	WDFY3-AS2	AKR1B15	AMPK
MIR8196	H3K9me1K14ac0	AMBRA1	CFT1A	S2RD1	hsa-miR-342-3p	LINC01270	TP53	S6
RRG1CP	H3K9me2	PRDM1	B6MR2L	PLOD1	hsa-miR-155	WAITR1-AS1	RBS3	MYO1A
AC244502.1	H3K9me3K14ac1	ERBB2	NOS3	CAMTA1	hsa-miR-28a	LINC02449	LHPP	STAT3
HSP1	H3K79me6	TFDP1	HISPAE	PLEKHB2	hsa-miR-195b	PWE1-AS	CCBL2	T8C2
PR3S1	H3K04me6	GULP1	68R	CDK11A	hsa-miR-221	AGAP3-AS1	ENDOB	MAP2K1
MIR439A	H3K27ac0K36me0	THDC6	ST6GALNAC4	GNB1	hsa-miR-1078	HCG11	PPP1R12A	JUN
DUJAP5	H3K9me2K14ac0	TMEM75	RIDA	MIFN2	hsa-miR-425-3p	SNHG15	ACSF3	AKT1
NPS9	H3K27me0K36me1	MIR3178	AMDTL2	SLC25E2B	hsa-miR-376	H19	STP8P8	FOX03
TGIF3LY	H3K27me0K36me3	SHRDCMT	PTGES2	UREL2	hsa-miR-15b	SNHG5	ATP5G2	FOPK1
MTAP	H3K27me0K36me1	WDR5	YIFF1	GNB1	hsa-miR-103a+hsa-miR-107	DUBR	FADB2	P27
TBC1D3J	H3K9me0K14me1	ARRB2	H1-4	CLSTN1	hsa-miR-218	LINC01189	MCALL2	RAF1
CDKN2A	H3K18ac0K23ac3	MGTE2P	CDC42EP3	RPL22	hsa-miR-148	LINC01869G	MAN2B2	MAPK14
SLDR	H3K36me6	LOC100t26531	PHPT1	CAMTA1	hsa-miR-766	LINC00472	ATXN1	BCATENIN
AIOUS	H3K9me1K14ac1	MGA	MTAP	CCBL2	hsa-miR-193b	MIR100HG	SRX11	YBK1
ACOT1	H3K18ac1K23ac3	WAV1	ANKRD37	GNB1	hsa-miR-1274a	LINC01639	IRF9	GSK3ALPHABETA
MIR3975	H3K18ac0K23ac1	MIR3772	SGLE	MINC91	hsa-miR-95	SNHG9	ZFYVE20	CHEK1
TBC1D3F	H3K9ac1K14ac1	SDNAP2	RPL38AL	LINC01238	hsa-miR-181b	STN1-AS1	SFK	MAPK8
MIR4375	H3K27me0K36me2	MFPS0	COA4	NECAP2	hsa-miR-9	SUCLG2-AS1	GMPR2	GSK3
OR11H1	H3K27ac1K36me0	SYDE1	PAOX	MRPL20	hsa-miR-183	UFF-AS2	NTSC2A	RPG3SH
ALAS2FR-1	H3K27me1K36me2	BTX76-NIPEPL1	NDUFAP	MINC81	hsa-miR-2b	SNHG12	PHLDB2	NDRG1
TRB-AS1	H3K18ac1K23me1	ZNF61X	CURBAP2	MIFN3	hsa-miR-182	Z8F718-AS1	CRASP1	MET
POTEH	H3K27me0K36me6	TME3014C	SH3BP1	PEX19	hsa-miR-695	LUCAT1	M8PR	SHC1
SIGLEO14	H3K27me1K36me1	FRYD1	AIR	ICMT	hsa-miR-620-8p	FOXO3-AS1	FG31	PHOSPHAMBETA6
CHES2	H3K27me0K36me1	DOB1	CRABINT	A3PT3AB	hsa-miR-395	SNHG27	OTUD4	EPH6

1097

Supplemental Table 3: Robustness of top features from CCLE transcriptomics in the NCI60 data for full data
models and Recon3D genes only. T-test and Kolomogorov-Smirnov test results from the top 1, 3, 5, 10, 50, and 100
CCLE features isolated in NCI60 data as compared to the whole data. Top features for predicting each metabolite
were tested against the whole data for the entire metabolite panel (pooled). Data were stacked to create a single
vector of feature importance scores across all metabolites for the top feature subsets as well as whole data. Feature
importances were used to assess the robustness of the random forest ML framework used to train the CCLE models.
Feature importances were significantly higher for all top feature subsets in the pooled analyses using
transcriptomics. Top features for individual metabolites also showed significantly higher correlations and feature
importance scores for a subset of metabolites (FDR p-value < 0.05). Only using the top 1 feature, CMP, sucrose,
GABA, anthranilic acid, N-carbamoyl-beta-alanine, and adenosine displayed conserved top features, which are all in
peripheral metabolism. * = significant (p-value < 0.05)

| | Whole-data | | Recon3D-only | |
| | T-test | KS-test | T-test | KS-test |
Top Features	P value	P value	P value	P value
1	2.54E-05*	6.50E-02	3.02E-02*	5.2E-02
3	2.51E-05*	7.30E-02	2.48E-02*	6.7E-02
5	2.22E-06*	4.30E-02*	1.73E-02*	2.9E-02*
10	4.23E-10*	2.90E-03*	4.12E-03*	2.4E-02*
50	1.44E-10*	4.60E-03*	8.05E-07*	4.28E-06*
100	2.92E-17*	3.68E-05*	3.69E-13*	1.52E-10*

1109
1110
1111 **Supplemental Table 4:** Robustness of top features from CCLE genomics, miRNA, proteomics, and
1112   phosphoproteomics in NCI60 models illustrated by T-test and Kolmogorov-Smirnov tests using pooled feature
1113   importances. * = significant (p-value < 0.05)

| | Genomics | | miRNA | | Proteomics | | Phosphoproteomics | |
| | T-test | KS-test | T-test | KS-test | T-test | KS-test | T-test | KS-test |
Top Features	P value	P value	P value	P value	P value	P value	P value	P value
1	1.16E-01	5.64E-01	8.80E-01	9.3E-01	2.35E-01	5.7E-01	8.5E-01	9.7E-01
3	2.40E-02*	3.39E-01	4.00E-01	6.6E-01	1.54E-01	4.2E-01	7.7E-2	3.5E-01
5	2.40E-02*	3.46E-01	7.10E-02	1.8E-01	6.80E-02	1.5E-01	9.70E-04*	1.99E-02*
10	1.80E-03*	2.06E-01	7.62E-07*	1.91E-05*	8.80E-02	2.96E-01	1.00E-04*	8.00E-03*
50	4.75E-07*	7.66E-03*	4.79E-07*	3.46E-23*	2.23E-02*	1.88E-01	2.05E-11*	5.75E-08*
100	5.06E-12*	6.10E-04*	2.06E-33*	6.75E-26*	2.60E-03*	1.05E-01	N/A	N/A

1114
1115

1116 **Supplemental Table 5:** Robustness of top features from CCLE transcriptomics in single-cell models illustrated by
1117 T-test and Kolmogorov-Smirnov tests using pooled feature importances. * = significant (p-value < 0.05)

Top Features	T-test	KS-test
	P value (pooled)	P value (pooled)
1	2.00E-01	4.9E-01
3	2.10E-02*	1.6E-01
5	6.00E-04*	2.6E-02*
10	6.06E-08*	3.80E-04*
50	8.74E-16*	3.80E-07*
100	2.12E-25*	4.36E-11*

1118
1119
1120 **Supplemental Table 6:** Count of occurrences of mutations in the top 20 features (calculated using mutual
1121 information) for each of 225 metabolites.

Mutation	Count
TP53 (7157)	20
APC (324)	19
CDKN2A (1029)	17
ARID1A (8289)	17
PTEN (5728)	16
KMT2D (8085)	15
TTN (7273)	14
RB1 (5925)	14
ACVR2A (92)	13
MUC16 (94025)	12
ZNF141 (7700)	9
KRTAP21-1 (337977)	8
PCLO (27445)	8
GPR98 (84059)	8
RPL22 (6146)	8
ASXL1 (171023)	8
NF2 (4771)	8
CCDC73 (493860)	8
TEAD2 (8463)	8
CREBBP (1387)	8

1122

1123   **Supplemental Table 7:** Predicted non-lipid metabolites for which mutations of interest were in the top 20 features.

Mutation	Predicted Metabolites
TP53	CMP, dimethylglycine, kynurenic acid, adenosine, deoxycytidine, carnitine, thymidine, GMP, creatinine, arachidonylcarnitine, adenine, creatine
APC	N-carbamoyl beta alanine, betaine, creatine, taurine, NMMA, arachidonylcarnitine, pantothenate, tyrosine, alpha-glycerophosphocholine, phosphocreatine, carnitine, acetylcarnitine, histidine, alpha-hydroxybutyrate, UMP, GABA
CDKN2A	Creatinine, NMMA, dimethylglycine, allantoin, UMP, xanthine, hexoses, CMP
PTEN	Phenylalanine, carnosine, oleylcarnitine, kynurenic acid, pantothenate, glutathione oxidized, anserine, threonine, leucine
RB1	Isoleucine, lactose, phosphocreatine, phenylalanine, creatine, arginine, adenosine
MUC16	Putrescine, hydroxybutyrate, isoleucine, dimethylglycine, glutathione reduced, creatine, NADP, acetylcarnitine, phosphocreatine

1124
1125

1126 **Supplemental Table 8:** High confidence lncRNAs and their respective predicted metabolite prioritized for further
1127 study. lncRNAs were selected based on feature importance (in the top 3 most important features) and confidence
1128 score (6 or greater).

Metabolite	lncRNA	Confidence Score
Cystathionine	LINC01588 (ENSG00000214900)	6
NADP	LINC00942 (ENSG00000249628)	6
2-hydroxyglutarate	LINC01164 (ENSG00000189275)	6
Inositol	SEC24B-AS1 (ENSG00000247950)	6
Lysine	MIR22HG (ENSG00000186594)	6
Proline	SNHG4 (ENSG00000281398)	7
N-carbamoyl-beta-alanine	MIR137HG (ENSG00000225206)	6
1-methylnicotinamide	ITGB1-DT (ENSG00000229656)	6
Carnitine	MIR3936HG (ENSG00000233006)	7
Acetylcarnitine	MIR3936HG (ENSG00000233006)	7
Propionylcarnitine	MIR3936HG (ENSG00000233006)	6
Malonylcarnitine	MIR3936HG (ENSG00000233006)	6
Valerylcarnitine	MIR3936HG (ENSG00000233006)	8
Hexanoylcarnitine	MIR3936HG (ENSG00000233006)	6
Beta-alanine	LUCAT1 (ENSG00000248323)	6
C34:4 PC	LINC00973 (ENSG00000240476)	6
C46:1 TAG	MIR22HG (ENSG00000186594)	6
C48:2 TAG	LINC01358 (ENSG00000237352)	7
C50:3 TAG	LINC01358 (ENSG00000237352)	7
C50:2 TAG	LINC01358 (ENSG00000237352)	7

1129

1130 **Supplemental Table 9:** Number of metabolite models trained on the Recon3D metabolic gene subset that had at
1131 least one top feature map to Recon3D for the reactions directly associated with or within one neighbor of the
1132 predicted metabolite (out of 95 common metabolites). Random analyses (last row) with transcriptomics data
1133 involved picking random transcripts per metabolite model and seeing if at least one feature mapped to Recon3D.
1134 This process was repeated 50 times, and the median number of overlaps was calculated for each category.

Recon3D Genes	Total Model Training Features	Top Feature		Top 2 Features		Top 5 Features	
		Primary Reaction	1st degree neighbor	Primary Reaction	1st degree neighbor	Primary Reaction	1st degree neighbor
Genomics	1,817	1	56	2	78	7	88
Methylation	1,578	8	60	10	75	16	90
Proteomics	1,375	20	74	22	83	27	89
Transcriptomics	1,659	16	77	23	90	32	92
Transcriptomics (random)	1,659	1	57	2	76	4	87

1135
1136

1137 **Supplemental Table 10:** Global feature importance ranks for the top 20 features from each omics class from
1138 random forests (except mutations, which used SVMs). Median rank was used to determine the most important
1139 features over all 225 metabolite models for each feature.

Global Average Rank	Gene	Mutations	Hist	Methyl	Transc	Splice	miRNA	lncRNA	Prot	Phos
1	LSP1P4	RB1	H3K9me3K14ac0	CAPN2	YAP1	EPHA2	hsa-miR-142-5p	MIR22HG	LAMC1	YAP1
2	FAM30A	CDKN2A	H3K27me2K36me0	OSM	GPX8	CLSTN1	hsa-miR-142-3p	TSPOAP1-AS1	PURA	PRKCA
3	ADAM6	ASXL1	H3.3K27me0K36me0	CTTN	SCD	PLEKHM2	hsa-miR-22	LINC02535	PLIN2	BRAF
4	FRG1CP	NF2	H3K27me1K36me3	MIR142	DGAT2	CLSTN1	hsa-miR-150	LINC01186	FTL	ACC
5	MTRNR2L6	ESRP1	H3K4me1	HDAC7	PPIC	DVL1	hsa-miR-100	LINC01636	TLK1	BAD
6	LILRA6	PHF6	H3K27me3K36me0	LOC101080498	IKZF1	ACOT7	hsa-miR-31	SNHG4	HBA1	RB
7	H3Y1	STK11	H3K27me1K36me2	YAP1	KLF13	KIF1B	hsa-miR-27a	MIR100HG	COL6A3	ESR1
8	VMO1	ZNRF3	H3K4me2	C5orf66-AS2	FAM162A	MINOS1	hsa-miR-375	SNHG19	TSPAN6	MYO1A
9	TIMP3	NLRP13	H3K9ac1K14ac0	INPP5D	CLCF1	EPHA2	hsa-miR-23a	SNHG6	SLC39A14	ERALPHA
10	FAM242B	FILIP1L	H3K27me1K36me0	AMOTL2	CAV1	SZRD1	hsa-miR-24	SNHG8	GLUL	ERBB2
11	CTSW	USP26	H3K27me2K36me2	SHROOM1	FAM53B	LZIC	hsa-miR-98	PINK1-AS	ZFYVE16	SRC
12	PRSS1	SCN7A	H3K9me1K14ac0	THOC6	PPARD	RPL22	hsa-miR-222	SNHG7	PSTPIP2	STAT3
13	FAM230C	PAX5	H3K18me0K23me1	NRROS	CCN1	GNB1	hsa-miR-183	FOXP4-AS1	ECHDC3	ERBB3
14	MIR4436A	MTA2	H3K9me2K14ac1	GTPBP2	EPHA2	RERE	hsa-miR-183b	LINC00528	TWSG1	MTOR
15	AC244502.1	CCDC73	H3K4ac1	HIST1H4G	TNFRSF12A	GNB1	hsa-miR-342-3p	LPP-AS2	LRPAP1	MAPK
16	PRSS2	COL13A1	H3K27me0K36me2	ERBB2	IRF2BP2	GNB1	hsa-miR-148a	FOXD2-AS1	NMES1	BCATENIN
17	LINC00221	SPEG	H3K79me1	LPCAT1	F12	MFN2	hsa-miR-674-3p	USP2-AS1	PTPN6	NFKBP65
18	CDKN2A	ANKRD11	H3K9me0K14ac0	ARRB2	WWTR1	RPL22	hsa-miR-29a	KTN1-AS1	RAB13	MAP2K1
19	DLXAP6	GNA13	H3K27me3K36me1	WIZ	POLR1H	ICMT	hsa-miR-221	OLMALINC	STX7	EIF4EBP1
20	SEMA6D	GLTSCR1	H3K79me0	ZFAT	BATF2	MINOS1	hsa-miR-455-3p	SNHG17	LAMB2	FOXO3

1140

1141 **Supplemental Table 11:** Correlation of two protein isoforms of YAP1 with metabolites for which ML found YAP1
1142 to be the top feature from transcriptomics inputs. Correlations were obtained from proteomic and metabolomic
1143 CCLE data using the DepMap portal[46]. Associations between YAP1 and the metabolites it predicted well from ML
1144 were assessed for statistical significance using a Bonferroni corrected P value (marked with a *).

Metabolite	YAP1 (P46937-7)		YAP1 (P46937-8)	
	Pearson's R	P value	Pearson's R	P value
aconitate	0.117	3.08E-02	0.159	2.11E-03
dCMP	-0.212	7.60E-05*	-0.302	2.97E-09*
hypoxanthine	-0.134	1.33E-02	-0.122	1.90E-02
oxalate	-0.325	7.02E-10*	-0.377	5.92E-14*
uracil	0.249	2.94E-06*	0.3	3.93E-09*
xanthine	0.34	1.04E-10*	0.437	1.13E-18*
3-methyladipate/pimelate	-0.112	3.80E-02	-0.136	8.75E-03
α-hydroxybutyrate	-0.204	1.39E-04*	-0.258	5.12E-07*
Met	0.241	6.61E-06*	0.356	1.59E-12*
Leu	0.245	4.27E-06*	0.363	5.34E-13*
Ile	0.187	4.94E-04*	0.297	5.97E-09*
Phe	0.231	1.59E-05*	0.338	2.41E-11*
Tyr	0.204	1.45E-04*	0.308	1.35E-09*
Trp	0.16	2.97E-03*	0.261	3.72E-07*
dimethylglycine	0.311	3.87E-09*	0.368	2.69E-13*
SDMA/ADMA	0.326	6.55E-10*	0.409	2.22E-16*
NMMA	0.333	2.68E-10*	0.403	7.12E-16*
allantoin	0.28	1.42E-07*	0.372	1.36E-13*
kynurenic acid	0.38	3.24E-13*	0.427	8.61E-18*
carnosine	0.352	1.85E-11*	0.39	6.57E-15*
creatinine	0.324	7.93E-10*	0.386	1.46E-14*
methionine sulfoxide	0.291	4.27E-08*	0.346	7.71E-12*
propionylcarnitine	0.188	4.67E-04*	0.179	5.52E-04*
C32:0 PC	-0.248	3.23E-06*	-0.222	1.68E-05*
C34:1 PC	-0.246	4.19E-06*	-0.28	4.46E-08*
C38:4 PC	-0.38	3.04E-13*	-0.352	2.99E-12*
C40:6 PC	-0.306	7.46E-09*	-0.298	5.25E-09*
C18:2 SM	-0.179	8.61E-04*	-0.238	3.74E-06*
C16:1 SM	0.213	7.07E-05*	0.252	9.27E-07*
C16:0 CE	0.399	1.62E-14*	0.453	4.09E-20*
C18:2 CE	0.317	1.82E-09*	0.401	9.55E-16*
C18:1 CE	0.349	3.06E-11*	0.416	7.07E-17*
C18:0 CE	0.224	2.75E-5*	0.255	6.51E-7*
C20:4 CE	0.334	2.25E-10*	0.424	1.43E-17*
C56:8 TAG	-0.313	3.19E-09*	-0.37	1.82E-13*

1145