1 **Title -** Species-specific interactions with apex carnivores yield unique benefits and burdens for
2 mesocarnivores
3

4 **Authors and Affiliations**:
5 Wesley Binder[1,2]*, Jack W. Rabe[2,3]*, Zoe Lowe[4], Gordon Scott[2], Claire Lacey[2], Eliza King[4],
6 Daniel R. Stahler[2], Stotra Chakrabarti[4]
7 *Contributed Equally
8
[1]Department of Fisheries, Wildlife, and Conservation Sciences, Oregon State University,
Corvallis, Oregon, USA
[2]Yellowstone Center for Resources, Yellowstone National Park, Mammoth Hot Springs,
Wyoming, USA
[3]Department of Fisheries, Wildlife, and Conservation Biology, University of Minnesota – Twin
Cities, Saint Paul, Minnesota, USA
[4]Department of Biology, Macalester College, Saint Paul, Minnesota, USA
16

Author Contributions:
WB, JWR, DRS, and SC conceptualized the study, raised resources, and oversaw project
completion. WB, DRS, CL, and GS conducted fieldwork and collected the data. WB, JWR, and
SC developed the methodology and analytical approaches. WB, JWR, ZL, GS, and EK curated
the data and performed the analyses, with inputs from SC and DRS. WB, JWR, and SC wrote the
manuscript, with help from GS. All authors reviewed subsequent drafts and approved the final
version.
24
Statement on Inclusion:
Our study brings together authors from diverse career stages, including scientists based in the
country where the study was carried out. All authors were engaged early on with the research to
ensure that the diverse sets of perspectives they represent was considered from the onset.
Whenever relevant, literature published by scientists from the region was cited.

Acknowledgements:
We are grateful to the numerous field technicians from the Yellowstone Wolf and Cougar
Projects who made data collection possible. We would also like to thank the fStop Foundation
for providing necessary equipment. We acknowledge Yellowstone Forever (and their
contributing donors), the National Science Foundation, Oregon State University, University of
Minnesota, National Park Service, US Geological Survey, and Macalester College for funding
support.
37

38 **Conflict of interest:**
39 The authors declare no conflicts of interest.
40

ABSTRACT

1. Mesocarnivores must balance the negative (competition) and positive (carrion

provisioning) effects of dominant apex carnivores to coexist in shared landscapes. How

mesocarnivores navigate such top-down effects have typically been examined through a

risk-reward lens. However, the risks accrued by mesocarnivores might not be even across

all apex-mesocarnivore pairings in multi-carnivore systems, and instead can be mediated

through species-specific interactions.

2. Using remote-camera surveys from 2021–2023, we estimated apex (wolf, cougar) and

mesocarnivore (red fox, coyote) spatial overlap, occupancy, temporal activity, and fine-

scale behaviors in northern Yellowstone National Park, USA. Additionally, we monitored

GPS-collared wolf and cougar predation from 2016–2023 and recorded red fox and

coyote mortalities and scavenging events at these locations.

3. Red foxes and coyotes were ubiquitous in our study area with high naïve occupancy,

despite abundant wolves and cougars. Coyote occupancy was positively associated with

wolf occupancy, but exhibited marginal avoidance of cougars. Photo-detection

probabilities of both mesocarnivores doubled in the 24 hours following apex carnivore

detections, with the exception of coyotes following cougars. Overall latency of coyote

detections was longest following cougar detections and shortest following wolves, while

foxes exhibited no such difference. Further, coyotes were most active when wolves and

cougars were collectively least active. Red foxes were nocturnal and appeared less

sensitive to wolf and cougar activities, but avoided coyote peak times.

4. Coyotes were more prevalent scavengers than red foxes at apex carnivore kills, but

scavenging often had lethal consequences. We found that when wolves kill coyotes, it

was typically when coyotes attempted to scavenge from wolf kills, while cougars killed

and consumed coyotes as prey. Thus wolves act as competitors for coyotes, while

cougars are both predators and competitors. This distinction possibly explains coyote

avoidance of cougars but attraction to wolves.

5. Mesocarnivores in Yellowstone leverage foraging benefits from associating with apex

carnivores. However, these benefits are burdened by associating with wolves and cougars

whose contrasting hunting modes and habitat preferences yield separate risks. Disparate

costs to mesocarnivores contingent upon apex carnivore behavior offer novel insights

into carnivore community ecology, and call for a more nuanced analysis of trophic

interactions.

KEYWORDS: community ecology, intra-guild competition, landscape of fear, mesopredator

release, occupancy analysis, resource facilitation, trophic-mediated behavior, Yellowstone

INTRODUCTION

Intraguild carnivore interactions have crucial bearings on the functional ecology of an ecosystem (Caro & Stoner, 2003; Schuette et al., 2013) and the management of multi-carnivore systems (Brook et al., 2012; Creel & Creel, 1996). Interactions between carnivores range from lethal confrontations and competitive exclusion to facilitation, and can proximately alter the abundance, distribution and behavior of all the species in the guild (Diserens et al., 2022; Palomares & Caro, 1999; Prugh & Sivy, 2020). Such interactions can ultimately affect the structure and function of entire ecosystems through trophic-mediated effects (Brashares et al., 2010; Elmhagen et al., 2010; Elmhagen & Rushton, 2007; Terborgh et al., 2001). Apex carnivores, through their direct (lethal) and indirect (exploitation competition and behavioral suppression) effects, often determine where, when, and how frequently subordinate/mesocarnivores use resources (Gordon et al., 2015; Newsome et al., 2017). For instance, African wild dogs (*Lyacon pictus*) are often excluded from areas of high preferred-prey density by larger, more dominant lions (*Panthera leo*) and spotted hyenas (*Crocuta crocuta*) due to both direct killing and kleptoparasitism (Creel & Creel, 1996; Mills & Gorman, 1997). Similar effects have also been observed in other carnivore systems (see Caro & Stoner, 2003; Curveira-Santos et al., 2021; Dyck et al., 2022; Fedriani et al., 2000).

Consequently, mesocarnivores avoid competition and confrontation through fine-scale spatiotemporal segregation and resource partitioning. Fedriani et al. (1999) found that when lynx (*Lynx lynx*) were more active, red fox (*Vulpes vulpes*) avoided habitats frequented by lynx. In Southern India, while leopards (*P. pardus*) and tigers (*P. tigris*) showed very little temporal partitioning in their activity patterns, leopards hunted in areas with significantly lower vegetation cover than tigers, and both selected different prey types and size-classes (Karanth & Sunquist,

1995). In contrast, fine-scale temporal avoidance of lions by cheetahs (*Acinonyx jubatus*) has

been observed in multiple systems of Tanzania to reduce kleptoparasitism and/or lethal

confrontations (see Durant, 2000; Swanson et al., 2014).

Quantifying the complex fine-scale strategies employed by mesocarnivores to coexist

with competitively dominant carnivores is of particular relevance in the Anthropocene because

humans have disproportionately affected apex carnivores. We have either caused severe

population declines and local extinctions (Ripple et al., 2014), or promoted the recovery of apex

carnivores through reintroduction and conservation efforts (Bangs & Fritts, 1996; Chapron et al.,

2014; Lotze et al., 2011). Such scenarios can have intriguing consequences—in some places

mesocarnivores have been released from competitive suppression (Crooks & Soulé, 1999; Prugh

et al., 2009), while in other areas they face rising competition from recovering apex carnivores

(Berger & Gese, 2007; Jiménez et al., 2019). However, while apex carnivores may limit

mesocarnivores through direct and indirect effects, they can also offer scavenging opportunities

from carcasses of killed prey (Brunet et al., 2022; Sivy et al., 2017; Wilmers et al., 2003). Thus,

the distribution and behavior of mesocarnivores with respect to their apex counterparts are

contingent upon the *risks* (interference competition) and *rewards* (carcass provisioning) of

associations in space and time (Ruprecht et al., 2021).

This risk-reward spectrum for mesocarnivores can be further nuanced by the type of

interactions with apex carnivores and seasonality. For instance, the magnitude of risks for a

mesocarnivore that primarily competes over resources with an apex carnivore is arguably

different than when the mesocarnivore, in addition to resource competition, is also hunted as

prey (Broekhuis, 2015; Carbone et al., 2005). Additionally, in mountainous environments,

ungulate migrations and snow conditions influence the distribution and availability of prey for

both apex and mesocarnivores. In these systems, ungulate birth pulses and changes to body

condition lead to seasonal variation in the average size of apex carnivore kills, with larger

carcasses (and thus opportunities for mesocarnivores to scavenge) more readily available during

winter months (Wilmers et al., 2003). Large carcass availability in winter might shift the

dynamics of spatiotemporal cohabitation within carnivore guilds. Thus, interactions between

apex and mesocarnivores are mediated by a suite of local ecological conditions and inter-specific

interactions.

Yellowstone National Park is the largest intact, temperate ecosystem in the USA that

hosts a diverse community of mammalian carnivores and ungulates. In the Northern Range of

Yellowstone National Park, apex carnivores such as wolves and cougars (*Puma concolor*) are

abundant, particularly in the winter when they are supported by large ungulate prey that

congregate at high densities (Anton, 2020; MacNulty et al., 2020; Smith et al., 2020). Given the

abundance of wolves and cougars and their potential dominating/top-down effects in the system,

we assessed two competing hypotheses with respect to mesocarnivore (coyote and red fox)

distribution and behavior:

1. Mesocarnivores will avoid apex carnivores, both spatially and temporally, to

139   minimize direct (lethal) risks (*risk aversion hypothesis*).

140   2. Mesocarnivores will be spatially and temporally attracted to apex carnivores to

141   leverage scavenging opportunities from apex carnivore kills (*facilitation*

142   *hypothesis*).

These hypotheses might not be mutually exclusive, instead mesocarnivores may employ a

mix of proactive and reactive strategies at different spatiotemporal scales that can partially align

with both hypotheses (Creel, 2018; Swanson et al., 2014). Furthermore, long-term anecdotes

from Yellowstone suggest that while coyotes and red foxes are killed by both wolves and

cougars, the mechanisms of such lethal actions differ. Cougars appear to *hunt* coyotes as prey,

while wolves kill them *opportunistically* at ungulate-carcass sites. Thus, the magnitude of risks

(versus rewards) for mesocarnivores in this system is expected to differ as a function of the apex

carnivore in question, arguably resulting in separate spatiotemporal and behavioral effects.

Alternatively, mesocarnivores may not be affected by top-down regulation from apex carnivores,

instead their distribution and behavior might be mediated by bottom-up processes such as habitat

structure and other eco-geographical covariates (Cano-Martínez et al., 2024). To understand

these complex relationships, we estimated spatial overlap, occupancy, diel activity patterns, and

fine-scale temporal responses between carnivore detections through winter camera-trapping

surveys. We also monitored predation events of GPS-collared wolves and cougars to quantify

mesocarnivore scavenging at apex carnivore kills, as well as direct killings of mesocarnivores.

MATERIALS AND METHODS

Study Area

We conducted this study in the Northern Range of Yellowstone National Park (Figure 1).

Our study area ranges in elevation between 1500–2400m and includes a mix of rugged and rocky

terrain along with open, rolling hills where the common vegetation types are Douglas-fir

(*Psuedotsuga menziesii*), sage brush (*Artemisia* spp.), and juniper (*Juniperus* spp.). The area

experiences long, cold winters with average snow-water-equivalents ranging from 2–30 cm

(Plumb et al., 2009). Wolves, grizzly bears (*Ursus arctos horribilis)*, black bears (*U.*

americanus), and cougars exist in relatively high numbers as apex carnivores, while coyotes, red

fox, bobcats, and martens (*Martes americana*) form the prominent mesocarnivore strata.

Camera Trap Surveys

Camera surveys were done in a 580-km^2 area delineated by a 5-km buffer around our

camera-trap grid that was originally developed to estimate cougar densities. We deployed camera

stations in a "checkerboard" pattern (to increase variability in the spacing between stations; Sun

et al., 2014) with a grid of 3.2-km^2 cells (Anton, 2020). Within each selected grid cell, we

deployed two camera stations at locations chosen to maximize detections, with a minimum

distance of 1 km between stations.

Camera stations were active for 15-week winter sampling periods (i.e., sessions) between

December and March of 2020-21, 2021-22, and 2022-23 (hereafter 2021, 2022, and 2023,

respectively). We used Browning Spec Ops trail cameras with motion-activated passive infrared

detectors. During 2021, we deployed 90 cameras across 60 stations, such that half of the stations

had one camera while the remaining stations had two opposite-facing cameras. In 2022 and

2023, two cameras were deployed at each station for a total of 120 cameras. Each camera was set

for long-range motion detection and recorded 20-second videos with one-minute delays between

each recording. We manually examined all videos, classified species, and extracted meta-data

using the *timelapse2* image analyzing software (Greenberg et al., 2019).

Data Analysis

Spatial Overlap

We assessed the spatial overlap between the four carnivore species (wolf, cougar, coyote,

and red fox) using Pianka's niche overlap index (Pianka, 1973). We calculated the mean relative

abundance index (RAI), also known as the photo-capture rate (Carbone et al., 2001) across all

three sessions for each species at each camera station. Detections of the same species within 30

minutes of one another were omitted to minimize duplicate records of the same individual

(Tanwar et al., 2021). Camera stations were treated as spatially independent to calculate RAI.

Pianka's index measures overlap in space use for each species pair and ranges from 0–1 (Justa &

Lyngdoh, 2023; Pianka, 1973). We also ran 100,000 bootstrap samples to generate 95%

confidence intervals (CI) around the mean.

Multi-Species Occupancy Models

To evaluate how habitat covariates and spatial interactions with other carnivores

influenced the occurrence of apex and mesocarnivores, we constructed multi-species occupancy

models (Rota et al., 2016) using the *unmarked R* package (Fiske & Chandler, 2011; R Core

Team, 2023). Occupancy models incorporate binary presence-absence (1/0) data through

likelihood-based approaches to estimate species occurrence at site i in a latent z (occupancy)

"state" model, while accounting for imperfect detections through a separate "detection" model.

Multi-species occupancy models are an extension of the single-species occupancy model

(MacKenzie, 2006) that apply a hierarchical modeling framework to estimate the effect of

covariates (first-order) and species co-occurrence (second-order) on occupancy.

We used 1-km^2 grid cells for modeling occupancy, with each cell containing no more

than one camera station. Because wolves, cougars, coyotes, and potentially red foxes typically

occupy ranges larger than 1 km^2, the close spacing of our cameras may violate assumptions of

spatial independence. Hence, our occupancy estimates more aptly describe patterns of "space

use" rather than distribution (Steenweg et al., 2018). For estimating detection probabilities of

each species, we considered three covariates. First, the number of active cameras at a station

(ranging from 0–2 as a product of how many cameras were active and for how long during week-

long occasions). As the number of active cameras increased, we expected an increase in

detection probability. Second, we used ratios of topographic roughness and percent tree cover at

camera stations by dividing fine-scale (30 m) values by the average values from a 1-km^2 buffer

around the station for each covariate. For estimating species occupancy, we included three

landscape covariates at a 1-km spatial resolution: topographic roughness, percent tree cover, and

average snow depth. These habitat covariates were included because they are known to affect the

occupancy, movement, and detection of carnivores. Please refer to Table S1 for covariate details.

We used a multi-stage build-up strategy to select the top multi-species occupancy models

(Twining et al., 2024) for each of our three sessions. Models only included univariate effects on

detection or occupancy, unless multiple covariates had significant effects ($p < 0.05$), at which

point we added additional significant covariates (no more than two). We began by constructing

single-species models to evaluate which covariates best explained detection probabilities while

keeping occupancy as constant ($\Psi(.)$). The top detection models were selected for each species

based on the significance of beta estimates and Akaike Information Criterion (AIC), after which

we included covariates on occupancy to find the top overall first-order model by again using

AIC. After selecting the top first-order model, we added second-order species interactions to

develop multi-species models that tested the effects of species co-occurrence on occupancy.

Since we were interested in ecological co-occurrence, we elected to report results from the

second-order multi-species occupancy models even if they performed worse than the top single-

species model. Please refer to Table S3 and S4 for a list and comparison of candidate models.

Temporal Activity Patterns

To maintain independence of detections, photo-captures of the same species within 30

minutes were omitted (Tanwar et al., 2021). We used the *activity* package in R to estimate diel

activity patterns for wolves, cougars, coyotes, and red foxes using non-parametric kernel density

estimation (Rowcliffe, 2023). To quantify temporal niche partitioning between species and

identify potential coexistence strategies that extend beyond broad-scale occupancy, we used the

overlap package in R to assess overlap of diel activity patterns between each species-pair (e.g.,

wolf-coyote, coyote-red fox) by calculating the area under the combined activity curves, $\widehat{\Delta_4}$

(Ridout & Linkie, 2009). We generated 10,000 bootstrap samples of the activity overlap

coefficient, $\widehat{\Delta_4}$, for each species-pair to estimate 95% confidence intervals of overlap.

Time-to-event

We examined fine-scale mesocarnivore activity in response to apex carnivore movements

using time-to-event analysis (Cusack et al., 2017; Davis et al., 2021). The elapsed time between a

mesocarnivore detection following an apex carnivore detection was recorded at each camera

station for a given session. Subsequently, we combined data across all sessions for each apex and

mesocarnivore pairing. We truncated apex carnivore detections to only include events that were

followed by a mesocarnivore detection (i.e., apex carnivore detections followed by another apex

carnivore detection were omitted). Detections of mesocarnivores were recorded for 5 days

following an apex carnivore detection, with elapsed time differences binned into 24-hour

periods. Thereafter, we estimated detection probabilities by dividing the number of

mesocarnivore detections in each 24-hour period by the total number of detections across all

sessions (Cusack et al., 2017; Davis et al., 2021). These detection probabilities were compared to

random iterations (n = 1,000) of mesocarnivore detections to derive expected detection

probabilities for each 24-hour period by using standard two-tailed permutation tests (Davis et al.,

2021). If the risk aversion hypothesis was true, we would expect coyote and red fox detection

probabilities to be significantly lower following an apex carnivore detection. The converse

would be in effect if the facilitation hypothesis was true.

Monitoring of Apex Carnivore Predation Events

To assess mesocarnivore scavenging at apex carnivore kills and direct killing of

mesocarnivores, we monitored predation by GPS-collared wolves (n = 26 from 8 packs) and

cougars (n = 17) during 30-day winter sampling periods (November to March) from 2016–2023.

Monitored wolves and cougars were fitted with Vectronics or Telonics GPS-Satellite collars that

were programmed to collect hourly fixes. We searched GPS-location clusters (Anderson &

Lindzey, 2003) to identify kill sites and conducted necropsies on prey remains. At each kill, we

ascertained the likely cause of death, the species, sex, age of prey when possible, and the

presence of scavengers through visual observations and sign (e.g., scat, tracks, hair).

RESULTS

Spatial Overlap

Mean spatial overlap between apex and mesocarnivores was 0.57 ± 0.02 for coyote-wolf,

0.24 ± 0.005 for coyote-cougar, 0.67 ± 0.01 for red fox-wolf, and 0.53 ± 0.01 for red fox-cougar

across the 3 sessions (Figure 2, Table S2). Overlap between mesocarnivores (red fox-coyote)

was relatively high across all sessions (0.54 ± 0.01), while overlap between apex carnivores

(wolf-cougar) was comparatively lower (0.33 ± 0.02).

Multi-Species Occupancy Models

Mean naive occupancy across all three sessions was 0.89 for red fox, 0.59 for coyote,

0.59 for wolf, and 0.52 for cougar (Figure 3a). In 2021, the best-performing model had poor

convergence with inflated standard errors, possibly due to high naive occupancy estimates

(Figure 3a). We re-fitted the top model by imposing an optimized penalty (0.5) on the calculated

likelihood through K-fold cross-validation. We acknowledge that adding a likelihood penalty

introduces bias to parameter estimates. Therefore, results from the 2021 model should be

interpreted with caution. Given this, cover ratio best explained variation in both red fox (β = -

0.19, $P = 0.02$) and cougar detection probability in 2021 ($\beta = $ -0.16, $P = 0.4$). The effect of cover

ratio on cougar detections became insignificant when moving from the detection-only model to

the full, penalized model. No predictors explained significant variation in coyote or wolf

detections. Topographic roughness had a significant positive effect on cougar occupancy

probability ($\beta = 1.71$, $P = 0.001$), but there were no covariates that influenced red fox, coyote, or

wolf occupancy.

In 2022 (Table S3 and S4, Figure S1), variation in red fox detection probability was best

explained by roughness ratio, with red fox detection increasing as roughness ratio increased ($\beta = $

-0.25, $P = 0.09$). Cover ratio had a significant, positive effect on coyote detection probability (β

$= 0.14$, $P = 0.01$) and a negative effect on cougar detection probability ($\beta = $ -0.11, $P = 0.37$).

Roughness ratio for red fox and cover ratio for cougars had significant effects in the detection-

only model. However, these effects on detection became insignificant once occupancy covariates

were included in the model. There was no covariate that explained variation in the probability of

wolf detection better than the null model. Habitat covariates only helped explain variation in

cougar occupancy, while the null model performed best for red fox, coyote, and wolf. Cougar

occupancy probability increased as mean snow depth decreased, ($\beta = $ -0.94, $P = 0.05$) and

roughness increased ($\beta = 1.16$, $P = 0.04$) (Figure S2).

In 2023 (Tables S3 and S4, Figure S1), active cameras were positively-associated ($\beta = $

0.63, $P = 0.005$) and cover ratio was negatively-associated ($\beta = $ -0.1, $P = 0.03$) with red fox

detection probability. As cover ratio increased, coyote detection probability also increased ($\beta = $

0.16, $P = 0.007$). Roughness ratio had a positive effect on wolf detection probability ($\beta = $ -0.78,

$P = 0.002$), while active cameras were positively-associated ($\beta = 2.57$, $P = 0.04$) and cover ratio

negatively-associated ($\beta = $ -0.34, $P = 0.006$) with cougar detection probability. Once again,

covariates only helped explain variation in cougar occupancy, with occupancy probability

increasing as roughness increased ($\beta = 1.6$, $P = 0.007$) (Figure S2) and cover decreased ($\beta = -$

1.1, $P = 0.06$).

After incorporating second-order species interactions on occupancy, we found that the

majority of species occurred independent of each other (Table S5). However, coyote occupancy

was conditional on wolf occupancy, with coyote occupancy being significantly higher where

wolves were present in 2021 ($\beta = 1.35$, $P = 0.006$) and 2022 ($\beta = 2.3$, $P = 0.03$), and marginally

significant in 2023 ($\beta = 1.23$, $P = 0.07$) (Figure 3b). Possibly due to bias from the penalized

model in 2021, we also found positive associations between red fox and coyote ($\beta = 1.47$, $P <$

0.0001), red fox and wolf ($\beta = 1.14$, $P = 0.001$), and red fox and cougar ($\beta = 1.2$, $P = 0.003$), and

a near significant negative association between coyote and cougar ($\beta = -1.0$, $P = 0.07$).

Temporal Activity Patterns

Cougars (n = 341 detections) exhibited unimodal diel activity, being most active around

dusk peaking at 1800 and least active from morning (0800) to midday (1400) (Figure 4). Wolves

(n = 443 detections) had bimodal, crepuscular activity patterns, with peak activity shortly after

dawn (~0800–0900) and a secondary peak at dusk (1700–1800) (Figure 4). Wolves were least

active during midday (~1100–1500). Coyotes (n = 541 detections) had more consistent activity

across the diel cycle and were more diurnal, with peak activity occurring from late morning

(0900) to midday (1400) and lowest activity before dawn (0500) and after dusk (2100) (Figure

4). Red foxes (n = 1520 detections), on the other hand, exhibited highly nocturnal patterns, being

most active shortly after dusk (1800–2100) and sustaining high activity until before dawn (2100–

0600) before plummeting to little or no activity during daylight hours (0800–1600) (Figure 4).

Temporal overlap between apex and mesocarnivores was $\widehat{\Delta_4} = 0.86$ (CI: $0.81 – 0.89$) for wolf-

coyote, 0.68 (CI: 0.61 – 0.7) for wolf-red fox, 0.81 (CI: 0.74 – 0.85) for cougar-coyote, and 0.72

(CI: 0.66 – 0.76) for cougar-red fox across the three sessions (Figure 4a). Overlap between

mesocarnivores (red fox-coyote) was $\widehat{\Delta_4} = 0.61$ (CI: 0.53 – 0.62), while overlap between apex

carnivores (wolf-cougar) was $\widehat{\Delta_4} = 0.86$ (CI: 0.8 – 0.91) (Figure 4b).

Time-to-event

Compared to the mean values derived from random iterations of detection probabilities,

red foxes and coyotes were both over twice as likely to be detected in the 24-hour period

following a wolf detection (red foxes: 0.034 compared to 0.015, $P = 0.001$; coyotes: 0.05

compared to 0.018; $P = 0.001$; Figure 5). However, their detection probabilities following cougar

detections differed. While red foxes were over twice as likely to be detected in the 24-hour

period following a cougar detection (0.024 compared to 0.011; $P = 0.001$), coyote detection rates

were not affected. Additionally, red foxes were more likely to be detected in the 72-hour period

following a wolf detection (0.01 compared to 0.007; $P = 0.04$) and the 96-hour period following

a cougar detection (0.009 compared to 0.004; $P = 0.001$). Overall, the time interval between

detections of a mesocarnivore subsequent to an apex carnivore detection was shortest for wolf-

coyote (median = 17.67 hours) and longest for cougar-coyote (median = 57.35 hours). Foxes

showed similar latency to wolf and cougar detections, with medians of 21.11 and 24.69 hours,

respectively (Figure 5 b).

Monitoring of Apex Carnivore Predation Events

We detected 327 wolf kills and 257 cougar kills during winter sampling periods. Coyote

presence was recorded at 222 (68%) wolf kills and 80 (31%) cougar kills (Figure 6a). Red fox

presence was recorded at 67 (20%) wolf kills and 46 (18%) cougar kills (Figure 6a). From this

predation monitoring, we documented 18 coyotes and one red fox killed by wolves (Figure 6b).

Of the coyotes killed by wolves, 11 (61%) occurred at wolf feeding sites (i.e., ungulates killed or

scavenged by wolves), while the red fox mortality did not occur at a wolf-killed prey site.

Cougars killed eight coyotes and three red foxes during this time (Figure 6b). None of these

mortalities were associated with cougar-killed prey sites, rather they were fully consumed and

generally found in cache piles - suggestive of cougar consumption.

DISCUSSION

Red foxes and coyotes, mesocarnivores that frequently scavenge from wolf and cougar

kills, were ubiquitous in our study area (Figure 1). Aside from coyote and wolf overlap with

cougars, we found a high degree of spatial overlap between mesocarnivores, as well as apex-

mesocarnivore pairs (Figure 2). Our multi-species occupancy models revealed no significant

effect of habitat covariates on mesocarnivore occupancy, an unsurprising result given their

relatively high naïve occupancy estimates (Figure 3). Our analysis suggests a positive

relationship between wolf and coyote occupancy for all years, as well as evidence of negative

impacts of cougar occupancy on coyote space-use (Table S4). Notably, coyotes and cougars also

had the lowest spatial overlap of all species pairs (Figure 2), indicating a likely avoidance of

cougars by coyotes in our system. However, potential pitfalls of this analysis should be

considered.

Our camera-trapping study was designed to estimate cougar density rather than apex and

mesocarnivore occupancy, leading to camera spacing closer than the average home range size of

each species, with the potential exception of red fox (Main et al., 2020). While similar camera

spacings have been used to assess occupancy of apex and mesocarnivores (Justa & Lyngdoh,

2023), this may violate assumptions of spatial independence, and hence our results are more

indicative of carnivore space-use patterns as opposed to occupancy (Steenweg et al., 2018).

Further, competition between species is often more nuanced than the binary "occupied" versus

"unoccupied." Instead, species may coexist in broad space by mitigating finer-scale overlap in

space and time. The fact that each apex and mesocarnivore species was detected at the majority

of sites during the course of our study (Figure 1) supports this notion and further demonstrates

the limitations of using occupancy analysis to study species interactions in saturated systems.

These analytical constraints, however, do not limit the scope of our conclusions which are drawn

at a very high resolution of space use for most of the species. In fact, they provide stronger

support for the saturation of this system. As such, fine-scale temporal approaches were required

to disentangle these complex intra-guild relationships.

Despite having considerable temporal overlap with wolves and cougars, diel activity plots

indicated that coyotes were most active mid-day when apex carnivores were collectively the least

active in our study area (Figure 4). This peak in activity differs from those of coyotes inhabiting

adjacent areas of the Greater Yellowstone Ecosystem with lower apex carnivore densities, where

coyotes were found to be more active in the early morning and at night (Smith et al., 2023). In

contrast, red foxes appeared less sensitive to the diel activity of apex carnivores. Not only did

their peak in activity closely align with that of cougars, but they also displayed the same

nocturnal patterns as red foxes across the Greater Yellowstone Ecosystem (Smith et al., 2023).

However, red foxes exhibited high temporal segregation from coyotes which was evident in

having the lowest temporal overlap of any species pair (Figure 4). Studies elsewhere have shown

that coyotes and red foxes compete severely, with coyotes competitively excluding and/or killing

red foxes (Fedriani et al., 2000; Levi & Wilmers, 2012). Thus, the nocturnal activity of red foxes

might also be to avoid coyote peak activity during the day that is in turn modulated by wolf and

cougar activity (Figure 4). Temporal sympatry amongst the carnivore guild appears to be layered

with apex carnivores affecting coyotes, and coyotes affecting red foxes.

Our findings add to a growing body of work that demonstrate how inferences concerning

apex and mesocarnivore interactions depend on the spatiotemporal scale at which analyses are

conducted. For example, Prugh et al. (2023) found that coyotes avoided areas with greater wolf

utilization when using step-selection analyses. Mesocarnivores may indeed avoid possible

encounters with apex carnivores when moving throughout their landscape, yet scavenging

opportunities from apex carnivore kills likely drive temporary attraction. In our study, both

coyote and red fox detection probabilities at camera stations doubled in the 24 hours following

wolf detections (Figure 5), and this pattern also held true for red fox detections following

cougars. In conjunction with frequent mesocarnivore scavenging signs recorded at wolf and

cougar kill sites (Figure 6a), these results suggest that the benefits from such scavenging

opportunities perhaps facilitate broad mesocarnivore space-use despite high densities of apex

carnivores in our study area.

It is, however, noteworthy that coyote detection probabilities were not affected by cougar

detections. This possible lack of attraction is corroborated by our occupancy and temporal

activity results that collectively indicate coyotes exhibit some proactive avoidance of cougars.

Additionally, our predation monitoring provides evidence of cougars killing and consuming

coyotes, a key distinction from wolves that typically kill, but do not consume coyotes attempting

to scavenge from their kills (Figure 6b). Based on our results and relevant earlier research from

the study system (Ruth et al., 2019), we propose that coyote avoidance of cougars could be a

result of coyotes serving as prey for cougars in addition to being a competitor, thereby facing

higher risks. This is unlike coyote-wolf interactions, where coyotes are only a competitor of

wolves. Furthermore, the ambush hunting style of cougars, combined with the forested and

topographically rough terrain they inhabit, could lead to coyotes being more vulnerable in those

habitats and/or at cougar kill sites (Perrig et al., 2023). In contrast, the open and flatter terrain

preferred by wolves could provide more visual and olfactory cues for coyotes to assess (and

escape) immediate risks of lethal encounters. Such disparate risk-effects might have also

manifested in coyote-scavenging to be over twice as prevalent at wolf kills than cougar kills in

our study area (Figure 6a).

Red fox detection probabilities were affected by both wolf and cougar detections,

suggesting attraction to apex carnivore movements. Coupled with their highly nocturnal activity

that is offset from peak coyote activity, these patterns suggest foxes may be operating along a

risk-reward continuum to seek scavenging opportunities from apex carnivore kills while

minimizing exposure to competitively-dominant coyotes. While relatively few fox mortalities

were detected through monitoring apex carnivore predation, we speculate similar relationships as

with coyotes; foxes are more likely to be preyed upon and consumed by cougars, whereas risk

from wolves is based on resource competition. Yet, foxes exhibited more attraction to cougars

compared to coyotes, indicating that they may be more adept at avoiding immediate risks of

cougar predation. Owing to their relatively large foot:body ratio, foxes are better at maneuvering

on top of snow (Van Etten et al., 2007), possibly explaining their increased ability to escape

cougar predation.

The availability of alternative resources may also influence these relationships. The

limitations of prey acquisition during winter months may increase mesocarnivore reliance on

scavenging opportunities (Jensen et al., 2022; Smith et al., 2023). However, lower detections of

fox sign at wolf and cougar kills in winter compared to coyotes suggests foxes may be able to

meet energetic demands from alternative food sources, presumably due to their smaller body

size. For example, compared to coyotes and wolves, foxes in the study area are more commonly

observed hunting subnivean rodents by pouncing through the snowpack during the winter (D.

Stahler *pers.comm.*). Therefore, the facilitative relationship between apex and mesocarnivores

through carcass scavenging might not be even between coyotes and foxes, nor as strong for both

in the summer/spring when rodent availability increases, as well as ungulate neonates,

vegetation, and insects (Jensen et al., 2022). Also, the presence of grizzly and black bears that

emerge in the summer after winter-long hibernations can complicate these relationships because

bears often limit carrion consumption for other scavengers (Allen et al., 2014). Collectively,

these factors demonstrate how multi-carnivore interactions often depend on both the availability

and necessity of scavenging opportunities.

Notably, while our study highlights the positive aspects of apex and mesocarnivore

interactions, we did not explore variations in red fox and coyote population trends. In fact,

previous research documented a 39% decline in coyote abundance following wolf reintroduction

in the Lamar Valley of Yellowstone, an area adjacent to the study area (Berger & Gese, 2007).

Additionally, Merkle et al. (2009) showed a decrease in wolf-coyote interactions over time

(between 1997 and 2007) in our study area, indicative of changes to coyote density and/or

behavior (e.g., wolf avoidance). In NE Oregon, coyotes balanced high scavenging rates from

cougar's kills with high predation rates, as cougars were responsible for 23% of coyote

mortalities (Ruprecht et al., 2021). Despite benefiting from carrion made available by wolves and

cougars, red fox and coyote populations may experience top-down regulation via direct killing

that maintain their populations at low densities (Sinclair, 2003). Long-term studies that estimate

population trends of mesocarnivores with respect to apex carnivore densities are required to test

such effects.

The magnitude and impact of trophic-mediated responses related to large carnivore

recoveries have become a contentious issue, perhaps nowhere more so than in Yellowstone

National Park (Brice et al., 2022; Hobbs et al., 2024; Ripple & Beschta, 2012). The

reintroduction of wolves and recolonization of cougars to this area likely changed mesocarnivore

distributions and abundance, yet the resulting populations of red foxes and coyotes may have

also benefited from living with their dominant counterparts. Overall, red foxes and coyotes in our

study area exhibited more attraction than avoidance to apex carnivores during winter months,

when small prey and alternative food resources are scant. Such attraction, possibly a

consequence of desperate foraging decisions in winter, can also prove fatal. However, risks to

mesocarnivores are uneven across dominant-subordinate carnivore pairings, with apex carnivore

behavior adding nuance to the *landscape of fear* for mesocarnivores, a novel insight from our

study. Mesocarnivores appear to walk a thin line between the risks and rewards of associating

with their dominant counterparts, which in turn is governed by the spatiotemporal dispersion of

resources and inter-carnivore relationships. Our results showcase the complexity of interspecific

relationships within a rich carnivore guild, and highlight key trophic links that can promote

community resilience in the wake of rapidly changing ecological conditions in the

Anthropocene.

REFERENCES

Allen, M. L., Elbroch, L. M., Wilmers, C. C., & Wittmer, H. U. (2014). Trophic Facilitation or
Limitation? Comparative Effects of Pumas and Black Bears on the Scavenger
Community. *PLoS ONE*, *9*(7), e102257. https://doi.org/10.1371/journal.pone.0102257

Anton, C. B. (2020). *The demography and comparative ethology of top predators in a multi-
carnivore system*. University of California-Santa Cruz.

Bangs, E. E., & Fritts, S. H. (1996). Reintroducing the Gray Wolf to Central Idaho and
Yellowstone National Park. *Wildlife Society Bulletin*, *24*(3), 402–413.

Berger, K. M., & Gese, E. M. (2007). Does interference competition with wolves limit the
distribution and abundance of coyotes? *Journal of Animal Ecology*, *76*(6), 1075–1085.
https://doi.org/10.1111/j.1365-2656.2007.01287.x

Brashares, J., Prugh, L. R., Stoner, C. J., & Epps, C. (2010). Ecological and conservation
implications of mesopredator release. In *Trophic cascades: Predators, prey, and the
changing dynamics of nature*.

Brice, E. M., Larsen, E. J., & MacNulty, D. R. (2022). Sampling bias exaggerates a textbook
example of a trophic cascade. *Ecology Letters*, *25*(1), 177–188.
https://doi.org/10.1111/ele.13915

Broekhuis, F. (2015). Cat eats cat: Leopard consumes adult cheetah, Maasai Mara Game
Reserve, Kenya. *CAT News*, *65*, 33–34.

Brook, L. A., Johnson, C. N., & Ritchie, E. G. (2012). Effects of predator control on behaviour
of an apex predator and indirect consequences for mesopredator suppression. *Journal of
Applied Ecology*, *49*(6), 1278–1286. https://doi.org/10.1111/j.1365-2664.2012.02207.x

Brunet, M. J., Monteith, K. L., Huggler, K. S., Clapp, J. G., Thompson, D. J., Burke, P. W.,

Zornes, M., Lionberger, P., Valdez, M., & Holbrook, J. D. (2022). Cats and dogs: A mesopredator navigating risk and reward provisioned by an apex predator. *Ecology and Evolution*, *12*(2), e8641. https://doi.org/10.1002/ece3.8641

Cano-Martínez, R., Thorsen, N. H., Hofmeester, T. R., Odden, J., Linnell, J., Devineau, O., Angoh, S. Y. J., & Odden, M. (2024). Bottom-up rather than top-down mechanisms determine mesocarnivore interactions in Norway. *Ecology and Evolution*, *14*(3), e11064. https://doi.org/10.1002/ece3.11064

Carbone, C., Christie, S., Conforti, K., Coulson, T., Franklin, N., Ginsberg, J. R., Griffiths, M., Holden, J., Kawanishi, K., Kinnaird, M., Laidlaw, R., Lynam, A., Macdonald, D. W., Martyr, D., McDougal, C., Nath, L., O'Brien, T., Seidensticker, J., Smith, D. J. L., … Wan Shahruddin, W. N. (2001). The use of photographic rates to estimate densities of tigers and other cryptic mammals. *Animal Conservation*, *4*(1), 75–79. https://doi.org/10.1017/S1367943001001081

Carbone, C., Frame, L., Frame, G., Malcolm, J., Fanshawe, J., FitzGibbon, C., Schaller, G., Gordon, I. J., Rowcliffe, J. M., & Toit, J. T. (2005). Feeding success of African wild dogs (*Lycaon pictus*) in the Serengeti: The effects of group size and kleptoparasitism. *Journal of Zoology*, *266*(2), 153–161. https://doi.org/10.1017/S0952836905006710

Caro, T. M., & Stoner, C. J. (2003). The potential for interspecific competition among African carnivores. *Biological Conservation*, *110*(1), 67–75. https://doi.org/10.1016/S0006-3207(02)00177-5

Chapron, G., Kaczensky, P., Linnell, J. D. C., Von Arx, M., Huber, D., Andrén, H., López-Bao, J. V., Adamec, M., Álvares, F., Anders, O., Balčiauskas, L., Balys, V., Bedő, P., Bego, F., Blanco, J. C., Breitenmoser, U., Brøseth, H., Bufka, L., Bunikyte, R., … Boitani, L.

(2014). Recovery of large carnivores in Europe's modern human-dominated landscapes.

Science, *346*(6216), 1517–1519. https://doi.org/10.1126/science.1257553

Creel, S. (2018). The control of risk hypothesis: Reactive vs. proactive antipredator responses

and stress-mediated vs. food-mediated costs of response. *Ecology Letters*, *21*(7), 947–

956. https://doi.org/10.1111/ele.12975

Creel, S., & Creel, N. M. (1996). Limitation of African Wild Dogs by Competition with Larger

Carnivores. *Conservation Biology*, *10*(2), 526–538. https://doi.org/10.1046/j.1523-

1739.1996.10020526.x

Crooks, K. R., & Soulé, M. E. (1999). Mesopredator release and avifaunal extinctions in a

fragmented system. *Nature*, *400*(6744), 563–566. https://doi.org/10.1038/23028

Curveira-Santos, G., Sutherland, C., Tenan, S., Fernández-Chacón, A., Mann, G. K. H., Pitman,

R. T., & Swanepoel, L. H. (2021). Mesocarnivore community structuring in the presence

of Africa's apex predator. *Proceedings of the Royal Society B: Biological Sciences*,

288(1946), 20202379. https://doi.org/10.1098/rspb.2020.2379

Cusack, J. J., Dickman, A. J., Kalyahe, M., Rowcliffe, J. M., Carbone, C., MacDonald, D. W., &

Coulson, T. (2017). Revealing kleptoparasitic and predatory tendencies in an African

mammal community using camera traps: A comparison of spatiotemporal approaches.

Oikos, *126*(6), 812–822. https://doi.org/10.1111/oik.03403

Davis, R. S., Yarnell, R. W., Gentle, L. K., Uzal, A., Mgoola, W. O., & Stone, E. L. (2021). Prey

availability and intraguild competition regulate the spatiotemporal dynamics of a

modified large carnivore guild. *Ecology and Evolution*, *11*(12), 7890–7904.

https://doi.org/10.1002/ece3.7620

Diserens, T. A., Churski, M., Bubnicki, J. W., Zalewski, A., Brzeziński, M., & Kuijper, D. P. J.

(2022). Wolf risk fails to inspire fear in two mesocarnivores suggesting facilitation

prevails. *Scientific Reports*, *12*(1), 16469. https://doi.org/10.1038/s41598-022-20725-3

Durant, S. M. (2000). Predator avoidance, breeding experience and reproductive success in

endangered cheetahs, Acinonyx jubatus. *Animal Behaviour*, *60*(1), 121–130.

https://doi.org/10.1006/anbe.2000.1433

Dyck, M. A., Wyza, E., & Popescu, V. D. (2022). When carnivores collide: A review of studies

exploring the competitive interactions between bobcats *Lynx rufus* and coyotes *Canis*

latrans. *Mammal Review*, *52*(1), 52–66. https://doi.org/10.1111/mam.12260

Elmhagen, B., Ludwig, G., Rushton, S. P., Helle, P., & Lindén, H. (2010). Top predators,

mesopredators and their prey: Interference ecosystems along bioclimatic productivity

gradients. *Journal of Animal Ecology*, *79*(4), 785–794. https://doi.org/10.1111/j.1365-

2656.2010.01678.x

Elmhagen, B., & Rushton, S. P. (2007). Trophic control of mesopredators in terrestrial

ecosystems: Top-down or bottom-up? *Ecology Letters*, *10*(3), 197–206.

https://doi.org/10.1111/j.1461-0248.2006.01010.x

Fedriani, J. M., Fuller, T. K., Sauvajot, R. M., & York, E. C. (2000). Competition and intraguild

predation among three sympatric carnivores. *Oecologia*, *125*(2), 258–270.

https://doi.org/10.1007/s004420000448

Fedriani, J. M., Palomares, F., & Delibes, M. (1999). Niche relations among three sympatric

Mediterranean carnivores. *Oecologia*, *121*(1), 138–148.

https://doi.org/10.1007/s004420050915

Fiske, I., & Chandler, R. (2011). **unmarked**: An *R* Package for Fitting Hierarchical Models of

Wildlife Occurrence and Abundance. *Journal of Statistical Software*, *43*(10).

https://doi.org/10.18637/jss.v043.i10

Gordon, C. E., Feit, A., Grüber, J., & Letnic, M. (2015). Mesopredator suppression by an apex

predator alleviates the risk of predation perceived by small prey. *Proceedings of the*

Royal Society B: Biological Sciences, *282*(1802), 20142870.

https://doi.org/10.1098/rspb.2014.2870

Greenberg, S., Godin, T., & Whittington, J. (2019). Design patterns for wildlife-related camera

trap image analysis. *Ecology and Evolution*, *9*(24), 13706–13730.

https://doi.org/10.1002/ece3.5767

Hobbs, N. T., Johnston, D. B., Marshall, K. N., Wolf, E. C., & Cooper, D. J. (2024). Does

restoring apex predators to food webs restore ecosystems? Large carnivores in Y

ellowstone as a model system. *Ecological Monographs*, *94*(2), e1598.

https://doi.org/10.1002/ecm.1598

Jensen, A. J., Marneweck, C. J., Kilgo, J. C., & Jachowski, D. S. (2022). Coyote diet in North

America: Geographic and ecological patterns during range expansion. *Mammal Review*,

52(4), 480–496. https://doi.org/10.1111/mam.12299

Jiménez, J., Nuñez-Arjona, J. C., Mougeot, F., Ferreras, P., González, L. M., García-Domínguez,

F., Muñoz-Igualada, J., Palacios, M. J., Pla, S., Rueda, C., Villaespesa, F., Nájera, F.,

Palomares, F., & López-Bao, J. V. (2019). Restoring apex predators can reduce

mesopredator abundances. *Biological Conservation*, *238*, 108234.

https://doi.org/10.1016/j.biocon.2019.108234

Justa, P., & Lyngdoh, S. (2023). Understanding carnivore interactions in a cold arid trans-

Himalayan landscape: What drives co-existence patterns within predator guild along

varying resource gradients? *Ecology and Evolution*, *13*(5), e10040.

https://doi.org/10.1002/ece3.10040

Karanth, K. U., & Sunquist, M. E. (1995). Prey Selection by Tiger, Leopard and Dhole in

Tropical Forests. *The Journal of Animal Ecology*, *64*(4), 439.

https://doi.org/10.2307/5647

Levi, T., & Wilmers, C. C. (2012). Wolves–coyotes–foxes: A cascade among carnivores.

Ecology, *93*(4), 921–929. https://doi.org/10.1890/11-0165.1

Lotze, H. K., Coll, M., Magera, A. M., Ward-Paige, C., & Airoldi, L. (2011). Recovery of

marine animal populations and ecosystems. *Trends in Ecology & Evolution*, *26*(11), 595–

605. https://doi.org/10.1016/j.tree.2011.07.008

MacKenzie, D. I. (Ed.). (2006). *Occupancy estimation and modeling: Inferring patterns and*

dynamics of species. Elsevier.

MacNulty, D. R., Stahler, D. R., Wyman, C. T., Ruprecht, J., Smith, L. M., Kohl, M. T., &

Smith, D. W. (2020). Population Dynamics of Northern Yellowstone Elk after Wolf

Reintroduction. In *D.W. Smith, D.R. Stahler, and D.R. MacNulty, editors. Yellowstone*

wolves: Science and discovery in the world's first national park (pp. 184–199).

University of Chicago Press.

Main, M. T., Davis, R. A., Blake, D., Mills, H., & Doherty, T. S. (2020). Human impact

overrides bioclimatic drivers of red fox home range size globally. *Diversity and*

Distributions, *26*(9), 1083–1092. https://doi.org/10.1111/ddi.13115

Merkle, J. A., Stahler, D. R., & Smith, D. W. (2009). Interference competition between gray

wolves and coyotes in Yellowstone National Park. *Canadian Journal of Zoology*, *87*(1),

56–63. https://doi.org/10.1139/Z08-136

Mills, M. G. L., & Gorman, M. L. (1997). Factors Affecting the Density and Distribution of

Wild Dogs in the Kruger National Park. *Conservation Biology*, *11*(6), 1397–1406.

https://doi.org/10.1046/j.1523-1739.1997.96252.x

Newsome, T. M., Greenville, A. C., Ćirović, D., Dickman, C. R., Johnson, C. N., Krofel, M.,

Letnic, M., Ripple, W. J., Ritchie, E. G., Stoyanov, S., & Wirsing, A. J. (2017). Top

predators constrain mesopredator distributions. *Nature Communications*, *8*(1), 15469.

https://doi.org/10.1038/ncomms15469

Palomares, F., & Caro, T. M. (1999). Interspecific Killing among Mammalian Carnivores. *The*

American Naturalist, *153*(5), 492–508. https://doi.org/10.1086/303189

Perrig, P. L., Lambertucci, S. A., Donadio, E., Smith, J. A., Middleton, A. D., & Pauli, J. N.

(2023). Risk effects cascade up to an obligate scavenger. *Ecology*, *104*(2), e3871.

https://doi.org/10.1002/ecy.3871

Pianka, E. R. (1973). The Structure of Lizard Communities. *Annual Review of Ecology and*

Systematics, *4*(1), 53–74. https://doi.org/10.1146/annurev.es.04.110173.000413

Plumb, G. E., White, P. J., Coughenour, M. B., & Wallen, R. L. (2009). Carrying capacity,

migration, and dispersal in Yellowstone bison. *Biological Conservation*, *142*(11), 2377–

2387. https://doi.org/10.1016/j.biocon.2009.05.019

Prugh, L. R., Cunningham, C. X., Windell, R. M., Kertson, B. N., Ganz, T. R., Walker, S. L., &

Wirsing, A. J. (2023). Fear of large carnivores amplifies human-caused mortality for

mesopredators. *Science*, *380*(6646), 754–758. https://doi.org/10.1126/science.adf2472

Prugh, L. R., & Sivy, K. J. (2020). Enemies with benefits: Integrating positive and negative

interactions among terrestrial carnivores. *Ecology Letters*, *23*(5), 902–918.

https://doi.org/10.1111/ele.13489

Prugh, L. R., Stoner, C. J., Epps, C. W., Bean, W. T., Ripple, W. J., Laliberte, A. S., &

Brashares, J. S. (2009). The Rise of the Mesopredator. *BioScience*, *59*(9), 779–791.

https://doi.org/10.1525/bio.2009.59.9.9

R Core Team. (2023). *R: A Language and Environment for Statistical Computing* [Computer

software]. R Foundation for Statistical Computing. https://www.r-project.org/

Ridout, M. S., & Linkie, M. (2009). Estimating overlap of daily activity patterns from camera

trap data. *Journal of Agricultural, Biological, and Environmental Statistics*, *14*(3), 322–

337. https://doi.org/10.1198/jabes.2009.08038

Ripple, W. J., & Beschta, R. L. (2012). Trophic cascades in Yellowstone: The first 15years after

wolf reintroduction. *Biological Conservation*, *145*(1), 205–213.

https://doi.org/10.1016/j.biocon.2011.11.005

Ripple, W. J., Estes, J. A., Beschta, R. L., Wilmers, C. C., Ritchie, E. G., Hebblewhite, M.,

Berger, J., Elmhagen, B., Letnic, M., Nelson, M. P., Schmitz, O. J., Smith, D. W.,

Wallach, A. D., & Wirsing, A. J. (2014). Status and Ecological Effects of the World's

Largest Carnivores. *Science*, *343*(6167), 1241484.

https://doi.org/10.1126/science.1241484

Rota, C. T., Ferreira, M. A. R., Kays, R. W., Forrester, T. D., Kalies, E. L., McShea, W. J.,

Parsons, A. W., & Millspaugh, J. J. (2016). A multispecies occupancy model for two or

more interacting species. *Methods in Ecology and Evolution*, *7*(10), 1164–1173.

https://doi.org/10.1111/2041-210X.12587

Rowcliffe, M. (2023). *activity: Animal Activity Statistics* (1.3.4) [R].

10.32614/CRAN.package.activity

Ruprecht, J., Eriksson, C. E., Forrester, T. D., Spitz, D. B., Clark, D. A., Wisdom, M. J., Bianco,

M., Rowland, M. M., Smith, J. B., Johnson, B. K., & Levi, T. (2021). Variable strategies

to solve risk–reward tradeoffs in carnivore communities. *Proceedings of the National*

Academy of Sciences, *118*(35), e2101614118. https://doi.org/10.1073/pnas.2101614118

Ruth, T., Buotte, P., & Hornocker, M. (2019). *Yellowstone Cougars: Ecology Before And During*

Wolf Restoration. University Press of Colorado. https://doi.org/10.5876/9781607328292

Schuette, P., Wagner, A. P., Wagner, M. E., & Creel, S. (2013). Occupancy patterns and niche

partitioning within a diverse carnivore community exposed to anthropogenic pressures.

Biological Conservation, *158*, 301–312. https://doi.org/10.1016/j.biocon.2012.08.008

Sinclair, A. R. E. (2003). Mammal population regulation, keystone processes and ecosystem

dynamics. *Philosophical Transactions of the Royal Society of London. Series B:*

Biological Sciences, *358*(1438), 1729–1740. https://doi.org/10.1098/rstb.2003.1359

Sivy, K. J., Pozzanghera, C. B., Grace, J. B., & Prugh, L. R. (2017). Fatal Attraction? Intraguild

Facilitation and Suppression among Predators. *The American Naturalist*, *190*(5), 663–

679. https://doi.org/10.1086/693996

Smith, A. B., Squires, J. R., Bjornlie, N. L., & Holbrook, J. D. (2023). Divergent or convergent:

How do forest carnivores use time in the Greater Yellowstone Ecosystem? *Journal of*

Mammalogy, gyad070. https://doi.org/10.1093/jmammal/gyad070

Smith, D. W., Cassidy, K. A., Stahler, D. R., MacNulty, D. R., Harrison, Q., Balmford, B.,

Stahler, E. E., Brandell, E. E., & Coulson, T. (2020). Population Dynamics and

Demography. In *D.W. Smith, D.R. Stahler, and D.R. MacNulty, editors. Yellowstone*

wolves: Science and discovery in the world's first national park (pp. 77–92). University

of Chicago Press.

Steenweg, R., Hebblewhite, M., Whittington, J., Lukacs, P., & McKelvey, K. (2018). Sampling

scales define occupancy and underlying occupancy–abundance relationships in animals.

Ecology, *99*(1), 172–183. https://doi.org/10.1002/ecy.2054

Sun, C. C., Fuller, A. K., & Royle, J. A. (2014). Trap Configuration and Spacing Influences

Parameter Estimates in Spatial Capture-Recapture Models. *PLOS ONE*, *9*(2), e88025.

https://doi.org/10.1371/journal.pone.0088025

Swanson, A., Caro, T., Davies-Mostert, H., Mills, M. G. L., Macdonald, D. W., Borner, M.,

Masenga, E., & Packer, C. (2014). Cheetahs and wild dogs show contrasting patterns of

suppression by lions. *Journal of Animal Ecology*, *83*(6), 1418–1427.

https://doi.org/10.1111/1365-2656.12231

Tanwar, K. S., Sadhu, A., & Jhala, Y. V. (2021). Camera trap placement for evaluating species

richness, abundance, and activity. *Scientific Reports*, *11*(1), 23050.

https://doi.org/10.1038/s41598-021-02459-w

Terborgh, J., Lopez, L., Nuñez, P., Rao, M., Shahabuddin, G., Orihuela, G., Riveros, M.,

Ascanio, R., Adler, G. H., Lambert, T. D., & Balbas, L. (2001). Ecological Meltdown in

Predator-Free Forest Fragments. *Science*, *294*(5548), 1923–1926.

https://doi.org/10.1126/science.1064397

Twining, J. P., Brazeal, J. L., Jensen, P. G., & Fuller, A. K. (2024). Intraguild interactions and

abiotic conditions mediate occupancy of mammalian carnivores: Co-occurrence of

coyotes–fishers–martens. *Oikos*, *2024*(6), e10577. https://doi.org/10.1111/oik.10577

Van Etten, K. W., Wilson, K. R., & Crabtree, R. L. (2007). Habitat Use of Red Foxes in

Yellowstone National Park Based on Snow Tracking and Telemetry. *Journal of*

Mammalogy, *88*(6), 1498–1507. https://doi.org/10.1644/07-MAMM-A-076.1

Wilmers, C. C., Crabtree, R. L., Smith, D. W., Murphy, K. M., & Getz, W. M. (2003). Trophic

facilitation by introduced top predators: Grey wolf subsidies to scavengers in

Yellowstone National Park. *Journal of Animal Ecology*, *72*(6), 909–916.

https://doi.org/10.1046/j.1365-2656.2003.00766.x

Figure 1 | Locations of camera stations, gradient of habitat covariates, and photo-capture rates of
mesocarnivores (red fox = red, coyote = blue) and apex carnivores (wolf = green, cougar = pink)
in northern Yellowstone National Park, USA.

Figure 2 | Pairwise comparisons of Pianka's spatial index overlap for study species, computed
from detections at camera stations. Estimates are yearly means and 95% CIs.

 Figure 3 | (a) Apex and mesocarnivore naive occupancy probabilities estimated from camera
 trapping surveys between 2021–2023. **(b)** Coyote occupancy probability for 2021–2023,
 conditional on wolf presence-absence. Conditional occupancy was based on the best-supported,
 second-order multi-species occupancy model for each session.

Figure 4 | Diel activity patterns with 95% bootstrapped CIs for study carnivores. Mean activity
overlap **(a)** between apex and mesocarnivores, and **(b)** between apex carnivores and between
mesocarnivores.

Figure 5 | (a) Daily binned temporal attraction-avoidance of mesocarnivores in response to apex
carnivore detections at camera stations. Apex-mesocarnivore pairs include wolf-coyote (top-left),
wolf-red fox (top-right), cougar-coyote (bottom-left), and cougar-red fox (bottom-right). Points
and lines represent observed mesocarnivore detection probabilities, while violin plots show the
distribution of expected mesocarnivore detection probabilities based on random movements (i.e.,
no attraction or avoidance). *denotes significant differences between observed and expected
detection probabilities. **(b)** Median time differences of mesocarnivore detections following apex
carnivore detections.

Figure 6 | Wolf (n = 327) and cougar (n = 257) kills, surveyed through GPS-cluster searches of
collared wolves and cougars between 2016-2023, bearing **a)** presence of coyote and red fox
scavenging expressed as a proportion, and **b)** counts of coyote and red fox killed. For methods
see main text. Wolf and elk carcass graphics by Kira Cassidy.

769 **SUPPLEMENTARY INFORMATION FOR:**

771 **Title:** Species-specific interactions with apex carnivores yield unique benefits and burdens for

772 mesocarnivores

773 **Table S1:** Covariate List for Detection and Occupancy Models & their Description and sources
774

Occupancy Covariates		
Variable	**Type, Range**	**Description**
Cover	Continuous	Values accrued from the Rangeland Analysis Platform's (RAP) annual percent tree cover landsat satellite data (Robinson et al., 2019). Originally at 30-m spatial resolution but aggregated to 1 km. We used the cover value associated with the cell that a given camera station lies within.
Topographic Roughness	Continuous	Values represent the sum of the absolute value of the difference in elevation between each grid cell and its surrounding eight neighbors (3 x 3 window) (Kohl et al., 2019; Ruth et al., 2019). A value of 0 represents a flat surface while the maximum value represents the roughest terrain in the study area. Originally at 30-m spatial resolution but aggregated to 1 km. We used the roughness value associated with the cell that a given camera station lies within.
Photo Capture Rates	Continuous	Number of independent photographs of a species captured at a camera station divided by the total number of operational camera days. We used photo-captures for each species of interest segregated by sessions.
Snow Depth	Continuous	Daily snow depth data at 1-km grid cell accrued from SNODAS. We used the average value of all days within each session, so that snow depth values in a grid varied between but not within sessions.
Detection Covariates		
Cover Ratio	Continuous	Values accrued from the Rangeland Analysis Platform's (RAP) annual percent tree cover landsat satellite data (Robinson et al., 2019), originally at 30 m spatial resolution. We used the 30 m cover value at a given camera station and divided it by the mean cover within a 1-km buffer around that camera station. High values indicate a large difference between cover at the camera station relative to the surrounding landscape
Topographic Roughness Ratio	Continuous	We used the 30-m roughness value (Kohl et al., 2019) at a given camera station and divided it by the mean roughness within a 1-km buffer around that camera station. High values indicate a large difference between roughness at the camera station relative to the broader surrounding landscape and may be suggestive of a "pinch point" that funnels wildlife through it.
Number of Cameras	Continuous, Range: 0-2	Number of active/operational cameras at a given station.

776 **Table S2 |** Pianka's spatial overlap index for each species pairwise comparison during each
777 session.
778

Species Pair	2021			2022			2023		
	Mean	SE	95% CI	Mean	SE	95% CI	Mean	SE	95% CI
Red Fox – Coyote	0.56	0.12	0.33 – 0.78	0.57	0.13	0.28 – 0.78	0.48	0.14	0.24 – 0.74
Red Fox – Wolf	0.74	0.08	0.58 – 0.87	0.58	0.11	0.34 – 0.78	0.7	0.1	0.48 – 0.87
Red Fox – Cougar	0.58	0.13	0.32 – 0.8	0.55	0.11	0.36 – 0.76	0.45	0.16	0.16 – 0.74
Coyote – Wolf	0.63	0.15	0.30 – 0.86	0.49	0.11	0.29 – 0.7	0.58	0.17	0.26 – 0.85
Coyote – Cougar	0.2	0.11	0.06 – 0.47	0.34	0.12	0.13 – 0.6	0.19	0.11	0.05 – 0.49
Wolf – Cougar	0.38	0.13	0.15 – 0.61	0.3	0.1	0.13 – 0.53	0.31	0.16	0.07 – 0.69

780 **Table S3 |** Detection and occupancy covariates in the best-supported first-order models for red
781 fox, coyote, wolf, and cougar in each session.
782

Session-Species	Detection model (p)	Top occupancy model (Ψ)
2021		
Red Fox	*Cover ratio*	*Null*
Coyote	*Null*	*Null*
Wolf	*Null*	*Null*
Cougar	*Cover ratio*	*Roughness*
2022		
Red Fox	*Roughness ratio*	*Null*
Coyote	*Cover ratio*	*Null*
Wolf	*Null*	*Null*
Cougar	*Cover ratio*	*Snow depth + Roughness*
2023		
Red Fox	*Active cameras + Cover ratio*	*Null*
Coyote	*Cover ratio*	*Null*
Wolf	*Roughness ratio*	*Null*
Cougar	*Active cameras + Cover ratio*	*Cover + Roughness*

784   **Table S4 |** AIC model comparison for all candidate first- and second-order multi-species
785   occupancy models considered during each session. Bolded rows indicate selected models that we
786   reported results for. K = number of model parameters, AIC = Akaike Information Criterion,
787   ΔAIC = difference in AIC from the top model.
788

Model	K	AIC	ΔAIC	AIC$_{weight}$
2021				
First-Order Detection + Occupancy	11	2870.8	0	0.54
Second-Order Detection Only	16	2871.5	0.7	0.39
Second-Order Detection + Occupancy (Penalized)	**17**	**2875**	**4.2**	**0.067**
First Order Detection Only	10	2885.7	14.9	0.0003
Second-Order Null	14	2889.3	18.5	5.2E-05
First-Order Null	8	2903.3	32.4	4.9E-08
2022				
Second-Order Detection + Occupancy	**19**	**2300**	**0**	**0.78**
First-Order Detection + Occupancy	13	2302.7	2.7	0.21
Second-Order Detection Only	17	2308.3	8.3	0.012
First Order Detection Only	11	2310.6	10.6	0.004
Second-Order Null	14	2314.8	14.8	0.0005
First-Order Null	8	2316.6	16.6	0.0002
2023				
Second-Order Detection + Occupancy	**22**	**2327.6**	**0**	**0.57**
First-Order Detection + Occupancy	16	2328.2	0.6	0.43
Second-Order Detection Only	20	2341.2	13.6	0.0007
First Order Detection Only	14	2344.3	16.7	0.0001
Second-Order Null	14	2376.4	48.7	1.5E-11
First-Order Null	8	2380.1	52.5	2.3E-12

790 **Table S5 |** Beta coefficient estimates for detection and occupancy covariates retained in the top
791 second-order multi-species occupancy model for each session.
792

Detection model (p)						Occupancy model (Ψ)				
[Species] Covariate	β	SE	z	P		[Species] Covariate	β	SE	z	P
2021										
[redfox] (Intercept)	-0.29	0.26	-1.14	0.253		[redfox] (Intercept)	0.78	0.58	1.35	0.178
[redfox] cover ratio	-0.19	0.08	-2.27	0.023		[coyote] (Intercept)	-0.79	0.39	-2.01	0.045
[coyote] (Intercept)	-1.11	0.20	-5.52	3.3E-08		[wolf] (Intercept)	-1.09	0.39	-2.83	0.005
[wolf] (Intercept)	-1.56	0.18	-8.45	3.0E-17		[cougar] (Intercept)	0.45	0.57	0.78	0.433
[cougar] (Intercept)	-1.54	0.39	-3.91	9.2E-05		[cougar] scale(roughness)	1.71	0.52	3.32	0.001
[cougar] cover ratio	-0.16	0.17	-0.92	0.357		[redfox:coyote] (Intercept)	1.47	0.31	4.68	2.9E-06
						[redfox:wolf] (Intercept)	1.14	0.34	3.33	0.001
						[redfox:cougar] (Intercept)	1.20	0.40	2.96	0.003
						[coyote:wolf] (Intercept)	1.35	0.49	2.74	0.006
						[coyote:cougar] (Intercept)	-1.00	0.55	-1.81	0.071
						[wolf:cougar] (Intercept)	-0.10	0.51	-0.19	0.847
2022										
[redfox] (Intercept)	-0.67	0.20	-3.33	8.7E-04		[redfox] (Intercept)	0.51	0.88	0.58	0.561
[redfox] roughness ratio	-0.25	0.15	-1.71	0.088		[coyote] (Intercept)	-1.32	1.69	-0.79	0.432
[coyote] (Intercept)	-2.33	0.28	-8.20	2.3E-16		[wolf] (Intercept)	-1.39	1.71	-0.81	0.417
[coyote] cover ratio	0.14	0.06	2.44	0.015		[cougar] (Intercept)	-2.04	2.44	-0.84	0.402
[wolf] (Intercept)	-2.03	0.18	11.49	1.5E-30		[cougar] scale(snow depth)	-0.94	0.49	-1.92	0.055
[cougar] (Intercept)	-1.64	0.28	-5.92	3.2E-09		[cougar] scale(roughness)	1.16	0.57	2.04	0.042
[cougar] cover ratio	-0.11	0.12	-0.90	0.369		[redfox:coyote] (Intercept)	0.51	1.80	0.28	0.778
						[redfox:wolf] (Intercept)	0.78	1.78	0.44	0.660
						[redfox:cougar] (Intercept)	3.04	2.41	1.26	0.208
						[coyote:wolf] (Intercept)	2.30	1.06	2.16	0.031
						[coyote:cougar] (Intercept)	0.07	1.08	0.06	0.949
						[wolf:cougar] (Intercept)	-0.54	1.12	-0.48	0.629
2023										
[redfox] (Intercept)	-1.57	0.43	-3.69	2.2E-04		[redfox] (Intercept)	0.37	0.55	0.68	0.497
[redfox] active cameras	0.63	0.22	2.80	0.005		[coyote] (Intercept)	-1.16	0.78	-1.48	0.138
[redfox] cover ratio	-0.11	0.05	-2.20	0.028		[wolf] (Intercept)	-1.69	0.94	-1.81	0.070
[coyote] (Intercept)	-1.73	0.23	-7.38	1.6E-13		[cougar] (Intercept)	-2.27	1.19	-1.92	0.055
[coyote] cover ratio	0.16	0.06	2.69	0.007		[cougar] scale(roughness)	1.60	0.59	2.69	0.007
[wolf] (Intercept)	-0.52	0.27	-1.89	0.058		[cougar] scale(cover)	-1.10	0.59	-1.87	0.061
[wolf] roughness ratio	-0.78	0.26	-3.03	0.002		[redfox:coyote] (Intercept)	0.55	0.87	0.64	0.525
[cougar] (Intercept)	-6.06	2.46	-2.47	0.014		[redfox:wolf] (Intercept)	1.23	0.97	1.27	0.205
[cougar] active cameras	2.57	1.23	2.09	0.036		[redfox:cougar] (Intercept)	1.74	1.17	1.49	0.137
[cougar] cover ratio	-0.34	0.12	-2.74	0.006		[coyote:wolf] (Intercept)	1.23	0.67	1.84	0.066
						[coyote:cougar] (Intercept)	0.34	0.74	0.46	0.648
						[wolf:cougar] (Intercept)	0.36	0.77	0.47	0.637

794 **Figure S1 |** Detection probabilities for each species as a function of covariates retained in the top
795 second-order multi-species occupancy model for each session. Shaded regions are 95%
796 confidence intervals.
797

799 **Figure S2 |** Marginal occupancy probability for cougar as a function of habitat covariates
800 retained in the top second-order multi-species occupancy model for each session. Shaded regions
801 are 95% confidence intervals.
802

REFERENCES

Kohl, M. T., Ruth, T. K., Metz, M. C., Stahler, D. R., Smith, D. W., White, P. J., & MacNulty,

D. R. (2019). Do prey select for vacant hunting domains to minimize a multi-predator

threat? *Ecology Letters*, *22*(11), 1724–1733. https://doi.org/10.1111/ele.13319

Robinson, N. P., Jones, M. O., Moreno, A., Erickson, T. A., Naugle, D. E., & Allred, B. W.

(2019). Rangeland Productivity Partitioned to Sub-Pixel Plant Functional Types. *Remote*

Sensing, *11*(12), 1427. https://doi.org/10.3390/rs11121427

Ruth, T., Buotte, P., & Hornocker, M. (2019). *Yellowstone Cougars: Ecology Before And During*

Wolf Restoration. University Press of Colorado. https://doi.org/10.5876/9781607328292

814