1   **Title:** Species-specific interactions with apex carnivores yield unique benefits and burdens for
2   mesocarnivores
3

4   **Authors and Affiliations**: Wesley Binder[1,2]*, Jack W. Rabe[2,3]*, Zoe Lowe[4], Gordon Scott[2],
5   Claire Lacey[2], Eliza King[4], Daniel R. Stahler[2], Stotra Chakrabarti[4]
6   *Contributed Equally
7
[1]Department of Fisheries, Wildlife, and Conservation Sciences, Oregon State University,
Corvallis, Oregon, USA
[2]Yellowstone Center for Resources, Yellowstone National Park, Mammoth Hot Springs,
Wyoming, USA
[3]Department of Fisheries, Wildlife, and Conservation Biology, University of Minnesota – Twin
Cities, Saint Paul, Minnesota, USA
[4]Department of Biology, Macalester College, Saint Paul, Minnesota, USA

15
Author Contributions: WB, JWR, DRS, and SC conceptualized the study, raised resources, and
oversaw project completion. WB, DRS, CL, and GS conducted fieldwork and collected the data.
WB, JWR, and SC developed the methodology and analytical approaches. WB, JWR, ZL, GS,
and EK curated the data and performed the analyses, with inputs from SC and DRS. WB, JWR,
and SC wrote the manuscript, with help from GS. All authors reviewed subsequent drafts and
approved the final version.
22

Acknowledgements: We are grateful to the numerous field technicians from the Yellowstone
Wolf and Cougar Projects who made data collection possible. We would also like to thank the
fStop Foundation for providing necessary equipment. We acknowledge Yellowstone Forever
(and their contributing donors), the National Science Foundation, Oregon State University,
University of Minnesota, National Park Service, US Geological Survey, and Macalester College
for funding support.

29
30   **Data Access Statement:** All data and code used for analyses can be found in the following
31   repository: https://zenodo.org/records/14852260
32
33   **Ethics Statement:** Wildlife captures and handling were conducted under National Park Service
34   protocols (IACUC permits IMR_YELL_Stahler_Cougar_2018_A1,
35   IMR_YELL_Smith_Wolf_2012).
36
37   **Conflict of Interest:** The authors declare no conflicts of interest.
38
39   **Corresponding Author:** Wesley Binder (wesley.binder@oregonstate.edu)
40

ABSTRACT

Mesocarnivores navigate a complex *risk-reward* continuum in ecosystems shared with their apex counterparts, balancing scavenging opportunities with risks of mortality. However, the risks to mesocarnivores in multi-carnivore systems are not uniform—often varying with specific apex-meso pairings. Using camera traps and GPS-telemetry, we examined space-use, temporal activity, fine-scale interactions, and scavenging behavior of apex (wolves, cougars) and mesocarnivores (red foxes, coyotes) in northern Yellowstone National Park, USA. Coyote space use was positively linked to wolf presence, while red fox space use was positively linked to cougar presence. Notably, photo-detection rates of mesocarnivores doubled within 24 hours of apex carnivore detections—except for coyotes following cougars. Coyotes were more frequent scavengers at wolf and cougar kills than red foxes, but they also suffered the highest mortality rates. Over 60% of wolf caused coyote mortalities were linked to wolf kill-sites, though wolves rarely consumed the coyotes themselves. In contrast, cougars hunted and consumed coyotes as prey, posing an additional risk of predation. These findings highlight how divergent hunting strategies and habitat preferences between canid and felid apex carnivores create separate risks for mesocarnivores. Our study offers new insights into the species-specific behavioral dynamics that can shape trophic interactions in multi-carnivore systems.

KEYWORDS: behavioral ecology, intra-guild competition, landscape of fear, mesopredator release, trophic-mediated responses, Yellowstone

INTRODUCTION

Intraguild carnivore interactions play a pivotal role in ecosystem function, shaping the abundance, distribution, and behavior of both carnivores and their prey through top-down effects [1]. These interactions, shaped by the complex strategies employed by carnivores across different trophic levels to coexist, are particularly important to understand given that human activities have disproportionately disrupted apex carnivores in the Anthropocene. Humans have both driven significant population declines and local extinctions of apex carnivores in some areas [1] and facilitated their recovery through reintroduction and conservation efforts in others [2–4]. These actions, in turn, impact carnivore communities, as apex carnivores can affect mesocarnivores both directly and indirectly. For instance, wolves (*Canis lupus*) can suppress coyote (*Canis latrans*) populations through both lethal removal and resource competition, which in turn reduces the top-down effects that coyotes exert on red fox (*Vulpes vulpes*) populations [5]. Thus, the extent to which mesocarnivores are affected by the presence of apex carnivores depends on their relative position within the trophic community.

However, the factors mediating the dynamics between apex and mesocarnivores are complex and often difficult to untangle. While apex carnivores can suppress mesocarnivore populations, they can also benefit them by providing scavenging opportunities from the carcasses of their prey [6–8]. As a result, mesocarnivore distributions and behavior in relation to their apex counterparts depend on a delicate balance of risks (interference competition) and rewards (carcass availability) that vary in both space and time [9].For instance, a global review revealed that 30% of mesocarnivore diets were composed of scavenged ungulate carcasses, yet a nearly identical proportion—about one-third—of their deaths were attributed to apex carnivores [10]. Larger mesocarnivores, in particular, were more reliant on scavenging opportunities, and this

facilitation appeared to be more costly in less productive environments, leading to greater

suppression from apex carnivores [10]. The relationships between mesocarnivores and apex

carnivores are thus shaped by several factors, including the mesocarnivores' reliance on

scavenging to supplement their diets, their ability to locate apex carnivore kill sites quickly, and

the extent to which apex carnivores influence mesocarnivore mortality risks.

The risks faced by mesocarnivores also depend on how they interact with specific apex

carnivores. For example, when mesocarnivores are primarily competing for resources—such as

carcasses—their risks mostly arise from direct encounters with apex carnivores at kill sites [10].

In contrast, if mesocarnivores are killed by apex carnivores independent from carcass sites, then

their risks are influenced by the habitat preferences and activity patterns of the apex carnivore

[11]. Such relationships likely drive significant variation in intraguild mortality rates across

carnivore family groups. For example, canid mesocarnivores experience mortality from canid

apex carnivores at rates five times greater than from felid apex carnivores [10]. Thus, the risks

and rewards mesocarnivores experience when associating with their apex counterparts should

vary with specific meso-apex pairings, yet the mechanisms driving such differences are poorly

understood. Such species-specific interactions can be critical in further mediating mesocarnivore

landscapes of fear, and in turn have effects on entire ecosystems.

In the Northern Range of Yellowstone National Park, wolves and cougars (*Puma*

concolor) are abundant during winter when they are supported by a diverse assemblage of

herbivores such as elk (*Cervus canadensis*), bison (*Bison bison*) and deer (*Odocoileus* spp.) that

congregate at high densities [12–14]. Wolves and cougars kill larger ungulate prey during these

months [15–17], and such large carcasses provide scavenging opportunities to a host of

mesocarnivores including coyotes and red foxes [8]. Such scavenging opportunities appear to be

critical for mesocarnivores in the winter because much of their regular prey (small mammals and

ungulate neonates, birds, fruits, insects) become scarce and/or subnivean [8]. Coyotes and red

foxes, however, also navigate the threat of lethal encounters with wolves and cougars while

accessing these carcasses, and in general. To examine how competitive and facilitative

relationships vary by specific apex-mesocarnivore pairings, we evaluated the degree to which

coyotes and red foxes exhibited spatial and temporal avoidance, or attraction, to wolves and

cougars in this system.

We expected each apex-mesocarnivore pairing to exhibit distinct associations for two

reasons. First, long-term anecdotes from Yellowstone suggest the mechanisms of lethal

interactions for mesocarnivores differ by apex carnivore species. Cougars appear to hunt

mesocarnivores as prey, while wolves are more likely to kill them at ungulate-carcass sites as a

form of resource defense. Therefore, we anticipate mesocarnivores to exhibit more avoidance of

cougars compared to wolves given the higher threats associated with cougar predation. Second,

we expected coyotes to exhibit more attraction to apex carnivores than red foxes as their greater

size and energetic demands should increase their reliance on scavenging kills when alternative

resources are scarce. To explore these relationships, we conducted winter camera-trapping

surveys to assess occupancy, diel activity patterns, and fine-scale spatiotemporal responses of

apex and mesocarnivores. We also monitored predation events of GPS-collared wolves and

cougars to quantify mesocarnivore scavenging at kills, as well as direct killings of

mesocarnivores.

MATERIALS AND METHODS

Study Area

We conducted the study in the Northern Range of Yellowstone National Park (Figure 1),

an area with elevations ranging from 1500 to 2400 meters. The landscape features a mix of

rugged, rocky terrain and open, rolling hills. Common vegetation types in the region include

Douglas-fir (*Pseudotsuga menziesii*), sagebrush (*Artemisia* spp.), and juniper (*Juniperus spp.*).

The area experiences long, cold winters, with average snow-water equivalents ranging from 2 to

30 cm [18]. Apex carnivores, such as wolves, grizzly bears (*Ursus arctos horribilis*), black bears

(*U. americanus*), and cougars, are present in relatively high numbers, while coyotes, red foxes,

bobcats, and martens (*Martes americana*) are the prominent mesocarnivores [14,17].

Camera Trap Surveys

Camera surveys were conducted within a 580 km² area, defined by a 5-km buffer

surrounding our camera-trap grid, which was designed to estimate cougar densities. We deployed

camera stations in a "checkerboard" pattern to increase the variability in station spacing [19],

using a grid of 3.2 km² cells [20]. In each selected grid cell, we set up two camera stations at

locations chosen to maximize detections, such as natural movement corridors or pinch points,

with a minimum distance of 1 km between stations.

Camera stations were active during 15-week winter sampling periods (referred to as

"sessions") from December to March in the years 2020-21, 2021-22, and 2022-23 (hereafter

referred to as 2021, 2022, and 2023, respectively). We used Browning Spec Ops trail cameras

equipped with motion-activated passive infrared sensors. In 2021, we deployed 90 cameras

across 60 stations, with half of the stations having a single camera and the other half having two

cameras placed opposite each other. For 2022 and 2023, all stations were equipped with two

cameras, bringing the total number of cameras to 120. Each camera was configured for long-

range motion detection and recorded 20-second videos with a one-minute delay between

recordings. We manually reviewed all videos, identified species, and extracted metadata using

the Timelapse2 image analysis software [21].

Data Analysis

Multi-Species Occupancy Models

To evaluate how habitat covariates and spatial interactions with other carnivores

influence the occurrence of apex and mesocarnivores, we constructed multi-species occupancy

models [22] using the *unmarked R* package [23,24]. Occupancy models incorporate binary

presence-absence (1/0) data through likelihood-based approaches to estimate species occurrence

at site i in a latent z (occupancy) "state" model, while accounting for imperfect detections

through a separate "detection" model. Multi-species occupancy models are an extension of the

single-species occupancy model [25] that apply a hierarchical modeling framework to estimate

the effect of covariates (first-order) and species co-occurrence (second-order) on occupancy.

We used 1-km^2 grid cells for modeling occupancy, with each cell containing no more

than one camera station. Because wolves, cougars, coyotes, and potentially red foxes typically

occupy ranges larger than 1 km^2, the close spacing of our cameras may violate assumptions of

spatial independence. Hence, our occupancy estimates more aptly describe patterns of "space

use" rather than distribution [26]. For estimating detection probabilities of each species, we

considered two covariates: the number of active cameras at a station and snow depth. As the

number of active cameras at a station (ranging from 0–2 as a product of how many cameras were

active and for how long during week-long occasions) increased, we expected an increase in

detection probability. We used mean snow depth at camera stations during each week-long

occasion at a 1-km^2 resolution using SNODAS [27]. We expected a negative relationship

between snow depth and detection probability, such that increasing snow depth as winter

sessions progressed would lead to fewer individuals moving through camera stations, lowering

detection probability. For estimating species occupancy, we included topographic roughness,

percent tree cover, and average snow depth at a 1-km^2 spatial resolution to account for bottom-up

processes, such as habitat and landscape characteristics, that influence carnivore distributions

[28]. These habitat covariates were included because they are known to affect the occupancy and

movement of carnivores [14]. Please refer to Table S1 for covariate details.

Our primary interest was in estimating species co-occurrence patterns and the effects of

landscape covariates on occupancy, rather than temporal occupancy dynamics. We therefore

adopted a 'stacked' data structure, where each site ✕ year combination was treated as an

independent site [29,30], resulting in 178 sites (58 in 2021, and 60 in 2022 and 2023). To

account for the non-independence of sites across years, we included year as a fixed effect on

both detection and occupancy in all models. Using a two-stage model selection approach [31],

we first constructed single-species models to evaluate which covariates best explained detection

probability while holding occupancy constant at the null model ($\Psi(.)$). For each species, we used

Akaike's Information Criterion (AIC) to select the best detection model from four candidate

models representing all possible additive combinations of the two detection covariates, including

the null model with only year as a fixed effect. Second, we carried over the top detection sub

model to select the best first-order model by considering all possible additive combinations of

covariates on occupancy (eight candidate models for each species). Following Arnold (2010), we

accounted for parameter redundancy by selecting more parsimonious models if they were within

2 AIC units of a model with more parameters [32]. After selecting the top first-order model, we

added second-order species interactions to develop multi-species models that tested the effects of

species co-occurrence on occupancy. Please refer to Table S3 and S4 for a list and comparison of

candidate models.

Temporal Activity Patterns

We used the *activity* package in R to estimate diel activity patterns for wolves, cougars,

coyotes, and red foxes using non-parametric kernel density estimation [33]. To maintain

independence of detections, photo-captures of the same species within 30 minutes were omitted

[34]. To quantify temporal niche partitioning between species and identify potential coexistence

strategies that extend beyond broad-scale occupancy, we used the *overlap* package in R to assess

overlap of diel activity patterns between each apex-meso pairing by calculating the area under

the combined activity curves, $\widehat{\Delta_4}$ [35]. We generated 10,000 bootstrap samples of the activity

overlap coefficient, $\widehat{\Delta_4}$, for each species-pair to estimate 95% confidence intervals of overlap.

Time-to-event

We examined fine-scale mesocarnivore activity in response to apex carnivore movements

using time-to-event analysis [36,37]. The elapsed time between a mesocarnivore detection

following an apex carnivore detection was recorded at each camera station for a given session.

Subsequently, we combined data across all sessions for each apex-meso pairing. We truncated

apex carnivore detections to only include events that were followed by a mesocarnivore

detection (i.e., apex carnivore detections followed by another apex carnivore detection were

omitted). Detections of mesocarnivores were recorded for 5 days following an apex carnivore

detection, with elapsed time differences binned into 24-hour periods. We estimated detection

probabilities by dividing the number of mesocarnivore detections in each 24-hour period by the

total number of detections across all sessions [36,37]. These detection probabilities were

compared to random iterations (n = 1,000) of mesocarnivore detections to derive expected

detection probabilities for each 24-hour period by using standard two-tailed permutation tests

[37]. If mesocarnivores were avoiding apex carnivores through fine-scale temporal segregation ,

we would expect coyote and red fox detection probabilities to be significantly lower following an

apex carnivore detection. The opposite would apply if they were instead tracking/following

wolves and cougars, likely to acquire food by scavenging apex carnivore kills .

Monitoring of Apex Carnivore Predation Events

To assess mesocarnivore scavenging at apex carnivore kills and direct killing of

mesocarnivores, we monitored predation by GPS-collared wolves (n = 26 from 8 packs) and

cougars (n = 17) during 30-day winter sampling periods (November to March) from 2016–2023.

Monitored wolves and cougars were fitted with Vectronics or Telonics GPS-Satellite collars that

were programmed to collect hourly fixes. We searched GPS-location clusters [38] to identify kill

sites and conducted necropsies on prey remains. At each kill, we ascertained the likely cause of

death, the species, sex, age of prey when possible, and the presence of scavengers through visual

observations and sign (e.g., scat, tracks, hair). To complement our camera-trap analysis, we also

filtered this long-term predation data by selecting scavenging and predation events that occurred

only during the three winter camera-trap sessions, and assessed whether the trends during those

years differed from our longer-term data (Figure S5).

RESULTS

Across all three 15-week winter camera-trapping sessions, we collectively monitored for

18,617 sampling days. Red foxes were detected at 88% of sites, coyotes at 56% of sites, wolves

at 53% of sites, and cougars at 47% of sites (Figure S2).

Multi-Species Occupancy Models

Detection probability

Variation in red fox detection probability was best explained by snow depth, with red fox

detection increasing as snow depth decreased (β = -0.21, 95% CI = -0.34 to -0.08, P = 0.0018).

Holding snow depth at its mean, red fox detection varied by year with marginal detection

probabilities of 0.33 (95% CI = 0.29 – 0.36) in 2021, 0.24 (95% CI = 0.21 – 0.28) in 2022, and

0.39 (95% CI = 0.34 – 0.43) in 2023. While neither the number of active cameras nor snow depth

significantly affected coyote or wolf detection probability, there was variation between sessions.

Coyote detection probability was 0.25 (95% CI = 0.21 – 0.28) in 2021, 0.14 (95% CI = 0.11 –

0.18) in 2022, and 0.22 (95% CI = 0.18 – 0.27) in 2023. Wolf detection probability was 0.17

(95% CI = 0.14 – 0.21) in 2021, 0.12 (95% CI = 0.09 – 0.16) in 2022, and 0.21 (95% CI = 0.18 –

0.26) in 2023. Cougar detection probability did not vary annually, but did increase significantly

as snow depth decreased (β = -0.27, 95% CI = -0.51 to -0.03, P = 0.03).

Occupancy probability

In the top-performing first-order model, habitat covariates only explained significant

variation in cougar occupancy, while the null "year-only" model performed best for red fox,

coyote, and wolf. Cougar occupancy probability increased as mean snow depth decreased, (β = -

1.06, 95% CI = -1.99 to -0.13, P = 0.025) and roughness increased (β = 1.4, 95% CI = 0.78 –

2.03, P < 0.0001; Figure S2). After incorporating second-order species interactions on

occupancy, we found that coyote occupancy was positively associated with wolf occupancy (β =

1.62, 95% CI = 0.81 – 2.42, P < 0.0001). Coyote occupancy was 0.75 (95% CI = 0.58 – 0.86) at

sites where wolves were present and 0.35 (95% CI = 0.20 – 0.55) where wolves were absent

(Figure 2b). Red fox occupancy was positively associated with cougar occupancy (β = 1.94, 95%

CI = 0.51 – 3.36, P = 0.0077) (Figure 2b). Where cougars occurred, red fox occupancy was 0.96

(95% CI = 0.86 – 0.99), compared to 0.82 (95% CI = 0.63 – 0.91) where cougars did not occur.

Temporal Activity Patterns

Cougars (n = 341 detections) exhibited unimodal diel activity, being most active around

dusk peaking at 1800 and least active from morning (0800) to midday (1400) (Figure 3). Wolves

(n = 443 detections) had bimodal, crepuscular activity patterns, with peak activity shortly after

dawn (~0800–0900) and a secondary peak at dusk (1700–1800) (Figure 3). Coyotes (n = 541

detections) had more consistent activity across the diel cycle and were more diurnal, with peak

activity occurring from late morning (0900) to midday (1400) and the lowest activity before

dawn (0500) and after dusk (2100) (Figure 3). Red foxes (n = 1520 detections), on the other

hand, exhibited highly nocturnal patterns, being most active shortly after dusk (1800–2100) and

sustaining high activity through the night (2100–0600) with little activity during daylight hours

(0800–1600) (Figure 3). Temporal overlap between apex and mesocarnivores was $\widehat{\mathit{\Delta}_4}$ = 0.86

(CI: 0.81 – 0.89) for wolf-coyote, 0.68 (CI: 0.61 – 0.7) for wolf-red fox, 0.81 (CI: 0.74 – 0.85)

for cougar-coyote, and 0.72 (CI: 0.66 – 0.76) for cougar-red fox across the three sessions (Figure

3a). Overlap between mesocarnivores (red fox-coyote) was $\widehat{\mathit{\Delta}_4}$ = 0.61 (CI: 0.53 – 0.62), while

overlap between apex carnivores (wolf-cougar) was $\widehat{\mathit{\Delta}_4}$ = 0.86 (CI: 0.8 – 0.91) (Figure 3b).

Time-to-event

Compared to the mean values derived from random iterations of detection probabilities,

red foxes and coyotes were both over twice as likely to be detected in the 24-hour period

following a wolf detection (red foxes: 0.034 compared to 0.015, P = 0.001; coyotes: 0.05

compared to 0.018; P = 0.001; Figure 4a). However, their detection probabilities following

cougar detections differed. While red foxes were over twice as likely to be detected in the 24-

hour period following a cougar detection (0.024 compared to 0.011; P = 0.001), coyote detection

rates were not affected. Additionally, red foxes were more likely to be detected in the 72-hour

292 period following a wolf detection (0.01 compared to 0.007; $P = 0.04$) and the 96-hour period

293 following a cougar detection (0.009 compared to 0.004; $P = 0.001$). Overall, the time interval

294 between detections of a mesocarnivore subsequent to an apex carnivore detection was shortest

295 for wolf-coyote (median = 17.67 hours) and longest for cougar-coyote (median = 57.35 hours).

296 Foxes showed similar latency to wolf and cougar detections, with medians of 21.11 and 24.69

297 hours, respectively (Figure 4b).

298 **Monitoring of Apex Carnivore Predation Events**

299  We detected 327 wolf kills and 257 cougar kills through winter predation monitoring

300 (2016–2023). Coyote presence was recorded at 222 (68%) wolf kills and 80 (31%) cougar kills

301 (Figure 5a). Red fox presence was recorded at 67 (20%) wolf kills and 46 (18%) cougar kills

302 (Figure 5a). From this predation monitoring, we documented 18 coyotes and one red fox killed

303 by wolves (Figure 5b). Of the coyotes killed by wolves, 11 (61%) occurred at wolf feeding sites

304 (i.e., ungulates killed or scavenged by wolves), while the red fox mortality did not occur at a

305 wolf-killed prey site. Cougars killed eight coyotes and three red foxes during this time (Figure

306 5b). None of these mortalities were associated with cougar-killed prey sites, rather they were

307 fully consumed and generally found in cache piles - suggestive of cougar consumption. We

308 found no major differences between the longer-term data and the data overlapping with our

309 camera-trapping study (see Figure S5 for filtered results).

310 **DISCUSSION**

311 Ecological communities are influenced by the top-down effects that carnivores exert on prey

312 populations, but these effects also extend to other carnivores [1]. Intraguild carnivore interactions

313 are complex in that the negative impacts of competition can be partially offset by the scavenging

opportunities competitors provide with their kills. While these tradeoffs are common between

apex and mesocarnivores, we found considerable variation across species-specific pairings.

Red foxes and coyotes were both ubiquitous in our study area and common scavengers at

wolf and cougar kills, but exhibited unique spatial and temporal relationships with each apex

carnivore. Our multi-species occupancy models revealed positive associations between the space

use of coyotes and wolves, as well as red foxes and cougars. Further, red foxes and coyotes

exhibited opposite diel activity patterns (Figure 3b), as the nocturnal activity of red foxes closely

aligned with that of cougars, while coyotes were most active mid-day when apex carnivores were

least active. Their fine-scale movements also differed in response to apex carnivore presence.

Red fox detection probabilities at camera stations doubled in the 24 hours following a wolf or

cougar detection, but coyotes only displayed this attraction to wolves, not cougars (Figure 4a).

Accordingly, we found that coyotes were more frequent scavengers at wolf kills compared to

cougar kills. However, scavenging on wolf kills was often costly, as over half the coyote

mortalities attributed to wolves during our study occurred at wolf kill sites. Interestingly, we did

not document wolves consuming the coyotes they killed, likely indicating that such lethal actions

resulted out of resource defense. Conversely, all coyotes killed by cougars occurred as

standalone predation events (i.e., independent of ungulate carcass sites) in which cougars

consumed the coyotes. The ambush hunting style of cougars, combined with the forested and

topographically rough terrain they inhabit, could lead to coyotes being more vulnerable in such

habitats [39]. In contrast, the open and flatter terrain preferred by wolves could provide more

visual and olfactory cues for coyotes to assess (and escape) immediate risks of lethal encounters.

Such disparities in apex carnivore hunting strategies seemingly tie back to the spatiotemporal

attraction of coyotes towards wolves but not for cougars, and highlight the nuances of such

competitive outcomes.

As mesocarnivores adapt their coexistence strategies to different apex carnivore species,

their behaviors are likely to shift more dramatically when multiple competitors are present. For

example, high winter apex carnivore densities in our study system may explain the midday peak

in activity for coyotes, which differed from the more crepuscular activity patterns of coyotes

inhabiting adjacent areas of the Greater Yellowstone Ecosystem with lower apex carnivore

densities [40]. In contrast, red foxes appeared less sensitive to the diel activity of apex

carnivores. Not only did their peak in activity align with that of cougars, but they also displayed

the same nocturnal patterns as red foxes across the Greater Yellowstone Ecosystem [40].

Notably, red foxes exhibited high temporal segregation from coyotes, having the lowest temporal

overlap of any species pair (Figure 4). Studies elsewhere have shown that coyotes and red foxes

compete severely, with coyotes competitively excluding and/or killing red foxes [5,41]. Thus, the

nocturnal activity of red foxes could be the result of avoiding peak coyote activity during the day

that is in turn mediated by wolf and cougar activity (Figure 4). Temporal sympatry amongst the

carnivore guild appears to be layered, with apex carnivores affecting coyotes, and coyotes

affecting red foxes.

Apex- and mesocarnivore relationships depend on competition that occurs within trophic

groups, as spatial and/or temporal responses to such competitors alters the availability of

ecological niche space for other carnivores in the community. For example, cougars in

Yellowstone changed their winter space use after wolves were reintroduced by increasing their

selection for areas with escape terrain (e.g., ruggedness, tree cover) [17]. This avoidance of

wolves could in turn facilitate coyote space use, providing habitats where encounters with

cougars are scarce. Our results support this, as coyote space use, fine-scale movements, and

scavenging rates were more strongly associated with wolves than cougars. Accordingly, if red

foxes experience the most severe competition from coyotes, then greater overlap with cougar

space use could alleviate such competition. While red foxes scavenged wolf and cougar kills at a

similar rate, our analyses revealed substantial overlap between cougar and red fox space use and

activity patterns. Mortalities from cougars still occurred, but we believe that foxes are more

adept at avoiding immediate risks of cougar predation during encounters through their increased

agility, as well as their relatively large foot:body ratio that allow for better maneuverability on

top of snow [42]. Coyotes being heavier, might not be as lucky when encountering cougars.

Thus, competition within trophic groups likely plays a key role in shaping the apex-

mesocarnivore interactions we observed, creating scenarios akin to "the enemy of my enemy is

my friend".

In contrast, competition between apex- and mesocarnivores is often driven by

scavenging, where apex carnivores take measures to reduce food loss (e.g., carcass defense,

caching, etc.) and mesocarnivores balance the need for such dietary subsidies with the associated

risks of confrontations at carcass sites [10]. Thus, the availability of alternative resources for

mesocarnivores should influence the extent to which they seek out scavenging opportunities and

the overall strength of their relationship with apex carnivores. In northern Yellowstone, the

limitations of prey acquisition during winter months may increase mesocarnivore reliance on

scavenging, particularly for coyotes who appear to depend on these subsidies more so than

smaller red foxes (Figure 5) [40,43]. Foxes may be better suited to meet their energetic demands

by hunting themselves, as they are commonly observed hunting subnivean rodents by pouncing

through the snowpack during the winter (D. Stahler *pers.comm.*). Therefore, apex carnivores

facilitate coyotes and red foxes differently, and these relationships are likely to change during the

spring/summer when the availability of rodents, neonatal ungulates, vegetation, and insects

increase [43]. The presence of grizzly and black bears that emerge in the summer after winter-

long hibernations can also complicate these relationships because bears often limit carrion

consumption for other scavengers [44].

While our study highlights the positive aspects of apex and mesocarnivore interactions

through scavenging, we did not explore variations in red fox and coyote population trends. In

fact, previous research documented a 39% decline in coyote abundance following wolf

reintroduction in the Lamar Valley of Yellowstone, an area adjacent to our study area [45].

Additionally, Merkle et al. (2009) showed a decrease in wolf-coyote interactions over time

(between 1997 and 2007) in our study area, indicative of changes to coyote density and/or

behavior (e.g., wolf avoidance) [46]. Despite benefiting from carrion made available by wolves

and cougars, red fox and coyote populations may still experience top-down population regulation

via direct killing that maintains them at low densities [47]. Long-term studies that estimate

population trends of mesocarnivores with respect to apex carnivore densities are required to test

such effects.

The magnitude and impact of trophic-mediated responses related to large carnivore

recoveries have become a contentious issue, perhaps nowhere more so than in Yellowstone

National Park [48–50]. Although the reintroduction of wolves and recolonization of cougars to

this area has likely altered mesocarnivore distributions and abundance, red foxes and coyotes

also benefit from living with their dominant counterparts. Overall, red foxes and coyotes in our

study area exhibited more attraction than avoidance to apex carnivores during winter months,

when small prey and alternative food resources are scant. Such attraction, possibly a

consequence of desperate foraging decisions in winter, can also prove fatal. However, risks to

mesocarnivores are uneven across dominant-subordinate carnivore pairings, with apex carnivore

behavior changing the drivers of these interactions, adding nuance to the *landscape of fear* for

mesocarnivores.

REFERENCES

1. Ripple WJ *et al.* 2014 Status and Ecological Effects of the World's Largest Carnivores.

Science **343**, 1241484. (doi:10.1126/science.1241484)

2. Bangs EE, Fritts SH. 1996 The Reintroduction of Gray Wolves to Yellowstone National Park

and Central Idaho - Final Environmental Impact Statement 1994. *Wildl. Soc. Bull.* , 414.

3. Chapron G *et al.* 2014 Recovery of large carnivores in Europe's modern human-dominated

landscapes. *Science* **346**, 1517–1519. (doi:10.1126/science.1257553)

4. Lotze HK, Coll M, Magera AM, Ward-Paige C, Airoldi L. 2011 Recovery of marine animal

populations and ecosystems. *Trends Ecol. Evol.* **26**, 595–605. (doi:10.1016/j.tree.2011.07.008)

5. Levi T, Wilmers CC. 2012 Wolves–coyotes–foxes: a cascade among carnivores. *Ecology* **93**,

921–929. (doi:10.1890/11-0165.1)

6. Brunet MJ *et al.* 2022 Cats and dogs: A mesopredator navigating risk and reward provisioned

by an apex predator. *Ecol. Evol.* **12**, e8641. (doi:10.1002/ece3.8641)

7. Sivy KJ, Pozzanghera CB, Grace JB, Prugh LR. 2017 Fatal Attraction? Intraguild Facilitation

and Suppression among Predators. *Am. Nat.* **190**, 663–679. (doi:10.1086/693996)

8. Wilmers CC, Crabtree RL, Smith DW, Murphy KM, Getz WM. 2003 Trophic facilitation by

introduced top predators: grey wolf subsidies to scavengers in Yellowstone National Park. *J.*

Anim. Ecol. **72**, 909–916. (doi:10.1046/j.1365-2656.2003.00766.x)

9. Ruprecht J *et al.* 2021 Variable strategies to solve risk–reward tradeoffs in carnivore

communities. *Proc. Natl. Acad. Sci.* **118**, e2101614118. (doi:10.1073/pnas.2101614118)

10. Prugh LR, Sivy KJ. 2020 Enemies with benefits: integrating positive and negative

interactions among terrestrial carnivores. *Ecol. Lett.* **23**, 902–918. (doi:10.1111/ele.13489)

11. Broekhuis F. 2015 Cat eats cat: leopard consumes adult cheetah, Maasai Mara Game

Reserve, Kenya. *CAT News* **65**, 33–34.

12. Houston DB. 1982 *The Northern Yellowstone Elk: Ecology and Management*. Macmillan.

13. White PJ, Wallen RL, Hallac D, Auttelet K, Jerret J. 2015 *Yellowstone Bison: Conserving an*

American Icon in Modern Society. Yellowstone Association.

14. Smith DW, Stahler DR, MacNulty DR. 2020 *Yellowstone Wolves: Science and Discovery in*

the World's First National Park. University of Chicago Press.

15. Metz MC, Smith DW, Vucetich JA, Stahler DR, Peterson RO. 2012 Seasonal patterns of

predation for gray wolves in the multi-prey system of Yellowstone National Park. *J. Anim.*

Ecol. **81**, 553–563. (doi:10.1111/j.1365-2656.2011.01945.x)

16. Metz MC. 2021 Estimating wolf predation metrics, patterns, and dynamics across time and

space in the multi-prey system of Yellowstone National Park. University of Montana. See

https://scholarworks.umt.edu/etd/11842/.

17. Ruth T, Buotte P, Hornocker M. 2019 *Yellowstone Cougars: Ecology Before And During*

Wolf Restoration. University Press of Colorado. (doi:10.5876/9781607328292)

18. Plumb GE, White PJ, Coughenour MB, Wallen RL. 2009 Carrying capacity, migration, and

dispersal in Yellowstone bison. *Biol. Conserv.* **142**, 2377–2387.

(doi:10.1016/j.biocon.2009.05.019)

19. Sun CC, Fuller AK, Royle JA. 2014 Trap Configuration and Spacing Influences Parameter

Estimates in Spatial Capture-Recapture Models. *PLOS ONE* **9**, e88025.

(doi:10.1371/journal.pone.0088025)

20. Anton C. 2020 The Demography and Comparative Ethology of Top Predators in a Multi-

Carnivore System. UC Santa Cruz. See https://escholarship.org/uc/item/3xd9379b.

21. Greenberg S, Godin T, Whittington J. 2019 Design patterns for wildlife-related camera trap

image analysis. *Ecol. Evol.* **9**, 13706–13730. (doi:10.1002/ece3.5767)

22. Rota CT, Ferreira MAR, Kays RW, Forrester TD, Kalies EL, McShea WJ, Parsons AW,

Millspaugh JJ. 2016 A multispecies occupancy model for two or more interacting species.

Methods Ecol. Evol. **7**, 1164–1173. (doi:10.1111/2041-210X.12587)

23. Fiske I, Chandler R. 2011 **unmarked** : An *R* Package for Fitting Hierarchical Models of

Wildlife Occurrence and Abundance. *J. Stat. Softw.* **43**. (doi:10.18637/jss.v043.i10)

24. R Core Team. 2023 R: A Language and Environment for Statistical Computing.

25. MacKenzie DI, editor. 2006 *Occupancy estimation and modeling: inferring patterns and*

dynamics of species. Amsterdam ; Boston: Elsevier.

26. Steenweg R, Hebblewhite M, Whittington J, Lukacs P, McKelvey K. 2018 Sampling scales

define occupancy and underlying occupancy–abundance relationships in animals. *Ecology* **99**,

172–183. (doi:10.1002/ecy.2054)

27. National Operational Hydrologic Remote Sensing Center. 2004 Snow Data Assimulation

System (SNODAS) Data Products at NSIDC. (doi:https://doi.org/10.7265/N5TB14TC)

28. Cano-Martínez R, Thorsen NH, Hofmeester TR, Odden J, Linnell J, Devineau O, Angoh

SYJ, Odden M. 2024 Bottom-up rather than top-down mechanisms determine mesocarnivore

interactions in Norway. *Ecol. Evol.* **14**, e11064. (doi:10.1002/ece3.11064)

29. Kéry M, Royle JA. 2016 *Distribution, Abundance, and Species Richness in Ecology*.

Elsevier. (doi:10.1016/B978-0-12-801378-6.00001-1)

30. Kéry M, Royle JA. 2021 *Applied Hierarchical Modeling in Ecology: Analysis of*

Distribution, Abundance and Species Richness in R and BUGS. Elsevier. (doi:10.1016/B978-

0-12-809585-0.00001-6)

31. Twining JP, Brazeal JL, Jensen PG, Fuller AK. 2024 Intraguild interactions and abiotic

conditions mediate occupancy of mammalian carnivores: co-occurrence of coyotes–fishers–

martens. *Oikos* **2024**, e10577. (doi:10.1111/oik.10577)

32. Arnold TW. 2010 Uninformative Parameters and Model Selection Using Akaike's

Information Criterion. *J. Wildl. Manag.* **74**, 1175–1178. (doi:10.1111/j.1937-

2817.2010.tb01236.x)

33. Rowcliffe M. 2023 **activity**: Animal Activity Statistics. 1.3.4

(url:10.32614/CRAN.package.activity)

34. Tanwar KS, Sadhu A, Jhala YV. 2021 Camera trap placement for evaluating species

richness, abundance, and activity. *Sci. Rep.* **11**, 23050. (doi:10.1038/s41598-021-02459-w)

35. Ridout MS, Linkie M. 2009 Estimating overlap of daily activity patterns from camera trap

data. *J. Agric. Biol. Environ. Stat.* **14**, 322–337. (doi:10.1198/jabes.2009.08038)

36. Cusack JJ, Dickman AJ, Kalyahe M, Rowcliffe JM, Carbone C, MacDonald DW, Coulson T.

2017 Revealing kleptoparasitic and predatory tendencies in an African mammal community

using camera traps: a comparison of spatiotemporal approaches. *Oikos* **126**, 812–822.

(doi:10.1111/oik.03403)

37. Davis RS, Yarnell RW, Gentle LK, Uzal A, Mgoola WO, Stone EL. 2021 Prey availability

and intraguild competition regulate the spatiotemporal dynamics of a modified large carnivore

guild. *Ecol. Evol.* **11**, 7890–7904. (doi:10.1002/ece3.7620)

38. Anderson CR, Lindzey FG. 2003 Estimating Cougar Predation Rates from GPS Location

Clusters. *J. Wildl. Manag.* **67**, 307. (doi:10.2307/3802772)

39. Perrig PL, Lambertucci SA, Donadio E, Smith JA, Middleton AD, Pauli JN. 2023 Risk

effects cascade up to an obligate scavenger. *Ecology* **104**, e3871. (doi:10.1002/ecy.3871)

40. Smith AB, Squires JR, Bjornlie NL, Holbrook JD. 2023 Divergent or convergent: how do

forest carnivores use time in the Greater Yellowstone Ecosystem? *J. Mammal.* , gyad070.

(doi:10.1093/jmammal/gyad070)

41. Fedriani JM, Fuller TK, Sauvajot RM, York EC. 2000 Competition and intraguild predation

among three sympatric carnivores. *Oecologia* **125**, 258–270. (doi:10.1007/s004420000448)

42. Van Etten KW, Wilson KR, Crabtree RL. 2007 Habitat Use of Red Foxes in Yellowstone

National Park Based on Snow Tracking and Telemetry. *J. Mammal.* **88**, 1498–1507.

(doi:10.1644/07-MAMM-A-076.1)

43. Jensen AJ, Marneweck CJ, Kilgo JC, Jachowski DS. 2022 Coyote diet in North America:

geographic and ecological patterns during range expansion. *Mammal Rev.* **52**, 480–496.

(doi:10.1111/mam.12299)

44. Allen ML, Elbroch LM, Wilmers CC, Wittmer HU. 2014 Trophic Facilitation or Limitation?

Comparative Effects of Pumas and Black Bears on the Scavenger Community. *PLoS ONE* **9**,

e102257. (doi:10.1371/journal.pone.0102257)

45. Berger KM, Gese EM. 2007 Does interference competition with wolves limit the distribution

and abundance of coyotes? *J. Anim. Ecol.* **76**, 1075–1085. (doi:10.1111/j.1365-

2656.2007.01287.x)

46. Merkle JA, Stahler DR, Smith DW. 2009 Interference competition between gray wolves and

coyotes in Yellowstone National Park. *Can. J. Zool.* **87**, 56–63. (doi:10.1139/Z08-136)

47. Sinclair ARE. 2003 Mammal population regulation, keystone processes and ecosystem

dynamics. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **358**, 1729–1740.

(doi:10.1098/rstb.2003.1359)

48. Brice EM, Larsen EJ, MacNulty DR. 2022 Sampling bias exaggerates a textbook example of

a trophic cascade. *Ecol. Lett.* **25**, 177–188. (doi:10.1111/ele.13915)

49. Hobbs NT, Johnston DB, Marshall KN, Wolf EC, Cooper DJ. 2024 Does restoring apex

predators to food webs restore ecosystems? Large carnivores in Y ellowstone as a model

system. *Ecol. Monogr.* **94**, e1598. (doi:10.1002/ecm.1598)

50. Ripple WJ, Beschta RL. 2012 Trophic cascades in Yellowstone: The first 15years after wolf

reintroduction. *Biol. Conserv.* **145**, 205–213. (doi:10.1016/j.biocon.2011.11.005)

551
552

Figure 2 | (a) Apex and mesocarnivore marginal occupancy probabilities estimated from camera
trapping surveys between 2021–2023. **(b)** Coyote occupancy probability for 2021–2023,
conditional on wolf presence-absence and red fox occupancy probability for 2021–2023,
conditional on cougar presence-absence. Occupancy probabilities were based on the best-
supported, second-order multi-species occupancy model.

Figure 3 | Diel activity patterns with 95% bootstrapped CIs for study carnivores. Mean activity
overlap **(a)** between apex and mesocarnivores, and **(b)** between apex carnivores and between
mesocarnivores.

Time of Day (Hours)

Figure 4 | (a) Daily binned temporal attraction-avoidance of mesocarnivores in response to apex carnivore detections at camera stations. Apex-mesocarnivore pairs include wolf-coyote (top-left), wolf-red fox (top-right), cougar-coyote (bottom-left), and cougar-red fox (bottom-right). Points and lines represent observed mesocarnivore detection probabilities, while violin plots show the distribution of expected mesocarnivore detection probabilities based on random movements (i.e., no attraction or avoidance). *denotes significant differences between observed and expected detection probabilities. **(b)** Median time differences of mesocarnivore detections following apex carnivore detections.

576 **Figure 5 |** Wolf (n = 327) and cougar (n = 257) kills, surveyed through GPS-cluster searches of
577 collared wolves and cougars between 2016-2023, bearing **a)** presence of coyote and red fox
578 scavenging expressed as a proportion, and **b)** counts of coyote and red fox killed. For methods
579 see main text. Wolf and elk carcass graphics by Kira Cassidy.

Figure 6 | Conceptual representation of results from analyses for each apex-mesocarnivore pairing. Mesocarnivores (Coyote = blue, Red fox = red) are shown in the center and apex carnivores around the perimeter (Wolf = Green, Cougar = Pink). For "Occupancy/ Broad Space Use," the multi-directional arrow refers to co-occurrence results from the top second-order multi-species occupancy model. "Scavenging" represents the proportion of ungulate kills made by an apex carnivore that were visited by the respective mesocarnivore. "+" represents a positive association/effect, "–" represents a negative association/effect, and "no effect" indicates an insignificant result. Effects are colored green or pink to indicate they are affiliated with wolf or cougar results, respectively. The size of the "+" or "–" is positively-related to the magnitude of the effect observed.

602 **SUPPLEMENTARY INFORMATION FOR:**

604 **Title:** Species-specific interactions with apex carnivores yield unique benefits and burdens for

605 mesocarnivores

606   **Table S1:** Covariate List for Detection and Occupancy Models & their Description and sources
607

Occupancy Covariates		
Variable	**Type, Range**	**Description**
Cover	Continuous	Values accrued from the Rangeland Analysis Platform's (RAP) annual percent tree cover landsat satellite data (Robinson et al., 2019). Originally at 30-m spatial resolution but aggregated to 1 km. We used the cover value associated with the cell that a given camera station lies within.
Topographic Roughness	Continuous	Values represent the sum of the absolute value of the difference in elevation between each grid cell and its surrounding eight neighbors (3 x 3 window) (Kohl et al., 2019; Ruth et al., 2019). A value of 0 represents a flat surface while the maximum value represents the roughest terrain in the study area. Originally at 30-m spatial resolution but aggregated to 1 km. We used the roughness value associated with the cell that a given camera station lies within.
Snow Depth	Continuous	Daily snow depth data at 1-km grid cell accrued from SNODAS. We used the average value of all days within each session, so that snow depth values in a grid varied between but not within sessions.
Year	Categorical	Values represent the three winter sessions of our study (2021, 2022, 2023).
Detection Covariates		
Snow Depth	Continuous	Daily snow depth data at 1-km grid cell accrued from SNODAS. We used the average value of all days within each weekly occasion, so that snow depth values at each station varied weekly.
Number of Cameras	Continuous, Range: 0-2	Number of active/operational cameras at a given station.
Year	Categorical	Values represent the three winter sessions of our study (2021, 2022, 2023).

610   **Table S2 |** Detection and occupancy covariates in the best-supported first-order models for red
611   fox, coyote, wolf, and cougar in each session.
612

Session-Species	Detection model (p)	Top occupancy model (Ψ)
Red Fox	*Year + Snow depth*	*Null (Year only)*
Coyote	*Null (Year only)*	*Null (Year only)*
Wolf	*Null (Year only)*	*Null (Year only)*
Cougar	*Year + Snow depth*	*Year + Snow depth + Roughness*

614 **Table S3 |** AIC model selection for all candidate first-order detection sub-models considered for
615 each species. Bolded rows indicate selected models that were carried over into the occupancy
616 model selection stage. K = number of model parameters, AIC = Akaike Information Criterion,
617 ΔAIC = difference in AIC from the top model.
618

First-order Detection Model	K	AIC	ΔAIC	AIC$_{weight}$
Red Fox				
p(Year + Snow depth) + Ψ(.)	**11**	**7611**	**0**	**0.714**
p(Year + Active cameras + Snow depth) + Ψ(.)	12	7612.9	1.96	0.268
p(Null (Year only)) + Ψ(.)	10	7619	8.01	0.013
p(Year + Active cameras) + Ψ(.)	11	7620.8	9.86	0.005
Coyote				
p(Null (Year only)) + Ψ(.)	**10**	**7612.1**	**0**	**0.42**
p(Year + Active cameras) + Ψ(.)	11	7612.7	0.6	0.31
p(Year + Snow depth) + Ψ(.)	11	7614.1	1.93	0.16
p(Year + Active cameras + Snow depth) + Ψ(.)	12	7614.6	2.51	0.12
Wolf				
p(Null (Year only)) + Ψ(.)	**10**	**7616.6**	**0**	**0.39**
p(Year + Snow depth) + Ψ(.)	11	7617.5	0.91	0.25
p(Year + Active cameras) + Ψ(.)	11	7617.8	1.17	0.22
p(Year + Active cameras + Snow depth) + Ψ(.)	12	7618.7	2.08	0.14
Cougar				
p(Year + Active cameras + Snow depth) + Ψ(.)	12	7624.8	0	0.479
p(Year + Snow depth) + Ψ(.)	**11**	**7625.1**	**0.23**	**0.427**
p(Null (Year only)) + Ψ(.)	10	7629.3	4.49	0.051
p(Year + Active cameras) + Ψ(.)	11	7629.6	4.8	0.043

Table S4 | AIC model selection for all candidate first-order occupancy models considered for
each species. Each model includes the full detection sub-model. Bolded rows indicate selected
models that were carried over into the occupancy model selection stage. K = number of model
parameters, AIC = Akaike Information Criterion, ΔAIC = difference in AIC from the top model.

First-order Detection + Occupancy Model	K	AIC	ΔAIC	AIC$_{weight}$
Red Fox				
p + Ψ(Null (Year only))	**20**	**7588.1**	**0**	**0.222**
p + Ψ(Year + Roughness)	21	7588.6	0.49	0.174
p + Ψ(Year + Snow depth)	21	7588.9	0.75	0.153
p + Ψ(Year + Roughness + Snow depth)	22	7589.2	1.01	0.134
p + Ψ(Year + Cover)	21	7589.6	1.48	0.106
p + Ψ(Year + Roughness + Cover)	22	7590	1.89	0.086
p + Ψ(Year + Snow depth + Cover)	22	7590.6	2.42	0.066
p + Ψ(Year + Roughness + Snow depth + Cover)	23	7590.8	2.66	0.059
Coyote				
p + Ψ(Year + Roughness + Cover)	22	7587.3	0	0.215
p + Ψ(Year + Cover)	21	7587.6	0.24	0.190
p + Ψ(Year + Roughness)	21	7588.2	0.84	0.141
p + Ψ(Null (Year only))	**20**	**7588.7**	**1.38**	**0.108**
p + Ψ(Year + Snow depth + Cover)	22	7588.8	1.41	0.106
p + Ψ(Year + Roughness + Snow depth + Cover)	23	7588.9	1.53	0.100
p + Ψ(Year + Roughness + Snow depth)	22	7589.5	2.19	0.072
p + Ψ(Year + Snow depth)	21	7589.6	2.27	0.069
Wolf				
p + Ψ(Null (Year only))	**20**	**7588.9**	**0**	**0.221**
p + Ψ(Year + Snow depth)	21	7589.4	0.59	0.165
p + Ψ(Year + Roughness + Snow depth)	22	7589.6	0.71	0.155
p + Ψ(Year + Roughness)	21	7589.6	0.74	0.153
p + Ψ(Year + Cover)	21	7590.6	1.71	0.094
p + Ψ(Year + Snow depth + Cover)	22	7590.9	2.08	0.078
p + Ψ(Year + Roughness + Snow depth + Cover)	23	7591.1	2.27	0.071
p + Ψ(Year + Roughness + Cover)	22	7591.4	2.52	0.063
Cougar				
p + Ψ(Year + Roughness + Snow depth + Cover)	23	7542.4	0	0.54

p + Ψ(Year + Roughness + Snow depth)	**22**	**7543.8**	**1.35**	**0.28**
p + Ψ(Year + Roughness + Cover)	22	7545.6	3.15	0.11
p + Ψ(Year + Roughness)	21	7546.6	4.22	0.066
p + Ψ(Year + Snow depth)	21	7578.8	36.39	6.8e-9
p + Ψ(Year + Snow depth + Cover)	22	7579.4	36.97	5.1e-9
p + Ψ(Null (Year only))	20	7586.2	43.79	1.7e-10
p + Ψ(Year + Cover)	21	7587.3	44.86	9.9e-11

626 **Table S5 |** AIC model comparison for all candidate first- and second-order multi-species
627 occupancy models considered. Bolded rows indicate selected models that we reported results for.
628 K = number of model parameters, AIC = Akaike Information Criterion, ΔAIC = difference in
629 AIC from the top model.
630

Model	K	AIC	ΔAIC	AIC$_{weight}$
Second-Order Detection + Occupancy	**34**	**7514**	**0**	**1**
First-Order Detection + Occupancy	28	7545	31	1.9e-7
First Order Detection Only	18	7588	74	8.0e-17
First-Order Null	8	7626	112	6.1e-25

632   **Table S6** | Beta coefficient estimates for detection and occupancy covariates retained in the top
633   second-order multi-species occupancy model for each session.

Detection model (p)						Occupancy model (Ψ)				
[Species] Covariate	β	SE	z	P		[Species] Covariate	β	SE	z	P
[redfox] (Intercept)	-0.71	0.08	-9.5	2.3e-21		[redfox] (Intercept)	0.87	0.73	1.19	0.23
[redfox] year2022	-0.44	0.12	-3.74	0.0002		[redfox] year2022	-0.60	0.79	-0.76	0.44
[redfox] year2023	0.25	0.13	1.85	0.06		[redfox] year2023	-0.67	0.75	-0.90	0.37
[redfox] scale(snowdepth)	-0.21	0.07	-3.13	0.002		[coyote] (Intercept)	-1.01	0.69	-1.47	0.14
[coyote] (Intercept)	-1.12	0.10	-11.3	1.0e-29		[coyote] year2022	-0.38	0.48	-0.79	0.43
[coyote] year2022	-0.68	0.18	-3.79	0.0002		[coyote] year2023	-1.16	0.45	-1.07	0.28
[coyote] year2023	-0.12	0.16	-0.79	0.43		[wolf] (Intercept)	-1.57	0.76	-2.07	0.04
[wolf] (Intercept)	-1.57	0.12	-13.0	1.4e-38		[wolf] year2022	-0.25	0.52	-0.48	0.63
[wolf] year2022	-0.45	0.21	-2.15	0.03		[wolf] year2023	-0.41	0.47	-0.87	0.38
[wolf] year2023	0.27	0.18	1.56	0.12		[cougar] (Intercept)	-1.03	0.80	-1.30	0.19
[cougar] (Intercept)	-1.89	0.15	-12.6	1.1e-36		[cougar] year2022	-0.93	0.68	-1.36	0.17
[cougar] year2022	-0.20	0.21	-0.94	0.35		[cougar] year2023	0.23	0.80	0.29	0.77
[cougar] year2023	0.53	0.27	1.94	0.052		[cougar] scale(snowdepth)	-1.06	0.47	-2.24	0.02
[cougar] scale(snowdepth)	-0.27	0.12	-2.16	0.03		[cougar] scale(roughness)	1.40	0.32	4.38	1.2e-5
						[redfox:coyote] (Intercept)	1.14	0.66	1.72	0.09
						[redfox:wolf] (Intercept)	1.32	0.70	1.89	0.06
						[redfox:cougar] (Intercept)	1.94	0.73	2.67	0.008
						[coyote:wolf] (Intercept)	1.62	0.41	3.94	8.1e-5
						[coyote:cougar] (Intercept)	-0.49	0.44	-1.12	0.26
						[wolf:cougar] (Intercept)	0.06	0.45	0.14	0.89

637   **Figure S1 |** Proportion of camera sites exhibiting each co-occurrence state (*z*) for each meso-
638   apex species pair.

641 **Figure S2 |** Naive occupancy probability (proportion of sites occupied) across all sites for each
642 species.
643

644
645

646   **Figure S3 |** Detection probabilities for each species as a function of significant covariates
647   retained in the top second-order multi-species occupancy model. Shaded regions are 95%
648   confidence intervals.
649

652   **Figure S4 |** Marginal occupancy probability for cougar as a function of habitat covariates
653   retained in the top second-order multi-species occupancy model. Shaded regions are 95%
654   confidence intervals.
655

658   **Figure S5 |** Wolf (n = 113) and cougar (n = 67) kills, surveyed through GPS-cluster searches of
659   collared wolves and cougars during our three camera-trap sessions (2021-2023), bearing **a)**
660   presence of coyote and red fox scavenging expressed as a proportion, and **b**) counts of coyote
661   and red fox killed. For methods see main text. Wolf and elk carcass graphics by Kira Cassidy.
662

663
664

REFERENCES

Kohl, M. T., Ruth, T. K., Metz, M. C., Stahler, D. R., Smith, D. W., White, P. J., & MacNulty,

   D. R. (2019). Do prey select for vacant hunting domains to minimize a multi-predator

   threat? *Ecology Letters*, *22*(11), 1724–1733. https://doi.org/10.1111/ele.13319

Robinson, N. P., Jones, M. O., Moreno, A., Erickson, T. A., Naugle, D. E., & Allred, B. W.

   (2019). Rangeland Productivity Partitioned to Sub-Pixel Plant Functional Types. *Remote

   Sensing*, *11*(12), 1427. https://doi.org/10.3390/rs11121427

Ruth, T., Buotte, P., & Hornocker, M. (2019). *Yellowstone Cougars: Ecology Before And During

   Wolf Restoration*. University Press of Colorado. https://doi.org/10.5876/9781607328292