Entropy production constrains information throughput in gene regulation

Maximilian Gehri,[1] Lukas Stelzl,[2] and Heinz Koeppl[1]

[1)]*Centre for Synthetic Biology, Technical University of Darmstadt, 64283 Darmstadt, Germany*

[2)]*Institute of Molecular Physiology, Johannes Gutenberg University Mainz, 55122 Mainz, Germany*

(*Electronic mail: heinz.koeppl@tu-darmstadt.de)

(Dated: 21 October 2024)

Biological signal processing typically requires energy, leading us to hypothesize that a cell's information processing capacity is constrained by its energy dissipation. Signals and their processing mechanisms are often modeled using Markovian chemical reaction networks (CRNs). To enable rigorous analysis, we review and reformulate stochastic thermodynamics for open CRNs, utilizing Kurtz's process-based formulation. In particular, we revisit the identification of the energy dissipation rate with the entropy production rate (EPR) at the non-equilibrium steady state (NESS). We also highlight potential inconsistencies in traditional formulations for generic Markov processes when applied to open CRNs, which may lead to erroneous conclusions about equilibrium, reversibility, and the EPR. Additionally, we review the concepts of mutual information (MI) and directed information (DI) between continuous-time trajectories of CRNs, which capture the transmission of spatiotemporal patterns. We generalize existing expressions for the MI, originally accounting for transmission between two species, to now include transmission between arbitrary subnetworks. A rigorous derivation of the DI between subnetworks is presented. Based on channel coding theorems for continuous-time channels with feedback, we argue that directed information is the appropriate metric for quantifying information throughput in cellular signal processing. To support our initial hypothesis within the context of gene regulation, we present two case studies involving small promoter models: a two-state nonequilibrium promoter and a three-state promoter featuring two activation levels. We provide analytical expressions of the directed information rate (DIR) and maximize them subject to an upper bound on the EPR. The maximum is shown to increase with the EPR.

I. INTRODUCTION

Energy consumption is essential for gene regulatory mechanisms and intracellular signal transduction in both prokaryotes and eukaryotes. For example, in eukaryotes, chromatin remodeling complexes use ATP to maintain gene expression profiles[1,2]. In prokaryotes, e.g., the conversion of ATP to cAMP provides a cofactor for regulatory transcription factors like the catabolite-activating protein (CAP)[3]. Gene regulatory networks are an essential part of signal processing in cells[4], motivating an information-theoretic analysis of the network and its single regulatory elements like promoter architectures. There is increasing evidence that cells represent information in the spatiotemporal variation of signaling molecules when processing internal or external stimuli[5–7]. The benefit of stochastic chemical reaction networks (CRNs) is that these models account for the discrete nature of reactions and molecule counts. Any static or dynamical representation of information is thus accurately captured in the trajectories of a CRN. In complement, accurate thermodynamic characterization is also possible with CRNs.

We hypothesize that the information throughput in gene regulatory networks is limited by the amount of energy required. Such trade-offs between energy dissipation and precision in cellular decision making and adaptation have been described as an emerging general design principle of biological systems[8]. Therefore, in this paper, we propose the directed information[9] (DI) between continuous-time stochastic processes to quantify the directed information throughput in a CRN. We then use stochastic thermodynamics for CRNs (sTD) to quantify the energy dissipation rate and at the non-equilibrium steady state (NESS) and identify it with the entropy production rate (EPR). Finally, we conduct information-theoretic case studies on minimal models to support our hypothesis. We envision that further developments in this area will help in the design of new information processing genetic circuits that optimally handle the trade-offs.

A. Previous related work

A powerful formalization of the description of Markovian chemical reaction networks was provided by Kurtz and Anderson[10]. They demonstrated how counting processes relevant to these networks can be represented using Poisson-type processes, leading to a random time-change representation. Therewith, they obtain a stochastic equation for continuous-time Markov jump processes, i.e., their trajectory-level description. Our introduction to CRNs largely follows their work, supplemented by some thermodynamically relevant concepts.

Information theory was pioneered by Shannon and laid the foundation for the digital age. One of his landmark contributions was the channel coding theorem[11], which determines the maximum rate at which information can be reliably transmitted despite noise corrupting the received signal. A key aspect of Shannon's work was his abstraction of noise corruption through the concept of a transmission channel, characterized by the conditional distribution of the output given the input. He introduced mutual information, a functional of the joint distribution of input and output, to quantify the stochastic dependency between input and output and thereby the information exchange between them. The channel coding theorem identifies the supremum of the mutual information for a given

transmission channel as the maximum transmission rate at which the error probability asymptotically approaches zero in the limit of infinite transmission duration. This limiting rate is known as the channel capacity. Marko developed a theory of bidirectional communication, highlighting that communication, unlike mere transmission, is intrinsically bidirectional[12]. Building on his ideas, Massey established a consistent theory of information transmission that incorporates the use of feedback by emphasizing the distinction between statistical and causal dependence[13]. Feedback is not an inherent feature of the channel itself; rather, it is integrated into the encoding of the signals transmitted through the channel. In particular, Massey introduced the concept of directed information and demonstrated that, when used in place of mutual information, it accurately extends the channel coding theorem to transmission with the use of feedback. For channels that are used without feedback, directed information reduces to mutual information. Weissman et al. further generalized the traditional notion of directed information between finite-length discrete-time sequences to a notion between continuous-time trajectories[9]. They also proved a channel coding theorem for a broad class of continuous-time channels, used with feedback, and derived the channel capacity via the directed information. Although it is not immediately clear whether the channel between trajectories of subnetworks within a CRN falls into this class, the inherent multidirectionality of causal dependencies in CRNs, coupled with their continuous-time description, suggests that directed information between continuous-time trajectories is the appropriate information-theoretic measure for quantifying directional information throughput in CRNs.

The use of information theory in systems biology is a relatively recent research field[14]. The initial applications focused on analyzing gene regulatory networks by characterizing the mutual information of the joint distribution of network elements at steady-state[15] or at individual time points[16]. Hansen and O'Shea first accounted for dynamic information transduction by considering particular dynamic features as amplitude and frequency[17]. Tostevin and ten Wolde first considered the mutual information between trajectories, recognizing that relevant information may be contained in the temporal dynamics of the input signal[18]. They provide the mutual information rate, i.e., the asymptotic differential information gain, under a linear-noise Gaussian approximation. Lestas et al. used the mutual information between trajectories as a tool to lower bound the cost of suppressing molecular fluctuations of the input species via feedback under a Langevin approximation of the input[19]. Nakahira et al. extended their work on biomolecular control and identified the directed information as the appropriate information measure[20].

A measure-theoretic account for the mutual information between the trajectories of two species of a class of biochemical reaction networks was first given by Duso and Zechner[21] and later generalized by Moor and Zechner[22]. To compute the mutual information they proposed several methods: a combination of stochastic simulation (SSA)[23] and moment closure approximation, a quasi-exact approach combining SSA and numerical integration and an analytical approximation. Moor and Zechner further showed that the trajectory mutual infor-

mation decomposes into a pair of transfer entropies. These transfer entropies are in fact identical to to DI from input to output and vice versa. In this paper, we further generalize the expression for the mutual information in[22] to encompass the information flow between subnetworks. Sinzger et al. and Gehri et al. advanced techniques for analytically and numerically computing the mutual information of the Poisson-type channel, which can be regarded as a minimal module of a CRN[24–26]. In the traditional definition of the Poisson-type channel[27], the mutual information equals directed information[9,26]. Finally, Reinhardt et al. developed an exact Monte Carlo method called path weight sampling to compute the mutual information between trajectories[28], which currently represents the state-of-the-art approach for calculating mutual information in large networks. The algorithm has recently been adapted for the computation of transfer entropies[29].

Schnakenberg first formulated steady-state thermodynamics for Markov jump processes, defined via their infinitesimal generator, and derived a cycle decomposition for the entropy production rate[30]. A first formulation of stochastic thermodynamics for CRNs beyond steady state was done by Schmiedl and Seifert[31]. Rao and Esposito reformulated sTD for CRNs with emphasis on topological properties and conservation laws[32]. By leveraging the CRNs topology, they proposed a new Gibbs-type potential for the CRN, leading to a more informative thermodynamic description. Our review of sTD mainly follows Schmiedl and Rao, without explicitly considering conservation laws or fluctuation theorems. In particular, we leverage the benefits of representing trajectory-wise thermodynamic quantities via formal counting process integrals.

B. Outline

The paper is organized as follows. In Sec. II we introduce the process-based description of microscopically reversible Markovian CRNs as developed by Kurtz and Anderson[10].

The concepts of mutual information and directed information between continuous-time processes are both introduced in a general measure-theoretic way in Sec. III. We then use a classical expression[33] that links the mutual information between jump processes to the characterization of the counting processes of the distinct reactions. To obtain the MI between arbitrary subnetworks of the CRN, we introduce a notion of indistinguishability of different reactions from the perspective of the subnetworks. We also novelly and rigorously derive the directed information between subnetworks. All expressions presented are also valid for reactions with simultaneous changes in input and output species.

The concepts of stochastic thermodynamics for closed and open chemical reaction networks are pedagogically introduced in Sec. IV. In particular, we express thermodynamic characteristics of random trajectories using counting process integrals, a common technique for CRNs. Then, in Sec. IV C, we review the common identification of the EPR of the NESS, as derived from the kinetic model, with the energy dissipa-

tion rate. In Sec. IV E we formulate time-reversed reaction counting processes to express the KL-divergence of the CRNs dynamics from its time-reversal. This divergence is commonly used to quantify the degree of thermodynamic irreversibility. Using these formal tools, we review the relation of irreversibility and entropy production in the context of CRNs. The discussion concludes with a listing of the four equivalent conditions for thermodynamic equilibrium in CRNs. Throughout, we highlight differences and inconsistencies in the traditional formulation of sTD for generic Markov processes, defined via their infinitesimal generator, in the context of CRNs.

Finally, in Sec. V, we showcase the combination of the tools provided throughout the paper and explore how energy dissipation constrains the maximum information throughput in gene regulation, using two minimal promoter models as case studies. The first is a nonequilibrium two-state promoter model with parallel reaction channels, and the second is a three-state promoter model. Both models describe the switching between a single transcriptionally active state and an inactive state. We derive analytical expressions for the directed information (DI) between the promoter's switching activity and the RNA copy number, and we optimize the directed information rate (DIR) under an upper bound on the promoter's entropy production rate.

In the following let $(\Omega, \mathscr{F}, \mathbb{P})$ be a probability space with complete filtration $\mathscr{F} = \{\mathscr{F}_t\}_{t \geq 0}$. With $\int_s^t$ we denote the integration over the interval $(s, t]$, where $s < t$. This convention is important when considering counting process integrals. In the information-theoretic context we refer to X and Y as the channel input and output, respectively. In the notation of the quantities in stochastic thermodynamics we proximally follow the IUPAC conventions[34].

II. MARKOVIAN STOCHASTIC CHEMICAL REACTION NETWORKS

A stochastic chemical reaction network (CRN) consists of $N \in \mathbb{N}$ chemical species $Z = (Z_1, \ldots, Z_N)$ and $M \in \mathbb{N}$ reaction channels $\mathscr{R}_1, \ldots, \mathscr{R}_M$ with stoichiometric balance equations

$$\mathscr{R}_m : \quad \sum_{n=1}^{N} a_n^{(m)} Z_n \underset{\lambda_m^-}{\overset{\lambda_m^+}{\rightleftarrows}} \sum_{n=1}^{N} b_n^{(m)} Z_n,$$

wherein $a^{(m)}, b^{(m)} \in \mathbb{Z}_{\geq 0}^N$ represent the number of reacting molecules of each species and the number of product molecules, respectively, for the forward reaction, and vice versa for the backward reaction. The difference of product and reactant species of $\mathscr{R}_m$ forms the stoichiometric change vector $v_m = b^{(m)} - a^{(m)}$ of the forward reaction. For each reaction channel, the backward reaction microscopically reverses the forward reaction. We assume that all reactions are elementary[35], i.e., there are no reaction intermediates and each reaction has only a single transition state in the reaction coordinate diagram as illustrated in Fig. 1 (a).

Let $\{Z(t) \in \mathbb{Z}_{\geq 0}^N\}_{t \geq 0}$ be the continuous-time Markov chain of the copy number vector of the chemical species Z.

FIG. 1: Exemplary reaction coordinate diagrams for (a) elementary reactions and (b) a non-elementary reaction with a single intermediate. E_a denotes the activation energy of the reaction, while $-\Delta g$ is amount of energy released by the reaction.

Throughout, we sometimes assume that the rates of the reactions, commonly referred to as propensity functions, obey stochastic mass action kinetics[10,32]

$$\lambda_m^+(z) = k_m^+ \prod_{n=1}^{N} \binom{z_n}{a_n^{(m)}}, \quad \lambda_m^-(z) = k_m^- \prod_{n=1}^{N} \binom{z_n}{b_n^{(m)}} \quad (1)$$

with $z \in \mathbb{Z}_{\geq 0}^N$ and stochastic rate constants $k_m^+ > 0$ and $k_m^- \geq 0$. A conversion of the discrete molecule numbers to concentrations in a finite solvent volume V gives rise to another pair of rate constants $\kappa_m^+ > 0$, $\kappa_m^- \geq 0$, such that

$$k_m^+ = \kappa_m^+ \frac{V \prod_{n=1}^{N} a_n^{(m)}!}{\prod_{n=1}^{N} V^{a_n^{(m)}}}, \qquad k_m^- = \kappa_m^- \frac{V \prod_{n=1}^{N} b_n^{(m)}!}{\prod_{n=1}^{N} V^{b_n^{(m)}}},$$

due to consistency of physical units in both the deterministic ODE model (of the concentrations) and the probability evolution equation of the stochastic model[36]. κ_m^+ and κ_m^- can be identified with the rate constants of the deterministic law of mass action by taking the macroscopic limit. The mass action assumption can be dropped in the following if it is not explicitly used. A reaction is called microscopically reversible if $k_m^- > 0$, or more generally if $\lambda_m^+(z) > 0$ implies $\lambda_m^-(z + v_m) > 0$ and vice versa for all z. A CRN is called microscopically reversible if all reactions are microscopically reversible. In the following, a CRN is assumed to be microscopically reversible unless otherwise noted.

For each reactions of channel $\mathscr{R}_m$ we denote the counting processes of reactions in the forward and reverse direction as $\{R_m^+(t) \in \mathbb{Z}_{\geq 0}\}_{t \geq 0}$ with $R_m^+(0) = 0$ and $\{R_m^-(t) \in \mathbb{Z}_{\geq 0}\}_{t \geq 0}$ with $R_m^-(0) = 0$, respectively. These can be represented via random time changes

$$R_m^\rho(t) = N_m^\rho \left(\int_0^t \lambda_m^\rho(Z(s)) \, ds \right)$$

of $2M$ independent unit-rate Poisson processes $\{N_m^\rho(t) \in \mathbb{Z}_{\geq 0}\}_{t \geq 0}$ with $N_m^\rho(0) = 0$ for $m \in \{1, \ldots, M\}$ and $\rho \in \{+, -\}$[10]. Then the random state $Z(t)$ of the CRN with random initial value $Z(0)$ obeys

$$Z(t) = Z(0) + \sum_{m=1}^{M} v_m (R_m^+(t) - R_m^-(t)).$$

The probability mass function $p(z,t) = \mathbb{P}(Z(t) = z)$ follows the chemical master equation[31,32,37]

$$\partial_t p(z,t) = \sum_{m=1}^{M} \lambda_m^+(z - \nu_m) p(z - \nu_m, t) \\ + \lambda_m^-(z + \nu_m) p(z + \nu_m, t) \\ - (\lambda_m^+(z) + \lambda_m^-(z)) p(z,t) \tag{2}$$

Before proceeding, we recapitulate the model assumptions made so far, as they are fundamental for stochastic thermodynamics of CRNs. Assuming a Markov process description means that the future dynamics of the mesoscopic system is sufficiently described by the current state and does not depend on the past. In particular, the future dynamics do also not depend on the particular microstate, the system is in, but just on the current mesostate z.

The above assumptions about the system state Z have implications for the environment in which the reaction network is embedded. Since the propensity functions are independent of time and space, the embedding environment must be seen as thermally equilibrated and spatially homogeneous from the perspective of the chemical species[38]. That is, the surrounding of each molecule of Z homogenizes spatially on a much faster time scale than the reactions occur. In addition, the temperature, pressure, and volume of the solvent are constant over time and space. Similarly, the molecular composition of the solvent, which can affect the rate constants (e.g., its acidity), remains constant over time.

According to Gillespie, in the light of collision theory the elementary reaction assumption is plausible up to bimolecular reactions and the dynamics is rigorously shown to follow stochastic mass action kinetics under the above assumptions on the environment[38]. In other cases, in particular when the chemical species represent different conformations of polymers as is common in gene expression models, a more careful justification of the elementary reaction assumption and the Markov assumption may be required. The relevance of the elementary reaction assumption for stochastic thermodynamics of CRNs is discussed at the end of Sec. IV A.

III.   INFORMATION EXCHANGE BETWEEN SUBNETWORKS

The MI between two $\mathscr{F}$-adapted[39] stochastic processes on a finite interval $[0,t]$, $t > 0$ is defined via the Radon-Nikodym derivative[39] between the joint probability measure $\mathbb{P}_{[0,t]}^{XY}$ of input and output trajectories $(X_{[0,t]}, Y_{[0,t]}) = \{X(s), Y(s)\}_{s \in [0,t]}$ and the product of the respective marginal measures $\mathbb{P}_{[0,t]}^{X}$, $\mathbb{P}_{[0,t]}^{Y}$ of $X_{[0,t]} = \{X(s)\}_{s \in [0,t]}$, $Y_{[0,t]} = \{Y(s)\}_{s \in [0,t]}$:

$$\mathbb{I}(X_{[0,t]}; Y_{[0,t]}) := \mathbb{E}\left[\ln \frac{d\mathbb{P}_{[0,t]}^{XY}}{d(\mathbb{P}_{[0,t]}^{X} \otimes \mathbb{P}_{[0,t]}^{Y})}\right]$$

The mutual information rate (MIR) is defined as its asymptotic slope

$$\bar{\mathbb{I}}(X; Y) := \lim_{t \to \infty} \frac{1}{t} \mathbb{I}(X_{[0,t]}; Y_{[0,t]}),$$

with $(X,Y) = \{X(t), Y(t)\}_{t \geq 0}$. In turn, adapting the definition from Weissman et al.[9], the DI on the closed interval $[0,t]$ is defined as

$$\mathbb{I}(X_{[0,t]} \to Y_{[0,t]}) := \inf_{t \in \mathscr{T}(0,t)} \sum_{i=1}^{n(t)} \mathbb{I}(Y_{[t_{i-1}, t_i]}; X_{[0,t_i]} \mid Y_{[0,t_{i-1}]}), \tag{3}$$

where $\mathscr{T}(t_0, t) = \{t = (t_i)_{i \in \{1,\ldots,n\}} : t_0 < t_1 < \cdots < t_n = t, n \in \mathbb{N}\}$ is the set of all finite sequences partitioning the interval $[t_0, t]$ for any $t_0 < t$ and $n(t)$ denotes the length of the sequence t. Consequently, the directed information rate (DIR) is

$$\bar{\mathbb{I}}(X \to Y) := \lim_{t \to \infty} \frac{1}{t} \mathbb{I}(X_{[0,t]} \to Y_{[0,t]}).$$

In Appendix B we show that under certain continuity conditions, the DI obeys

$$\mathbb{I}(X_{[0,t]} \to Y_{[0,t]}) \\ = \mathbb{I}(X_0; Y_0) + \int_0^t \lim_{h \searrow 0} \frac{1}{h} \mathbb{I}(Y_{[s,s+h]}; X_{[0,s+h]} \mid Y_{[0,s)}) \, ds, \tag{4}$$

which is easier to evaluate than the original definition in Equation (3). An expression similar to (4) was previously proposed by Weissman et al.[9], though without a rigorous proof or clearly stated formal requirements.

If the MI is finite, then mutual and directed information are related by the information conservation law[9]

$$\mathbb{I}(X_{[0,t)}; Y_{[0,t)}) = \mathbb{I}(X_{[0,t)} \to Y_{[0,t)}) + \lim_{\delta \searrow 0} \mathbb{I}(Y_{[0,t-\delta)} \to X_{[0,t)}),$$

where $\mathbb{I}(Y_{[0,t-\delta)} \to X_{[0,t)}) := \mathbb{I}(0_{[0,\delta)} Y_{[0,t-\delta)} \to X_{[0,t)}) = \inf_{t \in \mathscr{T}(\delta,t)} \sum_{i=1}^{n(t)} \mathbb{I}(X_{[t_{i-1}, t_i)}; Y_{[0,t_i-\delta)} \mid X_{[0,t_{i-1})})$ is the directed information for a delayed input trajectory with delay $\delta \in (0,t]$.

Let $\mathscr{F}^{XY} = \{\mathscr{F}_t^{XY}\}_{t \geq 0}$ and $\mathscr{F}^Y = \{\mathscr{F}_t^Y\}_{t \geq 0}$ be the internal filtrations of (X,Y), respectively, Y, where $\mathscr{F}_t^{XY} = \sigma((X(s), Y(s)), 0 \leq s \leq t)$ and $\mathscr{F}_t^Y = \sigma(Y(s), 0 \leq s \leq t)$. $\mathscr{F}^X$ be analogous. The evaluation of the MI between jump-like processes has long been linked to a filtering problem[33,40,41]. In the following, we adapt the expression for the MI between priecewise-constant jump processes on continuous state spaces by Boel et al. to suit our specific needs. We then argue by Equation (4) how the MI is decomposed into the two directed information expressions.

A.   Trajectory Information between Subnetworks

To simplify notation, we redefine the set of bidirectional reactions $\{1, \ldots, M\} \cup \{-m : k_m^- > 0\}$, which has cardinality M', as unidirectional reaction channels $\mathscr{R}_1', \ldots, \mathscr{R}_{M'}'$ with stoichiometric balanced equations

$$\mathscr{R}_{m'}' : \quad \sum_{n=1}^{N} \tilde{a}_n^{(m')} Z_n \quad \xrightarrow{\lambda_{m'}'} \quad \sum_{n=1}^{N} \tilde{b}_n^{(m')} Z_n$$

and change vectors $\nu_{m'}' = \tilde{b}^{(m')} - \tilde{a}^{(m')}$. For clarity, we will omit the prime notation in the remainder of this section.

Let $\mathscr{S}^X, \mathscr{S}^Y \subset \{1,\dots,N\}$ be arbitrary sets in the index set of chemical species of the core system. They represent the subsets of species between which we want to characterize information exchange. Hence, we have $X = \{Z_n(s) : n \in \mathscr{S}^X\}_{s\geq 0}$ and $Y = \{Z_n(s) : n \in \mathscr{S}^Y\}_{s\geq 0}$. Let

$$\mathscr{V} := \{v_1,\dots,v_M\} \setminus \{0\} \subset \mathbb{Z}^N$$

be the set of non-zero change vectors and

$$\pi_X : \mathscr{V} \to \mathbb{Z}^{|\mathscr{S}^X|}, \; v \mapsto v_X = (\tilde{b}_n^{(m)} - \tilde{a}_n^{(m)} : n \in \mathscr{S}_X)$$
$$\mathscr{V}_X := \{\pi_X(v) \mid v \in \mathscr{V}\} \setminus \{0\}$$

be the reduction to the coordinates of the subnetwork of X and the corresponding set of non-zero change vectors. For Y we analogously define π_Y and $\mathscr{V}_Y$.

Definition 1. *The reactions $\mathscr{R}_m$ and $\mathscr{R}_{\tilde{m}}$ are called X-indistinguishable if*

$$\pi_X(v_m) = \pi_X(v_{\tilde{m}})$$

and analogously for Y.

As an example, consider the set of species X, Y, A and the X-indistinguishable reactions $\emptyset \to X$, $X \to 2X$, $Y + A \to Y + X$, which all increase the amount of X by one. Definition 1 allow us to account for the indistinguishability of changes in the marginal subnetwork compared to the entire network. This is important for the definition of the reaction counting processes associated with the joint and marginal probability measures $\mathbb{P}_{[0,t]}^{XY}$, $\mathbb{P}_{[0,t]}^{X}$ and $\mathbb{P}_{[0,t]}^{Y}$.

Let $V : \{1,\dots,M\} \to \mathscr{V} \cup \{0\}$, $m \mapsto v_m$ bet the mapping from reaction index to change vector. We now distinguish between reaction counters of changes in X or Y. Set

$$R_s^X(v_X) := \sum_{v \in \pi_X^{-1}(v_X)} R_{V^{-1}(v)}(s),$$
$$R_s^Y(v_Y) := \sum_{v \in \pi_Y^{-1}(v_Y)} R_{V^{-1}(v)}(s)$$

for all $v_X \in \mathscr{V}_X$ and $v_Y \in \mathscr{V}_Y$. In this notation the time dependence is shifted to the lower index and the functional arguments are the respective marginal change vectors. The corresponding propensity functions of subsystem changes are

$$\lambda^X(v_X, z) := \sum_{v \in \pi_X^{-1}(v_X)} \lambda_{V^{-1}(v)}(z),$$
$$\lambda^Y(v_Y, z) := \sum_{v \in \pi_Y^{-1}(v_Y)} \lambda_{V^{-1}(v)}(z)$$

With this notation, we obtain for all $s \geq 0$ the process equations

$$X(s) = X(0) + \sum_{v_X \in \mathscr{V}_X} v_X\, R_s^X(v_X),$$
$$Y(s) = Y(0) + \sum_{v_Y \in \mathscr{V}_Y} v_Y\, R_s^Y(v_Y).$$

To compute the MI, we must represent the evolution of X and Y relative to the histories of the marginalized and joint subsystems, specifically $\mathscr{F}_s^X$, $\mathscr{F}_s^Y$, and $\mathscr{F}_s^{XY}$, rather than relying on a representation that depends on the current state of Z. This is done by changing the propensity functions of the reaction counters $R_s^X(v_X)$, $R_s^Y(v_Y)$ to stochastic processes, that depend on the history of the respective marginal system. Abusing notation, we say that $R_s^X(v_X)$ has the $\mathscr{F}^{XY}$-intensity process[39] $\lambda^X(v_X)$ and the $\mathscr{F}^X$-intensity $\hat{\lambda}^X(v_X)$ with

$$\lambda_s^X(v_X) = \mathbb{E}\left[\lambda^X(v_X, Z(s-)) \mid \mathscr{F}_{s-}^{XY}\right]$$
$$\hat{\lambda}_s^X(v_X) = \mathbb{E}\left[\lambda^X(v_X, Z(s-)) \mid \mathscr{F}_{s-}^{X}\right].$$

Analogously, $R_s^Y(v_Y)$ has the $\mathscr{F}^{XY}$-intensity $\lambda^Y(v_Y)$ and the $\mathscr{F}^Y$-intensity $\hat{\lambda}^Y(v_Y)$ with

$$\lambda_s^Y(v_Y) = \mathbb{E}\left[\lambda^Y(v_Y, Z(s-)) \mid \mathscr{F}_{s-}^{XY}\right]$$
$$\hat{\lambda}_s^Y(v_Y) = \mathbb{E}\left[\lambda^Y(v_Y, Z(s-)) \mid \mathscr{F}_{s-}^{Y}\right].$$

By definition, all jump processes are non-explosive and both X and Y are "fundamental processes" of the kind as presented in §2.13 of Ref. 33. All intensities are left-continuous processes and therefore predictable[33] w.r.t. their particular intensity. Their expected value is finite at all times $s \in [0,t]$. Given these assumptions, the following theorem provides an expression for the mutual information, similar to the previously stated[22].

Theorem 1. *If $\mathbb{I}(X_0; Y_0) < \infty$ and the intensities meet the requirement*

$$\mathbb{E}\left[\sum_{v_X \in \mathscr{V}_X} \int_0^t \ln\left(\lambda_s^X(v_X)/\hat{\lambda}_s^X(v_X)\right) \lambda_s^X(v_X)\mathrm{d}s\right] < \infty,$$

and analogously for Y, then the MI is finite and satisfies

$$\mathbb{I}(X_{[0,t]}; Y_{[0,t]})$$
$$= \mathbb{I}(X_0; Y_0) + \sum_{v_X \in \mathscr{V}_X} \int_0^t \mathbb{E}\left[\phi(\lambda_s^X(v_X)) - \phi(\hat{\lambda}_s^X(v_X))\right] \mathrm{d}s$$
$$+ \sum_{v_Y \in \mathscr{V}_Y} \int_0^t \mathbb{E}\left[\phi(\lambda_s^Y(v_Y)) - \phi(\hat{\lambda}_s^Y(v_Y))\right] \mathrm{d}s,$$

where $\phi(a) := \ln(a)\, \mathbb{1}_{(0,\infty)}(a)$ with $a \geq 0$ and $\mathbb{1}$ denoting the indicator function of the measurable set in its lower index.

Proof. Apply Theorem 4.2 from Ref. 33 with a slight modification, requiring Theorem 4.1 therein. The detailes are outlined in Appendix A. $\square$

The MI naturally decomposes into structurally equivalent parts, which we identify as the directed informations.

Theorem 2. *Under the conditions of Theorem 1, it holds*

$$\mathbb{I}(X_{[0,t]} \to Y_{[0,t]}) = \mathbb{I}(X_0; Y_0)$$
$$+ \sum_{v_Y \in \mathscr{V}_Y} \int_0^t \mathbb{E}\left[\phi(\lambda_s^Y(v_Y)) - \phi(\hat{\lambda}_s^Y(v_Y))\right] ds,$$

$$\mathbb{I}(Y_{[0,t]} \to X_{[0,t]}) = \mathbb{I}(X_0; Y_0)$$
$$+ \sum_{v_X \in \mathscr{V}_X} \int_0^t \mathbb{E}\left[\phi(\lambda_s^X(v_X)) - \phi(\hat{\lambda}_s^X(v_X))\right] ds.$$

$$(5)$$

In particular, the information conservation law is

$$\mathbb{I}(X_{[0,t]}; Y_{[0,t]}) = \mathbb{I}(X_{[0,t]} \to Y_{[0,t]}) + \mathbb{I}(Y_{[0,t]} \to X_{[0,t]}) - \mathbb{I}(X_0; Y_0).$$

A formal proof is given in Appendix C.

Our expressions particularly allow for the cases, where $\mathscr{S}^X \cap \mathscr{S}^Y \neq \emptyset$. Consider, for instance, the case, where $\mathscr{S}^X = \mathscr{S}^Y$. Then, $\mathscr{F}^{XY} = \mathscr{F}^X = \mathscr{F}^Y$ and consequently, $\mathbb{I}(X_{[0,t]}; Y_{[0,t]}) = 0$ as the intensities satisfy $\lambda_s^X(v_X) = \hat{\lambda}_s^X(v_X)$ and $\lambda_s^Y(v_Y) = \hat{\lambda}_s^Y(v_Y)$ for all $v_X \in \mathscr{V}_X$, $v_Y \in \mathscr{V}_Y$ and all $s \geq 0$.

Moreover, our expression accommodates simultaneous changes in X and Y, a scenario previously discussed as complex[21,22]. The challenge arises from the existence of different probability density representations for the joint process (X,Y). For $\mathscr{S}_X \cap \mathscr{S}_Y = \emptyset$ one might intuitively describe the joint process by projecting the change vectors to the coordinates of the joint subsystem, similar to the reduction described above Definition 1. This yields the following stochastic equation for the joint subsystems

$$\begin{pmatrix} X(t) \\ Y(t) \end{pmatrix} = \begin{pmatrix} X(0) \\ Y(0) \end{pmatrix} + \sum_{v_{XY} \in \mathscr{V}_{XY}} v_{XY} R_t^{XY}(v_{XY}).$$

However, the path probability density in this description may differ in the number of reaction counters and thereby in the dimension of trajectory space representation, compared to the product probability density. In the product measure, simultaneous changes in X and Y, for instance, those arising from a conversion $X \to Y$ are treated as individual changes in both X and Y. In contrast, the above representation of the joint system accounts for such a change only once, and not twice. The path probability densities of the joint and the product measure do not match in terms of the dimension of the path-space representations. To address this dimensionality mismatch in our derivation of the MI and DI expressions, we resolve the issue by splitting the single counting process for simultaneous changes into two separate counting processes, one for X and one for Y.

Interestingly, when reducing to marginal change vectors on the subsystems, reactions that simultaneously affect both X and Y can merge with those that affect only X or Y individually. For example, consider the species X, Y, A with the reactions $\emptyset \to X$, $Y \to X$, $A + X \to 2X$, $Y \to \emptyset$, $\emptyset \to Y$. We identify the process X with the dynamics of the X copy number, and analogously for Y. The reactions in the set $\{\emptyset \to X, Y \to X, A + X \to 2X\}$ are X-indistinguishable with $v_X = 1$ and those in $\{Y \to X, Y \to \emptyset\}$ are Y-indistinguishable with $v_Y = -1$. Consequently, $R^X(1)$ and $R^Y(-1)$ exhibit both simultaneous and individual jumps.

IV. STOCHASTIC THERMODYNAMICS

A. Closed Stochastic Chemical Reaction Networks

A chemical reaction network is considered closed if it does not exchange molecules with the external environment and if it does not change the number of molecules of any of the chemical species that make up the solvent. A closed thermodynamic system may exchange heat with its environment. We interpret the solvent as a heat bath which maintains a constant temperature T.

The zeroth law of thermodynamics for chemical reaction networks dictates that a closed, microscopically reversible CRN that accounts for all solute species will always relax to equilibrium[32]. The equilibrium distribution depends on the initial state of the system and differs between stoichiometric compatibility classes[42], the sets of all states that are connected via the reaction channels $1, \ldots, M$. In the following we assume that any initial distribution assigns positive mass only to states of a single stoichiometric compatibility class, since otherwise the equilibrium distribution can become a mixture of the equilibrium distrubutions of individual compatibility classes. Then equilibrium statistical mechanics further requires that the equilibrium distribution of a closed chemical reaction network is (i) a Gibbs-Boltzmann distribution $\pi(z) = \lim_{t \to \infty} \mathbb{P}(Z(t) = z) \propto e^{-\beta g(z)}$, where $g(z)$ is the Gibbs free energy of the state z and $\beta = (k_B T)^{-1}$, and (ii) it satisfies detailed balance

$$\lambda_m^+(z)\pi(z) = \lambda_m^-(z + v_m)\pi(z + v_m) \qquad (6)$$

for all admissible states z and all microscopically reversible reactions $\mathscr{R}_m$[32]. Plugging the Boltzmannn distribution into (6) yields the local detailed balance condition, or thermodynamic consistency relation, for closed CRNs[32]:

$$\ln\left(\frac{\lambda_m^+(z)}{\lambda_m^-(z + v_m)}\right) = -\beta(g(z + v_m) - g(z)) \qquad (7)$$

for all admissible states z and all microscopically reversible reactions $\mathscr{R}_m$. Equation (7) relates kinetic and energetic constants, and therefore it is also satisfied during the transient nonequilibrium and independent of the initial conditions.

However, the consistency relation can be broken if reactions are not elementary. The equilibrium Gibbs free energy $g(z)$ of a state z can also be derived using equilibrium statistical mechanics of ideal dilute solutions[32]. It is crucial to ensure that all solute species are accounted for in such derivations. The fundamental assumption in equilibrium statistical mechanics and stochastic thermodynamics of discrete-state CRNs is that the Gibbs energy of the system Z is sufficiently described by the mesoscopic state z. This implies that fluctuations in the microscopic states constituting the mesoscopic state z have a negligible impact on the Gibbs free energy of the system.

FIG. 2: Visualization of an open CRN, where the core system Z is coupled to a number of chemostat species C. Each horizontal layer can exchange energy with its vertically adjacent layers.

Significant changes in the system's energy occur only instantaneously due to reactions. Any reaction coordinate of a potentially non-elementary reaction that does not decay instantaneously must be accounted for as a species of the CRN. Perturbations of the well-mixed system due to reactions are disregarded by assuming instantaneous diffusion in a finite volume. Consequently, the model assumptions of stochastic thermodynamics align with the kinetic discrete-state Markov process model.

B. Open Chemical Reaction Networks – Coupling to Chemostats

An open CRN allows the exchange of molecules with external reservoirs. In the following, we assume that the core system Z is coupled to L chemostats $C_1, \ldots, C_L$ via the solvent, as illustrated in Figure 2. We assume that the copy number of chemostat molecules in the solvent is sufficiently high to be represented by their real-valued concentrations $c = (c_1, \ldots, c_L)$, which are held constant over time by an external agent. In our case, this external agent is a biological cell that strives to maintain homeostasis. Others have discussed chemostats that vary in time according to a predetermined deterministic protocol[31,32,43]. The coupled system follows the stoichiometric balance equations

$$\mathscr{R}_m: \quad \sum_{l=1}^{L} r_l^{(m)} C_l + \sum_{n=1}^{N} a_n^{(m)} Z_n \quad \overset{\lambda_m^+}{\underset{\lambda_m^-}{\rightleftarrows}} \quad \sum_{l=1}^{L} s_l^{(m)} C_l + \sum_{n=1}^{N} b_n^{(m)} Z_n,$$

where $\gamma_m = s^{(m)} - r^{(m)}$ is the stoichiometric change vector of the chemostat species associated with $\mathscr{R}_m$. Immediately after each reaction, the chemostat is assumed to correct the copy number of all chemostat species. Consequently, under the assumption of stochastic mass action kinetics, the propensity functions of the coupled reactions maintain the form (1) with

the adjustment that the stochastic rate constants now depend on the chemostat concentrations insofar as they incorporate a deterministic law of mass action

$$k_m^+(c) = \kappa_m^+ \frac{V \prod_{n=1}^{N} a_n^{(m)}!}{\prod_{n=1}^{N} V^{a_n^{(m)}}} \prod_{l=1}^{L} c_l^{r_l^{(m)}},$$

$$k_m^-(c) = \kappa_m^- \frac{V \prod_{n=1}^{N} b_n^{(m)}!}{\prod_{n=1}^{N} V^{b_n^{(m)}}} \prod_{l=1}^{L} c_l^{s_l^{(m)}},$$

where the units of the deterministic rate constants κ_m^+, κ_m^- are adapted accordingly. For any Markovian kinetic laws, c is a fixed parameter of the propensity functions. Therefore, the probability evolution is still described by (2). In the case of a time-varying protocol, the propensity functions become explicitly time-dependent via the concentration values, resulting in a time-inhomogeneous Markov jump process. The generalization to stochastic protocols by describing the chemostats via a continuous-state stochastic process is straightforward. However, the feedback from the core system to the chemostat species must be negligible to maintain the validity of the chemostat assumption.

To describe the energetics of each reaction $\mathscr{R}_m$ we denote by μ_l the chemical potential of C_l, i.e., the energy associated with a single molecule of species C_l.

Definition 2. *For each reaction channel $\mathscr{R}_m$ we introduce the forward and backward difference operator Δ_m^+ and Δ_m^-, such that for any function $f \colon \mathbb{Z}_{\geq 0}^N \to \mathbb{R}$ we define*

$$\Delta_m^\pm f \colon \{z \in \mathbb{Z}_{\geq 0}^N \mid z \pm \nu_m \in \mathbb{Z}_{\geq 0}^N\} \to \mathbb{R}$$
$$(\Delta_m^\pm f)(z) = \pm(f(z \pm \nu_m) - f(z)).$$

The definition implies $(\Delta_m^+ f)(z) = -(\Delta_m^- f)(z + \nu_m)$. Starting from state z in forward direction, the change in Gibbs free energy of the state associated with $\mathscr{R}_m$ is given by

$$\Delta_m^+ g(z) = g(z + \nu_m) - g(z) = -\Delta_m^- g(z + \nu_m)$$

and the chemical work $W_{\text{chem}}^{(m)}$ done on the system by the chemostat

$$W_{\text{chem}}^{(\pm m)} = \mp \sum_l (s_l^{(m)} - r_l^{(m)}) \mu_l,$$

where $(-m)$ denotes the reverse reaction. Hence the energy dissipated into the environment per reaction is

$$\Delta_m^\pm Q_r(z) := W_{\text{chem}}^{(\pm m)} - \Delta g^{(\pm m)}(z) \tag{8}$$

where we incorporate the operator into the definition since it satisfies the property $\Delta_m^+ Q_r(z) = -\Delta_m^- Q_r(z + \nu_m)$. Intuitively, the dissipated energy is the chemical work done on the system but not stored in the system's state. Let $0 < \tau_1^{(\pm m)} < \tau_2^{(\pm m)} < \cdots \leq t$ be the random times, when reaction $\mathscr{R}_m$ happens in forward or backward direction. Then the energy dissipated by the random trajectory $Z_{[0,t]} := \{Z(s)\}_{s \in [0,t]}$ of the open CRN

within the time interval $(0,t]$ is

$$q(t) = \sum_{m=1}^{M} \left\{ \left(\sum_{k:0<\tau_k^{(m)}\leq t} \Delta_m^+ Q_r(Z(\tau_k^{(m)}-)) \right) \right.$$
$$\left. + \left(\sum_{k:0<\tau_k^{(-m)}\leq t} \Delta_m^- Q_r(Z(\tau_k^{(-m)}-)) \right) \right\} \quad (9)$$

where $Z(t-) = \lim_{s\uparrow t} Z(s)$ denotes the left limit in time, such that $Z(\tau_k^{(m)}-)$ is the state of the CRN just before the k-th occurrence of reaction $\mathscr{R}_m$ in forward direction[31]. We rewrite (9) via stochastic integrals of the reaction counting processes

$$q(t) = \sum_{m=1}^{M} \left\{ \int_0^t \Delta_m^+ Q_r(Z(s-)) \, \mathrm{d}R_m^+(s) \right.$$
$$\left. + \int_0^t \Delta_m^- Q_r(Z(s-)) \, \mathrm{d}R_m^-(s) \right\}.$$

Thus the mean energy dissipation rate (also referred to as thermal power[34]) $\dot{Q}(t) = \frac{\mathrm{d}}{\mathrm{d}t} \mathbb{E}[q(t)]$ of the open CRN is

$$\dot{Q}(t) = \sum_z \sum_{m=1}^{M} \Delta_m^+ Q_r(z) \left(\lambda_m^+(z) p(z,t) - \lambda_m^-(z+v_m) p(z+v_m,t) \right),$$
$$(10)$$

as we show in the following proof.

Proof of (10). By the use of the Poisson-type process definition in[39] (Page 27) the expectations of the stochastic integrals can be rewritten as expectations of Lebesgue/Riemann integrals

$$\mathbb{E}[q(t)] = \sum_{m=1}^{M} \mathbb{E}\left[\int_0^t \Delta_m^+ Q(Z(s-)) \, \mathrm{d}R_m^+(s) \right.$$
$$\left. + \int_0^t \Delta_m^- Q_r(Z(s-)) \, \mathrm{d}R_m^-(s) \right]$$
$$= \sum_{m=1}^{M} \mathbb{E}\left[\int_0^t \Delta_m^+ Q_r(Z(s)) \lambda_m^+(Z(s)) \, \mathrm{d}s \right.$$
$$\left. + \int_0^t \Delta_m^- Q_r(Z(s)) \lambda_m^-(Z(s)) \, \mathrm{d}s \right],$$

where the expectation of q is taken over the distribution of all trajectories $Z_{[0,t]}$. By exchanging the expectation with the integral via Fubini's theorem and taking the derivative we obtain

$$\dot{Q}(t) = \sum_{m=1}^{M} \left\{ \mathbb{E}[\Delta_m^+ Q_r(Z(t)) \lambda_m^+(Z(t))] \right.$$
$$\left. + \mathbb{E}[\Delta_m^- Q_r(Z(t)) \lambda_m^-(Z(t))] \right\}.$$

Evaluating the expectations yields

$$\dot{Q}(t) = \sum_z \sum_{m=1}^{M} (\Delta_m^+ Q_r(z) \lambda_m^+(z) + \Delta_m^- Q_r(z) \lambda_m^-(z)) p(z,t).$$

Equation (10) then follows by shifting the order of summation of the last term and using the property of Equation (8). $\square$

Defining the forward, the backward, and the net probability fluxes $J_m^+(z,t) = \mathbb{P}(Z(t) = z) \lambda_m^+(z)$, $J_m^-(z,t) = \mathbb{P}(Z(t) = z+v_m) \lambda_m^-(z+v_m)$, and $J_m(z,t) = J_m^+(z,t) - J_m^-(z,t)$, respectively, of reaction $\mathscr{R}_m$, we can rewrite (10) as

$$\dot{Q}(t) = \sum_z \sum_{m=1}^{M} \Delta_m^+ Q_r(z) J_m(z,t).$$

In a similar fashion we represent the chemical work done on the trajectory $Z_{[0,t]}$ by

$$w(t) = \sum_{m=1}^{M} W_{\text{chem}}^{(m)} (R_m^+(t) - R_m^-(t)),$$

which gives rise to the mean rate of chemical work

$$\dot{W}(t) = \frac{\mathrm{d}}{\mathrm{d}t} \mathbb{E}[w(t)] = \sum_z \sum_{m=1}^{M} W_{\text{chem}}^{(m)} J_m(z,t),$$

by following the steps in the proof of (10). $\dot{W}(t)$ can also be understood as the amount of energy required to maintain the chemostat concentrations, i.e., the energy expenditure of the external agent.

In open CRNs the thermodynamic consistency relation

$$\ln\left(\frac{\lambda_m^+(z)}{\lambda_m^-(z+v_m)} \right) = \beta \, \Delta_m^+ Q(z) \quad (11)$$

for all admissible states z and all reversible reactions $\mathscr{R}_m$ generalizes the local detailed balance (7)[32]. To show consistency in the deterministic limit, we use the macroscopic limit of the Gibbs free energy of the state and the propensity functions under the mass action assumption for a large volume V. Let g be the potential of an ideal dilute solution, i.e.,

$$g(V\tilde{z}) \approx V \sum_{n=1}^{N} \int_0^{\tilde{z}_n} (g_n^\circ + \beta^{-1} \ln(y_n)) \, \mathrm{d}y_n$$
$$\Delta_m^+ g(V\tilde{z}) \approx v_m(V^{-1} \nabla_{\tilde{z}}) g(V\tilde{z})$$
$$= \sum_{n=1}^{N} (b_n^{(m)} - a_n^{(m)})(g_n^\circ + \beta^{-1} \ln(\tilde{z}_n))$$
$$\lambda_m^+(V\tilde{z}) \approx \kappa_m^+ V \left(\prod_{l=1}^{L} c_l^{r_l^{(m)}} \right) \left(\prod_{n=1}^{N} \tilde{z}_n^{a_n^{(m)}} \right),$$
$$\lambda_m^-(V\tilde{z}) \approx \kappa_m^- V \left(\prod_{l=1}^{L} c_l^{s_l^{(m)}} \right) \left(\prod_{n=1}^{N} \tilde{z}_n^{b_n^{(m)}} \right),$$

where g_n° is the standard Gibbs free energy of species n, $\tilde{z} = V^{-1}z$, and the difference operator Δ_m^+ be comes a differential operator with $\nabla_{\tilde{z}}$ denoting the gradient operator w.r.t. the components of $\tilde{z}$. For g we assumed independence of the chemical potentials of distinct species, i.e., . Further we identify $\mu_l = \mu_l^\circ + \beta^{-1} \ln(c_l)$, with standard potential energy μ_l°, and observe that $-W_{\text{chem}}^{(\pm m)}$ precisely represents the change in Gibbs free energy of the chemostat species in the macroscopic limit. Thus, if the species $C_1, \ldots, C_L$ and the core system were to form a closed system, the macroscopic limit of $-\Delta_m^+ Q(z)$

would exactly correspond to the change in macroscopic Gibbs free energy of the total system upon the occurrence of reaction $\mathscr{R}_m$ in forward direction. So in the macroscopic limit the thermodynamic consistency relation for open CRNs (11), i.e.,

$$\ln\left(\frac{\lambda_m^+(V\tilde{z})}{\lambda_m^-(V\tilde{z})}\right) = \beta\left(\sum_l (s_l^{(m)} - r_l^{(m)})\mu_l - \Delta_m^+ g(V\tilde{z})\right), \quad (12)$$

is consistent with (7). Substituting the definitions of $\Delta_m^+ g(V\tilde{z})$ and μ_l in (12) yields the well known relation for the macroscopic rate constants[34]

$$\begin{aligned}\ln(K_{\text{eq}}^{(m)}) &= \ln\left(\frac{\kappa_m^+}{\kappa_m^-}\right)\\ &= -\sum_{l=1}^{L}(s_l^{(m)} - r_l^{(m)})\beta\mu_l^{\circ} \quad (13)\\ &\quad - \sum_{n=1}^{N}(b_n^{(m)} - a_n^{(m)})\beta g_n^{\circ},\end{aligned}$$

where $K_{\text{eq}}^{(m)}$ is the equilibrium constant of reaction $\mathscr{R}_m$. Consistency in the mesoscopic limit can be established by dropping the chemostat assumptions on the chemostat species and noting that the sum of the potential of the chemostat species and g represents the mesoscopic Gibbs free energy of the total system. This total Gibbs free energy and the respective mesoscopic rates then satisfy Equation (7).

C. Relation of Entropy Production Rate and Energy Dissipation

Plugging (11) into (10) reexpresses the mean energy dissipation rate (in units of $k_{\text{B}}T$)

$$\beta\dot{Q}(t) = \sum_z \sum_{m=1}^{M} \ln\left(\frac{\lambda_m^+(z)}{\lambda_m^-(z+v_m)}\right)J_m(z,t) \quad (14)$$

based only on the kinetic parameters of the open CRN. Now (14) is further identified as the (Shannon) entropy flow into the environment[31,32] $\frac{d}{dt}H^{\text{env}}(t) = \beta\dot{Q}(t)$. The mean entropy production $H^{\text{tot}}(t)$ of the whole system, consisting of the core system and its environment, during the interval $[0,t]$ is given as

$$H^{\text{tot}}(t) := \int_0^t \left(\frac{d}{ds}H[Z(s)] + \frac{d}{ds}H^{\text{env}}(s)\right)ds, \quad (15)$$

where $H[Z(t)] = -\sum_z p(z,t)\ln(p(z,t))$ is the (Shannon) entropy of the core system at time $t \geq 0$. Using the CME (2) we express the derivative of the entropy of the core system as

$$\begin{aligned}\frac{d}{dt}H[Z(t)] &= -\sum_z (\ln(p(z,t)) + 1)\,\partial_t p(z,t)\\ &= \sum_z \sum_{m=1}^{M} \ln\left(\frac{p(z,t)}{p(z+v_m,t)}\right)J_m(z,t).\end{aligned}$$

Thus the entropy production rate is given by

$$\frac{d}{dt}H^{\text{tot}}(t) = \sum_z \sum_{m=1}^{M} \ln\left(\frac{J_m^+(z,t)}{J_m^-(z,t)}\right)J_m(z,t). \quad (16)$$

At stationarity, we have $\lim_{t\to\infty} H[Z(t)] = -\sum_z \pi(z)\ln(\pi(z))$, implying $\lim_{t\to\infty}\frac{d}{dt}H[Z(t)] = 0$. Consequently,

$$e_p := \lim_{t\to\infty}\frac{d}{dt}H^{\text{tot}}(t) = \lim_{t\to\infty}\frac{d}{dt}H^{\text{env}}(t) = \lim_{t\to\infty}\beta\dot{Q}(t),$$

i.e., the entropy production rate in the NESS quantifies the mean energy dissipation rate of the core chemical reaction network into the environment.

Thus far, we have provided the mean energy dissipation rate for an unspecified coupling to chemostat species and unspecified chemostat concentrations, based solely on the knowledge of the stochastic rate constants and the change vectors v_m of the core system. Since the deterministic rate constants are constrained by the change vectors γ_m and the concentrations c of the chemostat species, as described by Equation (13), the stochastic rate constants must also satisfy the relation

$$\frac{k_m^+}{k_m^-} = K_{\text{eq}}^{(m)}\left(\prod_{n=1}^{N} V^{b_n^{(m)} - a_n^{(m)}}\right)\left(\prod_{l=1}^{L} c^{r_l^{(m)} - s_l^{(m)}}\right).$$

D. Entropy Production Rate of a Finite Conversion Networks

Much of the literature on the entropy production rate uses a description based on the infinitesimal generator matrix Λ of a CTMC instead of a reaction network description[30,31,37]. A connection between the two pictures is established by using a one-hot coding of the reaction network, which then describes a set of microscopically reversible conversion reactions.

Let Z be N-dimensional with a finite state space $\mathscr{Z} = \{e_1, \dots, e_N\} \subset \{0,1\}^N$, where e_i is the unit vector with a one at position i, and the generator matrix $\Lambda \in \mathbb{R}_{\geq 0}^{N \times N}$. Further, the set of reaction channels be $\mathscr{R} := \{(i,j) \in \{1,\dots,N\}^2 : j > i, \Lambda_{ij} > 0\}$ with associated stoichiometric change vectors $v_{(i,j)} = e_j - e_i$. For $i < j$ we define the reversible conversion reaction

$$\begin{aligned}\mathscr{R}_{(i,j)}: \quad & Z_i \underset{\lambda_{(i,j)}^-}{\overset{\lambda_{(i,j)}^+}{\rightleftarrows}} Z_j \quad \text{with}\\ & \lambda_{(i,j)}^+(z) = \Lambda_{ij}\,\mathbf{1}(z = e_i),\\ & \lambda_{(i,j)}^-(z) = \Lambda_{ji}\,\mathbf{1}(z = e_j).\end{aligned} \quad (17)$$

In the context of an open CRN, this definition implies that each reaction $\mathscr{R}_{(i,j)}$ is associated with a unique change vector of chemostat species. By (16) we obtain

$$\begin{aligned}\frac{d}{dt}H^{\text{tot}}(t) = \sum_{i=1}^{N}\sum_{(k,j)\in\mathscr{R}} &\ln\left(\frac{p(e_i,t)\lambda_{(k,j)}^+(e_k)}{p(e_i + e_j - e_k, t)\lambda_{(k,j)}^-(e_j)}\right)\\ &\times(\lambda_{(k,j)}^+(e_k)p(e_i,t) - \lambda_{(k,j)}^-(e_j)p(e_i + e_j - e_k, t))\end{aligned}$$

and hence substituting (17) yields

$$\frac{d}{dt}H^{\mathrm{tot}}(t) = \sum_{i=1}^{N} \sum_{j:\, j>i} \ln\left(\frac{p(e_i,t)\Lambda_{ij}}{p(e_j,t)\Lambda_{ji}}\right) \times (\Lambda_{ij}\, p(e_i,t) - \Lambda_{ji}\, p(e_j,t)).$$

(18)

We emphasize that the description of the system via the generator matrix is less general than the reaction-based description. In particular, the traditional notion (18) of the entropy production rate is inconsistent with the case, where some reactions channels are parallel.

Definition 3. *Two reaction channels $\mathscr{R}_{m_1}$ and $\mathscr{R}_{m_2}$ are called parallel, if $v_{m_1} = v_{m_2}$ and $\gamma_{m_1} \neq \gamma_{m_2}$, or $v_{m_1} = -v_{m_2}$ and $\gamma_{m_1} \neq -\gamma_{m_2}$.*

The generator matrix merges the rates of the two reactions, as they are indistinguishable from the perspective of the core network only. For example, any two-state Markov process satisfies detailed balance w.r.t. the rates in the generator matrix, i.e., $\ln(\pi(1)\Lambda_{12}/\pi(2)\Lambda_{21}) = 0$ and $e_p = 0$ in the notion of (18). In Sec. V A, however, we describe a two-state Markov process exhibiting a NESS via the reaction-based description.

E. Relation of entropy production and time reversibility

Thus far we have considered the mean entropy production $H^{\mathrm{tot}}(t)$ by the whole system during the interval $[0,t]$ as defined in (15). There is, however, also a trajectory-wise notion of the entropy production. Noting that the entropy of the core system at time $t \geq 0$ can be written as $H[Z(t)] = \mathbb{E}[-\ln(p(Z(t),t))]$ we regard $-\ln(p(Z(t),t))$ as the entropy of Z at time $t \geq 0$. Consequently, the entropy change of the core system between initial and final state of the random trajectory $Z_{[0,t]}$ is

$$h^{\mathrm{sys}}(t) := \ln\left(\frac{p(Z(0),0)}{p(Z(t),t)}\right).$$

(19)

For later convenience we rewrite (19) via integrals. Since $-\ln(p(Z(t),t))$ evolves continuously during intervals, where Z is constant, and has stochastic updates upon reactions, we obtain

$$h^{\mathrm{sys}}(t) = -\int_0^t \partial_s \ln(p(Z(s),s))\, ds$$
$$+ \sum_{m=1}^{M} \int_0^t \ln\left(\frac{p(Z(s-),s)}{p(Z(s-)+v_m,s)}\right) dR_m^+(s)$$
$$+ \sum_{m=1}^{M} \int_0^t \ln\left(\frac{p(Z(s-),s)}{p(Z(s-)-v_m,s)}\right) dR_m^-(s),$$

where by the use of the CME (2) we may substitute

$$\int_0^t \partial_s \ln(p(Z(s),s))\, ds$$
$$= \sum_{m=1}^{M} \int_0^t \left\{ \lambda_m^+(Z(s)-v_m)\frac{p(Z(s)-v_m,s)}{p(Z(s),s)} \right.$$
$$- (\lambda_m^+(Z(s)) + \lambda_m^-(Z(s)))$$
$$\left. + \lambda_m^-(Z(s)+v_m)\frac{p(Z(s)+v_m,s)}{p(Z(s),s)} \right\} ds.$$

We further have $\mathbb{E}\left[\int_0^t \partial_s \ln(p(Z(s),s))\, ds\right] = 0$ as we show later in this Section. The trajectory-wise entropy flow into the environment during $[0,t]$ is defined as $h^{\mathrm{env}}(t) = \beta q(t)$ and substituting with (11), such that

$$h^{\mathrm{env}}(t) := \sum_{m=1}^{M} \left\{ \int_0^t \ln\left(\frac{\lambda_m^+(Z(s-))}{\lambda_m^-(Z(s-)+v_m)}\right) dR_m^+(s) \right.$$
$$\left. + \int_0^t \ln\left(\frac{\lambda_m^-(Z(s-))}{\lambda_m^+(Z(s-)-v_m)}\right) dR_m^-(s) \right\}.$$

Finally, the total entropy production of the trajectory $Z_{[0,t]}$ is

$$h^{\mathrm{tot}}(t) := h^{\mathrm{sys}}(t) + h^{\mathrm{env}}(t)$$

and following the steps in the proof of (10) yields

$$H^{\mathrm{tot}}(t) = \mathbb{E}[h^{\mathrm{tot}}(t)].$$

We now turn to the discussion of time-reversibility. Consider an interval $[0,t]$ and define the time reversed reaction counters $\tilde{R}_m^+(\tau) := R_m^-(t) - R_m^-(t-\tau)$ and $\tilde{R}_m^-(\tau) := R_m^+(t) - R_m^+(t-\tau)$ for all $\tau \in [0,t]$. The time reversed process $\tilde{Z}(\tau) = Z(t-\tau)$ then obeys

$$\tilde{Z}(\tau) = \tilde{Z}(0) + \sum_{m=1}^{M} v_m(\tilde{R}_m^+(\tau) - \tilde{R}_m^-(\tau))$$

for all $\tau \in [0,t]$. To derive the propensity functions for the time-reversed counting processes, we employ a useful trick by leveraging mesoscopic consistency. If we consider the entire system as a closed CRN and treat the chemostat assumption as an approximation, the combined CRN (Z,C), without the chemostat approximation, would be a Markov jump process devoid of parallel reaction channels. In this scenario, it is well known that the propensity functions for the time-reversed CRN are given by

$$\tilde{\lambda}_m^\pm(z,c,\tau)$$
$$= \lim_{h \searrow 0} \frac{1}{h}\mathbb{P}(\tilde{Z}(\tau+h) = z \pm v_m, \tilde{C}(\tau+h) = c \pm \frac{\gamma_m}{V} \mid \ldots$$
$$\ldots \tilde{Z}(\tau) = z, \tilde{C}(\tau) = c)$$
$$= \lim_{h \searrow 0} \frac{1}{h}\mathbb{P}(Z(t-\tau) = z, C(t-\tau) = c \mid \ldots$$
$$\ldots Z(t-\tau-h) = z \pm v_m, C(t-\tau-h) = c \pm \frac{\gamma_m}{V})$$
$$\times \frac{\mathbb{P}(Z(t-\tau-h) = z \pm v_m, C(t-\tau-h) = c \pm \frac{\gamma_m}{V})}{\mathbb{P}(Z(t-\tau) = z, C(t-\tau) = c)}$$
$$= \lambda_m^\mp(z \pm v_m, c \pm \frac{\gamma_m}{V}) \frac{\mathbb{P}(\tilde{Z}(\tau) = z \pm v_m, \tilde{C}(\tau) = c \pm \frac{\gamma_m}{V})}{\mathbb{P}(\tilde{Z}(\tau) = z, \tilde{C}(\tau) = c)}.$$

Reestablishing the chemostat assumptions then yields

$$\tilde{\lambda}_m^\pm(z,\tau) = \lambda^\mp(z \pm v_m) \frac{\mathbb{P}(\tilde{Z}(\tau) = z \pm v_m)}{\mathbb{P}(\tilde{Z}(\tau) = z)}.$$

We now turn to the concepts of reversibility.

Definition 4. *A stationary Markov process Z is called reversible, if $Z_{[0,t]} \overset{d}{=} \tilde{Z}_{[0,t]}$ (equality in distribution) for all $t \geq 0$.*

However, this definition is not entirely consistent with thermodynamics in the presence of parallel reaction channels, as the trajectories of Z lack information about which specific channel was utilized. It can therefore not account for the reversal of the interactions with a chemostat as we exemplify in Sec. V A. Therefore we generalize the definition.

Definition 5. *A stationary Markov CRN Z is called thermodynamically reversible, if $X_{[0,t]} \overset{d}{=} \tilde{X}_{[0,t]}$ (equality in distribution) for all $t \geq 0$, where*

$$X_{[0,t]} := \left\{ Z(0), (R_m^+(s), R_m^-(s)): m \in \{1, \ldots, M\} \right\}_{s \in [0,t]},$$
$$\tilde{X}_{[0,t]} := \left\{ \tilde{Z}(0), (\tilde{R}_m^+(s), \tilde{R}_m^-(s)): m \in \{1, \ldots, M\} \right\}_{s \in [0,t]}.$$

Since the KL-divergence quantifies the divergence of distributions and vanishes in the case of equality, it can be regarded as a quantification of the degree of irreversibility. It holds that

$$\mathbb{D}_{\mathrm{KL}}(X_{[0,t]} \| \tilde{X}_{[0,t]}) = 0$$

for all $t \geq 0$ if and only if Z is thermodynamically reversible. The KL-divergence between counting processes is defined via the corresponding Radon-Nikodym derivative[44] of the forward process w.r.t. the time reversed process, such that

$$\mathbb{D}_{\mathrm{KL}}(X_{[0,t]} \| \tilde{X}_{[0,t]})$$
$$= \mathbb{D}_{\mathrm{KL}}(Z(0) \| \tilde{Z}(0))$$
$$+ \mathbb{D}_{\mathrm{KL}}\left((R_m^+(s), R_m^-(s))_{s \in [0,t], m \in \mathscr{R}} \| (\tilde{R}_m^+(s), \tilde{R}_m^-(s))_{s \in [0,t], m \in \mathscr{R}}\right),$$

where $\mathbb{D}_{\mathrm{KL}}(Z(0) \| \tilde{Z}(0)) = \sum_z p(z,0) \ln\left(\frac{p(z,0)}{p(z,t)}\right)$ and

$$\mathbb{D}_{\mathrm{KL}}\left((R_m^+(s), R_m^-(s))_{s \in [0,t], m \in \mathscr{R}} \| (\tilde{R}_m^+(s), \tilde{R}_m^-(s))_{s \in [0,t], m \in \mathscr{R}}\right)$$
$$= \sum_{m=1}^{M} \mathbb{E}\left[\int_0^t \tilde{\lambda}_m^+(Z(s), s) + \tilde{\lambda}_m^-(Z(s), s) \right.$$
$$\left. - \lambda_m^+(Z(s)) - \lambda_m^-(Z(s)) \, ds \right]$$
$$+ \sum_{m=1}^{M} \mathbb{E}\left[\int_0^t \ln\left(\frac{\lambda_m^+(Z(s-))}{\tilde{\lambda}_m^+(Z(s-), s)} \right) dR_m^+(s) \right]$$
$$+ \sum_{m=1}^{M} \mathbb{E}\left[\int_0^t \ln\left(\frac{\lambda_m^-(Z(s-))}{\tilde{\lambda}_m^-(Z(s-), s)} \right) dR_m^-(s) \right]$$

Now note that

$$\int_0^t \partial_s \ln(p(Z(s), s)) \, ds$$
$$= \sum_{m=1}^{M} \int_0^t \tilde{\lambda}_m^+(Z(s), s) + \tilde{\lambda}_m^-(Z(s), s) - \lambda_m^+(Z(s)) - \lambda_m^-(Z(s)) \, ds,$$

but also

$$\mathbb{E}\left[\int_0^t \tilde{\lambda}_m^+(Z(s), s) + \tilde{\lambda}_m^-(Z(s), s) - \lambda_m^+(Z(s)) - \lambda_m^-(Z(s)) \, ds \right]$$
$$= \mathbb{E}\left[\tilde{R}_m^+(t) + \tilde{R}_m^-(t) - R_m^+(t) - R_m^+(t) \right] = 0,$$

which proves $\mathbb{E}\left[\int_0^t \partial_s \ln(p(Z(s), s)) \, ds \right] = 0$. Finally, we obtain

$$H^{\mathrm{tot}}(t) = \mathbb{E}[h^{\mathrm{tot}}(t)]$$
$$= \mathbb{D}_{\mathrm{KL}}\left((R_m^+(s), R_m^-(s))_{s \in [0,t], m \in \mathscr{R}} \| (\tilde{R}_m^+(s), \tilde{R}_m^-(s))_{s \in [0,t], m \in \mathscr{R}}\right)$$

by putting everything together and following the steps in the proof of (10). By this relation, some authors[45] have proposed a measure-theoretic definition of the entropy production rate, which can now be generalized for open CRNs.

Definition 6.

$$e_p(t) := \lim_{h \searrow 0} \frac{1}{h} \mathbb{D}_{KL}(X_{[t,t+h]} \| \tilde{X}_{[t,t+h]})$$

is called instantaneous entropy production rate of a CRN and

$$e_p := \lim_{t \to \infty} \frac{1}{t} \mathbb{D}_{KL}(X_{[0,t]} \| \tilde{X}_{[0,t]})$$

is called limiting entropy production rate of a CRN.

It is straight forward to show that these measure-theoretic definitions are consistent with the previously mentioned notions of the entropy production rate. Since $\mathbb{D}_{\mathrm{KL}}(X_{[t,t+h]} \| \tilde{X}_{[t,t+h]}) \geq 0$ for all $t, h \geq 0$, we can use the measure-theoretic definition to obtain the second law for the thermodynamics of open CRNs, i.e.,

$$e_p(t), e_p \geq 0.$$

We conclude by reviewing equivalent conditions for thermodynamic equilibrium of non-explosive open CRNs.

Theorem 3. *Let Z be stationary with stationary distribution $\pi(z)$. Then the following conditions are equivalent.*

(a) *Z is thermodynamically reversible. In particular, $\mathbb{D}_{KL}(X_{[0,t]} \| \tilde{X}_{[0,t]}) = 0$ for all $t \geq 0$ and the time reversed propensities satisfy $\tilde{\lambda}_m^{\pm}(z,t) = \lambda_m^{\pm}(z)$.*

(b) *$e_p = 0$*

(c) *Stoichiometric Kolmogorov (Wegscheider) cycle condition: Any admissible closed cycle of states*

$$z_0 \xrightarrow{\rho_1 m_1} z_1 \xrightarrow{\rho_2 m_2} \ldots \xrightarrow{\rho_j m_j} z_n \xrightarrow{\rho_{n+1} m_{n+1}} z_0,$$

originating from a sequence of reaction channel uses m_j in the directions $\rho_j \in \{+, -\}$, such that $\rho_j v_{m_j} = (z_j - z_{j-1})$, satisfies

$$\frac{\lambda_{m_1}^{\rho_1}(z_0) \lambda_{m_2}^{\rho_2}(z_1) \cdots \lambda_{m_{n+1}}^{\rho_{n+1}}(z_n)}{\lambda_{m_{n+1}}^{-\rho_{n+1}}(z_0) \lambda_{m_n}^{-\rho_n}(z_n) \cdots \lambda_{m_1}^{-\rho_1}(z_1)} = 1.$$

FIG. 3: State transition diagram of a two-state promoter model with two microscopically reversible reaction channels. The first reaction channel involves ATP hydrolysis and is therefore coupled to the chemostats of ATP, ADP and P_i.

FIG. 4: State transition diagram of a three-state promoter model with three microscopically reversible reaction channels. The upper path from OFF to ON involves ATP hydrolysis. Hence, OFF* is the intermediate of a non-elementary hydrolysis reaction.

(d) Z satisfies the detailed balance condition

$$\lambda_m^+(z)\pi(z) = \lambda_m^-(z+\nu_m)\pi(z+\nu_m)$$

for all z and all $m \in \mathscr{R}$.

The stoichiometric cycle condition, when combined with the thermodynamic consistency relation, is analogous to the path-independence property of conservative force fields. This relation essentially verifies whether the forces acting on the system originate from a potential or contain nonconservative component.

A proof for Theorem 3 is given in Appendix D. Note that the limiting distribution is not necessarily unique and that the above equivalent equilibrium conditions are still satisfied if the stationary distribution $\pi(z)$ is a mixture of equilibrium distributions on individual stoichiometric compatibility classes.

V. CASE STUDIES

The initial hypothesis was that a cell's information processing capacity is constrained by its energy dissipation. In Section IV C we demonstrated that the energy dissipation rate at the NESS equals the entropy production rate. Here, using two minimal nonequilibrium promoter models, we show that the maximum directed information rate (DIR) of the channel from the promoter's transcriptional activity X to the RNA copy number Y increases as the upper bound on the entropy production rate e_p is raised. This finding supports our initial hypothesis within the context of gene regulation.

A. Models

Consider a two-state Markov jump process with states OFF and ON, which modulate transcriptional activity by switching between the transcription rates 0 and k_{tx}. For simplicity, we set $k_{tx} = 1$ in our analysis, thereby fixing the system's time scale. The state transition diagram of this model, along with transition rates and thermodynamic assumptions, is shown in Fig. 3. The model features a pair of parallel reaction channels: the first channel involves ATP hydrolysis with $\Delta\mu = \mu_{ADP+P_i} - \mu_{ATP}$ and is coupled to corresponding chemostats, while the second reaction channel is thermally driven and not coupled to a chemostat. We denote the Gibbs free energy difference between the states by $\Delta g = g(ON) - g(OFF)$. The entropy production rate at the NESS is

$$e_p = J \ln\left(\frac{k_1^+ k_2^-}{k_2^+ k_1^-}\right) \tag{20}$$

with net probability flux $J = \frac{k_1^+ k_2^- - k_2^+ k_1^-}{k_1^+ + k_1^- + k_2^+ + k_2^-}$, as we show in appendix E.

The three-state promoter model in Figure 4 extends the two-state model by adding an intermediate state, OFF*, which represents a step in a non-elementary reaction involving ATP hydrolysis. In this model, the first reaction channel handles ATP binding and unbinding between OFF and OFF*, while the second channel accounts for the actual ATP hydrolysis, driving the transition from OFF* to ON. The third reaction allows direct transition between OFF and ON and is not coupled to any chemostat species. Thermodynamically, the three-state promoter is characterized by

$$\Delta Q^{(3)} = -\mu_{ADP+P_i} - \Delta g^{(3)} = \frac{1}{\beta}\ln\frac{k_3^+}{k_3^-},$$

$$\Delta g := \Delta g^{(2)} = \Delta g^{(1)} + \Delta g^{(3)} \text{ and}$$

$$\Delta Q^{(-2)} = \Delta g = \frac{1}{\beta}\ln\frac{k_2^-}{k_2^+}.$$

The EPR satisfies

$$e_p = J \ln \left(\frac{k_1^+ k_3^+ k_2^-}{k_1^- k_3^- k_2^+} \right)$$

with $J = k_2^- \pi(ON) - k_2^+ \pi(OFF)$, which is related to the general form of the EPR for circular state transition diagrams[37].

B. Maximum information under energy constraints

We now formulate the maximization problem and refer to the maximum DIR as the (information) capacity. For the two-state promoter we compute

$$C = \sup_{\substack{0 \leq \bar{k}^+, \bar{k}^- \\ 0 \leq e_p(\bar{k}^+, \bar{k}^-) \leq r}} \bar{\mathbb{I}}(X \to Y) \tag{21}$$

where $\bar{k}^+ = k_1^+ + k_2^+$, $\bar{k}^- = k_1^- + k_2^-$ are the effective rates of the transitions between OFF and ON.

The physical boundary conditions $-\Delta\mu, k_1^+, k_2^- > 0$ constrain the admissible region of the effective rates:

$$\bar{k}^+ > c_2 e^{-\beta \Delta g}, \quad c_2 > \bar{k}^+ e^{\beta(\Delta g + \Delta\mu)}$$

For this admissible region the EPR is thus expressed in terms of the effective rates via

$$e_p(\bar{k}^+, \bar{k}^-) = -\beta\Delta\mu \frac{\left(\bar{k}^+ - \bar{k}^- e^{-\beta\Delta g}\right)\left(\bar{k}^- e^{-\beta\Delta\mu} - \bar{k}^+ e^{-\beta\Delta g}\right)}{\left(\bar{k}^+ + \bar{k}^-\right)\left(e^{-\beta\Delta\mu} - 1\right)}, \tag{22}$$

as we show in appendix E.

The DIR for the model with a two-state promoter has been shown to satisfy[26]

$$\begin{aligned}
\bar{\mathbb{I}}(X \to Y) &= \frac{1}{\mathsf{E}[\tau]}(\ln(k_{tx}) + h(\tau) - 1) \\
&= k_{tx} \pi(ON)(\ln(k_{tx}) + h(\tau) - 1) \\
&= k_{tx} \frac{h(\tilde{\tau}) - 1}{\mathsf{E}[\tilde{\tau}]},
\end{aligned}$$

where $h(\tau)$ is the differential entropy of the time τ between transcription events in the dynamics of Y. $\tilde{\tau} = k_{tx}\tau$ is its dimensionless version. The probability density of τ is

$$p(\tau = t) = \frac{k_{tx} - w_1}{w_2 - w_1} w_2 e^{-w_2 t} + \frac{w_2 - k_{tx}}{w_2 - w_1} w_1 e^{-w_1 t},$$

which is an affine combination of exponentials and where $w_1, w_2 > 0$ are the roots of $w \mapsto (\bar{k}^- + k_{tx} - w)(\bar{k}^+ - w) - \bar{k}^+ \bar{k}^-$ with $w_1 < w_2$.

Figure 5 presents the numerical solution of Equation (21) for a fixed value of $-\beta\Delta\mu = 16$, corresponding to the chemical work performed by ATP hydrolysis[46], and a fixed ratio of $-\Delta g/\Delta\mu = 0.5$. In Figure 5a the information capacity is evaluated for different $r > 0$ subject given that $-\Delta g/\Delta\mu = 0.5$. Complementarily, Figure 5b displays a density plot of the e_p

in the $(\bar{k}^+, \bar{k}^-)$-plane together with its hyperbolic level sets of and the curve of DIR maximizing effective rates for varying $r > 0$.

In contrast, Figure 6 shows the information capacity for varying $-\Delta g/\Delta\mu$ and different level sets of e_p. In the range $-\Delta g/\Delta\mu \in [0,1]$ the DIR increases with raised e_p.

For $-\Delta g/\Delta\mu \notin [0,1]$ the level set $e_p = 0$ in the $(\bar{k}^+, \bar{k}^-)$-plane crosses the DIR maximizing ratio $\bar{k}^-/\bar{k}^+ = e - 1$. However, $-\Delta g/\Delta\mu < 0$ can be regarded as implausible since ON would then have less Gibbs free energy than OFF. $-\Delta g/\Delta\mu > 1$ on the other hand corresponds to $\Delta Q^{(1)} < 0$, i.e., a chemically driven reaction, which additionally absorbs heat from the environment. Both scenarios are biophysically implausible.

For the three-state promoter we define the capacity problem as

$$C = \sup_{\substack{0 \leq k_1^+, k_1^-, k_2^-, k_3^+ \\ 0 \leq e_p \leq r}} \bar{\mathbb{I}}(X \to Y),$$

where Δg, $\Delta\mu$ and k_{tx} are fixed. The DIR of this model can be shown to have the form[26]

$$\bar{\mathbb{I}}(X \to Y) = k_{tx} \ln(k_{tx}) \pi(ON) + \frac{h(\tau) - 1}{\mathsf{E}[\tau]}. \tag{23}$$

The corresponding information capacity curves for $k_{tx} = 1$ in Figure 7 are analogous to the curves of the two-state promoter.

VI. CONCLUSION AND OUTLOOK

In this paper, we explored the relationship between energy dissipation and information processing in cellular systems using Markovian chemical reaction networks. We provided a pedagogical introduction to stochastic thermodynamics for open CRNs in Kurtz's process-based formulation. Under the assumptions presented, the energy dissipation rate at the non-equilibrium steady state is identified with the entropy production rate of the reaction counting processes. Parallel reaction channels were identified as the source of discrepancies in this identification when traditional formulations of stochastic thermodynamics are applied to open CRNs.

We extended the mutual information and directed information to account for information transmission between arbitrary subnetworks within CRNs, providing the first rigorous derivation for this case. While suitable channel coding theorems for CRNs are still lacking, the DI is a promising candidate for future research.

Case studies in gene regulation demonstrated that energy dissipation constrains information throughput, with the directed information rate (DIR) increasing with the EPR, supporting the hypothesis that cellular information processing is limited by energy dissipation.

The insights gained here provide a foundation for further research on the energetic limits of information processing capabilities in biological systems, with implications for the design of energy-efficient biological networks.

(a) Maximum DIR for fixed $-\Delta g/\Delta\mu = 0.5$ as a function of the upper bound $r \geq e_p$.

(b) Maximum DIR as a function of $-\Delta g/\Delta\mu$ for different level sets of e_p.

FIG. 5: Maximum DIR for the two-state promoter model and fixed $-\beta\Delta\mu = 16$ and $-\Delta g/\Delta\mu = 0.5$.

FIG. 6: Maximum DIR of the two-state promoter model with $-\beta\Delta\mu = 16$ as a function of $-\Delta g/\Delta\mu$ for different level sets of e_p.

Appendix A: Mutual information between subnetworks of Markovian CRNs

In the following we provide the details for the proof of Theorem 1. For brevity we adopt the notation of Ref. 33 for this section. The term at $s = 0$ stems from the fact that the integrator μ specified in Theorem 4.1, is the product measure

$$\mu = \left(\sum_{x\in\mathbb{Z}_{\geq0}^{|\mathscr{S}^X|}} \delta_x \quad \otimes \sum_{y\in\mathbb{Z}_{\geq0}^{|\mathscr{S}^Y|}} \delta_y \right) \otimes (\delta_0 + \tilde{\mu})$$

where δ_x denotes a Dirac measure at $x \in \mathbb{Z}_{\geq0}^{|\mathscr{S}^X|}$, δ_0 a Dirac measure at $0 \in \mathbb{R}_{\geq0}$ and $\tilde{\mu}$ is the Lebesgue measure on $\mathbb{R}_{\geq0}$. The joint compensator process is absolutely continuous w.r.t. μ and has a $(\mathscr{F}^{XY}, \mathbb{P}^{XY})$-predictable density process of the form

$$\bar{g}_1(x,y,s) = \mathbb{P}(X_0 = x, Y_0 = y)\mathbb{1}\{s = 0\} \\ + \left(g_1(x,s) \times g_1'(y,s)\right)\mathbb{1}\{s > 0\}.$$

Thus, for all $s > 0$, the compensator can be written as a product measure over the state spaces of X and Y, yielding a product decomposition of the Radon-Nikodym derivative, apart from the initial value, where one factor accounts for changes

in X and the other for changes in Y. The $(\mathscr{F}^{XY}, \mathbb{P}^X \otimes \mathbb{P}^Y)$-predictable density has the form

$$\bar{g}(x,y,s) = \mathbb{P}(X_0 = x)\mathbb{P}(Y_0 = y)\mathbb{1}\{s = 0\} \\ + \left(\mathbb{E}[g_1(x,s) \mid \mathscr{F}_{s-}^X] \times \mathbb{E}[g_1'(y,s) \mid \mathscr{F}_{s-}^Y]\right)\mathbb{1}\{s > 0\}.$$

Appendix B: Integral version of directed information

Weissman *et al.*, in Proposition 2 of Ref. 9, provided conditions for the existence of the right derivative of the directed information (DI) for paths on $[0,t]$, along with an explicit expression. They informally stated an "integral version" of this Proposition, implying that $s \mapsto \mathbb{I}(X_{[0,s)} \to Y_{[0,s)})$ be absolutely continuous. However, even if the conditions for the right derivative are satisfied for all $s \in [0,t]$, discontinuities in the DI may still occur. In such cases, the DI is not absolutely continuous if the left derivative fails to exist somewhere in the interval. For example, consider a piecewise constant process with càdlàg paths and deterministic jump times: the right derivative exists and is zero everywhere, yet the left derivative does not exist at the jump times, violating the uniform convergence condition in an analogous left-sided version of Proposition 2 in Ref. 9. Additionally, if the random paths lack a fixed deterministic initial value, the DI may have a positive value at $s = 0$, in accordance with the fundamental theorem of calculus. In what follows, we build on the ideas in the original proofs to establish rigorous conditions for the existence of the common derivatives of $s \mapsto \mathbb{I}(X_{[0,s]} \to Y_{[0,s]})$.

For completeness we state the following lemma, which is analogous to Proposition 1 in Ref. 9 and its proof.

Lemma 1. *Let $t, t' \in \mathscr{T}(0,t)$ be partitions of $[0,t]$, such that t' is a refinement of t, i.e., exists a subsequence $(n_m)_{m\in\{1,\dots,n(t)\}}$ of $(1,\dots,n(t'))$, such that $t_m = t'_{n_m}$ for all m. Then*

$$\sum_{i=1}^{n(t')} \mathbb{I}(Y_{[t'_{i-1},t'_i]}; X_{[0,t'_i]} \mid Y_{[0,t'_{i-1}]}) \leq \sum_{i=1}^{n(t)} \mathbb{I}(Y_{[t_{i-1},t_i]}; X_{[0,t_i]} \mid Y_{[0,t_{i-1}]}).$$

(a) Maximum DIR for fixed $-\Delta g/\Delta\mu = 0.5$ as a function of the upper bound $r \geq e_p$.

(b) Maximum DIR as a function of $-\Delta g/\Delta\mu$ for different level sets of e_p.

FIG. 7: Maximum DIR for the three-state promoter model with fixed $-\beta\Delta\mu = 16$.

The following lemma resembles a chain rule for directed information.

Lemma 2. *Let* $0 < s < t$. *Then*

$$\mathbb{I}(X_{[0,t]} \to Y_{[0,t]}) = \mathbb{I}(X_{[0,s]} \to Y_{[0,s]})$$
$$+ \mathbb{I}(X_{[0,t]} \to Y_{[s,t]} \mid Y_{[0,s)})$$

with $\mathbb{I}(X_{[0,t]} \to Y_{[s,t]} \mid Y_{[0,s)}) = \inf_{t \in \mathscr{T}(s,t)} \sum_{i=1}^{n(t)} \mathbb{I}(Y_{[t_{i-1},t_i]}; X_{[0,t_i]} \mid Y_{[0,t_{i-1})})$.

Proof. Consider the function $\Phi \colon \mathscr{T}(0,s) \times \mathscr{T}(s,t) \to \mathscr{T}(0,t)$, $(\boldsymbol{a},\boldsymbol{b}) \mapsto (a_1,\ldots,a_{n(\boldsymbol{a})},b_1,\ldots,b_{n(\boldsymbol{b})})$ with image $\Phi(\mathscr{T}(0,s) \times \mathscr{T}(s,t))$. Then

$$\mathbb{I}(X_{[0,t]} \to Y_{[0,t]})$$

$$= \inf_{t \in \mathscr{T}(0,t)} \sum_{i=1}^{n(t)} \mathbb{I}(Y_{[t_{i-1},t_i]}; X_{[0,t_i]} \mid Y_{[0,t_{i-1})})$$

$$\leq \inf_{t \in \Phi(\mathscr{T}(0,s) \times \mathscr{T}(s,t))} \sum_{i=1}^{n(t)} \mathbb{I}(Y_{[t_{i-1},t_i]}; X_{[0,t_i]} \mid Y_{[0,t_{i-1})})$$

$$= \mathbb{I}(X_{[0,s]} \to Y_{[0,s]}) + \mathbb{I}(X_{[0,t]} \to Y_{[s,t]} \mid Y_{[0,s)}),$$

where the inequality follows from $\Phi(\mathscr{T}(0,s) \times \mathscr{T}(s,t)) \subseteq \mathscr{T}(0,t)$ and the last equality follows by the additive property of the infimum[47]. For the reverse inequality consider the partitions $\boldsymbol{t}',\boldsymbol{t} \in \mathscr{T}(0,t)$, such that $\boldsymbol{t}' = \boldsymbol{t}$ if there is a j such that $t_j = s$ and otherwise $n(\boldsymbol{t}') = n(\boldsymbol{t})+1$ and there is a j such that $t'_j = s, t'_i = t_i$ for all $i < j$ and $t'_{i+1} = t_i$ for all $i \geq j$. Now define $\Psi \colon \mathscr{T}(0,t) \to \mathscr{T}(0,s) \times \mathscr{T}(s,t)$ such that $\Psi(\boldsymbol{t}) = \boldsymbol{t}'$ and note that

$$\mathbb{I}(X_{[0,s]} \to Y_{[0,s]}) = \inf_{\boldsymbol{s} \in \mathscr{T}(0,s)} \sum_{i=1}^{n(s)} \mathbb{I}(Y_{[s_{i-1},s_i]}; X_{[0,s_i]} \mid Y_{[0,s_{i-1})})$$

$$= \inf_{\boldsymbol{t} \in \mathscr{T}(0,t)} \sum_{i=1}^{n(t')} \mathbb{I}(Y_{[t'_{i-1},t'_i]}; X_{[0,t'_i]} \mid Y_{[0,t'_{i-1})}) \mathbb{1}\{t'_i \leq s\} \text{ and}$$

$$\mathbb{I}(X_{[0,t]} \to Y_{[s,t]} \mid Y_{[0,s)}) = \inf_{\boldsymbol{s} \in \mathscr{T}(s,t)} \sum_{i=1}^{n(s)} \mathbb{I}(Y_{[s_{i-1},s_i]}; X_{[0,s_i]} \mid Y_{[0,s_{i-1})})$$

$$= \inf_{\boldsymbol{t} \in \mathscr{T}(0,t)} \sum_{i=1}^{n(t')} \mathbb{I}(Y_{[t'_{i-1},t'_i]}; X_{[0,t'_i]} \mid Y_{[0,t'_{i-1})}) \mathbb{1}\{t'_i > s\},$$

where we omitted explicitly writing Ψ for brevity. Thus we obtain

$$\mathbb{I}(X_{[0,s]} \to Y_{[0,s]}) + \mathbb{I}(X_{[0,t]} \to Y_{[s,t]} \mid Y_{[0,s)})$$

$$\leq \inf_{\boldsymbol{t} \in \mathscr{T}(0,t)} \sum_{i=1}^{n(\boldsymbol{t}')} \mathbb{I}(Y_{[t'_{i-1},t'_i]}; X_{[0,t'_i]} \mid Y_{[0,t'_{i-1})})$$

$$\leq \inf_{\boldsymbol{t} \in \mathscr{T}(0,t)} \sum_{i=1}^{n(\boldsymbol{t})} \mathbb{I}(Y_{[t_{i-1},t_i]}; X_{[0,t_i]} \mid Y_{[0,t_{i-1})})$$

$$= \mathbb{I}(X_{[0,t]} \to Y_{[0,t]}),$$

where for the first inequality we applied the well-known inequality for the infimum of the sum of real-valued bounded functions, and for the second we applied Lemma 1. $\square$

The central function for deriving a density of DI is $\xi \colon \mathscr{G}_0 \to \mathbb{R}_{\geq 0}$ with $\mathscr{G}_0 := \{(s,h) \in (0,t) \times \mathbb{R} \setminus \{0\} \colon 0 < s+h < t\}$ and

$$\xi(s,h) := \begin{cases} \frac{1}{h}\mathbb{I}(Y_{[s,s+h]}; X_{[0,s+h]} \mid Y_{[0,s)}) & \text{for } h > 0 \\ -\frac{1}{h}\mathbb{I}(Y_{[s+h,s]}; X_{[0,s]} \mid Y_{[0,s+h)}) & \text{for } h < 0. \end{cases}$$

The condition in $\mathscr{G}_0$ serves as a technical restriction on the range of h for different s, resulting in the set having a trapezoidal shape. The definition further implies $\xi(s,h) = \xi(s+h,-h)$ for all $(s,h) \in \mathscr{G}_0$. Let

$$\mathscr{D} := \{s \in (0,t) \colon \lim_{h \to 0} \xi(s,h) \text{ exists}\}$$

be the set, where the $h \mapsto \xi(s,h)$ has a continuous extension at $h = 0$. With this set we define $\mathscr{G} := \{(u,h) \in \mathscr{D} \times \mathbb{R} \colon 0 < u+h < t\}$ and the function $\overline{\xi} \colon \mathscr{G} \to \mathbb{R}_{\geq 0}$ with

$$\overline{\xi}(s,h) = \begin{cases} \xi(s,h) & \text{for } h \neq 0 \\ \lim_{h \to 0} \xi(s,h) & \text{for } h = 0. \end{cases}$$

We denote the open ball of radius $\varepsilon > 0$ around $(s,0) \in \mathscr{G}$ by $U_\varepsilon(s) := \{(u,h) \in (0,t) \times \mathbb{R} \colon \sqrt{h^2 + (u-s)^2} < \varepsilon\}$. Note that if $(s-\varepsilon',s+\varepsilon') \subseteq \mathscr{D}$ for some $\varepsilon' > 0$, then always exists $0 < \varepsilon < \varepsilon'$ with $U_\varepsilon(s) \subseteq \mathscr{G}$.

In simple words, the following theorem states that the derivative of the DI exists at $s \in (0,t)$ if the bivariate function $\overline{\xi}$ exists and is continuous in the neighborhood of $(s,0)$.

Theorem 4. *Let $s \in \mathscr{D}$. If $\varepsilon > 0$ exists, such that $(s - \varepsilon, s + \varepsilon) \subseteq \mathscr{D}$, $U_\varepsilon(s) \subseteq \mathscr{G}$ and $\overline{\xi}$ is continuous on $U_\varepsilon(s)$, then*

$$\frac{\mathrm{d}}{\mathrm{d}s} \mathbb{I}(X_{[0,s]} \to Y_{[0,s]}) = \overline{\xi}(s,0).$$

Proof. Let $h \in (-\varepsilon, \varepsilon) \setminus \{0\}$. Then $K(|h|) := \{(u,\delta) \in (0,t) \times \mathbb{R} : \sqrt{\delta^2 + (u-s)^2} \le |h|\} \subset U_\varepsilon(s)$ is a compact subset. By Lemma 2 the difference quotient of the DI at $s \in \mathscr{D}$, for any h, is given by

$$\frac{1}{h}\left(\mathbb{I}(X_{[0,s+h]} \to Y_{[0,s+h]}) - \mathbb{I}(X_{[0,s]} \to Y_{[0,s]})\right)$$
$$= \begin{cases} \frac{1}{|h|}\mathbb{I}(X_{[0,s+h]} \to Y_{[s,s+h]} \mid Y_{[0,s)}) & \text{for } h > 0 \\ \frac{1}{|h|}\mathbb{I}(X_{[0,s]} \to Y_{[s+h,s]} \mid Y_{[0,s+h)}) & \text{for } h < 0. \end{cases}$$

Then we have for $h \in (0,\varepsilon)$

$$\frac{1}{|h|}\mathbb{I}(X_{[0,s+h]} \to Y_{[s,s+h]} \mid Y_{[0,s)})$$
$$= \frac{1}{|h|} \inf_{t \in \mathscr{T}(s,s+h)} \sum_{i=1}^{n(t)} \mathbb{I}(Y_{[t_{i-1},t_i]}; X_{[0,t_i]} \mid Y_{[0,t_{i-1})}) \quad \text{(B1)}$$
$$= \frac{1}{|h|} \inf_{t \in \mathscr{T}(s,s+h)} \sum_{i=1}^{n(t)} (t_i - t_{i-1})\overline{\xi}(t_{i-1}, t_i - t_{i-1})$$

and for $h \in (-\varepsilon,0)$

$$\frac{1}{|h|}\mathbb{I}(X_{[0,s]} \to Y_{[s+h,s]} \mid Y_{[0,s+h)})$$
$$= \frac{1}{|h|} \inf_{t \in \mathscr{T}(s+h,s)} \sum_{i=1}^{n(t)} \mathbb{I}(Y_{[t_{i-1},t_i]}; X_{[0,t_i]} \mid Y_{[0,t_{i-1})}) \quad \text{(B2)}$$
$$= \frac{1}{|h|} \inf_{t \in \mathscr{T}(s+h,s)} \sum_{i=1}^{n(t)} (t_i - t_{i-1})\overline{\xi}(t_{i-1}, t_i - t_{i-1})$$

By the continuity of $\overline{\xi}$ on all compact sets $K(|h|)$, both $\max_{(\tau,\kappa) \in K(|h|)} \overline{\xi}(\tau,\kappa)$ and $\min_{(\tau,\kappa) \in K(|h|)} \overline{\xi}(\tau,\kappa)$ exist[47] (cf. Theorem 4.28 therein). Since supremum and infimum are taken on the set $K(|h|)$ it follows that $\lim_{h \to 0} \max_{(\tau,\kappa) \in K(|h|)} \overline{\xi}(\tau,\kappa) = \lim_{h \to 0} \min_{(\tau,\kappa) \in K(|h|)} \overline{\xi}(\tau,\kappa) = \overline{\xi}(s,0)$. Noting that $\mathscr{T}(s+\delta,s) = \mathscr{T}(s,s-\delta) = \emptyset$ for any $\delta \in (0,\varepsilon)$ and

using (B1) and (B2) we obtain for any $h \in (-\varepsilon, \varepsilon) \setminus \{0\}$

$$\min_{(\tau,\kappa) \in K(|h|)} \overline{\xi}(\tau,\kappa)$$
$$= \frac{1}{|h|} \inf_{t \in \mathscr{T}(s+h,s) \cup \mathscr{T}(s,s+h)} \sum_{i=1}^{n(t)} (t_i - t_{i-1}) \min_{(\tau,\kappa) \in K(|h|)} \overline{\xi}(\tau,\kappa)$$
$$\le \frac{1}{|h|} \inf_{t \in \mathscr{T}(s+h,s) \cup \mathscr{T}(s,s+h)} \sum_{i=1}^{n(t)} (t_i - t_{i-1})\overline{\xi}(t_{i-1}, t_i - t_{i-1})$$
$$= \frac{1}{h}\left(\mathbb{I}(X_{[0,s+h]} \to Y_{[0,s+h]}) - \mathbb{I}(X_{[0,s]} \to Y_{[0,s]})\right)$$
$$\le \frac{1}{|h|} \inf_{t \in \mathscr{T}(s+h,s) \cup \mathscr{T}(s,s+h)} \sum_{i=1}^{n(t)} (t_i - t_{i-1}) \max_{(\tau,\kappa) \in K(|h|)} \overline{\xi}(\tau,\kappa)$$
$$= \max_{(\tau,\kappa) \in K(|h|)} \overline{\xi}(\tau,\kappa).$$

Taking the limit on both sides of the inequality finishes the proof. $\qquad\square$

We complete the section by providing the desired "integral version".

Corollary 1. *If $\mathscr{D} = (0,t)$ and $\overline{\xi}$ is continuous on $\mathscr{G}$, then*

$$\mathbb{I}(X_{[0,t]} \to Y_{[0,t]})$$
$$= \mathbb{I}(Y_0; X_0) + \int_0^t \lim_{h \searrow 0} \frac{1}{h}\mathbb{I}(Y_{[s,s+h]}; X_{[0,s+h]} \mid Y_{[0,s)}) \, \mathrm{d}s \quad \text{(B3)}$$
$$+ \lim_{h \searrow 0} \mathbb{I}(Y_{[t-h,t]}; X_{[0,t]} \mid Y_{[0,t-h)}).$$

Proof. By Theorem 4 and the fundamental theorem of calculus, we directly obtain

$$\lim_{h \searrow 0}\left(\mathbb{I}(X_{[0,t-h]} \to Y_{[0,t-h]}) - \mathbb{I}(X_{[0,h]} \to Y_{[0,h]})\right)$$
$$= \int_0^t \lim_{h \searrow 0} \frac{1}{h}\mathbb{I}(Y_{[s,s+h]}; X_{[0,s+h]} \mid Y_{[0,s)}) \, \mathrm{d}s.$$

At the right boundary

$$\lim_{h \searrow 0}\left(\mathbb{I}(X_{[0,t]} \to Y_{[0,t]}) - \mathbb{I}(X_{[0,t-h]} \to Y_{[0,t-h]})\right)$$
$$= \lim_{h \searrow 0} \mathbb{I}(X_{[0,t]} \to Y_{[t-h,t]} \mid Y_{[0,t-h)})$$
$$= \lim_{h \searrow 0} \inf_{t \in \mathscr{T}(t-h,t)} \sum_{i=1}^{n(t)} \mathbb{I}(Y_{[t_{i-1},t_i]}; X_{[0,t_i]} \mid Y_{[0,t_{i-1})})$$
$$\ge \lim_{h \searrow 0} \inf_{t \in \mathscr{T}(t-h,t)} \mathbb{I}(Y_{[t_{n(t)-1},t]}; X_{[0,t]} \mid Y_{[0,t_{n(t)-1})})$$
$$= \lim_{h \searrow 0} \lim_{\delta \searrow 0} \mathbb{I}(Y_{[t-\delta,t]}; X_{[0,t]} \mid Y_{[0,t-\delta)}),$$

where the first equality follows with Lemma 2 and the last equality follows with the fact that $\delta \mapsto \mathbb{I}(Y_{[t-\delta,t]}; X_{[0,t]} \mid Y_{[0,t-\delta)})$ is a monotone decreasing function, as can be shown

with the chain rule for mutual information. On the other hand

$$\lim_{h \searrow 0} \inf_{\boldsymbol{t} \in \mathscr{T}(t-h,t)} \sum_{i=1}^{n(\boldsymbol{t})} \mathbb{I}(Y_{[t_{i-1},t_i]}; X_{[0,t_i]} \mid Y_{[0,t_{i-1}]})$$

$$\leq \lim_{h \searrow 0} \inf_{\boldsymbol{t} \in \mathscr{T}(t-h,t)} \mathbb{I}(Y_{[t-h,t]}; X_{[0,t]} \mid Y_{[0,t-h]})$$

$$= \lim_{h \searrow 0} \mathbb{I}(Y_{[t-h,t]}; X_{[0,t]} \mid Y_{[0,t-h]})$$

by Lemma 1, and setting $\boldsymbol{t} = t_1 = t$. At the left boundary we have

$$\lim_{h \searrow 0} \mathbb{I}(X_{[0,h]} \to Y_{[0,h]})$$

$$= \lim_{h \searrow 0} \inf_{\boldsymbol{t} \in \mathscr{T}(0,h)} \sum_{i=1}^{n(\boldsymbol{t})} \mathbb{I}(Y_{[t_{i-1},t_i]}; X_{[0,t_i]} \mid Y_{[0,t_{i-1}]})$$

$$\geq \lim_{h \searrow 0} \inf_{\boldsymbol{t} \in \mathscr{T}(0,h)} \mathbb{I}(Y_{[0,t_1]}; X_{[0,t_1]})$$

$$= \mathbb{I}(X_0; Y_0),$$

where we used that $\delta \mapsto \mathbb{I}(X_{[0,\delta]}; Y_{[0,\delta]})$ is monotone increasing by the chain rule for mutual information. By Lemma 1 and setting $\boldsymbol{t} = t_1 = h$ we obtain

$$\lim_{h \searrow 0} \inf_{\boldsymbol{t} \in \mathscr{T}(0,h)} \sum_{i=1}^{n(\boldsymbol{t})} \mathbb{I}(Y_{[t_{i-1},t_i]}; X_{[0,t_i]} \mid Y_{[0,t_{i-1}]})$$

$$\leq \lim_{h \searrow 0} \inf_{\boldsymbol{t} \in \mathscr{T}(0,h)} \mathbb{I}(Y_{[0,h]}; X_{[0,h]})$$

$$= \mathbb{I}(X_0; Y_0).$$

$\square$

Appendix C: Directed information between subnetworks of Markovian CRNs

In the following we constructively show that the conditions for Equation (B3) are satisfied and thereby verify the expressions (4) and (5) for the DI between subnetworks. We start by defining the relevant filtrations $\mathscr{F}^1$, $\mathscr{F}^{X,2}$, $\mathscr{F}^{Y,2}$ such that for any fixed $s \geq 0$ and all $\tau \geq 0$ they satisfy

$$\mathscr{F}_\tau^{X,2} = \mathscr{F}_{s-}^Y \vee \mathscr{F}_\tau^X$$

$$\mathscr{F}_\tau^{Y,2} = \begin{cases} \mathscr{F}_{s-}^Y & \text{for } \tau < s \\ \mathscr{F}_\tau^Y & \text{for } \tau \geq s, \end{cases}$$

$$\mathscr{F}_\tau^1 = \begin{cases} \mathscr{F}_\tau^{X,2} & \text{for } \tau < s \\ \mathscr{F}_\tau^{XY} & \text{for } \tau \geq s. \end{cases}$$

Here, $\vee$ denotes the join operation for sigma algebras. Let Σ_X and Σ_Y be the power sets of $\mathscr{X} := \mathbb{Z}_{\geq 0}^{|\mathscr{S}^X|}$ and $\mathscr{Y} := \mathbb{Z}_{\geq 0}^{|\mathscr{S}^Y|}$, respectively. The measurable space of trajectories of $(X,Y)_{[0,t]}$ is given by the product space $(\mathscr{X}^{[0,t]} \times \mathscr{Y}^{[0,t]}, \Sigma_X^{\otimes[0,t]} \otimes \Sigma_Y^{\otimes[0,t]})$. We define the conditional probability measures $\mathbb{P}_{\tau,s}^1$, $\mathbb{P}_{\tau,s}^2$ such that for all $\tau \geq s$, $A \in \Sigma_X^{\otimes[0,\tau]}$ and $B \in \Sigma_Y^{\otimes[s,\tau]}$ it holds that

$$\mathbb{P}_{\tau,s}^1(A \times B) = \mathbb{E}[\mathbb{1}\{X_{[0,\tau]} \in A, \ Y_{[s,\tau]} \in B\} \mid \mathscr{F}_{s-}^Y] \text{ and}$$

$$\mathbb{P}_{\tau,s}^2(A \times B) = \mathbb{E}[\mathbb{1}\{X_{[0,\tau]} \in A\} \mid \mathscr{F}_{s-}^Y] \otimes \mathbb{E}[\mathbb{1}\{Y_{[s,\tau]} \in B\} \mid \mathscr{F}_{s-}^Y].$$

Then, for $h > 0$ we obtain

$$\mathbb{I}(Y_{[s,s+h]}; X_{[0,s+h]} \mid Y_{[0,s)})$$

$$= \mathbb{E}\left[\mathbb{E}\left[\ln\left(\frac{d\mathbb{P}_{s+h,s}^1}{d\mathbb{P}_{s+h,s}^2}\right) \middle| \mathscr{F}_{s-}^Y\right]\right] \qquad (C1)$$

$$= \int_{\mathscr{Y}^{[0,s)}} \mathbb{E}\left[\ln\left(\frac{d\mathbb{P}_{s+h,s}^1}{d\mathbb{P}_{s+h,s}^2}\right) \middle| Y_{[0,s)} = y_{[0,s)}\right] d\mathbb{P}_{[0,s)}^Y(y_{[0,s)}).$$

With Theorem 4.1 in[33], the conditional expectation in (C1) obeys

$$\mathbb{E}\left[\ln\left(\frac{d\mathbb{P}_{s+h,s}^1}{d\mathbb{P}_{s+h,s}^2}\right) \middle| \mathscr{F}_s^Y\right]$$

$$= \sum_{v_X \in \mathscr{V}_X} \int_0^{s+h} \mathbb{E}\left[\phi(\lambda_u^{X,1}(v_X)) - \phi(\lambda_u^{X,2}(v_X)) \mid \mathscr{F}_{s-}^Y\right] du$$

$$+ \sum_{v_Y \in \mathscr{V}_Y} \int_s^{s+h} \mathbb{E}\left[\phi(\lambda_u^{Y,1}(v_Y)) - \phi(\lambda_u^{Y,2}(v_Y)) \mid \mathscr{F}_{s-}^Y\right] du,$$

with newly defined intensity processes

$$\lambda_u^{X,1}(v_X) = \mathbb{E}\left[\lambda^X(v_X, Z(u-)) \mid \mathscr{F}_{u-}^1\right]$$

$$\lambda_u^{X,2}(v_X) = \mathbb{E}\left[\lambda^X(v_X, Z(u-)) \mid \mathscr{F}_{u-}^{X,2}\right]$$

for all $u \geq 0$ and

$$\lambda_u^{Y,1}(v_Y) = \mathbb{E}\left[\lambda^Y(v_Y, Z(u-)) \mid \mathscr{F}_{u-}^1\right]$$

$$\lambda_u^{Y,2}(v_Y) = \mathbb{E}\left[\lambda^Y(v_Y, Z(u-)) \mid \mathscr{F}_{u-}^{Y,2}\right]$$

for all $u \geq s$. Now note that $\mathscr{F}_u^1 = \mathscr{F}_u^{X,2}$ for all $u < s$, such that all v_X the integrals vanish on $[0,s]$. Note also that $\lim_{h \searrow 0} \mathscr{F}_{s+h}^1 = \mathscr{F}_s^{X,2}$, such that $\mathbb{E}\left[\phi(\lambda_s^{X,1}(v_X)) - \phi(\lambda_s^{X,2}(v_X)) \mid \mathscr{F}_{s-}^Y\right] = 0$. Hence,

$$\lim_{h \searrow 0} \frac{1}{h} \int_s^{s+h} \mathbb{E}\left[\phi(\lambda_u^{X,1}(v_X)) - \phi(\lambda_u^{X,2}(v_X)) \mid \mathscr{F}_{s-}^Y\right] du = 0$$

Lastly, $\mathscr{F}_u^1 = \mathscr{F}_u^{XY}$ and $\mathscr{F}_u^{Y,2} = \mathscr{F}_u^Y$ for all $u \geq s$, which implies $\lambda_u^{Y,1}(v_Y) = \lambda_u^Y(v_Y)$ and $\lambda_u^{Y,2}(v_Y) = \hat{\lambda}_u^Y(v_Y)$. Concluding, we have

$$\lim_{h \searrow 0} \frac{1}{h} \mathbb{I}(Y_{[s,s+h]}; X_{[0,s+h]} \mid Y_{[0,s)})$$

$$= \sum_{v_Y \in \mathscr{V}_Y} \mathbb{E}\left[\phi(\lambda_s^Y(v_Y)) - \phi(\hat{\lambda}_s^Y(v_Y))\right],$$

the left-limit follows analogously and coincides with the right limit, and the function $\bar{\xi}$ is continuous on its whole domain. Equation (5) follows directly.

Appendix D: Equivalent equilibrium conditions

Proof of Thm. 3. We start showing the equivalence of (b), (c) and (d).

(d) $\implies$ (c): For all $j \in \{1, \ldots, n+1\}$ the condition in (d) can be rewritten as

$$\lambda_{m_j}^{\rho_j}(z_{j-1})\pi(z_{j-1}) = \lambda_m^{-\rho_j}(z_j)\pi(z_j)$$

with $z_0 = z_{n+1}$. The implication then simply follows by substituting $\lambda_{m_j}^{\rho_j}(z_{j-1})/\lambda_m^{-\rho_j}(z_j) = \pi(z_j)/\pi(z_{j-1})$.

(c) $\implies$ (b): Note that microscopic reversibility implies that the process is irreducible on each stoichiometric compatibility class, which consists of more than a single state. Note that cycles are not possible at absorbing states, so we can ignore this case. In turn, non-explosiveness, irreducibility and the existence of a stationary distribution imply that the process is positive recurrent on the compatibility class, i.e., the mean recurrence time of an arbitrary state is finite. Thus, for any non-absorbing initial state we can decompose the process equation in a sum over completed cycles and a remainder of an incomplete cycle. Let $\{\sigma(z_0, s) \in \mathbb{R}_{\geq 0}\}_{s \geq 0}$ be the stochastic process accounting for the time that has passed since the latest recurrence to z_0. Further let

$$A(z_0, \xi) := \sum_{j=1}^{n+1} \ln\left(\frac{\lambda_{m_j}^{\rho_j}(z_{j-1})}{\lambda_{m_j}^{-\rho_j}(z_j)}\right)$$

denote the cycle affinities, where the stoichiometric cycle is represented by its starting point z_0 and the sequence of directed reactions $\xi = (\rho_1 m_1, \ldots, \rho_{n+1} m_{n+1})$. Let $\{D_\xi(s)\}_{s \geq 0}$ be the counting process of cycles ξ, starting in z_0.

By Definition 6 we write the EPR as

$$e_p = \lim_{t \to \infty} \frac{1}{t}\left(\sum_{m=1}^{M} \mathbb{E}\left[\int_0^t \ln\left(\frac{\lambda_m^+(Z(s-))}{\tilde{\lambda}_m^+(Z(s-),s)}\right) dR_m^+(s)\right] \right.$$
$$\left. + \sum_{m=1}^{M} \mathbb{E}\left[\int_0^t \ln\left(\frac{\lambda_m^-(Z(s-))}{\tilde{\lambda}_m^-(Z(s-),s)}\right) dR_m^-(s)\right]\right)$$

Using the stationarity we decompose

$$\sum_{m=1}^{M} \mathbb{E}\left[\int_0^t \ln\left(\frac{\lambda_m^+(Z(s-))}{\tilde{\lambda}_m^+(Z(s-),s)}\right) dR_m^+(s)\right.$$
$$\left. + \int_0^t \ln\left(\frac{\lambda_m^-(Z(s-))}{\tilde{\lambda}_m^-(Z(s-),s)}\right) dR_m^-(s)\right]$$
$$= \sum_\xi \mathbb{E}\left[A(Z(0), \xi)D_\xi(t)\right]$$
$$+ \sum_{m=1}^{M} \mathbb{E}\left[\int_{t-\sigma(Z(0),t)}^t \ln\left(\frac{\lambda_m^+(Z(s-))}{\tilde{\lambda}_m^+(Z(s-),s)}\right) dR_m^+(s)\right.$$
$$\left. + \int_{t-\sigma(Z(0),t)}^t \ln\left(\frac{\lambda_m^-(Z(s-))}{\tilde{\lambda}_m^-(Z(s-),s)}\right) dR_m^-(s)\right].$$

By the cycle condition $A(Z(0), \xi) = 0$ for all admissible ξ. The contribution of the incomplete cycles vanishes in the limit since $\sigma(Z(0), t)$ is bounded by the recurrence time of $Z(0)$.

(b) $\implies$ (d): All terms in Equation (16) are non-negative as can be shown by distinguishing the cases $J_m(z,t) \geq 0$ and $J_m(z,t) < 0$. Thus, if $e_p = 0$, then all terms must vanish individually, i.e., $\lim_{t \to \infty} J_m(z,t) = 0$ for all z and $m \in \mathscr{R}$. Thus detailed balance is statisfied.

(a) $\implies$ (b): Apply Definition 6.

(d) $\implies$ (a): At stationarity, we have $\mathbb{D}_{\mathrm{KL}}\left(Z(0)\|\tilde{Z}(0)\right) = 0$. Further, the detailed balance condition implies $\tilde{\lambda}_m^\pm(z,t) = \lambda_m^\pm(z)$ for all $t \geq 0$. Hence, $\ln\left(\lambda_m^+(z)/\tilde{\lambda}_m^+(z,t)\right) = 0$ and, consequently, $\mathbb{D}_{\mathrm{KL}}\left((R_m^+(s), R_m^-(s))_{s \in [0,t], m \in \mathscr{R}}\|(\tilde{R}_m^+(s), \tilde{R}_m^-(s))_{s \in [0,t], m \in \mathscr{R}}\right) = 0$. $\square$

Appendix E: EPR of a non-equilibrium two-state model

To derive the EPR for the two-state model we use Equation (14) for the entropy exchange rate, which is equal to the EPR at the NESS:

$$e_p = \pi(OFF)k_1^+ \ln\left(\frac{k_1^+}{k_1^-}\right) + \pi(ON)k_1^- \ln\left(\frac{k_1^-}{k_1^+}\right)$$
$$+ \pi(OFF)k_2^+ \ln\left(\frac{k_2^+}{k_2^-}\right) + \pi(ON)k_2^- \ln\left(\frac{k_2^-}{k_2^+}\right)$$
$$= \ln\left(\frac{k_1^+}{k_1^-}\right)(\pi(OFF)k_1^+ - \pi(ON)k_1^-)$$
$$+ \ln\left(\frac{k_2^+}{k_2^-}\right)(\pi(OFF)k_2^+ - \pi(ON)k_2^-)$$
$$= \ln\left(\frac{k_1^+}{k_1^-}\right)\frac{(k_1^- + k_2^-)k_1^+ - (k_1^+ + k_2^+)k_1^-}{k_1^+ + k_1^- + k_2^+ + k_2^-}$$
$$+ \ln\left(\frac{k_2^+}{k_2^-}\right)\frac{(k_1^- + k_2^-)k_2^+ - (k_1^+ + k_2^+)k_2^-}{k_1^+ + k_1^- + k_2^+ + k_2^-}$$

with $\pi(OFF) = \frac{k_1^- + k_2^-}{k_1^+ + k_1^- + k_2^+ + k_2^-}$ and $\pi(ON) = \frac{k_1^+ + k_2^+}{k_1^+ + k_1^- + k_2^+ + k_2^-}$. The stationary distribution is the kernel of the generator matrix corresponding to the model. Simplifying the above expression yields (20).

To obtain the EPR as a function of the effective rates $\bar{k}^+ = k_1^+ + k_2^+$, $\bar{k}^- = k_1^- + k_2^-$, we we express the rates in the state transition diagram via the effective rates and the potential differences Δg, $\Delta\mu$. We have

$$k_1^- = k_1^+ e^{\beta(\Delta g + \Delta\mu)}, \quad k_2^+ = k_2^- e^{-\beta\Delta g},$$
$$\bar{k}^- = k_1^+ e^{\beta(\Delta g + \Delta\mu)} + k_2^-, \quad \bar{k}^+ = k_1^+ + k_2^- e^{-\beta\Delta g},$$

which implies

$$k_1^+ = \frac{\bar{k}^+ - \bar{k}^- e^{-\beta\Delta g}}{1 - e^{\Delta\mu}}, \quad k_2^- = \frac{\bar{k}^- - \bar{k}^+ e^{\beta(\Delta g + \Delta\mu)}}{1 - e^{\Delta\mu}}$$

and therefore also

$$k_1^- = \frac{\bar{k}^+ - \bar{k}^- e^{-\beta\Delta g}}{1 - e^{\Delta\mu}}e^{\beta(\Delta g + \Delta\mu)}$$
$$k_2^+ = \frac{\bar{k}^- - \bar{k}^+ e^{\beta(\Delta g + \Delta\mu)}}{1 - e^{\Delta\mu}}e^{-\beta\Delta g}$$

The physical boundary conditions $-\Delta\mu, k_1^+, k_2^- > 0$ constrain the admissible region of the effective rates to

$$\bar{k}^+ > \bar{k}^- e^{-\beta\Delta g}, \quad \bar{k}^- > \bar{k}^+ e^{\beta(\Delta g + \Delta\mu)}.$$

For this region we obtain (22) by substituting the original rates with the above expressions and simplifying.

Appendix F: Interarrival time between transcription events for the circular three-state promoter

The DI between the promoter dynamics and the RNA copy number has was linked to the differential entropy of the interarrival time in (23). Here, we apply Anderson's filtering theorem for semi Markov processes similarly as described in[26]. Define the state space $E := \{J, ON, OFF^*, OFF\}$, where J is the jump state, which mimics ON in terms of leaving transitions. The Laplace version of the semi-Markov kernel density is

$$q^*(s) = \begin{pmatrix} \frac{k_{tx}}{u_{ON}+s} & 0 & \frac{k_3^-}{u_{ON}+s} & \frac{k_2^-}{u_{ON}+s} \\ \frac{k_{tx}}{u_{ON}+s} & 0 & \frac{k_3^-}{u_{ON}+s} & \frac{k_2^-}{u_{ON}+s} \\ 0 & \frac{k_3^+}{u_{OFF^*}+s} & 0 & \frac{k_1^-}{u_{OFF^*}+s} \\ 0 & \frac{k_2^+}{u_{OFF}+s} & \frac{k_1^+}{u_{OFF}+s} & 0 \end{pmatrix},$$

where $u_{ON} = k_2^- + k_3^- + k_{tx}$, $u_{OFF} = k_1^+ + k_2^+$ We filter E to the reduced state space $\{J\}$ which yields the interarrival time distribution f_τ of the output. Since the geometric series of a 3×3-matrix is hard to evaluate, we do the filtering state by state, from right to left. In the first step we identify

$$a^*(s) = \begin{pmatrix} \frac{k_{tx}}{u_{ON}+s} & 0 & \frac{k_3^-}{u_{ON}+s} \\ \frac{k_{tx}}{u_{ON}+s} & 0 & \frac{k_3^-}{u_{ON}+s} \\ 0 & \frac{k_3^+}{u_{OFF^*}+s} & 0 \end{pmatrix}$$

$$b^*(s) = \left(\frac{k_2^-}{u_{ON}+s}, \frac{k_2^-}{u_{ON}+s}, \frac{k_1^-}{u_{OFF^*}+s} \right)^T,$$

$$c^*(s) = \left(0, \frac{k_2^+}{u_{OFF}+s}, \frac{k_1^+}{u_{OFF}+s} \right) \text{ and}$$

$$d^* = 0.$$

Thus

$$b^*(s)c^*(s) = \begin{pmatrix} 0 & \frac{k_2^+ k_2^-}{(u_{OFF}+s)(u_{ON}+s)} & \frac{k_1^+ k_2^-}{(u_{OFF}+s)(u_{ON}+s)} \\ 0 & \frac{k_2^+ k_2^-}{(u_{OFF}+s)(u_{ON}+s)} & \frac{k_1^+ k_2^-}{(u_{OFF}+s)(u_{ON}+s)} \\ 0 & \frac{k_1^- k_2^+}{(u_{OFF^*}+s)(u_{OFF}+s)} & \frac{k_1^+ k_1^-}{(u_{OFF^*}+s)(u_{OFF}+s)} \end{pmatrix}$$

and

$$\tilde{q}^*(s) = a^*(s) + b^*(s)c^*(s)$$
$$= \begin{pmatrix} \frac{k_{tx}}{u_{ON}+s} & \frac{k_2^+ k_2^-}{(u_{OFF}+s)(u_{ON}+s)} & \frac{k_1^+ k_2^- + k_3^-(u_{OFF}+s)}{(u_{OFF}+s)(u_{ON}+s)} \\ \frac{k_{tx}}{u_{ON}+s} & \frac{k_2^+ k_2^-}{(u_{OFF}+s)(u_{ON}+s)} & \frac{k_1^+ k_2^- + k_3^-(u_{OFF}+s)}{(u_{OFF}+s)(u_{ON}+s)} \\ 0 & \frac{k_1^- k_2^+ + k_3^+(u_{OFF}+s)}{(u_{OFF^*}+s)(u_{OFF}+s)} & \frac{k_1^+ k_1^-}{(u_{OFF^*}+s)(u_{OFF}+s)} \end{pmatrix}$$

In the second step we identify

$$a^*(s) = \begin{pmatrix} \frac{k_{tx}}{u_{ON}+s} & \frac{k_2^+ k_2^-}{(u_{OFF}+s)(u_{ON}+s)} \\ \frac{k_{tx}}{u_{ON}+s} & \frac{k_2^+ k_2^-}{(u_{OFF}+s)(u_{ON}+s)} \end{pmatrix},$$

$$b^*(s) = \frac{k_3^-(r_2+s)}{(u_{OFF}+s)(u_{ON}+s)}(1,1)^T,$$

$$c^*(s) = \frac{k_3^+(r_1+s)}{(u_{OFF^*}+s)(u_{OFF}+s)}(0,1),$$

$$\sum_{k=0}^{\infty}(d^*(s))^k = \frac{(u_{OFF^*}+s)(u_{OFF}+s)}{(u_{OFF^*}+s)(u_{OFF}+s)-k_1^+ k_1^-}$$

with $r_1 := \frac{k_1^- k_2^+}{k_3^+} + u_{OFF}$ and $r_2 := \frac{k_1^+ k_2^-}{k_3^-} + u_{OFF}$. We denote the roots of $s \mapsto (u_{OFF^*}-s)(u_{OFF}-s)-k_1^+ k_1^-$ by w_1, w_2. Then

$$b^*(s)\sum_{k=0}^{\infty}(d^*(s))^k c^*(s)$$
$$= \frac{k_3^+ k_3^-(r_1+s)(r_2+s)}{(u_{OFF}+s)(u_{ON}+s)(w_1+s)(w_2+s)}\begin{pmatrix} 0 & 1 \\ 0 & 1 \end{pmatrix}$$

$$\tilde{\tilde{q}}^*(s) = a^*(s) + b^*(s)\sum_{k=0}^{\infty}(d^*(s))^k c^*(s)$$
$$= \begin{pmatrix} \frac{k_{tx}}{u_{ON}+s} & \frac{k_3^+ k_3^-(r_1+s)(r_2+s)+k_2^+ k_2^-(w_1+s)(w_2+s)}{(u_{OFF}+s)(u_{ON}+s)(w_1+s)(w_2+s)} \\ \frac{k_{tx}}{u_{ON}+s} & \frac{k_3^+ k_3^-(r_1+s)(r_2+s)+k_2^+ k_2^-(w_1+s)(w_2+s)}{(u_{OFF}+s)(u_{ON}+s)(w_1+s)(w_2+s)} \end{pmatrix}.$$

Denoting the roots of $\mathbb{C} \ni s \mapsto k_3^+ k_3^-(r_1-s)(r_2-s)+k_2^+ k_2^-(w_1-s)(w_2-s)$ by q_1, q_2 we rewrite

$$\tilde{\tilde{q}}^*(s) = \begin{pmatrix} \frac{k_{tx}}{u_{ON}+s} & \frac{(k_3^+ k_3^- + k_2^+ k_2^-)(q_1+s)(q_2+s)}{(u_{OFF}+s)(u_{ON}+s)(w_1+s)(w_2+s)} \\ \frac{k_{tx}}{u_{ON}+s} & \frac{(k_3^+ k_3^- + k_2^+ k_2^-)(q_1+s)(q_2+s)}{(u_{OFF}+s)(u_{ON}+s)(w_1+s)(w_2+s)} \end{pmatrix}.$$

For the last filtering step we use that $\frac{a}{b}\frac{1}{1-\frac{a}{b}} = \frac{a}{b-a}$ and $1 + \frac{a}{b-a} = \frac{b}{b-a}$ for any $a \neq b$. We obtain

$$f_\tau^*(s) = \{k_{tx}(u_{OFF}+s)(w_1+s)(w_2+s)\}$$
$$\times \{u_{OFF}+s)(u_{ON}+s)(w_1+s)(w_2+s)$$
$$- (k_3^+ k_3^- + k_2^+ k_2^-)(q_1+s)(q_2+s)\}^{-1}.$$

Denote $R(s) := k_{tx}(u_{OFF}+s)(w_1+s)(w_2+s)$ and let $v_1, \ldots, v_4$ be the roots of $U(s) := (u_{OFF}+s)(u_{ON}+s)(w_1+s)(w_2+s) - (k_3^+ k_3^- + k_2^+ k_2^-)(q_1+s)(q_2+s)$. Then

$$f_\tau(t) = \sum_{k=1}^{4}\frac{R(v_k)}{U'(v_k)}e^{v_k t}.$$

[1]M. Magaña-Acosta and V. Valadez-Graham, "Chromatin Remodelers in the 3D Nuclear Compartment," Frontiers in Genetics **11**, 600615 (2020).

[2]S. L. Johnson and G. J. Narlikar, "Atp hydrolysis coordinates the activities of two motors in a dimeric chromatin remodeling enzyme," Journal of Molecular Biology **434**, 167653 (2022).

[3]W. S. Klug, M. R. Cummings, C. A. Spencer, M. A. Palladino, and D. J. Killian, "Concepts of genetics," (2020).

[4] U. Alon, *An introduction to systems biology: design principles of biological circuits* (Chapman and Hall/CRC, 2019).

[5] M. Behar and A. Hoffmann, "Understanding the temporal codes of intracellular signals," Current opinion in genetics & development **20**, 684–693 (2010).

[6] J. E. Purvis and G. Lahav, "Encoding and Decoding Cellular Information through Signaling Dynamics," Cell **152**, 945–956 (2013).

[7] D. Friedrich, L. Friedel, A. Finzel, A. Herrmann, S. Preibisch, and A. Loewer, "Stochastic transcription in the p53-mediated response to DNA damage is modulated by burst frequency," Molecular Systems Biology **15**, e9068 (2019).

[8] P. R. Ten Wolde, "The price of accuracy," Nature Physics **8**, 361–362 (2012).

[9] T. Weissman, Y.-H. Kim, and H. H. Permuter, "Directed information, causal estimation, and communication in continuous time," IEEE Transactions on Information Theory **59**, 1271–1287 (2012).

[10] D. F. Anderson and T. G. Kurtz, "Continuous time markov chain models for chemical reaction networks," in *Design and analysis of biomolecular circuits: engineering approaches to systems and synthetic biology* (Springer, 2011) pp. 3–42.

[11] C. E. Shannon, "A mathematical theory of communication," The Bell system technical journal **27**, 379–423 (1948).

[12] H. Marko, "The bidirectional communication theory-a generalization of information theory," IEEE Transactions on communications **21**, 1345–1351 (1973).

[13] J. Massey *et al.*, "Causality, feedback and directed information," in *Proc. Int. Symp. Inf. Theory Applic.(ISITA-90)*, Vol. 2 (1990).

[14] S. Uda, "Application of information theory in systems biology," Biophysical reviews **12**, 377–384 (2020).

[15] G. Tkačik and A. M. Walczak, "Information transmission in genetic regulatory networks: a review," Journal of Physics: Condensed Matter **23**, 153102 (2011).

[16] R. Cheong, A. Rhee, C. J. Wang, I. Nemenman, and A. Levchenko, "Information Transduction Capacity of Noisy Biochemical Signaling Networks," science **334**, 354–358 (2011).

[17] A. S. Hansen and E. K. O'Shea, "Limits on information transduction through amplitude and frequency regulation of transcription factor activity," Elife **4**, e06559 (2015).

[18] F. Tostevin and P. R. Ten Wolde, "Mutual Information between Input and Output Trajectories of Biochemical Networks," Physical review letters **102**, 218101 (2009).

[19] I. Lestas, G. Vinnicombe, and J. Paulsson, "Fundamental limits on the suppression of molecular fluctuations," Nature **467**, 174–178 (2010).

[20] Y. Nakahira, F. Xiao, V. Kostina, and J. C. Doyle, "Fundamental limits and achievable performance in biomolecular control," in *2018 Annual American Control Conference (ACC)* (IEEE, 2018) pp. 2707–2714.

[21] L. Duso and C. Zechner, "Path mutual information for a class of biochemical reaction networks," in *2019 IEEE 58th Conference on Decision and Control (CDC)* (IEEE, 2019) pp. 6610–6615.

[22] A.-L. Moor and C. Zechner, "Dynamic information transfer in stochastic biochemical networks," Physical Review Research **5**, 013032 (2023).

[23] D. T. Gillespie, "Exact stochastic simulation of coupled chemical reactions," The journal of physical chemistry **81**, 2340–2361 (1977).

[24] M. Sinzger, M. Gehri, and H. Koeppl, "Poisson channel with binary Markov input and average sojourn time constraint," in *2020 International Symposium on Information Theory (ISIT)* (IEEE, 2020) pp. 2873–2878.

[25] M. Sinzger-D'Angelo and H. Koeppl, "Counting Processes with Piecewise-Deterministic Markov Conditional Intensity: Asymptotic Analysis, Implementation, and Information-Theoretic Use," IEEE Transactions on Information Theory (2023).

[26] M. Gehri, N. Engelmann, and H. Koeppl, "Mutual information of a class of poisson-type channels using markov renewal theory," in *2024 IEEE International Symposium on Information Theory (ISIT)* (2024) pp. 1931–1936.

[27] Y. M. Kabanov, "The capacity of a channel of the Poisson type," Theory of Probability & Its Applications **23**, 143–147 (1978).

[28] M. Reinhardt, G. Tkačik, and P. R. Ten Wolde, "Path Weight Sampling: Exact Monte Carlo Computation of the Mutual Information between Stochastic Trajectories," Physical Review X **13**, 041017 (2023).

[29] A. Das and P. R. t. Wolde, "Exact computation of Transfer Entropy with Path Weight Sampling," arXiv preprint arXiv:2409.01650 (2024).

[30] J. Schnakenberg, "Network theory of microscopic and macroscopic behavior of master equation systems," Rev. Mod. Phys. **48**, 571–585 (1976).

[31] T. Schmiedl and U. Seifert, "Stochastic thermodynamics of chemical reaction networks," The Journal of Chemical Physics **126** (2007).

[32] R. Rao and M. Esposito, "Conservation laws and work fluctuation relations in chemical reaction networks," The Journal of Chemical Physics **149** (2018).

[33] R. Boel, P. Varaiya, and E. Wong, "Martingales on jump processes. II: Applications," SIAM Journal on Control **13**, 1022–1061 (1975).

[34] T. Renner, E. R. Cohen, T. Cvitas, J. G. Frey, B. Holström, K. Kuchitsu, R. Marquardt, I. Mills, F. Pavese, M. Quack, J. Stohner, H. L. Strauss, M. Takami, and A. J. Thor, *Quantities, units and symbols in physical chemistry* (The Royal Society of Chemistry, 2007).

[35] "Elementary reaction," (2019), doi:10.1351/goldbook.E02035.

[36] T. G. Kurtz, "The relationship between stochastic and deterministic models for chemical reactions," The Journal of Chemical Physics **57**, 2976–2978 (1972).

[37] H. Qian and H. Ge, *Stochastic Chemical Reaction Systems in Biology* (Springer International Publishing, 2021).

[38] D. T. Gillespie, "A rigorous derivation of the chemical master equation," Physica A: Statistical Mechanics and its Applications **188**, 404–425 (1992).

[39] P. M. Brémaud, *Point Processes and Queues: Martingale Dynamics*, Vol. 50 (Springer, 1981).

[40] P. M. Brémaud, *A Martingale Approach to Point-Processes*, Ph.D. thesis, EECS Department, University of California, Berkeley (1972).

[41] B. Grigelionis, "Mutual information for locally infinitely divisible random processes," Lithuanian mathematical transactions of the Academy of Sciences of the Lithuanian SSR **14**, 1–6 (1974).

[42] D. Cappelletti and C. Wiuf, "Product-form poisson-like distributions and complex balanced reaction systems," SIAM Journal on Applied Mathematics **76**, 411–432 (2016).

[43] U. Seifert, "Stochastic thermodynamics: From principles to the cost of precision," Physica A: Statistical Mechanics and its Applications **504**, 176–191 (2018).

[44] J. Jacod, "Multivariate Point Processes: Predictable Projection, Radon-Nikodym Derivatives, Representation of Martingales," Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete **31**, 235–253 (1975).

[45] D.-Q. Jiang and D. Jiang, *Mathematical Theory of Nonequilibrium Steady States: On the Frontier of Probability and Dynamical Systems* (Springer Science & Business Media, 2004).

[46] B. Alberts, A. Johnson, J. Lewis, M. Raff, K. Roberts, and P. Walter, *Molecular Biology of the Cell*, 6th ed. (Garland Science, New York, NY, 2017).

[47] T. M. Apostol, "Mathematical analysis," (1998).