Title: Berberine is a novel mitochondrial calcium uniporter (MCU) inhibitor that disrupts MCU-EMRE assembly

Authors:

Haixin Zhao[1, 3, 5], Siqi Chen[1, 2, 5], Nian Cao[1, 4, 5], Wenjun Wu[1, 5], Guangqin Liu[1,2], Jun Gao[1], Jiayi Chen[1], Ting Li[1], Dingyi Lu[1], Lingmin Zeng[1], Haizhen Zhu[1], Weina Zhang[1], Qing Xia[1], Teng Li[1], Tao Zhou[1], Xue-Min Zhang [1,2], Ai-Ling Li [1, 2, *], Xin Pan[1, 2, *]

Affiliations:

1. Nanhu Laboratory, National Center of Biomedical Analysis, 27 Tai-Ping Road, Beijing 100039, China

2. School of Basic Medical Sciences, Fudan University, Shanghai 200032, China

3. State Key Laboratory of Experimental Haematology, Fifth Medical Center of Chinese PLA General Hospital, Beijing 100071, China

4. Department of Cardiology, the Sixth Medical Centre, Chinese PLA General Hospital, Beijing, P.R. China

5. These authors contributed equally

*Correspondence: xpan@ncba.ac.cn ; alli@ncba.ac.cn

Abstract

The mitochondrial calcium uniporter (MCU) complex, localized in the inner

mitochondrial membrane, plays a crucial role in regulating mitochondrial Ca^{2+} influx,

thereby impacting cellular energy metabolism and apoptosis. Dysregulation of

mitochondrial calcium levels is implicated in various diseases, including

neurodegenerative disorders, cardiac diseases and cancer. Despite the critical roles of

MCU, developing specific and clinically viable inhibitors has been challenging. Here,

we identify Berberine, a well-established drug with a documented safety profile, as a

potent MCU inhibitor. Utilizing virtual screening of an FDA-approved drug library,

we discovered Berberine's significant inhibitory effect on MCU. Mechanistically,

Berberine localizes within mitochondria and binds near the MCU's juxtamembrane

loop (JML) domain. This binding disrupts the interaction of MCU with its essential

regulator, EMRE, thereby inhibiting rapid Ca^{2+} entry into the mitochondrial matrix.

Notably, Berberine pretreatment reduced mitochondrial Ca^{2+} overload and mitigated

ischemia/reperfusion-induced myocardial injury in a murine model. Our findings

establish Berberine as a potent MCU inhibitor, offering a potential therapeutic avenue

for diseases associated with dysregulated mitochondrial calcium homeostasis.

Significance Statement:

Mitochondrial calcium dysregulation is implicated in various diseases, highlighting

the crucial need for effective MCU inhibitors. However, current MCU inhibitors often

suffer from off-target effects and toxicity, limiting their clinical use. In this study, we

identify Berberine, a drug with an established safety profile, as a novel MCU inhibitor,

demonstrating its potential beyond traditional applications. Berberine directly binds to

the MCU, disrupting its interaction with EMRE, a critical regulator of calcium uptake.

This discovery not only identifies Berberine as the first MCU inhibitor targeting

MCU-EMRE assembly, but also introduces a new avenue for therapeutic intervention

in diseases associated with mitochondrial calcium dysregulation. By repurposing

Berberine, our work paves the way for safer and more effective treatments targeting

mitochondrial dysfunction.

Introduction

The entry of Ca^{2+} ions into mitochondria primarily occurs through a highly selective Ca^{2+} channels known as mitochondrial Ca^{2+} uniporter (MCU) (1). This process is crucial for regulating various cellular processes, including mitochondrial dynamics, energy metabolism and cell death (2). The MCU complex, along with its regulatory components MICU1, MICU2, and EMRE, ensures precise control of mitochondrial Ca2+ uptake, which is crucial for maintaining cellular homeostasis (3). However, under pathological conditions, dysregulation of mitochondrial Ca^{2+} homeostasis may lead to detrimental effects. Excessive Ca^{2+} accumulation in mitochondria, referred to as mitochondrial Ca^{2+} overload, triggers mitochondrial permeability transition pore opening, leading to mitochondrial membrane depolarization, release of pro-apoptotic factors, and ultimately leading to cell death. These processes contribute to various diseases, including neurodegenerative disorders, cardiac diseases, and ischemia/reperfusion injury (4, 5).

Despite the critical role of MCU in disease pathophysiology, developing effective therapeutic interventions targeting mitochondrial Ca^{2+} regulation remains challenging. Existing MCU inhibitors, such as ruthenium red and Ru360, exhibit poor delivery and toxicity issues (6, 7). More recent inhibitors, including DS16570511 and mitoxantrone, have shown promise in preclinical studies but are hampered by toxicity, limiting their clinical applicability (8, 9). Ideal MCU inhibitors that is suitable for clinical application must meet several criteria: 1) efficient penetration of cellular and mitochondrial membranes to effectively inhibit mitochondrial Ca^{2+} uptake; 2) selective modulatation of mitochondrial Ca^{2+} uptake without affecting cytosolic Ca^{2+} signaling, ensuring normal cellular functions; 3) preservation of mitochondrial membrane potential, which is vital for mitochondrial function; and 4) low toxicity at therapeutically relevant concentrations to ensure safety and minimal adverse effects.

In response to the demand for safe and effective MCU inhibitors, we employed virtual molecular docking followed by a functional screening of an FDA-approved

drug library. This led to the identification of Berberine, a natural alkaloid with

established safety profiles, as a novel MCU inhibitor. Berberine localizes to

mitochondria, directly binds to MCU, and disrupts the assembly of the MCU-EMRE

complex. Importantly, Berberine significantly reduces mitochondrial Ca^{2+} overload,

providing cardioprotection against ischemia/reperfusion-induced myocardial injury in

mice. This discovery positions Berberine as a promising candidate for further

development in the broader context of diseases involving mitochondrial Ca^{2+}

dysregulation.

Results

Screening for potential MCU inhibitors from an FDA-approved drug library

MCU functions as a major gate of Ca^{2+} flux from cytosol into the mitochondrial

matrix. Targeting MCU may serve as therapeutic strategies for many diseases, such as

pathological cardiac cell death (10-12). To closely emulate MCU-targeted clinical

treatment, we employed a screening approach using an FDA-approved drug library

comprising 2,816 compounds (Fig. 1*A*). In our initial step, we performed a virtual

screening via MCU structure-based molecular docking and identified 120 hits with a

docking score of < -7.5 kcal/mol, suggesting their potential MCU binding capacity

(Fig. 1*B*). To assess the impact of these hits on mitochondrial Ca^{2+} uptake, we

established HeLa cells expressing 4mt-GCaMP6, a fluorescent indicator reflecting

mitochondrial Ca^{2+} levels (13). Subsequently, we generated a screening system for

mitochondrial Ca^{2+} uptake activity using permeabilized Hela cells. Exposure to 100

μM $CaCl_2$ led to a rapid increase of the fluorescent signal, indicating mitochondrial

Ca^{2+} uptake (*SI Appendix*, Fig. S1*A*). Ru360, a well-known MCU inhibitor (6, 7),

effectively blocked Ca^{2+} entry into mitochondria (Fig 1*C*; *SI Appendix*, Fig. S1). In a

small-scale screen using the 120 hits, four small molecules demonstrated at least 50%

inhibition of mitochondrial Ca^{2+} uptake at a concentration of 10 μM (Fig 1*C*; *SI*

Appendix, Fig. S1*B*). Among them, Berberine exhibited the highest inhibitory effect,

reaching up to 87% (*SI Appendix*, Fig. S1*B*). To rule out the possibility that the

observed effect was a secondary consequence of mitochondrial membrane potential

disruption, the major driving force for Ca^{2+} uptake within the organelle (14, 15), we

performed TMRM staining analysis. The mitochondrial decoupler, CCCP, induced a

decrease in TMRM intensity. All tested drugs, except for Berberine, including

Trifarotene, Eltrombopag, and Tipranavir, significantly decreased the mitochondrial

membrane potential (Fig. 1*D*). Therefore, this screening process identified Berberine,

a natural compound derived from medicinal herbs, as a promising candidate for

further investigation as MCU inhibitor.

Berberine inhibits mitochondrial Ca^{2+} uptake

Given that MCU is a mitochondrial inner membrane protein, a suitable MCU inhibitor must traverse at least two membrane layers, the cellular membrane, and the mitochondrial outer membrane. Previous studies have documented the subcellular localization of Berberine within mitochondria (16, 17). Exploiting Berberine's autofluorescence, we observed its cellular penetration at 10 μM, co-localizing with MitoTracker stain (*SI Appendix*, Fig. S2 *A* and *B*). This finding is consistent with previous studies (18) and leads us to postulate that Berberine is capable of crossing intact cellular membranes to inhibit mitochondrial Ca^{2+} uptake.

To substantiate our hypothesis, we evaluated Berberine's inhibitory effects in intact cells by initiating intracellular Ca^{2+} signaling with histamine stimulation while monitoring cytosolic and mitochondrial Ca^{2+} dynamics, simultaneously. Berberine specifically inhibits mitochondrial Ca^{2+} uptake, as evidenced by green fluorescent indicator, 4mt-GCaMP6, without affecting histamine-induced cytosolic Ca^{2+} signals, as shown by the red fluorescent dye, Calbryte590 (Fig. 2 *A-E*). Similar results were obtained using other Ca^{2+} indicators, cyto-GCaMP6 or 4mt-GCaMP8 in Hela cells (*SI Appendix*, Fig. S2 *C-F*). Next, we assessed the mitochondrial Ca^{2+} uptake in cells permeabilized with digitonin. Under these experimental conditions, addition of CaCl$_2$ rapidly elevated cytosolic Ca^{2+} levels and induced a rapid Ca^{2+} uptake by mitochondria. Berberine showed a potent inhibition of mitochondrial Ca^{2+} uptake, in a dose-dependent manner (Fig. 2*F*). The IC$_{50}$ was determined to be 2.202 μM (Fig. 2*G*). These results collectively suggest that Berberine's inhibitory action within cells is likely due to its direct interaction with the MCU complex, without perturbing the cytoplasmic Ca^{2+} ion signaling.

To evaluate the safety of Berberine in cells, we treated the cells with increasing amount of Berberine, and confirmed that up to 10 μM Berberine treatment did not affect mitochondrial membrane potential by TMRM staining (*SI Appendix*, Fig. S2 *G* and *H*). We further investigated the cytotoxicity of Berberine to understand its

potential impact on cell growth. It showed that at lower doses, specifically between 5-

10 μM, which are sufficient to inhibit MCU, Berberine exhibited minimal cytotoxic

effects (*SI Appendix*, Fig. S2*I*). This indicates that Berberine is an effective MCU

inhibitor with low cytotoxicity, making it a relative safe option at these concentrations.

Direct binding of Berberine to the JML domain of MCU

To elucidate the mechanism underlying the inhibitory effect of Berberine on

MCU, we synthesized biotin-conjugated Berberine (Biotin-BBR) (Fig. 3*A*). The

Biotin-BBR exhibited similar mitochondrial localization and inhibition efficiency as

Berberine (Fig. 3*A*; *SI Appendix*, Fig. S3*A* and *B*), suggesting that the Biotin-BBR

retains the ability of interacting with the same molecular targets as the unmodified

Berberine. Molecular docking screening results suggest that BBR may directly bind to

the MCU monomer (Fig. 1 *A* and *B*). Consistent with these results, Biotin pull-down

assays demonstrated that Biotin-BBR could interact with both exogenous and

endogenous MCU in cell lysate (Fig. 3*B*; *SI Appendix*, Fig. S3*C*). *In vitro* pull-down

assays further confirmed that BBR directly binds to recombinant glutathione-S-

transferase (GST)-MCU (Fig. 3*C*). Notably, this binding was diminished when

unlabeled BBR competitively interacted with GST-MCU, indicating a consistent

binding site and confirming the specificity of the binding (Fig. 3*C*). Microscale

thermophoresis (MST) analysis further confirmed that the affinity between GST-MCU

and Berberine was ~8 μM (Fig. 3*D*).

Next, to explore the mechanism of Berberine's binding to MCU and its

functional impact, we first mapped MCU regions interacting with Berberine. We *in*

vitro translated MCU truncations based on its secondary structure (19), excluding the

mitochondrial targeting sequence (residues 1-74) due to its non-conservative nature

(20). We assessed their interactions with Berberine using Biotin pull-down assays.

Berberine showed a strong affinity for the MCU juxtamembrane loop (JML) segment

spanning residues 275-295 (Fig. 3 *E* and *F*), although minor binding to other areas

was also observed (Fig. 3*E*). This observation were consistent with molecular docking

simulations (Fig. 3*G*), which indicated that Berberine binds near JML domain, a

region critical for EMRE-dependent MCU opening (19, 21). Docking analyses

pinpointed key binding residues, including Tyr 281, Tyr 289, Tyr 291, and Ala 294

(Fig. 3*G*; *SI Appendix*, Fig. S3*D*). Notably, Tyr 281 and Tyr 289 are likely pivotal for

hydrogen bonds with Berberine as shown by molecular docking simulations (Fig. 3*G*;

SI Appendix, Fig. S3*D*).

To explore the interaction of Berberine with predicted binding sites within the

MCU, we performed site-directed mutagenesis. Sequence alignment of the JML

domain revealed four conserved residues (Y281, Y289, Y291, and A294) across

various MCU homologs (*SI Appendix*, Fig. S3*E*), known to play a role in maintaining

the open conformation of the MCU (21). It was reported that mutations of Y289 and

Y291 to alanine, as well as A294 to phenylalanine, do not alter the open conformation

of MCU and preserve robust mitochondrial Ca^{2+} uptake capacity. However, the

critical residue Y281, when mutated to alanine, induces a closed conformation of the

MCU (21). Under these circumstances, simultaneous mutations of Y281A, Y289A,

Y291A, and A294F significantly disrupted the interaction with the Berberine-MCU

JML domain (Fig. 3*G*), confirming Berberine's direct binding to these residues. To

determine whether the ability of Berberine to bind was independent of the MCU's

conformational state (open or closed), we introduced mutations Y281H and Y281R,

which maintain the open conformation (21) but disrupt potential hydrogen bonding

with Berberine. These mutations still abolished Berberine binding (Fig. 3*H*), negating

the hypothesis that Berberine's binding is confined to the open conformation of MCU.

Berberine disrupts MCU-EMRE interaction

Ca^{2+} transfer through the mitochondrial inner membrane is mediated by the

mitochondrial calcium uniporter (MCU) complex, consisting of MCU, EMRE,

MICU1, and MICU2 (3). We investigated Berberine's influence on the MCU complex

assembly via co-immunoprecipitation (IP) experiments. The components of MCU

complex, including MCU, EMRE, MICU1 and MICU2, were SNAP-tagged and co-

expressed with Flag-tagged MCU in 293T cells. Following Berberine treatment, only

the interaction between SNAP-tagged EMRE and Flag-tagged MCU was significantly

diminished, while the interactions of MCU with other MCU components remained

unaltered (Fig. 4*A*). These results suggest that Berberine selectively disrupts the

EMRE-MCU interaction, which was further confirmed by a reverse IP assay (*SI*

Appendix, Fig. 4*A*).

To directly assess Berberine's effect on MCU-EMRE interaction, we employed

in vitro binding assays. These assays confirmed Berberine's ability to directly interfere

with MCU-EMRE assembly (Fig. 4*B*). To understand the mechanism behind this

disruption, we conducted molecular dynamics simulations. The simulations revealed

that the MCU-EMRE complex rapidly reached equilibrium and maintained stability

during the simulation in the absence of Berberine. At 100 ns, the Rg value, a measure

of compactness, significantly increased from 3.734 nm to 4.982 nm upon Berberine

addition (Fig. 4*C*), indicating a more spread-out and less compact complex structure.

This observation was further corroborated by the increased RMSD values (reflecting

average atomic displacement) in the presence of Berberine (Fig. 4*D*). Together, these

results suggest that Berberine induces a looser and less stable complex structure (*SI*

Appendix, Fig. S4*B*). Furthermore, hydrogen bond analysis revealed a reduction in the

average number of bonds between MCU and EMRE from 10.540 to 6.287 upon

Berberine treatment (*SI Appendix*, Fig. S4 *C* and *D*), signifying a weakening of

interactions between the two proteins. This finding aligns with the significant

decrease in binding energy observed between MCU and EMRE in the presence of

Berberine (from -100.02 ± 17.85 kcal/mol to -29.52 ± 27.07 kcal/mol, *SI Appendix*,

Fig. S4*E*). Collectively, these data strongly support the conclusion that Berberine

disrupts the MCU-EMRE complex by weakening the interactions between MCU and

EMRE.

Berberine inhibits Ca^{2+} uptake by disrupting MCU-EMRE interaction

To investigate whether Berberine affects mitochondrial Ca^{2+} uptake by disrupting

the MCU-EMRE interaction, we need to identify an MCU mutant that resists Berberine binding while retaining EMRE binding capability and preserving normal mitochondrial Ca^{2+} uptake capacity. Previous studies have identified several MCU mutations that prevent Berberine binding (Fig. 3*H*). One of these mutations, MCU^{AAAF}, also eliminates EMRE binding and mitochondrial Ca^{2+} uptake capacity (21). Therefore, we selected the MCU^{HAAF} mutant for further investigation. Similar to the wild-type MCU (MCU^{WT}), the MCU^{HAAF} mutant robustly binds to EMRE. Howerver, unlike MCU^{WT}, the binding of MCU^{HAAF} to EMRE was not affected by Berberine (Fig. 4*E*). This suggests that the MCU^{HAAF} mutation does not interfere with the EMRE-binding interface of MCU.

By establishing cell lines stably reconstituted either MCU^{WT} and MCU^{HAAF} (Fig. 4*F*), we can compare the effects of Berberine on mitochondrial Ca^{2+} uptake in the two contexts. As expected, Berberine treatment reduced the interaction between MCU^{WT} and EMRE (Fig. 4*E*) and lowered mitochondrial Ca^{2+} uptake in both control and MCU^{WT} -reconstituted cells (Fig. 4 *G* and *H*). This confirms that Berberine disrupts the MCU-EMRE complex and impairs mitochondrial Ca^{2+} transport in cells expressing the wild-type MCU. In contrast, the MCU^{HAAF} mutant, which failed to bind to Berberine (Fig. 3*H*), preserved its mitochondrial Ca^{2+} uptake capabilities even after Berberine exposure (Fig. 4 *G* and *H*). These findings indicate that Berberine's modulation of mitochondrial Ca^{2+} uptake relies on its role in binding to MCU, subsequently disrupting the MCU-EMRE complex assembly and ultimately leading to the closure of the MCU channel, thereby preventing Ca^{2+} entry into mitochondria (Fig. 4*I*).

Berberine suppresses mitochondrial Ca^{2+} overload and protects against myocardial ischemia-reperfusion injury

We next sought to explore the physiological relevance of our findings. It's well established that excessive Ca^{2+} entry into mitochondria can lead to mitochondrial Ca^{2+} overload. This overload subsequently triggers the opening of the permeability

transition pore (PTP), causing mitochondrial swelling, and eventually, cell death (22).

Cardiac-specific conditional knockout of MCU could protect mitochondria by

preventing the opening of mPTP *in vitro* and *in vivo* (10, 11). The mPTP opening can

be monitored by measuring mitochondrial absorbance (23). In an *in vitro* context, we

observed that Berberine inhibited Ca^{2+} overload-induced mitochondrial swelling (Fig.

5*A*). This effect paralleled those of Ru360 and cyclosporin A (CsA) (24), a known

inhibitor of mPTP opening, suggesting that Berberine might confer cellular protection

under conditions of mitochondrial Ca^{2+} overload.

To validate this hypothesis, we first established a model of hypoxia/reoxygenation

(H/R) injury in neonatal rat cardiomyocytes and embryonic rat cardiomyocyte-derived

H9c2 cell line. Consistent with previous studies (25, 26), H/R injury led to a

significantly elevated mitochondrial Ca^{2+} levels, indicated by fluorescent intensity of

4mtGCaMP6 or Rhod2 (Fig. 5 *B* and *C*; *SI Appendix*, Fig. S5). Furthermore,

Berberine pretreatment alleviated this mitochondrial Ca^{2+} overload (Fig. 5 *B* and *C*; *SI*

Appendix, Fig. S5). To assess the potential cardioprotective effect of Berberine *in vivo*,

a mouse acute cardiac ischemia/reperfusion (I/R) model was established (Fig. 5*D*).

Pretreatment of Berberine (24h plus 15 min before reperfusion) significantly reduced

I/R induced myocardial infarct size (Fig. 5 *E* and *F*). Additionally, Berberine

ameliorated cardiomyocyte death, evidenced by decreased cTNl and LDH

concentration (Fig. 5 *G* and *H*). TUNEL staining experiments further confirmed the

protective effect of Berberine against myocadiac I/R injury (Fig. 5*I*). In summary,

these data indicate that Berberine exhibits therapeutic effects in myocardial

ischemia/reperfusion injury via counteracting mitochondrial Ca^{2+} overload.

Discussion:

Our virtual screening approach using an FDA-approved drug library identified Berberine, a natural alkaloid derived from medicinal plants, as a potent MCU inhibitor. Berberine exerts its inhibitory effect by directly binding to the MCU and disrupting the assembly of MCU-EMRE complex, consequently hindering Ca^{2+} entry into mitochondrial matrix. Notably, Berberine significantly reduces mitochondrial Ca^{2+} overload, thereby protecting cardiomyocytes from ischemia-reperfusion (I/R)-induced apoptosis *in vitro* and *in vivo*. These results underscore the potential of Berberine as a promising MCU-inhibiting drug for clinical application.

Berberine, a well-characterized plant alkaloid with a long history of traditional use, has been employed for treating various conditions, including diabetes, hyperlipidemia, and cardiovascular diseases (27). Its antimicrobial properties are also well-documented (28). Despite its extensive therapeutic application, Berberine's precise molecular targets remains elusive. Our study provides compelling evidence that Berberine targets MCU, effectively impeding mitochondrial Ca^{2+} uptake through direct binding. Importantly, Berberine may overcome limitations observed in existing MCU inhibitors such as Ru360, DS16570511 (8), mitoxantrone (9), and MCU-i11 (29)—a specific MICU1-targeting inhibitor—in animal experiments. DS16570511, for instance, exhibits off-target effects on mitochondrial membrane potential, thereby limiting its application (30). Similarly, mitoxantrone, a topoisomerase inhibitor, is known to have cardiotoxic effects (31). Berberine's well-established safety profile *in vivo* positions it as a strong candidate for further exploration in clinical settings.

Our studies, along with previous studies, consistently demonstrate Berberine's capacity to provide significant cardioprotection in the context of myocardial I/R injury (32). Clinical studies further support its potential, indicating that Berberine can modulate inflammatory markers in patients suffering from acute myocardial infarction (33). Berberine's protective effects are multifaceted, encompassing antioxidant, anti-inflammatory, and anti-apoptotic activities. A key aspect of this protection is

Berberine's ability to maintain mitochondrial integrity and high activity of the

respiratory complexes (34-36), as evidenced by its regulation of Bcl-2 and Bax

proteins, which are crucial for mitochondrial membrane stability and the prevention of

apoptotic factor release (34, 37). Our findings establish MCU as a key cardiac target

of Berberine, revealing a core mechanism for its protective effects in I/R injury. Given

that mitochondrial Ca^{2+} overload occurring rapidly within the initial 20 minutes post-

ischemia (26), early intervention with Berberine is of paramount importance.

Furthermore, Berberine's potential interactions with other targets and its slower-acting

mechanisms warrant further investigation to fully understand its therapeutic potential.

Our findings reveal that Berberine binds directly to the MCU near the

juxtamembrane loop (JML) domain, at residues Y281, Y289, Y291, and A294. The

JML is essential for channel gate formation and its interaction with EMRE is critical

for Ca^{2+} channel activation (19). Mutational analysis of Y281 and Y289 highlights

their significance in MCU-EMRE assembly and the EMRE-mediated conformational

opening of MCU (21). Y281A mutants significantly disrupts EMRE binding (21),

paralleling the loss of Ca^{2+} uptake observed with Berberine treatment. Notably, Y289,

a key hydrophobic residue within the JML, facilitates EMRE-dependent MCU

opening (21). Substituting Y289 with hydrophilic residues impairs channel opening,

underscoring the critical role of hydrophobic interactions at this site for EMRE

binding and MCU activation. Berberine's hydrophobic surface (38) might promote its

integration near the JML, disrupting interactions between critical residues like Y281

and Y289, and fostering a closed MCU conformation unfavorable to EMRE

association. While our molecular dynamics simulations support this hypothesis, the

complex mechanisms of MCU gating warrant further structural studies for full

elucidation.

In conclusion, our study identifies Berberine as a novel MCU inhibitor with

significant potential for clinical applications. Revealing its inhibitory effects on MCU

and providing insights into the complex Berberine-MCU-EMRE dynamics, our

findings lay a solid foundation for the development of targeted drug strategies aimed

335 at mitochondrial Ca^{2+} regulation. Structural analogs of Berberine could be engineered

336 for enhanced specificity and safety as MCU inhibitors. Furthermore, the role of MCU

337 dysfunction is becoming increasingly understood, not only in cardiac

338 ischemia/reperfusion injury but also for other diseases, such as the recently reported

339 dependency of venetoclax-resistant leukemic stem cells (LSCs) on mitochondrial

340 calcium for survival(39). These findings position Berberine as a valuable starting

341 point for developing novel interventions in diseases associated with dysregulated

342 mitochondrial Ca^{2+} homeostasis.

344   **Reference:**

345   1.   Kamer KJ & Mootha VK (2015) The molecular era of the mitochondrial calcium

346   uniporter. *Nat Rev Mol Cell Biol* 16(9):545-553.

347   2.   Giorgi C, Marchi S, & Pinton P (2018) The machineries, regulation and cellular

348   functions of mitochondrial calcium. *Nat Rev Mol Cell Biol* 19(11):713-730.

349   3.   Murphy E & Steenbergen C (2021) Regulation of Mitochondrial Ca(2+) Uptake. *Annu*

350   *Rev Physiol* 83:107-126.

351   4.   Bernardi P, Rasola A, Forte M, & Lippe G (2015) The Mitochondrial Permeability

352   Transition Pore: Channel Formation by F-ATP Synthase, Integration in Signal

353   Transduction, and Role in Pathophysiology. *Physiol Rev* 95(4):1111-1155.

354   5.   Bauer TM & Murphy E (2020) Role of Mitochondrial Calcium and the Permeability

355   Transition Pore in Regulating Cell Death. *Circ Res* 126(2):280-293.

356   6.   Moore CL (1971) Specific inhibition of mitochondrial Ca++ transport by ruthenium red.

357   *Biochem Biophys Res Commun* 42(2):298-305.

358   7.   Ying WL, Emerson J, Clarke MJ, & Sanadi DR (1991) Inhibition of mitochondrial

359   calcium ion transport by an oxo-bridged dinuclear ruthenium ammine complex.

360   *Biochemistry* 30(20):4949-4952.

361   8.   Kon N, *et al.* (2017) DS16570511 is a small-molecule inhibitor of the mitochondrial

362   calcium uniporter. *Cell Death Discov* 3:17045.

363   9.   Arduino DM, *et al.* (2017) Systematic Identification of MCU Modulators by Orthogonal

364   Interspecies Chemical Screening. *Mol Cell* 67(4):711-723 e717.

10.   Luongo TS, *et al.* (2015) The Mitochondrial Calcium Uniporter Matches Energetic

Supply with Cardiac Workload during Stress and Modulates Permeability Transition.

Cell Rep 12(1):23-34.

11.   Kwong JQ, *et al.* (2015) The Mitochondrial Calcium Uniporter Selectively Matches

Metabolic Output to Acute Contractile Stress in the Heart. *Cell Rep* 12(1):15-22.

12.   Lambert JP, *et al.* (2019) MCUB Regulates the Molecular Composition of the

Mitochondrial Calcium Uniporter Channel to Limit Mitochondrial Calcium Overload

During Stress. *Circulation* 140(21):1720-1733.

13.   Zhao H, *et al.* (2019) AMPK-mediated activation of MCU stimulates mitochondrial

Ca(2+) entry to promote mitotic progression. *Nat Cell Biol* 21(4):476-486.

14.   Baughman JM, *et al.* (2011) Integrative genomics identifies MCU as an essential

component of the mitochondrial calcium uniporter. *Nature* 476(7360):341-345.

15.   De Stefani D, Raffaello A, Teardo E, Szabo I, & Rizzuto R (2011) A forty-kilodalton

protein of the inner membrane is the mitochondrial calcium uniporter. *Nature*

476(7360):336-340.

16.   Agnarelli A, *et al.* (2018) Cell-specific pattern of berberine pleiotropic effects on

different human cell lines. *Sci Rep* 8(1):10599.

17.   Jin M, *et al.* (2020) Synthesis of a novel fluorescent berberine derivative convenient

for its subcellular localization study. *Bioorg Chem* 101:104021.

18.   Pereira GC, *et al.* (2007) Mitochondrially targeted effects of berberine [Natural Yellow

18, 5,6-dihydro-9,10-dimethoxybenzo(g)-1,3-benzodioxolo(5,6-a) quinolizinium] on

K1735-M2 mouse melanoma cells: comparison with direct effects on isolated

mitochondrial fractions. *J Pharmacol Exp Ther* 323(2):636-649.

19.   Wang Y, *et al.* (2019) Structural Mechanism of EMRE-Dependent Gating of the

Human Mitochondrial Calcium Uniporter. *Cell* 177(5):1252-1261 e1213.

20.   Lee SK, *et al.* (2016) Structural Insights into Mitochondrial Calcium Uniporter

Regulation by Divalent Cations. *Cell Chem Biol* 23(9):1157-1169.

21.   Van Keuren AM, *et al.* (2020) Mechanisms of EMRE-Dependent MCU Opening in the

Mitochondrial Calcium Uniporter Complex. *Cell Rep* 33(10):108486.

22.   Rasola A & Bernardi P (2011) Mitochondrial permeability transition in Ca(2+)-

dependent apoptosis and necrosis. *Cell Calcium* 50(3):222-233.

23.   Blattner JR, He L, & Lemasters JJ (2001) Screening assays for the mitochondrial

permeability transition using a fluorescence multiwell plate reader. *Anal Biochem*

295(2):220-226.

24.   Pan X, *et al.* (2013) The physiological role of mitochondrial calcium revealed by mice

lacking the mitochondrial calcium uniporter. *Nat Cell Biol* 15(12):1464-1472.

25.   Paillard M, *et al.* (2013) Depressing mitochondria-reticulum interactions protects

cardiomyocytes from lethal hypoxia-reoxygenation injury. *Circulation* 128(14):1555-

1565.

26.   Petersen CE, *et al.* (2023) Increased mitochondrial free Ca(2+) during ischemia is

suppressed, but not eliminated by, germline deletion of the mitochondrial Ca(2+)

uniporter. *Cell Rep* 42(7):112735.

27.   Gasmi A, *et al.* (2024) Berberine: Pharmacological Features in Health, Disease and

Aging. *Curr Med Chem* 31(10):1214-1234.

28.   Liang J, Huang X, & Ma G (2022) Antimicrobial activities and mechanisms of extract

and components of herbs in East Asia. *RSC Adv* 12(45):29197-29213.

29.   Di Marco G, *et al.* (2020) A High-Throughput Screening Identifies MICU1 Targeting

Compounds. *Cell Rep* 30(7):2321-2331 e2326.

30.   Payne R, Li C, Fernandez-Garcia E, Vais H, & Foskett K (2019) The MCU Inhibitor

Ds16570511 has Off-Target Effects on Mitochondrial Membrane Potential.

Biophysical Journal 116(3).

31.   Sawyer DB, Peng X, Chen B, Pentassuglia L, & Lim CC (2010) Mechanisms of

anthracycline cardiac injury: can we identify strategies for cardioprotection? *Prog*

Cardiovasc Dis 53(2):105-113.

32.   Liu DQ, *et al.* (2019) Berberine protects against ischemia-reperfusion injury: A review

of evidence from animal models and clinical studies. *Pharmacol Res* 148:104385.

33.   Qing Y, Dong X, Hongli L, & Yanhui L (2018) Berberine promoted myocardial

protection of postoperative patients through regulating myocardial autophagy. *Biomed*

Pharmacother 105:1050-1053.

34.   Wang Y, Liu J, Ma A, & Chen Y (2015) Cardioprotective effect of berberine against

myocardial ischemia/reperfusion injury via attenuating mitochondrial dysfunction and

apoptosis. *Int J Clin Exp Med* 8(8):14513-14519.

35.   Hu F, *et al.* (2024) Berberine inhibits excessive autophagy and protects myocardium

against ischemia/reperfusion injury via the RhoE/AMPK pathway. *Int J Mol Med* 53(5).

36.   Ding L, *et al.* (2021) Berberine improves dietary-induced cardiac remodeling by

upregulating Kruppel-like factor 4-dependent mitochondrial function. *Biol Chem*

402(7):795-803.

37.   Zhao GL, *et al.* (2016) Berberine protects rat heart from ischemia/reperfusion injury

433   via activating JAK2/STAT3 signaling and attenuating endoplasmic reticulum stress.

434   *Acta Pharmacol Sin* 37(3):354-367.

435   38.   Chu M, *et al.* (2018) Polypharmacology of Berberine Based on Multi-Target Binding

436   Motifs. *Front Pharmacol* 9:801.

437   39.   Sheth AI, *et al.* (2024) Targeting Acute Myeloid Leukemia Stem Cells Through

438   Perturbation of Mitochondrial Calcium. *Cancer Discov.*

Acknowledgments

The authors thank D.D. Stefani for providing 4mt-GCaMP6 plasmids, and K. Wang and X. Xu for assistance with microscopy assays. This work was supported by the National Natural Science Foundation of China (grant numbers 82025028, 92354304 and 82000107), the National Key R&D Program of China (2021YFA1300203 and 2020YFA0113300), China Postdoctoral Science Foundation (2022M723897).

Author Contributions

Xin Pan conceived the study. Haixin Zhao designed and carried out most of the experiments with the assistance from Siqi Chen, Nian Cao, and Wenjun Wu. Haixin Zhao, Siqi Chen, Guangqin Liu and Haizhen Zhu performed calcium measurements and biochemical experiments. Haixin Zhao, Nian Cao and Jun Gao contributed to mouse studies. Haixin Zhao and Wenjun Wu performed Screening and Data Analysis. Dingyi Lu and Lingmin Zeng performed immunostaining and flow cytometry analysis. Jiayi Chen, Teng Li, Ting Li, Weina Zhang, Qing Xia, Tao Zhou, Ai-Ling Li, Xue-Min Zhang, and Xin Pan contributed to interpreting the results. Xin Pan and Ai-Ling Li supervised the research. Xin Pan and Haixin Zhao wrote the paper.

Declaration of Interests

The authors declare no competing interests.

Supplementary Materials:

Materials and Methods

Supplementary Figures

Supplementary Figure Legends

References (40-42)

Figure Legends

Fig. 1. The discovery of Berberine as a novel MCU inhibitor

(A) Schematic of screening workflow.

(B) Histogram plot of the 2,816 FDA-approved drug library based on the free-energy

docking score with MCU.

(C) Graphical representation of drug screen results. 'Z-score' ranks drugs based on

maximal amplitudes of the traces in Supplementary Fig 1a.

(D) Measurement of mitochondrial membrane potential indicated by TMRM

fluorescence intensity. Data are shown as the mean ± s.e.m. n = 50 cells for each

condition. *P* value was analyzed using one-way ANOVA.

Fig. 2. Berberine inhibits mitochondrial Ca^{2+} uptake.

(A) Snapshot of cytosolic (left, red) and mitochondrial (right, green) Ca^{2+} transient

images from time-lapse movies of representative HeLa cells stained with

Calbryte590 and 4mtGCaMP6 following 10 μM Histamine (His) stimulation.

Cells were pretreated with various concentrations of Berberine. The time

displayed on the images is in seconds. Scale bar, 50 μm.

(B) Representative traces of mitochondrial Ca^{2+} level (indicated by 4mtGCaMP6) in

HeLa cells stimulated with 10 μM Histamine and pretreated with different

concentrations of Berberine. Data are shown as mean ± s.e.m. n = 40 cells for

each condition.

(C) Measurement of the maximal amplitudes of the mitochondrial Ca^{2+} traces in **B**.

Data are shown as the mean ± s.e.m. n = 40 cells for each condition. *P* value was

analyzed using one-way ANOVA.

(D) Representative traces of cytosolic Ca^{2+} level (indicated by Calbryte590) in HeLa

cells stimulated with 10 μM Histamine and pretreated with different

concentrations of Berberine. Data are shown as the mean ± s.e.m. n = 40 cells for

each condition.

(E) Measurement of the maximal amplitudes of the mitochondrial Ca^{2+} traces in **D**.

Data are shown as the mean ± s.e.m. n = 40 cells for each condition. *P* value was

analyzed using one-way ANOVA.

(F) Traces of mitochondrial Ca^{2+} level (indicated by fluorescence of 4mt-GCaMP6) in

digitonin-permeabilized HeLa cells upon the addition of Ca^{2+} under varying

concentrations of Berberine.

(G) Dose-dependent inhibition of mitochondrial Ca^{2+} uptake activity by Berberine

with an IC_{50} of 2.202 μM.

Fig. 3. Berberine directly targets the JML domain of MCU.

(A) Upper panel: Chemical structure of Biotin-Berberine. Lower panel:

Representative immunofluorescent images of HeLa cells treated with Biotin-

Berberine (green) and stained for mitochondria (COXIV, red). Scale bar, 10 μm.

(B) Western blot analysis from Biotin-Berberine pulldown assay using HEK293T cell

lysates. Cells were treated with Biotin-Berberine (10 μM, 2 h).

(C) Western Blot analysis from *in vitro* pulldown assay investigating interactions

between GST-MCU and Biotin-Berberine under varying concentrations of

Berberine.

(D) Binding ability of Berberine to GST-MCU in MST assays.

(E) Schematic representation of MCU truncation mutants utilized for domain-

mapping studies. Numerical values in parentheses correspond to the amino acid

sequences encompassed by each construct. Key structural domains are labeled:

NTD (N-terminal domain), CC (coiled-coil domain), OJMH (outer

juxtamembrane helix), TM (transmembrane helix), DIME (DIME motif, Ca^{2+}

filter), JML (juxtamembrane loop), and CTD (C-terminal domain).

(F) Western blot analysis of Biotin-Berberine pulldown assays using *in vitro*

transcribed and translated SNAP-tagged MCU truncation mutants as shown in **E**.

(G) Molecular docking model revealing the interaction between human MCU and

Berberine.

(H) Western blot analysis of biotin-berberine pulldown assays using *in vitro*

transcribed and translated SNAP-tagged MCU mutants (amino acids 275-295,

encompassing the JML region). Wild-type residues Y281, Y289, Y291, and A294

were mutated to AAAF, HAAF, and RAAF, respectively.

Fig. 4. Berberine inhibits Ca^{2+} uptake by disrupting MCU-EMRE complex

assembly.

(A) Western blot analysis depicting the co-immunoprecipitation of Flag-tagged MCU

with SNAP-tagged EMRE, MCU, MICU1, and MICU2 in HEK293T cells,

conducted with or without the presence of 10 μM Berberine for 2 hours.

(B) Western blot analysis of GST pulldown assays performed with *in vitro* transcribed

and translated Flag-tagged EMRE and recombinant GST-MCU, with and without

the addition of 2 μg Berberine.

(C) Molecular dynamics simulations showing the radius of gyration (Rg) of the MCU-

EMRE complex in the presence and absence of Berberine.

(D) Molecular dynamics simulations depicting the root-mean-square deviation

(RMSD) of the MCU-EMRE complex with and without Berberine.

(E) Western blot analysis of the co-precipitation of Flag-tagged MCU wild-type (WT)

or Flag-tagged MCU^{HAAF} mutant with SNAP-EMRE in HEK293T cells, treated

with 10 μM Berberine for 2 hours or untreated.

(F) Western blot analysis of endogenous MCU and exogenous MCU-mCherry

expression in HeLa stable cell lines with MCU wild-type (WT) and MCU^{HAAF}

mutants.

(G) Traces of mitochondrial Ca^{2+} levels, as indicated by the fluorescence intensity of

4mt-GCaMP6, in digitonin-permeabilized HeLa cells following Ca^{2+} addition.

The cells include control HeLa, and stable cell lines expressing mcherry-tagged

MCU wild-type (WT) and MCU^{HAAF} mutants, under conditions with and without

10 μM Berberine.

(H) Measurement of the maximal amplitudes of the mitochondrial Ca^{2+} traces in **G**.

Data are shown as the mean ± s.e.m. n = 45 cells for each condition. *P* value was

analyzed using two-way ANOVA.

(I) Proposed model for Berberine-mediated MCU gating. Left panel, EMRE binds to

MCU and stabilizes the luminal gate of MCU in the open conformation. Right

panel, Berberine integrates into the MCU, inducing a closed conformation that

disrupts the assembly of the MCU-EMRE complex and seals the MCU's luminal

gate.

Fig. 5. Berberine protects against myocardial ischemia/reperfusion injury

through inhibiting mitochondrial Ca^{2+} overload.

(A) Mitochondrial swelling in response to Ca^{2+} challenge (500 μM CaCl$_2$). Swelling is

inhibited with Berberine (10 μM), Ru360 (3 μM) or CsA (3 μM).

(B) Representative images of neonatal rat cardiomyocytes expressing 4mtGCaMP6.

Cells were either treated with 10 μM Berberine or not, then subjected to hypoxia

(3 h) followed by reoxygenation (12 h) (H/R). Fluorescence at ex488 (green)

represents the Ca^{2+}-dependent signal, while fluorescence at ex390 (blue) indicates

the Ca^{2+}-independent signal. The ratio (pseudocolour) of ex488/ex390 indicates

the level of mitochondrial Ca^{2+}. Scale bar, 10 μm.

(C) Measurement of mitochondrial Ca^{2+} level indicated by 4mtGCaMP6 fluorescence

ratio (ex488/ex405) in **B**. Data are shown as the mean ± s.e.m. n = 50 cells for

each condition. *P* value was analyzed using one-way ANOVA.

(D) The experimental scheme of the mice with acute I/R injury (45 minutes ischemia

followed by 24 hours reperfusion) with vehicle or Berberine pretreatment (24

hours and 15 minutes before the onset of ischemia).

(E) Evans Blue and 2, 3, 5-triphenyltetrazolium chloride (TTC) staining of

myocardium after I/R injury. The blue area indicates non-ischemic area. The red

indicates viable tissue at risk. The white indicates infarct area.

(F) Quantification of infarct size (IF), area at risk (AAR) of mice subjected to acute

I/R injury (as protocols in **D**). n = 9 or 10 mice.

(G-I). Cardiac cell death indexed by serum cTnI (**G**, n=12 [sham], 6 [sham+BBR], 11

[I/R], and 12 [I/R+BBR]), serum LDH concentration (**H**, n=13 [sham], 6

[sham+BBR], 11 [I/R], and 12 [I/R+BBR]), and myocardial TUNEL-positive cells (**I**,

n=8 [sham], 6 [sham+BBR], 11 [I/R], and 12 [I/R+BBR]) of the mice subjected to I/R

injury (as the protocols in **D**) with vehicle or Berberine treatment (5 mg/kg, ip.24 h

and 15min before I/R injury). *P* value was analyzed using one-way ANOVA. Scale

bar, 50 μm.

Fig. 1. The discovery of Berberine as a novel MCU inhibitor.

Fig. 2. Berberine inhibits mitochondrial Ca²⁺ uptake.

Fig. 3. Berberine directly targets the JML domain of MCU.

Fig. 4. Berberine inhibits Ca²⁺ uptake by disrupting MCU-EMRE complex assembly.

Fig. 5. Berberine protects against myocardial ischemia/reperfusion injury through inhibiting mitochondrial Ca²⁺ overload.

Supplementary Materials

Materials and Methods

Experimental animals

All animals used in this study were maintained in the pathogen-free barrier animal facility at the National Center of Biomedical Analysis. Animal care was monitored daily by certified veterinary staff and laboratory personnel. 10-week-old mice in the C57BL/6J background and pregnant Sprague-Dawley rats were purchased from the Beijing Vital River Laboratory Animal Technology Company.

Murine model of myocardial ischemia

The 10 week-male-C57BL/6 mice (>22 g) were randomly assigned to specific groups. Berberine (5 mg/kg) or vehicle was administered intraperitoneally twice (24 hours and 15 minutes prior) to ischemia-reperfusion (I/R) surgery as described (1). Briefly, pentobarbital (70 mg/kg i.p.) was used to induce and maintain anesthesia. Mice were then ventilated on a Harvard rodent respirator via a tracheostomy. A midline sternotomy was performed, with a reversible snare occlude placed around the left anterior descending coronary artery. I/R injury was initiated by tightening the snare for 45 minutes, then releasing it. After reperfusion for 24 hours, the mice were sacrificed for further detection and measurement.

For serum LDH concentration and cTnI level, blood samples were centrifuged at 3,000 rpm for 10 min and then detected using an LDH kit (Sigma-Aldrich, Cat# MAK066) and a mouse cardiac troponin I ELISA kit (ThermoFisher, Cat#EEL112) according to the manufacturer's instructions.

For the measurement of the area-at-risk (AAR) and infarct size, the heart was initially excised post-reocclusion of the coronary artery at the previous occlusion site. The heart ascending aorta was cannulated (distal to the sinus of Valsalva) and then perfused retrogradely with 0.05% Alcian blue to delineate the AAR. Following AAR visualization, the heart was frozen at -80 ℃ for 5 min, cut into ~1 mm slices, and the

unstained infarcted (IF) region was lucifugal visualized using 1% 2,3,5-triphenyl-

tetrazolium chloride at 37 ℃ for 15 min. Infarct and left ventricular areas were

determined by planimetry with Image J. The infarct size was calculated as infarct area

divided by area at risk (IF/AAR).

Molecular Docking Screening

The 3D structures of MCU (PDB: 6XJV) were retrieved from the Protein Data

Bank. A set of 2,816 FDA-approved drugs were converted to 3D structures using Open

Babel. Preparation and parameterization of the receptor protein and ligands were

conducted with AutoDock Tools (ADT3). Docking grid documents were created using

AutoGrid of sitemap, and docking simulation were performed with AutoDock Vina

(1.2.0) (2). The score was ranked according to the value of the binding energy of

protein-ligand complex. Finally, the protein-ligand interaction figure was generated by

PyMOL.

Cell culture

To establish HeLa cells stably expressing 4mt-GCaMP6s, we constructed a

lentiviral vector (pLVX-puro) containing the 4mt-GCaMP6s insert. This vector was

transfected into HEK293T cells along with the packaging plasmids pSPAX2 and pVSV-

G using a 4:3:1 ratio. Stable transfectants were selected using 1 μg/ml puromycin. HeLa

cells were cultured at 37°C in DMEM supplemented with 10% FBS and 1% penicillin-

streptomycin. A similar approach was employed to express MCU WT and MCU HAAF

in the HeLa cell line. Initially, HeLa cells were infected with the pLKO.1-shMCU

lentivirus and subsequently selected with 1 μg/ml puromycin. Knockout HeLa cells

were then infected with lentivirus carrying pLVX-mCherry-N1 vectors harboring either

MCU^{WT} or MCU^{HAAF}. Cells expressing mCherry were isolated using flow cytometry.

Knockout of endogenous MCU and expression of exogenous MCU-mCherry were

confirmed by Western blot analysis. All other mammalian cell lines used in this study

were from the American Type Culture Collection. All the cell lines were fully

authenticated and tested free of mycoplasma.

Mitochondrial Ca^{2+} modulator Screening

Hela cells expressing 4mt-GCaMP6s were permeabilized by a 1 min perfusion

with intracellular buffer (130 mM KCl, 10 mM NaCl, 2 mM K_2HPO_4, 5 mM Succinic

acid, 5 mM Malic acid, 1 mM Pyruvate, 0.5 mM ATP, 0.1 mM ADP, $MgCl_2$, 20 mM

HEPES, pH 7, 37 ℃) supplemented with 10 μM digitonin and 50 μM EGTA. Then the

buffer was changed to intracellular buffer supplemented with 10 μM compounds from

primary screen hits of FDA-approved drugs for 10 min. Mitochondrial Ca^{2+} uptake was

evoked by adding 100 μM $CaCl_2$ to the buffer. Ca^{2+} imaging was performed using a

DeltaVision Deconvolution microscope (GE Healthcare) equipped with a ×20 objective.

Exposure time was set to 25 ms for FITC. Images were acquired at 1 frame per 3

seconds with an EM-CCD camera. Images were analysed using Image J software. All

images were background-corrected by subtracting mean pixel values of a cell-free

region of interest.

Molecular dynamics (MD) simulations

Structural data for MCU and EMRE monomers were retrieved from the Protein

Data Bank (PDB) via the entry code 6XJV. The preprocessing phase employed

Gromacs 2018 as the simulation suite, coupled with the Amber99SB-ildn force field for

encompassing protein and small molecule interactions. A $10 \times 10 \times 10$ nm^3 water box,

utilizing the TIP3P model, was constructed to solvate the system, ensuring a minimal

1.2 nm buffer between the protein surfaces and the box boundaries. This was followed

by ion addition for electrostatic equilibrium.

For treating electrostatic interactions, we implemented the Particle-mesh Ewald

(PME) method. Energy minimization was executed through the steepest descent

method over 50,000 steps, setting both the Coulombic and van der Waals interaction

cutoffs at 1 nm. Equilibration phases under NVT (constant number of particles, volume,

and temperature) and NPT (constant number of of particles, pressure, and temperature)

settings preceded a 100 ns MD simulation, conducted under standard laboratory

conditions of temperature and pressure. Non-bonded interactions were applied with a

10 Å cutoff. Thermal and pressure stability were maintained at 300 K and 1 bar,

respectively, utilizing the V-rescale thermostat and the Berendsen barostat.

This rigorous preparation and execution strategy yielded a robust framework for

accurately simulating the complex's behavior in a physiological context, culminating in

the acquisition of data such as radius of gyration (Rg), root-mean-square deviation

(RMSD), number of hydrogen bonds (H-bonds), and total free energy, to assess

structural stability, conformational dynamics, interaction strength, and energetics

throughout the simulation period.

Ca^{2+} measurements

For imaging Ca^{2+} transients evoked by treatments with an agonist, cells

labelled with Calbryte590 or 4mt-GCaMP6 were incubated at 37 °C with Krebs-Ringer

modified buffer (KRB), which contains 125 mM NaCl, 5 mM KCl, 1 mM Na_3PO_4, 1

mM $MgSO_4$, 5.5 mM glucose and 20 mM HEPES (pH 7.4). After 10 μM histamine

stimulation, a series of images were captured every 3 s using a DeltaVision

Deconvolution microscope (GE Healthcare). The recorded images were analyzed and

quantified using Image J software (NIH).

For image analysis, background correction was first performed frame by frame by

subtracting the intensity of a nearby cell-free region from the signal of the imaged cell.

F_0 is calculated as the initial background-subtracted fluorescence intensity. F indicates

the background-subtracted fluorescence intensity at each time point. $\Delta F/F_0 = (F - F_0)/F_0$,

indicates the Ca^{2+} concentration change after stimulation.

Mitochondrial membrane potential measurements

For confocal microscopy, adherent HeLa cells were stained with

Tetramethylrhodamine (TMRM) at a final concentration of 100 nM in fresh culture

medium. After a 30-minute incubation period at 37°C to allow for adequate dye uptake,

cells were treated with the compounds of interest for an additional 30 minutes at the

same temperature. Subsequently, cells were washed to remove excess dye and unbound

compounds. Imaging was performed on a DeltaVision confocal microscope (GE

Healthcare). Acquired images were then processed and quantified using ImageJ

software.

For flow cytometric analysis, Hela cells were digested and stained with 100 nM

Tetramethylrhodamine (TMRM) in fresh medium for 30 min at 37℃. Compounds were added for 30 min at 37℃. Then cells were washed twice and resuspended in PBS containing 7AAD. Samples were analyzed using cytoFLEX flow cytometer (Beckman Coulter).

Western blot analysis.

For western blot analysis, cells were lysed in RIPA buffer (50 mM Tris-HCl (pH 8.0), 150 mM NaCl, 1% (v/v) Nonidet P-40, 0.5% sodium deoxycholate, 0.1% SDS, protease inhibitor cocktail). Protein samples were resolved by SDS-PAGE and transferred onto PVDF membranes. Membranes were immunoblotted with the indicated primary antibodies. Antibodies were used at the following concentrations: anti-MCU, rabbit polyclonal, Cell Signaling Technology, Cat# 14997 (1:1000); anti-SNAP, rabbit polyclonal, New England Biolabs, Cat# P9310S (1:1000).

Immunoprecipitation

HEK293T cells were transfected with the indicated plasmids using Lipofectamine^{TM} 2000 transfection reagent (ThermoFisher, Cat# 11668500). After transfection for 24 h, cells were treated with 10 μM Berberine for 2 hours before harvested. Cells were lysed with RIPA buffer (50 mM Tris-HCl (pH 8.0), 150 mM NaCl, 1% (v/v) Nonidet P-40, 0.5% sodium deoxycholate, protease inhibitor cocktail). Cell lysates were immunoprecipitated by anti-Flag M2 affinity gel (Selleck, Cat# B23102) overnight at 4 ℃. Beads were washed 5 times in RIPA buffer followed by eluting bound protein complexes in 2 × SDS dye. Whole cell lysates and co-precipitation samples were analyzed by western blot.

Biotin pulldown assay

For *in vivo* Biotin pulldown assay, HEK293T cells were treated with Biotin or Biotin-labeled Berberine (10 μM, 2 h). Cells were lysed and harvested with RIPA buffer，which containing 50 mM Tris-HCl, 150 mM NaCl, 1% NP-40 and 0.5% sodium deoxycholate. Balanced Streptavidin Agarose beads (ThermoFisher, Cat# 20357) were added and incubated overnight at 4 ℃. The beads were washed with RIPA buffer adding 0.03% Tween 5 minutes for 5 times. The indicated proteins were detected by

Immunoblotting.

For the *in vitro* biotin pulldown assay, biotin or biotin-labeled berberine (2 µg) was

pre-incubated with either recombinant protein (2 µg) or in vitro transcribed/translated

protein (Promega, Cat# L1170) for 8 hours at 4°C. The mixture was then incubated with

streptavidin agarose beads (Thermo Fisher, Cat# 20357) overnight at 4°C. The beads

were washed as previously described and the bound proteins were analyzed by

immunoblotting.

Immunofluorescence and confocal microscopy

For visualizing the intracellular distribution of Berberine, HeLa cells were

grown on coverslips, incubated with 50 nM MitoTracker^{TM} DeepRed (ThermoFisher,

Cat# M22426) in fresh medium for 15 min at 37 °C. 10 µM Berberine was added for

10 min. After treatment, the medium was replaced with phenol red-free DMEM

supplemental with 10% FBS. Then cells were visualized using a DeltaVision

Deconvolution microscope (GE Healthcare). Images of Berberine and MitoTracker^{TM}

DeepRed signals, acquired by FITC and Cy5 filters respectively, were merged to

visualize their co-localization in mitochondria.

For visualizing the intracellular distribution of Biotin-labeled Berberine,

HeLa cells were grown on coverslips, incubated with 10 µM Biotin-Berberine for 10

min at 37 °C. After treatment, cells were fixed in 4% paraformaldehyde for 5 min. Next,

the cells were incubated with 0.3% Triton X-100 in PBS for 10 min on ice and then

blocked in 3% BSA in PBS for at least 1 h. The primary antibody was incubated

overnight at 4°C followed by secondary antibody incubation. Finally, the cells were

mounted using 80% glycerol. Antibodies were used at the following concentrations:

anti-COXIV, rabbit polyclonal, Cell Signaling Technology, Cat# 4850 (1:200); Alexa

Fluor 488-conjugated streptavidin antibody, ThermoFisher, Cat# S11223 (1:400);

Alexa Fluor 546-conjugated anti-Rabbit Secondary Antibody, ThermoFisher, Cat#

A11035 (1:400). Images were acquired using a DeltaVision Deconvolution microscope.

Mitochondrial isolation and swelling assay

172   For isolating cardiac mitochondria, mouse hearts were minced in

mitochondrial isotonic buffer (225 mM mannitol, 75 mM sucrose, 5 mM MOPS, 0.5 mM EGTA and 2 mM taurine, pH 7.25) supplemented with protease inhibitor cocktail, and subsequently homogenized using a dounce homogenizer. The homogenate was initially centrifuged for 10 min at 700 $\times$ g and the resulting supernatant was next spun for 10 min at 12,000 $\times$ g to pellet the mitochondria.

Mitochondrial swelling was measured as previously described (3). In brief, isolated cardiac mitochondria (100 μg) were resuspended in swelling buffer (120 mM KCl, 10 mM Tris-HCl, 5 mM MOPS, 5 mM Na_2HPO_4, 10 mM glutamate, 2 mM malate and 0.1 mM EGTA) in a total volume of 200 μl. Pore opening was induced and detected by the addition of 500 μM of total $CaCl_2$ while monitoring absorbance at 540 nm. Where indicated, Berberine (10 μM), cyclosporin A (3 μM) or Ru360 (3 μM) was added.

Isolation and culture of primary neonatal rate cardiomyocytes (NRCMs)

NRCMs were isolated and cultured as described previously (4). Briefly, cardiomyocytes were isolated from the neonatal hearts of <24h-old Sprague-Dawley rats using PBS containing 0.03% trypsin and 0.04% collagenase type II. The digestion was halted by Dulbecco's modified Eagle's medium (DMEM), supplemented with 10% fetal bovine serum (FBS). After differential attachment for 90 min, the cells in suspension were then plated and cultured in DMEM with 10% FBS at 37°C with 5% CO_2 for 24 h.

Hypoxia/reoxygenation (H/R) injury model

The hypoxia/reoxygenation (H/R) injury of NRCMs or H9c2 cells were achieved in incubator (Thermo Fisher Scientific, Madison, WI, USA) filled with 94 % N_2, 5 % CO_2, and 1% O_2 in DMEM deprived of glucose and sodium pyruvate for 4h, followed by 8 h-reoxygenation by transferring the cells to a normoxia incubator with normal supplement of oxygen and glucose.

To examine the protective effect of BBR on H/R-induced Ca^{2+} overload, NRCMs infected with Ad-4mt-GCaMP6 or H9c2 cells stained with Rhod2 were treated with BBR at a final concentration 10 μM throughout the H/R period. Then cells were

visualized using a DeltaVision Deconvolution microscope (GE Healthcare). To assess

mitochondrial Ca^{2+} level using Rhod2, cells labeled with Rhod2 were illuminated at

546 nm, and fluorescence was collected through a TRITC filter. To assess mitochondrial

Ca^{2+} level using 4mt-GCaMP6, cells labeled with 4mt-GCaMP6 were illuminated at

488 and 390 nm, and fluorescence was collected through a FITC filter. Images were

analysed using Image J software. All images were background-corrected by subtracting

mean pixel values of a cell-free region of interest.

Statistics and reproducibility.

Statistical comparisons between only two groups were carried out using unpaired

Student's t test or the Mann-Whiney U test when a normal distribution could not be

assumed. One-way analysis of variance (ANOVA) was used for multiple-group

comparisons. Statistical calculations were carried out using GraphPad Prism 6.0. We

tested data for normality and variance, and considered a P value of less than 0.05 as

significant. All experiments were performed three or more times independently under

identical or similar conditions.

A

B

Compound (10 μM)	Inhibition (%)	*P* value
Berberine	87.21264	< 0.0001
Trifarotene	79.77925	< 0.0001
Ru360	71.36228	< 0.0001
Eltrombopag	67.76502	< 0.0001
Tipranavir	55.00618	< 0.0001
Oxyphenisatine	47.85387	< 0.0001
Ramatroban	44.67031	< 0.0001

Supplementary Figure 1. Screening identifies Berberine as an inhibitor of mitochondrial Ca^{2+} uptake.

Supplementary Figure 2. Berberine selectively inhibits mitochondrial Ca²⁺ uptake without impairing cytosolic Ca²⁺ signaling, mitochondrial membrane potential, or cell viability.

Supplementary Figure 3. Berberine directly targets MCU.

Supplementary Figure 4. Berberine disrupts MCU-EMRE assembly.

A

H/R	-	-	+	+
BBR	-	+	-	+

Rhod2

B

p=0.0008

Rel. Rhod2 fluorescence

H/R	-	-	+	+
BBR	-	+	-	+

Supplementary Figure 5. Berberine attenuates mitochondrial Ca^{2+} overload in cardiomyocytes under pathological conditions.

Supplementary Figure Legends

Supplementary Figure 1. Screening identifies Berberine as an inhibitor of mitochondrial Ca²⁺ uptake.

(A) Traces of mitochondrial Ca²⁺ level (indicated by fluorescence of 4mt-GCaMP6) in digitonin-permeabilized HeLa cells upon the addition of Ca²⁺ in the presence of DMSO, Ru360, or drugs.

(B) The list of the top 7 compounds with mitochondrial Ca²⁺ uptake inhibitory activity.

Supplementary Figure 2. Berberine selectively inhibits mitochondrial Ca²⁺ uptake without impairing cytosolic Ca²⁺ signaling, mitochondrial membrane potential, or cell viability.

(A) Representative images of Berberine (left panel) treated HeLa cells co-stained with MitoTracker DeepRed (middle panel). The merged image is shown in the right panel. Scale bar, 10 μm.

(B) Traces of normalized fluorescence intensity spatial profiles through the white line shown in **A**.

(C) Representative traces of cytosolic Ca²⁺ level (indicated by GCaMP6) in 10 μM Histamine-stimulated HeLa cells pretreated with different concentration of Berberine. Data are shown as the mean ± s.e.m. n = 50 cells for each condition.

(D) Measurement of the maximal amplitudes of the cytosolic Ca²⁺ traces in **C**. Data are shown as the mean ± s.e.m. n = 50 cells for each condition. *P* value was analyzed using one-way ANOVA.

(E) Representative traces of mitochondrial Ca²⁺ level (indicated by 4mt-GCaMP8) in 10 μM Histamine-stimulated HeLa cells pretreated with different concentration of

Berberine. Data are shown as the mean $\pm$ s.e.m. n = 50 cells for each condition.

(F) Measurement of the maximal amplitudes of the mitochondrial Ca^{2+} traces in **E**.

Data are shown as the mean $\pm$ s.e.m. n = 50 cells for each condition. *P* value was

analyzed using one-way ANOVA.

(G) Flow cytometry analysis of TMRM staining in HeLa cells pretreated with

different concentrations of Berberine and 50 μM CCCP. Shown is one experiment

that is representative of three similar experiments.

(H) Quantitative analysis of relative fluorescence of TMRM staining after different

concentrations of Berberine treatment and 50 μM CCCP treatment. Data are

shown as the mean $\pm$ s.e.m. n = 3 independent experiments.

(I) Cell viability of HeLa cells, U2OS cells and HEK293T cells after different

concentration of Berberine treatment was assessed at indicated time and

normalized to day 0 in each group. Data are shown as the mean $\pm$ s.e.m. n = 3.

Supplementary Figure 3. Berberine directly targets MCU.

(A) Representative traces of mitochondrial Ca^{2+} level (4mt-GCaMP6) dynamics in 10

μM Histamine-stimulated HeLa cells pretreated with Vehicle, 10 μM Biotin-

Berberine or 10 μM Berberine. Data are shown as the mean $\pm$ s.e.m. n = 40 cells

for each condition.

(B) Measurement of the maximal amplitudes of the mitochondrial Ca^{2+} traces in **A**.

Data are shown as the mean $\pm$ s.e.m. n = 40 cells for each condition. *P* value was

analyzed using one-way ANOVA.

(C) Western Blot analysis of Biotin-Berberine pulldown assay in MCU-Flag

overexpressed HEK293T cells. Cells were treated with 2 h 10 μM Biotin or 10

μM Biotin-Berberine.

(D) Two-dimensional diagrams illustrating the binding mode of berberine with MCU.

(E) Sequence alignment of the putative region (amino acids 275-295) surrounding the
    JML domain of MCU proteins from 16 diverse species.

Supplementary Figure 4. Berberine disrupts MCU-EMRE assembly.

(A) Western Blot analysis of co-precipitation of the Flag-EMRE with SNAP-MCU in
    HEK293T cells in the presence of 10 μM Berberine or not for 2 hours.
(B) Molecular dynamics simulations illustrating the assembly conformations of the
    MCU-EMRE complex in the presence and absence of Berberine.
(C) Molecular dynamics simulations reveal the hydrogen bonding interactions
    between MCU and EMRE under conditions with and without Berberine.
(D) Molecular dynamics simulations displaying the number of hydrogen bonds
    between MCU and EMRE in the presence and absence of Berberine.
(E) Molecular dynamics simulations demonstrating the binding free energy between
    MCU and EMRE with and without Berberine. The components of the binding free
    energy are detailed as follows: van der Waals energy (VDWAALS), electrostatic
    energy (EEL), polar solvation energy (EGB), non-polar solvation energy
    (ESURF), total gas phase free energy (GGAS, calculated as the sum of
    VDWAALS and EEL), total solvation free energy (GSOLV, calculated as the sum
    of EGB and ESURF), and the total free energy (TOTAL, calculated as the sum of
    GSOLV and GGAS).

Supplementary Figure 5. Berberine attenuates mitochondrial Ca^{2+} overload in cardiomyocytes under pathological conditions.

(A-B) Measurement of mitochondrial Ca^{2+} level (indicated by Rhod2) in H9C2 cells.
Cells were pretreated with 10 μM Berberine or not, and then were subjected to hypoxia
(3 h) followed by reoxygenation (12 h) (H/R). Relative fluorescence intensity of Rhod2
were measured in **B**. Data are shown as the mean ± s.e.m. n = 30 cells (Control and

295   BBR groups), n = 40 cells (H/R and H/R+BBR groups). *P* value was analyzed using

296   one-way ANOVA. Scale bar, 10 μm.

Reference (40-42):

40. Zhang J, *et al.* (2022) Novel CaMKII-delta Inhibitor Hesperadin Exerts Dual Functions to Ameliorate Cardiac Ischemia/Reperfusion Injury and Inhibit Tumor Growth. *Circulation* 145(15):1154-1168.

41. Trott O & Olson AJ (2010) AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. *J Comput Chem* 31(2):455-461.

42. Lan C, *et al.* (2022) Inhibition of DYRK1A, via histone modification, promotes cardiomyocyte cell cycle activation and cardiac repair after myocardial infarction. *EBioMedicine* 82:104139.